

RESULTS FROM THE 2014 NATIONAL SURVEY ON DRUG USE AND HEALTH: DETAILED TABLES

PREVALENCE ESTIMATES, STANDARD ERRORS, *P* VALUES, AND SAMPLE SIZES

- Section 1: Illicit Drug Use Tables – 1.1 to 1.92**
- Section 2: Tobacco Product and Alcohol Use Tables – 2.1 to 2.84**
- Section 3: Risk and Protective Factor Tables – 3.1 to 3.25**
- Section 4: Incidence Tables – 4.1 to 4.16**
- Section 5: Dependence, Abuse, and Treatment Tables – 5.1 to 5.56**
- Section 6: Miscellaneous Tables – 6.1 to 6.107**
- Section 7: Trend Tables – 7.1 to 7.80**
- Section 8: Sample Size and Population Tables – 8.1 to 8.13**

DISCLAIMER

SAMHSA provides links to other Internet sites as a service to its users and is not responsible for the availability or content of these external sites. SAMHSA, its employees, and contractors do not endorse, warrant, or guarantee the products, services, or information described or offered at these other Internet sites. Any reference to a commercial product, process, or service is not an endorsement or recommendation by SAMHSA, its employees, or contractors. For documents available from this server, the U.S. Government does not warrant or assume any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed.

Substance Abuse and Mental Health Services Administration
Center for Behavioral Health Statistics and Quality
Rockville, Maryland 20857

RESULTS FROM THE 2014 NATIONAL SURVEY ON DRUG USE AND HEALTH: DETAILED TABLES

PREVALENCE ESTIMATES, STANDARD ERRORS, *P* VALUES, AND SAMPLE SIZES

- Section 1: Illicit Drug Use Tables – 1.1 to 1.92
- Section 2: Tobacco Product and Alcohol Use Tables – 2.1 to 2.84
- Section 3: Risk and Protective Factor Tables – 3.1 to 3.25
- Section 4: Incidence Tables – 4.1 to 4.16
- Section 5: Dependence, Abuse, and Treatment Tables – 5.1 to 5.56
- Section 6: Miscellaneous Tables – 6.1 to 6.107
- Section 7: Trend Tables – 7.1 to 7.80
- Section 8: Sample Size and Population Tables – 8.1 to 8.13

Contract No. HHSS283201300001C
Project No. 0213757.003.107.003
Deliverable No. 29

For questions about these tables, please e-mail Peter.Tice@samhsa.hhs.gov.

Prepared for Substance Abuse and Mental Health Services Administration,
Rockville, Maryland

Prepared by RTI International, Research Triangle Park, North Carolina

September 10, 2015

Recommended Citation: Center for Behavioral Health Statistics and Quality.
(2015). *2014 National Survey on Drug Use and Health: Detailed Tables*.
Substance Abuse and Mental Health Services Administration, Rockville, MD.

Acknowledgments

These tables were prepared by the Center for Behavioral Health Statistics and Quality (CBHSQ), Substance Abuse and Mental Health Services Administration (SAMHSA), U.S. Department of Health and Human Services (HHS), and by RTI International, a registered trademark and a trade name of Research Triangle Institute, Research Triangle Park, North Carolina. Work by RTI was performed under Contract No. HHSS283201300001C. Contributors and reviewers at RTI listed alphabetically include Jeremy Aldworth, Katherine J. Asman, Stephanie N. Barnett, Ellen Bishop, Michael R. Bradshaw, Kristen Brown, Lisa A. Carpenter, Pinliang (Patrick) Chen, Chuchun Chien, Elizabeth Copello, Devon S. Cribb, Lanting Dai, Teresa R. Davis, Susan L. Edwards, Glynis S. Ewing, Barbara J. Felts, Misty S. Foster, Peter A. Frechtel, Julia M. Gable, Gina S. Geercken, Harper Gordek, Rebecca A. Granger, Wafa Handley, Rachel Harter, Erica L. Hirsch, David Hunter (Project Director), Phillip S. Kott, Larry A. Kroutil, Jeffrey S. Laufenberg, Philip K. Lee, Dan Liao, Peilan Chen Martin, Dexter McNutt, Andrew S. Moore, Lisa E. Packer, Michael R. Pemberton, Brenda K. Porter, Jeremy D. Porter, Rosanna S. Quiroz, Jessica Roycroft, Neeraja S. Sathe, Jennifer H. Schoden, Kathryn Spagnola, Paxton Syrek, Jiantong (Jean) Wang, Lauren K. Warren, Matthew J. Westlake, Cherie J. Winder, Carol L. Woodell, and Amy Zemanick. Contributors at SAMHSA listed alphabetically include Rebecca Ahrnsbrak, Peggy Barker (contractor), Jonaki Bose, Beth Han, Sarra L. Hedden, Elizabeth Hoeffel, Arthur Hughes, Joel Kennet, Rachel Lipari, Grace E. Medley, Pradip Muhuri, Dicy Painter, Kathryn Piscopo, Peter Tice (Project Officer), and Matthew Williams. At RTI, Farrah Bullock Mann and Brenda K. Porter formatted the tables; Melissa H. Hargraves, Laura James, Amber M. Rohloff, Margaret A. Smith, and Richard S. Straw provided editorial assistance; and Teresa F. Bass, Debbie F. Bond, Kimberly H. Cone, Daniel Occoquan, Pamela Couch Prevatt, Roxanne Snaauw, and Pamela Tuck provided web conversion and/or document production support.

Introduction

Results from the 2014 National Survey on Drug Use and Health: Detailed Tables is a collection of tables presenting national estimates from the National Survey on Drug Use and Health (NSDUH). These tables present information on drug, alcohol, and tobacco use, as well as drug and alcohol dependence and abuse and treatment. Measures of these behaviors and characteristics are presented by a variety of demographic, geographic, and other variables. The estimates in the tables include rates of the behaviors, numbers of persons engaging in these behaviors, and other measures.

Although the majority of these tables are trend tables presenting estimates from the 2013 and 2014 NSDUHs, some tables include only estimates from the 2014 NSDUH. Also, a number of tables contain annual averages that are generated by combining multiple years of data if sufficient data within a single year are unavailable to produce reliable estimates. These tables may provide annual averages combining data for (a) 2013 and 2014; (b) 2011 and 2012; and (c) 2011, 2012, 2013, and 2014. Also included are a number of tables that present data from earlier surveys in the series, including a section of tables presenting data mainly from 2002 to 2014 and a couple of tables within that section presenting data from 1971 to 2014.

Because of survey improvements in the 2002 NSDUH, the 2002 data constitute a new baseline for tracking trends in substance use and other measures. Therefore, estimates from the 2002 through 2014 surveys should not be compared with estimates from the 2001 or earlier surveys to examine changes over time. Methodology changes throughout NSDUH's history make it difficult to assess long-term trends from tables presenting data from 1971 to 2014. However, it is instructive to compare NSDUH estimates from 1971 to 2014 by "piecing together" the data from time periods for which data are comparable. Specifically, valid trend comparisons can be made for 1971-1998, 1999-2001, and 2002-2014. With this approach, comparisons between 1998 and 1999, and between 2001 and 2002, should be made with caution because they are potentially biased due to changes in methods. Nevertheless, when these data are combined in a single presentation, it often becomes clear that the effects of the changes in methods are small compared with the major shifts in substance use prevalence that have occurred over the past four decades.

For the 2014 NSDUH, an investigation was completed to look at missing data rates in the 2014 detailed tables. The investigation concluded that missing data are not a concern for most topics in the 2014 detailed tables. However, a few items do have some missing data (e.g., items on perceived availability of various illicit drugs). For example, the rates of missing data for these availability questions ranged from 3.5 to 7.4 percent in 2014. Respondents with missing data are generally excluded from the relevant analyses.

A description of the survey measurement issues and the sample design and estimation procedures used in the 2014 NSDUH can be found in an online report titled the *2014 National*

it would be to obtain these drugs are not likely to be interested in using them. Note that detailed tables with missing data include a note for reference.

A description of the survey measurement issues and the sample design and estimation procedures used in the 2014 NSDUH can be found in an online report titled the *2014 National Survey on Drug Use and Health: Methodological Summary and Definitions*.¹ Key definitions (i.e., the glossary) for many of the measures and terms used in these detailed tables and in other 2014 NSDUH documents can be located in Section C of that online report.² Where relevant, the glossary provides cross-references between terms and specific question wording for clarity.

Along with these 2014 detailed tables, tables presenting the prevalence of mental health problems, related treatment, and the co-occurrence of substance use and mental health problems are being released separately as the *Results from the 2014 National Survey on Drug Use and Health: Mental Health Detailed Tables*. These two sets of tables and four first release reports (FRRs) that are focused on specific behavioral health topics in the 2014 NSDUH, such as trends in substance use and mental health issues, are scheduled to be made available online by the end of September 2015 at <http://www.samhsa.gov/data/>.

Table Revisions

During regular data collection and processing checks for the 2011 NSDUH, data errors were identified. These errors affected the data for Pennsylvania (2006-2010) and Maryland (2008-2009). Cases with erroneous data were removed from the data files, and the remaining cases were reweighted to provide representative estimates. The errors had minimal impact on the national estimates and no effect on direct estimates for the other 48 states and the District of Columbia. In reports where model-based small area estimation techniques are used, estimates for all states may be affected, even though the errors were concentrated in only two states. In reports that do not use model-based estimates, the only estimates appreciably affected are estimates for Pennsylvania, Maryland, the mid-Atlantic division, and the Northeast region.

The 2014 detailed tables and 2014 FRRs do not include State-level or model-based estimates. However, the 2014 detailed tables do include estimates for the mid-Atlantic division and the Northeast region. Estimates based on 2006-2010 data may differ from previously published estimates. Tables and estimates based only on data since 2011 are unaffected by these data errors. All affected tables (i.e., tables with estimates based on 2006-2010 data) contain a note to indicate this to the user.

Caution is advised when comparing data from older reports with data from more recent reports that are based on corrected data files. As discussed above, comparisons of estimates for Pennsylvania, Maryland, the mid-Atlantic division, and the Northeast region are of most concern, while comparisons of national data or data for other states and regions are essentially still valid. The Center for Behavioral Health Statistics and Quality within the Substance Abuse

¹ Center for Behavioral Health Statistics and Quality. (2015). *2014 National Survey on Drug Use and Health: Methodological summary and definitions*. Retrieved from <http://www.samhsa.gov/data/>

² Prior to the 2010 NSDUH, one glossary of key definitions for both substance use and mental health terms and other topics appeared as an appendix in each year's national findings report. Between 2010 and 2013, separate glossaries appeared with the annual detailed tables and mental health detailed tables.

and Mental Health Services Administration does not recommend making comparisons between unrevised 2006-2010 estimates and estimates based on data for 2011 and subsequent years for the areas of greatest concern.

Table Numbering

The detailed tables are numbered using a three-part numbering scheme (e.g., 1.15A). The first part of the table number (1.15A) is the subject matter section to which a particular table belongs. The second part (1.15A) is the number of the table within a particular section. The third part (1.15A) is a table type indicator, an alphabetic letter appended to the table number. Each table number, as explained below, has multiple table types. Tables are numbered sequentially within each subject matter section. To the extent possible, identical tables are assigned the same table number each year except in the case where specific tables are removed or added.

The eight subject matter sections and the number of tables per section are as follows:

Section 1: Illicit Drug Use Tables – 1.1 to 1.92

Section 2: Tobacco Product and Alcohol Use Tables – 2.1 to 2.84

Section 3: Risk and Protective Factor Tables – 3.1 to 3.25

Section 4: Incidence Tables – 4.1 to 4.16

Section 5: Dependence, Abuse, and Treatment Tables – 5.1 to 5.56

Section 6: Miscellaneous Tables – 6.1 to 6.107

Section 7: Trend Tables – 7.1 to 7.80

Section 8: Sample Size and Population Tables – 8.1 to 8.13

The table type indicators are primarily defined as follows; however, some exceptions do exist and are noted in subsequent bullets.

Table Type Purpose of the Table

- A: Presents estimates of the numbers of persons exhibiting the specified behavior or characteristic (e.g., substance use) in the populations described by the column and row headings.
- B: Presents estimates of the percentages of persons exhibiting the specified behavior or characteristic (e.g., substance use) in the populations described by the column and row headings.
- C: Presents the standard error associated with each of the estimates in the "A" tables.
- D: Presents the standard error associated with each of the estimates in the "B" tables.
- N: Presents the number of cases in the specified NSDUH sample with the characteristics defined by the column and row headings.
- P: Presents the *p* values from tests of the statistical significance of differences between columns in the "B" tables.

The majority of tables within the detailed tables contain five table types (A, B, C, D, and P) as defined above. Note that table type N is used exclusively within Section 8 to display the sample size counts. Exceptions to this organization are noted as follows:

- Section 3 (Risk and Protective Factor Tables) includes tables for which only table types A, B, C, and D are produced (Tables 3.19 to 3.25). These tables present estimates for the 2014 NSDUH only; thus, these are not trend tables, so significance tests between years are not relevant and table type P is not applicable.
- Section 4 (Incidence Tables) contains both tables for which only table types A and C are produced and tables for which only table types B, D, and P are produced. Specifically, one subset of tables (Tables 4.9 to 4.12) includes table types A and C. This subset of tables presents the number of past year initiates in table type A and the associated standard errors in table type C. The remaining Section 4 tables include only table types B, D, and P. Tables 4.1 to 4.8 present numbers and percentages of past year initiates in table type B, associated standard errors for each of these estimates in table type D, and *p* values from tests of statistical differences between years in table type P. Tables 4.13 to 4.16 present the mean age for past year initiates in table type B, associated standard errors in table type D, and *p* values from tests of statistical differences between years in table type P.
- Section 5 (Dependence, Abuse, and Treatment Tables) includes tables for which only table types A, B, C, and D are produced (Tables 5.54 to 5.56). These tables present annual averages based on combined 2011-2014 NSDUH data; thus, these are not trend tables, so significance tests between years are not relevant and table type P is not applicable.
- Section 7 (Trend Tables) contains one subset of tables for which only table types B and D are produced and one subset of tables for which only table types B, D, and P are produced. Specifically, one subset of tables (Tables 7.79 and 7.80) are multiyear tables that present estimated percentages for 1971 to 2014 in table type B and associated standard errors in table type D. The estimated total number of users and associated standard errors are not presented, nor are between-year significance tests implemented; thus, table types A, C, and P are not used. The other subset of tables (Table 7.49) includes multiyear tables that involve only table types B, D, and P. Table 7.49 presents the mean age for past year initiates in table type B, associated standard errors for each of these estimates in table type D, and *p* values from tests of statistical differences between years in table type P.
- Section 8 (Sample Size and Population Tables) contains only table types A, C, and N. Population counts, standard errors, and sample sizes are displayed in table types A, C, and N, respectively. Percentages of the population and associated standard error tables are not provided in this section.

Locating a Table

The detailed tables consist of prevalence estimates (table types A and B), standard errors and *p* values (table types C, D, and P), and sample sizes (table type N). The tables (all types) are organized by subject matter sections. Eight subject matter sections are available for 2014, and

two methods can be used to locate and access a table of interest within the PDF version of the detailed tables.

Subject Matter Sections

If a user knows which of the eight subject matter sections is of interest, then he or she can click directly on the bookmark for that subject matter section. This will take the user to the index of tables for that section. This index can be used to identify the group of tables in which the desired table is likely to be located. Clicking on the appropriate link will take the user directly to the first table in the specified group where he or she then can scroll through the tables to locate the table of interest. The group links will provide the user access to the prevalence estimate tables. To access the corresponding standard error and p value tables, a user must click on the "SE" link associated with that group link. The eight subject matter sections are as follows:

Index of Section 1: Illicit Drug Use Tables (1.1 to 1.92)

Index of Section 2: Tobacco Product and Alcohol Use Tables (2.1 to 2.84)

Index of Section 3: Risk and Protective Factor Tables (3.1 to 3.25)

Index of Section 4: Incidence Tables (4.1 to 4.16)

Index of Section 5: Dependence, Abuse, and Treatment Tables (5.1 to 5.56)

Index of Section 6: Miscellaneous Tables (6.1 to 6.107)

Index of Section 7: Trend Tables (7.1 to 7.80)

Index of Section 8: Sample Size and Population Tables (8.1 to 8.13)

Reference Tools

If a user needs more detailed information on which subject matter section to enter, he or she can click on the "Reference Tools" bookmark to learn about various other tools available for locating a specific table of interest. These tools include a key to selected variables, key definitions for the 2014 NSDUH (i.e., the glossary), and a list of tables.

The key to selected variables defines key topics (e.g., demographics) used throughout the 2014 detailed tables. As mentioned earlier, the glossary defines and cross-references key measures used in the 2014 detailed tables and the 2014 FRRs. Both the key and the glossary may be helpful when used in conjunction with the table index in identifying tables that contain information for selected measures. However, not all measures and terms listed in the key and the glossary are specifically mentioned in the table index. The list of tables can be used to identify a specific table for one of the subject matter sections. Once the desired table has been located using the reference tools, the user then should follow the directions provided earlier in the section on the "Subject Matter Sections" to access the table.

Note that once a group of prevalence estimate tables is accessed, there is no direct link to the corresponding standard error and p value tables (and vice versa). If the user wishes to see the corresponding prevalence estimate, standard error and p value, or sample size and population

tables, it will be necessary first to click on the appropriate subject matter bookmark, then to select the new group that contains the table of interest.

Index of Section 1 Tables: Illicit Drug Use (1.1 to 1.92)

Illicit Drug Use Table	Table Number	Link to Corresponding Standard Error Table
Lifetime, past year, and past month illicit drug use, by selected age categories and gender	1.1 to 1.10	SE
Selected illicit drugs, by detailed age categories	1.11 to 1.18	SE
Selected illicit drugs, by demographic characteristics	1.19 to 1.58	
Illicit drugs	1.19 to 1.23	SE
Marijuana	1.24 to 1.28	SE
Cocaine	1.29 to 1.33	SE
Crack	1.34 to 1.38	SE
Hallucinogens	1.39 to 1.43	SE
Inhalants	1.44 to 1.48	SE
Nonmedical use of prescription-type psychotherapeutics	1.49 to 1.53	SE
Nonmedical use of pain relievers	1.54 to 1.58	SE
Selected illicit drugs, by geographic characteristics	1.59 to 1.87	
Illicit drugs	1.59 to 1.63	SE
Marijuana	1.64 to 1.68	SE
Cocaine	1.69 to 1.73	SE
Crack	1.74	SE
Hallucinogens	1.75	SE
Inhalants	1.76 to 1.77	SE
Nonmedical use of prescription-type psychotherapeutics	1.78 to 1.82	SE
Nonmedical use of pain relievers	1.83 to 1.87	SE
Specific hallucinogen, inhalant, needle, heroin, pain reliever, tranquilizer, stimulant, and sedative use, by selected age categories	1.88 to 1.92	SE

Index of Section 2 Tables: Tobacco Product and Alcohol Use (2.1 to 2.84)

Tobacco Product and Alcohol Use Table	Table Number	Link to Corresponding Standard Error Table
Lifetime, past year, and past month tobacco product and alcohol use, by selected age categories and gender	2.1 to 2.10	SE
Selected tobacco product and alcohol use, by detailed age categories	2.11 to 2.16	SE
Selected tobacco product and alcohol use, by demographic characteristics	2.17 to 2.46	
Tobacco products	2.17 to 2.21	SE
Cigarettes	2.22 to 2.26	SE
Smokeless tobacco	2.27 to 2.31	SE
Cigars	2.32 to 2.36	SE
Alcohol	2.37 to 2.46	SE
Selected tobacco product and alcohol use, by geographic characteristics	2.47 to 2.76	
Tobacco products	2.47 to 2.51	SE
Cigarettes	2.52 to 2.56	SE
Smokeless tobacco	2.57 to 2.61	SE
Cigars	2.62 to 2.66	SE
Alcohol	2.67 to 2.76	SE
Underage and legal drinkers' alcohol use, by demographic or geographic characteristics	2.77 to 2.84	SE

Index of Section 3 Tables: Risk and Protective Factors (3.1 to 3.25)

Risk and Protective Factor Table	Table Number	Link to Corresponding Standard Error Table
Perceived risk and availability of substances	3.1 to 3.5	
Perceived risk and availability, by selected age categories	3.1	SE
Perceived risk, by demographic and geographic characteristics	3.2 to 3.4	SE
Perceived availability, by demographic and geographic characteristics	3.5	SE
Risk and protective factors, by demographic and geographic characteristics	3.6 to 3.18	
Disapproval of substance use behavior	3.6 to 3.8	SE
Prevention programs/messages	3.9 to 3.10	SE
Delinquent behavior	3.11 to 3.12	SE
Selected youth behavior	3.13	SE
Religious beliefs	3.14	SE
Parental involvement in specific behaviors	3.15	SE
Student characteristics	3.16	SE
Adult involvement in specific school-related behaviors	3.17	SE
Substance use of students in grade level	3.18	SE
Past month substance use, by risk and protective factors	3.19 to 3.25	
Perceived risk and availability of substances	3.19 to 3.20	SE
Disapproval of substance use behavior	3.21 to 3.22	SE
Prevention programs/messages and delinquent behavior	3.23	SE
Selected youth behavior, religious beliefs, and parental involvement in specific behaviors	3.24	SE
Student characteristics, adult involvement in specific school-related behaviors, and substance use of students in grade level	3.25	SE

Index of Section 4 Tables: Incidence (4.1 to 4.16)

Incidence Table	Table Number	Link to Corresponding Standard Error Table
Past year initiation of substance use	4.1 to 4.16	
Initiation of selected substances among total population, persons at risk for initiation, and past year users, by selected age categories and gender	4.1 to 4.4	SE
Initiation among total population, persons at risk for initiation, and past year users	4.5 to 4.8	SE
Initiation among persons aged 12 or older, by gender	4.9	SE
Initiation among persons aged 12 or older who initiated use prior to a selected age, by gender	4.10 to 4.11	SE
Initiation among persons aged 12 to 49, by gender	4.12	SE
Mean age at first substance use among past year initiates aged 12 or older, by gender	4.13	SE
Mean age at first substance use among past year initiates who initiated use prior to a selected age, by gender	4.14 to 4.15	SE
Mean age at first substance use among past year initiates aged 12 to 49, by gender	4.16	SE

Index of Section 5 Tables: Dependence, Abuse, and Treatment (5.1 to 5.56)

Dependence, Abuse, and Treatment Table	Table Number	Link to Corresponding Standard Error Table
Substance dependence or abuse	5.1 to 5.13	SE
Substance dependence	5.14	SE
Received substance use treatment	5.15 to 5.26	
Treatment, by demographic or geographic characteristics	5.15 to 5.21	SE
Specific substance for last or current treatment	5.22	SE
Location of treatment	5.23 to 5.25	SE
Treatment, by dependence and/or abuse	5.26	SE
Received substance use treatment at specialty facility	5.27 to 5.42	
Treatment, by demographic or geographic characteristics	5.27 to 5.33	SE
Specific substance for last or current treatment	5.34	SE
Source of payment for last or current treatment	5.35 to 5.37	SE
Location of treatment	5.38 to 5.40	SE
Treatment, by dependence and/or abuse	5.41 to 5.42	SE
Needed and received treatment or perceived need for treatment	5.43 to 5.53	
Illicit drug	5.43 to 5.47	SE
Alcohol	5.48 to 5.50	SE
Illicit drug or alcohol	5.51 to 5.53	SE
Reasons for not receiving substance use treatment	5.54 to 5.56	SE

Index of Section 6 Tables: Miscellaneous (6.1 to 6.107)

Miscellaneous Table	Table Number	Link to Corresponding Standard Error Table
Frequency of use	6.1 to 6.6	SE
Illicit drugs, by cigarette use	6.7 to 6.10	SE
Illicit drugs, by alcohol use	6.11 to 6.18	SE
Illicit drugs, by cigarette and alcohol use	6.19 to 6.23	SE
Tobacco product and alcohol, by cigarette use	6.24 to 6.27	SE
Tobacco product, by alcohol use	6.28 to 6.31	SE
Prior year and past year marijuana use status, by selected age categories and gender	6.32	SE
Nicotine (cigarette) dependence	6.33 to 6.37	SE
Substance dependence or abuse, by age at first use and demographic characteristics	6.38 to 6.40	SE
Substance dependence, by age at first use and demographic characteristics	6.41 to 6.43	SE
Obtaining marijuana	6.44 to 6.46	SE
Source for most recent nonmedical drug use (pain relievers, tranquilizers, stimulants, methamphetamine, and sedatives)	6.47 to 6.66	SE
Social context, location, and source of underage drinking	6.67 to 6.70	SE
Pregnancy	6.71 to 6.78	SE
College enrollment	6.79 to 6.93	SE
Driving under the influence	6.94 to 6.97	SE
Probation	6.98 to 6.102	SE
Parole/supervised release	6.103 to 6.107	SE

Index of Section 7 Tables: Trends (7.1 to 7.80)

Trend Table	Table Number	Link to Corresponding Standard Error Table
Substance use	7.1 to 7.35	
Lifetime, past year, and past month illicit drug use, by selected age categories and gender	7.1 to 7.21	SE
Lifetime, past year, and past month special drug use, by selected age categories and gender (2006-2014)	7.22 to 7.28	SE
Past month tobacco product and alcohol use, by selected age categories and gender	7.29 to 7.33	SE
Past month daily cigarette use, by selected age categories	7.34	SE
Lifetime, past year, and past month alcohol use, by underage drinkers and gender	7.35	SE
Risk and protective factors	7.36 to 7.43	
Perceived risk and availability of substances, by selected age categories	7.36 to 7.39	SE
Disapproval of substance use behavior	7.40 to 7.41	SE
Prevention programs/messages	7.42 to 7.43	SE
Initiation of substance use	7.44 to 7.49	
Past year initiation, by selected age categories	7.44 to 7.47	SE
First illicit drug initiated in the past year	7.48	SE
Mean age at first substance use among past year initiates of substance use aged 12 to 49	7.49	SE
Substance dependence or abuse	7.50 to 7.57	
Dependence or abuse	7.50 to 7.53	SE
Dependence	7.54 to 7.57	SE
Substance treatment	7.58 to 7.78	
Received treatment at any location or at a specialty facility	7.58 to 7.61	SE
Specific substance for last or current treatment	7.62 to 7.65	SE
Location of treatment	7.66 to 7.69	SE
Need for and receipt of treatment at a specialty facility	7.70 to 7.73	SE
Type of substance treatment at any location	7.74	SE
Type of substance treatment at a specialty facility	7.75	SE
Receipt of illicit drug treatment at a specialty facility	7.76	SE
Receipt of alcohol treatment at a specialty facility	7.77	SE
Receipt of illicit drug or alcohol treatment at a specialty facility	7.78	SE
1971-2014 past month marijuana and past year cocaine use	7.79 to 7.80	SE

NOTE: Estimates for 2002-2014 are presented in these tables unless otherwise noted above.

Index of Section 8 Tables: Sample Size and Population Estimates (8.1 to 8.13)

Sample Size and Population Table	Table Number	Link to Corresponding Standard Error Table
Demographic characteristics	8.1 to 8.5	SE
Geographic characteristics	8.6 to 8.8	SE
Selected age categories and gender (2002-2014)	8.9	SE
Pregnancy	8.10	SE
College enrollment	8.11	SE
Probation	8.12	SE
Parole/supervised release	8.13	SE

Reference Tools

To assist a user in locating a specific table of interest within the *Results from the 2014 National Survey on Drug Use and Health: Detailed Tables*, reference tools have been developed. Note that the following tools are provided solely for guidance when examining the 2014 detailed tables because there are no hyperlinks within these tools that lead directly to the detailed tables:

- [a key to selected variables](#);
- Key Definitions for the 2014 National Survey on Drug Use and Health (i.e., the glossary);
- [List of Tables: Prevalence Estimates and Sample Sizes - Table Types A, B, and N](#); and
- [List of Tables: Standard Errors and P Values - Table Types C, D, and P](#).

Hyperlinks that lead directly to the tables can be found under the "Tables" bookmark. How to locate a table by selecting the "Tables" bookmark is explained in the "Locating a Table" section of the introduction.

Using the Key to Selected Variables. The key to selected variables defines key topics (e.g., demographics) used throughout the 2014 detailed tables. For example, a user interested in information about the prevalence of illicit drug use by level of education can first review the key to selected variables and see that education is categorized as a demographic characteristic. Next, a user can determine, by looking at the bookmarks for the subject matter sections, that tables pertaining to illicit drug use can be found in Section 1: Illicit Drug Use Tables. By selecting this bookmark, the user will be taken to the table index for the illicit drug use tables. Now the user can scan the links to learn that data on the prevalence of illicit drug use by demographic characteristics are located in the group of tables numbered 1.19 to 1.23. Selecting the link to the left of the table numbers will take the user directly to the first table in the specified group where the user then can scroll through the tables to locate the table(s) pertaining to the prevalence of illicit drug use by level of education (Tables 1.21 to 1.23).

Using the Key Definitions for the 2014 National Survey on Drug Use and Health. The glossary defines and cross-references commonly used measures and terms in the 2014 detailed tables and the 2014 first release reports.¹ For example, a user interested in information about the prevalence of alcohol use measures by demographic characteristics can first review the alcohol use entry in the glossary to see a cross-reference of additional selected alcohol use measures included in the tables and reports from the 2014 National Survey on Drug Use and Health. After reviewing the definitions for the various measures, the user can select measures of interest (e.g., "binge use of alcohol"). Next, a user can determine, by looking at the bookmarks for the subject matter sections, that tables pertaining to alcohol use can be found in Section 2: Tobacco Product and Alcohol Use Tables. By selecting this bookmark, the user will be taken to

¹ The glossary is available online as Section C of the following report: Center for Behavioral Health Statistics and Quality. (2015). *2014 National Survey on Drug Use and Health: Methodological summary and definitions*. Retrieved from <http://www.samhsa.gov/data/>

the table index for tobacco product and alcohol use tables. Now the user can scan the links to learn that data on the prevalence of alcohol use by demographics are located in the group of tables numbered 2.37 to 2.46. Selecting the link to the left of the table numbers will take the user directly to the first table in the specified group where the user can scroll through the tables to locate the table(s) pertaining to the prevalence of "binge use of alcohol" by demographic characteristics (Tables 2.42 to 2.46).

Using the List of Tables. The list of tables for both prevalence estimates and sample sizes (table types A, B, and N) and standard errors and p values (table types C, D, and P) can be used to identify a specific table for one of the subject matter sections. For example, a user interested in information about the prevalence of past month illicit drug use among youths aged 12 or 13 can review the list of tables in Section 1: Illicit Drug Use Tables and identify Table 1.2B as the table of interest.

Key to Selected Variables

The following key defines selected variables (e.g., demographics) used throughout the *Results from the 2014 National Survey on Drug Use and Health: Detailed Tables*. This may be helpful when used in conjunction with the index of tables in identifying tables that contain information for these selected characteristics. However, not all variables listed below are specifically mentioned in the index of tables.¹

Demographic Characteristics

Standard Age Categories: 12 or Older, 12-17, 18-25, 26 or Older, 18 or Older

7 Detailed Age Categories: 12-13, 14-15, 16-17, 18-20, 21-25, 26-34, 35 or Older

23 Detailed Age Categories: 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26-29, 30-34, 35-39, 40-44, 45-49, 50-54, 55-59, 60-64, 65 or Older

Underage/Legal Drinking Age Categories: 12-20, 21 or Older

Gender: Male, Female

Hispanic Origin and Race: Not Hispanic or Latino White, Not Hispanic or Latino Black or African American, Not Hispanic or Latino American Indian or Alaska Native, Not Hispanic or Latino Native Hawaiian or Other Pacific Islander, Not Hispanic or Latino Asian, Not Hispanic or Latino Two or More Races, Hispanic or Latino, Not Hispanic or Latino

Gender/Race/Hispanic Origin: Male White Not Hispanic, Female White Not Hispanic, Male Black Not Hispanic, Female Black Not Hispanic, Male Hispanic, Female Hispanic

Education: Less Than High School, High School Graduate, Some College, College Graduate
(Note: This variable is defined for persons aged 18 or older.)

Current Employment: Full-Time, Part-Time, Unemployed, Other
(Note: This variable is defined for persons aged 18 or older.)

Family Income: Less Than \$20,000; \$20,000-\$49,999; \$50,000-\$74,999; \$75,000 or More

¹ Definitions for many of the measures and terms used in the detailed tables and reports from the 2014 National Survey on Drug Use and Health (NSDUH) are available online in Section C of the following report: Center for Behavioral Health Statistics and Quality. (2015). *2014 National Survey on Drug Use and Health: Methodological summary and definitions*. Retrieved from <http://www.samhsa.gov/data/>

Geographic Characteristics

Region: Northeast, Midwest, South, West

Division: New England, Middle Atlantic, East North Central, West North Central, South Atlantic, East South Central, West South Central, Mountain, Pacific

County Type: Large Metro counties; Small Metro counties, including counties with a population of 250,000 up to 1 million and counties with a population of less than 250,000; Nonmetro, including urbanized counties, less urbanized counties, or completely rural counties

LIST OF TABLES

Table

- 1.1A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2013 and 2014
- 1.1B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Percentages, 2013 and 2014
- 1.2A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Numbers in Thousands, 2013 and 2014
- 1.2B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Percentages, 2013 and 2014
- 1.3A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Numbers in Thousands, 2013 and 2014
- 1.3B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Percentages, 2013 and 2014
- 1.4A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Numbers in Thousands, 2013 and 2014
- 1.4B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Percentages, 2013 and 2014
- 1.5A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Numbers in Thousands, 2013 and 2014
- 1.5B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Percentages, 2013 and 2014
- 1.6A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Numbers in Thousands, 2013 and 2014
- 1.6B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Percentages, 2013 and 2014
- 1.7A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Numbers in Thousands, 2013 and 2014
- 1.7B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Percentages, 2013 and 2014
- 1.8A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Numbers in Thousands, 2013 and 2014
- 1.8B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Percentages, 2013 and 2014
- 1.9A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Numbers in Thousands, 2013 and 2014
- 1.9B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Percentages, 2013 and 2014
- 1.10A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Numbers in Thousands, 2013 and 2014
- 1.10B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Percentages, 2013 and 2014
- 1.11A Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
- 1.11B Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
- 1.12A Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
- 1.12B Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 1.13A Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
- 1.13B Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
- 1.14A Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
- 1.14B Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
- 1.15A Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
- 1.15B Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
- 1.16A Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
- 1.16B Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
- 1.17A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
- 1.17B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
- 1.18A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
- 1.18B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
- 1.19A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.19B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.20A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.20B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.21A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.21B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.22A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.22B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.23A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.23B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.24A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.24B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 1.25A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.25B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.26A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.26B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.27A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.27B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.28A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.28B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.29A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.29B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.30A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.30B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.31A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.31B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.32A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.32B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.33A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.33B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.34A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.34B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.35A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.35B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
- 1.36A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.36B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

1.37A	Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.37B	Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.38A	Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.38B	Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.39A	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.39B	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.40A	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.40B	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
1.41A	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.41B	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
1.42A	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.42B	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.43A	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.43B	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.44A	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.44B	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.45A	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.45B	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
1.46A	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.46B	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
1.47A	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.47B	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.48A	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.48B	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

1.49A	Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.49B	Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.50A	Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.50B	Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
1.51A	Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.51B	Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
1.52A	Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.52B	Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.53A	Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.53B	Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.54A	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.54B	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.55A	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.55B	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
1.56A	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.56B	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
1.57A	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.57B	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.58A	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
1.58B	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
1.59A	Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
1.59B	Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
1.60A	Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
1.60B	Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 1.61A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.61B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.62A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.62B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.63A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.63B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.64A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.64B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.65A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.65B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.66A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.66B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.67A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.67B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.68A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.68B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.69A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.69B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.70A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.70B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.71A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.71B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.72A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.72B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 1.73A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.73B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.74A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.74B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.75A Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.75B Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.76A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.76B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.77A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.77B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.78A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.78B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.79A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.79B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.80A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.80B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.81A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.81B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.82A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.82B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.83A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.83B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
- 1.84A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 1.84B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

1.85A	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
1.85B	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
1.86A	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
1.86B	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
1.87A	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
1.87B	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
1.88A	Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Numbers in Thousands, 2013 and 2014
1.88B	Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Percentages, 2013 and 2014
1.89A	Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Numbers in Thousands, 2013 and 2014
1.89B	Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Percentages, 2013 and 2014
1.90A	Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Numbers in Thousands, 2013 and 2014
1.90B	Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Percentages, 2013 and 2014
1.91A	Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Numbers in Thousands, 2013 and 2014
1.91B	Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Percentages, 2013 and 2014
1.92A	Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Numbers in Thousands, 2013 and 2014
1.92B	Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Percentages, 2013 and 2014
2.1A	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2013 and 2014
2.1B	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Percentages, 2013 and 2014
2.2A	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Numbers in Thousands, 2013 and 2014
2.2B	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Percentages, 2013 and 2014
2.3A	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Numbers in Thousands, 2013 and 2014
2.3B	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Percentages, 2013 and 2014
2.4A	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Numbers in Thousands, 2013 and 2014
2.4B	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Percentages, 2013 and 2014
2.5A	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Numbers in Thousands, 2013 and 2014
2.5B	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

2.6A	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Numbers in Thousands, 2013 and 2014
2.6B	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Percentages, 2013 and 2014
2.7A	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Numbers in Thousands, 2013 and 2014
2.7B	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Percentages, 2013 and 2014
2.8A	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Numbers in Thousands, 2013 and 2014
2.8B	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Percentages, 2013 and 2014
2.9A	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Numbers in Thousands, 2013 and 2014
2.9B	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Percentages, 2013 and 2014
2.10A	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Numbers in Thousands, 2013 and 2014
2.10B	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Percentages, 2013 and 2014
2.11A	Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
2.11B	Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
2.12A	Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
2.12B	Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
2.13A	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
2.13B	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
2.14A	Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
2.14B	Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
2.15A	Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
2.15B	Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014
2.16A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
2.16B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Percentages, 2013 and 2014
2.17A	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.17B	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 2.18A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.18B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
- 2.19A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.19B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
- 2.20A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.20B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 2.21A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.21B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 2.22A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.22B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 2.23A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.23B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
- 2.24A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.24B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
- 2.25A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.25B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 2.26A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.26B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 2.27A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.27B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 2.28A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.28B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
- 2.29A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 2.29B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

2.30A	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.30B	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.31A	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.31B	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.32A	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.32B	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.33A	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.33B	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
2.34A	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.34B	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
2.35A	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.35B	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.36A	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.36B	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.37A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.37B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.38A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.38B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
2.39A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.39B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
2.40A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.40B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.41A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.41B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

2.42A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.42B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.43A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.43B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
2.44A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.44B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
2.45A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.45B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.46A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.46B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.47A	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.47B	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.48A	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.48B	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014
2.49A	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.49B	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
2.50A	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.50B	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.51A	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.51B	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.52A	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.52B	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.53A	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.53B	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

2.54A	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.54B	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
2.55A	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.55B	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.56A	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.56B	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.57A	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.57B	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.58A	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.58B	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014
2.59A	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.59B	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
2.60A	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.60B	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.61A	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.61B	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.62A	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.62B	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.63A	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.63B	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014
2.64A	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.64B	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
2.65A	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.65B	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

2.66A	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.66B	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.67A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.67B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.68A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.68B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014
2.69A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.69B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
2.70A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.70B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.71A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.71B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.72A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.72B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.73A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.73B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014
2.74A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.74B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
2.75A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.75B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.76A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.76B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.77A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.77B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

2.78A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.78B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.79A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.79B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Percentages, 2013 and 2014
2.80A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
2.80B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
2.81A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.81B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Percentages, 2013 and 2014
2.82A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.82B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
2.83A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.83B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Percentages, 2013 and 2014
2.84A	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
2.84B	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
3.1A	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances, by Age Group: Numbers in Thousands, 2013 and 2014
3.1B	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances, by Age Group: Percentages, 2013 and 2014
3.2A	Perceived Great Risk of Harm Associated with Cigarette and Alcohol Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
3.2B	Perceived Great Risk of Harm Associated with Cigarette and Alcohol Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
3.3A	Perceived Great Risk of Harm Associated with Marijuana and Cocaine Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
3.3B	Perceived Great Risk of Harm Associated with Marijuana and Cocaine Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
3.4A	Perceived Great Risk of Harm Associated with Heroin and LSD Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
3.4B	Perceived Great Risk of Harm Associated with Heroin and LSD Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
3.5A	Perceived That Obtaining Specific Illicit Drugs Would Be Fairly or Very Easy among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
3.5B	Perceived That Obtaining Specific Illicit Drugs Would Be Fairly or Very Easy among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 3.6A Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.6B Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.7A Youths Felt That Close Friends Would Strongly Disapprove or Somewhat Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.7B Youths Felt That Close Friends Would Strongly Disapprove or Somewhat Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.8A Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.8B Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.9A Participated in a Prevention Program or Group in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.9B Participated in a Prevention Program or Group in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.10A Participated in a Substance Use Program or Exposure to Substance Use Prevention Message in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.10B Participated in a Substance Use Program or Exposure to Substance Use Prevention Message in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.11A Participated in a Physical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.11B Participated in a Physical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.12A Participated in a Nonphysical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.12B Participated in a Nonphysical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.13A Selected Risk and Protective Factors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.13B Selected Risk and Protective Factors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.14A Youths Agreed or Strongly Agreed with Statements about Religious Beliefs among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.14B Youths Agreed or Strongly Agreed with Statements about Religious Beliefs among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.15A Parents Always or Sometimes Took Part in Specific Behaviors in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.15B Parents Always or Sometimes Took Part in Specific Behaviors in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.16A Measures of Student Characteristics in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.16B Measures of Student Characteristics in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 3.17A Adults Always or Sometimes Took Part in Specific School-Related Behaviors in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.17B Adults Always or Sometimes Took Part in Specific School-Related Behaviors in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.18A Youths Perceived That All or Most Students in Grade Level Use Specific Substances among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 3.18B Youths Perceived That All or Most Students in Grade Level Use Specific Substances among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014
- 3.19A Past Month Use of Selected Substances, by Perceived Risk of Harm Associated with Substance Use Behaviors among Persons Aged 12 to 17: Numbers in Thousands, 2014
- 3.19B Past Month Use of Selected Substances, by Perceived Risk of Harm Associated with Substance Use Behaviors among Persons Aged 12 to 17: Percentages, 2014
- 3.20A Past Month Use of Selected Substances, by Perceived Availability of Selected Illicit Drugs among Persons Aged 12 to 17: Numbers in Thousands, 2014
- 3.20B Past Month Use of Selected Substances, by Perceived Availability of Selected Illicit Drugs among Persons Aged 12 to 17: Percentages, 2014
- 3.21A Past Month Use of Selected Substances, by Youth's Perceptions of Parents' Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Numbers in Thousands, 2014
- 3.21B Past Month Use of Selected Substances, by Youth's Perceptions of Parents' Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Percentages, 2014
- 3.22A Past Month Use of Selected Substances, by Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Numbers in Thousands, 2014
- 3.22B Past Month Use of Selected Substances, by Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Percentages, 2014
- 3.23A Past Month Use of Selected Substances, by Measures of Youth's Exposure to Prevention Programs and Messages and Youth's Delinquency in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2014
- 3.23B Past Month Use of Selected Substances, by Measures of Youth's Exposure to Prevention Programs and Messages and Youth's Delinquency in the Past Year among Persons Aged 12 to 17: Percentages, 2014
- 3.24A Past Month Use of Selected Substances, by Measures of Risk and Protective Factors, Religious Beliefs, and Past Year Parental Involvement among Persons Aged 12 to 17: Numbers in Thousands, 2014
- 3.24B Past Month Use of Selected Substances, by Measures of Risk and Protective Factors, Religious Beliefs, and Past Year Parental Involvement among Persons Aged 12 to 17: Percentages, 2014
- 3.25A Past Month Use of Selected Substances, by Measures of Scholastic Environment among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year: Numbers in Thousands, 2014
- 3.25B Past Month Use of Selected Substances, by Measures of Scholastic Environment among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year: Percentages, 2014
- 4.1B Past Year Initiation of Illicit Drug Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Illicit Drug Use, and Past Year Illicit Drug Users Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014
- 4.2B Past Year Initiation of Marijuana Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Marijuana Use, and Past Year Marijuana Users Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014
- 4.3B Past Year Initiation of Cigarette Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Cigarette Use, and Past Year Cigarette Users Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 4.4B Past Year Initiation of Alcohol Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Alcohol Use, and Past Year Alcohol Users Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014
- 4.5B Past Year Initiation of Substance Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 or Older: Numbers in Thousands and Percentages, 2013 and 2014
- 4.6B Past Year Initiation of Substance Use among Persons Aged 12 to 17, Persons Aged 12 to 17 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 to 17: Numbers in Thousands and Percentages, 2013 and 2014
- 4.7B Past Year Initiation of Substance Use among Persons Aged 18 to 25, Persons Aged 18 to 25 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 18 to 25: Numbers in Thousands and Percentages, 2013 and 2014
- 4.8B Past Year Initiation of Substance Use among Persons Aged 26 or Older, Persons Aged 26 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 26 or Older: Numbers in Thousands and Percentages, 2013 and 2014
- 4.9A Past Year Initiation of Substance Use among Persons Aged 12 or Older, by Gender: Numbers in Thousands, 2013 and 2014
- 4.10A Past Year Initiation of Substance Use among Persons Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: Numbers in Thousands, 2013 and 2014
- 4.11A Past Year Initiation of Substance Use among Persons Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: Numbers in Thousands, 2013 and 2014
- 4.12A Past Year Initiation of Substance Use among Persons Aged 12 to 49, by Gender: Numbers in Thousands, 2013 and 2014
- 4.13B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older, by Gender: 2013 and 2014
- 4.14B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: 2013 and 2014
- 4.15B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: 2013 and 2014
- 4.16B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49, by Gender: 2013 and 2014
- 5.1A Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2013 and 2014
- 5.1B Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older: Percentages, 2013 and 2014
- 5.2A Substance Dependence or Abuse for Specific Substances in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
- 5.2B Substance Dependence or Abuse for Specific Substances in the Past Year, by Age Group: Percentages, 2013 and 2014
- 5.3A Substance Dependence or Abuse in the Past Year, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
- 5.3B Substance Dependence or Abuse in the Past Year, by Detailed Age Category: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

5.4A	Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.4B	Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
5.5A	Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.5B	Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
5.6A	Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.6B	Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
5.7A	Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.7B	Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
5.8A	Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.8B	Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
5.9A	Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
5.9B	Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
5.10A	Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
5.10B	Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014
5.11A	Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
5.11B	Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014
5.12A	Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
5.12B	Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
5.13A	Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
5.13B	Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
5.14A	Substance Dependence for Specific Substances in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.14B	Substance Dependence for Specific Substances in the Past Year, by Age Group: Percentages, 2013 and 2014
5.15A	Received Substance Use Treatment in the Past Year, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
5.15B	Received Substance Use Treatment in the Past Year, by Detailed Age Category: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

5.16A	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.16B	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
5.17A	Received Substance Use Treatment in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.17B	Received Substance Use Treatment in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
5.18A	Received Substance Use Treatment in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.18B	Received Substance Use Treatment in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
5.19A	Received Substance Use Treatment in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.19B	Received Substance Use Treatment in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
5.20A	Received Substance Use Treatment in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.20B	Received Substance Use Treatment in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
5.21A	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
5.21B	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
5.22A	Substances for Which Last or Current Treatment Was Received among Persons Who Received Substance Use Treatment in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.22B	Substances for Which Last or Current Treatment Was Received among Persons Who Received Substance Use Treatment in the Past Year, by Age Group: Percentages, 2013 and 2014
5.23A	Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.23B	Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment in the Past Year, by Age Group: Percentages, 2013 and 2014
5.24A	Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.24B	Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: Percentages, 2013 and 2014
5.25A	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.25B	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: Percentages, 2013 and 2014
5.26A	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Numbers in Thousands, 2013 and 2014
5.26B	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Percentages, 2013 and 2014
5.27A	Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
5.27B	Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

5.28A	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.28B	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
5.29A	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.29B	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014
5.30A	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.30B	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014
5.31A	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.31B	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
5.32A	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.32B	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
5.33A	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
5.33B	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
5.34A	Substances for Which Last or Current Treatment Was Received among Persons Who Received Their Last or Current Substance Use Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.34B	Substances for Which Last or Current Treatment Was Received among Persons Who Received Their Last or Current Substance Use Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014
5.35A	Source of Payment for Last or Current Illicit Drug Treatment among Persons Who Received Their Last or Current Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.35B	Source of Payment for Last or Current Illicit Drug Treatment among Persons Who Received Their Last or Current Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014
5.36A	Source of Payment for Last or Current Alcohol Treatment among Persons Who Received Their Last or Current Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.36B	Source of Payment for Last or Current Alcohol Treatment among Persons Who Received Their Last or Current Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014
5.37A	Source of Payment for Last or Current Illicit Drug or Alcohol Treatment among Persons Who Received Their Last or Current Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.37B	Source of Payment for Last or Current Illicit Drug or Alcohol Treatment among Persons Who Received Their Last or Current Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014
5.38A	Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.38B	Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

5.39A	Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.39B	Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014
5.40A	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
5.40B	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014
5.41A	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Numbers in Thousands, 2013 and 2014
5.41B	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Percentages, 2013 and 2014
5.42A	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse of Specific Substances: Numbers in Thousands, 2013 and 2014
5.42B	Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse of Specific Substances: Percentages, 2013 and 2014
5.43A	Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year, by Detailed Age Category: Numbers in Thousands and Percentages, 2013 and 2014
5.43B	Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year, by Detailed Age Category: Percentages, 2013 and 2014
5.44A	Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014
5.44B	Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
5.45A	Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014
5.45B	Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
5.46A	Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014
5.46B	Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
5.47A	Perceived Need for Illicit Drug Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug Problem, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
5.47B	Perceived Need for Illicit Drug Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug Problem, by Demographic Characteristics: Percentages, 2013 and 2014
5.48A	Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014
5.48B	Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 5.49A Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014
- 5.49B Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 5.50A Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 5.50B Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Percentages, 2013 and 2014
- 5.51A Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014
- 5.51B Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 5.52A Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014
- 5.52B Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 5.53A Perceived Need for Illicit Drug or Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug or Alcohol Problem, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 5.53B Perceived Need for Illicit Drug or Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug or Alcohol Problem, by Demographic Characteristics: Percentages, 2013 and 2014
- 5.54A Detailed Reasons for Not Receiving Illicit Drug Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Numbers in Thousands, Annual Averages Based on 2011-2014
- 5.54B Detailed Reasons for Not Receiving Illicit Drug Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Percentages, Annual Averages Based on 2011-2014
- 5.55A Detailed Reasons for Not Receiving Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Alcohol Treatment at a Specialty Facility and Who Felt a Need for Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Numbers in Thousands, Annual Averages Based on 2011-2014
- 5.55B Detailed Reasons for Not Receiving Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Alcohol Treatment at a Specialty Facility and Who Felt a Need for Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Percentages, Annual Averages Based on 2011-2014
- 5.56A Detailed Reasons for Not Receiving Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug or Alcohol Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug or Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Numbers in Thousands, Annual Averages Based on 2011-2014

LIST OF TABLES (Continued)

Table

5.56B	Detailed Reasons for Not Receiving Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug or Alcohol Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug or Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Percentages, Annual Averages Based on 2011-2014
6.1A	Number of Days Used Marijuana in the Past Year among Past Year Users and the Number of Days Used Marijuana in the Past Month among Past Month Users, by Age Group: Numbers in Thousands, 2013 and 2014
6.1B	Number of Days Used Marijuana in the Past Year among Past Year Users and the Number of Days Used Marijuana in the Past Month among Past Month Users, by Age Group: Percentages, 2013 and 2014
6.2A	Number of Days Used Cocaine in the Past Year among Past Year Users and the Number of Days Used Cocaine in the Past Month among Past Month Users, by Age Group: Numbers in Thousands, 2013 and 2014
6.2B	Number of Days Used Cocaine in the Past Year among Past Year Users and the Number of Days Used Cocaine in the Past Month among Past Month Users, by Age Group: Percentages, 2013 and 2014
6.3A	Number of Days Used Hallucinogens in the Past Year among Past Year Users and the Number Days Used Hallucinogens in the Past Month among Past Month Users, by Age Group: Numbers in Thousands, 2013 and 2014
6.3B	Number of Days Used Hallucinogens in the Past Year among Past Year Users and the Number Days Used Hallucinogens in the Past Month among Past Month Users, by Age Group: Percentages, 2013 and 2014
6.4A	Number of Days Used Inhalants in the Past Year among Past Year Users and the Number of Days Used Inhalants in the Past Month among Past Month Users, by Age Group: Numbers in Thousands, 2013 and 2014
6.4B	Number of Days Used Inhalants in the Past Year among Past Year Users and the Number of Days Used Inhalants in the Past Month among Past Month Users, by Age Group: Percentages, 2013 and 2014
6.5A	Number of Days Used Cigarettes in the Past Month among Past Month Users and the Number of Cigarettes Smoked Per Day among Daily Smokers, by Age Group: Numbers in Thousands, 2013 and 2014
6.5B	Number of Days Used Cigarettes in the Past Month among Past Month Users and the Number of Cigarettes Smoked Per Day among Daily Smokers, by Age Group: Percentages, 2013 and 2014
6.6A	Number of Days Used Alcohol in the Past Year among Past Year Users and the Number of Days Used Alcohol in the Past Month among Past Month Users, by Age Group: Numbers in Thousands, 2013 and 2014
6.6B	Number of Days Used Alcohol in the Past Year among Past Year Users and the Number of Days Used Alcohol in the Past Month among Past Month Users, by Age Group: Percentages, 2013 and 2014
6.7A	Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014
6.7B	Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2013 and 2014
6.8A	Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014
6.8B	Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2013 and 2014
6.9A	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014
6.9B	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2013 and 2014
6.10A	Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014
6.10B	Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 6.11A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.11B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014
- 6.12A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.12B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014
- 6.13A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.13B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014
- 6.14A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.14B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014
- 6.15A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Numbers in Thousands, 2013 and 2014
- 6.15B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Percentages, 2013 and 2014
- 6.16A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Numbers in Thousands, 2013 and 2014
- 6.16B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Percentages, 2013 and 2014
- 6.17A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2013 and 2014
- 6.17B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2013 and 2014
- 6.18A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2013 and 2014
- 6.18B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2013 and 2014
- 6.19A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.19B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Percentages, 2013 and 2014
- 6.20A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.20B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Percentages, 2013 and 2014
- 6.21A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.21B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 6.22A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.22B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Percentages, 2013 and 2014
- 6.23A Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Numbers in Thousands, 2013 and 2014
- 6.23B Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Percentages, 2013 and 2014
- 6.24A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014
- 6.24B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2013 and 2014
- 6.25A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014
- 6.25B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2013 and 2014
- 6.26A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014
- 6.26B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2013 and 2014
- 6.27A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014
- 6.27B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2013 and 2014
- 6.28A Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.28B Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014
- 6.29A Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.29B Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014
- 6.30A Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.30B Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014
- 6.31A Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014
- 6.31B Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014
- 6.32A Marijuana Use in the Prior Year and Past Year Marijuana Use Status among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 6.32B Marijuana Use in the Prior Year and Past Year Marijuana Use Status among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
- 6.33A Cigarette Use and Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Detailed Age Category: Numbers in Thousands and Percentages, 2013 and 2014
- 6.33B Cigarette Use and Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Detailed Age Category: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 6.34A Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 6.34B Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: Percentages, 2013 and 2014
- 6.35A Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Geographic Characteristics: Numbers in Thousands, 2013 and 2014
- 6.35B Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Geographic Characteristics: Percentages, 2013 and 2014
- 6.36A Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 6.36B Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes and Demographic Characteristics: Percentages, 2013 and 2014
- 6.37A Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes Daily and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 6.37B Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes Daily and Demographic Characteristics: Percentages, 2013 and 2014
- 6.38A Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 6.38B Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Percentages, 2013 and 2014
- 6.39A Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 6.39B Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Percentages, 2013 and 2014
- 6.40A Alcohol Dependence or Abuse in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Numbers in Thousands, 2013 and 2014
- 6.40B Alcohol Dependence or Abuse in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Percentages, 2013 and 2014
- 6.41A Illicit Drug Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 6.41B Illicit Drug Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Percentages, 2013 and 2014
- 6.42A Alcohol Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 6.42B Alcohol Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Percentages, 2013 and 2014
- 6.43A Alcohol Dependence in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Numbers in Thousands, 2013 and 2014
- 6.43B Alcohol Dependence in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Percentages, 2013 and 2014
- 6.44A Method of Obtaining Most Recently Used Marijuana among Past Year Marijuana Users, by Age Group: Numbers in Thousands, 2013 and 2014
- 6.44B Method of Obtaining Most Recently Used Marijuana among Past Year Marijuana Users, by Age Group: Percentages, 2013 and 2014
- 6.45A Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Bought Their Most Recently Used Marijuana, by Age Group: Numbers in Thousands, 2013 and 2014
- 6.45B Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Bought Their Most Recently Used Marijuana, by Age Group: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 6.46A Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Obtained for Free Their Most Recently Used Marijuana and Who Had Not Bought or Traded for Marijuana in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014
- 6.46B Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Obtained for Free Their Most Recently Used Marijuana and Who Had Not Bought or Traded for Marijuana in the Past Year, by Age Group: Percentages, 2013 and 2014
- 6.47A Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.47B Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.48A Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.48B Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.49A Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.49B Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.50A Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.50B Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.51A Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.51B Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.52A Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.52B Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.53A Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.53B Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.54A Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.54B Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

LIST OF TABLES (Continued)

Table

- 6.55A Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.55B Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.56A Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.56B Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.57A Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.57B Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.58A Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.58B Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.59A Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.59B Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.60A Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.60B Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.61A Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.61B Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.62A Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.62B Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.63A Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.63B Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

LIST OF TABLES (Continued)

Table

- 6.64A Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.64B Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.65A Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.65B Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.66A Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.66B Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.67A Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Numbers in Thousands, 2013 and 2014
- 6.67B Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Percentages, 2013 and 2014
- 6.68A Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Numbers in Thousands, 2013 and 2014
- 6.68B Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Percentages, 2013 and 2014
- 6.69A Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Numbers in Thousands, 2013 and 2014
- 6.69B Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Percentages, 2013 and 2014
- 6.70A Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Numbers in Thousands, 2013 and 2014
- 6.70B Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Percentages, 2013 and 2014
- 6.71A Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.71B Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.72A Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.72B Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.73A Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.73B Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.74A Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.74B Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

LIST OF TABLES (Continued)

Table

- 6.75A Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.75B Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.76A Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.76B Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.77A Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.77B Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.78A Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.78B Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.79A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014
- 6.79B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014
- 6.80A Types of Illicit Drug Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014
- 6.80B Types of Illicit Drug Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014
- 6.81A Types of Illicit Drug Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014
- 6.81B Types of Illicit Drug Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014
- 6.82A Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 6.82B Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014
- 6.83A Marijuana Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
- 6.83B Marijuana Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014
- 6.84A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014
- 6.84B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014
- 6.85A Tobacco Product and Alcohol Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014
- 6.85B Tobacco Product and Alcohol Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014
- 6.86A Tobacco Product and Alcohol Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014
- 6.86B Tobacco Product and Alcohol Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

6.87A	Cigarette Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.87B	Cigarette Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014
6.88A	Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.88B	Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014
6.89A	Binge Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.89B	Binge Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014
6.90A	Heavy Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.90B	Heavy Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014
6.91A	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.91B	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014
6.92A	Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.92B	Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014
6.93A	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.93B	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014
6.94A	Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year, by Detailed Age Category: Numbers in Thousands, 2013 and 2014
6.94B	Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year, by Detailed Age Category: Percentages, 2013 and 2014
6.95A	Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.95B	Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
6.96A	Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.96B	Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014
6.97A	Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014
6.97B	Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014
6.98A	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status: Numbers in Thousands, 2013 and 2014
6.98B	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status: Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

6.99A	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Probation Status: Numbers in Thousands, 2013 and 2014
6.99B	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Probation Status: Percentages, 2013 and 2014
6.100A	Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.100B	Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Percentages, 2013 and 2014
6.101A	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.101B	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Percentages, 2013 and 2014
6.102A	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.102B	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Percentages, 2013 and 2014
6.103A	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status: Numbers in Thousands, 2013 and 2014
6.103B	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status: Percentages, 2013 and 2014
6.104A	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Parole/Supervised Release Status: Numbers in Thousands, 2013 and 2014
6.104B	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Parole/Supervised Release Status: Percentages, 2013 and 2014
6.105A	Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.105B	Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Percentages, 2013 and 2014
6.106A	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.106B	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Percentages, 2013 and 2014
6.107A	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014
6.107B	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Percentages, 2013 and 2014
7.1A	Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014
7.1B	Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Percentages, 2002-2014
7.2A	Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014
7.2B	Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2014
7.3A	Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014
7.3B	Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Percentages, 2002-2014
7.4A	Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014
7.4B	Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Percentages, 2002-2014

LIST OF TABLES (Continued)

Table

7.5A	Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014
7.5B	Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2014
7.6A	Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014
7.6B	Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Percentages, 2002-2014
7.7A	Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014
7.7B	Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Percentages, 2002-2014
7.8A	Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014
7.8B	Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2014
7.9A	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014
7.9B	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Percentages, 2002-2014
7.10A	Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014
7.10B	Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Percentages, 2002-2014
7.11A	Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014
7.11B	Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2014
7.12A	Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014
7.12B	Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Percentages, 2002-2014
7.13A	Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Numbers in Thousands, 2002-2014
7.13B	Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Percentages, 2002-2014
7.14A	Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Numbers in Thousands, 2002-2014
7.14B	Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Percentages, 2002-2014
7.15A	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Numbers in Thousands, 2002-2014
7.15B	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Percentages, 2002-2014
7.16A	Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Numbers in Thousands, 2002-2014
7.16B	Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Percentages, 2002-2014
7.17A	Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Numbers in Thousands, 2002-2014
7.17B	Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Percentages, 2002-2014
7.18A	Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Numbers in Thousands, 2002-2014
7.18B	Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Percentages, 2002-2014
7.19A	Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Numbers in Thousands, 2002-2014
7.19B	Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Percentages, 2002-2014
7.20A	Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Numbers in Thousands, 2002-2014
7.20B	Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Percentages, 2002-2014

LIST OF TABLES (Continued)

Table

7.21A	Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Numbers in Thousands, 2002-2014
7.21B	Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Percentages, 2002-2014
7.22A	GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014
7.22B	GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014
7.23A	Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014
7.23B	Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014
7.24A	Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014
7.24B	Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014
7.25A	Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014
7.25B	Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014
7.26A	Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014
7.26B	Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014
7.27A	DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014
7.27B	DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014
7.28A	Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014
7.28B	Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014
7.29A	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Numbers in Thousands, 2002-2014
7.29B	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Percentages, 2002-2014
7.30A	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Numbers in Thousands, 2002-2014
7.30B	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Percentages, 2002-2014
7.31A	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Numbers in Thousands, 2002-2014
7.31B	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Percentages, 2002-2014
7.32A	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Numbers in Thousands, 2002-2014
7.32B	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Percentages, 2002-2014

LIST OF TABLES (Continued)

Table

7.33A	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Numbers in Thousands, 2002-2014
7.33B	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Percentages, 2002-2014
7.34A	Daily Cigarette Use in the Past Month among Past Month Users Aged 12 or Older and Smoked One or More Packs of Cigarettes Per Day among Daily Cigarette Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2002-2014
7.34B	Daily Cigarette Use in the Past Month among Past Month Users Aged 12 or Older and Smoked One or More Packs of Cigarettes Per Day among Daily Cigarette Users Aged 12 or Older, by Age Group: Percentages, 2002-2014
7.35A	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Gender: Numbers in Thousands, 2002-2014
7.35B	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Gender: Percentages, 2002-2014
7.36A	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014
7.36B	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 or Older: Percentages, 2002-2014
7.37A	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014
7.37B	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 to 17: Percentages, 2002-2014
7.38A	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014
7.38B	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 18 to 25: Percentages, 2002-2014
7.39A	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014
7.39B	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 26 or Older: Percentages, 2002-2014
7.40A	Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Youths Aged 12 to 17: Numbers in Thousands, 2002-2014
7.40B	Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Youths Aged 12 to 17: Percentages, 2002-2014
7.41A	Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Youths Aged 12 to 17: Numbers in Thousands, 2002-2014
7.41B	Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Youths Aged 12 to 17: Percentages, 2002-2014
7.42A	Exposure to Substance Use Prevention Program or Message in the Past Year among Youths Aged 12 to 17: Numbers in Thousands, 2002-2014
7.42B	Exposure to Substance Use Prevention Program or Message in the Past Year among Youths Aged 12 to 17: Percentages, 2002-2014
7.43A	Substance Use in the Past Month among Youths Aged 12 to 17, by Exposure to Substance Use Prevention Program or Message in the Past Year: Numbers in Thousands, 2002-2014
7.43B	Substance Use in the Past Month among Youths Aged 12 to 17, by Exposure to Substance Use Prevention Program or Message in the Past Year: Percentages, 2002-2014
7.44A	Past Year Initiation of Substance Use among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014
7.44B	Past Year Initiation of Substance Use among Persons Aged 12 or Older: Percentages, 2002-2014

LIST OF TABLES (Continued)

Table

7.45A	Past Year Initiation of Substance Use among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014
7.45B	Past Year Initiation of Substance Use among Persons Aged 12 to 17: Percentages, 2002-2014
7.46A	Past Year Initiation of Substance Use among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014
7.46B	Past Year Initiation of Substance Use among Persons Aged 18 to 25: Percentages, 2002-2014
7.47A	Past Year Initiation of Substance Use among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014
7.47B	Past Year Initiation of Substance Use among Persons Aged 26 or Older: Percentages, 2002-2014
7.48A	Past Year Initiation of Illicit Drugs and First Drug Initiated in the Past Year among Persons Aged 12 or Older Who Initiated Illicit Drug Use in the Past Year: Numbers in Thousands, 2002-2014
7.48B	Past Year Initiation of Illicit Drugs and First Drug Initiated in the Past Year among Persons Aged 12 or Older Who Initiated Illicit Drug Use in the Past Year: Percentages, 2002-2014
7.49B	Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49: 2002-2014
7.50A	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014
7.50B	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2014
7.51A	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014
7.51B	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2014
7.52A	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014
7.52B	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2014
7.53A	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014
7.53B	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2014
7.54A	Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014
7.54B	Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2014
7.55A	Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014
7.55B	Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2014
7.56A	Substance Dependence for Specific Substances in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014
7.56B	Substance Dependence for Specific Substances in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2014
7.57A	Substance Dependence for Specific Substances in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014
7.57B	Substance Dependence for Specific Substances in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2014
7.58A	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014
7.58B	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2014

LIST OF TABLES (Continued)

Table

7.59A	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014
7.59B	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2014
7.60A	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014
7.60B	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2014
7.61A	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014
7.61B	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2014
7.62A	Substances for Which Last or Current Treatment Was Received among Persons Aged 12 or Older Who Received Substance Use Treatment in the Past Year: Numbers in Thousands, 2002-2014
7.62B	Substances for Which Last or Current Treatment Was Received among Persons Aged 12 or Older Who Received Substance Use Treatment in the Past Year: Percentages, 2002-2014
7.63A	Substances for Which Last or Current Treatment Was Received among Persons Aged 12 to 17 Who Received Substance Use Treatment in the Past Year: Numbers in Thousands, 2002-2014
7.63B	Substances for Which Last or Current Treatment Was Received among Persons Aged 12 to 17 Who Received Substance Use Treatment in the Past Year: Percentages, 2002-2014
7.64A	Substances for Which Last or Current Treatment Was Received among Persons Aged 18 to 25 Who Received Substance Use Treatment in the Past Year: Numbers in Thousands, 2002-2014
7.64B	Substances for Which Last or Current Treatment Was Received among Persons Aged 18 to 25 Who Received Substance Use Treatment in the Past Year: Percentages, 2002-2014
7.65A	Substances for Which Last or Current Treatment Was Received among Persons Aged 26 or Older Who Received Substance Use Treatment in the Past Year: Numbers in Thousands, 2002-2014
7.65B	Substances for Which Last or Current Treatment Was Received among Persons Aged 26 or Older Who Received Substance Use Treatment in the Past Year: Percentages, 2002-2014
7.66A	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Numbers in Thousands, 2002-2014
7.66B	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Percentages, 2002-2014
7.67A	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 to 17 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Numbers in Thousands, 2002-2014
7.67B	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 to 17 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Percentages, 2002-2014
7.68A	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 18 to 25 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Numbers in Thousands, 2002-2014
7.68B	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 18 to 25 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Percentages, 2002-2014
7.69A	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 26 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Numbers in Thousands, 2002-2014
7.69B	Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 26 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Percentages, 2002-2014
7.70A	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014
7.70B	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2014

LIST OF TABLES (Continued)

Table

7.71A	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014
7.71B	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2014
7.72A	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014
7.72B	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2014
7.73A	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014
7.73B	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2014
7.74A	Type of Substance Treatment Received at Any Location in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use in the Past Year, by Age Group: Numbers in Thousands, 2002-2014
7.74B	Type of Substance Treatment Received at Any Location in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use in the Past Year, by Age Group: Percentages, 2002-2014
7.75A	Type of Substance Treatment Received at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2002-2014
7.75B	Type of Substance Treatment Received at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use at a Specialty Facility in the Past Year, by Age Group: Percentages, 2002-2014
7.76A	Receipt of Illicit Drug Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug Treatment, by Age Group: Numbers in Thousands, 2002-2014
7.76B	Receipt of Illicit Drug Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug Treatment, by Age Group: Percentages, 2002-2014
7.77A	Receipt of Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Alcohol Treatment, by Age Group: Numbers in Thousands, 2002-2014
7.77B	Receipt of Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Alcohol Treatment, by Age Group: Percentages, 2002-2014
7.78A	Receipt of Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment, by Age Group: Numbers in Thousands, 2002-2014
7.78B	Receipt of Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment, by Age Group: Percentages, 2002-2014
7.79B	Marijuana Use in the Past Month, by Age Group: Percentages, 1971-2014
7.80B	Cocaine Use in the Past Year, by Age Group: Percentages, 1971-2014
8.1A	Numbers (in Thousands) of Persons Aged 12 or Older, by Gender and Detailed Age Category: 2013 and 2014
8.1N	Survey Sample Size for Respondents Aged 12 or Older, by Gender and Detailed Age Category: 2013 and 2014
8.2A	Numbers (in Thousands) of Persons Aged 12 or Older, by Age Group and Demographic Characteristics: 2013 and 2014
8.2N	Survey Sample Size for Respondents Aged 12 or Older, by Age Group and Demographic Characteristics: 2013 and 2014

LIST OF TABLES (Continued)

Table

8.3A	Numbers (in Thousands) of Persons Aged 18 or Older, by Age Group and Demographic Characteristics: 2013 and 2014
8.3N	Survey Sample Size for Respondents Aged 18 or Older, by Age Group and Demographic Characteristics: 2013 and 2014
8.4A	Numbers (in Thousands) of Persons Aged 12 or Older, by Youth and Adult Age Groups and Demographic Characteristics: 2013 and 2014
8.4N	Survey Sample Size for Respondents Aged 12 or Older, by Youth and Adult Age Groups and Demographic Characteristics: 2013 and 2014
8.5A	Numbers (in Thousands) of Persons Aged 12 or Older, by Underage and Legal Drinking Age Groups and Demographic Characteristics: 2013 and 2014
8.5N	Survey Sample Size for Respondents Aged 12 or Older, by Underage and Legal Drinking Age Groups and Demographic Characteristics: 2013 and 2014
8.6A	Numbers (in Thousands) of Persons Aged 12 or Older, by Age Group and Geographic Characteristics: 2013 and 2014
8.6N	Survey Sample Size for Respondents Aged 12 or Older, by Age Group and Geographic Characteristics: 2013 and 2014
8.7A	Numbers (in Thousands) of Persons Aged 18 or Older, by Age Group and Geographic Characteristics: 2013 and 2014
8.7N	Survey Sample Size for Respondents Aged 18 or Older, by Age Group and Geographic Characteristics: 2013 and 2014
8.8A	Numbers (in Thousands) of Persons Aged 12 or Older, by Underage and Legal Drinking Age Groups and Geographic Characteristics: 2013 and 2014
8.8N	Survey Sample Size for Respondents Aged 12 or Older, by Underage and Legal Drinking Age Groups and Geographic Characteristics: 2013 and 2014
8.9A	Numbers (in Thousands) of Persons Aged 12 or Older, by Youth and Adult Age Groups and Gender: 2002-2014
8.9N	Survey Sample Size for Respondents Aged 12 or Older, by Youth and Adult Age Groups and Gender: 2002-2014
8.10A	Numbers (in Thousands) of Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Annual Averages Based on 2011-2012 and 2013-2014
8.10N	Survey Sample Size for Female Respondents Aged 15 to 44, by Pregnancy and Demographic Characteristics: 2011-2012 and 2013-2014
8.11A	Numbers (in Thousands) of Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: 2013 and 2014
8.11N	Survey Sample Size for Respondents Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: 2013 and 2014
8.12A	Numbers (in Thousands) of Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: 2013 and 2014
8.12N	Survey Sample Size for Respondents Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: 2013 and 2014
8.13A	Numbers (in Thousands) of Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: 2013 and 2014
8.13N	Survey Sample Size for Respondents Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: 2013 and 2014

LIST OF TABLES

Table

- 1.1C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.1D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014
- 1.1P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.2C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.2D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2013 and 2014
- 1.2P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.3C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.3D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2013 and 2014
- 1.3P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.4C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.4D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2013 and 2014
- 1.4P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.5C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.5D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2013 and 2014
- 1.5P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.6C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.6D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2013 and 2014
- 1.6P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.7C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.7D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2013 and 2014
- 1.7P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.8C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.8D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2013 and 2014
- 1.8P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 1.9C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.9D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014
- 1.9P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.10C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.10D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014
- 1.10P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.11C Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.11D Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 1.11P Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.12C Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.12D Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 1.12P Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.13C Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.13D Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 1.13P Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.14C Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.14D Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 1.14P Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.15C Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.15D Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 1.15P Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.16C Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.16D Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 1.16P Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.17C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.17D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 1.17P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 1.18C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.18D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 1.18P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.19C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.19D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.19P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.20C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.20D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.20P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.21C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.21D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.21P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.22C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.22D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.22P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.23C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.23D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.23P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.24C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.24D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.24P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.25C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.25D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.25P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.26C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.26D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.26P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 1.27C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.27D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.27P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.28C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.28D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.28P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.29C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.29D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.29P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.30C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.30D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.30P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.31C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.31D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.31P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.32C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.32D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.32P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.33C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.33D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.33P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.34C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.34D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.34P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.35C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.35D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.35P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 1.36C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.36D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.36P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.37C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.37D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.37P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.38C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.38D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.38P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.39C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.39D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.39P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.40C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.40D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.40P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.41C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.41D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.41P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.42C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.42D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.42P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.43C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.43D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.43P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.44C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.44D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.44P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 1.45C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.45D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.45P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.46C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.46D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.46P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.47C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.47D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.47P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.48C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.48D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.48P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.49C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.49D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.49P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.50C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.50D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.50P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.51C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.51D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.51P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.52C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.52D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 1.52P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.53C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.53D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.53P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.54C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.54D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.54P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.55C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.55D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.55P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.56C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.56D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.56P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.57C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.57D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.57P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.58C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.58D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.58P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.59C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.59D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.59P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.60C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 1.60D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.60P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.61C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.61D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.61P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.62C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.62D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.62P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.63C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.63D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.63P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.64C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.64D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.64P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.65C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.65D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.65P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.66C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.66D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.66P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.67C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.67D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.67P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.68C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.68D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.68P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

1.69C	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.69D	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.69P	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.70C	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.70D	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.70P	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.71C	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.71D	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.71P	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.72C	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.72D	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.72P	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.73C	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.73D	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.73P	Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.74C	Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.74D	Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.74P	Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.75C	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.75D	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.75P	Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.76C	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.76D	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.76P	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.77C	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.77D	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.77P	Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 1.78C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.78D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.78P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.79C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.79D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.79P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.80C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.80D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.80P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.81C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.81D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.81P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.82C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.82D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.82P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.83C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.83D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.83P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 1.84C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 1.84D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 1.84P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

1.85C	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.85D	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.85P	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.86C	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.86D	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.86P	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.87C	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
1.87D	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
1.87P	Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.88C	Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
1.88D	Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Percentages, 2013 and 2014
1.88P	Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.89C	Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
1.89D	Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Percentages, 2013 and 2014
1.89P	Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.90C	Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
1.90D	Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Percentages, 2013 and 2014
1.90P	Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.91C	Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
1.91D	Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Percentages, 2013 and 2014
1.91P	Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
1.92C	Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
1.92D	Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Percentages, 2013 and 2014
1.92P	Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.1C	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014
2.1D	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 2.1P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.2C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.2D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2013 and 2014
- 2.2P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.3C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.3D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2013 and 2014
- 2.3P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.4C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.4D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2013 and 2014
- 2.4P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.5C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.5D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2013 and 2014
- 2.5P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.6C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.6D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2013 and 2014
- 2.6P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.7C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.7D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2013 and 2014
- 2.7P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.8C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.8D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2013 and 2014
- 2.8P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.9C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.9D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014
- 2.9P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.10C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014

LIST OF TABLES (Continued)

Table

2.10D	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014
2.10P	Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.11C	Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
2.11D	Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
2.11P	Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.12C	Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
2.12D	Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
2.12P	Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.13C	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
2.13D	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
2.13P	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.14C	Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
2.14D	Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
2.14P	Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.15C	Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
2.15D	Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
2.15P	Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.16C	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
2.16D	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
2.16P	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.17C	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.17D	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.17P	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.18C	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.18D	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.18P	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

2.19C	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.19D	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.19P	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.20C	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.20D	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.20P	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.21C	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.21D	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.21P	Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.22C	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.22D	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.22P	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.23C	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.23D	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.23P	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.24C	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.24D	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.24P	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.25C	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.25D	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.25P	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.26C	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.26D	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.26P	Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.27C	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.27D	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.27P	Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 2.28C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.28D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.28P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.29C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.29D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.29P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.30C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.30D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.30P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.31C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.31D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.31P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.32C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.32D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.32P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.33C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.33D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.33P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.34C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.34D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.34P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.35C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.35D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.35P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.36C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.36D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

2.36P	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.37C	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.37D	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.37P	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.38C	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.38D	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.38P	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.39C	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.39D	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.39P	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.40C	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.40D	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.40P	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.41C	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.41D	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.41P	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.42C	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.42D	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.42P	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.43C	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.43D	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.43P	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.44C	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.44D	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.44P	Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 2.45C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.45D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.45P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.46C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.46D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.46P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.47C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.47D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.47P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.48C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.48D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.48P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.49C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.49D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.49P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.50C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.50D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.50P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.51C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.51D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.51P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.52C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.52D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.52P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.53C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.53D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

2.62P	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.63C	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.63D	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.63P	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.64C	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.64D	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.64P	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.65C	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.65D	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.65P	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.66C	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.66D	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.66P	Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.67C	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.67D	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.67P	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.68C	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.68D	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.68P	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.69C	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.69D	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.69P	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.70C	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.70D	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
2.70P	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
2.71C	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
2.71D	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 2.71P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.72C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.72D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.72P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.73C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.73D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.73P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.74C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.74D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.74P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.75C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.75D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.75P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.76C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.76D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.76P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.77C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.77D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.77P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.78C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.78D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.78P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.79C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.79D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.79P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 2.80C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.80D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.80P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.81C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.81D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.81P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.82C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.82D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.82P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.83C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.83D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.83P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 2.84C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 2.84D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 2.84P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.1C Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.1D Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 3.1P Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.2C Perceived Great Risk of Harm Associated with Cigarette and Alcohol Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.2D Perceived Great Risk of Harm Associated with Cigarette and Alcohol Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.2P Perceived Great Risk of Harm Associated with Cigarette and Alcohol Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.3C Perceived Great Risk of Harm Associated with Marijuana and Cocaine Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.3D Perceived Great Risk of Harm Associated with Marijuana and Cocaine Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.3P Perceived Great Risk of Harm Associated with Marijuana and Cocaine Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 3.4C Perceived Great Risk of Harm Associated with Heroin and LSD Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.4D Perceived Great Risk of Harm Associated with Heroin and LSD Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.4P Perceived Great Risk of Harm Associated with Heroin and LSD Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.5C Perceived That Obtaining Specific Illicit Drugs Would Be Fairly or Very Easy among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.5D Perceived That Obtaining Specific Illicit Drugs Would Be Fairly or Very Easy among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.5P Perceived That Obtaining Specific Illicit Drugs Would Be Fairly or Very Easy among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.6C Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.6D Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.6P Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.7C Youths Felt That Close Friends Would Strongly Disapprove or Somewhat Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.7D Youths Felt That Close Friends Would Strongly Disapprove or Somewhat Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.7P Youths Felt That Close Friends Would Strongly Disapprove or Somewhat Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.8C Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.8D Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.8P Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.9C Participated in a Prevention Program or Group in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.9D Participated in a Prevention Program or Group in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.9P Participated in a Prevention Program or Group in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.10C Participated in a Substance Use Program or Exposure to Substance Use Prevention Message in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.10D Participated in a Substance Use Program or Exposure to Substance Use Prevention Message in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 3.10P Participated in a Substance Use Program or Exposure to Substance Use Prevention Message in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.11C Participated in a Physical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.11D Participated in a Physical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.11P Participated in a Physical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.12C Participated in a Nonphysical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.12D Participated in a Nonphysical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.12P Participated in a Nonphysical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.13C Selected Risk and Protective Factors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.13D Selected Risk and Protective Factors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.13P Selected Risk and Protective Factors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.14C Youths Agreed or Strongly Agreed with Statements about Religious Beliefs among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.14D Youths Agreed or Strongly Agreed with Statements about Religious Beliefs among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.14P Youths Agreed or Strongly Agreed with Statements about Religious Beliefs among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.15C Parents Always or Sometimes Took Part in Specific Behaviors in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.15D Parents Always or Sometimes Took Part in Specific Behaviors in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.15P Parents Always or Sometimes Took Part in Specific Behaviors in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.16C Measures of Student Characteristics in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.16D Measures of Student Characteristics in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.16P Measures of Student Characteristics in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.17C Adults Always or Sometimes Took Part in Specific School-Related Behaviors in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.17D Adults Always or Sometimes Took Part in Specific School-Related Behaviors in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 3.17P Adults Always or Sometimes Took Part in Specific School-Related Behaviors in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.18C Youths Perceived That All or Most Students in Grade Level Use Specific Substances among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 3.18D Youths Perceived That All or Most Students in Grade Level Use Specific Substances among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 3.18P Youths Perceived That All or Most Students in Grade Level Use Specific Substances among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 3.19C Past Month Use of Selected Substances, by Perceived Risk of Harm Associated with Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014
- 3.19D Past Month Use of Selected Substances, by Perceived Risk of Harm Associated with Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Percentages, 2014
- 3.20C Past Month Use of Selected Substances, by Perceived Availability of Selected Illicit Drugs among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014
- 3.20D Past Month Use of Selected Substances, by Perceived Availability of Selected Illicit Drugs among Persons Aged 12 to 17: Standard Errors of Percentages, 2014
- 3.21C Past Month Use of Selected Substances, by Youth's Perceptions of Parents' Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014
- 3.21D Past Month Use of Selected Substances, by Youth's Perceptions of Parents' Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Percentages, 2014
- 3.22C Past Month Use of Selected Substances, by Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014
- 3.22D Past Month Use of Selected Substances, by Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Percentages, 2014
- 3.23C Past Month Use of Selected Substances, by Measures of Youth's Exposure to Prevention Programs and Messages and Youth's Delinquency in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014
- 3.23D Past Month Use of Selected Substances, by Measures of Youth's Exposure to Prevention Programs and Messages and Youth's Delinquency in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2014
- 3.24C Past Month Use of Selected Substances, by Measures of Risk and Protective Factors, Religious Beliefs, and Past Year Parental Involvement among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014
- 3.24D Past Month Use of Selected Substances, by Measures of Risk and Protective Factors, Religious Beliefs, and Past Year Parental Involvement among Persons Aged 12 to 17: Standard Errors of Percentages, 2014
- 3.25C Past Month Use of Selected Substances, by Measures of Scholastic Environment among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year: Standard Errors of Numbers in Thousands, 2014
- 3.25D Past Month Use of Selected Substances, by Measures of Scholastic Environment among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year: Standard Errors of Percentages, 2014
- 4.1D Past Year Initiation of Illicit Drug Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Illicit Drug Use, and Past Year Illicit Drug Users Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 4.1P Past Year Initiation of Illicit Drug Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Illicit Drug Use, and Past Year Illicit Drug Users Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 4.2D Past Year Initiation of Marijuana Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Marijuana Use, and Past Year Marijuana Users Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 4.2P Past Year Initiation of Marijuana Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Marijuana Use, and Past Year Marijuana Users Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 4.3D Past Year Initiation of Cigarette Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Cigarette Use, and Past Year Cigarette Users Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 4.3P Past Year Initiation of Cigarette Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Cigarette Use, and Past Year Cigarette Users Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 4.4D Past Year Initiation of Alcohol Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Alcohol Use, and Past Year Alcohol Users Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 4.4P Past Year Initiation of Alcohol Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Alcohol Use, and Past Year Alcohol Users Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 4.5D Past Year Initiation of Substance Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 or Older: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 4.5P Past Year Initiation of Substance Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 4.6D Past Year Initiation of Substance Use among Persons Aged 12 to 17, Persons Aged 12 to 17 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 to 17: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 4.6P Past Year Initiation of Substance Use among Persons Aged 12 to 17, Persons Aged 12 to 17 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 4.7D Past Year Initiation of Substance Use among Persons Aged 18 to 25, Persons Aged 18 to 25 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 18 to 25: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 4.7P Past Year Initiation of Substance Use among Persons Aged 18 to 25, Persons Aged 18 to 25 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 18 to 25: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 4.8D Past Year Initiation of Substance Use among Persons Aged 26 or Older, Persons Aged 26 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 26 or Older: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 4.8P Past Year Initiation of Substance Use among Persons Aged 26 or Older, Persons Aged 26 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 26 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 4.9C Past Year Initiation of Substance Use among Persons Aged 12 or Older, by Gender: Standard Errors of Numbers in Thousands, 2013 and 2014
- 4.10C Past Year Initiation of Substance Use among Persons Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: Standard Errors of Numbers in Thousands, 2013 and 2014
- 4.11C Past Year Initiation of Substance Use among Persons Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: Standard Errors of Numbers in Thousands, 2013 and 2014

LIST OF TABLES (Continued)

Table

4.12C	Past Year Initiation of Substance Use among Persons Aged 12 to 49, by Gender: Standard Errors of Numbers in Thousands, 2013 and 2014
4.13D	Standard Errors of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older, by Gender: 2013 and 2014
4.13P	<i>P</i> Values from Tests of Differences of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older, by Gender: 2014 Versus 2013
4.14D	Standard Errors of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: 2013 and 2014
4.14P	<i>P</i> Values from Tests of Differences of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: 2014 Versus 2013
4.15D	Standard Errors of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: 2013 and 2014
4.15P	<i>P</i> Values from Tests of Differences of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: 2014 Versus 2013
4.16D	Standard Errors of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49, by Gender: 2013 and 2014
4.16P	<i>P</i> Values from Tests of Differences of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49, by Gender: 2014 Versus 2013
5.1C	Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014
5.1D	Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014
5.1P	Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.2C	Substance Dependence or Abuse for Specific Substances in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
5.2D	Substance Dependence or Abuse for Specific Substances in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
5.2P	Substance Dependence or Abuse for Specific Substances in the Past Year, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.3C	Substance Dependence or Abuse in the Past Year, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
5.3D	Substance Dependence or Abuse in the Past Year, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
5.3P	Substance Dependence or Abuse in the Past Year, by Detailed Age Category: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.4C	Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
5.4D	Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
5.4P	Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.5C	Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
5.5D	Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
5.5P	Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.6C	Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
5.6D	Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 5.6P Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.7C Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.7D Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.7P Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.8C Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.8D Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.8P Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.9C Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.9D Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.9P Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.10C Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.10D Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.10P Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.11C Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.11D Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.11P Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.12C Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.12D Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.12P Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.13C Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.13D Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.13P Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.14C Substance Dependence for Specific Substances in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.14D Substance Dependence for Specific Substances in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 5.14P Substance Dependence for Specific Substances in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.15C Received Substance Use Treatment in the Past Year, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

LIST OF TABLES (Continued)

Table

5.15D	Received Substance Use Treatment in the Past Year, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
5.15P	Received Substance Use Treatment in the Past Year, by Detailed Age Category: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.16C	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
5.16D	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
5.16P	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.17C	Received Substance Use Treatment in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
5.17D	Received Substance Use Treatment in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
5.17P	Received Substance Use Treatment in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.18C	Received Substance Use Treatment in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
5.18D	Received Substance Use Treatment in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
5.18P	Received Substance Use Treatment in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.19C	Received Substance Use Treatment in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
5.19D	Received Substance Use Treatment in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
5.19P	Received Substance Use Treatment in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.20C	Received Substance Use Treatment in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
5.20D	Received Substance Use Treatment in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
5.20P	Received Substance Use Treatment in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.21C	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
5.21D	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
5.21P	Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.22C	Substances for Which Last or Current Treatment Was Received among Persons Who Received Substance Use Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
5.22D	Substances for Which Last or Current Treatment Was Received among Persons Who Received Substance Use Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
5.22P	Substances for Which Last or Current Treatment Was Received among Persons Who Received Substance Use Treatment in the Past Year, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
5.23C	Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
5.23D	Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
5.23P	Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment in the Past Year, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 5.24C Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.24D Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 5.24P Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.25C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.25D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 5.25P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.26C Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.26D Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Percentages, 2013 and 2014
- 5.26P Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.27C Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.27D Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 5.27P Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.28C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.28D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.28P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.29C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.29D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.29P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.30C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.30D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.30P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.31C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.31D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.31P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 5.32C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.32D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.32P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.33C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.33D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.33P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.34C Substances for Which Last or Current Treatment Was Received among Persons Who Received Their Last or Current Substance Use Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.34D Substances for Which Last or Current Treatment Was Received among Persons Who Received Their Last or Current Substance Use Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 5.34P Substances for Which Last or Current Treatment Was Received among Persons Who Received Their Last or Current Substance Use Treatment at a Specialty Facility in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.35C Source of Payment for Last or Current Illicit Drug Treatment among Persons Who Received Their Last or Current Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.35D Source of Payment for Last or Current Illicit Drug Treatment among Persons Who Received Their Last or Current Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 5.35P Source of Payment for Last or Current Illicit Drug Treatment among Persons Who Received Their Last or Current Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.36C Source of Payment for Last or Current Alcohol Treatment among Persons Who Received Their Last or Current Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.36D Source of Payment for Last or Current Alcohol Treatment among Persons Who Received Their Last or Current Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 5.36P Source of Payment for Last or Current Alcohol Treatment among Persons Who Received Their Last or Current Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.37C Source of Payment for Last or Current Illicit Drug or Alcohol Treatment among Persons Who Received Their Last or Current Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.37D Source of Payment for Last or Current Illicit Drug or Alcohol Treatment among Persons Who Received Their Last or Current Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 5.37P Source of Payment for Last or Current Illicit Drug or Alcohol Treatment among Persons Who Received Their Last or Current Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.38C Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 5.38D Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 5.38P Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.39C Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.39D Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 5.39P Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.40C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.40D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 5.40P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.41C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.41D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Percentages, 2013 and 2014
- 5.41P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.42C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse of Specific Substances: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.42D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse of Specific Substances: Standard Errors of Percentages, 2013 and 2014
- 5.42P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse of Specific Substances: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.43C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year, by Detailed Age Category: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 5.43D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 5.43P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.44C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 5.44D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 5.44P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.45C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 5.45D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.45P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.46C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 5.46D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.46P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.47C Perceived Need for Illicit Drug Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug Problem, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.47D Perceived Need for Illicit Drug Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug Problem, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.47P Perceived Need for Illicit Drug Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug Problem, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.48C Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 5.48D Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.48P Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.49C Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 5.49D Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.49P Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.50C Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.50D Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 5.50P Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.51C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 5.51D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.51P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.52C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 5.52D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.52P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.53C Perceived Need for Illicit Drug or Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug or Alcohol Problem, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 5.53D Perceived Need for Illicit Drug or Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug or Alcohol Problem, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 5.53P Perceived Need for Illicit Drug or Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug or Alcohol Problem, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 5.54C Detailed Reasons for Not Receiving Illicit Drug Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2014
- 5.54D Detailed Reasons for Not Receiving Illicit Drug Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2011-2014
- 5.55C Detailed Reasons for Not Receiving Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Alcohol Treatment at a Specialty Facility and Who Felt a Need for Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2014
- 5.55D Detailed Reasons for Not Receiving Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Alcohol Treatment at a Specialty Facility and Who Felt a Need for Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2011-2014
- 5.56C Detailed Reasons for Not Receiving Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug or Alcohol Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug or Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2014

LIST OF TABLES (Continued)

Table

- 5.56D Detailed Reasons for Not Receiving Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug or Alcohol Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug or Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2011-2014
- 6.1C Number of Days Used Marijuana in the Past Year among Past Year Users and the Number of Days Used Marijuana in the Past Month among Past Month Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.1D Number of Days Used Marijuana in the Past Year among Past Year Users and the Number of Days Used Marijuana in the Past Month among Past Month Users, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.1P Number of Days Used Marijuana in the Past Year among Past Year Users and the Number of Days Used Marijuana in the Past Month among Past Month Users, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.2C Number of Days Used Cocaine in the Past Year among Past Year Users and the Number of Days Used Cocaine in the Past Month among Past Month Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.2D Number of Days Used Cocaine in the Past Year among Past Year Users and the Number of Days Used Cocaine in the Past Month among Past Month Users, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.2P Number of Days Used Cocaine in the Past Year among Past Year Users and the Number of Days Used Cocaine in the Past Month among Past Month Users, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.3C Number of Days Used Hallucinogens in the Past Year among Past Year Users and the Number Days Used Hallucinogens in the Past Month among Past Month Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.3D Number of Days Used Hallucinogens in the Past Year among Past Year Users and the Number Days Used Hallucinogens in the Past Month among Past Month Users, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.3P Number of Days Used Hallucinogens in the Past Year among Past Year Users and the Number Days Used Hallucinogens in the Past Month among Past Month Users, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.4C Number of Days Used Inhalants in the Past Year among Past Year Users and the Number of Days Used Inhalants in the Past Month among Past Month Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.4D Number of Days Used Inhalants in the Past Year among Past Year Users and the Number of Days Used Inhalants in the Past Month among Past Month Users, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.4P Number of Days Used Inhalants in the Past Year among Past Year Users and the Number of Days Used Inhalants in the Past Month among Past Month Users, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.5C Number of Days Used Cigarettes in the Past Month among Past Month Users and the Number of Cigarettes Smoked Per Day among Daily Smokers, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.5D Number of Days Used Cigarettes in the Past Month among Past Month Users and the Number of Cigarettes Smoked Per Day among Daily Smokers, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.5P Number of Days Used Cigarettes in the Past Month among Past Month Users and the Number of Cigarettes Smoked Per Day among Daily Smokers, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.6C Number of Days Used Alcohol in the Past Year among Past Year Users and the Number of Days Used Alcohol in the Past Month among Past Month Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.6D Number of Days Used Alcohol in the Past Year among Past Year Users and the Number of Days Used Alcohol in the Past Month among Past Month Users, by Age Group: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 6.6P Number of Days Used Alcohol in the Past Year among Past Year Users and the Number of Days Used Alcohol in the Past Month among Past Month Users, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.7C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.7D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014
- 6.7P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.8C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.8D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014
- 6.8P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.9C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.9D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014
- 6.9P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.10C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.10D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014
- 6.10P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.11C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.11D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014
- 6.11P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.12C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.12D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014
- 6.12P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.13C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.13D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014
- 6.13P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.14C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.14D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014
- 6.14P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.15C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 6.15D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2013 and 2014
- 6.15P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.16C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.16D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2013 and 2014
- 6.16P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.17C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.17D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.17P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.18C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.18D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.18P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.19C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.19D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2013 and 2014
- 6.19P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.20C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.20D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2013 and 2014
- 6.20P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.21C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.21D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2013 and 2014
- 6.21P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.22C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.22D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2013 and 2014
- 6.22P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

6.23C	Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
6.23D	Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Percentages, 2013 and 2014
6.23P	Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.24C	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014
6.24D	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014
6.24P	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.25C	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014
6.25D	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014
6.25P	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.26C	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014
6.26D	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014
6.26P	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.27C	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014
6.27D	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014
6.27P	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.28C	Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
6.28D	Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014
6.28P	Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.29C	Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
6.29D	Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014
6.29P	Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.30C	Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
6.30D	Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014
6.30P	Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.31C	Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014
6.31D	Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014
6.31P	Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 6.32C Marijuana Use in the Prior Year and Past Year Marijuana Use Status among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.32D Marijuana Use in the Prior Year and Past Year Marijuana Use Status among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.32P Marijuana Use in the Prior Year and Past Year Marijuana Use Status among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.33C Cigarette Use and Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014
- 6.33D Cigarette Use and Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 6.33P Cigarette Use and Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.34C Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.34D Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.34P Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.35C Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.35D Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.35P Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.36C Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.36D Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.36P Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.37C Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes Daily and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.37D Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes Daily and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.37P Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes Daily and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.38C Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.38D Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.38P Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.39C Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.39D Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

LIST OF TABLES (Continued)

Table

- 6.39P Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.40C Alcohol Dependence or Abuse in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.40D Alcohol Dependence or Abuse in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.40P Alcohol Dependence or Abuse in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.41C Illicit Drug Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.41D Illicit Drug Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.41P Illicit Drug Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.42C Alcohol Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.42D Alcohol Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.42P Alcohol Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.43C Alcohol Dependence in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.43D Alcohol Dependence in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.43P Alcohol Dependence in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.44C Method of Obtaining Most Recently Used Marijuana among Past Year Marijuana Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.44D Method of Obtaining Most Recently Used Marijuana among Past Year Marijuana Users, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.44P Method of Obtaining Most Recently Used Marijuana among Past Year Marijuana Users, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.45C Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Bought Their Most Recently Used Marijuana, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.45D Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Bought Their Most Recently Used Marijuana, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.45P Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Bought Their Most Recently Used Marijuana, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.46C Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Obtained for Free Their Most Recently Used Marijuana and Who Had Not Bought or Traded for Marijuana in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.46D Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Obtained for Free Their Most Recently Used Marijuana and Who Had Not Bought or Traded for Marijuana in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.46P Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Obtained for Free Their Most Recently Used Marijuana and Who Had Not Bought or Traded for Marijuana in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.47C Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

LIST OF TABLES (Continued)

Table

- 6.47D Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.47P Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.48C Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.48D Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.48P Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.49C Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.49D Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.49P Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.50C Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.50D Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.50P Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.51C Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.51D Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.51P Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.52C Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.52D Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.52P Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.53C Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

LIST OF TABLES (Continued)

Table

- 6.53D Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.53P Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.54C Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.54D Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.54P Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.55C Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.55D Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.55P Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.56C Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.56D Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.56P Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.57C Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.57D Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.57P Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.58C Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.58D Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.58P Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.59C Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

LIST OF TABLES (Continued)

Table

- 6.59D Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.59P Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.60C Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.60D Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.60P Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.61C Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.61D Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.61P Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.62C Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.62D Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.62P Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.63C Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.63D Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.63P Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.64C Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.64D Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.64P Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

LIST OF TABLES (Continued)

Table

- 6.65C Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.65D Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.65P Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.66C Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.66D Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.66P Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.67C Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.67D Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Standard Errors of Percentages, 2013 and 2014
- 6.67P Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.68C Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.68D Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.68P Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.69C Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.69D Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Standard Errors of Percentages, 2013 and 2014
- 6.69P Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.70C Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.70D Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Standard Errors of Percentages, 2013 and 2014
- 6.70P Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.71C Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
- 6.71D Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
- 6.71P Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
- 6.72C Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

LIST OF TABLES (Continued)

Table

6.72D	Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
6.72P	Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
6.73C	Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
6.73D	Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
6.73P	Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
6.74C	Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
6.74D	Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
6.74P	Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: <i>P</i> Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
6.75C	Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
6.75D	Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
6.75P	Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
6.76C	Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
6.76D	Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
6.76P	Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
6.77C	Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
6.77D	Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
6.77P	Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
6.78C	Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014
6.78D	Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014
6.78P	Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012
6.79C	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014
6.79D	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014
6.79P	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.80C	Types of Illicit Drug Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014

LIST OF TABLES (Continued)

Table

6.80D	Types of Illicit Drug Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014
6.80P	Types of Illicit Drug Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.81C	Types of Illicit Drug Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014
6.81D	Types of Illicit Drug Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014
6.81P	Types of Illicit Drug Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.82C	Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
6.82D	Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
6.82P	Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.83C	Marijuana Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
6.83D	Marijuana Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
6.83P	Marijuana Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.84C	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014
6.84D	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014
6.84P	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.85C	Tobacco Product and Alcohol Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014
6.85D	Tobacco Product and Alcohol Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014
6.85P	Tobacco Product and Alcohol Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.86C	Tobacco Product and Alcohol Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014
6.86D	Tobacco Product and Alcohol Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014
6.86P	Tobacco Product and Alcohol Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.87C	Cigarette Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
6.87D	Cigarette Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
6.87P	Cigarette Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.88C	Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
6.88D	Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
6.88P	Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013

LIST OF TABLES (Continued)

Table

- 6.89C Binge Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.89D Binge Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.89P Binge Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.90C Heavy Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.90D Heavy Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.90P Heavy Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.91C Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.91D Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.91P Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.92C Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.92D Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.92P Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.93C Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.93D Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.93P Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.94C Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.94D Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014
- 6.94P Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.95C Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.95D Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.95P Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.96C Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
- 6.96D Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
- 6.96P Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013
- 6.97C Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

LIST OF TABLES (Continued)

Table

6.97D	Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014
6.97P	Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.98C	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status: Standard Errors of Numbers in Thousands, 2013 and 2014
6.98D	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status: Standard Errors of Percentages, 2013 and 2014
6.98P	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.99C	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Probation Status: Standard Errors of Numbers in Thousands, 2013 and 2014
6.99D	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Probation Status: Standard Errors of Percentages, 2013 and 2014
6.99P	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Probation Status: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.100C	Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
6.100D	Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
6.100P	Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.101C	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
6.101D	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
6.101P	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.102C	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
6.102D	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
6.102P	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.103C	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status: Standard Errors of Numbers in Thousands, 2013 and 2014
6.103D	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status: Standard Errors of Percentages, 2013 and 2014
6.103P	Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.104C	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Parole/Supervised Release Status: Standard Errors of Numbers in Thousands, 2013 and 2014
6.104D	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Parole/Supervised Release Status: Standard Errors of Percentages, 2013 and 2014
6.104P	Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Parole/Supervised Release Status: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.105C	Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

LIST OF TABLES (Continued)

Table

6.105D	Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
6.105P	Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.106C	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
6.106D	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
6.106P	Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
6.107C	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014
6.107D	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014
6.107P	Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2013
7.1C	Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.1D	Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014
7.1P	Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.2C	Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.2D	Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014
7.2P	Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.3C	Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.3D	Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014
7.3P	Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.4C	Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
7.4D	Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014
7.4P	Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.5C	Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
7.5D	Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014
7.5P	Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

LIST OF TABLES (Continued)

Table

- 7.6C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
- 7.6D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014
- 7.6P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.7C Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014
- 7.7D Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014
- 7.7P Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.8C Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014
- 7.8D Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014
- 7.8P Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.9C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014
- 7.9D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014
- 7.9P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.10C Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014
- 7.10D Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014
- 7.10P Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.11C Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014
- 7.11D Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014
- 7.11P Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.12C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014
- 7.12D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014
- 7.12P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.13C Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Standard Errors of Numbers in Thousands, 2002-2014
- 7.13D Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Standard Errors of Percentages, 2002-2014
- 7.13P Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.14C Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Standard Errors of Numbers in Thousands, 2002-2014
- 7.14D Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Standard Errors of Percentages, 2002-2014
- 7.14P Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

LIST OF TABLES (Continued)

Table

- 7.15C Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Standard Errors of Numbers in Thousands, 2002-2014
- 7.15D Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Standard Errors of Percentages, 2002-2014
- 7.15P Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.16C Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
- 7.16D Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Standard Errors of Percentages, 2002-2014
- 7.16P Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.17C Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
- 7.17D Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Standard Errors of Percentages, 2002-2014
- 7.17P Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.18C Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
- 7.18D Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Standard Errors of Percentages, 2002-2014
- 7.18P Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.19C Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
- 7.19D Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Standard Errors of Percentages, 2002-2014
- 7.19P Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.20C Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
- 7.20D Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Standard Errors of Percentages, 2002-2014
- 7.20P Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.21C Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
- 7.21D Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Standard Errors of Percentages, 2002-2014
- 7.21P Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.22C GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014
- 7.22D GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014
- 7.22P GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2006-2013
- 7.23C Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014
- 7.23D Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014
- 7.23P Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2006-2013

LIST OF TABLES (Continued)

Table

- 7.24C Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014
- 7.24D Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014
- 7.24P Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2006-2013
- 7.25C Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014
- 7.25D Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014
- 7.25P Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2006-2013
- 7.26C Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014
- 7.26D Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014
- 7.26P Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2006-2013
- 7.27C DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014
- 7.27D DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014
- 7.27P DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2006-2013
- 7.28C Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014
- 7.28D Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014
- 7.28P Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2006-2013
- 7.29C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2014
- 7.29D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Percentages, 2002-2014
- 7.29P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.30C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Numbers in Thousands, 2002-2014
- 7.30D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Percentages, 2002-2014
- 7.30P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.31C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Numbers in Thousands, 2002-2014
- 7.31D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Percentages, 2002-2014
- 7.31P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.32C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2014

LIST OF TABLES (Continued)

Table

7.32D	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Percentages, 2002-2014
7.32P	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.33C	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2014
7.33D	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Percentages, 2002-2014
7.33P	Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.34C	Daily Cigarette Use in the Past Month among Past Month Users Aged 12 or Older and Smoked One or More Packs of Cigarettes Per Day among Daily Cigarette Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014
7.34D	Daily Cigarette Use in the Past Month among Past Month Users Aged 12 or Older and Smoked One or More Packs of Cigarettes Per Day among Daily Cigarette Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2002-2014
7.34P	Daily Cigarette Use in the Past Month among Past Month Users Aged 12 or Older and Smoked One or More Packs of Cigarettes Per Day among Daily Cigarette Users Aged 12 or Older, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.35C	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Gender: Standard Errors of Numbers in Thousands, 2002-2014
7.35D	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Gender: Standard Errors of Percentages, 2002-2014
7.35P	Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Gender: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.36C	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.36D	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014
7.36P	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.37C	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
7.37D	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014
7.37P	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 to 17: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.38C	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014
7.38D	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014
7.38P	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 18 to 25: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.39C	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.39D	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014
7.39P	Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 26 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.40C	Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Youths Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
7.40D	Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Youths Aged 12 to 17: Standard Errors of Percentages, 2002-2014

LIST OF TABLES (Continued)

Table

- 7.40P Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Youths Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.41C Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Youths Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
- 7.41D Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Youths Aged 12 to 17: Standard Errors of Percentages, 2002-2014
- 7.41P Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Youths Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.42C Exposure to Substance Use Prevention Program or Message in the Past Year among Youths Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
- 7.42D Exposure to Substance Use Prevention Program or Message in the Past Year among Youths Aged 12 to 17: Standard Errors of Percentages, 2002-2014
- 7.42P Exposure to Substance Use Prevention Program or Message in the Past Year among Youths Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.43C Substance Use in the Past Month among Youths Aged 12 to 17, by Exposure to Substance Use Prevention Program or Message in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014
- 7.43D Substance Use in the Past Month among Youths Aged 12 to 17, by Exposure to Substance Use Prevention Program or Message in the Past Year: Standard Errors of Percentages, 2002-2014
- 7.43P Substance Use in the Past Month among Youths Aged 12 to 17, by Exposure to Substance Use Prevention Program or Message in the Past Year: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.44C Past Year Initiation of Substance Use among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014
- 7.44D Past Year Initiation of Substance Use among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014
- 7.44P Past Year Initiation of Substance Use among Persons Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.45C Past Year Initiation of Substance Use among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
- 7.45D Past Year Initiation of Substance Use among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014
- 7.45P Past Year Initiation of Substance Use among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.46C Past Year Initiation of Substance Use among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014
- 7.46D Past Year Initiation of Substance Use among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014
- 7.46P Past Year Initiation of Substance Use among Persons Aged 18 to 25: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.47C Past Year Initiation of Substance Use among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014
- 7.47D Past Year Initiation of Substance Use among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014
- 7.47P Past Year Initiation of Substance Use among Persons Aged 26 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.48C Past Year Initiation of Illicit Drugs and First Drug Initiated in the Past Year among Persons Aged 12 or Older Who Initiated Illicit Drug Use in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014
- 7.48D Past Year Initiation of Illicit Drugs and First Drug Initiated in the Past Year among Persons Aged 12 or Older Who Initiated Illicit Drug Use in the Past Year: Standard Errors of Percentages, 2002-2014
- 7.48P Past Year Initiation of Illicit Drugs and First Drug Initiated in the Past Year among Persons Aged 12 or Older Who Initiated Illicit Drug Use in the Past Year: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

LIST OF TABLES (Continued)

Table

7.49D	Standard Errors of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49: 2002-2014
7.49P	<i>P</i> Values from Tests of Differences of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49: 2014 Versus 2002-2013
7.50C	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.50D	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014
7.50P	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.51C	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
7.51D	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014
7.51P	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 to 17: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.52C	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014
7.52D	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014
7.52P	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 18 to 25: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.53C	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.53D	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014
7.53P	Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 26 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.54C	Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.54D	Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014
7.54P	Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.55C	Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
7.55D	Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014
7.55P	Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 to 17: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.56C	Substance Dependence for Specific Substances in the Past Year among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014
7.56D	Substance Dependence for Specific Substances in the Past Year among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014
7.56P	Substance Dependence for Specific Substances in the Past Year among Persons Aged 18 to 25: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.57C	Substance Dependence for Specific Substances in the Past Year among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.57D	Substance Dependence for Specific Substances in the Past Year among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014
7.57P	Substance Dependence for Specific Substances in the Past Year among Persons Aged 26 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

LIST OF TABLES (Continued)

Table

7.58C	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.58D	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014
7.58P	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.59C	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
7.59D	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014
7.59P	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 to 17: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.60C	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014
7.60D	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014
7.60P	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 18 to 25: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.61C	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.61D	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014
7.61P	Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 26 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.62C	Substances for Which Last or Current Treatment Was Received among Persons Aged 12 or Older Who Received Substance Use Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014
7.62D	Substances for Which Last or Current Treatment Was Received among Persons Aged 12 or Older Who Received Substance Use Treatment in the Past Year: Standard Errors of Percentages, 2002-2014
7.62P	Substances for Which Last or Current Treatment Was Received among Persons Aged 12 or Older Who Received Substance Use Treatment in the Past Year: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.63C	Substances for Which Last or Current Treatment Was Received among Persons Aged 12 to 17 Who Received Substance Use Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014
7.63D	Substances for Which Last or Current Treatment Was Received among Persons Aged 12 to 17 Who Received Substance Use Treatment in the Past Year: Standard Errors of Percentages, 2002-2014
7.63P	Substances for Which Last or Current Treatment Was Received among Persons Aged 12 to 17 Who Received Substance Use Treatment in the Past Year: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.64C	Substances for Which Last or Current Treatment Was Received among Persons Aged 18 to 25 Who Received Substance Use Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014
7.64D	Substances for Which Last or Current Treatment Was Received among Persons Aged 18 to 25 Who Received Substance Use Treatment in the Past Year: Standard Errors of Percentages, 2002-2014
7.64P	Substances for Which Last or Current Treatment Was Received among Persons Aged 18 to 25 Who Received Substance Use Treatment in the Past Year: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.65C	Substances for Which Last or Current Treatment Was Received among Persons Aged 26 or Older Who Received Substance Use Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014

LIST OF TABLES (Continued)

Table

- 7.65D Substances for Which Last or Current Treatment Was Received among Persons Aged 26 or Older Who Received Substance Use Treatment in the Past Year: Standard Errors of Percentages, 2002-2014
- 7.65P Substances for Which Last or Current Treatment Was Received among Persons Aged 26 or Older Who Received Substance Use Treatment in the Past Year: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.66C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014
- 7.66D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Percentages, 2002-2014
- 7.66P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.67C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 to 17 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014
- 7.67D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 to 17 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Percentages, 2002-2014
- 7.67P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 to 17 Who Received Illicit Drug or Alcohol Treatment in the Past Year: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.68C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 18 to 25 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014
- 7.68D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 18 to 25 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Percentages, 2002-2014
- 7.68P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 18 to 25 Who Received Illicit Drug or Alcohol Treatment in the Past Year: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.69C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 26 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014
- 7.69D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 26 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Percentages, 2002-2014
- 7.69P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 26 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.70C Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014
- 7.70D Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014
- 7.70P Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
- 7.71C Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014
- 7.71D Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014
- 7.71P Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

LIST OF TABLES (Continued)

Table

7.72C	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014
7.72D	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014
7.72P	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 18 to 25: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.73C	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014
7.73D	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014
7.73P	Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 26 or Older: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.74C	Type of Substance Treatment Received at Any Location in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014
7.74D	Type of Substance Treatment Received at Any Location in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use in the Past Year, by Age Group: Standard Errors of Percentages, 2002-2014
7.74P	Type of Substance Treatment Received at Any Location in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use in the Past Year, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.75C	Type of Substance Treatment Received at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014
7.75D	Type of Substance Treatment Received at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2002-2014
7.75P	Type of Substance Treatment Received at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use at a Specialty Facility in the Past Year, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.76C	Receipt of Illicit Drug Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug Treatment, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014
7.76D	Receipt of Illicit Drug Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug Treatment, by Age Group: Standard Errors of Percentages, 2002-2014
7.76P	Receipt of Illicit Drug Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug Treatment, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.77C	Receipt of Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Alcohol Treatment, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014
7.77D	Receipt of Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Alcohol Treatment, by Age Group: Standard Errors of Percentages, 2002-2014
7.77P	Receipt of Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Alcohol Treatment, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013
7.78C	Receipt of Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014
7.78D	Receipt of Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment, by Age Group: Standard Errors of Percentages, 2002-2014
7.78P	Receipt of Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment, by Age Group: <i>P</i> Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

LIST OF TABLES (Continued)

Table

7.79D	Marijuana Use in the Past Month, by Age Group: Standard Errors of Percentages, 1971-2014
7.80D	Cocaine Use in the Past Year, by Age Group: Standard Errors of Percentages, 1971-2014
8.1C	Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Gender and Detailed Age Category: 2013 and 2014
8.2C	Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Age Group and Demographic Characteristics: 2013 and 2014
8.3C	Standard Errors of Numbers (in Thousands) of Persons Aged 18 or Older, by Age Group and Demographic Characteristics: 2013 and 2014
8.4C	Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Youth and Adult Age Groups and Demographic Characteristics: 2013 and 2014
8.5C	Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Underage and Legal Drinking Age Groups and Demographic Characteristics: 2013 and 2014
8.6C	Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Age Group and Geographic Characteristics: 2013 and 2014
8.7C	Standard Errors of Numbers (in Thousands) of Persons Aged 18 or Older, by Age Group and Geographic Characteristics: 2013 and 2014
8.8C	Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Underage and Legal Drinking Age Groups and Geographic Characteristics: 2013 and 2014
8.9C	Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Youth and Adult Age Groups and Gender: 2002-2014
8.10C	Standard Errors of Numbers (in Thousands) of Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Annual Averages Based on 2011-2012 and 2013-2014
8.11C	Standard Errors of Numbers (in Thousands) of Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: 2013 and 2014
8.12C	Standard Errors of Numbers (in Thousands) of Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: 2013 and 2014
8.13C	Standard Errors of Numbers (in Thousands) of Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: 2013 and 2014

Table 1.1A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	127,458 ^a	130,332	41,591 ^b	44,157	24,573 ^b	26,983
Marijuana and Hashish	114,712	117,213	32,952 ^b	35,124	19,810 ^b	22,188
Cocaine	37,634	39,200	4,182	4,553	1,549	1,530
Crack	8,870	9,424	632	773	377	354
Heroin	4,812	4,813	681 ^a	914	289 ^a	435
Hallucinogens	39,736	39,647	4,430	4,250	1,333	1,173
LSD	24,770	25,035	1,111	1,290	229	287
PCP	6,462	6,388	90	90	33	*
Ecstasy	17,815	17,548	2,588	2,342	660	609
Inhalants	21,068	21,293	1,533	1,617	496	546
Nonmedical Use of Psychotherapeutics ^{2,3}	53,172	54,395	15,348	14,966	6,484	6,537
Pain Relievers	35,473	36,064	11,082	10,337	4,521	4,325
OxyContin [®]	6,973	7,031	1,442	1,237	492	356
Tranquilizers	23,493	24,851	5,269	5,202	1,705	1,875
Stimulants ³	21,656	22,530	3,492	3,715	1,365	1,595
Methamphetamine ³	12,257	12,943	1,186	1,301	595	569
Sedatives	7,480	7,826	639	775	251	330
ILLCIT DRUGS OTHER THAN MARIJUANA¹	78,076	80,119	19,868	19,719	8,665	8,719

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.1B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	48.6	49.2	15.9 ^a	16.7	9.4 ^b	10.2
Marijuana and Hashish	43.7	44.2	12.6 ^a	13.2	7.5 ^b	8.4
Cocaine	14.3	14.8	1.6	1.7	0.6	0.6
Crack	3.4	3.6	0.2	0.3	0.1	0.1
Heroin	1.8	1.8	0.3 ^a	0.3	0.1 ^a	0.2
Hallucinogens	15.1	15.0	1.7	1.6	0.5	0.4
LSD	9.4	9.4	0.4	0.5	0.1	0.1
PCP	2.5	2.4	0.0	0.0	0.0	*
Ecstasy	6.8	6.6	1.0	0.9	0.3	0.2
Inhalants	8.0	8.0	0.6	0.6	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{2,3}	20.3	20.5	5.8	5.6	2.5	2.5
Pain Relievers	13.5	13.6	4.2 ^a	3.9	1.7	1.6
OxyContin [®]	2.7	2.7	0.5	0.5	0.2	0.1
Tranquilizers	9.0	9.4	2.0	2.0	0.6	0.7
Stimulants ³	8.3	8.5	1.3	1.4	0.5	0.6
Methamphetamine ³	4.7	4.9	0.5	0.5	0.2	0.2
Sedatives	2.9	3.0	0.2	0.3	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA¹	29.8	30.2	7.6	7.4	3.3	3.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.2A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	772	822	466	556	207	266
Marijuana and Hashish	240	215	197	186	81	89
Cocaine	9	10	8	1	2	*
Crack	2	1	2	*	2	*
Heroin	6	2	5	2	1	*
Hallucinogens	58	47	35	25	11	10
LSD	15	8	9	2	5	*
PCP	6	6	2	2	1	*
Ecstasy	19	8	18	8	1	3
Inhalants	336	342	145	162	47	55
Nonmedical Use of Psychotherapeutics ^{2,3}	330 ^a	422	187 ^b	283	100	144
Pain Relievers	267	316	150 ^a	209	75	103
OxyContin [®]	22	17	13	13	6	6
Tranquilizers	47	70	31	36	16	6
Stimulants ³	49	80	25 ^a	59	14	35
Methamphetamine ³	12	20	7	12	4	8
Sedatives	28	54	18	29	12	13
ILLICIT DRUGS OTHER THAN MARIJUANA¹	612	691	322 ^b	431	143 ^a	200

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.2B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	9.7	10.4	5.8	7.0	2.6 ^a	3.4
Marijuana and Hashish	3.0	2.7	2.5	2.4	1.0	1.1
Cocaine	0.1	0.1	0.1	0.0	0.0	*
Crack	0.0	0.0	0.0	*	0.0	*
Heroin	0.1	0.0	0.1	0.0	0.0	*
Hallucinogens	0.7	0.6	0.4	0.3	0.1	0.1
LSD	0.2	0.1	0.1	0.0	0.1	*
PCP	0.1	0.1	0.0	0.0	0.0	*
Ecstasy	0.2	0.1	0.2	0.1	0.0	0.0
Inhalants	4.2	4.3	1.8	2.1	0.6	0.7
Nonmedical Use of Psychotherapeutics ^{2,3}	4.1 ^a	5.3	2.3 ^b	3.6	1.3	1.8
Pain Relievers	3.3	4.0	1.9 ^a	2.6	0.9	1.3
OxyContin [®]	0.3	0.2	0.2	0.2	0.1	0.1
Tranquilizers	0.6	0.9	0.4	0.5	0.2	0.1
Stimulants ³	0.6	1.0	0.3 ^a	0.7	0.2	0.4
Methamphetamine ³	0.1	0.3	0.1	0.2	0.1	0.1
Sedatives	0.4	0.7	0.2	0.4	0.1	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA¹	7.7	8.7	4.0 ^b	5.5	1.8 ^a	2.5

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.3A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	1,862	1,837	1,343	1,367	664	664
Marijuana and Hashish	1,210	1,168	975	955	492	468
Cocaine	37	41	25	30	12	11
Crack	5	9	2	5	*	1
Heroin	10	2	9	2	5	2
Hallucinogens	151	158	99	113	37	30
LSD	45	75	30 ^a	60	6	13
PCP	17	28	10	18	4	*
Ecstasy	71	70	39	47	10	10
Inhalants	552	541	197	238	50	81
Nonmedical Use of Psychotherapeutics ^{2,3}	694	681	477	483	184	217
Pain Relievers	590	567	402	388	158	173
OxyContin [®]	68	74	48	57	9	14
Tranquilizers	165	171	95	117	30	28
Stimulants ³	123	119	77	89	18 ^a	53
Methamphetamine ³	25	46	13 ^a	35	2	16
Sedatives	48	72	22	45	10	7
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1,104	1,099	649	686	244	296

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.3B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	21.9	21.7	15.8	16.2	7.8	7.9
Marijuana and Hashish	14.2	13.8	11.4	11.3	5.8	5.5
Cocaine	0.4	0.5	0.3	0.4	0.1	0.1
Crack	0.1	0.1	0.0	0.1	*	0.0
Heroin	0.1	0.0	0.1	0.0	0.1	0.0
Hallucinogens	1.8	1.9	1.2	1.3	0.4	0.4
LSD	0.5	0.9	0.3 ^a	0.7	0.1	0.2
PCP	0.2	0.3	0.1	0.2	0.0	*
Ecstasy	0.8	0.8	0.5	0.6	0.1	0.1
Inhalants	6.5	6.4	2.3	2.8	0.6	1.0
Nonmedical Use of Psychotherapeutics ^{2,3}	8.1	8.0	5.6	5.7	2.2	2.6
Pain Relievers	6.9	6.7	4.7	4.6	1.8	2.1
OxyContin [®]	0.8	0.9	0.6	0.7	0.1	0.2
Tranquilizers	1.9	2.0	1.1	1.4	0.3	0.3
Stimulants ³	1.4	1.4	0.9	1.1	0.2 ^a	0.6
Methamphetamine ³	0.3	0.5	0.2 ^a	0.4	0.0	0.2
Sedatives	0.6	0.8	0.3	0.5	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA¹	13.0	13.0	7.6	8.1	2.9	3.5

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.4A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	3,160	3,131	2,477	2,413	1,326	1,407
Marijuana and Hashish	2,643	2,708	2,174	2,120	1,189	1,273
Cocaine	170	171	98	136	30	29
Crack	15	24	7	15	1	7
Heroin	23	32	17	25	6	14
Hallucinogens	493 ^a	405	348	283	106	97
LSD	160	208	111	152	40	52
PCP	23	27	17	14	2	3
Ecstasy	280 ^a	218	176 ^a	126	37	26
Inhalants	439	424	139	112	24	13
Nonmedical Use of Psychotherapeutics ^{2,3}	1,165	1,187	778	788	264	293
Pain Relievers	949	920	605	568	193	191
OxyContin [®]	184	161	89	92	14	15
Tranquilizers	360	425	221	271	45	69
Stimulants ³	265	323	172	213	52	81
Methamphetamine ³	82	105	53	59	16	21
Sedatives	49	81	24	42	3 ^a	20
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1,563	1,532	1,008	958	370	389

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.4B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	37.6	36.8	29.5	28.3	15.8	16.5
Marijuana and Hashish	31.5	31.8	25.9	24.9	14.2	15.0
Cocaine	2.0	2.0	1.2	1.6	0.4	0.3
Crack	0.2	0.3	0.1	0.2	0.0	0.1
Heroin	0.3	0.4	0.2	0.3	0.1	0.2
Hallucinogens	5.9 ^a	4.8	4.1	3.3	1.3	1.1
LSD	1.9	2.4	1.3	1.8	0.5	0.6
PCP	0.3	0.3	0.2	0.2	0.0	0.0
Ecstasy	3.3 ^a	2.6	2.1 ^a	1.5	0.4	0.3
Inhalants	5.2	5.0	1.7	1.3	0.3	0.1
Nonmedical Use of Psychotherapeutics ^{2,3}	13.9	13.9	9.3	9.3	3.1	3.4
Pain Relievers	11.3	10.8	7.2	6.7	2.3	2.2
OxyContin [®]	2.2	1.9	1.1	1.1	0.2	0.2
Tranquilizers	4.3	5.0	2.6	3.2	0.5	0.8
Stimulants ³	3.1	3.8	2.0	2.5	0.6	0.9
Methamphetamine ³	1.0	1.2	0.6	0.7	0.2	0.2
Sedatives	0.6	0.9	0.3	0.5	0.0 ^a	0.2
ILLICIT DRUGS OTHER THAN MARIJUANA¹	18.6	18.0	12.0	11.3	4.4	4.6

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.5A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	6,819	6,795	5,061	4,943	2,978	2,973
Marijuana and Hashish	6,135	6,067	4,524	4,495	2,684	2,735
Cocaine	875	752	450	419	125	148
Crack	96	83	24	40	4	4
Heroin	126	124	53	67	23	20
Hallucinogens	1,736	1,545	927	834	281 ^b	172
LSD	606	677	301	319	70	37
PCP	77	87	15	20	2	2
Ecstasy	1,220 ^a	1,009	532 ^a	417	112	86
Inhalants	866 ^a	721	218	160	51	34
Nonmedical Use of Psychotherapeutics ^{2,3}	2,892	2,719	1,602	1,428	584	483
Pain Relievers	2,302 ^a	2,046	1,179 ^a	989	389 ^a	293
OxyContin [®]	540	443	150	133	44	22
Tranquilizers	1,224	1,190	545	475	143	131
Stimulants ³	890	769	460	427	152	120
Methamphetamine ³	220	225	110	104	35	32
Sedatives	168 ^a	106	60	51	16	14
ILLCIT DRUGS OTHER THAN MARIJUANA¹	3,700	3,610	2,207	2,023	879 ^a	721

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.5B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	51.7	51.8	38.4	37.7	22.6	22.7
Marijuana and Hashish	46.5	46.3	34.3	34.3	20.3	20.9
Cocaine	6.6	5.7	3.4	3.2	0.9	1.1
Crack	0.7	0.6	0.2	0.3	0.0	0.0
Heroin	1.0	0.9	0.4	0.5	0.2	0.2
Hallucinogens	13.2	11.8	7.0	6.4	2.1 ^b	1.3
LSD	4.6	5.2	2.3	2.4	0.5	0.3
PCP	0.6	0.7	0.1	0.2	0.0	0.0
Ecstasy	9.2 ^a	7.7	4.0 ^a	3.2	0.8	0.7
Inhalants	6.6 ^a	5.5	1.6	1.2	0.4	0.3
Nonmedical Use of Psychotherapeutics ^{2,3}	21.9	20.7	12.1	10.9	4.4	3.7
Pain Relievers	17.4 ^a	15.6	8.9 ^a	7.5	3.0 ^a	2.2
OxyContin [®]	4.1	3.4	1.1	1.0	0.3	0.2
Tranquilizers	9.3	9.1	4.1	3.6	1.1	1.0
Stimulants ³	6.7	5.9	3.5	3.3	1.1	0.9
Methamphetamine ³	1.7	1.7	0.8	0.8	0.3	0.2
Sedatives	1.3 ^a	0.8	0.5	0.4	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA¹	28.0	27.5	16.7	15.4	6.7 ^a	5.5

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.6A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	13,008	13,422	7,388	7,654	4,513	4,703
Marijuana and Hashish	11,915	12,322	6,476	6,659	3,952	4,101
Cocaine	3,166	3,128	1,093	1,185	273	325
Crack	467	463	71	87	32	25
Heroin	493	559	191	200	69	62
Hallucinogens	4,386	4,240	1,411	1,371	336	330
LSD	1,639	1,774	407	481	35 ^a	82
PCP	250	218	11	16	2	3
Ecstasy	3,216	3,191	847	802	207	183
Inhalants	1,746	1,723	260	219	55	46
Nonmedical Use of Psychotherapeutics ^{2,3}	6,347	6,478	2,643	2,704	1,077	1,069
Pain Relievers	4,949	5,003	1,881	1,751	743	685
OxyContin [®]	1,349	1,340	323	293	63	80
Tranquilizers	2,961	3,172	1,023	972	291	285
Stimulants ³	2,389	2,361	841	932	310	286
Methamphetamine ³	821	815	208	246	85	54
Sedatives	271	354	78	102	18	42
ILLCIT DRUGS OTHER THAN MARIJUANA¹	8,153	8,252	3,647	3,757	1,459	1,527

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.6B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	60.2	61.5	34.2	35.1	20.9	21.5
Marijuana and Hashish	55.2	56.5	30.0	30.5	18.3	18.8
Cocaine	14.7	14.3	5.1	5.4	1.3	1.5
Crack	2.2	2.1	0.3	0.4	0.2	0.1
Heroin	2.3	2.6	0.9	0.9	0.3	0.3
Hallucinogens	20.3	19.4	6.5	6.3	1.6	1.5
LSD	7.6	8.1	1.9	2.2	0.2 ^a	0.4
PCP	1.2	1.0	0.1	0.1	0.0	0.0
Ecstasy	14.9	14.6	3.9	3.7	1.0	0.8
Inhalants	8.1	7.9	1.2	1.0	0.3	0.2
Nonmedical Use of Psychotherapeutics ^{2,3}	29.4	29.7	12.2	12.4	5.0	4.9
Pain Relievers	22.9	22.9	8.7	8.0	3.4	3.1
OxyContin [®]	6.2	6.1	1.5	1.3	0.3	0.4
Tranquilizers	13.7	14.5	4.7	4.5	1.4	1.3
Stimulants ³	11.1	10.8	3.9	4.3	1.4	1.3
Methamphetamine ³	3.8	3.7	1.0	1.1	0.4	0.2
Sedatives	1.3	1.6	0.4	0.5	0.1	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA¹	37.8	37.8	16.9	17.2	6.8	7.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.7A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	23,057	23,187	9,501	9,452	5,699	5,728
Marijuana and Hashish	21,263	21,114	7,834	7,756	4,721	4,822
Cocaine	7,053	6,775	1,274	1,140	480	376
Crack	1,767	1,693	151	133	83	69
Heroin	1,151	1,185	249	346	119	172
Hallucinogens	8,766	8,268	917	1,026	238	311
LSD	4,653 ^b	3,922	174	211	41	78
PCP	728	635	13	4	*	*
Ecstasy	6,398	6,069	651	624	181	195
Inhalants	4,578	4,112	228	334	84	135
Nonmedical Use of Psychotherapeutics ^{2,3}	11,462	11,117	3,843 ^a	3,337	1,659	1,539
Pain Relievers	8,735	8,283	2,665	2,276	1,115	983
OxyContin [®]	2,389	2,477	386	284	107	62
Tranquilizers	5,671	5,573	1,523	1,333	521	554
Stimulants ³	4,811	4,582	1,076	920	422	362
Methamphetamine ³	2,531	2,333	295	322	159	108
Sedatives	804	719	175	136	86	35
ILLCIT DRUGS OTHER THAN MARIJUANA¹	15,592	14,994	4,770	4,375	2,109	1,981

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.7B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	61.7	61.2	25.4	25.0	15.3	15.1
Marijuana and Hashish	56.9	55.7	21.0	20.5	12.6	12.7
Cocaine	18.9	17.9	3.4	3.0	1.3	1.0
Crack	4.7	4.5	0.4	0.4	0.2	0.2
Heroin	3.1	3.1	0.7	0.9	0.3	0.5
Hallucinogens	23.5	21.8	2.5	2.7	0.6	0.8
LSD	12.5 ^b	10.4	0.5	0.6	0.1	0.2
PCP	1.9	1.7	0.0	0.0	*	*
Ecstasy	17.1	16.0	1.7	1.6	0.5	0.5
Inhalants	12.3 ^a	10.9	0.6	0.9	0.2	0.4
Nonmedical Use of Psychotherapeutics ^{2,3}	30.7	29.3	10.3 ^a	8.8	4.4	4.1
Pain Relievers	23.4	21.9	7.1 ^a	6.0	3.0	2.6
OxyContin [®]	6.4	6.5	1.0	0.8	0.3	0.2
Tranquilizers	15.2	14.7	4.1	3.5	1.4	1.5
Stimulants ³	12.9	12.1	2.9	2.4	1.1	1.0
Methamphetamine ³	6.8	6.2	0.8	0.9	0.4	0.3
Sedatives	2.2	1.9	0.5	0.4	0.2	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA¹	41.7	39.6	12.8	11.6	5.6	5.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.8A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	78,781 ^a	81,137	15,355 ^b	17,771	9,186 ^b	11,241
Marijuana and Hashish	71,305 ^a	73,618	10,772 ^b	12,952	6,691 ^b	8,699
Cocaine	26,324 ^a	28,322	1,233 ^a	1,641	628	642
Crack	6,517	7,149	376	494	255	249
Heroin	3,004	2,909	157	272	66 ^a	165
Hallucinogens	24,145	24,984	693	598	324	225
LSD	17,651	18,370	79	64	31	25
PCP	5,362	5,388	22	16	22	*
Ecstasy	6,611	6,982	326	318	110	105
Inhalants	12,551	13,430	347	392	185	182
Nonmedical Use of Psychotherapeutics ^{2,3}	30,283	31,792	5,819	5,943	2,615	2,792
Pain Relievers	17,682	18,929	4,199	4,156	1,849	1,896
OxyContin [®]	2,421	2,518	433	364	249	157
Tranquilizers	13,065	14,251	1,831	1,998	659	802
Stimulants ³	13,129 ^a	14,297	841	1,075	397 ^a	659
Methamphetamine ³	8,565	9,397	500	523	294	331
Sedatives	6,112	6,442	261	370	105	198
ILLICIT DRUGS OTHER THAN MARIJUANA¹	47,353 ^a	49,943	7,265	7,488	3,462	3,605

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.8B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	47.6	48.5	9.3 ^b	10.6	5.6 ^b	6.7
Marijuana and Hashish	43.1	44.0	6.5 ^b	7.7	4.0 ^b	5.2
Cocaine	15.9 ^a	16.9	0.7 ^a	1.0	0.4	0.4
Crack	3.9	4.3	0.2	0.3	0.2	0.1
Heroin	1.8	1.7	0.1	0.2	0.0 ^a	0.1
Hallucinogens	14.6	14.9	0.4	0.4	0.2	0.1
LSD	10.7	11.0	0.0	0.0	0.0	0.0
PCP	3.2	3.2	0.0	0.0	0.0	*
Ecstasy	4.0	4.2	0.2	0.2	0.1	0.1
Inhalants	7.6	8.0	0.2	0.2	0.1	0.1
Nonmedical Use of Psychotherapeutics ^{2,3}	18.3	19.0	3.5	3.5	1.6	1.7
Pain Relievers	10.7	11.3	2.5	2.5	1.1	1.1
OxyContin [®]	1.5	1.5	0.3	0.2	0.2	0.1
Tranquilizers	7.9	8.5	1.1	1.2	0.4	0.5
Stimulants ³	7.9	8.5	0.5	0.6	0.2 ^a	0.4
Methamphetamine ³	5.2	5.6	0.3	0.3	0.2	0.2
Sedatives	3.7	3.8	0.2	0.2	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	28.6	29.8	4.4	4.5	2.1	2.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.9A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	67,412 ^a	69,492	23,694 ^b	25,483	14,671 ^b	16,423
Marijuana and Hashish	61,743 ^a	63,633	19,713 ^a	21,196	12,300 ^b	14,033
Cocaine	22,612	23,917	2,855	2,965	1,038	991
Crack	5,504	6,163	390	479	218	206
Heroin	3,110	3,451	448 ^a	622	173 ^a	284
Hallucinogens	23,835	24,029	2,809	2,756	902	748
LSD	15,592	15,950	726	863	156	195
PCP	4,116	4,306	52	43	7	10
Ecstasy	10,040	10,513	1,554	1,429	439	349
Inhalants	13,675	14,058	1,025	1,082	342	390
Nonmedical Use of Psychotherapeutics ^{2,3}	27,490 ^a	28,978	7,772	7,852	3,340	3,382
Pain Relievers	18,918 ^a	20,356	5,834	5,585	2,398	2,369
OxyContin [®]	4,207	4,517	948	716	337	222
Tranquilizers	11,942 ^a	12,971	2,602	2,636	833	970
Stimulants ³	11,872	12,606	1,831	2,040	742	830
Methamphetamine ³	7,366	8,013	793	765	432	299
Sedatives	4,176	4,590	286	391	121	183
ILLCIT DRUGS OTHER THAN MARIJUANA¹	42,406 ^a	44,070	10,870	10,919	4,855	4,778

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.9B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	53.0	54.1	18.6 ^a	19.8	11.5 ^b	12.8
Marijuana and Hashish	48.6	49.5	15.5 ^a	16.5	9.7 ^b	10.9
Cocaine	17.8	18.6	2.2	2.3	0.8	0.8
Crack	4.3	4.8	0.3	0.4	0.2	0.2
Heroin	2.4	2.7	0.4 ^a	0.5	0.1	0.2
Hallucinogens	18.7	18.7	2.2	2.1	0.7	0.6
LSD	12.3	12.4	0.6	0.7	0.1	0.2
PCP	3.2	3.4	0.0	0.0	0.0	0.0
Ecstasy	7.9	8.2	1.2	1.1	0.3	0.3
Inhalants	10.8	10.9	0.8	0.8	0.3	0.3
Nonmedical Use of Psychotherapeutics ^{2,3}	21.6	22.6	6.1	6.1	2.6	2.6
Pain Relievers	14.9 ^a	15.9	4.6	4.3	1.9	1.8
OxyContin [®]	3.3	3.5	0.7 ^a	0.6	0.3	0.2
Tranquilizers	9.4	10.1	2.0	2.1	0.7	0.8
Stimulants ³	9.3	9.8	1.4	1.6	0.6	0.6
Methamphetamine ³	5.8	6.2	0.6	0.6	0.3	0.2
Sedatives	3.3	3.6	0.2	0.3	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	33.4	34.3	8.6	8.5	3.8	3.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.10A Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	60,046	60,839	17,897	18,674	9,903	10,560
Marijuana and Hashish	52,969	53,580	13,239	13,927	7,510	8,155
Cocaine	15,022	15,283	1,327	1,588	511	539
Crack	3,366	3,261	242	294	160	149
Heroin	1,703	1,362	233	292	116	151
Hallucinogens	15,902	15,617	1,621	1,494	431	426
LSD	9,178	9,085	386	426	73	93
PCP	2,346	2,082	38	47	26	*
Ecstasy	7,774 ^a	7,035	1,034	913	221	260
Inhalants	7,393	7,235	508	535	153	155
Nonmedical Use of Psychotherapeutics ^{2,3}	25,682	25,417	7,576	7,114	3,144	3,155
Pain Relievers	16,555	15,709	5,248	4,752	2,123	1,956
OxyContin [®]	2,766	2,514	495	521	155	134
Tranquilizers	11,551	11,880	2,667	2,566	872	905
Stimulants ³	9,784	9,924	1,661	1,676	623	765
Methamphetamine ³	4,890	4,930	393	536	163	270
Sedatives	3,304	3,237	353	384	130	146
ILLCIT DRUGS OTHER THAN MARIJUANA¹	35,670	36,049	8,997	8,800	3,810	3,941

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.10B Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	44.4	44.5	13.2	13.7	7.3	7.7
Marijuana and Hashish	39.2	39.2	9.8	10.2	5.6	6.0
Cocaine	11.1	11.2	1.0	1.2	0.4	0.4
Crack	2.5	2.4	0.2	0.2	0.1	0.1
Heroin	1.3	1.0	0.2	0.2	0.1	0.1
Hallucinogens	11.8	11.4	1.2	1.1	0.3	0.3
LSD	6.8	6.6	0.3	0.3	0.1	0.1
PCP	1.7	1.5	0.0	0.0	0.0	*
Ecstasy	5.7 ^a	5.1	0.8	0.7	0.2	0.2
Inhalants	5.5	5.3	0.4	0.4	0.1	0.1
Nonmedical Use of Psychotherapeutics ^{2,3}	19.0	18.6	5.6	5.2	2.3	2.3
Pain Relievers	12.2	11.5	3.9	3.5	1.6	1.4
OxyContin [®]	2.0	1.8	0.4	0.4	0.1	0.1
Tranquilizers	8.5	8.7	2.0	1.9	0.6	0.7
Stimulants ³	7.2	7.3	1.2	1.2	0.5	0.6
Methamphetamine ³	3.6	3.6	0.3	0.4	0.1	0.2
Sedatives	2.4	2.4	0.3	0.3	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA¹	26.4	26.4	6.7	6.4	2.8	2.9

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.11A Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	127,458 ^a	130,332	41,591 ^b	44,157	24,573 ^b	26,983
12	299	320	191	203	87	99
13	473	502	275 ^a	354	119 ^a	167
14	802	712	567	530	291	242
15	1,060	1,125	776	837	373	423
16	1,439	1,412	1,127	1,109	646	641
17	1,721	1,719	1,350	1,303	680	766
18	2,258	2,080	1,765	1,604	993	963
19	2,120	2,229	1,591	1,674	948	971
20	2,441	2,485	1,705	1,665	1,036	1,039
21	2,524	2,607	1,646	1,734	1,022	1,016
22	2,633	2,837	1,558	1,728	980	1,068
23	2,708	2,702	1,455	1,478	847	998
24	2,646	2,659	1,460	1,362	921 ^a	772
25	2,496	2,619	1,268	1,352	742	849
26-29	10,631	10,406	4,808	4,759	2,743	2,875
30-34	12,425	12,781	4,692	4,693	2,957	2,853
35-39	10,149	10,955	2,826	3,268	1,780	2,056
40-44	11,565	10,697	2,855	2,926	1,649	1,937
45-49	11,631	11,815	2,578	2,679	1,454	1,724
50-54	14,530	14,420	2,877	3,037	1,812	1,885
55-59	11,704	12,194	1,757 ^b	2,817	1,165 ^b	1,860
60-64	8,861	9,447	1,150 ^a	1,597	692	947
65 or Older	10,340 ^a	11,608	1,311	1,446	634	832

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.11B Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	48.6	49.2	15.9 ^a	16.7	9.4 ^b	10.2
12	8.0	8.4	5.1	5.3	2.3	2.6
13	11.2	12.3	6.5 ^a	8.7	2.8 ^a	4.1
14	18.3	17.1	12.9	12.7	6.6	5.8
15	25.6	26.2	18.8	19.5	9.0	9.9
16	33.7	32.5	26.4	25.6	15.1	14.8
17	41.7	41.2	32.7	31.2	16.5	18.3
18	47.7	45.2	37.3	34.8	21.0	20.9
19	50.9	53.8	38.2	40.4	22.8	23.5
20	56.8	57.0	39.7	38.2	24.1	23.8
21	59.6	60.0	38.8	39.9	24.1	23.4
22	58.8 ^a	62.9	34.8	38.3	21.9	23.7
23	61.9	61.5	33.2	33.6	19.4 ^a	22.7
24	60.2	61.5	33.2	31.5	21.0 ^a	17.8
25	60.7	61.6	30.9	31.8	18.1	20.0
26-29	61.8	61.1	27.9	28.0	15.9	16.9
30-34	61.7	61.3	23.3	22.5	14.7	13.7
35-39	53.3	56.0	14.8	16.7	9.3	10.5
40-44	53.9	53.0	13.3	14.5	7.7 ^a	9.6
45-49	58.2	57.5	12.9	13.0	7.3	8.4
50-54	63.0	60.7	12.5	12.8	7.9	7.9
55-59	57.0	58.4	8.6 ^b	13.5	5.7 ^b	8.9
60-64	49.8	53.8	6.5 ^a	9.1	3.9	5.4
65 or Older	23.8	25.8	3.0	3.2	1.5	1.9

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.12A Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	114,712	117,213	32,952 ^b	35,124	19,810 ^b	22,188
12	72	51	59	46	21	20
13	168	165	138	140	60	69
14	445	390	371	328	196	156
15	765	778	604	628	296	312
16	1,182	1,201	974	957	568	572
17	1,460	1,507	1,200	1,163	621	702
18	2,004	1,801	1,566	1,432	897	872
19	1,906	2,009	1,436	1,524	873	895
20	2,226	2,257	1,523	1,540	915	969
21	2,338	2,387	1,486	1,547	919	891
22	2,420	2,578	1,350	1,494	860	917
23	2,448	2,461	1,244	1,290	734	876
24	2,435	2,476	1,285	1,161	788	691
25	2,274	2,419	1,111	1,168	650	726
26-29	9,816	9,430	3,984	3,990	2,363	2,444
30-34	11,447	11,684	3,849	3,766	2,358	2,378
35-39	9,075	9,808	1,997 ^b	2,512	1,332 ^a	1,711
40-44	10,049	9,534	2,008	2,049	1,208	1,468
45-49	10,658	10,783	1,740	2,073	1,044 ^a	1,352
50-54	13,787	13,331	2,215	2,008	1,272	1,251
55-59	10,965	11,510	1,347 ^b	2,272	910 ^b	1,578
60-64	8,264	8,818	842	1,166	548	763
65 or Older	8,508 ^a	9,835	623	871	377	575

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.12B Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	43.7	44.2	12.6 ^a	13.2	7.5 ^b	8.4
12	1.9	1.3	1.6	1.2	0.6	0.5
13	4.0	4.0	3.3	3.4	1.4	1.7
14	10.2	9.3	8.5	7.9	4.5	3.7
15	18.5	18.1	14.6	14.6	7.1	7.3
16	27.7	27.7	22.8	22.1	13.3	13.2
17	35.4	36.1	29.1	27.9	15.0	16.8
18	42.3	39.1	33.1	31.1	18.9	18.9
19	45.7	48.5	34.5	36.8	20.9	21.6
20	51.8	51.7	35.5	35.3	21.3	22.2
21	55.2	54.9	35.1	35.6	21.7	20.5
22	54.1	57.1	30.2	33.1	19.2	20.3
23	55.9	56.0	28.4	29.3	16.8 ^a	19.9
24	55.4	57.3	29.3	26.8	18.0	16.0
25	55.3	56.9	27.0	27.5	15.8	17.1
26-29	57.0	55.4	23.1	23.4	13.7	14.4
30-34	56.9	56.0	19.1	18.1	11.7	11.4
35-39	47.6	50.1	10.5 ^a	12.8	7.0 ^a	8.7
40-44	46.8	47.3	9.4	10.2	5.6 ^a	7.3
45-49	53.3	52.5	8.7	10.1	5.2 ^a	6.6
50-54	59.7	56.1	9.6	8.5	5.5	5.3
55-59	53.4	55.2	6.6 ^b	10.9	4.4 ^b	7.6
60-64	46.5	50.3	4.7 ^a	6.6	3.1	4.4
65 or Older	19.6	21.9	1.4	1.9	0.9	1.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.13A Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	37,634	39,200	4,182	4,553	1,549	1,530
12	3	7	3	*	*	*
13	6	3	4	1	2	*
14	10	10	8	9	8	3
15	27	32	18	21	4	8
16	68	78	37	65	10	14
17	101	93	62	71	20	15
18	186	161	115	105	33	24
19	283	245	159	120	46	44
20	405	346	176	194	45	80
21	452	531	208	237	47	80
22	649	582	262	293	79	71
23	680	636	228	261	40	64
24	727	629	227	162	56	37
25	658	750	169	231	52	72
26-29	3,158	2,892	615	628	189	215
30-34	3,895	3,884	660	512	292	161
35-39	2,707 ^a	3,275	319	394	132	164
40-44	3,632	3,390	167 ^a	303	68	105
45-49	4,649	4,651	216	261	103	119
50-54	6,849	6,689	294	352	181	194
55-59	4,125 ^a	5,093	113	193	48	40
60-64	2,669	3,042	99	59	73	17
65 or Older	1,693	2,181	26	79	22	3

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.13B Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	14.3	14.8	1.6	1.7	0.6	0.6
12	0.1	0.2	0.1	*	*	*
13	0.1	0.1	0.1	0.0	0.0	*
14	0.2	0.2	0.2	0.2	0.2	0.1
15	0.6	0.7	0.4	0.5	0.1	0.2
16	1.6	1.8	0.9	1.5	0.2	0.3
17	2.5	2.2	1.5	1.7	0.5	0.4
18	3.9	3.5	2.4	2.3	0.7	0.5
19	6.8	5.9	3.8	2.9	1.1	1.1
20	9.4	7.9	4.1	4.4	1.0	1.8
21	10.7	12.2	4.9	5.4	1.1	1.8
22	14.5	12.9	5.8	6.5	1.8	1.6
23	15.5	14.5	5.2	6.0	0.9	1.5
24	16.5	14.5	5.2	3.7	1.3	0.9
25	16.0	17.7	4.1	5.4	1.3	1.7
26-29	18.4	17.0	3.6	3.7	1.1	1.3
30-34	19.3	18.6	3.3	2.5	1.4	0.8
35-39	14.2 ^a	16.7	1.7	2.0	0.7	0.8
40-44	16.9	16.8	0.8 ^a	1.5	0.3	0.5
45-49	23.3	22.7	1.1	1.3	0.5	0.6
50-54	29.7	28.2	1.3	1.5	0.8	0.8
55-59	20.1 ^b	24.4	0.5	0.9	0.2	0.2
60-64	15.0	17.3	0.6	0.3	0.4	0.1
65 or Older	3.9	4.8	0.1	0.2	0.1	0.0

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.14A Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8,870	9,424	632	773	377	354
12	*	*	*	*	*	*
13	2	1	2	*	2	*
14	1	1	*	1	*	*
15	5	8	2	4	*	1
16	6	8	3	2	*	1
17	9	17	4	14	1	6
18	29	12	8	8	2	1
19	27	33	10	12	2	3
20	40	37	6	20	*	1
21	49	64	11	18	6	12
22	54	84	11	18	5	2
23	111	109	10	16	1	0
24	119	91	9	15	3	5
25	134	116	30	19	17	5
26-29	809	644	49	85	17	49
30-34	957	1,049	101	48	67	20
35-39	784	1,008	35	69	20	35
40-44	939	1,084	44	81	37	29
45-49	1,217	1,349	40 ^a	116	21 ^a	66
50-54	1,760	1,578	121	137	103	85
55-59	1,128	1,223	81	87	24	30
60-64	386	590	29	3	29	3
65 or Older	303	318	26	*	21	*

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.14B Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	3.4	3.6	0.2	0.3	0.1	0.1
12	*	*	*	*	*	*
13	0.0	0.0	0.0	*	0.0	*
14	0.0	0.0	*	0.0	*	*
15	0.1	0.2	0.1	0.1	*	0.0
16	0.1	0.2	0.1	0.0	*	0.0
17	0.2	0.4	0.1	0.3	0.0	0.1
18	0.6	0.3	0.2	0.2	0.0	0.0
19	0.7	0.8	0.2	0.3	0.1	0.1
20	0.9	0.9	0.1	0.5	*	0.0
21	1.2	1.5	0.3	0.4	0.2	0.3
22	1.2	1.9	0.3	0.4	0.1	0.0
23	2.5	2.5	0.2	0.4	0.0	0.0
24	2.7	2.1	0.2	0.4	0.1	0.1
25	3.3	2.7	0.7	0.4	0.4	0.1
26-29	4.7	3.8	0.3	0.5	0.1	0.3
30-34	4.8	5.0	0.5	0.2	0.3	0.1
35-39	4.1	5.2	0.2	0.4	0.1	0.2
40-44	4.4	5.4	0.2	0.4	0.2	0.1
45-49	6.1	6.6	0.2 ^a	0.6	0.1 ^a	0.3
50-54	7.6	6.6	0.5	0.6	0.4	0.4
55-59	5.5	5.9	0.4	0.4	0.1	0.1
60-64	2.2	3.4	0.2	0.0	0.2	0.0
65 or Older	0.7	0.7	0.1	*	0.0	*

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.15A Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	39,736	39,647	4,430	4,250	1,333	1,173
12	25	24	16	10	3	6
13	34	23	19	16	8	4
14	54	42	38	25	18	10
15	97	116	61	88	19	20
16	200	194	140	141	32	48
17	293 ^b	211	208 ^a	142	74	49
18	486	436	305	275	123 ^b	50
19	564	486	309	227	93	50
20	687	623	313	332	65	72
21	785	759	357	328	65	77
22	908	914	333	359	88	90
23	936	833	303	252	62	64
24	975 ^a	815	251	187	79	53
25	783	919	167	244	42	47
26-29	3,863	3,431	473	579	147	192
30-34	4,903	4,837	444	447	90	118
35-39	3,741	4,023	218	248	117	48
40-44	3,778	3,423	153	173	21	62
45-49	3,395	3,727	57	68	12	48
50-54	5,215	4,863	87	65	77	50
55-59	3,804	4,080	8	11	*	9
60-64	2,542	2,812	13	13	*	*
65 or Older	1,671	2,057	157	20	97	8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.15B Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	15.1	15.0	1.7	1.6	0.5	0.4
12	0.7	0.6	0.4	0.3	0.1	0.1
13	0.8	0.6	0.5	0.4	0.2	0.1
14	1.2	1.0	0.9	0.6	0.4	0.2
15	2.4	2.7	1.5	2.1	0.5	0.5
16	4.7	4.5	3.3	3.2	0.7	1.1
17	7.1 ^b	5.0	5.0 ^a	3.4	1.8	1.2
18	10.3	9.5	6.4	6.0	2.6 ^b	1.1
19	13.5	11.7	7.4 ^a	5.5	2.2	1.2
20	16.0	14.3	7.3	7.6	1.5	1.7
21	18.5	17.5	8.4	7.6	1.5	1.8
22	20.3	20.3	7.4	8.0	2.0	2.0
23	21.4	19.0	6.9	5.7	1.4	1.4
24	22.2 ^a	18.8	5.7	4.3	1.8	1.2
25	19.1	21.6	4.1	5.8	1.0	1.1
26-29	22.4	20.2	2.7	3.4	0.9	1.1
30-34	24.4	23.2	2.2	2.1	0.4	0.6
35-39	19.6	20.6	1.1	1.3	0.6	0.2
40-44	17.6	17.0	0.7	0.9	0.1	0.3
45-49	17.0	18.1	0.3	0.3	0.1	0.2
50-54	22.6	20.5	0.4	0.3	0.3	0.2
55-59	18.5	19.6	0.0	0.1	*	0.0
60-64	14.3	16.0	0.1	0.1	*	*
65 or Older	3.8	4.6	0.4	0.0	0.2	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.16A Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	21,068	21,293	1,533	1,617	496	546
12	139	140	71	58	27	31
13	197	202	74	104	20	24
14	265	238	90	119	18	40
15	287	302	107	120	31	41
16	215	228	63	71	12	10
17	224	195	76 ^a	41	13 ^a	3
18	280	240	74	72	16	21
19	277	221	66	40	15	2
20	309	259	78	47	20	11
21	341	337	84 ^a	41	16 ^a	4
22	376	345	54	40	17	7
23	367	366	48	60	6	16
24	386	300	45	36	7	5
25	277 ^a	375	30	42	10	14
26-29	1,809	1,605	113	127	22	45
30-34	2,768	2,507	115	207	62	90
35-39	2,108	2,422	43	69	19	9
40-44	2,299	2,145	77	123	51	80
45-49	2,167	2,106	74	43	55	30
50-54	2,888	2,613	50	52	31	35
55-59	1,391 ^a	1,964	93	69	19	26
60-64	971	1,206	7	23	7	1
65 or Older	727	975	3	13	3	*

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.16B Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8.0	8.0	0.6	0.6	0.2	0.2
12	3.7	3.7	1.9	1.5	0.7	0.8
13	4.7	5.0	1.7	2.6	0.5	0.6
14	6.0	5.7	2.0	2.8	0.4 ^a	1.0
15	6.9	7.0	2.6	2.8	0.8	1.0
16	5.0	5.3	1.5	1.6	0.3	0.2
17	5.4	4.7	1.8 ^a	1.0	0.3 ^a	0.1
18	5.9	5.2	1.6	1.6	0.3	0.4
19	6.7	5.3	1.6	1.0	0.4	0.0
20	7.2	5.9	1.8	1.1	0.5	0.3
21	8.0	7.8	2.0 ^a	0.9	0.4 ^a	0.1
22	8.4	7.6	1.2	0.9	0.4	0.2
23	8.4	8.3	1.1	1.4	0.1	0.4
24	8.8	6.9	1.0	0.8	0.2	0.1
25	6.7 ^a	8.8	0.7	1.0	0.2	0.3
26-29	10.5	9.4	0.7	0.7	0.1	0.3
30-34	13.7	12.0	0.6	1.0	0.3	0.4
35-39	11.1	12.4	0.2	0.4	0.1	0.0
40-44	10.7	10.6	0.4	0.6	0.2	0.4
45-49	10.8	10.3	0.4	0.2	0.3	0.1
50-54	12.5	11.0	0.2	0.2	0.1	0.1
55-59	6.8 ^a	9.4	0.5	0.3	0.1	0.1
60-64	5.5	6.9	0.0	0.1	0.0	0.0
65 or Older	1.7	2.2	0.0	0.0	0.0	*

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.17A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	53,172	54,395	15,348	14,966	6,484	6,537
12	130	167	88	114	50	55
13	199	255	99 ^b	170	50 ^a	89
14	338 ^a	262	229	173	96	81
15	356	419	248	310	89 ^a	136
16	549	541	371	376	140	159
17	615	646	406	412	124	134
18	904	900	527	509	182	188
19	871	820	490	439	165	113
20	1,117	998	584	480	236	181
21	1,178	1,227	589	619	227	229
22	1,289	1,343	603	681	234	290
23	1,340	1,321	535	514	243	212
24	1,324	1,253	507	411	215 ^a	139
25	1,216	1,334	409	478	159	200
26-29	5,448	5,085	1,877	1,642	745	816
30-34	6,014	6,032	1,966	1,695	913	723
35-39	4,593	4,870	1,185	1,264	493	533
40-44	4,765	4,477	1,113	1,044	590	473
45-49	4,319	4,459	1,026	834	419	449
50-54	6,168	6,023	967	1,029	482	506
55-59	3,999 ^a	4,916	542	715	247	355
60-64	2,792	3,339	324	513	95	201
65 or Older	3,648	3,707	662	544	290	276

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.17B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	20.3	20.5	5.8	5.6	2.5	2.5
12	3.5	4.4	2.3	3.0	1.3	1.4
13	4.7 ^a	6.2	2.3 ^b	4.2	1.2 ^a	2.2
14	7.7	6.3	5.2	4.1	2.2	2.0
15	8.6	9.8	6.0	7.2	2.1 ^a	3.2
16	12.8	12.5	8.7	8.7	3.3	3.7
17	14.9	15.5	9.8	9.9	3.0	3.2
18	19.1	19.5	11.1	11.1	3.9	4.1
19	20.9	19.8	11.8	10.6	4.0	2.7
20	26.0	22.9	13.6 ^a	11.0	5.5	4.2
21	27.8	28.2	13.9	14.3	5.4	5.3
22	28.8	29.8	13.5	15.1	5.2	6.4
23	30.6	30.1	12.2	11.7	5.5	4.8
24	30.1	29.0	11.6	9.5	4.9 ^a	3.2
25	29.6	31.4	9.9	11.3	3.9	4.7
26-29	31.7	29.9	10.9	9.6	4.3	4.8
30-34	29.9	28.9	9.8	8.1	4.5	3.5
35-39	24.1	24.9	6.2	6.5	2.6	2.7
40-44	22.2	22.2	5.2	5.2	2.7	2.3
45-49	21.6	21.7	5.1	4.1	2.1	2.2
50-54	26.7	25.4	4.2	4.3	2.1	2.1
55-59	19.5 ^a	23.6	2.6	3.4	1.2	1.7
60-64	15.7 ^a	19.0	1.8	2.9	0.5	1.1
65 or Older	8.4	8.2	1.5	1.2	0.7	0.6

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.18A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	35,473	36,064	11,082	10,337	4,521	4,325
12	99	119	64	78	34	37
13	168	197	86 ^a	131	42	66
14	294 ^a	221	198 ^a	136	88	62
15	296	345	204	252	70 ^a	112
16	450	420	297	268	106	107
17	499	500	308	300	87	85
18	705	660	392	358	116	127
19	702	618	362	319	100	74
20	895	768	425 ^a	312	174 ^b	93
21	926	971	407	407	144	151
22	1,039	1,000	440	434	155	187
23	1,066	1,052	377	341	183 ^a	120
24	1,014	949	357	271	148	95
25	904	1,031	300	297	113	132
26-29	4,092	3,842	1,266	1,087	484	495
30-34	4,643	4,440	1,400	1,189	631	488
35-39	3,291	3,463	854	887	344	349
40-44	3,373	3,108	847	743	414	308
45-49	2,660	2,700	708 ^a	515	306	310
50-54	3,185	3,476	769	828	318	419
55-59	2,043 ^b	3,004	392	525	133	245
60-64	1,445	1,502	196	309	69	113
65 or Older	1,687	1,675	432	350	264	153

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.18B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	13.5	13.6	4.2 ^a	3.9	1.7	1.6
12	2.6	3.1	1.7	2.0	0.9	1.0
13	4.0	4.8	2.0 ^a	3.2	1.0	1.6
14	6.7	5.3	4.5 ^a	3.3	2.0	1.5
15	7.2	8.1	4.9	5.9	1.7 ^a	2.6
16	10.5	9.7	6.9	6.2	2.5	2.5
17	12.1	12.0	7.5	7.2	2.1	2.0
18	14.9	14.3	8.3	7.8	2.4	2.7
19	16.8	14.9	8.7	7.7	2.4	1.8
20	20.8 ^a	17.6	9.9 ^a	7.2	4.0 ^b	2.1
21	21.8	22.3	9.6	9.4	3.4	3.5
22	23.2	22.2	9.8	9.6	3.5	4.1
23	24.3	23.9	8.6	7.8	4.2 ^a	2.7
24	23.1	21.9	8.1	6.3	3.4	2.2
25	22.0	24.3	7.3	7.0	2.7	3.1
26-29	23.8	22.6	7.4	6.4	2.8	2.9
30-34	23.1	21.3	7.0	5.7	3.1	2.3
35-39	17.3	17.7	4.5	4.5	1.8	1.8
40-44	15.7	15.4	3.9	3.7	1.9	1.5
45-49	13.3	13.1	3.5 ^a	2.5	1.5	1.5
50-54	13.8	14.6	3.3	3.5	1.4	1.8
55-59	9.9 ^b	14.4	1.9	2.5	0.6	1.2
60-64	8.1	8.6	1.1	1.8	0.4	0.6
65 or Older	3.9	3.7	1.0	0.8	0.6	0.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.19A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	127,458 ^a	130,332	41,591 ^b	44,157	24,573 ^b	26,983
AGE						
12-17	5,794	5,791	4,287	4,336	2,197	2,338
18-25	19,826	20,217	12,448	12,597	7,490	7,676
26 or Older	101,838	104,324	24,856 ^b	27,223	14,886 ^b	16,969
GENDER						
Male	67,412 ^a	69,492	23,694 ^b	25,483	14,671 ^b	16,423
Female	60,046	60,839	17,897	18,674	9,903	10,560
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	111,317 ^a	113,792	35,368 ^b	37,552	20,930 ^b	23,191
White	89,899	91,533	27,450	28,691	16,204 ^a	17,661
Black or African American	14,784	15,079	5,240 ^b	6,164	3,274 ^b	3,928
American Indian or Alaska Native	853	877	375	346	173	215
Native Hawaiian or Other Pacific Islander	420	555	181	217	122	159
Asian	2,860	3,188	942	1,102	410	564
Two or More Races	2,500	2,559	1,179	1,031	748	663
Hispanic or Latino	16,141	16,540	6,222	6,605	3,643	3,792

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.19B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	48.6	49.2	15.9 ^a	16.7	9.4 ^b	10.2
AGE						
12-17	23.3	23.3	17.2	17.4	8.8	9.4
18-25	57.0	57.9	35.8	36.1	21.5	22.0
26 or Older	50.2	50.8	12.3 ^a	13.3	7.3 ^b	8.3
GENDER						
Male	53.0	54.1	18.6 ^a	19.8	11.5 ^b	12.8
Female	44.4	44.5	13.2	13.7	7.3	7.7
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	50.3	51.1	16.0 ^a	16.9	9.5 ^b	10.4
White	52.9	53.8	16.1	16.9	9.5 ^a	10.4
Black or African American	47.3	47.6	16.8 ^b	19.5	10.5 ^a	12.4
American Indian or Alaska Native	60.9	60.8	26.8	24.0	12.3	14.9
Native Hawaiian or Other Pacific Islander	48.3	54.4	20.8	21.3	14.0	15.6
Asian	21.4	23.0	7.1	8.0	3.1	4.1
Two or More Races	58.3	57.9	27.5	23.3	17.4	15.0
Hispanic or Latino	39.1	38.9	15.1	15.6	8.8	8.9

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.20A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	5,794	5,791	4,287	4,336	2,197	2,338
GENDER						
Male	3,056	2,898	2,263	2,112	1,220	1,222
Female	2,738	2,893	2,023 ^a	2,224	977 ^a	1,116
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4,476	4,367	3,329	3,267	1,715	1,746
White	3,031	2,959	2,315	2,288	1,196	1,260
Black or African American	963	965	691	667	361	342
American Indian or Alaska Native	62	48	44	39	*	18
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	190	163	113	112	47	43
Two or More Races	193	206	143	148	78	77
Hispanic or Latino	1,318	1,424	958	1,069	482 ^a	591
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	1,595	1,515	1,215	1,152	660	689
Female, White, Not Hispanic	1,437	1,444	1,100	1,135	536	571
Male, Black, Not Hispanic	498	466	352	313	209	173
Female, Black, Not Hispanic	465	498	339	354	151	169
Male, Hispanic	710	684	536	486	267	282
Female, Hispanic	608 ^a	740	422 ^b	583	215 ^a	309

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.20B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	23.3	23.3	17.2	17.4	8.8	9.4
GENDER						
Male	24.1	22.8	17.8	16.7	9.6	9.6
Female	22.5	23.7	16.6 ^a	18.2	8.0 ^a	9.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	23.2	22.7	17.2	17.0	8.9	9.1
White	22.2	21.9	17.0	17.0	8.8	9.3
Black or African American	28.0	27.7	20.1	19.2	10.5	9.8
American Indian or Alaska Native	38.7	29.0	27.8	23.2	*	10.6
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	15.2	13.3	9.1	9.1	3.8	3.5
Two or More Races	25.7	26.8	19.0	19.3	10.3	10.0
Hispanic or Latino	23.7	25.2	17.2	18.9	8.7 ^a	10.5
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	22.9	22.0	17.5	16.7	9.5	10.0
Female, White, Not Hispanic	21.5	21.9	16.5	17.2	8.0	8.7
Male, Black, Not Hispanic	28.4	26.4	20.1	17.7	11.9	9.8
Female, Black, Not Hispanic	27.5	29.0	20.1	20.6	9.0	9.8
Male, Hispanic	25.0	23.8	18.9	16.9	9.4	9.8
Female, Hispanic	22.3 ^a	26.7	15.5 ^b	21.0	7.9 ^a	11.1

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.21A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	19,826	20,217	12,448	12,597	7,490	7,676
GENDER						
Male	10,571	10,639	7,097	7,002	4,522	4,449
Female	9,255	9,578	5,351	5,596	2,969	3,228
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	16,003	16,365	10,194	10,241	6,171	6,338
White	11,897	12,105	7,577	7,422	4,650	4,569
Black or African American	2,682	2,836	1,719	1,907	1,047	1,210
American Indian or Alaska Native	150	138	87	82	48	42
Native Hawaiian or Other Pacific Islander	*	*	*	*	38	*
Asian	673	672	419	418	193	238
Two or More Races	490	504	335	330	195	220
Hispanic or Latino	3,824	3,852	2,254	2,356	1,319	1,338
EDUCATION						
< High School	3,168 ^a	2,826	2,119 ^a	1,860	1,407 ^a	1,192
High School Graduate	6,615	6,674	4,179	4,240	2,496	2,687
Some College	7,065	7,469	4,454	4,727	2,713	2,846
College Graduate	2,978	3,248	1,696	1,770	874	951
CURRENT EMPLOYMENT						
Full-Time	8,283 ^a	8,960	4,713 ^a	5,164	2,769 ^b	3,240
Part-Time	5,439	5,433	3,726	3,667	2,262	2,172
Unemployed	2,564	2,319	1,832 ^a	1,588	1,147	1,053
Other ¹	3,541	3,505	2,177	2,178	1,313	1,212

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.21B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	57.0	57.9	35.8	36.1	21.5	22.0
GENDER						
Male	60.5	60.7	40.6	39.9	25.9	25.4
Female	53.4	55.0	30.9	32.2	17.1	18.5
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	58.1	59.4	37.0	37.1	22.4	23.0
White	61.0	62.5	38.9	38.3	23.9	23.6
Black or African American	53.6	56.3	34.3 ^a	37.9	20.9 ^a	24.0
American Indian or Alaska Native	65.3	68.2	37.7	40.2	20.9	20.7
Native Hawaiian or Other Pacific Islander	*	*	*	*	19.0	*
Asian	36.0	33.4	22.4	20.8	10.3	11.8
Two or More Races	66.9	64.3	45.7	42.1	26.6	28.1
Hispanic or Latino	52.7	52.3	31.1	32.0	18.2	18.2
EDUCATION						
< High School	56.7	57.1	37.9	37.6	25.2	24.1
High School Graduate	55.3	55.7	34.9	35.4	20.9	22.4
Some College	58.0	60.0	36.6	38.0	22.3	22.9
College Graduate	59.0	58.3	33.6	31.8	17.3	17.1
CURRENT EMPLOYMENT						
Full-Time	61.3	62.9	34.9	36.2	20.5 ^a	22.7
Part-Time	56.3	57.3	38.5	38.7	23.4	22.9
Unemployed	61.7	59.6	44.1	40.8	27.6	27.1
Other ¹	47.5	47.9	29.2	29.8	17.6	16.6

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.22A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	101,838	104,324	24,856 ^b	27,223	14,886 ^b	16,969
GENDER						
Male	53,784 ^a	55,956	14,333 ^b	16,370	8,929 ^b	10,752
Female	48,053	48,368	10,522	10,854	5,957	6,217
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	90,838	93,060	21,845 ^b	24,044	13,044 ^b	15,107
White	74,971	76,469	17,559	18,982	10,357 ^b	11,832
Black or African American	11,139	11,279	2,830 ^b	3,590	1,866 ^a	2,376
American Indian or Alaska Native	641	691	244	226	106	156
Native Hawaiian or Other Pacific Islander	*	*	*	122	*	*
Asian	1,998	2,353	410	572	169	283
Two or More Races	1,817	1,849	702	553	475	366
Hispanic or Latino	10,999	11,264	3,010	3,179	1,842	1,863
EDUCATION						
< High School	9,790	10,761	3,334	3,787	2,392	2,429
High School Graduate	27,323	27,682	6,721	7,047	4,411	4,835
Some College	28,846	30,039	6,767 ^b	8,117	4,120 ^b	5,207
College Graduate	35,879	35,842	8,033	8,273	3,963	4,499
CURRENT EMPLOYMENT						
Full-Time	60,321	62,383	13,824 ^b	16,075	8,030 ^b	9,781
Part-Time	13,353	12,449	3,719	3,536	2,352	2,187
Unemployed	4,529	4,229	1,782	1,532	1,002	967
Other ¹	23,635	25,262	5,530	6,080	3,502	4,034

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.22B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	50.2	50.8	12.3 ^a	13.3	7.3 ^b	8.3
GENDER						
Male	55.5	57.0	14.8 ^b	16.7	9.2 ^b	10.9
Female	45.4	45.2	9.9	10.1	5.6	5.8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	52.1	52.9	12.5 ^b	13.7	7.5 ^b	8.6
White	54.8	55.7	12.8 ^a	13.8	7.6 ^b	8.6
Black or African American	48.9	48.7	12.4 ^b	15.5	8.2 ^a	10.3
American Indian or Alaska Native	63.4	64.4	24.1	21.0	10.5	14.5
Native Hawaiian or Other Pacific Islander	*	*	*	16.4	*	*
Asian	19.5	22.2	4.0	5.4	1.7	2.7
Two or More Races	64.8	64.5	25.0	19.3	17.0	12.8
Hispanic or Latino	38.7	38.2	10.6	10.8	6.5	6.3
EDUCATION						
< High School	36.8	39.4	12.5	13.9	9.0	8.9
High School Graduate	47.1	48.6	11.6	12.4	7.6	8.5
Some College	56.3	56.5	13.2 ^b	15.3	8.0 ^b	9.8
College Graduate	53.6	52.7	12.0	12.2	5.9	6.6
CURRENT EMPLOYMENT						
Full-Time	57.2	57.4	13.1 ^b	14.8	7.6 ^b	9.0
Part-Time	55.4	52.9	15.4	15.0	9.8	9.3
Unemployed	59.2	61.3	23.3	22.2	13.1	14.0
Other ¹	36.1	38.2	8.4	9.2	5.3	6.1

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.23A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	121,664 ^a	124,541	37,304 ^b	39,821	22,376 ^b	24,646
GENDER						
Male	64,356 ^a	66,595	21,431 ^b	23,371	13,451 ^b	15,201
Female	57,308	57,946	15,873	16,449	8,926	9,445
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	106,841 ^a	109,425	32,039 ^b	34,285	19,215 ^b	21,445
White	86,868	88,574	25,136	26,403	15,008 ^a	16,401
Black or African American	13,821	14,114	4,549 ^b	5,497	2,913 ^b	3,586
American Indian or Alaska Native	791	829	331	307	154	198
Native Hawaiian or Other Pacific Islander	383	529	158	203	*	152
Asian	2,670	3,025	829	990	362 ^a	521
Two or More Races	2,307	2,354	1,036	883	670	587
Hispanic or Latino	14,823	15,116	5,264	5,536	3,161	3,201
EDUCATION						
< High School	12,958	13,587	5,453	5,647	3,799	3,620
High School Graduate	33,938	34,356	10,900	11,288	6,907	7,522
Some College	35,911	37,508	11,221 ^b	12,844	6,833 ^b	8,053
College Graduate	38,856	39,090	9,729	10,042	4,837	5,450
CURRENT EMPLOYMENT						
Full-Time	68,604	71,344	18,537 ^b	21,239	10,799 ^b	13,021
Part-Time	18,792	17,881	7,446	7,203	4,613	4,359
Unemployed	7,093	6,548	3,614 ^a	3,121	2,149	2,020
Other ¹	27,176	28,768	7,707	8,258	4,815	5,246

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.23B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	51.2	51.8	15.7 ^a	16.6	9.4 ^b	10.3
GENDER						
Male	56.2	57.5	18.7 ^b	20.2	11.8 ^b	13.1
Female	46.6	46.5	12.9	13.2	7.3	7.6
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	52.9	53.8	15.9 ^a	16.9	9.5 ^b	10.5
White	55.6	56.5	16.1	16.8	9.6 ^a	10.5
Black or African American	49.7	50.1	16.4 ^b	19.5	10.5 ^b	12.7
American Indian or Alaska Native	63.7	65.0	26.7	24.1	12.4	15.5
Native Hawaiian or Other Pacific Islander	49.6	57.6	20.5	22.2	*	16.6
Asian	22.1	24.0	6.8	7.8	3.0	4.1
Two or More Races	65.2	64.5	29.3	24.2	19.0	16.1
Hispanic or Latino	41.5	41.0	14.7	15.0	8.9	8.7
EDUCATION						
< High School	40.3	42.2	17.0	17.5	11.8	11.2
High School Graduate	48.5	49.9	15.6	16.4	9.9 ^a	10.9
Some College	56.6	57.2	17.7 ^b	19.6	10.8 ^b	12.3
College Graduate	54.0	53.2	13.5	13.7	6.7	7.4
CURRENT EMPLOYMENT						
Full-Time	57.7	58.0	15.6 ^b	17.3	9.1 ^b	10.6
Part-Time	55.7	54.2	22.1	21.8	13.7	13.2
Unemployed	60.1	60.7	30.6	28.9	18.2	18.7
Other ¹	37.2	39.1	10.6	11.2	6.6	7.1

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.24A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	114,712	117,213	32,952 ^b	35,124	19,810 ^b	22,188
AGE						
12-17	4,093	4,091	3,346	3,262	1,762	1,830
18-25	18,051	18,390	11,000	11,154	6,636	6,836
26 or Older	92,568	94,732	18,606 ^b	20,707	11,412 ^b	13,521
GENDER						
Male	61,743 ^a	63,633	19,713 ^a	21,196	12,300 ^b	14,033
Female	52,969	53,580	13,239	13,927	7,510	8,155
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	101,254	103,532	28,538 ^a	30,316	17,147 ^b	19,343
White	82,425	83,907	21,984 ^a	23,335	13,168 ^b	14,844
Black or African American	13,241	13,511	4,433 ^a	4,978	2,727 ^b	3,275
American Indian or Alaska Native	722	777	275	261	151	171
Native Hawaiian or Other Pacific Islander	398	505	167	167	117	123
Asian	2,238	2,452	695	729	300	384
Two or More Races	2,229	2,380	985	846	684	546
Hispanic or Latino	13,459	13,681	4,414	4,807	2,663	2,845

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.24B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	43.7	44.2	12.6 ^a	13.2	7.5 ^b	8.4
AGE						
12-17	16.4	16.4	13.4	13.1	7.1	7.4
18-25	51.9	52.6	31.6	31.9	19.1	19.6
26 or Older	45.7	46.1	9.2 ^b	10.1	5.6 ^b	6.6
GENDER						
Male	48.6	49.5	15.5 ^a	16.5	9.7 ^b	10.9
Female	39.2	39.2	9.8	10.2	5.6	6.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	45.8	46.5	12.9 ^a	13.6	7.8 ^b	8.7
White	48.5	49.3	12.9 ^a	13.7	7.7 ^b	8.7
Black or African American	42.4	42.6	14.2	15.7	8.7 ^a	10.3
American Indian or Alaska Native	51.5	53.9	19.6	18.1	10.8	11.8
Native Hawaiian or Other Pacific Islander	45.7	49.5	19.2	16.3	13.4	12.1
Asian	16.8	17.7	5.2	5.3	2.2	2.8
Two or More Races	52.0	53.9	23.0	19.1	16.0	12.4
Hispanic or Latino	32.6	32.2	10.7	11.3	6.5	6.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.25A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	4,093	4,091	3,346	3,262	1,762	1,830
GENDER						
Male	2,188	2,117	1,793	1,675	1,004	1,000
Female	1,905	1,975	1,554	1,587	758	831
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	3,141	3,074	2,596	2,468	1,361	1,382
White	2,192	2,172	1,836	1,773	971	1,040
Black or African American	640	622	506	474	255	240
American Indian or Alaska Native	44	39	32	33	15	16
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	103	94	88	76	45	30
Two or More Races	134	128	117	100	65	53
Hispanic or Latino	953	1,017	751	794	401	448
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	1,169	1,147	986	935	546	577
Female, White, Not Hispanic	1,023	1,025	851	838	425	463
Male, Black, Not Hispanic	344	330	273	253	157	144
Female, Black, Not Hispanic	295	292	232	221	97	96
Male, Hispanic	524	493	417	374	230	219
Female, Hispanic	429	524	334	420	171	229

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.25B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	16.4	16.4	13.4	13.1	7.1	7.4
GENDER						
Male	17.2	16.7	14.1	13.2	7.9	7.9
Female	15.6	16.2	12.8	13.0	6.2	6.8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	16.2	16.0	13.4	12.8	7.0	7.2
White	16.1	16.1	13.5	13.2	7.1	7.7
Black or African American	18.6	17.9	14.7	13.6	7.4	6.9
American Indian or Alaska Native	27.8	23.7	20.1	20.2	9.1	9.4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	8.3	7.7	7.1	6.2	3.6	2.5
Two or More Races	17.8	16.6	15.5	13.0	8.6	6.9
Hispanic or Latino	17.1	18.0	13.5	14.1	7.2	7.9
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	16.8	16.6	14.2	13.6	7.8	8.4
Female, White, Not Hispanic	15.3	15.5	12.7	12.7	6.4	7.0
Male, Black, Not Hispanic	19.6	18.7	15.6	14.3	9.0	8.1
Female, Black, Not Hispanic	17.5	17.0	13.8	12.9	5.8	5.6
Male, Hispanic	18.4	17.2	14.7	13.0	8.1	7.6
Female, Hispanic	15.8	18.9	12.3	15.2	6.3	8.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.26A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	18,051	18,390	11,000	11,154	6,636	6,836
GENDER						
Male	9,790	9,791	6,422	6,343	4,111	4,064
Female	8,260 ^a	8,599	4,577	4,812	2,525	2,772
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	14,611	14,989	9,076	9,125	5,472	5,689
White	10,953	11,158	6,737	6,646	4,142	4,069
Black or African American	2,436	2,589	1,575	1,728	936 ^a	1,151
American Indian or Alaska Native	129	133	72	69	36	33
Native Hawaiian or Other Pacific Islander	*	*	*	*	37	*
Asian	542	539	347	321	144	183
Two or More Races	447	477	289	296	177	203
Hispanic or Latino	3,439	3,400	1,924	2,029	1,164	1,147
EDUCATION						
< High School	2,844 ^a	2,502	1,887 ^a	1,662	1,269 ^a	1,083
High School Graduate	5,973	6,059	3,656	3,741	2,209	2,418
Some College	6,504	6,860	3,964	4,239	2,426	2,542
College Graduate	2,729	2,969	1,493	1,512	732	793
CURRENT EMPLOYMENT						
Full-Time	7,613 ^a	8,246	4,128	4,491	2,421 ^b	2,885
Part-Time	4,963	4,941	3,318	3,282	2,044	1,941
Unemployed	2,356	2,125	1,667 ^a	1,450	1,037	928
Other ¹	3,118	3,077	1,887	1,931	1,134	1,082

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.26B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	51.9	52.6	31.6	31.9	19.1	19.6
GENDER						
Male	56.1	55.9	36.8	36.2	23.5	23.2
Female	47.7	49.4	26.4	27.6	14.6	15.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	53.1	54.4	33.0	33.1	19.9	20.6
White	56.2	57.6	34.6	34.3	21.2	21.0
Black or African American	48.7	51.4	31.5	34.3	18.7 ^b	22.9
American Indian or Alaska Native	56.3	65.6	31.5	33.9	15.7	16.3
Native Hawaiian or Other Pacific Islander	*	*	*	*	18.2	*
Asian	29.0	26.8	18.5	16.0	7.7	9.1
Two or More Races	61.0	60.8	39.5	37.7	24.2	25.9
Hispanic or Latino	47.4	46.2	26.5	27.5	16.1	15.6
EDUCATION						
< High School	50.9	50.6	33.8	33.6	22.7	21.9
High School Graduate	49.9	50.6	30.5	31.2	18.5	20.2
Some College	53.4	55.1	32.5	34.1	19.9	20.4
College Graduate	54.1	53.3	29.6	27.1	14.5	14.2
CURRENT EMPLOYMENT						
Full-Time	56.3	57.8	30.5	31.5	17.9 ^b	20.2
Part-Time	51.3	52.1	34.3	34.6	21.1	20.5
Unemployed	56.7	54.7	40.1	37.3	24.9	23.9
Other ¹	41.9	42.1	25.3	26.4	15.2	14.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.27A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	92,568	94,732	18,606 ^b	20,707	11,412 ^b	13,521
GENDER						
Male	49,764 ^a	51,725	11,497 ^b	13,178	7,185 ^b	8,969
Female	42,804	43,007	7,108	7,529	4,227	4,553
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	83,502	85,469	16,867 ^b	18,723	10,314 ^b	12,272
White	69,280	70,577	13,411 ^a	14,917	8,055 ^b	9,735
Black or African American	10,165	10,300	2,352	2,777	1,537	1,884
American Indian or Alaska Native	548	605	170	159	100	122
Native Hawaiian or Other Pacific Islander	*	*	*	89	*	69
Asian	1,594	1,820	259	331	111	171
Two or More Races	1,648	1,776	579	450	442	290
Hispanic or Latino	9,067	9,263	1,739	1,984	1,098	1,250
EDUCATION						
< High School	8,402	9,133	2,394	2,623	1,705	1,846
High School Graduate	24,969	25,326	5,103	5,423	3,427	3,970
Some College	26,538	27,541	5,044 ^b	6,287	3,206 ^b	4,130
College Graduate	32,660	32,732	6,064	6,374	3,075	3,575
CURRENT EMPLOYMENT						
Full-Time	55,692	57,234	10,576 ^b	12,387	6,467 ^b	7,864
Part-Time	12,099	11,358	2,826	2,749	1,876	1,781
Unemployed	4,263	3,970	1,392	1,151	731	718
Other ¹	20,514	22,169	3,812	4,420	2,338 ^b	3,158

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.27B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	45.7	46.1	9.2 ^b	10.1	5.6 ^b	6.6
GENDER						
Male	51.3	52.7	11.9 ^b	13.4	7.4 ^b	9.1
Female	40.5	40.2	6.7	7.0	4.0	4.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	47.9	48.6	9.7 ^b	10.6	5.9 ^b	7.0
White	50.6	51.4	9.8 ^a	10.9	5.9 ^b	7.1
Black or African American	44.6	44.5	10.3	12.0	6.7	8.1
American Indian or Alaska Native	54.2	56.3	16.8	14.8	9.9	11.4
Native Hawaiian or Other Pacific Islander	*	*	*	12.0	*	9.4
Asian	15.6	17.1	2.5	3.1	1.1	1.6
Two or More Races	58.8	61.9	20.6	15.7	15.8	10.1
Hispanic or Latino	31.9	31.4	6.1	6.7	3.9	4.2
EDUCATION						
< High School	31.6	33.5	9.0	9.6	6.4	6.8
High School Graduate	43.0	44.5	8.8	9.5	5.9 ^a	7.0
Some College	51.8	51.8	9.8 ^b	11.8	6.3 ^b	7.8
College Graduate	48.8	48.2	9.1	9.4	4.6	5.3
CURRENT EMPLOYMENT						
Full-Time	52.8	52.7	10.0 ^b	11.4	6.1 ^b	7.2
Part-Time	50.2	48.3	11.7	11.7	7.8	7.6
Unemployed	55.7	57.5	18.2	16.7	9.6	10.4
Other ¹	31.3 ^a	33.5	5.8	6.7	3.6 ^b	4.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.28A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	110,619	113,121	29,606 ^b	31,862	18,048 ^b	20,357
GENDER						
Male	59,555 ^a	61,516	17,920 ^b	19,521	11,296 ^b	13,033
Female	51,064	51,605	11,686	12,341	6,752	7,324
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	98,113	100,458	25,942 ^b	27,849	15,786 ^b	17,960
White	80,233	81,735	20,148 ^a	21,562	12,197 ^b	13,804
Black or African American	12,601	12,889	3,927 ^a	4,505	2,473 ^b	3,035
American Indian or Alaska Native	678	738	243	228	136	155
Native Hawaiian or Other Pacific Islander	370	485	*	155	*	119
Asian	2,136	2,358	606	653	256	354
Two or More Races	2,095	2,253	868	746	620	493
Hispanic or Latino	12,506	12,664	3,663	4,013	2,262	2,397
EDUCATION						
< High School	11,247	11,635	4,281	4,285	2,974	2,929
High School Graduate	30,941	31,385	8,759	9,165	5,636 ^a	6,389
Some College	33,042	34,401	9,008 ^b	10,526	5,632 ^b	6,672
College Graduate	35,389	35,701	7,557	7,886	3,806	4,368
CURRENT EMPLOYMENT						
Full-Time	63,305	65,480	14,704 ^b	16,878	8,889 ^b	10,749
Part-Time	17,062	16,299	6,144	6,032	3,920	3,723
Unemployed	6,620	6,095	3,058 ^a	2,601	1,768	1,646
Other ¹	23,632	25,246	5,699	6,351	3,472 ^b	4,239

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.28B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	46.6	47.1	12.5 ^a	13.3	7.6 ^b	8.5
GENDER						
Male	52.1	53.1	15.7 ^a	16.9	9.9 ^b	11.3
Female	41.5	41.4	9.5	9.9	5.5	5.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	48.6	49.4	12.9 ^a	13.7	7.8 ^b	8.8
White	51.3	52.1	12.9 ^a	13.8	7.8 ^b	8.8
Black or African American	45.3	45.7	14.1 ^a	16.0	8.9 ^a	10.8
American Indian or Alaska Native	54.6	57.8	19.5	17.9	11.0	12.1
Native Hawaiian or Other Pacific Islander	47.8	52.9	*	16.9	*	12.9
Asian	17.6	18.7	5.0	5.2	2.1	2.8
Two or More Races	59.3	61.7	24.5	20.4	17.5	13.5
Hispanic or Latino	35.0	34.4	10.3	10.9	6.3	6.5
EDUCATION						
< High School	35.0	36.1	13.3	13.3	9.2	9.1
High School Graduate	44.2	45.5	12.5	13.3	8.1 ^b	9.3
Some College	52.1	52.5	14.2 ^b	16.0	8.9 ^a	10.2
College Graduate	49.2	48.6	10.5	10.7	5.3	5.9
CURRENT EMPLOYMENT						
Full-Time	53.2	53.3	12.4 ^b	13.7	7.5 ^b	8.7
Part-Time	50.6	49.4	18.2	18.3	11.6	11.3
Unemployed	56.1	56.5	25.9	24.1	15.0	15.3
Other ¹	32.4 ^a	34.3	7.8	8.6	4.8 ^b	5.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.29A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	37,634	39,200	4,182	4,553	1,549	1,530
AGE						
12-17	215	222	131	168	43	39
18-25	4,041	3,881	1,543	1,604	398	473
26 or Older	33,378	35,097	2,508	2,782	1,108	1,018
GENDER						
Male	22,612	23,917	2,855	2,965	1,038	991
Female	15,022	15,283	1,327	1,588	511	539
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	32,901	34,410	3,520	3,735	1,313	1,249
White	28,674	29,617	2,961	2,916	1,036	938
Black or African American	2,748	3,020	401	565	246	190
American Indian or Alaska Native	197	327	16	45	1	26
Native Hawaiian or Other Pacific Islander	*	133	13	15	*	3
Asian	383	536	77	106	17	66
Two or More Races	752	777	53	89	10	26
Hispanic or Latino	4,733	4,790	662	817	236	281

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.29B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	14.3	14.8	1.6	1.7	0.6	0.6
AGE						
12-17	0.9	0.9	0.5	0.7	0.2	0.2
18-25	11.6	11.1	4.4	4.6	1.1	1.4
26 or Older	16.5	17.1	1.2	1.4	0.5	0.5
GENDER						
Male	17.8	18.6	2.2	2.3	0.8	0.8
Female	11.1	11.2	1.0	1.2	0.4	0.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	14.9	15.5	1.6	1.7	0.6	0.6
White	16.9	17.4	1.7	1.7	0.6	0.6
Black or African American	8.8	9.5	1.3	1.8	0.8	0.6
American Indian or Alaska Native	14.0 ^a	22.7	1.1	3.1	0.1	1.8
Native Hawaiian or Other Pacific Islander	*	13.0	1.5	1.5	*	0.3
Asian	2.9	3.9	0.6	0.8	0.1	0.5
Two or More Races	17.5	17.6	1.2	2.0	0.2	0.6
Hispanic or Latino	11.5	11.3	1.6	1.9	0.6	0.7

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.30A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	215	222	131	168	43	39
GENDER						
Male	121	105	74	87	19	16
Female	94	117	58	81	24	23
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	145	143	96	101	29	18
White	132	120	86	88	25	17
Black or African American	4	16	4	7	4	*
American Indian or Alaska Native	0	1	*	*	*	*
Native Hawaiian or Other Pacific Islander	2	*	*	*	*	*
Asian	1	*	*	*	*	*
Two or More Races	6	5	5	4	*	0
Hispanic or Latino	70	80	35	67	14	21
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	70	57	47	46	9	7
Female, White, Not Hispanic	62	63	40	43	16	11
Male, Black, Not Hispanic	4	9	4	7	4	*
Female, Black, Not Hispanic	0	7	0	*	*	*
Male, Hispanic	45	38	21	33	7	9
Female, Hispanic	25	42	14	34	8	12

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.30B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9	0.9	0.5	0.7	0.2	0.2
GENDER						
Male	1.0	0.8	0.6	0.7	0.2	0.1
Female	0.8	1.0	0.5	0.7	0.2	0.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.8	0.7	0.5	0.5	0.2	0.1
White	1.0	0.9	0.6	0.7	0.2	0.1
Black or African American	0.1	0.5	0.1	0.2	0.1	*
American Indian or Alaska Native	0.3	0.9	*	*	*	*
Native Hawaiian or Other Pacific Islander	1.6	*	*	*	*	*
Asian	0.1	*	*	*	*	*
Two or More Races	0.9	0.7	0.6	0.6	*	0.0
Hispanic or Latino	1.3	1.4	0.6	1.2	0.3	0.4
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	1.0	0.8	0.7	0.7	0.1	0.1
Female, White, Not Hispanic	0.9	1.0	0.6	0.6	0.2	0.2
Male, Black, Not Hispanic	0.2	0.5	0.2	0.4	0.2	*
Female, Black, Not Hispanic	0.0	0.4	0.0	*	*	*
Male, Hispanic	1.6	1.3	0.7	1.2	0.2	0.3
Female, Hispanic	0.9	1.5	0.5	1.2	0.3	0.4

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.31A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	4,041	3,881	1,543	1,604	398	473
GENDER						
Male	2,408	2,323	994	1,008	263	268
Female	1,633	1,558	549	596	134 ^a	205
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	3,189	3,051	1,267	1,228	322	376
White	2,845	2,692	1,143	1,049	284	314
Black or African American	106	156	47	84	23	27
American Indian or Alaska Native	25	24	8	9	1	4
Native Hawaiian or Other Pacific Islander	15	*	*	*	*	*
Asian	84	66	29	43	6	21
Two or More Races	113	96	33	38	5	8
Hispanic or Latino	852	830	276	376	76	97
EDUCATION						
< High School	725 ^a	581	215	176	54	47
High School Graduate	1,300	1,256	454	515	131	148
Some College	1,365	1,409	566	614	147	191
College Graduate	651	635	308	299	66	86
CURRENT EMPLOYMENT						
Full-Time	1,914	2,005	701	762	183	200
Part-Time	981	873	448	440	121	170
Unemployed	522	479	169	182	32	41
Other ¹	624	525	225	221	62	61

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.31B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	11.6	11.1	4.4	4.6	1.1	1.4
GENDER						
Male	13.8	13.3	5.7	5.7	1.5	1.5
Female	9.4	9.0	3.2	3.4	0.8 ^a	1.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	11.6	11.1	4.6	4.5	1.2	1.4
White	14.6	13.9	5.9	5.4	1.5	1.6
Black or African American	2.1	3.1	0.9	1.7	0.5	0.5
American Indian or Alaska Native	11.1	11.7	3.6	4.6	0.6	1.8
Native Hawaiian or Other Pacific Islander	7.7	*	*	*	*	*
Asian	4.5	3.3	1.6	2.2	0.3	1.0
Two or More Races	15.4	12.3	4.6	4.8	0.6	1.0
Hispanic or Latino	11.8	11.3	3.8	5.1	1.1	1.3
EDUCATION						
< High School	13.0	11.7	3.9	3.6	1.0	1.0
High School Graduate	10.9	10.5	3.8	4.3	1.1	1.2
Some College	11.2	11.3	4.6	4.9	1.2	1.5
College Graduate	12.9	11.4	6.1	5.4	1.3	1.5
CURRENT EMPLOYMENT						
Full-Time	14.2	14.1	5.2	5.3	1.4	1.4
Part-Time	10.2	9.2	4.6	4.6	1.2	1.8
Unemployed	12.6	12.3	4.1	4.7	0.8	1.1
Other ¹	8.4	7.2	3.0	3.0	0.8	0.8

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.32A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	33,378	35,097	2,508	2,782	1,108	1,018
GENDER						
Male	20,083 ^a	21,489	1,788	1,870	756	708
Female	13,295	13,608	720	911	352	310
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	29,566	31,216	2,156	2,407	962	855
White	25,697	26,805	1,731	1,779	727	607
Black or African American	2,637	2,848	350	473	219	163
American Indian or Alaska Native	171	302	8	35	*	21
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	298	470	48	62	10	45
Two or More Races	632	675	14	47	5	18
Hispanic or Latino	3,811	3,881	351	374	145	163
EDUCATION						
< High School	3,861	4,207	386	388	220	109
High School Graduate	9,139	9,979	569	787	256	303
Some College	9,868	10,888	818	850	356	334
College Graduate	10,510	10,024	734	756	276	273
CURRENT EMPLOYMENT						
Full-Time	19,922	21,309	1,260	1,517	396	533
Part-Time	4,881	4,182	418	474	237	149
Unemployed	1,722	1,738	243	201	109	116
Other ¹	6,853 ^a	7,868	586	591	365	220

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.32B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	16.5	17.1	1.2	1.4	0.5	0.5
GENDER						
Male	20.7	21.9	1.8	1.9	0.8	0.7
Female	12.6	12.7	0.7	0.9	0.3	0.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	17.0	17.8	1.2	1.4	0.6	0.5
White	18.8	19.5	1.3	1.3	0.5	0.4
Black or African American	11.6	12.3	1.5	2.0	1.0	0.7
American Indian or Alaska Native	16.9 ^a	28.1	0.7	3.2	*	2.0
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.9	4.4	0.5	0.6	0.1	0.4
Two or More Races	22.5	23.5	0.5	1.6	0.2	0.6
Hispanic or Latino	13.4	13.2	1.2	1.3	0.5	0.6
EDUCATION						
< High School	14.5	15.4	1.5	1.4	0.8	0.4
High School Graduate	15.8 ^a	17.5	1.0	1.4	0.4	0.5
Some College	19.3	20.5	1.6	1.6	0.7	0.6
College Graduate	15.7	14.8	1.1	1.1	0.4	0.4
CURRENT EMPLOYMENT						
Full-Time	18.9	19.6	1.2	1.4	0.4	0.5
Part-Time	20.3	17.8	1.7	2.0	1.0	0.6
Unemployed	22.5	25.2	3.2	2.9	1.4	1.7
Other ¹	10.5 ^a	11.9	0.9	0.9	0.6	0.3

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.33A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	37,419	38,978	4,051	4,385	1,505	1,491
GENDER						
Male	22,491	23,812	2,781	2,878	1,019	975
Female	14,928	15,166	1,269	1,507	487	515
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	32,755	34,267	3,424	3,635	1,284	1,231
White	28,542	29,497	2,874	2,828	1,011	921
Black or African American	2,744	3,004	397	557	242	190
American Indian or Alaska Native	196	326	16	44	1	25
Native Hawaiian or Other Pacific Islander	*	133	11	15	*	3
Asian	382	536	77	106	17	66
Two or More Races	745	771	48	85	10	26
Hispanic or Latino	4,663	4,711	627	751	222	260
EDUCATION						
< High School	4,586	4,788	602	564	274	156
High School Graduate	10,439	11,235	1,023 ^a	1,302	387	451
Some College	11,233	12,297	1,384	1,464	503	525
College Graduate	11,161	10,658	1,042	1,056	341	359
CURRENT EMPLOYMENT						
Full-Time	21,835	23,314	1,961	2,278	579	733
Part-Time	5,862 ^a	5,055	866	913	358	319
Unemployed	2,244	2,217	412	382	142	157
Other ¹	7,477	8,393	811	811	427	281

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.33B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	15.8	16.2	1.7	1.8	0.6	0.6
GENDER						
Male	19.7	20.6	2.4	2.5	0.9	0.8
Female	12.1	12.2	1.0	1.2	0.4	0.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	16.2	16.8	1.7	1.8	0.6	0.6
White	18.3	18.8	1.8	1.8	0.6	0.6
Black or African American	9.9	10.7	1.4	2.0	0.9	0.7
American Indian or Alaska Native	15.8 ^a	25.5	1.3	3.4	0.1	2.0
Native Hawaiian or Other Pacific Islander	*	14.5	1.5	1.6	*	0.3
Asian	3.2	4.2	0.6	0.8	0.1	0.5
Two or More Races	21.1	21.1	1.4	2.3	0.3	0.7
Hispanic or Latino	13.1	12.8	1.8	2.0	0.6	0.7
EDUCATION						
< High School	14.3	14.9	1.9	1.8	0.9	0.5
High School Graduate	14.9 ^a	16.3	1.5 ^a	1.9	0.6	0.7
Some College	17.7	18.8	2.2	2.2	0.8	0.8
College Graduate	15.5	14.5	1.4	1.4	0.5	0.5
CURRENT EMPLOYMENT						
Full-Time	18.4	19.0	1.6	1.9	0.5	0.6
Part-Time	17.4 ^a	15.3	2.6	2.8	1.1	1.0
Unemployed	19.0	20.5	3.5	3.5	1.2	1.5
Other ¹	10.2	11.4	1.1	1.1	0.6	0.4

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.34A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8,870	9,424	632	773	377	354
AGE						
12-17	23	35	11	20	3	8
18-25	563	546	95	127	36	29
26 or Older	8,284	8,843	526	627	338	317
GENDER						
Male	5,504	6,163	390	479	218	206
Female	3,366	3,261	242	294	160	149
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	7,868	8,466	536	627	303	304
White	6,002	6,483	371	400	204	182
Black or African American	1,449	1,496	154	188	94	103
American Indian or Alaska Native	76	128	4	2	1	1
Native Hawaiian or Other Pacific Islander	*	17	1	*	*	*
Asian	73	110	4	29	4	18
Two or More Races	181	231	2	8	1	*
Hispanic or Latino	1,002	957	96	146	74	51

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.34B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	3.4	3.6	0.2	0.3	0.1	0.1
AGE						
12-17	0.1	0.1	0.0	0.1	0.0	0.0
18-25	1.6	1.6	0.3	0.4	0.1	0.1
26 or Older	4.1	4.3	0.3	0.3	0.2	0.2
GENDER						
Male	4.3	4.8	0.3	0.4	0.2	0.2
Female	2.5	2.4	0.2	0.2	0.1	0.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	3.6	3.8	0.2	0.3	0.1	0.1
White	3.5	3.8	0.2	0.2	0.1	0.1
Black or African American	4.6	4.7	0.5	0.6	0.3	0.3
American Indian or Alaska Native	5.4	8.9	0.3	0.2	0.1	0.1
Native Hawaiian or Other Pacific Islander	*	1.7	0.1	*	*	*
Asian	0.5	0.8	0.0	0.2	0.0	0.1
Two or More Races	4.2	5.2	0.1	0.2	0.0	*
Hispanic or Latino	2.4	2.3	0.2	0.3	0.2	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.35A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	23	35	11	20	3	8
GENDER						
Male	15	17	7	12	2	7
Female	7	17	4	8	*	1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	14	24	8	10	1	4
White	13	22	6	9	1	4
Black or African American	*	*	*	*	*	*
American Indian or Alaska Native	0	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	*	*	*	*	*	*
Two or More Races	0	1	*	1	*	*
Hispanic or Latino	8	11	4	10	1	4
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	11	9	5	4	1	3
Female, White, Not Hispanic	2	13	2	4	*	1
Male, Black, Not Hispanic	*	*	*	*	*	*
Female, Black, Not Hispanic	*	*	*	*	*	*
Male, Hispanic	3	7	1	7	1	4
Female, Hispanic	5	4	2	3	0	*

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.35B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1	0.1	0.0	0.1	0.0	0.0
GENDER						
Male	0.1	0.1	0.1	0.1	0.0	0.1
Female	0.1	0.1	0.0	0.1	*	0.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1	0.1	0.0	0.1	0.0	0.0
White	0.1	0.2	0.0	0.1	0.0	0.0
Black or African American	*	*	*	*	*	*
American Indian or Alaska Native	0.1	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	*	*	*	*	*	*
Two or More Races	0.0	0.1	*	0.1	*	*
Hispanic or Latino	0.2	0.2	0.1	0.2	0.0	0.1
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.2	0.1	0.1	0.1	0.0	0.0
Female, White, Not Hispanic	0.0 ^a	0.2	0.0	0.1	*	0.0
Male, Black, Not Hispanic	*	*	*	*	*	*
Female, Black, Not Hispanic	*	*	*	*	*	*
Male, Hispanic	0.1	0.3	0.0	0.2	0.0	0.1
Female, Hispanic	0.2	0.1	0.1	0.1	0.0	*

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.36A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	563	546	95	127	36	29
GENDER						
Male	343	308	59	79	24	21
Female	221	238	36	47	13	8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	448	477	70	107	21	21
White	414	441	64	94	19	20
Black or African American	10	21	2	12	*	*
American Indian or Alaska Native	3	5	2	1	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	9	1	*	*	*	*
Two or More Races	9	9	2	*	1	*
Hispanic or Latino	115	69	24	20	15	8
EDUCATION						
< High School	175 ^a	110	21	25	8	8
High School Graduate	238	233	47	53	23	10
Some College	131	169	20	42	6	11
College Graduate	20	34	6	6	*	*
CURRENT EMPLOYMENT						
Full-Time	265	274	38	56	18	7
Part-Time	102	100	21	25	7	19
Unemployed	113	86	15	30	3	1
Other ¹	84	86	22	17	8	3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.36B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1.6	1.6	0.3	0.4	0.1	0.1
GENDER						
Male	2.0	1.8	0.3	0.5	0.1	0.1
Female	1.3	1.4	0.2	0.3	0.1	0.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1.6	1.7	0.3	0.4	0.1	0.1
White	2.1	2.3	0.3	0.5	0.1	0.1
Black or African American	0.2	0.4	0.0	0.2	*	*
American Indian or Alaska Native	1.1	2.5	0.9	0.6	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	0.5	0.0	*	*	*	*
Two or More Races	1.2	1.1	0.3	*	0.1	*
Hispanic or Latino	1.6	0.9	0.3	0.3	0.2	0.1
EDUCATION						
< High School	3.1	2.2	0.4	0.5	0.1	0.2
High School Graduate	2.0	1.9	0.4	0.4	0.2	0.1
Some College	1.1	1.4	0.2	0.3	0.0	0.1
College Graduate	0.4	0.6	0.1	0.1	*	*
CURRENT EMPLOYMENT						
Full-Time	2.0	1.9	0.3	0.4	0.1	0.0
Part-Time	1.1	1.1	0.2	0.3	0.1	0.2
Unemployed	2.7	2.2	0.4	0.8	0.1	0.0
Other ¹	1.1	1.2	0.3	0.2	0.1	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.37A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8,284	8,843	526	627	338	317
GENDER						
Male	5,146	5,837	324	388	192	178
Female	3,137	3,005	202	239	146	139
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	7,405	7,965	458	511	281	279
White	5,576	6,020	300	298	183	158
Black or African American	1,439	1,475	153	176	94	103
American Indian or Alaska Native	73	*	2	1	*	*
Native Hawaiian or Other Pacific Islander	*	16	*	*	*	*
Asian	64	110	4	29	4	18
Two or More Races	172	222	*	8	*	*
Hispanic or Latino	878	877	68	117	57	38
EDUCATION						
< High School	1,678	1,749	150	151	69	60
High School Graduate	2,767	3,127	144	248	78	170
Some College	2,685	2,526	164	159	134	55
College Graduate	1,154	1,441	67	69	57	32
CURRENT EMPLOYMENT						
Full-Time	4,093	4,497	99	201	66	74
Part-Time	1,105	1,039	64	143	41	93
Unemployed	646	549	68	63	53	50
Other ¹	2,440	2,758	295	220	177	100

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.37B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	4.1	4.3	0.3	0.3	0.2	0.2
GENDER						
Male	5.3	5.9	0.3	0.4	0.2	0.2
Female	3.0	2.8	0.2	0.2	0.1	0.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4.2	4.5	0.3	0.3	0.2	0.2
White	4.1	4.4	0.2	0.2	0.1	0.1
Black or African American	6.3	6.4	0.7	0.8	0.4	0.4
American Indian or Alaska Native	7.2	*	0.2	0.1	*	*
Native Hawaiian or Other Pacific Islander	*	2.2	*	*	*	*
Asian	0.6	1.0	0.0	0.3	0.0	0.2
Two or More Races	6.1	7.7	*	0.3	*	*
Hispanic or Latino	3.1	3.0	0.2	0.4	0.2	0.1
EDUCATION						
< High School	6.3	6.4	0.6	0.6	0.3	0.2
High School Graduate	4.8	5.5	0.2	0.4	0.1	0.3
Some College	5.2	4.8	0.3	0.3	0.3	0.1
College Graduate	1.7	2.1	0.1	0.1	0.1	0.0
CURRENT EMPLOYMENT						
Full-Time	3.9	4.1	0.1	0.2	0.1	0.1
Part-Time	4.6	4.4	0.3	0.6	0.2	0.4
Unemployed	8.4	8.0	0.9	0.9	0.7	0.7
Other ¹	3.7	4.2	0.4	0.3	0.3	0.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.38A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8,847	9,389	621	754	375	347
GENDER						
Male	5,489	6,146	383	467	215	199
Female	3,358	3,243	238	286	159	148
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	7,854	8,443	529	617	302	300
White	5,989	6,461	364	391	203	179
Black or African American	1,449	1,496	154	188	94	103
American Indian or Alaska Native	76	128	4	2	1	1
Native Hawaiian or Other Pacific Islander	*	17	*	*	*	*
Asian	73	110	4	29	4	18
Two or More Races	181	231	2	8	1	*
Hispanic or Latino	993	946	92	136	72	46
EDUCATION						
< High School	1,853	1,859	171	176	77	69
High School Graduate	3,005	3,360	192	302	101	179
Some College	2,815	2,695	185	201	140	66
College Graduate	1,174	1,475	73	74	57	32
CURRENT EMPLOYMENT						
Full-Time	4,358	4,771	137 ^a	256	85	80
Part-Time	1,207	1,139	85	168	49	112
Unemployed	759	635	83	93	56	51
Other ¹	2,523	2,843	316	237	185	103

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.38B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	3.7	3.9	0.3	0.3	0.2	0.1
GENDER						
Male	4.8	5.3	0.3	0.4	0.2	0.2
Female	2.7	2.6	0.2	0.2	0.1	0.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	3.9	4.2	0.3	0.3	0.1	0.1
White	3.8	4.1	0.2	0.2	0.1	0.1
Black or African American	5.2	5.3	0.6	0.7	0.3	0.4
American Indian or Alaska Native	6.1	10.0	0.3	0.1	0.1	0.1
Native Hawaiian or Other Pacific Islander	*	1.9	*	*	*	*
Asian	0.6	0.9	0.0	0.2	0.0	0.1
Two or More Races	5.1	6.3	0.1	0.2	0.0	*
Hispanic or Latino	2.8	2.6	0.3	0.4	0.2	0.1
EDUCATION						
< High School	5.8	5.8	0.5	0.5	0.2	0.2
High School Graduate	4.3	4.9	0.3	0.4	0.1	0.3
Some College	4.4	4.1	0.3	0.3	0.2	0.1
College Graduate	1.6	2.0	0.1	0.1	0.1	0.0
CURRENT EMPLOYMENT						
Full-Time	3.7	3.9	0.1 ^a	0.2	0.1	0.1
Part-Time	3.6	3.4	0.3	0.5	0.1	0.3
Unemployed	6.4	5.9	0.7	0.9	0.5	0.5
Other ¹	3.5	3.9	0.4	0.3	0.3	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.39A Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	39,736	39,647	4,430	4,250	1,333	1,173
AGE						
12-17	702	610	482	422	154	136
18-25	6,122	5,784	2,338	2,205	617	502
26 or Older	32,912	33,253	1,610	1,624	562	535
GENDER						
Male	23,835	24,029	2,809	2,756	902	748
Female	15,902	15,617	1,621	1,494	431	426
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	35,681	35,776	3,794	3,613	1,158	952
White	31,114	31,021	3,081	2,948	873	798
Black or African American	2,472	2,528	326	297	131	67
American Indian or Alaska Native	345	361	72	43	11	5
Native Hawaiian or Other Pacific Islander	128	110	8	21	3	3
Asian	716	762	129	156	44	36
Two or More Races	905	994	179	147	*	42
Hispanic or Latino	4,055	3,871	636	637	175	221

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.39B Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	15.1	15.0	1.7	1.6	0.5	0.4
AGE						
12-17	2.8	2.5	1.9	1.7	0.6	0.5
18-25	17.6	16.6	6.7	6.3	1.8	1.4
26 or Older	16.2	16.2	0.8	0.8	0.3	0.3
GENDER						
Male	18.7	18.7	2.2	2.1	0.7	0.6
Female	11.8	11.4	1.2	1.1	0.3	0.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	16.1	16.1	1.7	1.6	0.5	0.4
White	18.3	18.2	1.8	1.7	0.5	0.5
Black or African American	7.9	8.0	1.0	0.9	0.4	0.2
American Indian or Alaska Native	24.6	25.0	5.1	3.0	0.8	0.4
Native Hawaiian or Other Pacific Islander	14.7	10.8	0.9	2.0	0.4	0.3
Asian	5.4	5.5	1.0	1.1	0.3	0.3
Two or More Races	21.1	22.5	4.2	3.3	*	1.0
Hispanic or Latino	9.8	9.1	1.5	1.5	0.4	0.5

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.40A Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	702	610	482	422	154	136
GENDER						
Male	412 ^a	335	295 ^a	226	98	85
Female	291	274	186	195	57	51
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	555 ^a	464	381	328	133	101
White	437	390	302	278	102	86
Black or African American	45	36	26	17	12	7
American Indian or Alaska Native	*	5	*	4	*	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	25 ^a	4	21	4	8	*
Two or More Races	29	27	21	22	5	6
Hispanic or Latino	147	145	100	93	21	35
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	256	218	181	153	62	58
Female, White, Not Hispanic	182	171	121	125	40	28
Male, Black, Not Hispanic	27	21	15	9	6	7
Female, Black, Not Hispanic	18	15	12	8	6	0
Male, Hispanic	91	78	71	48	14	14
Female, Hispanic	56	67	30	45	7	21

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.40B Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	2.8	2.5	1.9	1.7	0.6	0.5
GENDER						
Male	3.2 ^a	2.6	2.3 ^a	1.8	0.8	0.7
Female	2.4	2.3	1.5	1.6	0.5	0.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2.9	2.4	2.0	1.7	0.7	0.5
White	3.2	2.9	2.2	2.1	0.7	0.6
Black or African American	1.3	1.0	0.8	0.5	0.3	0.2
American Indian or Alaska Native	*	3.1	*	2.5	*	0.9
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.0 ^a	0.4	1.7	0.4	0.6	*
Two or More Races	3.9	3.5	2.8	2.9	0.7	0.7
Hispanic or Latino	2.7	2.6	1.8	1.6	0.4	0.6
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	3.7	3.2	2.6	2.2	0.9	0.8
Female, White, Not Hispanic	2.7	2.6	1.8	1.9	0.6	0.4
Male, Black, Not Hispanic	1.5	1.2	0.8	0.5	0.3	0.4
Female, Black, Not Hispanic	1.1	0.9	0.7	0.5	0.3	0.0
Male, Hispanic	3.2	2.7	2.5	1.7	0.5	0.5
Female, Hispanic	2.1	2.4	1.1	1.6	0.2	0.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.41A Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	6,122	5,784	2,338	2,205	617	502
GENDER						
Male	3,679	3,422	1,550	1,394	441 ^b	293
Female	2,444	2,362	788	811	177	209
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5,058	4,740	1,989	1,821	559 ^b	380
White	4,191	3,887	1,678	1,492	463 ^b	311
Black or African American	439	430	145	133	52 ^a	17
American Indian or Alaska Native	53	39	11	*	4	2
Native Hawaiian or Other Pacific Islander	31	*	5	*	*	*
Asian	185	173	91	101	28	27
Two or More Races	159	184	58	69	12	21
Hispanic or Latino	1,064	1,044	349	384	59 ^a	122
EDUCATION						
< High School	1,009 ^a	832	302	258	95	66
High School Graduate	1,967	1,847	685	663	195	156
Some College	2,244	2,197	932	891	245	202
College Graduate	903	908	418	393	82	77
CURRENT EMPLOYMENT						
Full-Time	2,634	2,734	860	932	220	202
Part-Time	1,614	1,503	742	685	209 ^a	137
Unemployed	847 ^a	688	341	277	63	77
Other ¹	1,028	859	395	311	125	86

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.41B Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	17.6	16.6	6.7	6.3	1.8	1.4
GENDER						
Male	21.1	19.5	8.9	8.0	2.5 ^b	1.7
Female	14.1	13.6	4.5	4.7	1.0	1.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	18.4	17.2	7.2	6.6	2.0 ^b	1.4
White	21.5	20.1	8.6	7.7	2.4 ^b	1.6
Black or African American	8.8	8.5	2.9	2.6	1.0 ^a	0.3
American Indian or Alaska Native	23.3	19.2	4.6	*	1.5	1.2
Native Hawaiian or Other Pacific Islander	15.3	*	2.6	*	*	*
Asian	9.9	8.6	4.9	5.0	1.5	1.3
Two or More Races	21.7	23.4	8.0	8.8	1.6	2.7
Hispanic or Latino	14.7	14.2	4.8	5.2	0.8 ^a	1.7
EDUCATION						
< High School	18.1	16.8	5.4	5.2	1.7	1.3
High School Graduate	16.4	15.4	5.7	5.5	1.6	1.3
Some College	18.4	17.7	7.7	7.2	2.0	1.6
College Graduate	17.9	16.3	8.3	7.1	1.6	1.4
CURRENT EMPLOYMENT						
Full-Time	19.5	19.2	6.4	6.5	1.6	1.4
Part-Time	16.7	15.9	7.7	7.2	2.2	1.4
Unemployed	20.4	17.7	8.2	7.1	1.5	2.0
Other ¹	13.8	11.7	5.3	4.3	1.7	1.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.42A Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	32,912	33,253	1,610	1,624	562	535
GENDER						
Male	19,744	20,271	964	1,136	364	370
Female	13,167	12,981	647	488	198	165
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	30,068	30,571	1,424	1,463	466	471
White	26,485	26,745	1,101	1,178	308	401
Black or African American	1,989	2,062	154	147	68	43
American Indian or Alaska Native	276	317	*	17	4	1
Native Hawaiian or Other Pacific Islander	*	80	*	*	*	*
Asian	507	584	16	51	8	10
Two or More Races	717	784	*	57	*	16
Hispanic or Latino	2,843	2,681	187	161	96	64
EDUCATION						
< High School	3,052	2,951	122	95	26	47
High School Graduate	8,150	8,820	384	297	160	97
Some College	10,388	10,208	439	517	85	173
College Graduate	11,321	11,273	665	715	291	219
CURRENT EMPLOYMENT						
Full-Time	19,713	20,698	793 ^a	1,113	241	395
Part-Time	4,940	4,228	380	212	148	69
Unemployed	1,759	1,479	153	114	35	27
Other ¹	6,500	6,847	285	184	138	44

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.42B Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	16.2	16.2	0.8	0.8	0.3	0.3
GENDER						
Male	20.4	20.6	1.0	1.2	0.4	0.4
Female	12.4	12.1	0.6	0.5	0.2	0.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	17.3	17.4	0.8	0.8	0.3	0.3
White	19.4	19.5	0.8	0.9	0.2	0.3
Black or African American	8.7	8.9	0.7	0.6	0.3	0.2
American Indian or Alaska Native	27.2	29.5	*	1.6	0.4	0.1
Native Hawaiian or Other Pacific Islander	*	10.8	*	*	*	*
Asian	4.9	5.5	0.2	0.5	0.1	0.1
Two or More Races	25.6	27.3	*	2.0	*	0.5
Hispanic or Latino	10.0	9.1	0.7	0.5	0.3	0.2
EDUCATION						
< High School	11.5	10.8	0.5	0.3	0.1	0.2
High School Graduate	14.1 ^a	15.5	0.7	0.5	0.3	0.2
Some College	20.3	19.2	0.9	1.0	0.2	0.3
College Graduate	16.9	16.6	1.0	1.1	0.4	0.3
CURRENT EMPLOYMENT						
Full-Time	18.7	19.0	0.8 ^a	1.0	0.2	0.4
Part-Time	20.5	18.0	1.6	0.9	0.6	0.3
Unemployed	23.0	21.4	2.0	1.7	0.5	0.4
Other ¹	9.9	10.3	0.4	0.3	0.2	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.43A Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	39,034	39,037	3,948	3,829	1,179	1,037
GENDER						
Male	23,423	23,694	2,514	2,530	804	663
Female	15,611	15,343	1,435	1,299	375	374
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	35,126	35,312	3,412	3,285	1,025	850
White	30,677	30,632	2,779	2,670	771	712
Black or African American	2,428	2,492	299	280	119	60
American Indian or Alaska Native	329	356	*	39	8	4
Native Hawaiian or Other Pacific Islander	*	107	5	18	1	2
Asian	691	757	108	152	36	36
Two or More Races	876	967	158	125	*	37
Hispanic or Latino	3,908	3,725	536	544	154	187
EDUCATION						
< High School	4,061	3,783	424	353	121	114
High School Graduate	10,117	10,667	1,069	959	355	252
Some College	12,632	12,405	1,371	1,408	331	375
College Graduate	12,224	12,182	1,084	1,109	372	296
CURRENT EMPLOYMENT						
Full-Time	22,347	23,432	1,653 ^a	2,045	461	597
Part-Time	6,554 ^a	5,731	1,121	898	357	207
Unemployed	2,605 ^a	2,167	494	391	98	104
Other ¹	7,528	7,707	680	496	263 ^a	130

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.43B Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	16.4	16.2	1.7	1.6	0.5	0.4
GENDER						
Male	20.5	20.5	2.2	2.2	0.7	0.6
Female	12.7	12.3	1.2	1.0	0.3	0.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	17.4	17.4	1.7	1.6	0.5	0.4
White	19.6	19.5	1.8	1.7	0.5	0.5
Black or African American	8.7	8.8	1.1	1.0	0.4	0.2
American Indian or Alaska Native	26.5	27.9	*	3.1	0.6	0.3
Native Hawaiian or Other Pacific Islander	*	11.7	0.7	2.0	0.1	0.2
Asian	5.7	6.0	0.9	1.2	0.3	0.3
Two or More Races	24.8	26.5	4.5	3.4	*	1.0
Hispanic or Latino	10.9	10.1	1.5	1.5	0.4	0.5
EDUCATION						
< High School	12.6	11.7	1.3	1.1	0.4	0.4
High School Graduate	14.5	15.5	1.5	1.4	0.5	0.4
Some College	19.9	18.9	2.2	2.1	0.5	0.6
College Graduate	17.0	16.6	1.5	1.5	0.5	0.4
CURRENT EMPLOYMENT						
Full-Time	18.8	19.1	1.4 ^a	1.7	0.4	0.5
Part-Time	19.4	17.4	3.3	2.7	1.1	0.6
Unemployed	22.1	20.1	4.2	3.6	0.8	1.0
Other ¹	10.3	10.5	0.9	0.7	0.4 ^a	0.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.44A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	21,068	21,293	1,533	1,617	496	546
AGE						
12-17	1,327	1,307	480	512	121	149
18-25	2,613	2,444	478	379	106	80
26 or Older	17,129	17,543	575	725	269	316
GENDER						
Male	13,675	14,058	1,025	1,082	342	390
Female	7,393	7,235	508	535	153	155
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	18,646	18,847	1,285	1,275	436	398
White	16,639	16,667	1,051	1,015	341	301
Black or African American	875	1,038	142	141	52	46
American Indian or Alaska Native	135	170	6	10	3	4
Native Hawaiian or Other Pacific Islander	*	62	*	3	*	2
Asian	326	371	49	58	28	23
Two or More Races	587	539	37	47	12	23
Hispanic or Latino	2,422	2,446	248	342	60	148

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.44B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8.0	8.0	0.6	0.6	0.2	0.2
AGE						
12-17	5.3	5.3	1.9	2.1	0.5	0.6
18-25	7.5	7.0	1.4	1.1	0.3	0.2
26 or Older	8.4	8.5	0.3	0.4	0.1	0.2
GENDER						
Male	10.8	10.9	0.8	0.8	0.3	0.3
Female	5.5	5.3	0.4	0.4	0.1	0.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8.4	8.5	0.6	0.6	0.2	0.2
White	9.8	9.8	0.6	0.6	0.2	0.2
Black or African American	2.8	3.3	0.5	0.4	0.2	0.1
American Indian or Alaska Native	9.6	11.8	0.4	0.7	0.2	0.3
Native Hawaiian or Other Pacific Islander	*	6.1	*	0.3	*	0.1
Asian	2.4	2.7	0.4	0.4	0.2	0.2
Two or More Races	13.7	12.2	0.9	1.1	0.3	0.5
Hispanic or Latino	5.9	5.8	0.6	0.8	0.1	0.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.45A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,327	1,307	480	512	121	149
GENDER						
Male	683	595	237	217	59	70
Female	644	712	243	296	62	79
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,025	981	384	377	96	106
White	698	662	265	287	62	72
Black or African American	215	207	90 ^b	47	28	19
American Indian or Alaska Native	14	10	3	5	1	1
Native Hawaiian or Other Pacific Islander	2	*	*	*	*	*
Asian	63	42	13	15	1	2
Two or More Races	34	54	13	22	4	10
Hispanic or Latino	302	325	96	136	25	43
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	364	309	138	120	29	33
Female, White, Not Hispanic	333	353	126	167	33	39
Male, Black, Not Hispanic	103	82	32	18	10	8
Female, Black, Not Hispanic	112	125	58 ^a	28	17	12
Male, Hispanic	149	149	58	60	18	28
Female, Hispanic	153	176	38 ^a	76	7	15

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.45B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	5.3	5.3	1.9	2.1	0.5	0.6
GENDER						
Male	5.4	4.7	1.9	1.7	0.5	0.6
Female	5.3	5.8	2.0	2.4	0.5	0.6
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5.3	5.1	2.0	2.0	0.5	0.6
White	5.1	4.9	1.9	2.1	0.5	0.5
Black or African American	6.2	5.9	2.6 ^b	1.3	0.8	0.6
American Indian or Alaska Native	8.5	5.8	1.7	2.8	0.8	0.7
Native Hawaiian or Other Pacific Islander	2.0	*	*	*	*	*
Asian	5.1	3.4	1.1	1.2	0.1	0.2
Two or More Races	4.5	7.1	1.7	2.8	0.5	1.2
Hispanic or Latino	5.4	5.8	1.7	2.4	0.4	0.8
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	5.2	4.5	2.0	1.7	0.4	0.5
Female, White, Not Hispanic	5.0	5.4	1.9	2.5	0.5	0.6
Male, Black, Not Hispanic	5.9	4.6	1.8	1.0	0.6	0.4
Female, Black, Not Hispanic	6.6	7.3	3.5 ^a	1.7	1.0	0.7
Male, Hispanic	5.2	5.2	2.0	2.1	0.6	1.0
Female, Hispanic	5.6	6.3	1.4 ^a	2.7	0.3	0.5

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.46A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	2,613	2,444	478	379	106	80
GENDER						
Male	1,525	1,500	343 ^a	242	82	54
Female	1,088	944	134	137	24	27
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,095	1,970	374 ^a	284	91	66
White	1,757	1,689	305	242	67	48
Black or African American	162	117	32 ^a	13	10	7
American Indian or Alaska Native	22	23	2	3	1	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	80	54	12	7	4	1
Two or More Races	67	76	22	17	*	10
Hispanic or Latino	517	474	104	95	15	14
EDUCATION						
< High School	453	449	56	61	12	13
High School Graduate	921	791	189 ^b	104	40	25
Some College	879	834	159	146	40	29
College Graduate	360	369	74	68	14	14
CURRENT EMPLOYMENT						
Full-Time	1,074	1,054	139	115	35	22
Part-Time	707	668	148	123	30	26
Unemployed	391	294	104	53	27	17
Other ¹	441	428	86	89	14	14

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.46B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	7.5	7.0	1.4	1.1	0.3	0.2
GENDER						
Male	8.7	8.6	2.0 ^a	1.4	0.5	0.3
Female	6.3 ^a	5.4	0.8	0.8	0.1	0.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	7.6	7.1	1.4 ^a	1.0	0.3	0.2
White	9.0	8.7	1.6	1.2	0.3	0.2
Black or African American	3.2	2.3	0.6 ^a	0.3	0.2	0.1
American Indian or Alaska Native	9.4	11.2	1.0	1.5	0.5	0.3
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	4.3	2.7	0.7	0.4	0.2	0.0
Two or More Races	9.2	9.7	2.9	2.2	*	1.2
Hispanic or Latino	7.1	6.4	1.4	1.3	0.2	0.2
EDUCATION						
< High School	8.1	9.1	1.0	1.2	0.2	0.3
High School Graduate	7.7	6.6	1.6 ^b	0.9	0.3	0.2
Some College	7.2	6.7	1.3	1.2	0.3	0.2
College Graduate	7.1	6.6	1.5	1.2	0.3	0.3
CURRENT EMPLOYMENT						
Full-Time	7.9	7.4	1.0	0.8	0.3	0.2
Part-Time	7.3	7.1	1.5	1.3	0.3	0.3
Unemployed	9.4	7.6	2.5	1.4	0.6	0.4
Other ¹	5.9	5.9	1.2	1.2	0.2	0.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.47A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	17,129	17,543	575	725	269	316
GENDER						
Male	11,467	11,963	444	623	201	267
Female	5,661	5,580	130	103	68	50
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	15,526	15,895	527	614	249	226
White	14,184	14,316	482	486	211	181
Black or African American	498	714	19 ^a	82	14	20
American Indian or Alaska Native	100	138	1	2	*	2
Native Hawaiian or Other Pacific Islander	*	45	*	*	*	*
Asian	183	275	23	35	23	20
Two or More Races	486	408	2	8	*	3
Hispanic or Latino	1,603	1,648	48	111	21	90
EDUCATION						
< High School	1,411	1,572	34	63	19	18
High School Graduate	4,064	4,095	110	101	67	27
Some College	5,277	5,185	142	176	72	87
College Graduate	6,377	6,691	289	385	111	184
CURRENT EMPLOYMENT						
Full-Time	10,746	11,201	385	467	175	235
Part-Time	2,397	2,201	63	100	43	38
Unemployed	887	740	44	51	*	21
Other ¹	3,100	3,401	83	107	51	23

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.47B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8.4	8.5	0.3	0.4	0.1	0.2
GENDER						
Male	11.8	12.2	0.5	0.6	0.2	0.3
Female	5.4	5.2	0.1	0.1	0.1	0.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8.9	9.0	0.3	0.3	0.1	0.1
White	10.4	10.4	0.4	0.4	0.2	0.1
Black or African American	2.2	3.1	0.1	0.4	0.1	0.1
American Indian or Alaska Native	9.9	12.9	0.1	0.2	*	0.2
Native Hawaiian or Other Pacific Islander	*	6.1	*	*	*	*
Asian	1.8	2.6	0.2	0.3	0.2	0.2
Two or More Races	17.3	14.2	0.1	0.3	*	0.1
Hispanic or Latino	5.6	5.6	0.2	0.4	0.1	0.3
EDUCATION						
< High School	5.3	5.8	0.1	0.2	0.1	0.1
High School Graduate	7.0	7.2	0.2	0.2	0.1	0.0
Some College	10.3	9.8	0.3	0.3	0.1	0.2
College Graduate	9.5	9.8	0.4	0.6	0.2	0.3
CURRENT EMPLOYMENT						
Full-Time	10.2	10.3	0.4	0.4	0.2	0.2
Part-Time	10.0	9.3	0.3	0.4	0.2	0.2
Unemployed	11.6	10.7	0.6	0.7	*	0.3
Other ¹	4.7	5.1	0.1	0.2	0.1	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.48A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	19,741	19,986	1,053	1,104	375	397
GENDER						
Male	12,992	13,463	788	865	284	320
Female	6,749	6,524	265	239	92	77
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	17,621	17,865	901	898	340	292
White	15,941	16,005	787	728	278	229
Black or African American	661	831	51	95	25	26
American Indian or Alaska Native	121	161	3	5	1	3
Native Hawaiian or Other Pacific Islander	*	56	*	2	*	*
Asian	263	329	36	43	27	21
Two or More Races	553	484	24	26	8	13
Hispanic or Latino	2,120	2,121	152	207	35	105
EDUCATION						
< High School	1,864	2,022	89	124	31	31
High School Graduate	4,985	4,886	300	205	107	52
Some College	6,156	6,019	302	322	111	116
College Graduate	6,737	7,060	362	453	125	198
CURRENT EMPLOYMENT						
Full-Time	11,819	12,255	524	582	210	257
Part-Time	3,104	2,869	211	223	74	64
Unemployed	1,278	1,034	148	104	27	38
Other ¹	3,541	3,829	169	196	65	37

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.48B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8.3	8.3	0.4	0.5	0.2	0.2
GENDER						
Male	11.4	11.6	0.7	0.7	0.2	0.3
Female	5.5	5.2	0.2	0.2	0.1	0.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8.7	8.8	0.4	0.4	0.2	0.1
White	10.2	10.2	0.5	0.5	0.2	0.1
Black or African American	2.4	2.9	0.2	0.3	0.1	0.1
American Indian or Alaska Native	9.8	12.6	0.3	0.4	0.1	0.2
Native Hawaiian or Other Pacific Islander	*	6.1	*	0.2	*	*
Asian	2.2	2.6	0.3	0.3	0.2	0.2
Two or More Races	15.6	13.3	0.7	0.7	0.2	0.4
Hispanic or Latino	5.9	5.8	0.4	0.6	0.1	0.3
EDUCATION						
< High School	5.8	6.3	0.3	0.4	0.1	0.1
High School Graduate	7.1	7.1	0.4	0.3	0.2	0.1
Some College	9.7	9.2	0.5	0.5	0.2	0.2
College Graduate	9.4	9.6	0.5	0.6	0.2	0.3
CURRENT EMPLOYMENT						
Full-Time	9.9	10.0	0.4	0.5	0.2	0.2
Part-Time	9.2	8.7	0.6	0.7	0.2	0.2
Unemployed	10.8	9.6	1.3	1.0	0.2	0.4
Other ¹	4.9	5.2	0.2	0.3	0.1	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.49A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	53,172	54,395	15,348	14,966	6,484	6,537
AGE						
12-17	2,188	2,290	1,441	1,554	549 ^a	655
18-25	9,239	9,197	4,245	4,133	1,661	1,551
26 or Older	41,745	42,909	9,662	9,280	4,274	4,331
GENDER						
Male	27,490 ^a	28,978	7,772	7,852	3,340	3,382
Female	25,682	25,417	7,576	7,114	3,144	3,155
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	46,329	47,747	12,783	12,515	5,295	5,461
White	39,412	40,171	10,527	9,886	4,325	4,192
Black or African American	4,128	4,640	1,377 ^a	1,686	716	843
American Indian or Alaska Native	344	371	139	134	29 ^a	69
Native Hawaiian or Other Pacific Islander	212	196	43	85	10 ^a	64
Asian	1,110	1,305	313	397	110	153
Two or More Races	1,124	1,063	384	327	106	140
Hispanic or Latino	6,843	6,648	2,565	2,451	1,189	1,076

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.49B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	20.3	20.5	5.8	5.6	2.5	2.5
AGE						
12-17	8.8	9.2	5.8	6.2	2.2 ^a	2.6
18-25	26.6	26.3	12.2	11.8	4.8	4.4
26 or Older	20.6	20.9	4.8	4.5	2.1	2.1
GENDER						
Male	21.6	22.6	6.1	6.1	2.6	2.6
Female	19.0	18.6	5.6	5.2	2.3	2.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	21.0	21.4	5.8	5.6	2.4	2.5
White	23.2	23.6	6.2	5.8	2.5	2.5
Black or African American	13.2	14.6	4.4	5.3	2.3	2.7
American Indian or Alaska Native	24.5	25.7	9.9	9.3	2.1 ^a	4.8
Native Hawaiian or Other Pacific Islander	24.4	19.2	5.0	8.4	1.1 ^a	6.2
Asian	8.3	9.4	2.3	2.9	0.8	1.1
Two or More Races	26.2	24.1	9.0	7.4	2.5	3.2
Hispanic or Latino	16.6	15.7	6.2	5.8	2.9	2.5

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.50A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	2,188	2,290	1,441	1,554	549 ^a	655
GENDER						
Male	1,105	1,062	728	686	257	295
Female	1,083 ^a	1,227	713 ^b	867	292	360
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,700	1,739	1,131	1,178	429	496
White	1,214	1,211	834	839	293	336
Black or African American	306	377	188 ^a	247	102	117
American Indian or Alaska Native	23	20	18	15	3	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	48	46	22	28	5	17
Two or More Races	91	80	55	46	19	14
Hispanic or Latino	488	551	310	376	120	159
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	615	573	425	374	146	160
Female, White, Not Hispanic	599	638	409	465	147	176
Male, Black, Not Hispanic	147	158	85	96	49	46
Female, Black, Not Hispanic	159 ^a	219	103 ^a	151	54	71
Male, Hispanic	247	256	157	168	48	64
Female, Hispanic	241	295	153	208	71	94

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.50B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8.8	9.2	5.8	6.2	2.2 ^a	2.6
GENDER						
Male	8.7	8.4	5.7	5.4	2.0	2.3
Female	8.9 ^a	10.1	5.9 ^b	7.1	2.4	3.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8.8	9.0	5.9	6.1	2.2	2.6
White	8.9	9.0	6.1	6.2	2.2	2.5
Black or African American	8.9	10.8	5.5 ^a	7.1	3.0	3.4
American Indian or Alaska Native	14.2	12.1	11.2	8.8	1.9	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	3.9	3.7	1.7	2.3	0.4	1.4
Two or More Races	12.0	10.4	7.2	6.0	2.5	1.8
Hispanic or Latino	8.8	9.8	5.6	6.7	2.1	2.8
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	8.8	8.3	6.1	5.4	2.1	2.3
Female, White, Not Hispanic	9.0	9.7	6.1	7.0	2.2	2.7
Male, Black, Not Hispanic	8.4	9.0	4.9	5.4	2.8	2.6
Female, Black, Not Hispanic	9.4 ^a	12.7	6.1 ^a	8.8	3.2	4.1
Male, Hispanic	8.7	8.9	5.5	5.9	1.7	2.2
Female, Hispanic	8.8	10.6	5.6	7.5	2.6	3.4

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.51A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	9,239	9,197	4,245	4,133	1,661	1,551
GENDER						
Male	4,958	4,806	2,358	2,243	884	811
Female	4,280	4,390	1,886	1,890	777	740
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	7,605	7,609	3,534	3,430	1,398	1,266
White	6,108	5,958	2,864	2,633	1,109	989
Black or African American	912	1,023	408	499	194	157
American Indian or Alaska Native	64	72	28	35	17	17
Native Hawaiian or Other Pacific Islander	41	*	16	*	2	*
Asian	255	267	113	126	41	44
Two or More Races	224	231	105	100	36	33
Hispanic or Latino	1,634	1,588	711	703	263	285
EDUCATION						
< High School	1,585	1,390	710	630	331 ^a	248
High School Graduate	3,107	2,964	1,432	1,340	571	522
Some College	3,210	3,285	1,508	1,559	535	557
College Graduate	1,337	1,557	596	604	224	224
CURRENT EMPLOYMENT						
Full-Time	3,877	4,160	1,621	1,774	649	700
Part-Time	2,489	2,377	1,274	1,182	454	391
Unemployed	1,230	1,076	606	493	263	196
Other ¹	1,643	1,583	743	684	295	264

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.51B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	26.6	26.3	12.2	11.8	4.8	4.4
GENDER						
Male	28.4	27.4	13.5	12.8	5.1	4.6
Female	24.7	25.2	10.9	10.9	4.5	4.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	27.6	27.6	12.8	12.4	5.1	4.6
White	31.3	30.8	14.7	13.6	5.7	5.1
Black or African American	18.2	20.3	8.2	9.9	3.9	3.1
American Indian or Alaska Native	27.6	35.4	12.2	17.5	7.2	8.6
Native Hawaiian or Other Pacific Islander	20.6	*	7.8	*	0.9	*
Asian	13.7	13.3	6.0	6.3	2.2	2.2
Two or More Races	30.6	29.4	14.4	12.8	4.9	4.3
Hispanic or Latino	22.5	21.6	9.8	9.5	3.6	3.9
EDUCATION						
< High School	28.4	28.1	12.7	12.7	5.9	5.0
High School Graduate	26.0	24.8	12.0	11.2	4.8	4.4
Some College	26.4	26.4	12.4	12.5	4.4	4.5
College Graduate	26.5	27.9	11.8	10.8	4.4	4.0
CURRENT EMPLOYMENT						
Full-Time	28.7	29.2	12.0	12.4	4.8	4.9
Part-Time	25.7	25.1	13.2	12.5	4.7	4.1
Unemployed	29.6	27.7	14.6	12.7	6.3	5.0
Other ¹	22.1	21.6	10.0	9.4	4.0	3.6

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.52A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	41,745	42,909	9,662	9,280	4,274	4,331
GENDER						
Male	21,427 ^a	23,109	4,686	4,923	2,200	2,276
Female	20,318	19,799	4,976	4,357	2,075	2,055
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	37,024	38,400	8,118	7,908	3,468	3,699
White	32,089	33,002	6,829	6,415	2,923	2,867
Black or African American	2,910	3,241	781	940	419	569
American Indian or Alaska Native	257	279	93	83	9 ^a	42
Native Hawaiian or Other Pacific Islander	*	133	12	46	1	*
Asian	806	992	179	243	64	92
Two or More Races	809	753	224	180	51	92
Hispanic or Latino	4,721	4,509	1,544	1,372	807	632
EDUCATION						
< High School	4,284	4,927	1,378	1,624	827	842
High School Graduate	10,639	11,164	2,591	2,250	1,252	1,094
Some College	12,617	12,895	2,902	2,644	1,209	1,334
College Graduate	14,205	13,922	2,791	2,762	986	1,061
CURRENT EMPLOYMENT						
Full-Time	23,833 ^a	25,715	5,068	5,328	1,978	2,297
Part-Time	6,264 ^a	5,229	1,477	1,121	668	529
Unemployed	2,145	1,970	780	555	385	277
Other ¹	9,504	9,995	2,338	2,276	1,244	1,227

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.52B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	20.6	20.9	4.8	4.5	2.1	2.1
GENDER						
Male	22.1 ^a	23.5	4.8	5.0	2.3	2.3
Female	19.2	18.5	4.7 ^a	4.1	2.0	1.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	21.2	21.8	4.7	4.5	2.0	2.1
White	23.4	24.0	5.0	4.7	2.1	2.1
Black or African American	12.8	14.0	3.4	4.1	1.8	2.5
American Indian or Alaska Native	25.4	26.0	9.2	7.8	0.9 ^a	3.9
Native Hawaiian or Other Pacific Islander	*	18.0	2.2	6.1	0.2	*
Asian	7.9	9.3	1.7	2.3	0.6	0.9
Two or More Races	28.8	26.3	8.0	6.3	1.8	3.2
Hispanic or Latino	16.6	15.3	5.4	4.7	2.8	2.1
EDUCATION						
< High School	16.1	18.1	5.2	6.0	3.1	3.1
High School Graduate	18.3	19.6	4.5	4.0	2.2	1.9
Some College	24.6	24.3	5.7	5.0	2.4	2.5
College Graduate	21.2	20.5	4.2	4.1	1.5	1.6
CURRENT EMPLOYMENT						
Full-Time	22.6	23.7	4.8	4.9	1.9	2.1
Part-Time	26.0 ^b	22.2	6.1	4.8	2.8	2.2
Unemployed	28.0	28.5	10.2	8.0	5.0	4.0
Other ¹	14.5	15.1	3.6	3.4	1.9	1.9

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.53A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	50,984	52,105	13,907	13,412	5,935	5,882
GENDER						
Male	26,385 ^a	27,916	7,044	7,165	3,083	3,087
Female	24,599	24,190	6,862	6,247	2,852	2,795
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	44,629	46,009	11,652	11,337	4,866	4,965
White	38,198	38,960	9,693	9,047	4,032	3,856
Black or African American	3,823	4,263	1,189	1,440	613	726
American Indian or Alaska Native	321	351	121	119	26 ^a	60
Native Hawaiian or Other Pacific Islander	*	191	28 ^a	82	3 ^a	61
Asian	1,061	1,259	292	369	105	136
Two or More Races	1,033	984	330	280	87	126
Hispanic or Latino	6,355	6,097	2,254	2,075	1,070	917
EDUCATION						
< High School	5,869	6,317	2,088	2,254	1,158	1,090
High School Graduate	13,745	14,129	4,023	3,591	1,823	1,616
Some College	15,827	16,180	4,409	4,203	1,744	1,892
College Graduate	15,543	15,479	3,386	3,365	1,210	1,285
CURRENT EMPLOYMENT						
Full-Time	27,710 ^b	29,875	6,689	7,102	2,627	2,997
Part-Time	8,753 ^b	7,606	2,751 ^a	2,303	1,122	921
Unemployed	3,374	3,046	1,386 ^a	1,047	647	473
Other ¹	11,147	11,578	3,081	2,960	1,539	1,491

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.53B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	21.5	21.7	5.9	5.6	2.5	2.4
GENDER						
Male	23.1	24.1	6.2	6.2	2.7	2.7
Female	20.0	19.4	5.6 ^a	5.0	2.3	2.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	22.1	22.6	5.8	5.6	2.4	2.4
White	24.4	24.9	6.2	5.8	2.6	2.5
Black or African American	13.8	15.1	4.3	5.1	2.2	2.6
American Indian or Alaska Native	25.8	27.5	9.7	9.3	2.1 ^a	4.7
Native Hawaiian or Other Pacific Islander	*	20.9	3.6	8.9	0.4 ^a	6.6
Asian	8.8	10.0	2.4	2.9	0.9	1.1
Two or More Races	29.2	26.9	9.3	7.7	2.5	3.4
Hispanic or Latino	17.8	16.6	6.3	5.6	3.0	2.5
EDUCATION						
< High School	18.2	19.6	6.5	7.0	3.6	3.4
High School Graduate	19.6	20.5	5.7	5.2	2.6	2.3
Some College	25.0	24.7	7.0	6.4	2.8	2.9
College Graduate	21.6	21.1	4.7	4.6	1.7	1.7
CURRENT EMPLOYMENT						
Full-Time	23.3	24.3	5.6	5.8	2.2	2.4
Part-Time	25.9 ^b	23.0	8.2 ^a	7.0	3.3	2.8
Unemployed	28.6	28.2	11.7	9.7	5.5	4.4
Other ¹	15.3	15.8	4.2	4.0	2.1	2.0

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.54A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	35,473	36,064	11,082	10,337	4,521	4,325
AGE						
12-17	1,805	1,803	1,157	1,165	425	467
18-25	7,251	7,049	3,061 ^a	2,740	1,132	978
26 or Older	26,417	27,211	6,865	6,432	2,964	2,879
GENDER						
Male	18,918 ^a	20,356	5,834	5,585	2,398	2,369
Female	16,555	15,709	5,248	4,752	2,123	1,956
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	30,514	31,458	9,223	8,633	3,733	3,568
White	25,502	25,907	7,380 ^a	6,618	2,968	2,612
Black or African American	3,108	3,548	1,120	1,384	573	691
American Indian or Alaska Native	230	253	97	81	19	26
Native Hawaiian or Other Pacific Islander	177	157	34	70	9	53
Asian	716	839	244	246	74	86
Two or More Races	780	754	349	236	90	101
Hispanic or Latino	4,959	4,607	1,859	1,703	788	757

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.54B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	13.5	13.6	4.2 ^a	3.9	1.7	1.6
AGE						
12-17	7.3	7.3	4.6	4.7	1.7	1.9
18-25	20.8	20.2	8.8 ^a	7.8	3.3	2.8
26 or Older	13.0	13.3	3.4	3.1	1.5	1.4
GENDER						
Male	14.9 ^a	15.9	4.6	4.3	1.9	1.8
Female	12.2	11.5	3.9	3.5	1.6	1.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	13.8	14.1	4.2	3.9	1.7	1.6
White	15.0	15.2	4.3 ^a	3.9	1.7	1.5
Black or African American	10.0	11.2	3.6	4.4	1.8	2.2
American Indian or Alaska Native	16.4	17.6	6.9	5.6	1.4	1.8
Native Hawaiian or Other Pacific Islander	20.4	15.3	3.9	6.8	1.1	5.2
Asian	5.4	6.1	1.8	1.8	0.6	0.6
Two or More Races	18.2	17.1	8.1	5.3	2.1	2.3
Hispanic or Latino	12.0	10.8	4.5	4.0	1.9	1.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.55A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,805	1,803	1,157	1,165	425	467
GENDER						
Male	902	856	575	508	189	205
Female	904	948	582	657	236	262
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,410	1,368	916	881	332	354
White	988	928	664	611	220	228
Black or African American	268	322	160	203	86	93
American Indian or Alaska Native	20	18	16	13	3	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	37	34	18	17	3	9
Two or More Races	79	63	46	37	14	13
Hispanic or Latino	395	435	241	283	93	113
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	511	470	340	277	109	107
Female, White, Not Hispanic	477	458	324	334	111	121
Male, Black, Not Hispanic	125	131	65	71	37	34
Female, Black, Not Hispanic	143	190	94	132	49	59
Male, Hispanic	184	202	114	129	34	48
Female, Hispanic	211	233	126	155	59	65

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.55B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	7.3	7.3	4.6	4.7	1.7	1.9
GENDER						
Male	7.1	6.7	4.5	4.0	1.5	1.6
Female	7.4	7.8	4.8	5.4	1.9	2.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	7.3	7.1	4.7	4.6	1.7	1.8
White	7.2	6.9	4.9	4.5	1.6	1.7
Black or African American	7.8	9.2	4.6	5.8	2.5	2.7
American Indian or Alaska Native	12.6	10.7	9.9	7.7	1.9	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	3.0	2.8	1.4	1.4	0.3	0.7
Two or More Races	10.5	8.2	6.1	4.8	1.8	1.7
Hispanic or Latino	7.1	7.7	4.3	5.0	1.7	2.0
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	7.3	6.8	4.9	4.0	1.6	1.6
Female, White, Not Hispanic	7.1	6.9	4.9	5.1	1.7	1.8
Male, Black, Not Hispanic	7.1	7.4	3.7	4.0	2.1	1.9
Female, Black, Not Hispanic	8.5	11.1	5.6	7.7	2.9	3.4
Male, Hispanic	6.5	7.0	4.0	4.5	1.2	1.7
Female, Hispanic	7.8	8.4	4.6	5.6	2.2	2.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.56A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	7,251	7,049	3,061 ^a	2,740	1,132	978
GENDER						
Male	4,020	3,881	1,759 ^a	1,533	631	529
Female	3,231	3,169	1,302	1,207	501	450
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5,973	5,834	2,533 ^a	2,280	941 ^a	777
White	4,743	4,533	1,997 ^a	1,720	710	581
Black or African American	770	849	335	383	160	119
American Indian or Alaska Native	56	55	24	18	13	8
Native Hawaiian or Other Pacific Islander	35	*	15	*	1	*
Asian	170	172	71	70	29	23
Two or More Races	198	172	91	64	27	27
Hispanic or Latino	1,278	1,216	527	460	191	201
EDUCATION						
< High School	1,311	1,162	565	507	237	189
High School Graduate	2,578	2,375	1,164 ^a	961	457	370
Some College	2,454	2,513	1,007	1,002	328	328
College Graduate	909	999	325	270	110	91
CURRENT EMPLOYMENT						
Full-Time	3,084	3,234	1,193	1,168	455	466
Part-Time	1,913	1,759	869	748	298 ^a	195
Unemployed	983	858	486 ^a	357	185	142
Other ¹	1,271	1,199	512	467	194	176

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.56B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	20.8	20.2	8.8 ^a	7.8	3.3	2.8
GENDER						
Male	23.0	22.1	10.1 ^a	8.7	3.6	3.0
Female	18.6	18.2	7.5	6.9	2.9	2.6
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	21.7	21.2	9.2 ^a	8.3	3.4 ^a	2.8
White	24.3	23.4	10.2 ^a	8.9	3.6 ^a	3.0
Black or African American	15.4	16.9	6.7	7.6	3.2	2.4
American Indian or Alaska Native	24.4	27.0	10.6	8.8	5.6	3.8
Native Hawaiian or Other Pacific Islander	17.6	*	7.6	*	0.6	*
Asian	9.1	8.6	3.8	3.5	1.5	1.2
Two or More Races	27.1	22.0	12.5	8.2	3.7	3.5
Hispanic or Latino	17.6	16.5	7.3	6.2	2.6	2.7
EDUCATION						
< High School	23.5	23.5	10.1	10.3	4.2	3.8
High School Graduate	21.5	19.8	9.7 ^a	8.0	3.8	3.1
Some College	20.1	20.2	8.3	8.1	2.7	2.6
College Graduate	18.0	17.9	6.4	4.8	2.2	1.6
CURRENT EMPLOYMENT						
Full-Time	22.8	22.7	8.8	8.2	3.4	3.3
Part-Time	19.8	18.6	9.0	7.9	3.1 ^a	2.1
Unemployed	23.6	22.1	11.7 ^a	9.2	4.4	3.6
Other ¹	17.1	16.4	6.9	6.4	2.6	2.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.57A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	26,417	27,211	6,865	6,432	2,964	2,879
GENDER						
Male	13,997 ^b	15,620	3,501	3,545	1,578	1,635
Female	12,421	11,592	3,364	2,888	1,386	1,244
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	23,131	24,255	5,774	5,472	2,460	2,437
White	19,771	20,446	4,719	4,288	2,037	1,803
Black or African American	2,070	2,378	626	798	327	479
American Indian or Alaska Native	154	181	57	50	3	8
Native Hawaiian or Other Pacific Islander	*	99	6	42	1	*
Asian	509	633	155	159	42	54
Two or More Races	503	519	211	135	49	61
Hispanic or Latino	3,286	2,956	1,091	960	504	443
EDUCATION						
< High School	2,860 ^b	3,625	996	1,309	576	668
High School Graduate	7,249	7,459	2,030	1,702	907	744
Some College	8,170	8,337	2,038	1,817	883	886
College Graduate	8,138	7,791	1,801	1,605	598	582
CURRENT EMPLOYMENT						
Full-Time	15,230 ^a	16,748	3,677	3,728	1,424	1,569
Part-Time	3,853 ^a	3,080	937	738	386	333
Unemployed	1,529	1,476	591	419	311	214
Other ¹	5,806	5,908	1,659	1,548	843	763

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.57B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	13.0	13.3	3.4	3.1	1.5	1.4
GENDER						
Male	14.4 ^a	15.9	3.6	3.6	1.6	1.7
Female	11.7	10.8	3.2	2.7	1.3	1.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	13.3	13.8	3.3	3.1	1.4	1.4
White	14.4	14.9	3.4	3.1	1.5	1.3
Black or African American	9.1	10.3	2.7	3.4	1.4	2.1
American Indian or Alaska Native	15.2	16.8	5.6	4.7	0.3	0.8
Native Hawaiian or Other Pacific Islander	*	13.4	1.0	5.7	0.2	*
Asian	5.0	6.0	1.5	1.5	0.4	0.5
Two or More Races	17.9	18.1	7.5	4.7	1.7	2.1
Hispanic or Latino	11.6	10.0	3.8	3.3	1.8	1.5
EDUCATION						
< High School	10.8 ^a	13.3	3.7	4.8	2.2	2.4
High School Graduate	12.5	13.1	3.5	3.0	1.6	1.3
Some College	16.0	15.7	4.0	3.4	1.7	1.7
College Graduate	12.2	11.5	2.7	2.4	0.9	0.9
CURRENT EMPLOYMENT						
Full-Time	14.4	15.4	3.5	3.4	1.4	1.4
Part-Time	16.0 ^a	13.1	3.9	3.1	1.6	1.4
Unemployed	20.0	21.4	7.7	6.1	4.1	3.1
Other ¹	8.9	8.9	2.5	2.3	1.3	1.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.58A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	33,668	34,261	9,925	9,172	4,096	3,858
GENDER						
Male	18,017 ^a	19,500	5,259	5,077	2,209	2,164
Female	15,651	14,761	4,666	4,095	1,887	1,694
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	29,104	30,089	8,307	7,752	3,401	3,214
White	24,514	24,979	6,716 ^a	6,007	2,748	2,384
Black or African American	2,840	3,226	961	1,181	488	598
American Indian or Alaska Native	210	235	81	68	16	16
Native Hawaiian or Other Pacific Islander	*	153	21	68	2 ^a	51
Asian	679	805	226	229	70	77
Two or More Races	701	691	303	199	76	88
Hispanic or Latino	4,564	4,172	1,618	1,420	695	644
EDUCATION						
< High School	4,170 ^a	4,787	1,561	1,816	814	857
High School Graduate	9,827	9,834	3,193 ^a	2,663	1,364	1,114
Some College	10,623	10,849	3,045	2,819	1,210	1,214
College Graduate	9,047	8,790	2,126	1,875	708	673
CURRENT EMPLOYMENT						
Full-Time	18,314 ^a	19,981	4,870	4,896	1,880	2,036
Part-Time	5,766 ^b	4,839	1,806	1,486	684	528
Unemployed	2,511	2,333	1,078 ^b	775	495	356
Other ¹	7,077	7,107	2,172	2,014	1,037	938

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.58B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	14.2	14.3	4.2 ^a	3.8	1.7	1.6
GENDER						
Male	15.7 ^a	16.8	4.6	4.4	1.9	1.9
Female	12.7	11.9	3.8 ^a	3.3	1.5	1.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	14.4	14.8	4.1	3.8	1.7	1.6
White	15.7	15.9	4.3 ^a	3.8	1.8	1.5
Black or African American	10.2	11.4	3.5	4.2	1.8	2.1
American Indian or Alaska Native	16.9	18.5	6.5	5.3	1.3	1.3
Native Hawaiian or Other Pacific Islander	*	16.6	2.7	7.4	0.3 ^a	5.6
Asian	5.6	6.4	1.9	1.8	0.6	0.6
Two or More Races	19.8	18.9	8.6	5.5	2.2	2.4
Hispanic or Latino	12.8	11.3	4.5	3.9	1.9	1.7
EDUCATION						
< High School	13.0 ^a	14.9	4.9	5.6	2.5	2.7
High School Graduate	14.0	14.3	4.6 ^a	3.9	1.9	1.6
Some College	16.8	16.5	4.8	4.3	1.9	1.9
College Graduate	12.6	12.0	3.0	2.5	1.0	0.9
CURRENT EMPLOYMENT						
Full-Time	15.4	16.3	4.1	4.0	1.6	1.7
Part-Time	17.1 ^b	14.7	5.4	4.5	2.0	1.6
Unemployed	21.3	21.6	9.1 ^a	7.2	4.2	3.3
Other ¹	9.7	9.7	3.0	2.7	1.4	1.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.59A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	127,458 ^a	130,332	41,591 ^b	44,157	24,573 ^b	26,983
GEOGRAPHIC DIVISION						
Northeast	22,722 ^a	24,072	7,461 ^b	8,668	4,362 ^b	5,364
New England	6,509	7,027	2,230 ^a	2,651	1,397 ^a	1,741
Middle Atlantic	16,212	17,045	5,232 ^a	6,016	2,965 ^b	3,623
Midwest	27,289	28,300	8,466	8,830	4,914	5,209
East North Central	19,168 ^a	20,168	6,141	6,299	3,629	3,796
West North Central	8,121	8,132	2,324	2,531	1,285	1,413
South	45,007	46,098	13,976	14,823	8,066	8,811
South Atlantic	25,276	25,637	8,100	8,260	4,584	5,040
East South Central	6,565	6,810	2,005	2,176	1,220	1,242
West South Central	13,167	13,651	3,871 ^a	4,387	2,261	2,529
West	32,440	31,862	11,688	11,836	7,232	7,600
Mountain	10,042	9,942	3,493	3,484	2,124	2,211
Pacific	22,398	21,920	8,195	8,352	5,108	5,389
COUNTY TYPE						
Large Metro	69,365 ^a	73,045	22,999 ^b	25,484	13,565 ^b	15,553
Small Metro	39,350	38,489	13,018	12,799	7,694	7,897
250K – 1 Mil. Pop.	27,024	26,525	8,646	8,738	5,064	5,384
< 250K Pop.	12,326	11,964	4,372	4,062	2,630	2,513
Nonmetro	18,743	18,798	5,574	5,874	3,315	3,534
Urbanized	8,814	8,587	2,859	2,802	1,696	1,662
Less Urbanized	8,227	8,674	2,258	2,714	1,311	1,660
Completely Rural	1,702	1,538	457	358	308	212

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.59B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	48.6	49.2	15.9 ^a	16.7	9.4 ^b	10.2
GEOGRAPHIC DIVISION						
Northeast	47.9 ^a	50.5	15.7 ^b	18.2	9.2 ^b	11.3
New England	52.1	55.9	17.9 ^a	21.1	11.2 ^a	13.9
Middle Atlantic	46.4	48.6	15.0 ^a	17.2	8.5 ^b	10.3
Midwest	48.5	50.1	15.1	15.6	8.7	9.2
East North Central	49.2 ^a	51.6	15.8	16.1	9.3	9.7
West North Central	47.1	46.9	13.5	14.6	7.5	8.1
South	46.2	46.6	14.3	15.0	8.3	8.9
South Atlantic	49.1	49.1	15.7	15.8	8.9	9.7
East South Central	42.4	43.7	12.9	14.0	7.9	8.0
West South Central	43.1	43.9	12.7	14.1	7.4	8.1
West	52.9	51.2	19.1	19.0	11.8	12.2
Mountain	53.7	52.3	18.7	18.3	11.4	11.6
Pacific	52.6	50.8	19.2	19.3	12.0	12.5
COUNTY TYPE						
Large Metro	49.2	49.8	16.3 ^a	17.4	9.6 ^b	10.6
Small Metro	49.9	49.7	16.5	16.5	9.8	10.2
250K – 1 Mil. Pop.	50.6	50.6	16.2	16.7	9.5	10.3
< 250K Pop.	48.3	47.8	17.1	16.2	10.3	10.0
Nonmetro	44.2	45.7	13.1	14.3	7.8	8.6
Urbanized	46.4	49.8	15.0	16.2	8.9	9.6
Less Urbanized	43.3	44.0	11.9	13.8	6.9	8.4
Completely Rural	38.1	36.5	10.2	8.5	6.9	5.0

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.60A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	5,794	5,791	4,287	4,336	2,197	2,338
GEOGRAPHIC DIVISION						
Northeast	946	921	758	732	391	377
New England	253	264	204	216	113	96
Middle Atlantic	693	657	554	516	278	281
Midwest	1,171	1,149	879	854	445	435
East North Central	848	837	649	641	324	341
West North Central	322	312	231	213	121	93
South	2,163	2,193	1,517	1,585	739	849
South Atlantic	1,170	1,111	830	834	393	417
East South Central	308	286	230	196	111	111
West South Central	686 ^a	795	457 ^a	554	236 ^a	321
West	1,514	1,528	1,132	1,166	622	677
Mountain	508	480	375	365	217	203
Pacific	1,006	1,048	757	801	405	474
COUNTY TYPE						
Large Metro	3,136	3,218	2,371	2,427	1,223	1,338
Small Metro	1,785	1,747	1,319	1,294	665	670
250K – 1 Mil. Pop.	1,210	1,219	896	913	454	478
< 250K Pop.	575	528	423	381	211	192
Nonmetro	872	826	597	614	309	329
Urbanized	374	359	271	268	142	149
Less Urbanized	423	397	275	300	141	155
Completely Rural	75	69	51	46	26	25

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.60B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	23.3	23.3	17.2	17.4	8.8	9.4
GEOGRAPHIC DIVISION						
Northeast	22.6	22.2	18.1	17.6	9.3	9.1
New England	23.2	24.3	18.7	19.9	10.4	8.8
Middle Atlantic	22.4	21.4	17.9	16.8	9.0	9.1
Midwest	21.7	21.4	16.3	15.9	8.2	8.1
East North Central	22.6	22.4	17.3	17.2	8.6	9.1
West North Central	19.7	19.0	14.1	13.0	7.3	5.7
South	23.1	23.3	16.2	16.8	7.9	9.0
South Atlantic	25.1	23.7	17.8	17.8	8.4	8.9
East South Central	20.9	19.4	15.6	13.3	7.5	7.6
West South Central	21.3	24.4	14.2 ^a	17.0	7.3 ^a	9.9
West	25.4	25.7	19.0	19.6	10.5	11.4
Mountain	27.1	25.4	20.0	19.3	11.6	10.7
Pacific	24.7	25.9	18.6	19.8	9.9	11.7
COUNTY TYPE						
Large Metro	23.1	22.9	17.5	17.3	9.0	9.5
Small Metro	24.0	24.7	17.7	18.3	8.9	9.5
250K – 1 Mil. Pop.	23.6	25.1	17.5	18.8	8.9	9.8
< 250K Pop.	24.7	23.9	18.2	17.3	9.1	8.7
Nonmetro	22.4	21.9	15.4	16.3	8.0	8.7
Urbanized	22.5	23.4	16.3	17.4	8.5	9.7
Less Urbanized	23.7	21.2	15.4	16.0	7.9	8.3
Completely Rural	16.9	18.9	11.5	12.5	6.0	6.9

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.61A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	19,826	20,217	12,448	12,597	7,490	7,676
GEOGRAPHIC DIVISION						
Northeast	3,587	3,609	2,361	2,467	1,425	1,522
New England	1,083	1,081	790	776	491	517
Middle Atlantic	2,504	2,529	1,571	1,691	935	1,005
Midwest	4,311	4,463	2,648	2,755	1,602	1,611
East North Central	2,954	3,094	1,872	1,905	1,114	1,156
West North Central	1,357	1,369	776	850	488	455
South	6,955	7,191	4,160	4,324	2,412	2,578
South Atlantic	3,669	3,796	2,277	2,330	1,323	1,430
East South Central	1,059	1,074	614	629	365	352
West South Central	2,227	2,322	1,269	1,365	724	797
West	4,973	4,953	3,280	3,052	2,051	1,965
Mountain	1,496	1,483	918	920	582	600
Pacific	3,476	3,470	2,362	2,132	1,470	1,364
COUNTY TYPE						
Large Metro	10,832	11,240	7,071	7,189	4,344	4,393
Small Metro	6,235	6,112	3,778	3,748	2,200	2,296
250K – 1 Mil. Pop.	4,031	3,946	2,441	2,439	1,399	1,504
< 250K Pop.	2,203	2,165	1,337	1,308	800	792
Nonmetro	2,760	2,865	1,599	1,661	947	988
Urbanized	1,341	1,401	807	855	499	491
Less Urbanized	1,207	1,293	680	710	392	447
Completely Rural	211	171	112	96	56	50

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.61B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	57.0	57.9	35.8	36.1	21.5	22.0
GEOGRAPHIC DIVISION						
Northeast	58.3	58.7	38.4	40.1	23.2	24.8
New England	66.5	65.7	48.5	47.2	30.1	31.5
Middle Atlantic	55.4	56.1	34.8	37.5	20.7	22.3
Midwest	58.2	60.1	35.7	37.1	21.6	21.7
East North Central	57.9	60.6	36.7	37.3	21.8	22.6
West North Central	58.9	59.0	33.7	36.7	21.2	19.6
South	54.1	55.6	32.4	33.4	18.8	19.9
South Atlantic	55.9	57.6	34.7	35.3	20.2	21.7
East South Central	51.9	52.4	30.1	30.7	17.9	17.2
West South Central	52.3	54.0	29.8	31.7	17.0	18.5
West	59.4	58.9	39.2	36.3	24.5	23.4
Mountain	59.1	58.0	36.3	36.0	23.0	23.5
Pacific	59.5	59.3	40.4	36.4	25.2	23.3
COUNTY TYPE						
Large Metro	57.8	58.6	37.7	37.5	23.2	22.9
Small Metro	57.3	57.5	34.8	35.2	20.2	21.6
250K – 1 Mil. Pop.	57.7	57.4	34.9	35.5	20.0	21.9
< 250K Pop.	56.8	57.6	34.5	34.8	20.6	21.1
Nonmetro	53.4	55.9	30.9	32.4	18.3	19.3
Urbanized	54.2	58.5	32.6	35.7	20.2	20.5
Less Urbanized	54.3	55.2	30.6	30.3	17.6	19.1
Completely Rural	44.6	44.1	23.7	24.6	11.8	12.8

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.62A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	101,838	104,324	24,856 ^b	27,223	14,886 ^b	16,969
GEOGRAPHIC DIVISION						
Northeast	18,188 ^a	19,542	4,342 ^b	5,469	2,546 ^b	3,465
New England	5,172	5,682	1,236 ^a	1,659	793 ^a	1,128
Middle Atlantic	13,016	13,860	3,107 ^a	3,810	1,753 ^a	2,337
Midwest	21,807	22,688	4,939	5,222	2,867	3,163
East North Central	15,365 ^a	16,236	3,621	3,753	2,190	2,298
West North Central	6,442	6,451	1,318	1,469	677	864
South	35,889	36,714	8,299	8,915	4,915	5,384
South Atlantic	20,437	20,730	4,993	5,096	2,869	3,194
East South Central	5,198	5,450	1,160	1,352	745	779
West South Central	10,254	10,533	2,146	2,467	1,302	1,411
West	25,953	25,381	7,276	7,618	4,558	4,958
Mountain	8,038	7,979	2,200	2,199	1,325	1,407
Pacific	17,916	17,402	5,076	5,419	3,233	3,551
COUNTY TYPE						
Large Metro	55,396 ^a	58,586	13,557 ^b	15,867	7,998 ^b	9,822
Small Metro	31,330	30,630	7,921	7,757	4,829	4,930
250K – 1 Mil. Pop.	21,783	21,359	5,309	5,385	3,210	3,401
< 250K Pop.	9,548	9,271	2,611	2,372	1,619	1,529
Nonmetro	15,111	15,107	3,378	3,599	2,059	2,217
Urbanized	7,099	6,826	1,781	1,679	1,056	1,022
Less Urbanized	6,597	6,984	1,303	1,703	778	1,059
Completely Rural	1,416	1,297	294	217	225	136

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.62B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	50.2	50.8	12.3 ^a	13.3	7.3 ^b	8.3
GEOGRAPHIC DIVISION						
Northeast	49.1 ^a	52.4	11.7 ^b	14.7	6.9 ^b	9.3
New England	53.0	57.8	12.7 ^a	16.9	8.1 ^a	11.5
Middle Atlantic	47.7	50.4	11.4 ^a	13.9	6.4 ^a	8.5
Midwest	50.2	52.0	11.4	12.0	6.6	7.2
East North Central	51.0	53.6	12.0	12.4	7.3	7.6
West North Central	48.5	48.2	9.9	11.0	5.1	6.5
South	47.7	48.0	11.0	11.7	6.5	7.0
South Atlantic	50.8	50.7	12.4	12.5	7.1	7.8
East South Central	43.4	45.2	9.7	11.2	6.2	6.5
West South Central	44.4	44.8	9.3	10.5	5.6	6.0
West	55.3	53.1	15.5	15.9	9.7	10.4
Mountain	56.2	54.8	15.4	15.1	9.3	9.7
Pacific	54.9	52.3	15.5	16.3	9.9	10.7
COUNTY TYPE						
Large Metro	50.9	51.7	12.5 ^b	14.0	7.4 ^b	8.7
Small Metro	51.7	51.3	13.1	13.0	8.0	8.3
250K – 1 Mil. Pop.	52.8	52.6	12.9	13.3	7.8	8.4
< 250K Pop.	49.5	48.6	13.5	12.4	8.4	8.0
Nonmetro	45.2	46.8	10.1	11.2	6.2	6.9
Urbanized	47.8	51.3	12.0	12.6	7.1	7.7
Less Urbanized	44.0	45.1	8.7	11.0	5.2	6.8
Completely Rural	39.9	37.5	8.3	6.3	6.4	3.9

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.63A Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	121,664 ^a	124,541	37,304 ^b	39,821	22,376 ^b	24,646
GEOGRAPHIC DIVISION						
Northeast	21,775 ^a	23,151	6,703 ^b	7,935	3,971 ^b	4,987
New England	6,256	6,763	2,026 ^a	2,435	1,283 ^a	1,645
Middle Atlantic	15,519	16,388	4,677 ^b	5,500	2,688 ^b	3,342
Midwest	26,118	27,151	7,586	7,977	4,469	4,774
East North Central	18,320 ^a	19,331	5,493	5,658	3,304	3,455
West North Central	7,798	7,820	2,094	2,319	1,164	1,319
South	42,844	43,905	12,459	13,238	7,327	7,962
South Atlantic	24,106	24,526	7,269	7,426	4,191	4,624
East South Central	6,257	6,524	1,775	1,980	1,110	1,131
West South Central	12,481	12,855	3,414	3,832	2,026	2,208
West	30,926	30,334	10,556	10,670	6,610	6,922
Mountain	9,534	9,462	3,118	3,119	1,907	2,007
Pacific	21,392	20,872	7,438	7,551	4,703	4,915
COUNTY TYPE						
Large Metro	66,228 ^a	69,827	20,627 ^b	23,056	12,342 ^b	14,215
Small Metro	37,565	36,742	11,699	11,505	7,028	7,226
250K – 1 Mil. Pop.	25,814	25,305	7,751	7,824	4,609	4,906
< 250K Pop.	11,751	11,437	3,949	3,680	2,419	2,320
Nonmetro	17,871	17,972	4,977	5,260	3,006	3,205
Urbanized	8,441	8,228	2,588	2,534	1,554	1,513
Less Urbanized	7,804	8,277	1,983	2,414	1,170	1,506
Completely Rural	1,627	1,468	406	313	281	186

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.63B Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	51.2	51.8	15.7 ^a	16.6	9.4 ^b	10.3
GEOGRAPHIC DIVISION						
Northeast	50.4 ^a	53.3	15.5 ^b	18.3	9.2 ^b	11.5
New England	54.9	58.9	17.8	21.2	11.3 ^a	14.3
Middle Atlantic	48.8	51.2	14.7 ^a	17.2	8.4 ^b	10.4
Midwest	51.4	53.1	14.9	15.6	8.8	9.3
East North Central	52.0 ^a	54.6	15.6	16.0	9.4	9.8
West North Central	50.0	49.8	13.4	14.8	7.5	8.4
South	48.6	49.1	14.1	14.8	8.3	8.9
South Atlantic	51.5	51.7	15.5	15.6	9.0	9.7
East South Central	44.6	46.2	12.7	14.0	7.9	8.0
West South Central	45.7	46.2	12.5	13.8	7.4	7.9
West	55.9	53.9	19.1	19.0	11.9	12.3
Mountain	56.7	55.3	18.5	18.2	11.3	11.7
Pacific	55.6	53.3	19.3	19.3	12.2	12.6
COUNTY TYPE						
Large Metro	52.0	52.7	16.2 ^a	17.4	9.7 ^a	10.7
Small Metro	52.6	52.2	16.4	16.4	9.8	10.3
250K – 1 Mil. Pop.	53.5	53.3	16.1	16.5	9.5	10.3
< 250K Pop.	50.7	50.1	17.0	16.1	10.4	10.2
Nonmetro	46.3	48.1	12.9	14.1	7.8	8.6
Urbanized	48.7	52.4	14.9	16.1	9.0	9.6
Less Urbanized	45.4	46.4	11.5	13.5	6.8	8.4
Completely Rural	40.4	38.2	10.1	8.1	7.0	4.8

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.64A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	114,712	117,213	32,952 ^b	35,124	19,810 ^b	22,188
GEOGRAPHIC DIVISION						
Northeast	21,007 ^a	22,277	6,156 ^b	7,043	3,598 ^b	4,505
New England	6,081	6,576	1,944	2,282	1,263	1,544
Middle Atlantic	14,927	15,701	4,212 ^a	4,761	2,336 ^b	2,961
Midwest	24,956	25,818	6,772	7,017	4,007	4,324
East North Central	17,537	18,336	4,932	4,993	2,969	3,130
West North Central	7,419	7,482	1,840	2,023	1,038	1,195
South	39,685	40,497	10,602	11,387	6,274 ^a	7,018
South Atlantic	22,283	22,958	6,244	6,509	3,748	4,072
East South Central	6,013	5,895	1,531	1,618	870	956
West South Central	11,389	11,645	2,828 ^a	3,260	1,656 ^a	1,990
West	29,064	28,620	9,421	9,677	5,932	6,341
Mountain	9,034	8,825	2,742	2,798	1,764	1,862
Pacific	20,030	19,795	6,679	6,879	4,168	4,479
COUNTY TYPE						
Large Metro	62,270 ^a	65,985	18,403 ^b	20,362	10,941 ^b	12,673
Small Metro	35,494	34,405	10,386	10,014	6,200	6,472
250K – 1 Mil. Pop.	24,559	23,622	6,869	6,847	4,068	4,414
< 250K Pop.	10,935	10,782	3,517	3,167	2,132	2,059
Nonmetro	16,948	16,823	4,163	4,748	2,669	3,043
Urbanized	8,037	7,688	2,164	2,221	1,423	1,386
Less Urbanized	7,369	7,785	1,657 ^b	2,244	1,014 ^b	1,469
Completely Rural	1,542	1,350	342	283	232	187

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.64B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	43.7	44.2	12.6 ^a	13.2	7.5 ^b	8.4
GEOGRAPHIC DIVISION						
Northeast	44.3 ^a	46.8	13.0 ^b	14.8	7.6 ^b	9.5
New England	48.7	52.3	15.6	18.2	10.1	12.3
Middle Atlantic	42.8	44.8	12.1	13.6	6.7 ^b	8.4
Midwest	44.4	45.7	12.0	12.4	7.1	7.7
East North Central	45.0	46.9	12.7	12.8	7.6	8.0
West North Central	43.0	43.1	10.7	11.7	6.0	6.9
South	40.7	41.0	10.9	11.5	6.4	7.1
South Atlantic	43.3	44.0	12.1	12.5	7.3	7.8
East South Central	38.8	37.8	9.9	10.4	5.6	6.1
West South Central	37.3	37.5	9.3	10.5	5.4	6.4
West	47.4	46.0	15.4	15.6	9.7	10.2
Mountain	48.3	46.4	14.7	14.7	9.4	9.8
Pacific	47.0	45.8	15.7	15.9	9.8	10.4
COUNTY TYPE						
Large Metro	44.1	45.0	13.0 ^a	13.9	7.8 ^b	8.6
Small Metro	45.0	44.4	13.2	12.9	7.9	8.4
250K – 1 Mil. Pop.	46.0	45.1	12.9	13.1	7.6	8.4
< 250K Pop.	42.9	43.1	13.8	12.6	8.4	8.2
Nonmetro	39.9	40.9	9.8 ^a	11.5	6.3 ^a	7.4
Urbanized	42.3	44.6	11.4	12.9	7.5	8.0
Less Urbanized	38.8	39.5	8.7 ^b	11.4	5.3 ^b	7.5
Completely Rural	34.5	32.0	7.7	6.7	5.2	4.5

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.65A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	4,093	4,091	3,346	3,262	1,762	1,830
GEOGRAPHIC DIVISION						
Northeast	740	679	624	583	337	317
New England	203	211	176	190	102	87
Middle Atlantic	537	467	448	393	235	230
Midwest	842	805	700	634	367	331
East North Central	619	581	514	467	268	255
West North Central	223	224	187	167	98	75
South	1,404	1,441	1,120	1,099	540	609
South Atlantic	769	747	619	594	290	306
East South Central	197	179	165	125	80	72
West South Central	438	515	336	379	170 ^a	231
West	1,107	1,167	902	946	519	573
Mountain	387	357	316	288	186	167
Pacific	720	811	586	658	333	406
COUNTY TYPE						
Large Metro	2,228	2,272	1,846	1,853	997	1,063
Small Metro	1,270	1,270	1,049	975	533	522
250K – 1 Mil. Pop.	869	892	711	694	378	368
< 250K Pop.	401	378	338	282	155	154
Nonmetro	595	550	451	434	232	245
Urbanized	265	242	206	197	112	111
Less Urbanized	281	263	207	203	99	112
Completely Rural	49	44	38	35	21	22

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.65B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	16.4	16.4	13.4	13.1	7.1	7.4
GEOGRAPHIC DIVISION						
Northeast	17.7	16.3	14.9	14.0	8.0	7.6
New England	18.6	19.5	16.1	17.5	9.3	8.0
Middle Atlantic	17.4	15.2	14.5	12.8	7.6	7.5
Midwest	15.6	15.0	13.0	11.8	6.8	6.2
East North Central	16.5	15.6	13.7	12.5	7.1	6.8
West North Central	13.6	13.6	11.4	10.2	6.0	4.6
South	15.0	15.3	12.0	11.7	5.8	6.5
South Atlantic	16.5	16.0	13.3	12.7	6.2	6.5
East South Central	13.3	12.1	11.2	8.5	5.4	4.9
West South Central	13.6	15.8	10.4	11.6	5.3 ^a	7.1
West	18.6	19.7	15.2	15.9	8.7	9.7
Mountain	20.6	18.8	16.9	15.2	9.9	8.8
Pacific	17.7	20.0	14.4	16.3	8.2	10.0
COUNTY TYPE						
Large Metro	16.4	16.2	13.6	13.2	7.4	7.6
Small Metro	17.0	18.0	14.1	13.8	7.2	7.4
250K – 1 Mil. Pop.	17.0	18.3	13.9	14.3	7.4	7.6
< 250K Pop.	17.2	17.1	14.6	12.8	6.7	7.0
Nonmetro	15.3	14.6	11.6	11.5	6.0	6.5
Urbanized	16.0	15.7	12.4	12.8	6.7	7.2
Less Urbanized	15.8	14.1	11.6	10.8	5.6	6.0
Completely Rural	11.1	12.1	8.7	9.4	4.7	6.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.66A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	18,051	18,390	11,000	11,154	6,636	6,836
GEOGRAPHIC DIVISION						
Northeast	3,312	3,309	2,082	2,191	1,255	1,364
New England	1,022	998	727	707	462	473
Middle Atlantic	2,290	2,311	1,355	1,484	793	891
Midwest	3,956	4,029	2,349	2,440	1,434	1,430
East North Central	2,709	2,774	1,661	1,680	998	1,013
West North Central	1,247	1,255	688	760	436	416
South	6,210	6,467	3,650	3,774	2,093	2,297
South Atlantic	3,331	3,447	2,023	2,056	1,162	1,284
East South Central	932	948	541	535	308	303
West South Central	1,947	2,073	1,086	1,183	623	710
West	4,573	4,584	2,920	2,749	1,854	1,745
Mountain	1,385	1,382	825	823	514	536
Pacific	3,187	3,202	2,094	1,926	1,340	1,209
COUNTY TYPE						
Large Metro	9,893	10,242	6,298	6,345	3,856	3,870
Small Metro	5,676	5,544	3,315	3,304	1,927	2,065
250K – 1 Mil. Pop.	3,707	3,569	2,135	2,159	1,229	1,352
< 250K Pop.	1,968	1,975	1,180	1,144	699	713
Nonmetro	2,482	2,603	1,387	1,506	853	901
Urbanized	1,189	1,300	709	785	456	452
Less Urbanized	1,109	1,145	582	641	344	403
Completely Rural	184	158	95	79	52	47

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.66B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	51.9	52.6	31.6	31.9	19.1	19.6
GEOGRAPHIC DIVISION						
Northeast	53.9	53.8	33.9	35.6	20.4	22.2
New England	62.7	60.7	44.6	43.0	28.3	28.8
Middle Atlantic	50.7	51.3	30.0	32.9	17.6	19.8
Midwest	53.4	54.2	31.7	32.8	19.4	19.2
East North Central	53.1	54.3	32.5	32.9	19.6	19.8
West North Central	54.2	54.1	29.9	32.8	18.9	18.0
South	48.3	50.0	28.4	29.2	16.3	17.7
South Atlantic	50.8	52.3	30.8	31.2	17.7	19.5
East South Central	45.7	46.2	26.5	26.1	15.1	14.8
West South Central	45.8	48.2	25.5	27.5	14.6	16.5
West	54.6	54.5	34.9	32.7	22.1	20.7
Mountain	54.7	54.0	32.6	32.2	20.3	20.9
Pacific	54.6	54.7	35.9	32.9	22.9	20.7
COUNTY TYPE						
Large Metro	52.8	53.4	33.6	33.1	20.6	20.2
Small Metro	52.2	52.1	30.5	31.1	17.7	19.4
250K – 1 Mil. Pop.	53.0	51.9	30.5	31.4	17.6	19.7
< 250K Pop.	50.7	52.5	30.4	30.4	18.0	19.0
Nonmetro	48.0	50.8	26.8	29.4	16.5	17.6
Urbanized	48.1 ^a	54.3	28.7	32.8	18.5	18.9
Less Urbanized	49.9	48.9	26.2	27.4	15.5	17.2
Completely Rural	38.8	40.7	20.1	20.4	11.1	12.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.67A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	92,568	94,732	18,606 ^b	20,707	11,412 ^b	13,521
GEOGRAPHIC DIVISION						
Northeast	16,956 ^a	18,289	3,450 ^b	4,268	2,006 ^b	2,824
New England	4,856	5,366	1,041 ^a	1,385	699 ^a	984
Middle Atlantic	12,099	12,923	2,409	2,883	1,308 ^b	1,840
Midwest	20,157	20,984	3,723	3,943	2,206	2,564
East North Central	14,208	14,982	2,758	2,847	1,702	1,861
West North Central	5,949	6,002	966	1,097	503	703
South	32,071	32,589	5,833	6,513	3,641	4,112
South Atlantic	18,183	18,763	3,601	3,859	2,296	2,482
East South Central	4,884	4,768	825	958	481	581
West South Central	9,004	9,058	1,406	1,697	864	1,049
West	23,384	22,869	5,599	5,982	3,560	4,022
Mountain	7,262	7,086	1,601	1,687	1,065	1,159
Pacific	16,123	15,783	3,998	4,295	2,495	2,863
COUNTY TYPE						
Large Metro	50,148 ^a	53,471	10,259 ^b	12,164	6,088 ^b	7,739
Small Metro	28,549	27,590	6,022	5,735	3,739	3,885
250K – 1 Mil. Pop.	19,983	19,161	4,024	3,994	2,461	2,694
< 250K Pop.	8,566	8,429	1,998	1,741	1,278	1,192
Nonmetro	13,871	13,670	2,325	2,808	1,585	1,896
Urbanized	6,583	6,146	1,250	1,239	854	823
Less Urbanized	5,979	6,377	868 ^b	1,400	571 ^b	955
Completely Rural	1,309	1,147	208	169	159	118

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.67B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	45.7	46.1	9.2 ^b	10.1	5.6 ^b	6.6
GEOGRAPHIC DIVISION						
Northeast	45.8 ^a	49.0	9.3 ^b	11.4	5.4 ^b	7.6
New England	49.8	54.6	10.7 ^a	14.1	7.2 ^a	10.0
Middle Atlantic	44.3	47.0	8.8	10.5	4.8 ^b	6.7
Midwest	46.4	48.1	8.6	9.0	5.1	5.9
East North Central	47.2	49.5	9.2	9.4	5.7	6.1
West North Central	44.8	44.8	7.3	8.2	3.8	5.3
South	42.6	42.6	7.7	8.5	4.8	5.4
South Atlantic	45.2	45.9	9.0	9.4	5.7	6.1
East South Central	40.8	39.5	6.9	7.9	4.0	4.8
West South Central	39.0	38.5	6.1	7.2	3.7	4.5
West	49.8	47.8	11.9	12.5	7.6	8.4
Mountain	50.8	48.7	11.2	11.6	7.4	8.0
Pacific	49.4	47.4	12.2	12.9	7.6	8.6
COUNTY TYPE						
Large Metro	46.1	47.2	9.4 ^b	10.7	5.6 ^b	6.8
Small Metro	47.1	46.2	9.9	9.6	6.2	6.5
250K – 1 Mil. Pop.	48.4	47.2	9.7	9.8	6.0	6.6
< 250K Pop.	44.4	44.2	10.4	9.1	6.6	6.2
Nonmetro	41.5	42.4	7.0 ^a	8.7	4.7	5.9
Urbanized	44.3	46.2	8.4	9.3	5.7	6.2
Less Urbanized	39.9	41.2	5.8 ^b	9.0	3.8 ^b	6.2
Completely Rural	36.9	33.2	5.9	4.9	4.5	3.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.68A Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	110,619	113,121	29,606 ^b	31,862	18,048 ^b	20,357
GEOGRAPHIC DIVISION						
Northeast	20,267 ^a	21,598	5,532 ^b	6,460	3,262 ^b	4,188
New England	5,878	6,365	1,768	2,092	1,161 ^a	1,457
Middle Atlantic	14,389	15,234	3,764 ^a	4,368	2,101 ^b	2,731
Midwest	24,114	25,013	6,072	6,383	3,640	3,994
East North Central	16,917	17,756	4,418	4,527	2,701	2,874
West North Central	7,197	7,258	1,654	1,856	939	1,120
South	38,281	39,057	9,483	10,288	5,734 ^a	6,409
South Atlantic	21,514	22,210	5,624	5,915	3,458	3,766
East South Central	5,816	5,716	1,366	1,493	789	884
West South Central	10,951	11,131	2,492	2,880	1,486	1,759
West	27,957	27,453	8,519	8,731	5,413	5,767
Mountain	8,647	8,468	2,426	2,510	1,579	1,695
Pacific	19,310	18,985	6,093	6,221	3,835	4,073
COUNTY TYPE						
Large Metro	60,041 ^a	63,713	16,557 ^b	18,509	9,944 ^b	11,610
Small Metro	34,225	33,135	9,337	9,038	5,667	5,950
250K – 1 Mil. Pop.	23,691	22,731	6,158	6,153	3,690	4,046
< 250K Pop.	10,534	10,404	3,178	2,885	1,977	1,905
Nonmetro	16,353	16,273	3,712	4,314	2,437	2,798
Urbanized	7,772	7,446	1,959	2,025	1,311	1,275
Less Urbanized	7,088	7,522	1,450 ^b	2,041	915 ^b	1,357
Completely Rural	1,493	1,305	303	248	211	165

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.68B Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	46.6	47.1	12.5 ^a	13.3	7.6 ^b	8.5
GEOGRAPHIC DIVISION						
Northeast	46.9 ^a	49.7	12.8 ^b	14.9	7.5 ^b	9.6
New England	51.6	55.4	15.5	18.2	10.2	12.7
Middle Atlantic	45.2	47.6	11.8 ^a	13.7	6.6 ^b	8.5
Midwest	47.5	49.0	11.9	12.5	7.2	7.8
East North Central	48.0	50.2	12.5	12.8	7.7	8.1
West North Central	46.2	46.2	10.6	11.8	6.0	7.1
South	43.4	43.7	10.8	11.5	6.5	7.2
South Atlantic	46.0	46.8	12.0	12.5	7.4	7.9
East South Central	41.5	40.5	9.7	10.6	5.6	6.3
West South Central	40.1	40.0	9.1	10.3	5.4	6.3
West	50.5	48.8	15.4	15.5	9.8	10.3
Mountain	51.4	49.5	14.4	14.7	9.4	9.9
Pacific	50.2	48.5	15.8	15.9	10.0	10.4
COUNTY TYPE						
Large Metro	47.1	48.1	13.0 ^a	14.0	7.8 ^b	8.8
Small Metro	47.9	47.1	13.1	12.8	7.9	8.5
250K – 1 Mil. Pop.	49.1	47.8	12.8	13.0	7.6	8.5
< 250K Pop.	45.4	45.6	13.7	12.6	8.5	8.3
Nonmetro	42.4	43.5	9.6 ^b	11.5	6.3 ^a	7.5
Urbanized	44.8	47.4	11.3	12.9	7.6	8.1
Less Urbanized	41.2	42.2	8.4 ^b	11.5	5.3 ^b	7.6
Completely Rural	37.1	34.0	7.5	6.5	5.2	4.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.69A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	37,634	39,200	4,182	4,553	1,549	1,530
GEOGRAPHIC DIVISION						
Northeast	7,016	7,617	873	1,077	261 ^a	393
New England	2,081	2,196	306	293	77	94
Middle Atlantic	4,935	5,421	566 ^a	785	184	299
Midwest	6,897	7,587	738	739	250	238
East North Central	4,852	5,401	591	551	195	185
West North Central	2,045	2,186	147	188	55	53
South	12,806	12,795	1,355	1,444	590	440
South Atlantic	7,122	7,183	841	858	410	302
East South Central	1,808	1,655	141	182	58	51
West South Central	3,876	3,957	373	404	122	86
West	10,915	11,201	1,216	1,292	448	459
Mountain	3,220	3,326	360	379	137	158
Pacific	7,695	7,876	856	913	311	301
COUNTY TYPE						
Large Metro	20,349 ^a	22,385	2,529	2,865	961	996
Small Metro	11,711	11,823	1,246	1,338	412	446
250K – 1 Mil. Pop.	8,113	8,339	893	934	312	325
< 250K Pop.	3,598	3,484	353	403	101	120
Nonmetro	5,574	4,992	407	350	175	88
Urbanized	2,742	2,309	198	176	49	39
Less Urbanized	2,289	2,286	150	151	77	38
Completely Rural	543	397	59	23	49	11

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.69B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	14.3	14.8	1.6	1.7	0.6	0.6
GEOGRAPHIC DIVISION						
Northeast	14.8	16.0	1.8	2.3	0.6	0.8
New England	16.7	17.5	2.5	2.3	0.6	0.8
Middle Atlantic	14.1	15.5	1.6 ^a	2.2	0.5	0.9
Midwest	12.3	13.4	1.3	1.3	0.4	0.4
East North Central	12.4	13.8	1.5	1.4	0.5	0.5
West North Central	11.9	12.6	0.9	1.1	0.3	0.3
South	13.1	12.9	1.4	1.5	0.6	0.4
South Atlantic	13.8	13.8	1.6	1.6	0.8	0.6
East South Central	11.7	10.6	0.9	1.2	0.4	0.3
West South Central	12.7	12.7	1.2	1.3	0.4	0.3
West	17.8	18.0	2.0	2.1	0.7	0.7
Mountain	17.2	17.5	1.9	2.0	0.7	0.8
Pacific	18.1	18.2	2.0	2.1	0.7	0.7
COUNTY TYPE						
Large Metro	14.4	15.3	1.8	2.0	0.7	0.7
Small Metro	14.8	15.3	1.6	1.7	0.5	0.6
250K – 1 Mil. Pop.	15.2	15.9	1.7	1.8	0.6	0.6
< 250K Pop.	14.1	13.9	1.4	1.6	0.4	0.5
Nonmetro	13.1	12.1	1.0	0.9	0.4	0.2
Urbanized	14.4	13.4	1.0	1.0	0.3	0.2
Less Urbanized	12.1	11.6	0.8	0.8	0.4	0.2
Completely Rural	12.2	9.4	1.3	0.6	1.1	0.3

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.70A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	215	222	131	168	43	39
GEOGRAPHIC DIVISION						
Northeast	25	33	15	20	6	6
New England	5	5	3	1	0	*
Middle Atlantic	20	27	12	19	5	6
Midwest	31	35	16	26	4	8
East North Central	20	28	9	19	2	4
West North Central	11	7	8	7	2	3
South	85	67	48	44	15	19
South Atlantic	53	29	30	18	10	3
East South Central	9	7	6	4	3	*
West South Central	24	31	12	22	2	17
West	75	88	52	77	19	6
Mountain	38	25	23	18	6	5
Pacific	37	62	29	59	14 ^a	2
COUNTY TYPE						
Large Metro	103	139	62 ^a	109	20	21
Small Metro	84 ^b	42	57 ^a	26	19	11
250K – 1 Mil. Pop.	48	29	36	21	10	7
< 250K Pop.	36 ^a	13	22 ^a	5	9	4
Nonmetro	29	42	12	33	4	7
Urbanized	16	20	6	18	*	2
Less Urbanized	11	18	6	11	4	3
Completely Rural	1	4	*	4	*	*

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.70B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9	0.9	0.5	0.7	0.2	0.2
GEOGRAPHIC DIVISION						
Northeast	0.6	0.8	0.4	0.5	0.1	0.1
New England	0.4	0.5	0.3	0.1	0.0	*
Middle Atlantic	0.7	0.9	0.4	0.6	0.2	0.2
Midwest	0.6	0.7	0.3	0.5	0.1	0.1
East North Central	0.5	0.7	0.2	0.5	0.0	0.1
West North Central	0.7	0.5	0.5	0.4	0.1	0.2
South	0.9	0.7	0.5	0.5	0.2	0.2
South Atlantic	1.1	0.6	0.7	0.4	0.2	0.1
East South Central	0.6	0.5	0.4	0.3	0.2	*
West South Central	0.7	0.9	0.4	0.7	0.1	0.5
West	1.3	1.5	0.9	1.3	0.3	0.1
Mountain	2.0	1.3	1.2	1.0	0.3	0.3
Pacific	0.9	1.5	0.7	1.5	0.3 ^a	0.0
COUNTY TYPE						
Large Metro	0.8	1.0	0.5 ^a	0.8	0.1	0.2
Small Metro	1.1 ^b	0.6	0.8 ^a	0.4	0.3	0.2
250K – 1 Mil. Pop.	0.9	0.6	0.7	0.4	0.2	0.1
< 250K Pop.	1.5 ^a	0.6	0.9 ^a	0.2	0.4	0.2
Nonmetro	0.7	1.1	0.3	0.9	0.1	0.2
Urbanized	1.0	1.3	0.4	1.2	*	0.1
Less Urbanized	0.6	0.9	0.3	0.6	0.2	0.1
Completely Rural	0.3	1.1	*	1.1	*	*

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.71A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	4,041	3,881	1,543	1,604	398	473
GEOGRAPHIC DIVISION						
Northeast	759	668	332	308	77	78
New England	220	202	113	105	20	27
Middle Atlantic	539	465	218	203	57	51
Midwest	766	676	270	300	67	89
East North Central	504	467	206	208	50	64
West North Central	262	209	64	92	17	25
South	1,340	1,420	440	495	113	139
South Atlantic	675	754	235	301	63	101
East South Central	187	203	56	73	13	15
West South Central	478	462	149	122	37	23
West	1,175	1,117	502	500	140	166
Mountain	386	367	122	141	39	51
Pacific	789	750	380	359	101	115
COUNTY TYPE						
Large Metro	2,204	2,103	887	902	234	272
Small Metro	1,219	1,233	485	539	130	175
250K – 1 Mil. Pop.	832	827	320	338	90	121
< 250K Pop.	387	406	165	201	39	54
Nonmetro	618	544	171	163	34	26
Urbanized	328 ^a	237	108	75	18	12
Less Urbanized	249	264	54	76	14	14
Completely Rural	41	43	9	*	2	0

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.71B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	11.6	11.1	4.4	4.6	1.1	1.4
GEOGRAPHIC DIVISION						
Northeast	12.4	10.9	5.4	5.0	1.3	1.3
New England	13.5	12.3	7.0	6.4	1.2	1.7
Middle Atlantic	11.9	10.3	4.8	4.5	1.3	1.1
Midwest	10.3	9.1	3.6	4.0	0.9	1.2
East North Central	9.9	9.1	4.0	4.1	1.0	1.3
West North Central	11.4	9.0	2.8	4.0	0.7	1.1
South	10.4	11.0	3.4	3.8	0.9	1.1
South Atlantic	10.3	11.4	3.6	4.6	1.0	1.5
East South Central	9.2	9.9	2.7	3.5	0.6	0.7
West South Central	11.2	10.8	3.5	2.8	0.9	0.5
West	14.0	13.3	6.0	5.9	1.7	2.0
Mountain	15.3	14.3	4.8	5.5	1.6	2.0
Pacific	13.5	12.8	6.5	6.1	1.7	2.0
COUNTY TYPE						
Large Metro	11.8	11.0	4.7	4.7	1.2	1.4
Small Metro	11.2	11.6	4.5	5.1	1.2	1.6
250K – 1 Mil. Pop.	11.9	12.0	4.6	4.9	1.3	1.8
< 250K Pop.	10.0	10.8	4.3	5.3	1.0	1.4
Nonmetro	12.0	10.6	3.3	3.2	0.7	0.5
Urbanized	13.3 ^a	9.9	4.4	3.2	0.7	0.5
Less Urbanized	11.2	11.3	2.4	3.3	0.6	0.6
Completely Rural	8.6	11.1	2.0	*	0.5	0.1

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.72A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	33,378	35,097	2,508	2,782	1,108	1,018
GEOGRAPHIC DIVISION						
Northeast	6,231	6,917	526 ^a	749	178 ^a	309
New England	1,856	1,988	190	186	56	67
Middle Atlantic	4,375	4,928	336 ^b	563	122 ^a	242
Midwest	6,100 ^a	6,876	452	413	179	142
East North Central	4,328 ^a	4,906	377	324	144	116
West North Central	1,772	1,970	75	89	36	25
South	11,381	11,307	868	905	462	281
South Atlantic	6,394	6,399	576	539	337	199
East South Central	1,612	1,445	80	105	42	36
West South Central	3,374	3,463	212	261	83	46
West	9,665	9,997	662	715	289	286
Mountain	2,796	2,933	215	221	92	102
Pacific	6,870	7,063	447	494	197	185
COUNTY TYPE						
Large Metro	18,042 ^a	20,143	1,581	1,854	707	703
Small Metro	10,409	10,549	704	773	263	259
250K – 1 Mil. Pop.	7,233	7,483	538	576	211	197
< 250K Pop.	3,175	3,066	166	197	52	62
Nonmetro	4,927	4,406	223	154	137	55
Urbanized	2,397	2,051	83	82	31	26
Less Urbanized	2,029	2,004	90	64	59	22
Completely Rural	501	350	49	8	47	8

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.72B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	16.5	17.1	1.2	1.4	0.5	0.5
GEOGRAPHIC DIVISION						
Northeast	16.8	18.5	1.4 ^a	2.0	0.5 ^a	0.8
New England	19.0	20.2	1.9	1.9	0.6	0.7
Middle Atlantic	16.0	17.9	1.2 ^a	2.0	0.4 ^a	0.9
Midwest	14.1 ^a	15.7	1.0	0.9	0.4	0.3
East North Central	14.4	16.2	1.3	1.1	0.5	0.4
West North Central	13.3	14.7	0.6	0.7	0.3	0.2
South	15.1	14.8	1.2	1.2	0.6	0.4
South Atlantic	15.9	15.7	1.4	1.3	0.8	0.5
East South Central	13.5	12.0	0.7	0.9	0.3	0.3
West South Central	14.6	14.7	0.9	1.1	0.4	0.2
West	20.6	20.9	1.4	1.5	0.6	0.6
Mountain	19.6	20.2	1.5	1.5	0.6	0.7
Pacific	21.0	21.2	1.4	1.5	0.6	0.6
COUNTY TYPE						
Large Metro	16.6	17.8	1.5	1.6	0.7	0.6
Small Metro	17.2	17.7	1.2	1.3	0.4	0.4
250K – 1 Mil. Pop.	17.5	18.4	1.3	1.4	0.5	0.5
< 250K Pop.	16.5	16.1	0.9	1.0	0.3	0.3
Nonmetro	14.8	13.7	0.7	0.5	0.4	0.2
Urbanized	16.1	15.4	0.6	0.6	0.2	0.2
Less Urbanized	13.5	12.9	0.6	0.4	0.4	0.1
Completely Rural	14.1	10.1	1.4	0.2	1.3	0.2

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.73A Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	37,419	38,978	4,051	4,385	1,505	1,491
GEOGRAPHIC DIVISION						
Northeast	6,991	7,584	858	1,057	255 ^a	387
New England	2,076	2,191	303	291	76	94
Middle Atlantic	4,914	5,394	554 ^a	766	179	293
Midwest	6,866	7,552	722	713	246	231
East North Central	4,832	5,374	583	532	194	181
West North Central	2,034	2,179	139	181	53	50
South	12,721	12,727	1,307	1,400	575	420
South Atlantic	7,069	7,154	811	840	400	299
East South Central	1,799	1,648	136	177	55	51
West South Central	3,853	3,926	361	383	120	70
West	10,840	11,114	1,164	1,215	429	452
Mountain	3,182	3,300	337	361	131	153
Pacific	7,658	7,813	827	854	298	299
COUNTY TYPE						
Large Metro	20,246 ^a	22,246	2,468	2,756	941	975
Small Metro	11,628	11,782	1,189	1,312	393	434
250K – 1 Mil. Pop.	8,066	8,310	858	914	302	318
< 250K Pop.	3,562	3,472	331	398	91	116
Nonmetro	5,545	4,950	394	318	171	81
Urbanized	2,726	2,288	192	158	49	37
Less Urbanized	2,278	2,268	144	140	73	36
Completely Rural	542	393	59	20	49	8

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.73B Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	15.8	16.2	1.7	1.8	0.6	0.6
GEOGRAPHIC DIVISION						
Northeast	16.2	17.4	2.0	2.4	0.6	0.9
New England	18.2	19.1	2.7	2.5	0.7	0.8
Middle Atlantic	15.4	16.9	1.7 ^a	2.4	0.6	0.9
Midwest	13.5	14.8	1.4	1.4	0.5	0.5
East North Central	13.7	15.2	1.7	1.5	0.5	0.5
West North Central	13.0	13.9	0.9	1.2	0.3	0.3
South	14.4	14.2	1.5	1.6	0.7	0.5
South Atlantic	15.1	15.1	1.7	1.8	0.9	0.6
East South Central	12.8	11.7	1.0	1.3	0.4	0.4
West South Central	14.1	14.1	1.3	1.4	0.4	0.3
West	19.6	19.8	2.1	2.2	0.8	0.8
Mountain	18.9	19.3	2.0	2.1	0.8	0.9
Pacific	19.9	20.0	2.1	2.2	0.8	0.8
COUNTY TYPE						
Large Metro	15.9	16.8	1.9	2.1	0.7	0.7
Small Metro	16.3	16.7	1.7	1.9	0.6	0.6
250K – 1 Mil. Pop.	16.7	17.5	1.8	1.9	0.6	0.7
< 250K Pop.	15.4	15.2	1.4	1.7	0.4	0.5
Nonmetro	14.4	13.2	1.0	0.8	0.4	0.2
Urbanized	15.7	14.6	1.1	1.0	0.3	0.2
Less Urbanized	13.2	12.7	0.8	0.8	0.4	0.2
Completely Rural	13.5	10.2	1.5	0.5	1.2	0.2

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.74A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8,870	9,424	632	773	377	354
GEOGRAPHIC DIVISION						
Northeast	1,229	1,561	100 ^b	244	66	98
New England	412	496	29	72	23	22
Middle Atlantic	818	1,065	71 ^a	171	42	75
Midwest	1,814	1,924	178	137	98	73
East North Central	1,204	1,283	153	96	80	50
West North Central	610	641	25	41	18	23
South	3,421	3,555	254	314	146	138
South Atlantic	2,005	1,992	160	151	90	95
East South Central	579	581	44	41	33	28
West South Central	838	982	50	122	22	15
West	2,405	2,384	100	79	68	46
Mountain	781	859	50	36	36	21
Pacific	1,624	1,526	50	43	31	25
COUNTY TYPE						
Large Metro	4,435	4,872	252 ^b	496	189	239
Small Metro	2,783	3,094	252	220	123	80
250K – 1 Mil. Pop.	1,993	2,112	208	151	101	69
< 250K Pop.	791	982	44	68	22	11
Nonmetro	1,651	1,457	128	58	65	35
Urbanized	679	561	42	25	14	7
Less Urbanized	780	721	42	25	24	20
Completely Rural	193	175	44	8	28	8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.74B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	3.4	3.6	0.2	0.3	0.1	0.1
GEOGRAPHIC DIVISION						
Northeast	2.6	3.3	0.2 ^b	0.5	0.1	0.2
New England	3.3	3.9	0.2	0.6	0.2	0.2
Middle Atlantic	2.3	3.0	0.2 ^a	0.5	0.1	0.2
Midwest	3.2	3.4	0.3	0.2	0.2	0.1
East North Central	3.1	3.3	0.4	0.2	0.2	0.1
West North Central	3.5	3.7	0.1	0.2	0.1	0.1
South	3.5	3.6	0.3	0.3	0.1	0.1
South Atlantic	3.9	3.8	0.3	0.3	0.2	0.2
East South Central	3.7	3.7	0.3	0.3	0.2	0.2
West South Central	2.7	3.2	0.2	0.4	0.1	0.0
West	3.9	3.8	0.2	0.1	0.1	0.1
Mountain	4.2	4.5	0.3	0.2	0.2	0.1
Pacific	3.8	3.5	0.1	0.1	0.1	0.1
COUNTY TYPE						
Large Metro	3.1	3.3	0.2 ^b	0.3	0.1	0.2
Small Metro	3.5	4.0	0.3	0.3	0.2	0.1
250K – 1 Mil. Pop.	3.7	4.0	0.4	0.3	0.2	0.1
< 250K Pop.	3.1	3.9	0.2	0.3	0.1	0.0
Nonmetro	3.9	3.5	0.3	0.1	0.2	0.1
Urbanized	3.6	3.3	0.2	0.1	0.1	0.0
Less Urbanized	4.1	3.7	0.2	0.1	0.1	0.1
Completely Rural	4.3	4.2	1.0	0.2	0.6	0.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.75A Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	39,736	39,647	4,430	4,250	1,333	1,173
GEOGRAPHIC DIVISION						
Northeast	7,321	7,293	780	861	249	192
New England	2,263	2,347	224	236	43	44
Middle Atlantic	5,059	4,946	557	626	206	148
Midwest	7,966	8,446	831	794	171	194
East North Central	5,434	5,733	603	584	110	135
West North Central	2,532	2,713	228	209	61	59
South	12,694	12,466	1,393	1,199	437	360
South Atlantic	7,192	7,187	834	692	271	215
East South Central	1,734	1,627	166	156	67	66
West South Central	3,767	3,652	393	350	99	79
West	11,756	11,442	1,425	1,396	476	427
Mountain	3,598	3,491	450	388	95	105
Pacific	8,158	7,951	976	1,008	381	322
COUNTY TYPE						
Large Metro	22,074	22,801	2,597	2,624	709	765
Small Metro	12,190	11,768	1,294	1,173	414	299
250K – 1 Mil. Pop.	8,460	8,304	922	788	304	206
< 250K Pop.	3,731	3,464	372	385	110	93
Nonmetro	5,472	5,078	539	453	210	109
Urbanized	2,760	2,476	334	243	140	53
Less Urbanized	2,263	2,244	159	190	43	53
Completely Rural	449	357	46	21	28	3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.75B Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	15.1	15.0	1.7	1.6	0.5	0.4
GEOGRAPHIC DIVISION						
Northeast	15.4	15.3	1.6	1.8	0.5	0.4
New England	18.1	18.7	1.8	1.9	0.3	0.4
Middle Atlantic	14.5	14.1	1.6	1.8	0.6	0.4
Midwest	14.2	15.0	1.5	1.4	0.3	0.3
East North Central	13.9	14.7	1.5	1.5	0.3	0.3
West North Central	14.7	15.6	1.3	1.2	0.4	0.3
South	13.0	12.6	1.4	1.2	0.4	0.4
South Atlantic	14.0	13.8	1.6	1.3	0.5	0.4
East South Central	11.2	10.4	1.1	1.0	0.4	0.4
West South Central	12.3	11.7	1.3	1.1	0.3	0.3
West	19.2	18.4	2.3	2.2	0.8	0.7
Mountain	19.2	18.4	2.4	2.0	0.5	0.6
Pacific	19.2	18.4	2.3	2.3	0.9	0.7
COUNTY TYPE						
Large Metro	15.7	15.6	1.8	1.8	0.5	0.5
Small Metro	15.5	15.2	1.6	1.5	0.5	0.4
250K – 1 Mil. Pop.	15.8	15.9	1.7	1.5	0.6	0.4
< 250K Pop.	14.6	13.8	1.5	1.5	0.4	0.4
Nonmetro	12.9	12.3	1.3	1.1	0.5	0.3
Urbanized	14.5	14.4	1.8	1.4	0.7	0.3
Less Urbanized	11.9	11.4	0.8	1.0	0.2	0.3
Completely Rural	10.1	8.5	1.0	0.5	0.6	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.76A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	21,068	21,293	1,533	1,617	496	546
GEOGRAPHIC DIVISION						
Northeast	4,135	4,312	247	329	97	141
New England	1,162	1,144	64	65	20	18
Middle Atlantic	2,973	3,167	183	263	77	123
Midwest	4,141	4,490	234	296	81	80
East North Central	2,902	3,241	181	219	71	69
West North Central	1,239	1,249	53	77	10	12
South	6,707	6,749	530	508	130	189
South Atlantic	3,748	3,988	320	272	71	100
East South Central	1,015	1,046	82	124	28	65
West South Central	1,945	1,715	128	112	31	24
West	6,085	5,742	522	484	187	135
Mountain	2,017	1,834	189 ^a	101	60	25
Pacific	4,068	3,908	332	382	127	110
COUNTY TYPE						
Large Metro	11,790	12,136	873	1,019	280	363
Small Metro	6,312	6,381	433	423	136	104
250K – 1 Mil. Pop.	4,526	4,638	294	283	76	78
< 250K Pop.	1,786	1,743	138	140	61	26
Nonmetro	2,966	2,777	227	176	80	79
Urbanized	1,513	1,261	121	77	48	37
Less Urbanized	1,229	1,264	89	86	20	37
Completely Rural	224	252	17	13	11	4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.76B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8.0	8.0	0.6	0.6	0.2	0.2
GEOGRAPHIC DIVISION						
Northeast	8.7	9.1	0.5	0.7	0.2	0.3
New England	9.3	9.1	0.5	0.5	0.2	0.1
Middle Atlantic	8.5	9.0	0.5	0.8	0.2	0.4
Midwest	7.4	8.0	0.4	0.5	0.1	0.1
East North Central	7.4	8.3	0.5	0.6	0.2	0.2
West North Central	7.2	7.2	0.3	0.4	0.1	0.1
South	6.9	6.8	0.5	0.5	0.1	0.2
South Atlantic	7.3	7.6	0.6	0.5	0.1	0.2
East South Central	6.5	6.7	0.5	0.8	0.2	0.4
West South Central	6.4	5.5	0.4	0.4	0.1	0.1
West	9.9	9.2	0.9	0.8	0.3	0.2
Mountain	10.8	9.6	1.0 ^a	0.5	0.3	0.1
Pacific	9.6	9.1	0.8	0.9	0.3	0.3
COUNTY TYPE						
Large Metro	8.4	8.3	0.6	0.7	0.2	0.2
Small Metro	8.0	8.2	0.5	0.5	0.2	0.1
250K – 1 Mil. Pop.	8.5	8.9	0.6	0.5	0.1	0.1
< 250K Pop.	7.0	7.0	0.5	0.6	0.2	0.1
Nonmetro	7.0	6.7	0.5	0.4	0.2	0.2
Urbanized	8.0	7.3	0.6	0.4	0.3	0.2
Less Urbanized	6.5	6.4	0.5	0.4	0.1	0.2
Completely Rural	5.0	6.0	0.4	0.3	0.2	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.77A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,327	1,307	480	512	121	149
GEOGRAPHIC DIVISION						
Northeast	163	171	71	71	18	22
New England	36	43	14	15	5	6
Middle Atlantic	127	128	58	56	13	17
Midwest	251	277	98	115	25	33
East North Central	176	217	77	92	19	32
West North Central	75	60	21	23	6 ^a	0
South	573	496	212	191	62	67
South Atlantic	291	223	109	83	36	35
East South Central	91	68	29	35	4	12
West South Central	191	205	74	72	23	20
West	339	364	99	135	16	27
Mountain	110	104	40	30	7	7
Pacific	229	260	59 ^a	105	8	19
COUNTY TYPE						
Large Metro	699	747	256	279	52	72
Small Metro	429	381	150	153	45	49
250K – 1 Mil. Pop.	275	249	99	99	26	37
< 250K Pop.	154	132	51	54	19	13
Nonmetro	199	179	74	80	24	27
Urbanized	78	74	28	28	8	8
Less Urbanized	103	80	41	42	16	15
Completely Rural	18	25	5	11	0	4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.77B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	5.3	5.3	1.9	2.1	0.5	0.6
GEOGRAPHIC DIVISION						
Northeast	3.9	4.1	1.7	1.7	0.4	0.5
New England	3.3	3.9	1.2	1.4	0.4	0.5
Middle Atlantic	4.1	4.2	1.9	1.8	0.4	0.5
Midwest	4.7	5.2	1.8	2.1	0.5	0.6
East North Central	4.7	5.8	2.1	2.5	0.5	0.9
West North Central	4.6	3.6	1.3	1.4	0.3 ^a	0.0
South	6.1	5.3	2.3	2.0	0.7	0.7
South Atlantic	6.2	4.8	2.3	1.8	0.8	0.7
East South Central	6.2	4.6	1.9	2.4	0.2	0.8
West South Central	5.9	6.3	2.3	2.2	0.7	0.6
West	5.7	6.1	1.7	2.3	0.3	0.4
Mountain	5.9	5.5	2.1	1.6	0.4	0.4
Pacific	5.6	6.4	1.4 ^a	2.6	0.2	0.5
COUNTY TYPE						
Large Metro	5.2	5.3	1.9	2.0	0.4	0.5
Small Metro	5.8	5.4	2.0	2.2	0.6	0.7
250K – 1 Mil. Pop.	5.4	5.1	1.9	2.0	0.5	0.8
< 250K Pop.	6.6	6.0	2.2	2.5	0.8	0.6
Nonmetro	5.1	4.7	1.9	2.1	0.6	0.7
Urbanized	4.7	4.8	1.7	1.8	0.5	0.5
Less Urbanized	5.8	4.3	2.3	2.2	0.9	0.8
Completely Rural	4.1	6.8	1.1	2.9	0.1	1.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.78A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	53,172	54,395	15,348	14,966	6,484	6,537
GEOGRAPHIC DIVISION						
Northeast	8,415 ^a	9,321	2,424	2,712	915 ^a	1,175
New England	2,486	2,790	641	709	197 ^a	322
Middle Atlantic	5,929	6,530	1,783	2,004	719	853
Midwest	10,627	11,146	3,002	3,028	1,275	1,274
East North Central	7,199 ^a	7,867	2,163	2,124	921	919
West North Central	3,428	3,279	839	904	354	355
South	19,315	19,690	5,874	5,555	2,528	2,332
South Atlantic	10,455	10,381	3,197	2,841	1,263	1,213
East South Central	2,908	2,980	906	955	456	380
West South Central	5,952	6,329	1,770	1,759	810	738
West	14,814	14,238	4,048	3,670	1,765	1,757
Mountain	4,450	4,297	1,272	1,125	464	507
Pacific	10,364	9,941	2,776	2,545	1,301	1,250
COUNTY TYPE						
Large Metro	28,830	30,457	8,342	8,474	3,526	3,770
Small Metro	16,499	16,508	4,722	4,360	2,104	1,927
250K – 1 Mil. Pop.	11,325	11,234	3,047	2,927	1,391	1,294
< 250K Pop.	5,174	5,274	1,675	1,432	713	632
Nonmetro	7,844	7,430	2,284	2,132	853	841
Urbanized	3,881	3,444	1,127	1,079	397	435
Less Urbanized	3,286	3,365	975	917	383	339
Completely Rural	677	621	182	136	74	67

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.78B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	20.3	20.5	5.8	5.6	2.5	2.5
GEOGRAPHIC DIVISION						
Northeast	17.8 ^a	19.6	5.1	5.7	1.9 ^a	2.5
New England	19.9	22.2	5.1	5.6	1.6 ^a	2.6
Middle Atlantic	17.0	18.6	5.1	5.7	2.1	2.4
Midwest	18.9	19.7	5.3	5.4	2.3	2.3
East North Central	18.5	20.1	5.5	5.4	2.4	2.3
West North Central	19.9	18.9	4.9	5.2	2.1	2.0
South	19.8	19.9	6.0	5.6	2.6	2.4
South Atlantic	20.3	19.9	6.2	5.4	2.5	2.3
East South Central	18.8	19.1	5.8	6.1	2.9	2.4
West South Central	19.5	20.4	5.8	5.7	2.6	2.4
West	24.2	22.9	6.6	5.9	2.9	2.8
Mountain	23.8	22.6	6.8	5.9	2.5	2.7
Pacific	24.3	23.0	6.5	5.9	3.1	2.9
COUNTY TYPE						
Large Metro	20.4	20.8	5.9	5.8	2.5	2.6
Small Metro	20.9	21.3	6.0	5.6	2.7	2.5
250K – 1 Mil. Pop.	21.2	21.5	5.7	5.6	2.6	2.5
< 250K Pop.	20.3	21.1	6.6	5.7	2.8	2.5
Nonmetro	18.5	18.1	5.4	5.2	2.0	2.0
Urbanized	20.4	20.0	5.9	6.3	2.1	2.5
Less Urbanized	17.3	17.1	5.1	4.7	2.0	1.7
Completely Rural	15.2	14.8	4.1	3.2	1.7	1.6

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.79A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	2,188	2,290	1,441	1,554	549 ^a	655
GEOGRAPHIC DIVISION						
Northeast	272	294	192	213	64	79
New England	70	64	51	45	18	14
Middle Atlantic	202	231	141	168	46	64
Midwest	441	444	285	303	118	124
East North Central	304	325	195	223	81	88
West North Central	137	120	90	80	37	37
South	864	959	553	650	213 ^a	292
South Atlantic	401	446	264	313	96	134
East South Central	141	134	91	76	30	40
West South Central	323	379	198	262	87	118
West	611	592	411	388	154	160
Mountain	197	169	137	108	40	46
Pacific	414	423	274	280	114	115
COUNTY TYPE						
Large Metro	1,106	1,207	752	820	280	330
Small Metro	717	686	478	446	183	199
250K – 1 Mil. Pop.	472	483	308	311	116	142
< 250K Pop.	245	203	171	135	67	57
Nonmetro	364	397	211 ^a	288	86 ^a	126
Urbanized	163	182	103	136	40	61
Less Urbanized	164	184	87 ^a	132	41	58
Completely Rural	37	31	21	19	5	7

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.79B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8.8	9.2	5.8	6.2	2.2 ^a	2.6
GEOGRAPHIC DIVISION						
Northeast	6.5	7.1	4.6	5.1	1.5	1.9
New England	6.4	5.9	4.6	4.1	1.6	1.3
Middle Atlantic	6.5	7.5	4.6	5.5	1.5	2.1
Midwest	8.2	8.3	5.3	5.6	2.2	2.3
East North Central	8.1	8.7	5.2	6.0	2.1	2.3
West North Central	8.3	7.3	5.5	4.9	2.3	2.2
South	9.2	10.2	5.9	6.9	2.3 ^a	3.1
South Atlantic	8.6	9.5	5.7	6.7	2.1	2.9
East South Central	9.6	9.1	6.2	5.1	2.1	2.7
West South Central	10.0	11.6	6.1	8.0	2.7	3.6
West	10.3	10.0	6.9	6.5	2.6	2.7
Mountain	10.5	8.9	7.3	5.7	2.1	2.4
Pacific	10.2	10.5	6.7	6.9	2.8	2.8
COUNTY TYPE						
Large Metro	8.2	8.6	5.5	5.8	2.1	2.4
Small Metro	9.6	9.7	6.4	6.3	2.5	2.8
250K – 1 Mil. Pop.	9.2	9.9	6.0	6.4	2.3	2.9
< 250K Pop.	10.5	9.2	7.3	6.1	2.9	2.6
Nonmetro	9.4	10.5	5.4 ^b	7.6	2.2 ^a	3.3
Urbanized	9.8	11.8	6.2 ^a	8.9	2.4	3.9
Less Urbanized	9.2	9.8	4.9 ^a	7.1	2.3	3.1
Completely Rural	8.4	8.5	4.7	5.2	1.1	2.0

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.80A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	9,239	9,197	4,245	4,133	1,661	1,551
GEOGRAPHIC DIVISION						
Northeast	1,522	1,540	780	732	279	274
New England	414	419	253	194	72	83
Middle Atlantic	1,108	1,121	527	539	208	190
Midwest	1,970	2,083	900	977	380	385
East North Central	1,328	1,466	636	686	263	277
West North Central	642	617	264	291	118	108
South	3,355	3,519	1,523	1,606	615	562
South Atlantic	1,648	1,797	788	818	293	246
East South Central	549	566	242	292	117	112
West South Central	1,158	1,156	493	497	205	204
West	2,392 ^b	2,055	1,042 ^b	816	387	330
Mountain	746	657	303	252	118	88
Pacific	1,645 ^a	1,398	738 ^a	565	268	243
COUNTY TYPE						
Large Metro	4,978	4,986	2,283	2,305	895	896
Small Metro	2,874	2,886	1,353	1,283	522	440
250K – 1 Mil. Pop.	1,808	1,866	861	835	330	304
< 250K Pop.	1,066	1,019	492	448	192	137
Nonmetro	1,387	1,325	609	544	244	215
Urbanized	691	632	312	267	127	111
Less Urbanized	591	602	254	250	112	94
Completely Rural	105	91	43	26	6	10

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.80B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	26.6	26.3	12.2	11.8	4.8	4.4
GEOGRAPHIC DIVISION						
Northeast	24.8	25.0	12.7	11.9	4.5	4.5
New England	25.4	25.5	15.5	11.8	4.4	5.1
Middle Atlantic	24.5	24.9	11.7	12.0	4.6	4.2
Midwest	26.6	28.0	12.1	13.2	5.1	5.2
East North Central	26.0	28.7	12.5	13.4	5.1	5.4
West North Central	27.9	26.6	11.5	12.6	5.1	4.7
South	26.1	27.2	11.8	12.4	4.8	4.3
South Atlantic	25.1	27.3	12.0	12.4	4.5	3.7
East South Central	26.9	27.6	11.9	14.2	5.7	5.5
West South Central	27.2	26.9	11.6	11.5	4.8	4.7
West	28.6 ^b	24.4	12.4 ^b	9.7	4.6	3.9
Mountain	29.5	25.7	12.0	9.8	4.7	3.4
Pacific	28.2 ^a	23.9	12.6 ^a	9.6	4.6	4.1
COUNTY TYPE						
Large Metro	26.6	26.0	12.2	12.0	4.8	4.7
Small Metro	26.4	27.1	12.4	12.1	4.8	4.1
250K – 1 Mil. Pop.	25.9	27.1	12.3	12.1	4.7	4.4
< 250K Pop.	27.5	27.1	12.7	11.9	4.9	3.6
Nonmetro	26.8	25.9	11.8	10.6	4.7	4.2
Urbanized	28.0	26.4	12.6	11.2	5.1	4.6
Less Urbanized	26.6	25.7	11.4	10.7	5.0	4.0
Completely Rural	22.1	23.5	9.0	6.8	1.2	2.6

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.81A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	41,745	42,909	9,662	9,280	4,274	4,331
GEOGRAPHIC DIVISION						
Northeast	6,622 ^a	7,487	1,452	1,767	572 ^a	822
New England	2,002	2,308	337	470	108 ^a	224
Middle Atlantic	4,620	5,179	1,115	1,297	465	598
Midwest	8,217	8,619	1,817	1,748	777	765
East North Central	5,567	6,076	1,332	1,215	578	554
West North Central	2,650	2,543	485	533	199	210
South	15,095	15,212	3,798	3,299	1,700	1,478
South Atlantic	8,406	8,138	2,146	1,710	874	834
East South Central	2,219	2,280	573	587	309	228
West South Central	4,471	4,794	1,079	1,001	518	416
West	11,812	11,591	2,595	2,466	1,225	1,266
Mountain	3,507	3,470	832	765	306	373
Pacific	8,304	8,121	1,763	1,701	919	893
COUNTY TYPE						
Large Metro	22,746	24,265	5,307	5,349	2,352	2,544
Small Metro	12,907	12,936	2,891	2,631	1,399	1,287
250K – 1 Mil. Pop.	9,044	8,885	1,879	1,781	945	849
< 250K Pop.	3,863	4,051	1,013	849	454	438
Nonmetro	6,093	5,708	1,464	1,300	523	500
Urbanized	3,026	2,630	711	675	230	263
Less Urbanized	2,531	2,579	634	535	230	187
Completely Rural	535	499	118	90	63	49

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.81B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	20.6	20.9	4.8	4.5	2.1	2.1
GEOGRAPHIC DIVISION						
Northeast	17.9 ^a	20.1	3.9	4.7	1.5 ^a	2.2
New England	20.5	23.5	3.5	4.8	1.1 ^a	2.3
Middle Atlantic	16.9	18.8	4.1	4.7	1.7	2.2
Midwest	18.9	19.7	4.2	4.0	1.8	1.8
East North Central	18.5	20.1	4.4	4.0	1.9	1.8
West North Central	19.9	19.0	3.6	4.0	1.5	1.6
South	20.0	19.9	5.0	4.3	2.3	1.9
South Atlantic	20.9	19.9	5.3	4.2	2.2	2.0
East South Central	18.5	18.9	4.8	4.9	2.6	1.9
West South Central	19.4	20.4	4.7	4.3	2.2	1.8
West	25.2	24.2	5.5	5.2	2.6	2.6
Mountain	24.5	23.8	5.8	5.3	2.1	2.6
Pacific	25.4	24.4	5.4	5.1	2.8	2.7
COUNTY TYPE						
Large Metro	20.9	21.4	4.9	4.7	2.2	2.2
Small Metro	21.3	21.7	4.8	4.4	2.3	2.2
250K – 1 Mil. Pop.	21.9	21.9	4.6	4.4	2.3	2.1
< 250K Pop.	20.0	21.2	5.2	4.5	2.4	2.3
Nonmetro	18.2	17.7	4.4	4.0	1.6	1.5
Urbanized	20.4	19.8	4.8	5.1	1.5	2.0
Less Urbanized	16.9	16.7	4.2	3.5	1.5	1.2
Completely Rural	15.1	14.4	3.3	2.6	1.8	1.4

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.82A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	50,984	52,105	13,907	13,412	5,935	5,882
GEOGRAPHIC DIVISION						
Northeast	8,144 ^a	9,026	2,232	2,499	851	1,096
New England	2,416	2,727	591	664	179 ^a	307
Middle Atlantic	5,728	6,300	1,642	1,836	672	789
Midwest	10,186	10,702	2,717	2,726	1,157	1,150
East North Central	6,895	7,542	1,968	1,901	840	831
West North Central	3,292	3,160	749	824	317	318
South	18,451	18,731	5,321	4,905	2,316	2,040
South Atlantic	10,054	9,935	2,933	2,528	1,167	1,080
East South Central	2,767	2,846	815	879	426	340
West South Central	5,629	5,950	1,572	1,498	723	620
West	14,203	13,646	3,637	3,282	1,612	1,596
Mountain	4,254	4,128	1,135	1,017	425	461
Pacific	9,949	9,519	2,502	2,265	1,187	1,135
COUNTY TYPE						
Large Metro	27,723	29,251	7,590	7,654	3,247	3,440
Small Metro	15,781	15,822	4,244	3,914	1,921	1,727
250K – 1 Mil. Pop.	10,853	10,751	2,739	2,616	1,275	1,153
< 250K Pop.	4,929	5,071	1,505	1,298	646	575
Nonmetro	7,479	7,033	2,073	1,844	768	715
Urbanized	3,718	3,262	1,024	943	357	374
Less Urbanized	3,122	3,181	888	785	342	281
Completely Rural	640	590	161	117	69	59

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.82B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	21.5	21.7	5.9	5.6	2.5	2.4
GEOGRAPHIC DIVISION						
Northeast	18.9 ^a	20.8	5.2	5.7	2.0	2.5
New England	21.2	23.8	5.2	5.8	1.6 ^a	2.7
Middle Atlantic	18.0	19.7	5.2	5.7	2.1	2.5
Midwest	20.0	20.9	5.3	5.3	2.3	2.3
East North Central	19.6	21.3	5.6	5.4	2.4	2.3
West North Central	21.1	20.1	4.8	5.2	2.0	2.0
South	20.9	20.9	6.0	5.5	2.6	2.3
South Atlantic	21.5	20.9	6.3	5.3	2.5	2.3
East South Central	19.7	20.2	5.8	6.2	3.0	2.4
West South Central	20.6	21.4	5.8	5.4	2.6	2.2
West	25.7	24.3	6.6	5.8	2.9	2.8
Mountain	25.3	24.1	6.7	5.9	2.5	2.7
Pacific	25.8	24.3	6.5	5.8	3.1	2.9
COUNTY TYPE						
Large Metro	21.7	22.1	6.0	5.8	2.5	2.6
Small Metro	22.1	22.5	5.9	5.6	2.7	2.5
250K – 1 Mil. Pop.	22.5	22.6	5.7	5.5	2.6	2.4
< 250K Pop.	21.3	22.2	6.5	5.7	2.8	2.5
Nonmetro	19.4	18.8	5.4	4.9	2.0	1.9
Urbanized	21.4	20.8	5.9	6.0	2.1	2.4
Less Urbanized	18.1	17.8	5.2	4.4	2.0	1.6
Completely Rural	15.9	15.3	4.0	3.0	1.7	1.5

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.83A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	35,473	36,064	11,082	10,337	4,521	4,325
GEOGRAPHIC DIVISION						
Northeast	5,164 ^b	6,172	1,569	1,732	643	751
New England	1,530 ^a	1,938	393	432	115	193
Middle Atlantic	3,634 ^a	4,234	1,176	1,300	528	559
Midwest	7,182	7,553	2,153	2,194	879	895
East North Central	5,031	5,440	1,557	1,589	644	669
West North Central	2,151	2,113	596	605	235	226
South	13,149	12,901	4,276	3,857	1,777	1,558
South Atlantic	7,065	6,703	2,282	1,976	877	810
East South Central	2,073	2,054	712	627	368	267
West South Central	4,011	4,144	1,282	1,254	532	481
West	9,979	9,438	3,084 ^a	2,554	1,222	1,122
Mountain	3,117	2,753	989	792	280	320
Pacific	6,862	6,685	2,095	1,762	942	802
COUNTY TYPE						
Large Metro	19,343	20,121	5,952	5,665	2,473	2,399
Small Metro	10,810	10,945	3,371	3,120	1,486	1,360
250K – 1 Mil. Pop.	7,379	7,431	2,231	2,071	1,008	895
< 250K Pop.	3,431	3,514	1,139	1,048	478	465
Nonmetro	5,320	4,998	1,759	1,552	563	566
Urbanized	2,646	2,297	916	815	259	325
Less Urbanized	2,242	2,269	700	637	238	201
Completely Rural	432	433	143	99	66	40

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.83B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	13.5	13.6	4.2 ^a	3.9	1.7	1.6
GEOGRAPHIC DIVISION						
Northeast	10.9 ^b	13.0	3.3	3.6	1.4	1.6
New England	12.3 ^a	15.4	3.1	3.4	0.9	1.5
Middle Atlantic	10.4 ^a	12.1	3.4	3.7	1.5	1.6
Midwest	12.8	13.4	3.8	3.9	1.6	1.6
East North Central	12.9	13.9	4.0	4.1	1.7	1.7
West North Central	12.5	12.2	3.5	3.5	1.4	1.3
South	13.5	13.1	4.4	3.9	1.8	1.6
South Atlantic	13.7	12.8	4.4	3.8	1.7	1.6
East South Central	13.4	13.2	4.6	4.0	2.4	1.7
West South Central	13.1	13.3	4.2	4.0	1.7	1.5
West	16.3	15.2	5.0 ^a	4.1	2.0	1.8
Mountain	16.7	14.5	5.3	4.2	1.5	1.7
Pacific	16.1	15.5	4.9	4.1	2.2	1.9
COUNTY TYPE						
Large Metro	13.7	13.7	4.2	3.9	1.8	1.6
Small Metro	13.7	14.1	4.3	4.0	1.9	1.8
250K – 1 Mil. Pop.	13.8	14.2	4.2	4.0	1.9	1.7
< 250K Pop.	13.4	14.0	4.5	4.2	1.9	1.9
Nonmetro	12.5	12.1	4.1	3.8	1.3	1.4
Urbanized	13.9	13.3	4.8	4.7	1.4	1.9
Less Urbanized	11.8	11.5	3.7	3.2	1.3	1.0
Completely Rural	9.7	10.3	3.2	2.4	1.5	1.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.84A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,805	1,803	1,157	1,165	425	467
GEOGRAPHIC DIVISION						
Northeast	206	237	135	165	48	70
New England	50	45	35	31	15	13
Middle Atlantic	156	191	100	134	33	58
Midwest	365	366	230	228	85	79
East North Central	258	275	160	179	61	61
West North Central	107	90	70	48	24	18
South	714	729	452	468	171	202
South Atlantic	334	345	214	226	79	92
East South Central	108	115	68	67	26	33
West South Central	272	270	170	175	66	76
West	520	472	340	304	121	116
Mountain	163	145	114	93	27	36
Pacific	357	327	226	211	94	81
COUNTY TYPE						
Large Metro	898	909	596	578	213	220
Small Metro	596	571	382	362	139	153
250K – 1 Mil. Pop.	394	405	241	256	84	112
< 250K Pop.	202	166	141	105	55	40
Nonmetro	312	324	179	225	73	95
Urbanized	141	148	87	114	33	50
Less Urbanized	139	147	76	97	37	41
Completely Rural	31	29	16	14	2	4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.84B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	7.3	7.3	4.6	4.7	1.7	1.9
GEOGRAPHIC DIVISION						
Northeast	4.9	5.7	3.2	4.0	1.2	1.7
New England	4.6	4.2	3.2	2.9	1.4	1.2
Middle Atlantic	5.1	6.2	3.2	4.4	1.1	1.9
Midwest	6.8	6.8	4.3	4.2	1.6	1.5
East North Central	6.9	7.4	4.3	4.8	1.6	1.6
West North Central	6.5	5.5	4.3	2.9	1.5	1.1
South	7.6	7.7	4.8	5.0	1.8	2.1
South Atlantic	7.2	7.4	4.6	4.8	1.7	2.0
East South Central	7.3	7.8	4.6	4.5	1.8	2.2
West South Central	8.4	8.3	5.3	5.4	2.0	2.3
West	8.7	8.0	5.7	5.1	2.0	2.0
Mountain	8.7	7.7	6.1	4.9	1.5	1.9
Pacific	8.8	8.1	5.5	5.2	2.3	2.0
COUNTY TYPE						
Large Metro	6.6	6.5	4.4	4.1	1.6	1.6
Small Metro	8.0	8.1	5.1	5.1	1.9	2.2
250K – 1 Mil. Pop.	7.7	8.3	4.7	5.3	1.6	2.3
< 250K Pop.	8.7	7.5	6.0	4.8	2.4	1.8
Nonmetro	8.0	8.6	4.6	6.0	1.9	2.5
Urbanized	8.5	9.6	5.2	7.4	2.0	3.2
Less Urbanized	7.8	7.8	4.3	5.2	2.1	2.2
Completely Rural	7.0	7.9	3.5	3.9	0.5	1.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.85A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	7,251	7,049	3,061 ^a	2,740	1,132	978
GEOGRAPHIC DIVISION						
Northeast	1,095	1,158	499	458	189	148
New England	266	292	143	115	36	46
Middle Atlantic	829	865	356	343	154	103
Midwest	1,563	1,586	640	651	232	247
East North Central	1,043	1,125	438	466	154	181
West North Central	520	462	201	185	78	66
South	2,643	2,646	1,120	1,095	419	359
South Atlantic	1,273	1,316	556	518	195	148
East South Central	459	458	182	220	79	82
West South Central	911	873	382	356	146	130
West	1,950 ^b	1,659	801 ^b	536	291	223
Mountain	596	536	224	167	77	60
Pacific	1,353 ^a	1,122	578 ^b	369	214	163
COUNTY TYPE						
Large Metro	3,827	3,738	1,626	1,490	611	567
Small Metro	2,291	2,251	990	868	361 ^a	267
250K – 1 Mil. Pop.	1,462	1,432	650	557	237	179
< 250K Pop.	829	819	340	312	124	88
Nonmetro	1,133	1,060	445	381	160	144
Urbanized	564	494	236	170	82	65
Less Urbanized	493	488	176	190	72	73
Completely Rural	75	78	33	20	5	7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.85B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	20.8	20.2	8.8 ^a	7.8	3.3	2.8
GEOGRAPHIC DIVISION						
Northeast	17.8	18.8	8.1	7.4	3.1	2.4
New England	16.3	17.8	8.8	7.0	2.2	2.8
Middle Atlantic	18.4	19.2	7.9	7.6	3.4	2.3
Midwest	21.1	21.4	8.6	8.8	3.1	3.3
East North Central	20.4	22.0	8.6	9.1	3.0	3.5
West North Central	22.6	19.9	8.7	8.0	3.4	2.9
South	20.6	20.4	8.7	8.5	3.3	2.8
South Atlantic	19.4	20.0	8.5	7.9	3.0	2.2
East South Central	22.5	22.3	8.9	10.7	3.9	4.0
West South Central	21.4	20.3	9.0	8.3	3.4	3.0
West	23.3 ^b	19.7	9.6 ^b	6.4	3.5	2.7
Mountain	23.6	21.0	8.8	6.5	3.0	2.4
Pacific	23.2 ^a	19.2	9.9 ^b	6.3	3.7	2.8
COUNTY TYPE						
Large Metro	20.4	19.5	8.7	7.8	3.3	3.0
Small Metro	21.1	21.2	9.1	8.2	3.3 ^a	2.5
250K – 1 Mil. Pop.	20.9	20.8	9.3	8.1	3.4	2.6
< 250K Pop.	21.4	21.8	8.8	8.3	3.2	2.3
Nonmetro	21.9	20.7	8.6	7.4	3.1	2.8
Urbanized	22.8	20.6	9.6	7.1	3.3	2.7
Less Urbanized	22.2	20.8	7.9	8.1	3.3	3.1
Completely Rural	15.9	20.1	6.9	5.3	1.1	1.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.86A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	26,417	27,211	6,865	6,432	2,964	2,879
GEOGRAPHIC DIVISION						
Northeast	3,862 ^b	4,778	934	1,108	406	532
New England	1,214 ^a	1,600	214	286	64	134
Middle Atlantic	2,648 ^a	3,178	719	822	342	398
Midwest	5,254	5,601	1,283	1,316	562	568
East North Central	3,730	4,040	958	944	430	426
West North Central	1,524	1,561	325	372	132	142
South	9,792	9,526	2,704	2,295	1,187	997
South Atlantic	5,457	5,042	1,513	1,232	603	570
East South Central	1,506	1,482	462	340	263	153
West South Central	2,828	3,001	730	723	321	274
West	7,509	7,307	1,943	1,714	810	782
Mountain	2,358	2,071	651	532	176	224
Pacific	5,151	5,236	1,292	1,182	634	558
COUNTY TYPE						
Large Metro	14,619	15,474	3,730	3,597	1,648	1,612
Small Metro	7,923	8,123	1,999	1,890	986	940
250K – 1 Mil. Pop.	5,523	5,594	1,340	1,258	687	604
< 250K Pop.	2,400	2,529	659	632	299	336
Nonmetro	3,875	3,614	1,135	945	330	327
Urbanized	1,940	1,655	593	531	144	211
Less Urbanized	1,609	1,634	448	350	128	87
Completely Rural	325	325	95	65	58	29

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.86B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	13.0	13.3	3.4	3.1	1.5	1.4
GEOGRAPHIC DIVISION						
Northeast	10.4 ^b	12.8	2.5	3.0	1.1	1.4
New England	12.4 ^a	16.3	2.2	2.9	0.7	1.4
Middle Atlantic	9.7 ^a	11.6	2.6	3.0	1.3	1.4
Midwest	12.1	12.8	3.0	3.0	1.3	1.3
East North Central	12.4	13.3	3.2	3.1	1.4	1.4
West North Central	11.5	11.7	2.4	2.8	1.0	1.1
South	13.0	12.5	3.6	3.0	1.6	1.3
South Atlantic	13.6	12.3	3.8	3.0	1.5	1.4
East South Central	12.6	12.3	3.9	2.8	2.2	1.3
West South Central	12.3	12.8	3.2	3.1	1.4	1.2
West	16.0	15.3	4.1	3.6	1.7	1.6
Mountain	16.5	14.2	4.6	3.7	1.2	1.5
Pacific	15.8	15.7	4.0	3.6	1.9	1.7
COUNTY TYPE						
Large Metro	13.4	13.7	3.4	3.2	1.5	1.4
Small Metro	13.1	13.6	3.3	3.2	1.6	1.6
250K – 1 Mil. Pop.	13.4	13.8	3.2	3.1	1.7	1.5
< 250K Pop.	12.4	13.3	3.4	3.3	1.5	1.8
Nonmetro	11.6	11.2	3.4	2.9	1.0	1.0
Urbanized	13.1	12.4	4.0	4.0	1.0	1.6
Less Urbanized	10.7	10.6	3.0	2.3	0.9	0.6
Completely Rural	9.2	9.4	2.7	1.9	1.6	0.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.87A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	33,668	34,261	9,925	9,172	4,096	3,858
GEOGRAPHIC DIVISION						
Northeast	4,957 ^b	5,936	1,433	1,566	595	681
New England	1,480 ^a	1,892	358	401	100	180
Middle Atlantic	3,477 ^a	4,043	1,075	1,166	495	501
Midwest	6,817	7,187	1,923	1,967	794	816
East North Central	4,773	5,165	1,396	1,409	583	607
West North Central	2,044	2,023	527	557	211	208
South	12,435	12,172	3,825	3,389	1,606	1,356
South Atlantic	6,731	6,358	2,069	1,750	798	718
East South Central	1,965	1,940	644	560	341	235
West South Central	3,739	3,875	1,112	1,079	466	404
West	9,459	8,966	2,744 ^a	2,251	1,101	1,005
Mountain	2,954	2,608	875	699	253	284
Pacific	6,505	6,359	1,869	1,551	848	721
COUNTY TYPE						
Large Metro	18,445	19,212	5,356	5,087	2,259	2,179
Small Metro	10,214	10,374	2,989	2,758	1,347	1,207
250K – 1 Mil. Pop.	6,984	7,026	1,990	1,815	924	783
< 250K Pop.	3,229	3,348	999	943	423	424
Nonmetro	5,009	4,674	1,580	1,327	490	472
Urbanized	2,505	2,149	829	701	226	276
Less Urbanized	2,103	2,122	623	540	201	160
Completely Rural	401	403	128	85	63	36

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.87B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	14.2	14.3	4.2 ^a	3.8	1.7	1.6
GEOGRAPHIC DIVISION						
Northeast	11.5 ^b	13.7	3.3	3.6	1.4	1.6
New England	13.0 ^a	16.5	3.1	3.5	0.9	1.6
Middle Atlantic	10.9 ^a	12.6	3.4	3.6	1.6	1.6
Midwest	13.4	14.1	3.8	3.8	1.6	1.6
East North Central	13.6	14.6	4.0	4.0	1.7	1.7
West North Central	13.1	12.9	3.4	3.5	1.4	1.3
South	14.1	13.6	4.3	3.8	1.8	1.5
South Atlantic	14.4	13.4	4.4	3.7	1.7	1.5
East South Central	14.0	13.7	4.6	4.0	2.4	1.7
West South Central	13.7	13.9	4.1	3.9	1.7	1.5
West	17.1	15.9	5.0 ^a	4.0	2.0	1.8
Mountain	17.6	15.2	5.2	4.1	1.5	1.7
Pacific	16.9	16.2	4.9	4.0	2.2	1.8
COUNTY TYPE						
Large Metro	14.5	14.5	4.2	3.8	1.8	1.6
Small Metro	14.3	14.7	4.2	3.9	1.9	1.7
250K – 1 Mil. Pop.	14.5	14.8	4.1	3.8	1.9	1.6
< 250K Pop.	13.9	14.7	4.3	4.1	1.8	1.9
Nonmetro	13.0	12.5	4.1	3.5	1.3	1.3
Urbanized	14.4	13.7	4.8	4.5	1.3	1.8
Less Urbanized	12.2	11.9	3.6	3.0	1.2	0.9
Completely Rural	10.0	10.5	3.2	2.2	1.6	0.9

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.88A Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Numbers in Thousands, 2013 and 2014

Drug or Method of Administration	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
HALLUCINOGENS	39,736	39,647	702	610	6,122	5,784	32,912	33,253
PCP (Angel Dust, Phencyclidine)	6,462	6,388	46	60	326	305	6,090	6,023
LSD (Acid)	24,770	25,035	221 ^a	291	2,246	2,451	22,303	22,293
Peyote	5,759	5,684	37	36	332 ^b	210	5,390	5,438
Mescaline	8,116	7,955	23	20	324 ^a	227	7,769	7,708
Psilocybin (Mushrooms)	22,850	22,447	295	260	3,347	3,138	19,209	19,049
Ecstasy (MDMA)	17,815	17,548	369 ^a	297	4,436	4,200	13,009	13,051
INHALANTS	21,068	21,293	1,327	1,307	2,613	2,444	17,129	17,543
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	7,384	7,379	176	195	521	443	6,687	6,741
Correction Fluid, Degreaser, or Cleaning Fluid	1,350	1,495	277	256	216	219	857	1,020
Gasoline or Lighter Fluid	2,686	2,784	397	425	325	378	1,965	1,981
Glue, Shoe Polish, or Toluene	2,882	2,874	471	510	353	306	2,059	2,058
Halothane, Ether, or Other Anesthetics	803	980	50	37	99	101	653	842
Lacquer Thinner or Other Paint Solvents	1,359	1,374	171	200	183	200	1,005	973
Lighter Gases (Butane, Propane)	758	653	110	117	109	116	539	419
Nitrous Oxide or Whippits	12,368	12,671	156	166	1,529	1,450	10,683	11,055
Spray Paints	1,662	1,607	355 ^a	441	292 ^a	219	1,015	947
Other Aerosol Sprays	1,979	2,005	244	261	483	434	1,252	1,311
NEEDLE USE^{1,2,3}	4,442	4,256	48	49	473	467	3,920	3,740
Heroin Needle Use ^{1,2}	2,555	2,315	19	22	322	300	2,213	1,993
Cocaine Needle Use ^{1,2}	2,287	2,277	2	14	157	149	2,128	2,114
Stimulant Needle Use ^{1,2,3}	2,347	2,184	15	25	195	203	2,137	1,957
Methamphetamine Needle Use ^{1,2,3}	1,891	1,798	14	23	144	157	1,733	1,617
HEROIN	4,812	4,813	38	36	619	683	4,155	4,095
Smoke Heroin ²	1,650	1,643	15	23	295	322	1,341	1,298
Sniff or Snort Heroin ²	3,257	3,501	14	20	437	512	2,806	2,969

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

³ Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.88B Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Percentages, 2013 and 2014

Drug or Method of Administration	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
HALLUCINOGENS	15.1	15.0	2.8	2.5	17.6	16.6	16.2	16.2
PCP (Angel Dust, Phencyclidine)	2.5	2.4	0.2	0.2	0.9	0.9	3.0	2.9
LSD (Acid)	9.4	9.4	0.9 ^a	1.2	6.5	7.0	11.0	10.9
Peyote	2.2	2.1	0.2	0.1	1.0 ^b	0.6	2.7	2.6
Mescaline	3.1	3.0	0.1	0.1	0.9 ^a	0.6	3.8	3.8
Psilocybin (Mushrooms)	8.7	8.5	1.2	1.0	9.6	9.0	9.5	9.3
Ecstasy (MDMA)	6.8	6.6	1.5 ^a	1.2	12.8	12.0	6.4	6.4
INHALANTS	8.0	8.0	5.3	5.3	7.5	7.0	8.4	8.5
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	2.8	2.8	0.7	0.8	1.5	1.3	3.3	3.3
Correction Fluid, Degreaser, or Cleaning Fluid	0.5	0.6	1.1	1.0	0.6	0.6	0.4	0.5
Gasoline or Lighter Fluid	1.0	1.0	1.6	1.7	0.9	1.1	1.0	1.0
Glue, Shoe Polish, or Toluene	1.1	1.1	1.9	2.0	1.0	0.9	1.0	1.0
Halothane, Ether, or Other Anesthetics	0.3	0.4	0.2	0.1	0.3	0.3	0.3	0.4
Lacquer Thinner or Other Paint Solvents	0.5	0.5	0.7	0.8	0.5	0.6	0.5	0.5
Lighter Gases (Butane, Propane)	0.3	0.2	0.4	0.5	0.3	0.3	0.3	0.2
Nitrous Oxide or Whippits	4.7	4.8	0.6	0.7	4.4	4.1	5.3	5.4
Spray Paints	0.6	0.6	1.4 ^a	1.8	0.8 ^a	0.6	0.5	0.5
Other Aerosol Sprays	0.8	0.8	1.0	1.0	1.4	1.2	0.6	0.6
NEEDLE USE^{1,2,3}	1.7	1.6	0.2	0.2	1.4	1.3	1.9	1.8
Heroin Needle Use ^{1,2}	1.0	0.9	0.1	0.1	0.9	0.9	1.1	1.0
Cocaine Needle Use ^{1,2}	0.9	0.9	0.0	0.1	0.5	0.4	1.0	1.0
Stimulant Needle Use ^{1,2,3}	0.9	0.8	0.1	0.1	0.6	0.6	1.1	1.0
Methamphetamine Needle Use ^{1,2,3}	0.7	0.7	0.1	0.1	0.4	0.4	0.9	0.8
HEROIN	1.8	1.8	0.2	0.1	1.8	2.0	2.0	2.0
Smoke Heroin ²	0.6	0.6	0.1	0.1	0.8	0.9	0.7	0.6
Sniff or Snort Heroin ²	1.2	1.3	0.1	0.1	1.3	1.5	1.4	1.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

³ Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.89A Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Numbers in Thousands, 2013 and 2014

Pain Reliever	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	16,907	17,195	723	810	2,465	2,400	13,719	13,985
Percocet [®] , Percodan [®] , or Tylox [®]	13,953 ^a	15,099	288	305	2,822	2,698	10,843 ^a	12,096
Vicodin [®] , Lortab [®] , or Lorcet [®]	22,315	22,075	691	620	4,950 ^a	4,531	16,673	16,924
Codeine	7,264	6,873	271	306	1,746	1,625	5,247	4,942
Demerol [®]	2,316	2,118	30	45	225	250	2,060	1,824
Dilaudid [®]	1,730	1,850	17	27	342	344	1,371	1,479
Fioricet [®]	570	483	3	16	66	61	501	406
Fiorinal [®]	401	241	11	15	34	34	356	192
Hydrocodone	11,588	11,230	368	329	2,787	2,709	8,433	8,192
Methadone	2,488	2,250	66 ^a	116	680 ^a	547	1,742	1,587
Morphine	3,792	3,680	154	167	986	958	2,652	2,555
OxyContin [®]	6,973	7,031	274	252	1,889	1,783	4,811	4,996
Phenaphen [®] with Codeine	752	646	30	35	193	163	530	448
Propoxyphene	225	300	11	17	58	43	155	240
SK-65 [®]	50	89	18	12	15	36	18	41
Stadol [®]	169	121	2	8	11	21	157	92
Talacen [®]	42	63	8	6	12	13	22	44
Talwin [®]	239	237	4	5	7	11	228	222
Talwin [®] NX	44	51	4	3	16	16	24	32
Tramadol	3,197	3,040	89	109	928	868	2,181	2,063
Ultram [®]	1,645	1,317	27	22	271	209	1,348	1,086
SELECTED GROUPS OF DRUGS¹								
Propoxyphene or Codeine Products ²	19,182	19,370	888	977	3,290	3,156	15,004	15,238
Oxycodone Products ³	15,997	17,031	476	452	3,449	3,316	12,072 ^a	13,262
Hydrocodone Products ⁴	24,413	24,330	856	785	5,538 ^a	5,181	18,019	18,365
Tramadol Products ⁵	3,786	3,527	105	124	983	905	2,699	2,498

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.89B Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Percentages, 2013 and 2014

Pain Reliever	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	6.4	6.5	2.9	3.3	7.1	6.9	6.8	6.8
Percocet [®] , Percodan [®] , or Tylox [®]	5.3	5.7	1.2	1.2	8.1	7.7	5.3 ^a	5.9
Vicodin [®] , Lortab [®] , or Lorcet [®]	8.5	8.3	2.8	2.5	14.2 ^a	13.0	8.2	8.2
Codeine	2.8	2.6	1.1	1.2	5.0	4.7	2.6	2.4
Demerol [®]	0.9	0.8	0.1	0.2	0.6	0.7	1.0	0.9
Dilaudid [®]	0.7	0.7	0.1	0.1	1.0	1.0	0.7	0.7
Fioricet [®]	0.2	0.2	0.0	0.1	0.2	0.2	0.2	0.2
Fiorinal [®]	0.2	0.1	0.0	0.1	0.1	0.1	0.2	0.1
Hydrocodone	4.4	4.2	1.5	1.3	8.0	7.8	4.2	4.0
Methadone	0.9	0.8	0.3 ^a	0.5	2.0 ^a	1.6	0.9	0.8
Morphine	1.4	1.4	0.6	0.7	2.8	2.7	1.3	1.2
OxyContin [®]	2.7	2.7	1.1	1.0	5.4	5.1	2.4	2.4
Phenaphen [®] with Codeine	0.3	0.2	0.1	0.1	0.6	0.5	0.3	0.2
Propoxyphene	0.1	0.1	0.0	0.1	0.2	0.1	0.1	0.1
SK-65 [®]	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.0
Stadol [®]	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.0
Talacen [®]	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Talwin [®]	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.1
Talwin [®] NX	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Tramadol	1.2	1.1	0.4	0.4	2.7	2.5	1.1	1.0
Ultram [®]	0.6 ^a	0.5	0.1	0.1	0.8	0.6	0.7	0.5
SELECTED GROUPS OF DRUGS¹								
Propoxyphene or Codeine Products ²	7.3	7.3	3.6	3.9	9.5	9.0	7.4	7.4
Oxycodone Products ³	6.1	6.4	1.9	1.8	9.9	9.5	6.0	6.5
Hydrocodone Products ⁴	9.3	9.2	3.4	3.2	15.9 ^a	14.8	8.9	8.9
Tramadol Products ⁵	1.4	1.3	0.4	0.5	2.8	2.6	1.3	1.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.90A Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Numbers in Thousands, 2013 and 2014

Tranquilizer	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Klonopin® or Clonazepam	6,112	6,557	143	143	1,538	1,545	4,431	4,869
Xanax®, Alprazolam, Ativan®, or Lorazepam	15,529	16,209	405 ^a	527	3,594	3,685	11,531	11,997
Valium® or Diazepam	12,471	13,483	129	149	1,600	1,600	10,742	11,735
Atarax®	241	229	11	11	34	42	197	176
BuSpar®	570	748	14	16	155	165	401	568
Equanil®	87	65	3	11	24	11	61	43
Flexeril®	2,878	3,066	28	41	489	453	2,361	2,572
Librium®	638	437	10	16	56	34	571	387
Limbitrol®	38	29	9	5	13	12	16	12
Meprobamate	125	206	7	16	16	28	102	163
Miltown®	76	96	5	3	18	10	53	82
Rohypnol®	348	206	12	6	64	70	272	131
Serax®	62	65	2	14	14	14	45	37
Soma®	3,312	2,978	69	41	693 ^a	540	2,550	2,397
Tranxene®	73	124	4	*	15	17	53	107
Vistaril®	329	365	9	18	78	69	241	279
SELECTED GROUPS OF DRUGS¹								
Benzodiazepines ²	22,096	23,235	498 ^a	612	4,043	4,171	17,555	18,453
Meprobamate Products ³	234	318	11	25	41	39	183	253
Muscle Relaxants ⁴	5,099	4,887	89	82	954	812	4,055	3,993

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan® or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

³ Includes Equanil®, meprobamate, and Miltown®.

⁴ Includes Flexeril® and Soma®.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.90B Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Percentages, 2013 and 2014

Tranquilizer	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Klonopin® or Clonazepam	2.3	2.5	0.6	0.6	4.4	4.4	2.2	2.4
Xanax®, Alprazolam, Ativan®, or Lorazepam	5.9	6.1	1.6 ^a	2.1	10.3	10.5	5.7	5.8
Valium® or Diazepam	4.8	5.1	0.5	0.6	4.6	4.6	5.3	5.7
Atarax®	0.1	0.1	0.0	0.0	0.1	0.1	0.1	0.1
BuSpar®	0.2	0.3	0.1	0.1	0.4	0.5	0.2	0.3
Equanil®	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
Flexeril®	1.1	1.2	0.1	0.2	1.4	1.3	1.2	1.3
Librium®	0.2	0.2	0.0	0.1	0.2	0.1	0.3	0.2
Limbitrol®	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Meprobamate	0.0	0.1	0.0	0.1	0.0	0.1	0.1	0.1
Miltown®	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
Rohypnol®	0.1	0.1	0.0	0.0	0.2	0.2	0.1	0.1
Serax®	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
Soma®	1.3	1.1	0.3	0.2	2.0 ^a	1.5	1.3	1.2
Tranxene®	0.0	0.0	0.0	*	0.0	0.0	0.0	0.1
Vistaril®	0.1	0.1	0.0	0.1	0.2	0.2	0.1	0.1
SELECTED GROUPS OF DRUGS¹								
Benzodiazepines ²	8.4	8.8	2.0 ^a	2.5	11.6	11.9	8.7	9.0
Meprobamate Products ³	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1
Muscle Relaxants ⁴	1.9	1.8	0.4	0.3	2.7 ^a	2.3	2.0	1.9

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan® or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

³ Includes Equanil®, meprobamate, and Miltown®.

⁴ Includes Flexeril® and Soma®.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.91A Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Numbers in Thousands, 2013 and 2014

Stimulant	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Methamphetamine, Desoxyn®, or Methedrine ¹	12,257	12,943	119 ^a	171	1,042	1,040	11,096	11,731
Prescription Diet Pills ²	7,175	7,691	94	124	594 ^a	722	6,487	6,844
Ritalin® or Methylphenidate	6,397	6,645	217	211	1,824	1,642	4,355	4,792
Cylert®	147	155	8	19	16	30	124	107
Dexedrine®	1,922	1,956	11	15	69	104	1,841	1,837
Dextroamphetamine	993	1,222	31	32	166 ^a	248	796	942
Didrex®	76	165	3	8	22	10	51 ^a	146
Eskatrol®	86	149	3	6	15	10	69	133
Ionamin®	313	227	9	10	11	8	294	210
Mazanor®	16	48	3	20	3	3	11	26
Obedrin-LA®	15	18	3	4	9	2	*	12
Plegine®	27	27	5 ^a	27	3	*	19	*
Preludin®	223	327	*	10	14	6	210	311
Sanorex®	*	113	2 ^b	23	5	11	*	79
Tenuate®	99	147	4 ^a	21	8	9	86	117
SELECTED GROUPS OF DRUGS³								
Amphetamine, Dextroamphetamine, and Phentermine Products ⁴	9,819	10,372	199	224	1,467	1,544	8,153	8,605
Mazindol Products ⁵	22 ^b	134	4 ^b	33	7	14	11 ^b	87
Methylphenidate or Dexmethylphenidate Products ⁶	6,447	6,674	224	212	1,853	1,650	4,370	4,812

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn®, or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine®, Biphedamine®, Fastin®, or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine®, dextroamphetamine, and Ionamin®.

⁵ Includes Mazanor® and Sanorex®.

⁶ Includes Ritalin®, other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.91B Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Percentages, 2013 and 2014

Stimulant	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Methamphetamine, Desoxyn®, or Methedrine ¹	4.7	4.9	0.5 ^a	0.7	3.0	3.0	5.5	5.7
Prescription Diet Pills ²	2.7	2.9	0.4	0.5	1.7 ^a	2.1	3.2	3.3
Ritalin® or Methylphenidate	2.4	2.5	0.9	0.8	5.2	4.7	2.1	2.3
Cylert®	0.1	0.1	0.0	0.1	0.0	0.1	0.1	0.1
Dexedrine®	0.7	0.7	0.0	0.1	0.2	0.3	0.9	0.9
Dextroamphetamine	0.4	0.5	0.1	0.1	0.5 ^a	0.7	0.4	0.5
Didrex®	0.0	0.1	0.0	0.0	0.1	0.0	0.0 ^a	0.1
Eskatrol®	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1
Ionamin®	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.1
Mazanor®	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
Obedrin-LA®	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0
Plegine®	0.0	0.0	0.0 ^a	0.1	0.0	*	0.0	*
Preludin®	0.1	0.1	*	0.0	0.0	0.0	0.1	0.2
Sanorex®	*	0.0	0.0 ^b	0.1	0.0	0.0	*	0.0
Tenuate®	0.0	0.1	0.0 ^a	0.1	0.0	0.0	0.0	0.1
SELECTED GROUPS OF DRUGS³								
Amphetamine, Dextroamphetamine, and Phentermine Products ⁴	3.7	3.9	0.8	0.9	4.2	4.4	4.0	4.2
Mazindol Products ⁵	0.0 ^b	0.1	0.0 ^b	0.1	0.0	0.0	0.0 ^b	0.0
Methylphenidate or Dexmethylphenidate Products ⁶	2.5	2.5	0.9	0.9	5.3 ^a	4.7	2.2	2.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn®, or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine®, Biphedamine®, Fastin®, or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine®, dextroamphetamine, and Ionamin®.

⁵ Includes Mazanor® and Sanorex®.

⁶ Includes Ritalin®, other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.92A Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Numbers in Thousands, 2013 and 2014

Sedative	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Methaqualone, Sopor [®] , or Quaalude [®]	4,811	4,906	21	36	80	63	4,710	4,808
Barbiturates ¹	1,979	2,071	11 ^a	34	75	56	1,892	1,980
Restoril [®] or Temazepam	910	1,039	31	37	143	152	737	850
Amytal [®]	125	190	7	14	22	6	96	170
Butisol [®]	26 ^a	95	4	10	9	8	13 ^a	77
Chloral Hydrate	99	132	5	9	10	*	84	123
Dalmane [®]	251	170	4	14	16	3	230	154
Halcion [®]	459	343	8	15	29	38	422	290
Phenobarbital	848	921	16	22	71	60	762	839
Placidyl [®]	542	527	3	10	7	3	532	514
Tuinal [®]	701	649	5	7	4	*	691	641
SELECTED GROUPS OF DRUGS²								
Temazepam, Flurazepam, or Triazolam ³	1,436	1,415	42	57	170	181	1,225	1,177
Any Barbiturates ⁴	2,466	2,715	38 ^a	69	126	101	2,303	2,546

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.92B Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Percentages, 2013 and 2014

Sedative	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Methaqualone, Sopor [®] , or Quaalude [®]	1.8	1.9	0.1	0.1	0.2	0.2	2.3	2.3
Barbiturates ¹	0.8	0.8	0.0 ^a	0.1	0.2	0.2	0.9	1.0
Restoril [®] or Temazepam	0.3	0.4	0.1	0.1	0.4	0.4	0.4	0.4
Amytal [®]	0.0	0.1	0.0	0.1	0.1	0.0	0.0	0.1
Butisol [®]	0.0 ^a	0.0	0.0	0.0	0.0	0.0	0.0 ^a	0.0
Chloral Hydrate	0.0	0.0	0.0	0.0	0.0	*	0.0	0.1
Dalmane [®]	0.1	0.1	0.0	0.1	0.0	0.0	0.1	0.1
Halcion [®]	0.2	0.1	0.0	0.1	0.1	0.1	0.2	0.1
Phenobarbital	0.3	0.3	0.1	0.1	0.2	0.2	0.4	0.4
Placidyl [®]	0.2	0.2	0.0	0.0	0.0	0.0	0.3	0.3
Tuinal [®]	0.3	0.2	0.0	0.0	0.0	*	0.3	0.3
SELECTED GROUPS OF DRUGS²								
Temazepam, Flurazepam, or Triazolam ³	0.5	0.5	0.2	0.2	0.5	0.5	0.6	0.6
Any Barbiturates ⁴	0.9	1.0	0.2 ^a	0.3	0.4	0.3	1.1	1.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.1C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	1,015	873	677	595	509	476
Marijuana and Hashish	1,017	872	577	538	442	424
Cocaine	693	632	189	186	133	105
Crack	364	317	77	81	64	55
Heroin	281	213	70	75	43	55
Hallucinogens	707	636	204	159	132	88
LSD	580	508	81	76	37	40
PCP	346	274	21	16	15	*
Ecstasy	464	384	153	121	91	65
Inhalants	519	428	105	95	64	65
Nonmedical Use of Psychotherapeutics ^{2,3}	785	658	418	321	256	222
Pain Relievers	640	553	343	253	206	174
OxyContin [®]	280	220	119	83	81	50
Tranquilizers	576	443	230	190	121	115
Stimulants ³	501	441	178	159	116	116
Methamphetamine ³	393	363	114	104	85	78
Sedatives	377	324	84	75	67	54
ILLCIT DRUGS OTHER THAN MARIJUANA¹	923	803	469	371	306	261

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.1D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.39	0.33	0.26	0.22	0.19	0.18
Marijuana and Hashish	0.39	0.33	0.22	0.20	0.17	0.16
Cocaine	0.26	0.24	0.07	0.07	0.05	0.04
Crack	0.14	0.12	0.03	0.03	0.02	0.02
Heroin	0.11	0.08	0.03	0.03	0.02	0.02
Hallucinogens	0.27	0.24	0.08	0.06	0.05	0.03
LSD	0.22	0.19	0.03	0.03	0.01	0.02
PCP	0.13	0.10	0.01	0.01	0.01	*
Ecstasy	0.18	0.14	0.06	0.05	0.03	0.02
Inhalants	0.20	0.16	0.04	0.04	0.02	0.02
Nonmedical Use of Psychotherapeutics ^{2,3}	0.30	0.25	0.16	0.12	0.10	0.08
Pain Relievers	0.24	0.21	0.13	0.10	0.08	0.07
OxyContin [®]	0.11	0.08	0.05	0.03	0.03	0.02
Tranquilizers	0.22	0.17	0.09	0.07	0.05	0.04
Stimulants ³	0.19	0.17	0.07	0.06	0.04	0.04
Methamphetamine ³	0.15	0.14	0.04	0.04	0.03	0.03
Sedatives	0.14	0.12	0.03	0.03	0.03	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.35	0.30	0.18	0.14	0.12	0.10

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.1P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.2511		0.0189		0.0022	
Marijuana and Hashish	0.3327		0.0214		0.0004	
Cocaine	0.2134		0.2191		0.8370	
Crack	0.3420		0.2313		0.7509	
Heroin	0.8887		0.0295		0.0413	
Hallucinogens	0.5990		0.3865		0.2769	
LSD	0.9928		0.1356		0.3098	
PCP	0.7509		0.9760		*	
Ecstasy	0.4555		0.1639		0.6122	
Inhalants	0.9936		0.6320		0.6223	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.5162		0.3072		0.9662	
Pain Relievers	0.7945		0.0451		0.3709	
OxyContin [®]	0.9676		0.1321		0.1404	
Tranquilizers	0.1285		0.6857		0.3646	
Stimulants ³	0.3336		0.4362		0.1908	
Methamphetamine ³	0.2996		0.5088		0.7826	
Sedatives	0.5922		0.2542		0.3799	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.3178		0.5555		0.9303	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.2C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	43	47	33	39	20	25
Marijuana and Hashish	23	23	21	22	12	15
Cocaine	4	6	3	1	1	*
Crack	1	1	1	*	1	*
Heroin	4	1	3	1	1	*
Hallucinogens	11	10	8	7	4	4
LSD	5	4	4	2	3	*
PCP	3	3	2	2	1	*
Ecstasy	6	3	6	3	1	2
Inhalants	26	29	16	19	9	11
Nonmedical Use of Psychotherapeutics ^{2,3}	27	32	19	26	14	19
Pain Relievers	24	27	17	22	12	16
OxyContin [®]	6	6	5	6	4	5
Tranquilizers	9	12	7	8	6	3
Stimulants ³	10	14	6	13	4	11
Methamphetamine ³	4	7	3	6	2	5
Sedatives	7	11	6	8	5	5
ILLCIT DRUGS OTHER THAN MARIJUANA¹	37	41	26	32	17	22

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.2D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.49	0.55	0.39	0.47	0.24	0.31
Marijuana and Hashish	0.28	0.29	0.26	0.28	0.15	0.19
Cocaine	0.05	0.07	0.04	0.02	0.02	*
Crack	0.02	0.02	0.02	*	0.02	*
Heroin	0.05	0.02	0.04	0.02	0.01	*
Hallucinogens	0.14	0.12	0.11	0.09	0.05	0.05
LSD	0.06	0.04	0.05	0.02	0.04	*
PCP	0.04	0.04	0.02	0.02	0.01	*
Ecstasy	0.08	0.04	0.08	0.04	0.01	0.03
Inhalants	0.31	0.35	0.20	0.24	0.12	0.14
Nonmedical Use of Psychotherapeutics ^{2,3}	0.33	0.40	0.23	0.33	0.17	0.24
Pain Relievers	0.30	0.34	0.21	0.28	0.15	0.20
OxyContin [®]	0.08	0.08	0.06	0.07	0.05	0.06
Tranquilizers	0.11	0.15	0.09	0.11	0.07	0.04
Stimulants ³	0.12	0.17	0.07	0.16	0.05	0.14
Methamphetamine ³	0.05	0.09	0.03	0.07	0.03	0.07
Sedatives	0.09	0.15	0.08	0.10	0.06	0.07
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.44	0.50	0.31	0.40	0.21	0.27

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.2P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.3337		0.0522		0.0471	
Marijuana and Hashish	0.4822		0.7596		0.6560	
Cocaine	0.8939		0.0766		*	
Crack	0.7690		*		*	
Heroin	0.2611		0.3912		*	
Hallucinogens	0.4685		0.3946		0.8019	
LSD	0.2467		0.1253		*	
PCP	0.9206		0.7964		*	
Ecstasy	0.1409		0.1739		0.2497	
Inhalants	0.8163		0.4459		0.5216	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0198		0.0020		0.0576	
Pain Relievers	0.1446		0.0259		0.1575	
OxyContin [®]	0.6217		0.9469		0.9554	
Tranquilizers	0.1171		0.5632		0.1202	
Stimulants ³	0.0678		0.0134		0.0699	
Methamphetamine ³	0.3139		0.3796		0.4787	
Sedatives	0.0528		0.2948		0.8544	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.1102		0.0049		0.0310	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.3C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	67	63	55	57	37	39
Marijuana and Hashish	51	51	45	46	31	32
Cocaine	7	10	7	9	5	6
Crack	3	5	2	3	*	1
Heroin	4	2	4	2	3	2
Hallucinogens	15	20	12	17	8	8
LSD	8	15	6	13	3	6
PCP	5	8	4	7	3	*
Ecstasy	10	14	7	12	4	5
Inhalants	37	35	19	24	11	14
Nonmedical Use of Psychotherapeutics ^{2,3}	40	39	33	35	20	24
Pain Relievers	38	37	31	31	20	21
OxyContin [®]	11	14	9	12	3	6
Tranquilizers	18	21	12	17	7	7
Stimulants ³	16	18	11	16	5	14
Methamphetamine ³	6	11	4	10	1	7
Sedatives	9	15	6	12	4	4
ILLICIT DRUGS OTHER THAN MARIJUANA¹	52	50	37	41	23	28

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.3D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.68	0.67	0.60	0.61	0.42	0.44
Marijuana and Hashish	0.55	0.56	0.50	0.51	0.36	0.36
Cocaine	0.09	0.12	0.08	0.11	0.06	0.07
Crack	0.03	0.06	0.02	0.04	*	0.01
Heroin	0.05	0.02	0.05	0.02	0.04	0.02
Hallucinogens	0.18	0.23	0.14	0.20	0.10	0.10
LSD	0.09	0.17	0.07	0.16	0.03	0.07
PCP	0.06	0.09	0.05	0.08	0.03	*
Ecstasy	0.12	0.17	0.08	0.14	0.05	0.06
Inhalants	0.42	0.41	0.23	0.28	0.12	0.16
Nonmedical Use of Psychotherapeutics ^{2,3}	0.44	0.45	0.37	0.40	0.24	0.28
Pain Relievers	0.42	0.42	0.34	0.36	0.23	0.25
OxyContin [®]	0.13	0.17	0.11	0.14	0.04	0.07
Tranquilizers	0.21	0.24	0.14	0.20	0.08	0.08
Stimulants ³	0.18	0.21	0.13	0.19	0.06	0.17
Methamphetamine ³	0.07	0.13	0.05	0.12	0.01	0.08
Sedatives	0.10	0.17	0.07	0.15	0.04	0.05
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.56	0.56	0.41	0.46	0.27	0.32

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.3P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.8876		0.6440		0.9287	
Marijuana and Hashish	0.6118		0.8316		0.6416	
Cocaine	0.6910		0.6525		0.9197	
Crack	0.5622		0.5820		*	
Heroin	0.0880		0.0909		0.4474	
Hallucinogens	0.7553		0.4688		0.5374	
LSD	0.0692		0.0399		0.3263	
PCP	0.2497		0.2792		*	
Ecstasy	1.0000		0.5028		0.9399	
Inhalants	0.8840		0.1609		0.0662	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.8765		0.8505		0.2681	
Pain Relievers	0.7074		0.7789		0.5472	
OxyContin [®]	0.7269		0.5586		0.4589	
Tranquilizers	0.7978		0.2727		0.8903	
Stimulants ³	0.9127		0.5135		0.0187	
Methamphetamine ³	0.0781		0.0447		0.0563	
Sedatives	0.1607		0.0873		0.5927	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.9669		0.4279		0.1316	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.4C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	78	89	67	77	49	60
Marijuana and Hashish	70	83	63	72	46	57
Cocaine	17	22	13	21	8	8
Crack	4	7	3	6	1	4
Heroin	7	8	5	8	3	6
Hallucinogens	29	30	25	26	14	17
LSD	17	23	14	20	9	13
PCP	5	7	5	6	1	2
Ecstasy	22	22	17	19	8	9
Inhalants	28	32	17	17	6	4
Nonmedical Use of Psychotherapeutics ^{2,3}	48	53	38	45	23	29
Pain Relievers	44	46	34	38	19	23
OxyContin [®]	18	18	12	14	4	5
Tranquilizers	26	34	19	28	9	14
Stimulants ³	22	30	17	25	10	14
Methamphetamine ³	12	17	9	13	6	8
Sedatives	8	15	6	11	1	8
ILLCIT DRUGS OTHER THAN MARIJUANA¹	56	60	44	50	26	33

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.4D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.74	0.78	0.69	0.73	0.54	0.64
Marijuana and Hashish	0.72	0.76	0.67	0.69	0.51	0.61
Cocaine	0.20	0.25	0.15	0.24	0.09	0.10
Crack	0.05	0.08	0.03	0.07	0.01	0.05
Heroin	0.08	0.10	0.06	0.09	0.04	0.07
Hallucinogens	0.34	0.34	0.29	0.30	0.17	0.20
LSD	0.20	0.26	0.17	0.23	0.11	0.15
PCP	0.06	0.09	0.06	0.06	0.02	0.02
Ecstasy	0.26	0.26	0.20	0.22	0.09	0.10
Inhalants	0.32	0.38	0.20	0.20	0.07	0.05
Nonmedical Use of Psychotherapeutics ^{2,3}	0.52	0.56	0.42	0.49	0.27	0.33
Pain Relievers	0.48	0.50	0.38	0.42	0.23	0.26
OxyContin [®]	0.21	0.20	0.14	0.16	0.05	0.06
Tranquilizers	0.31	0.38	0.22	0.32	0.11	0.16
Stimulants ³	0.25	0.34	0.20	0.29	0.11	0.17
Methamphetamine ³	0.14	0.20	0.11	0.15	0.07	0.09
Sedatives	0.10	0.18	0.07	0.13	0.02	0.09
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.57	0.62	0.48	0.54	0.30	0.38

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.4P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.4397		0.2569		0.3728	
Marijuana and Hashish	0.7329		0.3163		0.3115	
Cocaine	0.9770		0.1305		0.8726	
Crack	0.2639		0.1944		0.1552	
Heroin	0.4402		0.4557		0.2813	
Hallucinogens	0.0205		0.0506		0.6460	
LSD	0.1054		0.1047		0.4688	
PCP	0.6637		0.6466		0.8118	
Ecstasy	0.0354		0.0390		0.3284	
Inhalants	0.6214		0.2329		0.0958	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.9154		1.0000		0.4688	
Pain Relievers	0.4852		0.3535		0.9015	
OxyContin [®]	0.3123		0.9199		0.9050	
Tranquilizers	0.1494		0.1606		0.1601	
Stimulants ³	0.1296		0.1892		0.1063	
Methamphetamine ³	0.2917		0.7633		0.6157	
Sedatives	0.0728		0.1627		0.0277	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.4723		0.3086		0.7406	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.5C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	169	180	143	154	103	117
Marijuana and Hashish	160	169	134	145	97	111
Cocaine	51	52	39	37	18	21
Crack	16	16	6	12	2	2
Heroin	20	21	13	17	9	9
Hallucinogens	75	73	56	55	33	25
LSD	41	48	31	33	15	10
PCP	13	18	5	7	1	2
Ecstasy	63	58	40	40	17	18
Inhalants	49	47	27	25	14	11
Nonmedical Use of Psychotherapeutics ^{2,3}	98	109	74	73	42	41
Pain Relievers	87	91	63	57	35	32
OxyContin [®]	42	37	26	19	13	6
Tranquilizers	61	69	41	38	18	20
Stimulants ³	52	53	37	40	21	19
Methamphetamine ³	25	28	16	18	8	10
Sedatives	22	20	13	12	6	7
ILLCIT DRUGS OTHER THAN MARIJUANA¹	113	121	89	87	54	51

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.5D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.79	0.90	0.76	0.87	0.64	0.73
Marijuana and Hashish	0.78	0.90	0.74	0.84	0.62	0.71
Cocaine	0.37	0.39	0.29	0.28	0.13	0.16
Crack	0.12	0.12	0.05	0.09	0.02	0.01
Heroin	0.15	0.16	0.10	0.13	0.07	0.07
Hallucinogens	0.52	0.53	0.40	0.41	0.24	0.19
LSD	0.31	0.36	0.23	0.25	0.12	0.08
PCP	0.10	0.14	0.04	0.06	0.01	0.01
Ecstasy	0.45	0.44	0.29	0.30	0.13	0.14
Inhalants	0.35	0.35	0.20	0.19	0.11	0.08
Nonmedical Use of Psychotherapeutics ^{2,3}	0.62	0.72	0.50	0.51	0.31	0.30
Pain Relievers	0.58	0.62	0.44	0.41	0.26	0.23
OxyContin [®]	0.31	0.28	0.19	0.14	0.10	0.05
Tranquilizers	0.44	0.50	0.30	0.28	0.14	0.15
Stimulants ³	0.37	0.39	0.26	0.30	0.16	0.15
Methamphetamine ³	0.18	0.21	0.12	0.14	0.06	0.07
Sedatives	0.17	0.15	0.10	0.09	0.05	0.05
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.68	0.77	0.58	0.60	0.38	0.37

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.5P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.8918		0.5799		0.9036	
Marijuana and Hashish	0.8608		0.9952		0.5779	
Cocaine	0.0991		0.5988		0.3667	
Crack	0.5755		0.2330		0.8933	
Heroin	0.9515		0.4853		0.8694	
Hallucinogens	0.0657		0.2498		0.0079	
LSD	0.2277		0.6595		0.0673	
PCP	0.6208		0.5879		0.9456	
Ecstasy	0.0134		0.0431		0.3105	
Inhalants	0.0322		0.1228		0.3506	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.2140		0.0836		0.0849	
Pain Relievers	0.0313		0.0205		0.0422	
OxyContin [®]	0.0875		0.6309		0.1371	
Tranquilizers	0.7694		0.2114		0.6883	
Stimulants ³	0.1002		0.5594		0.2748	
Methamphetamine ³	0.8692		0.8245		0.8216	
Sedatives	0.0396		0.6169		0.8476	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.6296		0.1221		0.0301	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.6C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	252	260	182	192	127	149
Marijuana and Hashish	233	251	170	181	119	138
Cocaine	105	116	63	72	29	35
Crack	37	38	14	19	12	9
Heroin	39	45	23	26	13	14
Hallucinogens	134	137	76	75	35	36
LSD	74	82	39	44	9	16
PCP	27	25	4	6	2	2
Ecstasy	111	117	57	59	26	27
Inhalants	73	84	30	28	12	12
Nonmedical Use of Psychotherapeutics ^{2,3}	162	169	105	106	59	63
Pain Relievers	134	143	87	80	49	51
OxyContin [®]	69	71	34	32	13	18
Tranquilizers	104	114	62	59	29	31
Stimulants ³	94	96	55	60	32	30
Methamphetamine ³	53	53	26	31	17	12
Sedatives	29	34	16	18	6	12
ILLCIT DRUGS OTHER THAN MARIJUANA¹	190	196	124	129	69	79

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.6D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.58	0.67	0.59	0.64	0.51	0.56
Marijuana and Hashish	0.59	0.69	0.57	0.63	0.47	0.54
Cocaine	0.43	0.47	0.28	0.31	0.13	0.16
Crack	0.17	0.17	0.07	0.09	0.05	0.04
Heroin	0.18	0.20	0.11	0.12	0.06	0.06
Hallucinogens	0.50	0.53	0.33	0.32	0.16	0.16
LSD	0.32	0.35	0.18	0.20	0.04	0.07
PCP	0.13	0.11	0.02	0.03	0.01	0.01
Ecstasy	0.44	0.46	0.25	0.26	0.12	0.12
Inhalants	0.31	0.36	0.14	0.12	0.06	0.06
Nonmedical Use of Psychotherapeutics ^{2,3}	0.55	0.60	0.41	0.43	0.26	0.27
Pain Relievers	0.49	0.54	0.36	0.34	0.22	0.22
OxyContin [®]	0.30	0.31	0.15	0.14	0.06	0.08
Tranquilizers	0.42	0.46	0.27	0.26	0.13	0.14
Stimulants ³	0.38	0.39	0.24	0.27	0.14	0.14
Methamphetamine ³	0.24	0.23	0.12	0.14	0.08	0.05
Sedatives	0.13	0.15	0.07	0.08	0.03	0.06
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.58	0.64	0.47	0.50	0.30	0.33

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.6P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	0.1595		0.3290		0.3921	
Marijuana and Hashish	0.1633		0.5459		0.4977	
Cocaine	0.6013		0.3879		0.2794	
Crack	0.8610		0.5175		0.5955	
Heroin	0.2993		0.8374		0.6640	
Hallucinogens	0.2200		0.5782		0.8411	
LSD	0.2616		0.2298		0.0111	
PCP	0.3506		0.4863		0.8005	
Ecstasy	0.6629		0.4920		0.4802	
Inhalants	0.6866		0.2746		0.5885	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.7280		0.8062		0.8023	
Pain Relievers	1.0000		0.1606		0.3364	
OxyContin [®]	0.8034		0.4638		0.4758	
Tranquilizers	0.1907		0.4510		0.8073	
Stimulants ³	0.6456		0.2948		0.5179	
Methamphetamine ³	0.8351		0.3809		0.1145	
Sedatives	0.0722		0.3386		0.0837	
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.9612		0.6400		0.5938	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.7C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	350	246	291	217	254	184
Marijuana and Hashish	352	259	267	207	231	170
Cocaine	255	196	119	88	83	54
Crack	137	97	35	28	27	22
Heroin	110	83	46	48	30	33
Hallucinogens	283	206	108	86	54	50
LSD	216	147	42	39	18	30
PCP	87	56	11	3	*	*
Ecstasy	259	183	96	63	45	40
Inhalants	206	153	46	49	30	34
Nonmedical Use of Psychotherapeutics ^{2,3}	316	216	206	139	149	107
Pain Relievers	275	194	169	108	123	78
OxyContin [®]	174	119	64	38	34	15
Tranquilizers	252	180	143	89	79	63
Stimulants ³	234	157	124	79	79	50
Methamphetamine ³	169	120	59	51	48	29
Sedatives	112	68	54	30	47	15
ILLICIT DRUGS OTHER THAN MARIJUANA¹	348	242	219	154	163	121

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.7D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.94	0.65	0.78	0.57	0.68	0.49
Marijuana and Hashish	0.94	0.68	0.72	0.55	0.62	0.45
Cocaine	0.68	0.52	0.32	0.23	0.22	0.14
Crack	0.37	0.26	0.09	0.07	0.07	0.06
Heroin	0.29	0.22	0.12	0.13	0.08	0.09
Hallucinogens	0.76	0.54	0.29	0.23	0.14	0.13
LSD	0.58	0.39	0.11	0.10	0.05	0.08
PCP	0.23	0.15	0.03	0.01	*	*
Ecstasy	0.69	0.48	0.26	0.17	0.12	0.11
Inhalants	0.55	0.40	0.12	0.13	0.08	0.09
Nonmedical Use of Psychotherapeutics ^{2,3}	0.85	0.57	0.55	0.37	0.40	0.28
Pain Relievers	0.74	0.51	0.45	0.29	0.33	0.21
OxyContin [®]	0.47	0.32	0.17	0.10	0.09	0.04
Tranquilizers	0.67	0.48	0.38	0.24	0.21	0.17
Stimulants ³	0.63	0.42	0.33	0.21	0.21	0.13
Methamphetamine ³	0.45	0.32	0.16	0.13	0.13	0.08
Sedatives	0.30	0.18	0.14	0.08	0.13	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.93	0.64	0.59	0.41	0.44	0.32

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.7P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.6450		0.6147		0.8673	
Marijuana and Hashish	0.3049		0.5781		0.9074	
Cocaine	0.2432		0.3075		0.2638	
Crack	0.5612		0.6677		0.6501	
Heroin	0.8985		0.1628		0.2568	
Hallucinogens	0.0778		0.4963		0.3494	
LSD	0.0026		0.5418		0.3011	
PCP	0.3218		0.4545		*	
Ecstasy	0.1902		0.7513		0.8519	
Inhalants	0.0410		0.1329		0.2846	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.1889		0.0256		0.4392	
Pain Relievers	0.0902		0.0354		0.3142	
OxyContin [®]	0.8010		0.1561		0.2170	
Tranquilizers	0.5649		0.2156		0.8005	
Stimulants ³	0.2962		0.2471		0.4869	
Methamphetamine ³	0.2643		0.7619		0.3459	
Sedatives	0.4627		0.4985		0.2938	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0555		0.0866		0.4411	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.8C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	896	754	508	444	383	350
Marijuana and Hashish	912	740	412	381	324	299
Cocaine	626	567	125	136	98	84
Crack	332	295	66	73	57	49
Heroin	247	187	41	48	25	40
Hallucinogens	604	542	139	76	111	52
LSD	527	458	50	19	25	13
PCP	327	266	15	10	15	*
Ecstasy	334	268	95	57	72	38
Inhalants	460	379	78	66	50	45
Nonmedical Use of Psychotherapeutics ^{2,3}	680	567	317	246	190	178
Pain Relievers	540	464	264	198	156	145
OxyContin [®]	191	155	81	63	69	42
Tranquilizers	481	378	162	137	81	84
Stimulants ³	437	391	107	115	71	97
Methamphetamine ³	357	329	85	81	65	70
Sedatives	359	301	61	60	48	48
ILLCIT DRUGS OTHER THAN MARIJUANA¹	798	698	360	285	240	211

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.8D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.54	0.45	0.31	0.27	0.23	0.21
Marijuana and Hashish	0.55	0.44	0.25	0.23	0.20	0.18
Cocaine	0.38	0.34	0.08	0.08	0.06	0.05
Crack	0.20	0.18	0.04	0.04	0.03	0.03
Heroin	0.15	0.11	0.02	0.03	0.01	0.02
Hallucinogens	0.37	0.32	0.08	0.05	0.07	0.03
LSD	0.32	0.27	0.03	0.01	0.02	0.01
PCP	0.20	0.16	0.01	0.01	0.01	*
Ecstasy	0.20	0.16	0.06	0.03	0.04	0.02
Inhalants	0.28	0.23	0.05	0.04	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{2,3}	0.41	0.34	0.19	0.15	0.12	0.11
Pain Relievers	0.33	0.28	0.16	0.12	0.09	0.09
OxyContin [®]	0.12	0.09	0.05	0.04	0.04	0.02
Tranquilizers	0.29	0.23	0.10	0.08	0.05	0.05
Stimulants ³	0.26	0.23	0.06	0.07	0.04	0.06
Methamphetamine ³	0.22	0.20	0.05	0.05	0.04	0.04
Sedatives	0.22	0.18	0.04	0.04	0.03	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.48	0.42	0.22	0.17	0.14	0.13

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.8P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	0.2452		0.0011		0.0002	
Marijuana and Hashish	0.2298		0.0003		0.0000	
Cocaine	0.0500		0.0351		0.9590	
Crack	0.2193		0.2527		0.9021	
Heroin	0.6727		0.0770		0.0364	
Hallucinogens	0.5111		0.5175		0.4019	
LSD	0.4779		0.7622		0.8292	
PCP	0.9241		0.7571		*	
Ecstasy	0.5051		0.9105		0.9348	
Inhalants	0.2291		0.6934		0.9294	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.2056		0.8996		0.5848	
Pain Relievers	0.1533		0.7744		0.9090	
OxyContin [®]	0.7878		0.4704		0.2480	
Tranquilizers	0.0972		0.5024		0.2531	
Stimulants ³	0.0892		0.1559		0.0343	
Methamphetamine ³	0.1385		0.8879		0.7299	
Sedatives	0.5916		0.2204		0.1827	
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.0619		0.7742		0.7550	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.9C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	702	569	472	428	363	360
Marijuana and Hashish	717	570	416	399	321	320
Cocaine	508	470	154	145	101	84
Crack	264	253	59	63	47	41
Heroin	228	182	58	63	33	45
Hallucinogens	524	461	167	129	119	67
LSD	447	381	57	59	24	32
PCP	267	225	14	12	3	4
Ecstasy	339	286	116	93	82	45
Inhalants	416	346	95	83	56	57
Nonmedical Use of Psychotherapeutics ^{2,3}	529	460	276	228	180	159
Pain Relievers	434	401	227	184	146	133
OxyContin [®]	205	186	104	63	73	42
Tranquilizers	373	320	159	137	79	82
Stimulants ³	380	325	123	112	86	80
Methamphetamine ³	322	275	97	74	78	51
Sedatives	268	241	59	57	48	43
ILLCIT DRUGS OTHER THAN MARIJUANA¹	624	548	330	273	231	192

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.9D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.55	0.44	0.37	0.33	0.29	0.28
Marijuana and Hashish	0.56	0.44	0.33	0.31	0.25	0.25
Cocaine	0.40	0.37	0.12	0.11	0.08	0.07
Crack	0.21	0.20	0.05	0.05	0.04	0.03
Heroin	0.18	0.14	0.05	0.05	0.03	0.04
Hallucinogens	0.41	0.36	0.13	0.10	0.09	0.05
LSD	0.35	0.30	0.04	0.05	0.02	0.02
PCP	0.21	0.18	0.01	0.01	0.00	0.00
Ecstasy	0.27	0.22	0.09	0.07	0.06	0.03
Inhalants	0.33	0.27	0.07	0.06	0.04	0.04
Nonmedical Use of Psychotherapeutics ^{2,3}	0.42	0.36	0.22	0.18	0.14	0.12
Pain Relievers	0.34	0.31	0.18	0.14	0.11	0.10
OxyContin [®]	0.16	0.14	0.08	0.05	0.06	0.03
Tranquilizers	0.29	0.25	0.13	0.11	0.06	0.06
Stimulants ³	0.30	0.25	0.10	0.09	0.07	0.06
Methamphetamine ³	0.25	0.21	0.08	0.06	0.06	0.04
Sedatives	0.21	0.19	0.05	0.04	0.04	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.49	0.43	0.26	0.21	0.18	0.15

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.9P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLICIT DRUGS¹	0.1268		0.0160		0.0019	
Marijuana and Hashish	0.1730		0.0273		0.0004	
Cocaine	0.1234		0.7044		0.6614	
Crack	0.1013		0.3261		0.8209	
Heroin	0.2924		0.0498		0.0514	
Hallucinogens	0.9436		0.7006		0.2366	
LSD	0.7371		0.1157		0.3573	
PCP	0.6732		0.6365		0.5521	
Ecstasy	0.4076		0.3473		0.3199	
Inhalants	0.6564		0.7158		0.5786	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0877		1.0000		0.9761	
Pain Relievers	0.0367		0.2940		0.7867	
OxyContin [®]	0.3381		0.0495		0.1608	
Tranquilizers	0.0671		0.9702		0.2631	
Stimulants ³	0.2243		0.2578		0.5002	
Methamphetamine ³	0.1812		0.7671		0.1436	
Sedatives	0.3071		0.2184		0.3446	
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.1418		0.8839		0.6762	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.10C Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	698	600	433	355	343	268
Marijuana and Hashish	685	583	357	296	294	229
Cocaine	439	372	109	105	84	60
Crack	238	171	51	53	43	40
Heroin	152	101	37	39	27	30
Hallucinogens	438	355	117	76	57	47
LSD	348	286	47	39	27	19
PCP	214	153	15	11	15	*
Ecstasy	286	205	96	65	40	41
Inhalants	289	227	44	48	29	30
Nonmedical Use of Psychotherapeutics ^{2,3}	556	445	284	212	179	149
Pain Relievers	434	346	241	173	147	115
OxyContin [®]	177	119	53	49	31	26
Tranquilizers	409	303	159	123	86	77
Stimulants ³	349	286	126	98	74	75
Methamphetamine ³	251	212	55	62	29	49
Sedatives	247	198	60	46	48	31
ILLCIT DRUGS OTHER THAN MARIJUANA¹	630	513	303	237	195	168

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.10D Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.52	0.44	0.32	0.26	0.25	0.20
Marijuana and Hashish	0.51	0.43	0.26	0.22	0.22	0.17
Cocaine	0.32	0.27	0.08	0.08	0.06	0.04
Crack	0.18	0.12	0.04	0.04	0.03	0.03
Heroin	0.11	0.07	0.03	0.03	0.02	0.02
Hallucinogens	0.32	0.26	0.09	0.06	0.04	0.03
LSD	0.26	0.21	0.03	0.03	0.02	0.01
PCP	0.16	0.11	0.01	0.01	0.01	*
Ecstasy	0.21	0.15	0.07	0.05	0.03	0.03
Inhalants	0.21	0.17	0.03	0.03	0.02	0.02
Nonmedical Use of Psychotherapeutics ^{2,3}	0.41	0.33	0.21	0.16	0.13	0.11
Pain Relievers	0.32	0.25	0.18	0.13	0.11	0.08
OxyContin [®]	0.13	0.09	0.04	0.04	0.02	0.02
Tranquilizers	0.30	0.22	0.12	0.09	0.06	0.06
Stimulants ³	0.26	0.21	0.09	0.07	0.05	0.06
Methamphetamine ³	0.19	0.16	0.04	0.05	0.02	0.04
Sedatives	0.18	0.14	0.04	0.03	0.04	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.47	0.38	0.22	0.17	0.14	0.12

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.10P Types of Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
ILLCIT DRUGS¹	0.8633		0.2969		0.2071	
Marijuana and Hashish	0.9550		0.2410		0.1321	
Cocaine	0.8595		0.1061		0.8269	
Crack	0.6339		0.4994		0.8301	
Heroin	0.0508		0.2926		0.4015	
Hallucinogens	0.4259		0.3068		0.8881	
LSD	0.6743		0.5490		0.5761	
PCP	0.2748		0.6736		*	
Ecstasy	0.0206		0.2587		0.5215	
Inhalants	0.5238		0.7355		0.9927	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.4551		0.1298		0.9266	
Pain Relievers	0.0681		0.0649		0.3135	
OxyContin [®]	0.1910		0.7723		0.5852	
Tranquilizers	0.6864		0.5236		0.8331	
Stimulants ³	0.9345		0.9866		0.2002	
Methamphetamine ³	0.9713		0.0933		0.0662	
Sedatives	0.7483		0.7212		0.8000	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.9971		0.4511		0.7256	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.11C Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,015	873	677	595	509	476
12	25	28	20	22	13	16
13	31	36	24	30	14	19
14	46	38	37	33	26	23
15	45	52	39	47	26	32
16	54	59	46	52	35	41
17	57	64	49	55	34	43
18	86	100	78	88	56	70
19	85	89	72	76	55	59
20	96	101	77	85	61	65
21	93	103	74	85	57	63
22	92	109	68	81	55	65
23	97	96	72	69	52	58
24	91	97	67	67	51	49
25	89	100	67	70	46	56
26-29	369	259	238	174	185	135
30-34	404	292	257	172	206	140
35-39	333	283	177	155	137	119
40-44	379	274	184	144	130	117
45-49	359	291	168	128	122	108
50-54	617	459	296	203	225	157
55-59	511	418	189	212	142	170
60-64	447	374	138	151	93	115
65 or Older	443	398	201	154	131	121

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.11D Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.33	0.26	0.22	0.19	0.18
12	0.65	0.70	0.53	0.57	0.35	0.41
13	0.68	0.82	0.54	0.69	0.34	0.46
14	0.92	0.84	0.79	0.74	0.58	0.53
15	0.93	1.04	0.86	0.96	0.61	0.69
16	1.01	1.06	0.93	0.99	0.74	0.84
17	1.06	1.16	1.01	1.10	0.77	0.95
18	1.25	1.49	1.19	1.45	1.00	1.23
19	1.31	1.46	1.26	1.41	1.14	1.25
20	1.33	1.50	1.31	1.47	1.17	1.23
21	1.23	1.35	1.25	1.31	1.12	1.17
22	1.25	1.41	1.22	1.38	1.07	1.21
23	1.23	1.33	1.27	1.26	1.07	1.13
24	1.21	1.32	1.21	1.23	1.03	1.01
25	1.29	1.39	1.27	1.26	1.03	1.13
26-29	1.32	0.94	1.14	0.84	0.97	0.69
30-34	1.21	0.83	1.05	0.72	0.91	0.62
35-39	1.19	0.98	0.87	0.72	0.68	0.57
40-44	1.16	0.96	0.80	0.65	0.58	0.55
45-49	1.18	0.92	0.79	0.58	0.59	0.51
50-54	1.49	1.19	1.17	0.78	0.92	0.64
55-59	1.61	1.32	0.89	0.97	0.68	0.79
60-64	1.79	1.49	0.75	0.82	0.52	0.64
65 or Older	1.02	0.88	0.46	0.34	0.30	0.27

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.11P Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2511		0.0189		0.0022	
12	0.6788		0.7946		0.6224	
13	0.3017		0.0147		0.0255	
14	0.3358		0.8419		0.2803	
15	0.6765		0.5685		0.3671	
16	0.4408		0.5514		0.7645	
17	0.7531		0.3229		0.1274	
18	0.1938		0.1922		0.9621	
19	0.1306		0.2365		0.6853	
20	0.9376		0.4442		0.8668	
21	0.8330		0.5624		0.6460	
22	0.0327		0.0587		0.2739	
23	0.8333		0.8309		0.0308	
24	0.4974		0.3076		0.0302	
25	0.6379		0.5893		0.2076	
26-29	0.6846		0.9923		0.4257	
30-34	0.7767		0.5305		0.3622	
35-39	0.0783		0.0975		0.1914	
40-44	0.5753		0.2423		0.0165	
45-49	0.6682		0.8779		0.1481	
50-54	0.2349		0.8221		0.9402	
55-59	0.4773		0.0002		0.0019	
60-64	0.0858		0.0182		0.0670	
65 or Older	0.1381		0.7318		0.3332	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.12C Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,017	872	577	538	442	424
12	12	11	12	11	7	8
13	18	21	17	20	10	13
14	32	29	29	27	20	18
15	38	42	34	39	24	26
16	49	55	43	49	33	38
17	51	61	46	52	33	41
18	81	92	72	84	52	66
19	80	86	68	72	53	56
20	91	96	73	82	56	63
21	90	96	69	79	54	57
22	86	105	63	76	52	61
23	88	92	67	65	49	53
24	87	93	63	61	48	45
25	87	96	64	66	43	55
26-29	353	249	217	163	175	128
30-34	383	280	227	158	181	123
35-39	313	264	139	134	116	107
40-44	347	259	149	120	112	102
45-49	339	278	133	114	102	97
50-54	604	451	272	168	202	130
55-59	494	407	162	193	122	160
60-64	424	363	111	127	77	104
65 or Older	425	376	146	117	110	96

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.12D Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.33	0.22	0.20	0.17	0.16
12	0.33	0.28	0.31	0.27	0.20	0.21
13	0.42	0.50	0.39	0.48	0.23	0.32
14	0.68	0.67	0.65	0.61	0.45	0.43
15	0.82	0.89	0.76	0.83	0.55	0.59
16	0.96	1.03	0.88	0.94	0.71	0.80
17	1.03	1.16	0.95	1.07	0.74	0.91
18	1.21	1.48	1.14	1.44	0.95	1.19
19	1.31	1.47	1.25	1.38	1.11	1.21
20	1.35	1.51	1.28	1.46	1.11	1.21
21	1.29	1.34	1.22	1.30	1.08	1.09
22	1.25	1.44	1.17	1.33	1.02	1.15
23	1.23	1.35	1.24	1.22	1.01	1.06
24	1.24	1.35	1.19	1.20	0.99	0.95
25	1.31	1.41	1.24	1.24	0.98	1.12
26-29	1.33	0.95	1.07	0.79	0.91	0.67
30-34	1.20	0.87	0.96	0.68	0.83	0.56
35-39	1.21	0.97	0.70	0.64	0.59	0.52
40-44	1.18	0.97	0.66	0.56	0.50	0.48
45-49	1.17	0.92	0.64	0.52	0.50	0.46
50-54	1.54	1.25	1.09	0.67	0.84	0.54
55-59	1.68	1.31	0.77	0.89	0.59	0.74
60-64	1.74	1.46	0.61	0.71	0.43	0.58
65 or Older	0.98	0.84	0.34	0.26	0.25	0.21

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.12P Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3327		0.0214		0.0004	
12	0.1789		0.3599		0.8879	
13	0.9459		0.7821		0.4959	
14	0.3960		0.4926		0.2394	
15	0.7682		0.9791		0.8715	
16	0.9933		0.5736		0.9117	
17	0.6375		0.3999		0.1317	
18	0.0926		0.2766		0.9928	
19	0.1582		0.2075		0.6852	
20	0.9684		0.9384		0.5824	
21	0.8880		0.7674		0.4446	
22	0.1106		0.0984		0.4731	
23	0.9679		0.5921		0.0316	
24	0.3208		0.1527		0.1506	
25	0.4033		0.7994		0.3993	
26-29	0.3122		0.8316		0.5802	
30-34	0.5796		0.3694		0.7576	
35-39	0.1086		0.0132		0.0257	
40-44	0.7671		0.3517		0.0180	
45-49	0.5920		0.0903		0.0450	
50-54	0.0671		0.3717		0.8059	
55-59	0.3989		0.0002		0.0010	
60-64	0.0972		0.0415		0.0807	
65 or Older	0.0769		0.2364		0.2140	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.13C Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	693	632	189	186	133	105
12	2	5	2	*	*	*
13	3	2	2	1	1	*
14	5	4	4	4	4	3
15	6	9	5	8	3	5
16	11	15	8	14	4	6
17	14	15	10	14	6	6
18	21	24	17	18	8	8
19	30	30	23	21	10	11
20	37	33	26	25	12	16
21	36	44	26	31	12	17
22	47	49	31	35	15	17
23	48	47	29	33	11	15
24	48	48	27	23	13	11
25	43	54	24	28	14	17
26-29	184	145	81	61	44	35
30-34	229	159	95	60	73	33
35-39	162	153	61	56	41	38
40-44	199	154	43	46	24	28
45-49	220	188	47	45	32	28
50-54	436	299	65	74	53	56
55-59	271	271	36	50	22	17
60-64	229	216	43	30	40	14
65 or Older	216	197	21	47	21	2

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.13D Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.26	0.24	0.07	0.07	0.05	0.04
12	0.06	0.14	0.06	*	*	*
13	0.06	0.05	0.05	0.03	0.03	*
14	0.11	0.10	0.10	0.10	0.10	0.06
15	0.14	0.22	0.12	0.19	0.06	0.12
16	0.26	0.34	0.19	0.32	0.10	0.14
17	0.34	0.36	0.25	0.32	0.15	0.14
18	0.44	0.52	0.36	0.38	0.17	0.16
19	0.71	0.72	0.54	0.49	0.24	0.27
20	0.79	0.74	0.57	0.56	0.26	0.37
21	0.79	0.91	0.59	0.68	0.28	0.37
22	0.96	1.00	0.67	0.73	0.34	0.37
23	0.99	0.97	0.64	0.71	0.24	0.33
24	0.98	1.01	0.61	0.52	0.29	0.25
25	0.99	1.10	0.57	0.63	0.34	0.40
26-29	0.91	0.75	0.46	0.35	0.25	0.20
30-34	1.01	0.68	0.46	0.28	0.36	0.16
35-39	0.82	0.71	0.32	0.28	0.21	0.19
40-44	0.84	0.69	0.20	0.22	0.11	0.14
45-49	0.94	0.79	0.23	0.22	0.16	0.13
50-54	1.56	1.08	0.28	0.31	0.23	0.23
55-59	1.21	1.16	0.18	0.24	0.11	0.08
60-64	1.21	1.10	0.24	0.17	0.23	0.08
65 or Older	0.50	0.44	0.05	0.10	0.05	0.00

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.13P Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2134		0.2191		0.8370	
12	0.5112		*		*	
13	0.3948		0.2371		*	
14	0.9441		0.8084		0.3722	
15	0.7326		0.7425		0.5075	
16	0.6252		0.0819		0.6501	
17	0.6360		0.6201		0.5370	
18	0.5228		0.7690		0.4508	
19	0.3784		0.2119		0.9085	
20	0.1675		0.6531		0.0854	
21	0.1999		0.5444		0.1234	
22	0.2432		0.5133		0.7090	
23	0.4427		0.4434		0.1741	
24	0.1551		0.0757		0.2870	
25	0.2642		0.1152		0.3920	
26-29	0.2454		0.8353		0.6131	
30-34	0.5526		0.1257		0.0818	
35-39	0.0202		0.4215		0.6123	
40-44	0.9178		0.0154		0.2393	
45-49	0.6256		0.5518		0.7657	
50-54	0.4233		0.6162		0.9283	
55-59	0.0100		0.1998		0.7584	
60-64	0.1556		0.4546		0.1909	
65 or Older	0.1522		0.3048		0.3504	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.14C Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	364	317	77	81	64	55
12	*	*	*	*	*	*
13	1	1	1	*	1	*
14	1	1	*	1	*	*
15	3	5	2	3	*	1
16	3	3	2	1	*	1
17	3	6	2	6	1	4
18	9	6	3	4	1	1
19	8	11	5	8	2	3
20	11	10	3	8	*	1
21	11	15	4	8	3	7
22	10	16	4	8	3	2
23	18	19	5	8	1	0
24	18	17	3	6	2	3
25	19	18	12	8	11	4
26-29	93	60	17	24	10	20
30-34	102	77	30	15	25	10
35-39	90	86	19	28	16	20
40-44	94	89	17	24	16	15
45-49	113	95	14	29	11	20
50-54	223	154	43	48	40	35
55-59	163	127	31	32	14	14
60-64	70	94	20	3	20	3
65 or Older	88	73	21	*	21	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.14D Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.14	0.12	0.03	0.03	0.02	0.02
12	*	*	*	*	*	*
13	0.03	0.03	0.03	*	0.03	*
14	0.02	0.01	*	0.01	*	*
15	0.07	0.12	0.05	0.07	*	0.02
16	0.07	0.07	0.05	0.03	*	0.03
17	0.08	0.14	0.04	0.13	0.02	0.10
18	0.19	0.12	0.07	0.08	0.03	0.02
19	0.19	0.26	0.11	0.20	0.04	0.06
20	0.25	0.22	0.07	0.18	*	0.02
21	0.25	0.35	0.09	0.18	0.07	0.16
22	0.23	0.35	0.09	0.18	0.07	0.05
23	0.41	0.42	0.12	0.18	0.02	0.01
24	0.41	0.38	0.08	0.14	0.05	0.08
25	0.47	0.43	0.29	0.20	0.26	0.08
26-29	0.52	0.35	0.10	0.14	0.06	0.12
30-34	0.49	0.36	0.15	0.07	0.13	0.05
35-39	0.48	0.43	0.10	0.14	0.08	0.10
40-44	0.43	0.43	0.08	0.12	0.08	0.08
45-49	0.55	0.45	0.07	0.14	0.05	0.10
50-54	0.93	0.62	0.18	0.20	0.17	0.15
55-59	0.76	0.59	0.15	0.15	0.07	0.07
60-64	0.39	0.53	0.11	0.02	0.11	0.02
65 or Older	0.20	0.16	0.05	*	0.05	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.14P Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3420		0.2313		0.7509	
12	*		*		*	
13	0.7863		*		*	
14	0.8596		*		*	
15	0.5571		0.7429		*	
16	0.6548		0.5348		*	
17	0.2989		0.0928		0.2187	
18	0.1158		0.9863		0.5596	
19	0.6386		0.8730		0.9183	
20	0.8474		0.0833		*	
21	0.4753		0.4107		0.4526	
22	0.1200		0.4360		0.4976	
23	0.9382		0.4941		0.5863	
24	0.2647		0.3752		0.7756	
25	0.4094		0.4228		0.2974	
26-29	0.1462		0.2104		0.1430	
30-34	0.6507		0.1050		0.0798	
35-39	0.1068		0.3314		0.5817	
40-44	0.1011		0.1741		0.7915	
45-49	0.4981		0.0217		0.0490	
50-54	0.3786		0.8424		0.6921	
55-59	0.7008		0.9284		0.7802	
60-64	0.0693		0.1853		0.1853	
65 or Older	0.9701		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.15C Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	707	636	204	159	132	88
12	7	7	6	4	2	3
13	8	7	5	6	3	2
14	10	10	8	7	6	4
15	12	17	9	15	5	7
16	19	21	15	19	8	12
17	23	22	19	19	12	12
18	37	41	30	33	22	13
19	42	39	33	26	19	11
20	46	44	32	35	14	17
21	52	50	38	35	14	18
22	55	59	35	36	16	18
23	58	52	33	29	14	16
24	55	54	29	27	17	14
25	51	61	25	33	14	13
26-29	215	158	72	70	41	40
30-34	256	178	84	56	35	30
35-39	198	174	58	46	50	20
40-44	215	157	50	38	15	20
45-49	189	171	26	23	7	21
50-54	368	270	43	31	42	30
55-59	269	244	8	8	*	7
60-64	208	209	13	10	*	*
65 or Older	228	180	87	12	76	8

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.15D Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.27	0.24	0.08	0.06	0.05	0.03
12	0.20	0.19	0.16	0.12	0.06	0.09
13	0.18	0.16	0.12	0.14	0.08	0.05
14	0.23	0.24	0.19	0.17	0.14	0.10
15	0.28	0.38	0.22	0.35	0.13	0.17
16	0.43	0.48	0.35	0.43	0.19	0.27
17	0.54	0.50	0.46	0.44	0.29	0.29
18	0.75	0.85	0.62	0.69	0.44	0.28
19	0.90	0.92	0.75	0.62	0.45	0.27
20	0.95	0.93	0.70	0.76	0.32	0.38
21	1.05	1.04	0.82	0.77	0.32	0.41
22	1.10	1.09	0.75	0.75	0.35	0.39
23	1.13	1.03	0.70	0.64	0.32	0.35
24	1.08	1.09	0.65	0.60	0.39	0.33
25	1.08	1.26	0.58	0.73	0.35	0.31
26-29	1.05	0.80	0.41	0.40	0.23	0.23
30-34	1.06	0.74	0.41	0.26	0.18	0.14
35-39	0.96	0.79	0.30	0.24	0.26	0.10
40-44	0.89	0.70	0.23	0.19	0.07	0.10
45-49	0.83	0.75	0.13	0.11	0.04	0.10
50-54	1.37	1.03	0.19	0.13	0.18	0.13
55-59	1.23	1.09	0.04	0.04	*	0.03
60-64	1.08	1.08	0.07	0.06	*	*
65 or Older	0.52	0.40	0.20	0.03	0.18	0.02

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.15P Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5990		0.3865		0.2769	
12	0.9570		0.4100		0.6015	
13	0.3208		0.6887		0.3657	
14	0.5191		0.2835		0.3172	
15	0.4727		0.1524		0.9944	
16	0.7461		0.9560		0.2761	
17	0.0056		0.0105		0.1336	
18	0.4813		0.6124		0.0040	
19	0.1671		0.0470		0.0511	
20	0.1994		0.7548		0.7871	
21	0.4635		0.4417		0.6714	
22	0.9910		0.6294		0.9625	
23	0.1153		0.2126		0.9360	
24	0.0289		0.1156		0.2522	
25	0.1229		0.0730		0.8535	
26-29	0.0835		0.2533		0.4093	
30-34	0.3689		0.8942		0.6047	
35-39	0.4566		0.7455		0.1915	
40-44	0.5764		0.6286		0.0830	
45-49	0.2974		0.7789		0.1139	
50-54	0.2150		0.6487		0.5819	
55-59	0.5253		0.7872		*	
60-64	0.2587		0.9924		*	
65 or Older	0.2701		0.1185		0.2427	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.16C Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	519	428	105	95	64	65
12	16	19	11	12	7	9
13	19	22	11	15	6	6
14	28	22	14	15	7	9
15	24	28	14	19	8	11
16	20	23	11	13	4	4
17	20	22	13	10	4	2
18	28	29	16	17	7	9
19	27	24	13	11	7	1
20	32	28	17	15	10	7
21	32	33	19	10	5	2
22	37	35	11	11	7	4
23	33	37	12	15	4	9
24	34	31	12	11	4	3
25	25	35	11	12	7	7
26-29	136	103	30	26	13	15
30-34	185	124	35	42	28	31
35-39	162	123	23	29	18	5
40-44	160	122	27	37	23	32
45-49	148	124	30	17	29	15
50-54	293	207	27	24	24	20
55-59	160	171	53	33	14	19
60-64	142	134	7	22	7	1
65 or Older	145	120	3	13	3	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.16D Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.20	0.16	0.04	0.04	0.02	0.02
12	0.43	0.48	0.30	0.30	0.19	0.23
13	0.42	0.53	0.27	0.36	0.14	0.15
14	0.62	0.52	0.30	0.36	0.15	0.21
15	0.56	0.63	0.35	0.42	0.19	0.25
16	0.46	0.53	0.26	0.31	0.09	0.09
17	0.47	0.51	0.31	0.24	0.10	0.05
18	0.58	0.63	0.33	0.36	0.14	0.19
19	0.62	0.58	0.32	0.27	0.17	0.03
20	0.70	0.63	0.39	0.34	0.23	0.15
21	0.73	0.73	0.44	0.23	0.12	0.05
22	0.77	0.74	0.25	0.24	0.15	0.08
23	0.71	0.79	0.28	0.35	0.09	0.19
24	0.73	0.69	0.26	0.26	0.08	0.06
25	0.60	0.79	0.27	0.28	0.16	0.17
26-29	0.73	0.55	0.18	0.15	0.08	0.09
30-34	0.82	0.55	0.17	0.20	0.14	0.15
35-39	0.80	0.58	0.12	0.15	0.09	0.03
40-44	0.71	0.57	0.12	0.18	0.11	0.16
45-49	0.69	0.56	0.15	0.08	0.14	0.07
50-54	1.17	0.81	0.12	0.10	0.10	0.09
55-59	0.75	0.78	0.26	0.16	0.07	0.09
60-64	0.77	0.73	0.04	0.12	0.04	0.01
65 or Older	0.33	0.27	0.01	0.03	0.01	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.16P Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9936		0.6320		0.6223	
12	0.9293		0.3732		0.7152	
13	0.6691		0.0725		0.5754	
14	0.6881		0.0915		0.0422	
15	0.9013		0.7009		0.5123	
16	0.7422		0.6656		0.6410	
17	0.2913		0.0291		0.0440	
18	0.4085		0.9772		0.6492	
19	0.1230		0.1464		0.0812	
20	0.1834		0.1589		0.4536	
21	0.7870		0.0394		0.0415	
22	0.4772		0.3561		0.1958	
23	0.9574		0.5429		0.2745	
24	0.0663		0.5961		0.7225	
25	0.0348		0.4911		0.6838	
26-29	0.2376		0.7100		0.2397	
30-34	0.0819		0.1056		0.5428	
35-39	0.1843		0.5076		0.5794	
40-44	0.9313		0.2616		0.4107	
45-49	0.5164		0.3495		0.4275	
50-54	0.2867		0.9781		0.9203	
55-59	0.0149		0.6890		0.7803	
60-64	0.1844		0.4785		0.4331	
65 or Older	0.2461		0.4661		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.17C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	785	658	418	321	256	222
12	16	21	13	16	10	11
13	21	25	14	20	10	15
14	29	23	24	19	16	14
15	26	32	21	28	12	19
16	32	36	25	31	17	20
17	35	39	28	32	15	19
18	52	62	39	43	23	29
19	53	49	42	36	22	18
20	59	61	44	41	28	23
21	60	66	42	48	25	29
22	64	73	45	50	26	34
23	69	67	42	41	26	27
24	62	65	39	39	26	21
25	57	67	36	39	20	25
26-29	246	171	147	100	95	78
30-34	289	190	166	103	121	69
35-39	229	191	114	95	72	59
40-44	237	175	111	85	81	60
45-49	204	173	100	68	61	52
50-54	398	285	149	117	96	83
55-59	283	256	111	98	69	71
60-64	231	219	76	90	36	55
65 or Older	329	245	146	92	83	72

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.17D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.30	0.25	0.16	0.12	0.10	0.08
12	0.42	0.53	0.34	0.42	0.26	0.30
13	0.49	0.58	0.32	0.48	0.23	0.37
14	0.62	0.54	0.52	0.45	0.36	0.33
15	0.60	0.69	0.49	0.63	0.28	0.43
16	0.70	0.74	0.56	0.66	0.39	0.45
17	0.76	0.85	0.65	0.71	0.36	0.45
18	0.99	1.19	0.78	0.85	0.48	0.59
19	1.05	1.10	0.90	0.84	0.51	0.43
20	1.11	1.22	0.91	0.87	0.62	0.51
21	1.13	1.21	0.89	0.97	0.58	0.64
22	1.22	1.29	0.93	1.00	0.56	0.71
23	1.27	1.25	0.86	0.89	0.58	0.59
24	1.16	1.22	0.83	0.84	0.58	0.47
25	1.20	1.24	0.79	0.82	0.48	0.58
26-29	1.16	0.84	0.81	0.57	0.54	0.44
30-34	1.15	0.73	0.75	0.46	0.57	0.32
35-39	1.05	0.85	0.59	0.46	0.37	0.29
40-44	0.95	0.75	0.51	0.41	0.38	0.29
45-49	0.88	0.75	0.49	0.32	0.30	0.25
50-54	1.43	1.03	0.62	0.48	0.41	0.34
55-59	1.26	1.11	0.54	0.47	0.33	0.34
60-64	1.22	1.13	0.43	0.51	0.20	0.31
65 or Older	0.76	0.54	0.33	0.20	0.19	0.16

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.17P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5162		0.3072		0.9662	
12	0.1809		0.2436		0.8242	
13	0.0452		0.0016		0.0217	
14	0.0880		0.1118		0.6393	
15	0.2157		0.1223		0.0478	
16	0.7083		0.9735		0.5114	
17	0.6209		0.9717		0.7041	
18	0.7779		0.9446		0.7638	
19	0.4687		0.3515		0.0645	
20	0.0603		0.0393		0.0978	
21	0.7951		0.7905		0.9184	
22	0.5932		0.2417		0.1923	
23	0.7545		0.6772		0.3879	
24	0.4859		0.0826		0.0240	
25	0.2919		0.2466		0.2692	
26-29	0.2126		0.2010		0.5086	
30-34	0.4893		0.0631		0.1027	
35-39	0.5614		0.7466		0.7774	
40-44	1.0000		0.9855		0.4030	
45-49	0.9236		0.0675		0.8213	
50-54	0.4355		0.8566		0.9377	
55-59	0.0149		0.2667		0.2947	
60-64	0.0457		0.0972		0.1024	
65 or Older	0.8669		0.4233		0.8293	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.18C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	640	553	343	253	206	174
12	14	17	11	13	8	9
13	20	21	13	18	9	13
14	28	21	23	17	16	12
15	24	29	19	25	11	17
16	29	31	23	26	15	15
17	32	33	24	27	12	15
18	46	55	34	37	18	24
19	48	44	36	32	17	14
20	54	52	38	33	26	16
21	53	59	34	38	20	23
22	57	61	39	41	21	29
23	56	60	35	32	24	20
24	53	56	33	29	22	18
25	49	60	32	30	17	22
26-29	206	152	109	79	69	54
30-34	256	165	138	80	104	54
35-39	195	161	96	77	60	49
40-44	201	149	99	71	68	44
45-49	165	137	80	51	53	41
50-54	269	222	134	108	75	76
55-59	218	211	95	84	43	59
60-64	168	150	60	66	31	38
65 or Older	208	161	111	77	80	57

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.18D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.24	0.21	0.13	0.10	0.08	0.07
12	0.35	0.45	0.28	0.34	0.21	0.23
13	0.46	0.50	0.30	0.42	0.22	0.31
14	0.60	0.50	0.50	0.40	0.35	0.29
15	0.56	0.65	0.45	0.56	0.25	0.39
16	0.64	0.66	0.53	0.56	0.34	0.35
17	0.71	0.73	0.57	0.61	0.29	0.36
18	0.89	1.07	0.68	0.74	0.38	0.51
19	1.00	1.00	0.79	0.75	0.41	0.33
20	1.07	1.10	0.81	0.72	0.57	0.36
21	1.08	1.12	0.76	0.79	0.46	0.51
22	1.13	1.14	0.82	0.84	0.46	0.62
23	1.12	1.17	0.74	0.70	0.53	0.45
24	1.05	1.14	0.72	0.64	0.49	0.40
25	1.05	1.17	0.71	0.67	0.41	0.50
26-29	1.02	0.78	0.62	0.45	0.40	0.31
30-34	1.05	0.68	0.64	0.37	0.50	0.26
35-39	0.92	0.75	0.49	0.38	0.32	0.25
40-44	0.84	0.68	0.45	0.35	0.32	0.22
45-49	0.76	0.63	0.39	0.24	0.26	0.20
50-54	1.09	0.85	0.56	0.44	0.32	0.32
55-59	1.00	0.94	0.46	0.40	0.21	0.28
60-64	0.91	0.82	0.34	0.37	0.17	0.22
65 or Older	0.48	0.36	0.25	0.17	0.19	0.13

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.18P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.7945		0.0451		0.3709	
12	0.4006		0.4390		0.8247	
13	0.2036		0.0232		0.1060	
14	0.0756		0.0497		0.2532	
15	0.2902		0.1979		0.0476	
16	0.3629		0.3199		0.9796	
17	0.9143		0.7391		0.8706	
18	0.6916		0.6126		0.6309	
19	0.1765		0.3586		0.2387	
20	0.0351		0.0123		0.0048	
21	0.7542		0.8242		0.9063	
22	0.5099		0.8577		0.3752	
23	0.7982		0.3986		0.0378	
24	0.4593		0.0539		0.0609	
25	0.1502		0.7574		0.5675	
26-29	0.3440		0.2052		0.8501	
30-34	0.1587		0.0895		0.1571	
35-39	0.7206		0.9350		0.9545	
40-44	0.7773		0.6484		0.2972	
45-49	0.8750		0.0248		0.9512	
50-54	0.5475		0.8334		0.3939	
55-59	0.0013		0.3201		0.1387	
60-64	0.7213		0.1917		0.3640	
65 or Older	0.7914		0.4771		0.2327	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.19C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,015	873	677	595	509	476
AGE						
12-17	97	108	86	95	62	73
18-25	168	190	166	185	143	160
26 or Older	978	818	622	515	472	415
GENDER						
Male	702	569	472	428	363	360
Female	698	600	433	355	343	268
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	938	800	623	552	471	443
White	823	686	554	481	419	381
Black or African American	326	275	219	189	175	160
American Indian or Alaska Native	93	100	54	47	30	30
Native Hawaiian or Other Pacific Islander	65	76	43	39	38	35
Asian	240	187	92	91	58	58
Two or More Races	180	128	122	71	104	51
Hispanic or Latino	395	333	243	210	197	158

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.19D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.33	0.26	0.22	0.19	0.18
AGE						
12-17	0.39	0.43	0.35	0.38	0.25	0.30
18-25	0.48	0.54	0.48	0.53	0.41	0.46
26 or Older	0.48	0.40	0.31	0.25	0.23	0.20
GENDER						
Male	0.55	0.44	0.37	0.33	0.29	0.28
Female	0.52	0.44	0.32	0.26	0.25	0.20
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.42	0.36	0.28	0.25	0.21	0.20
White	0.48	0.40	0.33	0.28	0.25	0.22
Black or African American	1.04	0.87	0.70	0.60	0.56	0.50
American Indian or Alaska Native	3.76	3.25	3.15	2.50	1.92	1.93
Native Hawaiian or Other Pacific Islander	5.46	4.73	4.33	3.67	3.96	3.33
Asian	1.59	1.26	0.70	0.68	0.44	0.42
Two or More Races	2.64	1.88	2.37	1.47	2.14	1.14
Hispanic or Latino	0.96	0.78	0.59	0.49	0.48	0.37

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.19P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2511		0.0189		0.0022	
AGE						
12-17	0.9952		0.6826		0.1397	
18-25	0.2296		0.7006		0.4747	
26 or Older	0.3587		0.0120		0.0029	
GENDER						
Male	0.1268		0.0160		0.0019	
Female	0.8633		0.2969		0.2071	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1678		0.0205		0.0012	
White	0.1620		0.1025		0.0116	
Black or African American	0.8465		0.0037		0.0113	
American Indian or Alaska Native	0.9925		0.4878		0.3394	
Native Hawaiian or Other Pacific Islander	0.3977		0.9309		0.7621	
Asian	0.4346		0.3569		0.0997	
Two or More Races	0.9168		0.1390		0.3202	
Hispanic or Latino	0.8870		0.5410		0.8699	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.20C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	97	108	86	95	62	73
GENDER						
Male	72	76	65	66	46	52
Female	68	71	60	65	41	50
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	84	93	73	82	54	64
White	84	93	70	80	47	59
Black or African American	47	53	38	40	28	29
American Indian or Alaska Native	13	10	11	9	*	6
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	28	22	18	18	11	10
Two or More Races	19	18	14	16	10	11
Hispanic or Latino	52	52	46	48	32	39
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	58	65	49	54	35	42
Female, White, Not Hispanic	54	61	47	54	31	38
Male, Black, Not Hispanic	33	33	27	26	21	20
Female, Black, Not Hispanic	30	38	24	29	16	20
Male, Hispanic	45	44	40	37	25	28
Female, Hispanic	39	44	31	40	22	30

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.20D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.43	0.35	0.38	0.25	0.30
GENDER						
Male	0.57	0.60	0.51	0.52	0.36	0.41
Female	0.56	0.58	0.50	0.53	0.34	0.41
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.44	0.49	0.38	0.43	0.28	0.33
White	0.49	0.58	0.43	0.51	0.32	0.41
Black or African American	1.06	1.14	0.90	0.94	0.73	0.76
American Indian or Alaska Native	5.05	4.42	4.94	4.29	*	2.78
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.91	1.63	1.37	1.40	0.91	0.84
Two or More Races	2.06	2.03	1.73	1.79	1.25	1.29
Hispanic or Latino	0.93	0.93	0.83	0.85	0.58	0.69
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.67	0.81	0.61	0.71	0.47	0.58
Female, White, Not Hispanic	0.70	0.79	0.64	0.72	0.44	0.55
Male, Black, Not Hispanic	1.58	1.52	1.36	1.29	1.11	1.03
Female, Black, Not Hispanic	1.45	1.72	1.24	1.35	0.89	1.08
Male, Hispanic	1.28	1.32	1.21	1.18	0.85	0.94
Female, Hispanic	1.24	1.30	1.08	1.24	0.78	1.00

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.20P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9952		0.6826		0.1397	
GENDER						
Male	0.1455		0.1124		0.9520	
Female	0.1172		0.0249		0.0328	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4922		0.6890		0.6314	
White	0.7025		0.9875		0.2728	
Black or African American	0.8460		0.4804		0.5249	
American Indian or Alaska Native	0.1509		0.4867		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4332		0.9848		0.8024	
Two or More Races	0.6832		0.8929		0.8596	
Hispanic or Latino	0.2436		0.1501		0.0449	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.3754		0.4319		0.4925	
Female, White, Not Hispanic	0.7147		0.4407		0.3700	
Male, Black, Not Hispanic	0.3525		0.2066		0.1559	
Female, Black, Not Hispanic	0.5083		0.7593		0.5358	
Male, Hispanic	0.5298		0.2504		0.7294	
Female, Hispanic	0.0164		0.0009		0.0113	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.21C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	168	190	166	185	143	160
GENDER						
Male	114	127	116	128	107	115
Female	110	127	105	122	87	102
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	141	166	142	164	125	144
White	247	256	194	197	138	151
Black or African American	108	115	80	87	58	71
American Indian or Alaska Native	19	18	13	13	9	9
Native Hawaiian or Other Pacific Islander	*	*	*	*	10	*
Asian	55	64	39	48	26	33
Two or More Races	38	36	31	30	22	25
Hispanic or Latino	85	92	75	83	63	66
EDUCATION						
< High School	100	103	80	81	65	64
High School Graduate	157	168	119	134	91	109
Some College	184	198	141	154	102	116
College Graduate	122	129	89	95	54	68
CURRENT EMPLOYMENT						
Full-Time	183	192	135	145	95	113
Part-Time	148	164	121	132	90	102
Unemployed	89	88	77	69	61	57
Other ¹	126	128	97	103	71	70

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.21D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.48	0.54	0.48	0.53	0.41	0.46
GENDER						
Male	0.66	0.72	0.67	0.73	0.61	0.66
Female	0.64	0.73	0.61	0.70	0.50	0.59
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.51	0.60	0.51	0.60	0.46	0.52
White	0.60	0.67	0.62	0.69	0.54	0.62
Black or African American	1.14	1.27	1.09	1.26	0.91	1.14
American Indian or Alaska Native	4.37	4.40	4.09	4.87	3.10	3.86
Native Hawaiian or Other Pacific Islander	*	*	*	*	4.41	*
Asian	2.27	2.30	1.72	1.99	1.27	1.51
Two or More Races	2.83	2.68	2.89	2.90	2.52	2.63
Hispanic or Latino	1.17	1.24	1.04	1.13	0.88	0.90
EDUCATION						
< High School	1.14	1.32	1.11	1.30	1.00	1.10
High School Graduate	0.79	0.90	0.72	0.83	0.66	0.77
Some College	0.75	0.87	0.76	0.88	0.66	0.76
College Graduate	1.25	1.32	1.21	1.26	0.94	1.02
CURRENT EMPLOYMENT						
Full-Time	0.73	0.74	0.74	0.74	0.61	0.66
Part-Time	0.85	0.97	0.87	0.99	0.78	0.88
Unemployed	1.26	1.52	1.34	1.41	1.23	1.29
Other ¹	1.09	1.12	0.97	1.06	0.81	0.80

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.21P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2296		0.7006		0.4747	
GENDER						
Male	0.8813		0.4747		0.5622	
Female	0.0960		0.1728		0.0688	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1161		0.8769		0.4052	
White	0.0969		0.5658		0.7576	
Black or African American	0.1064		0.0334		0.0328	
American Indian or Alaska Native	0.6432		0.6920		0.9707	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4321		0.5411		0.4364	
Two or More Races	0.5051		0.3897		0.6783	
Hispanic or Latino	0.7997		0.5532		0.9827	
EDUCATION						
< High School	0.7977		0.8604		0.4654	
High School Graduate	0.6884		0.6481		0.1155	
Some College	0.0796		0.2213		0.5508	
College Graduate	0.6915		0.2895		0.8558	
CURRENT EMPLOYMENT						
Full-Time	0.1337		0.1989		0.0129	
Part-Time	0.4112		0.9087		0.6841	
Unemployed	0.3018		0.0988		0.7733	
Other ¹	0.8042		0.7040		0.3568	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.22C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	978	818	622	515	472	415
GENDER						
Male	679	543	438	384	337	322
Female	673	567	407	304	326	230
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	908	758	577	483	438	388
White	1,320	1,053	574	465	414	366
Black or African American	492	390	205	194	162	161
American Indian or Alaska Native	87	93	53	41	26	27
Native Hawaiian or Other Pacific Islander	*	*	*	34	*	*
Asian	227	168	83	72	50	45
Two or More Races	173	121	118	62	101	44
Hispanic or Latino	364	309	217	177	177	137
EDUCATION						
< High School	394	360	222	203	186	152
High School Graduate	683	542	307	261	246	220
Some College	733	558	334	284	267	227
College Graduate	942	709	420	302	263	213
CURRENT EMPLOYMENT						
Full-Time	1,121	874	504	394	371	309
Part-Time	490	368	261	191	211	152
Unemployed	258	196	147	105	104	85
Other ¹	688	600	318	271	232	213

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.22D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.48	0.40	0.31	0.25	0.23	0.20
GENDER						
Male	0.70	0.55	0.45	0.39	0.35	0.33
Female	0.64	0.53	0.39	0.28	0.31	0.21
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.52	0.43	0.33	0.27	0.25	0.22
White	0.58	0.47	0.38	0.31	0.29	0.25
Black or African American	1.40	1.11	0.86	0.74	0.70	0.63
American Indian or Alaska Native	4.92	4.01	4.39	3.13	2.42	2.44
Native Hawaiian or Other Pacific Islander	*	*	*	4.34	*	*
Asian	2.02	1.54	0.82	0.71	0.50	0.44
Two or More Races	3.92	2.77	3.55	2.04	3.22	1.54
Hispanic or Latino	1.28	1.05	0.76	0.60	0.62	0.47
EDUCATION						
< High School	1.23	1.08	0.78	0.69	0.66	0.54
High School Graduate	0.88	0.69	0.50	0.43	0.41	0.37
Some College	0.92	0.70	0.61	0.49	0.50	0.40
College Graduate	0.89	0.70	0.58	0.42	0.38	0.30
CURRENT EMPLOYMENT						
Full-Time	0.63	0.51	0.44	0.32	0.33	0.27
Part-Time	1.36	1.08	0.99	0.75	0.82	0.61
Unemployed	2.30	1.83	1.81	1.33	1.33	1.13
Other ¹	0.83	0.70	0.47	0.39	0.35	0.32

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.22P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3587		0.0120		0.0029	
GENDER						
Male	0.0933		0.0017		0.0003	
Female	0.7403		0.6994		0.6467	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2412		0.0084		0.0010	
White	0.2426		0.0435		0.0061	
Black or African American	0.9151		0.0068		0.0280	
American Indian or Alaska Native	0.8765		0.5654		0.2437	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.2988		0.2027		0.1265	
Two or More Races	0.9533		0.1622		0.2426	
Hispanic or Latino	0.7936		0.8293		0.8464	
EDUCATION						
< High School	0.1103		0.1999		0.9102	
High School Graduate	0.1783		0.2311		0.1069	
Some College	0.8558		0.0088		0.0063	
College Graduate	0.4396		0.8133		0.1502	
CURRENT EMPLOYMENT						
Full-Time	0.8132		0.0020		0.0012	
Part-Time	0.1412		0.7330		0.6443	
Unemployed	0.4872		0.6236		0.6048	
Other ¹	0.0530		0.2211		0.1123	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.23C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	999	857	666	581	503	462
GENDER						
Male	691	559	461	420	356	351
Female	689	595	430	348	342	263
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	924	789	617	539	467	431
White	1,366	1,120	622	517	443	400
Black or African American	525	433	225	224	176	184
American Indian or Alaska Native	90	99	52	44	28	29
Native Hawaiian or Other Pacific Islander	63	74	42	37	*	35
Asian	237	182	91	88	56	56
Two or More Races	178	127	121	69	103	50
Hispanic or Latino	385	327	234	204	192	153
EDUCATION						
< High School	416	379	243	220	199	163
High School Graduate	707	575	328	299	259	250
Some College	765	595	368	328	286	259
College Graduate	967	729	432	320	269	223
CURRENT EMPLOYMENT						
Full-Time	1,150	909	528	430	384	336
Part-Time	517	394	285	235	228	188
Unemployed	272	219	165	128	121	102
Other ¹	702	615	331	288	244	222

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.23D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.42	0.36	0.28	0.24	0.21	0.19
GENDER						
Male	0.60	0.48	0.40	0.36	0.31	0.30
Female	0.56	0.48	0.35	0.28	0.28	0.21
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.46	0.39	0.31	0.26	0.23	0.21
White	0.52	0.43	0.35	0.30	0.27	0.24
Black or African American	1.17	0.96	0.76	0.66	0.61	0.56
American Indian or Alaska Native	4.11	3.49	3.51	2.77	2.08	2.14
Native Hawaiian or Other Pacific Islander	5.90	5.05	4.83	3.92	*	3.64
Asian	1.76	1.37	0.77	0.73	0.48	0.45
Two or More Races	3.18	2.28	2.88	1.77	2.59	1.36
Hispanic or Latino	1.08	0.89	0.65	0.55	0.54	0.42
EDUCATION						
< High School	1.05	0.96	0.70	0.63	0.58	0.50
High School Graduate	0.75	0.58	0.44	0.39	0.36	0.34
Some College	0.76	0.61	0.54	0.46	0.43	0.37
College Graduate	0.84	0.66	0.55	0.40	0.36	0.29
CURRENT EMPLOYMENT						
Full-Time	0.57	0.46	0.40	0.31	0.30	0.25
Part-Time	1.00	0.80	0.75	0.63	0.63	0.52
Unemployed	1.55	1.30	1.33	1.03	0.99	0.86
Other ¹	0.76	0.65	0.44	0.38	0.33	0.30

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.23P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2681		0.0193		0.0036	
GENDER						
Male	0.0962		0.0072		0.0016	
Female	0.9798		0.4811		0.3382	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1579		0.0160		0.0012	
White	0.1612		0.0953		0.0151	
Black or African American	0.8284		0.0020		0.0075	
American Indian or Alaska Native	0.8186		0.5660		0.3039	
Native Hawaiian or Other Pacific Islander	0.2996		0.7851		*	
Asian	0.3945		0.3489		0.0845	
Two or More Races	0.8453		0.1313		0.3259	
Hispanic or Latino	0.7330		0.7393		0.8107	
EDUCATION						
< High School	0.1877		0.5452		0.4502	
High School Graduate	0.1566		0.1756		0.0346	
Some College	0.5708		0.0076		0.0081	
College Graduate	0.4351		0.8402		0.1349	
CURRENT EMPLOYMENT						
Full-Time	0.6220		0.0008		0.0001	
Part-Time	0.2378		0.8037		0.5685	
Unemployed	0.7707		0.3128		0.6971	
Other ¹	0.0559		0.2453		0.2277	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.24C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,017	872	577	538	442	424
AGE						
12-17	86	95	78	82	56	67
18-25	170	197	163	186	135	157
26 or Older	986	811	519	451	407	361
GENDER						
Male	717	570	416	399	321	320
Female	685	583	357	296	294	229
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	930	805	545	494	419	397
White	823	685	479	430	368	346
Black or African American	320	273	203	168	157	136
American Indian or Alaska Native	85	89	42	33	28	27
Native Hawaiian or Other Pacific Islander	64	75	42	34	37	30
Asian	200	165	70	70	46	45
Two or More Races	167	125	110	65	103	47
Hispanic or Latino	380	314	192	185	152	138

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.24D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.33	0.22	0.20	0.17	0.16
AGE						
12-17	0.35	0.38	0.31	0.33	0.23	0.27
18-25	0.49	0.56	0.47	0.53	0.39	0.45
26 or Older	0.49	0.39	0.26	0.22	0.20	0.18
GENDER						
Male	0.56	0.44	0.33	0.31	0.25	0.25
Female	0.51	0.43	0.26	0.22	0.22	0.17
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.42	0.36	0.25	0.22	0.19	0.18
White	0.48	0.40	0.28	0.25	0.22	0.20
Black or African American	1.02	0.86	0.65	0.53	0.50	0.43
American Indian or Alaska Native	3.81	3.57	2.55	1.93	1.83	1.79
Native Hawaiian or Other Pacific Islander	5.38	5.06	4.30	3.21	3.94	2.83
Asian	1.39	1.17	0.54	0.52	0.35	0.33
Two or More Races	2.61	1.90	2.21	1.38	2.12	1.07
Hispanic or Latino	0.92	0.74	0.47	0.44	0.37	0.33

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.24P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3327		0.0214		0.0004	
AGE						
12-17	0.9930		0.4690		0.4269	
18-25	0.3159		0.6654		0.4081	
26 or Older	0.4483		0.0073		0.0004	
GENDER						
Male	0.1730		0.0273		0.0004	
Female	0.9550		0.2410		0.1321	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2005		0.0325		0.0004	
White	0.2046		0.0410		0.0011	
Black or African American	0.8487		0.0701		0.0153	
American Indian or Alaska Native	0.6539		0.6431		0.6761	
Native Hawaiian or Other Pacific Islander	0.6113		0.5947		0.7803	
Asian	0.6087		0.9373		0.2794	
Two or More Races	0.5526		0.1443		0.1312	
Hispanic or Latino	0.7302		0.3313		0.6193	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.25C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	86	95	78	82	56	67
GENDER						
Male	62	66	57	59	42	48
Female	60	62	54	56	37	43
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	72	80	65	70	48	57
White	67	78	61	68	42	53
Black or African American	37	41	32	33	22	24
American Indian or Alaska Native	10	10	8	9	5	6
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	18	16	16	15	11	8
Two or More Races	14	13	13	11	9	8
Hispanic or Latino	46	47	42	42	29	35
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	47	55	42	48	31	39
Female, White, Not Hispanic	44	49	40	45	28	33
Male, Black, Not Hispanic	27	27	23	23	17	18
Female, Black, Not Hispanic	24	29	20	22	12	15
Male, Hispanic	38	37	35	32	23	24
Female, Hispanic	32	38	28	34	19	26

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.25D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.35	0.38	0.31	0.33	0.23	0.27
GENDER						
Male	0.48	0.52	0.45	0.46	0.33	0.38
Female	0.49	0.51	0.44	0.46	0.30	0.35
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.37	0.42	0.34	0.36	0.25	0.30
White	0.43	0.51	0.40	0.45	0.29	0.37
Black or African American	0.92	1.00	0.83	0.83	0.59	0.65
American Indian or Alaska Native	4.22	4.25	3.52	4.28	2.58	2.78
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.36	1.25	1.26	1.15	0.90	0.69
Two or More Races	1.65	1.60	1.59	1.43	1.15	1.05
Hispanic or Latino	0.82	0.84	0.76	0.74	0.52	0.62
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.60	0.72	0.55	0.64	0.42	0.55
Female, White, Not Hispanic	0.60	0.66	0.56	0.61	0.40	0.48
Male, Black, Not Hispanic	1.34	1.29	1.22	1.15	0.94	0.93
Female, Black, Not Hispanic	1.25	1.48	1.12	1.13	0.70	0.84
Male, Hispanic	1.17	1.17	1.09	1.05	0.77	0.83
Female, Hispanic	1.13	1.23	0.99	1.10	0.69	0.88

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.25P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9930		0.4690		0.4269	
GENDER						
Male	0.4576		0.1657		0.9710	
Female	0.4300		0.6811		0.2033	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6438		0.2320		0.7074	
White	0.9614		0.5933		0.2157	
Black or African American	0.5897		0.3528		0.5502	
American Indian or Alaska Native	0.4929		0.9848		0.9305	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7466		0.5997		0.3208	
Two or More Races	0.6279		0.2460		0.2754	
Hispanic or Latino	0.4496		0.5914		0.3645	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.8703		0.4818		0.4485	
Female, White, Not Hispanic	0.8024		0.9689		0.3014	
Male, Black, Not Hispanic	0.6052		0.4408		0.5341	
Female, Black, Not Hispanic	0.8035		0.5795		0.8540	
Male, Hispanic	0.4432		0.2769		0.6969	
Female, Hispanic	0.0605		0.0537		0.0815	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.26C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	170	197	163	186	135	157
GENDER						
Male	115	131	115	127	103	112
Female	110	128	104	118	82	100
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	144	172	142	160	120	141
White	233	248	182	183	130	142
Black or African American	103	108	77	83	55	69
American Indian or Alaska Native	16	18	12	11	7	8
Native Hawaiian or Other Pacific Islander	*	*	*	*	10	*
Asian	47	56	34	38	20	29
Two or More Races	36	36	28	29	21	24
Hispanic or Latino	80	92	72	81	60	63
EDUCATION						
< High School	91	96	73	76	60	61
High School Graduate	147	155	110	126	85	104
Some College	177	189	131	147	96	109
College Graduate	112	123	86	87	50	62
CURRENT EMPLOYMENT						
Full-Time	174	183	127	136	88	108
Part-Time	137	156	113	126	88	96
Unemployed	85	82	73	65	59	53
Other ¹	118	120	89	98	66	66

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.26D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.49	0.56	0.47	0.53	0.39	0.45
GENDER						
Male	0.66	0.75	0.66	0.72	0.59	0.64
Female	0.63	0.74	0.60	0.68	0.48	0.57
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.52	0.62	0.51	0.58	0.43	0.51
White	0.61	0.71	0.62	0.67	0.53	0.60
Black or African American	1.16	1.26	1.08	1.26	0.89	1.12
American Indian or Alaska Native	4.32	4.42	4.03	4.49	2.64	3.49
Native Hawaiian or Other Pacific Islander	*	*	*	*	4.37	*
Asian	2.15	2.22	1.62	1.69	1.04	1.39
Two or More Races	2.96	2.72	2.75	2.84	2.45	2.52
Hispanic or Latino	1.11	1.25	1.00	1.10	0.83	0.86
EDUCATION						
< High School	1.10	1.33	1.05	1.26	0.95	1.08
High School Graduate	0.78	0.88	0.70	0.82	0.62	0.75
Some College	0.78	0.90	0.74	0.88	0.63	0.73
College Graduate	1.30	1.35	1.23	1.21	0.87	0.96
CURRENT EMPLOYMENT						
Full-Time	0.74	0.76	0.73	0.73	0.57	0.65
Part-Time	0.89	0.98	0.85	0.98	0.76	0.85
Unemployed	1.34	1.51	1.33	1.37	1.22	1.24
Other ¹	1.06	1.13	0.92	1.05	0.77	0.78

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.26P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3159		0.6654		0.4081	
GENDER						
Male	0.8272		0.5391		0.6798	
Female	0.0763		0.1769		0.0702	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1095		0.8582		0.2567	
White	0.1236		0.8012		0.7700	
Black or African American	0.1052		0.0868		0.0036	
American Indian or Alaska Native	0.1303		0.6813		0.8883	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4797		0.2755		0.4229	
Two or More Races	0.9530		0.6526		0.6267	
Hispanic or Latino	0.4514		0.4978		0.6909	
EDUCATION						
< High School	0.8608		0.9243		0.5672	
High School Graduate	0.5504		0.5157		0.0746	
Some College	0.1470		0.1875		0.5942	
College Graduate	0.6706		0.1567		0.8363	
CURRENT EMPLOYMENT						
Full-Time	0.1558		0.3553		0.0073	
Part-Time	0.5479		0.8099		0.5620	
Unemployed	0.3147		0.1418		0.5390	
Other ¹	0.8918		0.4455		0.6911	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.27C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	986	811	519	451	407	361
GENDER						
Male	699	542	387	348	297	282
Female	661	546	329	245	276	193
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	908	757	498	423	388	344
White	1,242	1,004	484	403	357	327
Black or African American	461	374	183	165	141	133
American Indian or Alaska Native	81	83	36	28	26	24
Native Hawaiian or Other Pacific Islander	*	*	*	29	*	27
Asian	195	146	61	54	40	33
Two or More Races	162	119	106	57	100	40
Hispanic or Latino	352	290	163	142	134	112
EDUCATION						
< High School	358	335	174	165	148	124
High School Graduate	647	528	260	222	214	192
Some College	699	529	278	249	225	200
College Graduate	895	667	357	263	230	191
CURRENT EMPLOYMENT						
Full-Time	1,085	847	420	339	323	265
Part-Time	474	365	227	167	193	136
Unemployed	254	189	129	88	88	70
Other ¹	625	567	260	230	182	185

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.27D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.49	0.39	0.26	0.22	0.20	0.18
GENDER						
Male	0.72	0.55	0.40	0.35	0.31	0.29
Female	0.63	0.51	0.31	0.23	0.26	0.18
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.52	0.43	0.29	0.24	0.22	0.20
White	0.59	0.47	0.32	0.28	0.25	0.23
Black or African American	1.37	1.10	0.78	0.66	0.61	0.53
American Indian or Alaska Native	5.03	4.53	3.18	2.30	2.39	2.25
Native Hawaiian or Other Pacific Islander	*	*	*	3.79	*	3.53
Asian	1.80	1.39	0.60	0.52	0.39	0.32
Two or More Races	3.89	2.79	3.28	1.91	3.17	1.40
Hispanic or Latino	1.24	0.98	0.57	0.48	0.47	0.38
EDUCATION						
< High School	1.16	1.05	0.62	0.58	0.53	0.44
High School Graduate	0.87	0.68	0.43	0.37	0.36	0.32
Some College	0.93	0.71	0.52	0.44	0.42	0.36
College Graduate	0.88	0.69	0.51	0.37	0.34	0.27
CURRENT EMPLOYMENT						
Full-Time	0.64	0.52	0.37	0.29	0.29	0.23
Part-Time	1.37	1.12	0.87	0.66	0.75	0.55
Unemployed	2.29	1.85	1.64	1.17	1.14	0.96
Other ¹	0.80	0.69	0.39	0.34	0.28	0.27

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.27P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.4483		0.0073		0.0004	
GENDER						
Male	0.1410		0.0036		0.0000	
Female	0.6942		0.4244		0.4235	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3102		0.0097		0.0004	
White	0.3214		0.0130		0.0004	
Black or African American	0.9352		0.1031		0.0877	
American Indian or Alaska Native	0.7537		0.6059		0.6579	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4925		0.4594		0.2997	
Two or More Races	0.5097		0.1938		0.1030	
Hispanic or Latino	0.7852		0.4072		0.5291	
EDUCATION						
< High School	0.2323		0.4743		0.6113	
High School Graduate	0.1938		0.1981		0.0276	
Some College	0.9904		0.0035		0.0064	
College Graduate	0.5675		0.6132		0.1259	
CURRENT EMPLOYMENT						
Full-Time	0.8505		0.0036		0.0032	
Part-Time	0.2619		0.9594		0.8136	
Unemployed	0.5520		0.4492		0.5741	
Other ¹	0.0375		0.0945		0.0021	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.28C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,008	860	565	528	437	414
GENDER						
Male	710	563	408	392	316	314
Female	679	577	354	290	291	224
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	922	796	537	485	416	388
White	1,285	1,070	527	452	386	360
Black or African American	490	416	205	190	155	152
American Indian or Alaska Native	83	88	40	31	27	26
Native Hawaiian or Other Pacific Islander	62	73	*	32	*	30
Asian	200	161	68	67	44	43
Two or More Races	166	124	110	64	102	46
Hispanic or Latino	371	309	184	179	148	135
EDUCATION						
< High School	372	354	193	180	162	135
High School Graduate	669	554	280	260	229	225
Some College	728	565	312	299	246	233
College Graduate	919	685	370	280	234	199
CURRENT EMPLOYMENT						
Full-Time	1,115	883	446	376	337	293
Part-Time	498	390	251	213	211	170
Unemployed	267	210	146	111	106	87
Other ¹	640	580	273	249	197	194

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.28D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.42	0.36	0.24	0.22	0.18	0.17
GENDER						
Male	0.62	0.49	0.36	0.34	0.28	0.27
Female	0.55	0.46	0.29	0.23	0.24	0.18
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.46	0.39	0.27	0.24	0.21	0.19
White	0.52	0.43	0.30	0.27	0.23	0.22
Black or African American	1.15	0.96	0.71	0.59	0.55	0.48
American Indian or Alaska Native	4.17	3.91	2.84	2.09	2.03	2.01
Native Hawaiian or Other Pacific Islander	5.85	5.31	*	3.45	*	3.16
Asian	1.54	1.26	0.59	0.55	0.37	0.35
Two or More Races	3.17	2.30	2.68	1.67	2.56	1.28
Hispanic or Latino	1.04	0.84	0.52	0.49	0.41	0.37
EDUCATION						
< High School	0.99	0.94	0.57	0.54	0.48	0.42
High School Graduate	0.74	0.57	0.38	0.35	0.32	0.31
Some College	0.77	0.61	0.46	0.43	0.37	0.34
College Graduate	0.83	0.65	0.49	0.36	0.32	0.26
CURRENT EMPLOYMENT						
Full-Time	0.58	0.47	0.34	0.28	0.27	0.23
Part-Time	1.01	0.83	0.67	0.59	0.58	0.48
Unemployed	1.55	1.32	1.22	0.93	0.88	0.76
Other ¹	0.73	0.63	0.37	0.33	0.27	0.26

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.28P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3598		0.0141		0.0006	
GENDER						
Male	0.1646		0.0146		0.0004	
Female	0.9571		0.2592		0.1817	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2037		0.0197		0.0004	
White	0.2220		0.0316		0.0016	
Black or African American	0.8039		0.0452		0.0106	
American Indian or Alaska Native	0.5735		0.6319		0.6816	
Native Hawaiian or Other Pacific Islander	0.5204		*		*	
Asian	0.6046		0.8394		0.1785	
Two or More Races	0.5315		0.1928		0.1627	
Hispanic or Latino	0.6289		0.3705		0.7564	
EDUCATION						
< High School	0.4014		0.9889		0.8045	
High School Graduate	0.1577		0.1322		0.0063	
Some College	0.7324		0.0034		0.0101	
College Graduate	0.5556		0.7105		0.1145	
CURRENT EMPLOYMENT						
Full-Time	0.9518		0.0020		0.0003	
Part-Time	0.3656		0.9428		0.6518	
Unemployed	0.8446		0.2401		0.8128	
Other ¹	0.0419		0.0945		0.0074	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.29C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	693	632	189	186	133	105
AGE						
12-17	19	26	15	23	9	10
18-25	108	120	72	81	34	40
26 or Older	674	611	172	165	129	98
GENDER						
Male	508	470	154	145	101	84
Female	439	372	109	105	84	60
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	652	588	178	168	126	95
White	609	515	165	139	119	82
Black or African American	212	183	64	80	51	40
American Indian or Alaska Native	34	64	6	18	1	15
Native Hawaiian or Other Pacific Islander	*	38	7	11	*	2
Asian	64	88	25	31	8	27
Two or More Races	110	86	13	17	4	9
Hispanic or Latino	266	214	66	83	44	47

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.29D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.26	0.24	0.07	0.07	0.05	0.04
AGE						
12-17	0.08	0.10	0.06	0.09	0.04	0.04
18-25	0.31	0.34	0.21	0.23	0.10	0.11
26 or Older	0.33	0.30	0.08	0.08	0.06	0.05
GENDER						
Male	0.40	0.37	0.12	0.11	0.08	0.07
Female	0.32	0.27	0.08	0.08	0.06	0.04
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.29	0.26	0.08	0.08	0.06	0.04
White	0.36	0.30	0.10	0.08	0.07	0.05
Black or African American	0.68	0.58	0.21	0.25	0.16	0.12
American Indian or Alaska Native	2.31	3.46	0.42	1.23	0.07	1.01
Native Hawaiian or Other Pacific Islander	*	3.38	0.78	1.05	*	0.19
Asian	0.48	0.63	0.19	0.22	0.06	0.19
Two or More Races	2.31	1.75	0.30	0.39	0.10	0.21
Hispanic or Latino	0.64	0.50	0.16	0.20	0.11	0.11

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.29P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2134		0.2191		0.8370	
AGE						
12-17	0.8252		0.1860		0.7663	
18-25	0.2725		0.6188		0.1629	
26 or Older	0.1591		0.3133		0.5229	
GENDER						
Male	0.1234		0.7044		0.6614	
Female	0.8595		0.1061		0.8269	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1461		0.4367		0.6436	
White	0.2592		0.8218		0.4907	
Black or African American	0.4094		0.1276		0.3637	
American Indian or Alaska Native	0.0379		0.1297		0.0966	
Native Hawaiian or Other Pacific Islander	*		0.9872		*	
Asian	0.2094		0.5320		0.0793	
Two or More Races	0.9825		0.1067		0.1156	
Hispanic or Latino	0.8133		0.2055		0.5562	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.30C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	19	26	15	23	9	10
GENDER						
Male	15	17	11	16	6	6
Female	13	19	10	15	7	8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	16	19	13	17	8	6
White	16	18	12	16	7	6
Black or African American	4	6	4	4	4	*
American Indian or Alaska Native	0	1	*	*	*	*
Native Hawaiian or Other Pacific Islander	1	*	*	*	*	*
Asian	1	*	*	*	*	*
Two or More Races	3	3	2	3	*	0
Hispanic or Latino	11	16	7	15	4	8
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	12	11	9	11	3	4
Female, White, Not Hispanic	10	13	9	11	6	5
Male, Black, Not Hispanic	4	4	4	4	4	*
Female, Black, Not Hispanic	0	5	0	*	*	*
Male, Hispanic	9	11	5	10	3	5
Female, Hispanic	6	11	5	11	3	6

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.30D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.08	0.10	0.06	0.09	0.04	0.04
GENDER						
Male	0.12	0.13	0.09	0.13	0.05	0.05
Female	0.10	0.15	0.09	0.13	0.06	0.06
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.08	0.10	0.07	0.09	0.04	0.03
White	0.11	0.13	0.09	0.12	0.05	0.05
Black or African American	0.11	0.18	0.11	0.11	0.11	*
American Indian or Alaska Native	0.18	0.65	*	*	*	*
Native Hawaiian or Other Pacific Islander	1.14	*	*	*	*	*
Asian	0.08	*	*	*	*	*
Two or More Races	0.36	0.36	0.32	0.35	*	0.02
Hispanic or Latino	0.19	0.28	0.13	0.27	0.08	0.14
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.17	0.16	0.13	0.15	0.05	0.06
Female, White, Not Hispanic	0.15	0.20	0.13	0.17	0.09	0.07
Male, Black, Not Hispanic	0.22	0.23	0.22	0.21	0.22	*
Female, Black, Not Hispanic	0.01	0.29	0.01	*	*	*
Male, Hispanic	0.31	0.37	0.19	0.36	0.10	0.18
Female, Hispanic	0.23	0.41	0.17	0.38	0.12	0.23

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.30P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.8252		0.1860		0.7663	
GENDER						
Male	0.4863		0.4983		0.6893	
Female	0.3129		0.2206		0.9565	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.9341		0.8250		0.2815	
White	0.6571		0.8845		0.4123	
Black or African American	0.1278		0.5584		*	
American Indian or Alaska Native	0.3495		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	*		*		*	
Two or More Races	0.7655		0.9096		*	
Hispanic or Latino	0.6530		0.0604		0.4579	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.4528		0.9565		0.7056	
Female, White, Not Hispanic	0.9112		0.7941		0.4615	
Male, Black, Not Hispanic	0.4261		0.5268		*	
Female, Black, Not Hispanic	0.1614		*		*	
Male, Hispanic	0.6162		0.2816		0.6837	
Female, Hispanic	0.2249		0.1017		0.5326	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.31C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	108	120	72	81	34	40
GENDER						
Male	81	91	56	62	27	29
Female	68	68	44	45	20	25
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	99	103	67	67	29	35
White	102	104	66	63	29	32
Black or African American	17	28	12	19	10	9
American Indian or Alaska Native	6	5	3	4	1	2
Native Hawaiian or Other Pacific Islander	6	*	*	*	*	*
Asian	16	16	7	15	4	11
Two or More Races	19	15	8	10	2	3
Hispanic or Latino	47	59	29	43	17	17
EDUCATION						
< High School	45	46	27	25	12	12
High School Graduate	61	69	37	43	19	21
Some College	69	77	42	53	21	28
College Graduate	50	52	35	37	13	19
CURRENT EMPLOYMENT						
Full-Time	78	86	47	54	22	26
Part-Time	56	59	39	42	20	25
Unemployed	42	40	23	26	8	11
Other ¹	46	43	30	27	13	14

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.31D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.31	0.34	0.21	0.23	0.10	0.11
GENDER						
Male	0.47	0.52	0.32	0.36	0.15	0.17
Female	0.39	0.39	0.26	0.26	0.12	0.14
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.36	0.37	0.24	0.24	0.11	0.13
White	0.46	0.48	0.32	0.31	0.14	0.16
Black or African American	0.34	0.53	0.24	0.37	0.19	0.19
American Indian or Alaska Native	2.40	2.57	1.42	1.87	0.41	0.93
Native Hawaiian or Other Pacific Islander	2.98	*	*	*	*	*
Asian	0.86	0.81	0.39	0.72	0.20	0.53
Two or More Races	2.48	1.82	1.08	1.19	0.23	0.37
Hispanic or Latino	0.65	0.81	0.41	0.59	0.23	0.23
EDUCATION						
< High School	0.72	0.88	0.47	0.50	0.22	0.24
High School Graduate	0.49	0.55	0.31	0.35	0.16	0.17
Some College	0.52	0.56	0.34	0.40	0.17	0.22
College Graduate	0.90	0.83	0.66	0.61	0.25	0.34
CURRENT EMPLOYMENT						
Full-Time	0.50	0.56	0.33	0.36	0.17	0.18
Part-Time	0.55	0.57	0.39	0.42	0.21	0.26
Unemployed	0.97	0.96	0.54	0.66	0.19	0.27
Other ¹	0.60	0.55	0.39	0.37	0.17	0.18

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.31P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2725		0.6188		0.1629	
GENDER						
Male	0.4383		0.9017		0.9331	
Female	0.3932		0.4885		0.0290	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3210		0.6612		0.2440	
White	0.3021		0.3111		0.4466	
Black or African American	0.1236		0.0952		0.7597	
American Indian or Alaska Native	0.8464		0.6513		0.2240	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.3160		0.4757		0.2176	
Two or More Races	0.3066		0.8669		0.3732	
Hispanic or Latino	0.6366		0.0677		0.4067	
EDUCATION						
< High School	0.2755		0.6637		0.9641	
High School Graduate	0.6144		0.2746		0.5484	
Some College	0.8762		0.5828		0.2302	
College Graduate	0.2167		0.4186		0.5559	
CURRENT EMPLOYMENT						
Full-Time	0.8967		0.7581		0.8262	
Part-Time	0.2353		0.9852		0.0959	
Unemployed	0.8563		0.4805		0.4016	
Other ¹	0.1383		0.9959		0.9908	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.32C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	674	611	172	165	129	98
GENDER						
Male	502	461	139	128	97	80
Female	425	358	99	95	82	53
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	633	573	161	155	122	90
White	680	589	151	124	115	76
Black or African American	229	203	64	81	50	39
American Indian or Alaska Native	33	64	5	17	*	15
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	62	87	24	27	7	24
Two or More Races	107	85	8	15	4	10
Hispanic or Latino	257	203	59	65	41	43
EDUCATION						
< High School	227	216	68	60	55	27
High School Graduate	381	338	78	96	53	58
Some College	405	347	108	87	81	53
College Graduate	497	345	99	78	70	48
CURRENT EMPLOYMENT						
Full-Time	591	485	114	111	64	67
Part-Time	299	227	89	74	76	41
Unemployed	152	140	50	39	30	34
Other ¹	354	306	91	78	77	44

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.32D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.33	0.30	0.08	0.08	0.06	0.05
GENDER						
Male	0.52	0.47	0.14	0.13	0.10	0.08
Female	0.40	0.33	0.09	0.09	0.08	0.05
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.36	0.33	0.09	0.09	0.07	0.05
White	0.43	0.36	0.11	0.09	0.08	0.05
Black or African American	0.92	0.77	0.28	0.34	0.22	0.17
American Indian or Alaska Native	3.22	4.40	0.47	1.60	*	1.34
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	0.61	0.82	0.24	0.26	0.07	0.23
Two or More Races	3.44	2.66	0.30	0.51	0.14	0.34
Hispanic or Latino	0.90	0.69	0.21	0.22	0.14	0.15
EDUCATION						
< High School	0.80	0.74	0.25	0.21	0.20	0.10
High School Graduate	0.60	0.54	0.14	0.17	0.09	0.10
Some College	0.71	0.58	0.21	0.16	0.16	0.10
College Graduate	0.66	0.46	0.15	0.11	0.10	0.07
CURRENT EMPLOYMENT						
Full-Time	0.49	0.39	0.11	0.10	0.06	0.06
Part-Time	1.07	0.86	0.37	0.31	0.31	0.17
Unemployed	1.80	1.70	0.64	0.55	0.39	0.48
Other ¹	0.51	0.43	0.14	0.12	0.12	0.07

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.32P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1591		0.3133		0.5229	
GENDER						
Male	0.0956		0.7566		0.6486	
Female	0.7963		0.1868		0.6366	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1081		0.3047		0.4457	
White	0.1954		0.8316		0.3708	
Black or African American	0.5483		0.2504		0.3489	
American Indian or Alaska Native	0.0397		0.1373		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1389		0.7381		0.1778	
Two or More Races	0.8183		0.0582		0.2211	
Hispanic or Latino	0.8432		0.9061		0.8397	
EDUCATION						
< High School	0.4103		0.9281		0.0595	
High School Graduate	0.0289		0.0619		0.5121	
Some College	0.1835		0.9934		0.7170	
College Graduate	0.2345		0.9317		0.9323	
CURRENT EMPLOYMENT						
Full-Time	0.2545		0.1734		0.1824	
Part-Time	0.0682		0.5677		0.3254	
Unemployed	0.2844		0.7485		0.6898	
Other ¹	0.0330		0.9903		0.0954	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.33C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	692	627	189	184	132	105
GENDER						
Male	508	467	154	144	100	84
Female	438	371	109	105	84	59
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	650	585	177	168	125	95
White	698	606	164	141	119	83
Black or African American	231	205	65	82	51	40
American Indian or Alaska Native	34	64	6	18	1	15
Native Hawaiian or Other Pacific Islander	*	38	7	11	*	2
Asian	64	88	25	31	8	27
Two or More Races	110	86	12	17	4	9
Hispanic or Latino	264	214	65	82	44	46
EDUCATION						
< High School	235	223	73	68	56	29
High School Graduate	383	345	87	105	56	63
Some College	413	357	114	103	84	61
College Graduate	503	349	104	86	71	51
CURRENT EMPLOYMENT						
Full-Time	599	498	123	127	66	72
Part-Time	304	238	96	86	78	48
Unemployed	158	147	55	47	31	35
Other ¹	356	306	96	80	78	46

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.33D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.29	0.26	0.08	0.08	0.06	0.04
GENDER						
Male	0.44	0.40	0.13	0.12	0.09	0.07
Female	0.36	0.30	0.09	0.08	0.07	0.05
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.32	0.29	0.09	0.08	0.06	0.05
White	0.39	0.33	0.11	0.09	0.08	0.05
Black or African American	0.76	0.64	0.23	0.29	0.18	0.14
American Indian or Alaska Native	2.62	3.79	0.47	1.39	0.07	1.14
Native Hawaiian or Other Pacific Islander	*	3.72	0.87	1.17	*	0.21
Asian	0.53	0.69	0.21	0.24	0.07	0.21
Two or More Races	2.80	2.11	0.36	0.47	0.12	0.25
Hispanic or Latino	0.74	0.58	0.18	0.22	0.12	0.13
EDUCATION						
< High School	0.68	0.64	0.23	0.21	0.17	0.09
High School Graduate	0.50	0.45	0.13	0.15	0.08	0.09
Some College	0.59	0.49	0.18	0.15	0.13	0.09
College Graduate	0.62	0.43	0.14	0.12	0.10	0.07
CURRENT EMPLOYMENT						
Full-Time	0.44	0.36	0.10	0.10	0.06	0.06
Part-Time	0.79	0.64	0.28	0.26	0.23	0.14
Unemployed	1.22	1.17	0.46	0.43	0.26	0.32
Other ¹	0.47	0.39	0.13	0.11	0.11	0.06

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.33P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2311		0.2783		0.8494	
GENDER						
Male	0.1274		0.7616		0.6736	
Female	0.9093		0.1415		0.8238	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1555		0.4536		0.6879	
White	0.2675		0.8036		0.5200	
Black or African American	0.4320		0.1360		0.3953	
American Indian or Alaska Native	0.0352		0.1372		0.1044	
Native Hawaiian or Other Pacific Islander	*		0.9118		*	
Asian	0.2140		0.5394		0.0800	
Two or More Races	0.9874		0.0998		0.1208	
Hispanic or Latino	0.7721		0.3275		0.6268	
EDUCATION						
< High School	0.5203		0.6959		0.0598	
High School Graduate	0.0411		0.0290		0.4068	
Some College	0.1799		0.8358		0.9652	
College Graduate	0.1785		0.9466		0.9076	
CURRENT EMPLOYMENT						
Full-Time	0.2808		0.1591		0.1739	
Part-Time	0.0430		0.5983		0.7328	
Unemployed	0.3659		0.9361		0.5416	
Other ¹	0.0530		0.9640		0.1021	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.34C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	364	317	77	81	64	55
AGE						
12-17	5	9	4	7	1	4
18-25	40	40	16	22	12	9
26 or Older	359	313	75	79	63	54
GENDER						
Male	264	253	59	63	47	41
Female	238	171	51	53	43	40
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	346	298	72	72	57	54
White	295	251	59	54	50	42
Black or African American	165	141	40	42	28	29
American Indian or Alaska Native	23	49	2	1	1	1
Native Hawaiian or Other Pacific Islander	*	8	1	*	*	*
Asian	29	34	4	21	4	18
Two or More Races	48	46	1	6	1	*
Hispanic or Latino	123	92	29	43	28	19

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.34D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.14	0.12	0.03	0.03	0.02	0.02
AGE						
12-17	0.02	0.03	0.02	0.03	0.01	0.02
18-25	0.11	0.11	0.05	0.06	0.03	0.03
26 or Older	0.18	0.15	0.04	0.04	0.03	0.03
GENDER						
Male	0.21	0.20	0.05	0.05	0.04	0.03
Female	0.18	0.12	0.04	0.04	0.03	0.03
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.16	0.13	0.03	0.03	0.03	0.02
White	0.17	0.15	0.03	0.03	0.03	0.02
Black or African American	0.53	0.44	0.13	0.13	0.09	0.09
American Indian or Alaska Native	1.58	3.10	0.15	0.08	0.06	0.05
Native Hawaiian or Other Pacific Islander	*	0.80	0.12	*	*	*
Asian	0.22	0.25	0.03	0.15	0.03	0.13
Two or More Races	1.10	1.03	0.03	0.13	0.02	*
Hispanic or Latino	0.30	0.22	0.07	0.10	0.07	0.05

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.34P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3420		0.2313		0.7509	
AGE						
12-17	0.2332		0.2519		0.2562	
18-25	0.7324		0.2483		0.6272	
26 or Older	0.3458		0.3901		0.7624	
GENDER						
Male	0.1013		0.3261		0.8209	
Female	0.6339		0.4994		0.8301	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2352		0.3927		0.9830	
White	0.2236		0.7227		0.7367	
Black or African American	0.9049		0.5943		0.8455	
American Indian or Alaska Native	0.3181		0.5504		0.9153	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4577		0.2427		0.4548	
Two or More Races	0.4994		0.3398		*	
Hispanic or Latino	0.6376		0.3635		0.4636	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.35C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	5	9	4	7	1	4
GENDER						
Male	4	6	3	5	1	4
Female	3	6	3	4	*	1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4	7	3	4	1	3
White	4	7	3	4	1	3
Black or African American	*	*	*	*	*	*
American Indian or Alaska Native	0	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	*	*	*	*	*	*
Two or More Races	0	1	*	1	*	*
Hispanic or Latino	3	5	2	5	1	3
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	4	4	2	3	1	3
Female, White, Not Hispanic	2	6	2	3	*	1
Male, Black, Not Hispanic	*	*	*	*	*	*
Female, Black, Not Hispanic	*	*	*	*	*	*
Male, Hispanic	2	4	1	4	1	3
Female, Hispanic	3	3	2	3	0	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.35D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.02	0.03	0.02	0.03	0.01	0.02
GENDER						
Male	0.03	0.05	0.02	0.04	0.01	0.03
Female	0.03	0.05	0.02	0.03	*	0.01
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.02	0.04	0.02	0.02	0.01	0.01
White	0.03	0.05	0.02	0.03	0.01	0.02
Black or African American	*	*	*	*	*	*
American Indian or Alaska Native	0.15	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	*	*	*	*	*	*
Two or More Races	0.00	0.07	*	0.07	*	*
Hispanic or Latino	0.06	0.09	0.04	0.09	0.02	0.06
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.05	0.06	0.03	0.04	0.02	0.04
Female, White, Not Hispanic	0.03	0.08	0.03	0.04	*	0.01
Male, Black, Not Hispanic	*	*	*	*	*	*
Female, Black, Not Hispanic	*	*	*	*	*	*
Male, Hispanic	0.07	0.15	0.03	0.15	0.03	0.11
Female, Hispanic	0.10	0.11	0.08	0.11	0.01	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.35P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2332		0.2519		0.2562	
GENDER						
Male	0.7764		0.3857		0.2950	
Female	0.1571		0.4527		*	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2351		0.6627		0.4137	
White	0.2141		0.6450		0.4122	
Black or African American	*		*		*	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	*		*		*	
Two or More Races	0.3100		*		*	
Hispanic or Latino	0.6930		0.2505		0.4328	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.7787		0.9135		0.5761	
Female, White, Not Hispanic	0.0488		0.4048		*	
Male, Black, Not Hispanic	*		*		*	
Female, Black, Not Hispanic	*		*		*	
Male, Hispanic	0.3755		0.1611		0.3661	
Female, Hispanic	0.6829		0.9082		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.36C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	40	40	16	22	12	9
GENDER						
Male	31	32	14	19	11	8
Female	23	25	7	12	5	5
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	34	38	11	20	6	8
White	33	38	11	19	6	8
Black or African American	5	10	2	7	*	*
American Indian or Alaska Native	1	2	1	1	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	5	1	*	*	*	*
Two or More Races	3	4	1	*	1	*
Hispanic or Latino	21	14	11	9	10	5
EDUCATION						
< High School	21	17	6	9	4	5
High School Graduate	28	27	13	13	11	4
Some College	18	23	5	12	3	7
College Graduate	7	11	4	6	*	*
CURRENT EMPLOYMENT						
Full-Time	29	29	12	14	11	4
Part-Time	16	15	6	9	3	8
Unemployed	17	18	4	13	1	1
Other ¹	13	15	7	6	5	3

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.36D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.11	0.11	0.05	0.06	0.03	0.03
GENDER						
Male	0.18	0.18	0.08	0.11	0.06	0.04
Female	0.13	0.14	0.04	0.07	0.03	0.03
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.12	0.14	0.04	0.07	0.02	0.03
White	0.17	0.19	0.05	0.10	0.03	0.04
Black or African American	0.09	0.19	0.03	0.14	*	*
American Indian or Alaska Native	0.59	0.94	0.57	0.41	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	0.28	0.03	*	*	*	*
Two or More Races	0.38	0.47	0.19	*	0.11	*
Hispanic or Latino	0.29	0.19	0.16	0.13	0.14	0.07
EDUCATION						
< High School	0.38	0.34	0.11	0.18	0.07	0.10
High School Graduate	0.23	0.23	0.11	0.11	0.09	0.03
Some College	0.15	0.18	0.04	0.10	0.02	0.05
College Graduate	0.13	0.20	0.09	0.10	*	*
CURRENT EMPLOYMENT						
Full-Time	0.21	0.20	0.09	0.10	0.08	0.03
Part-Time	0.17	0.16	0.06	0.09	0.03	0.09
Unemployed	0.40	0.45	0.10	0.34	0.03	0.03
Other ¹	0.17	0.21	0.09	0.08	0.06	0.03

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.36P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.7324		0.2483		0.6272	
GENDER						
Male	0.4192		0.3984		0.8425	
Female	0.6331		0.4037		0.5148	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.5767		0.1124		0.9866	
White	0.5479		0.1691		0.9250	
Black or African American	0.3093		0.1570		*	
American Indian or Alaska Native	0.2201		0.6577		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0941		*		*	
Two or More Races	0.9680		*		*	
Hispanic or Latino	0.0611		0.7340		0.5370	
EDUCATION						
< High School	0.0729		0.5256		0.8535	
High School Graduate	0.9023		0.7400		0.2648	
Some College	0.2163		0.1155		0.4466	
College Graduate	0.3579		0.9106		*	
CURRENT EMPLOYMENT						
Full-Time	0.9108		0.4219		0.2880	
Part-Time	0.9885		0.6665		0.1870	
Unemployed	0.4087		0.2450		0.3910	
Other ¹	0.8750		0.6039		0.2898	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.37C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	359	313	75	79	63	54
GENDER						
Male	262	249	56	61	46	40
Female	236	168	50	52	43	40
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	342	295	70	70	57	53
White	296	255	58	52	50	41
Black or African American	174	149	40	42	28	29
American Indian or Alaska Native	23	*	2	1	*	*
Native Hawaiian or Other Pacific Islander	*	8	*	*	*	*
Asian	29	34	4	21	4	18
Two or More Races	48	46	*	6	*	*
Hispanic or Latino	121	91	27	42	26	18
EDUCATION						
< High School	147	134	39	41	23	18
High School Graduate	225	184	35	53	26	46
Some College	213	161	41	38	39	18
College Graduate	145	126	36	27	35	21
CURRENT EMPLOYMENT						
Full-Time	256	214	27	47	24	23
Part-Time	140	109	23	44	19	38
Unemployed	97	76	24	23	22	22
Other ¹	216	183	62	45	50	26

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.37D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.18	0.15	0.04	0.04	0.03	0.03
GENDER						
Male	0.27	0.25	0.06	0.06	0.05	0.04
Female	0.22	0.16	0.05	0.05	0.04	0.04
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.20	0.17	0.04	0.04	0.03	0.03
White	0.21	0.18	0.04	0.04	0.04	0.03
Black or African American	0.73	0.60	0.18	0.18	0.12	0.13
American Indian or Alaska Native	2.21	*	0.17	0.06	*	*
Native Hawaiian or Other Pacific Islander	*	1.10	*	*	*	*
Asian	0.28	0.32	0.04	0.20	0.04	0.17
Two or More Races	1.68	1.59	*	0.19	*	*
Hispanic or Latino	0.43	0.31	0.09	0.14	0.09	0.06
EDUCATION						
< High School	0.53	0.47	0.14	0.15	0.09	0.07
High School Graduate	0.38	0.31	0.06	0.09	0.04	0.08
Some College	0.40	0.30	0.08	0.07	0.08	0.03
College Graduate	0.21	0.19	0.05	0.04	0.05	0.03
CURRENT EMPLOYMENT						
Full-Time	0.23	0.19	0.03	0.04	0.02	0.02
Part-Time	0.57	0.45	0.09	0.18	0.08	0.16
Unemployed	1.21	1.04	0.32	0.33	0.28	0.31
Other ¹	0.32	0.27	0.09	0.07	0.08	0.04

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.37P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3458		0.3901		0.7624	
GENDER						
Male	0.0870		0.4730		0.7873	
Female	0.5559		0.6364		0.8816	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2782		0.6291		0.9539	
White	0.2701		0.9629		0.6916	
Black or African American	0.9553		0.7261		0.8499	
American Indian or Alaska Native	*		0.5357		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.3415		0.2422		0.4547	
Two or More Races	0.4926		*		*	
Hispanic or Latino	0.8342		0.3565		0.5212	
EDUCATION						
< High School	0.8915		0.9524		0.7351	
High School Graduate	0.1407		0.0868		0.0731	
Some College	0.3306		0.8440		0.0590	
College Graduate	0.1591		1.0000		0.5307	
CURRENT EMPLOYMENT						
Full-Time	0.3972		0.0708		0.8783	
Part-Time	0.8085		0.1009		0.2113	
Unemployed	0.7557		0.9563		0.9466	
Other ¹	0.2893		0.3112		0.1711	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.38C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	364	316	77	81	64	55
GENDER						
Male	264	253	58	63	47	41
Female	237	170	50	53	43	40
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	345	298	71	72	57	54
White	301	257	59	53	50	42
Black or African American	174	149	40	43	28	29
American Indian or Alaska Native	23	49	2	1	1	1
Native Hawaiian or Other Pacific Islander	*	8	*	*	*	*
Asian	29	34	4	21	4	18
Two or More Races	48	46	1	6	1	*
Hispanic or Latino	123	92	29	43	28	19
EDUCATION						
< High School	150	136	39	42	24	19
High School Graduate	227	185	37	54	27	46
Some College	214	163	42	40	39	19
College Graduate	145	126	36	28	35	21
CURRENT EMPLOYMENT						
Full-Time	258	217	30	49	26	24
Part-Time	141	110	23	45	19	39
Unemployed	100	78	25	27	22	22
Other ¹	217	183	62	45	51	26

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.38D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.15	0.13	0.03	0.03	0.03	0.02
GENDER						
Male	0.23	0.22	0.05	0.05	0.04	0.04
Female	0.19	0.14	0.04	0.04	0.04	0.03
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.17	0.15	0.04	0.04	0.03	0.03
White	0.19	0.16	0.04	0.03	0.03	0.03
Black or African American	0.59	0.50	0.14	0.15	0.10	0.10
American Indian or Alaska Native	1.79	3.46	0.17	0.08	0.07	0.06
Native Hawaiian or Other Pacific Islander	*	0.89	*	*	*	*
Asian	0.24	0.27	0.03	0.17	0.03	0.14
Two or More Races	1.34	1.26	0.04	0.15	0.02	*
Hispanic or Latino	0.35	0.25	0.08	0.12	0.08	0.05
EDUCATION						
< High School	0.44	0.40	0.12	0.13	0.07	0.06
High School Graduate	0.32	0.26	0.05	0.08	0.04	0.07
Some College	0.33	0.25	0.07	0.06	0.06	0.03
College Graduate	0.20	0.17	0.05	0.04	0.05	0.03
CURRENT EMPLOYMENT						
Full-Time	0.21	0.17	0.02	0.04	0.02	0.02
Part-Time	0.41	0.33	0.07	0.13	0.06	0.12
Unemployed	0.81	0.69	0.21	0.24	0.18	0.20
Other ¹	0.29	0.24	0.09	0.06	0.07	0.04

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.38P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3655		0.2640		0.7016	
GENDER						
Male	0.1071		0.3569		0.7602	
Female	0.6018		0.5317		0.8215	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2512		0.4065		0.9541	
White	0.2391		0.7458		0.7050	
Black or African American	0.9059		0.5948		0.8461	
American Indian or Alaska Native	0.3159		0.3953		0.9161	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4634		0.2426		0.4556	
Two or More Races	0.5103		0.3845		*	
Hispanic or Latino	0.6175		0.4308		0.4154	
EDUCATION						
< High School	0.9906		0.9381		0.7862	
High School Graduate	0.1564		0.0801		0.1309	
Some College	0.4185		0.8603		0.0812	
College Graduate	0.1513		0.9887		0.5269	
CURRENT EMPLOYMENT						
Full-Time	0.4205		0.0469		0.8434	
Part-Time	0.8095		0.0908		0.1356	
Unemployed	0.6098		0.6196		1.0000	
Other ¹	0.2783		0.2875		0.1454	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.39C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	707	636	204	159	132	88
AGE						
12-17	35	37	29	31	17	19
18-25	133	139	89	90	48	47
26 or Older	678	601	177	120	123	74
GENDER						
Male	524	461	167	129	119	67
Female	438	355	117	76	57	47
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	673	593	188	147	116	83
White	628	530	157	136	81	77
Black or African American	194	151	52	40	37	19
American Indian or Alaska Native	58	55	38	13	6	2
Native Hawaiian or Other Pacific Islander	40	30	3	12	2	2
Asian	89	94	24	31	18	15
Two or More Races	124	84	75	19	*	11
Hispanic or Latino	237	179	69	61	47	37

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.39D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.27	0.24	0.08	0.06	0.05	0.03
AGE						
12-17	0.14	0.15	0.12	0.13	0.07	0.08
18-25	0.38	0.40	0.25	0.26	0.14	0.13
26 or Older	0.33	0.29	0.09	0.06	0.06	0.04
GENDER						
Male	0.41	0.36	0.13	0.10	0.09	0.05
Female	0.32	0.26	0.09	0.06	0.04	0.03
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.30	0.27	0.08	0.07	0.05	0.04
White	0.37	0.31	0.09	0.08	0.05	0.05
Black or African American	0.62	0.48	0.17	0.13	0.12	0.06
American Indian or Alaska Native	3.54	3.00	2.63	0.90	0.41	0.15
Native Hawaiian or Other Pacific Islander	4.24	2.88	0.39	1.13	0.26	0.20
Asian	0.68	0.68	0.18	0.22	0.13	0.11
Two or More Races	2.52	1.70	1.68	0.43	*	0.25
Hispanic or Latino	0.58	0.42	0.17	0.14	0.11	0.09

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.39P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5990		0.3865		0.2769	
AGE						
12-17	0.0681		0.1625		0.4918	
18-25	0.0590		0.2608		0.0780	
26 or Older	0.9291		0.9728		0.8146	
GENDER						
Male	0.9436		0.7006		0.2366	
Female	0.4259		0.3068		0.8881	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.8673		0.3879		0.1346	
White	0.8669		0.5082		0.4959	
Black or African American	0.9350		0.6167		0.1167	
American Indian or Alaska Native	0.9352		0.4450		0.3088	
Native Hawaiian or Other Pacific Islander	0.4407		0.3384		0.8601	
Asian	0.8908		0.5719		0.6818	
Two or More Races	0.6443		0.6335		*	
Hispanic or Latino	0.3166		0.8508		0.4942	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.40C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	35	37	29	31	17	19
GENDER						
Male	28	26	24	21	14	15
Female	22	26	18	22	10	12
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	32	32	27	28	16	16
White	28	31	23	27	13	16
Black or African American	10	8	7	5	5	3
American Indian or Alaska Native	*	2	*	2	*	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	8	3	8	3	5	*
Two or More Races	6	6	5	6	3	3
Hispanic or Latino	17	20	15	16	6	10
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	21	22	18	18	10	12
Female, White, Not Hispanic	18	21	15	19	9	10
Male, Black, Not Hispanic	8	6	6	4	4	3
Female, Black, Not Hispanic	5	5	4	4	3	0
Male, Hispanic	14	14	13	11	5	6
Female, Hispanic	10	14	8	12	4	8

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.40D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.14	0.15	0.12	0.13	0.07	0.08
GENDER						
Male	0.22	0.20	0.19	0.17	0.11	0.12
Female	0.18	0.21	0.15	0.18	0.08	0.10
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.16	0.17	0.14	0.14	0.08	0.09
White	0.20	0.23	0.17	0.19	0.10	0.12
Black or African American	0.28	0.23	0.21	0.15	0.15	0.10
American Indian or Alaska Native	*	1.18	*	1.04	*	0.57
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	0.67	0.25	0.63	0.25	0.38	*
Two or More Races	0.76	0.77	0.64	0.75	0.40	0.35
Hispanic or Latino	0.31	0.35	0.27	0.29	0.11	0.18
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.30	0.31	0.25	0.26	0.14	0.18
Female, White, Not Hispanic	0.27	0.32	0.22	0.28	0.13	0.15
Male, Black, Not Hispanic	0.45	0.35	0.34	0.22	0.23	0.20
Female, Black, Not Hispanic	0.31	0.30	0.26	0.21	0.19	0.01
Male, Hispanic	0.49	0.49	0.43	0.39	0.18	0.22
Female, Hispanic	0.36	0.49	0.27	0.42	0.13	0.28

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.40P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0681		0.1625		0.4918	
GENDER						
Male	0.0485		0.0328		0.5388	
Female	0.6255		0.7579		0.7446	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0524		0.1854		0.1739	
White	0.2952		0.5519		0.4811	
Black or African American	0.4596		0.2871		0.4025	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0220		0.0523		*	
Two or More Races	0.6814		0.9221		0.9709	
Hispanic or Latino	0.8723		0.6937		0.2432	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.2431		0.2901		0.8367	
Female, White, Not Hispanic	0.7692		0.8000		0.3865	
Male, Black, Not Hispanic	0.5337		0.4392		0.9470	
Female, Black, Not Hispanic	0.6942		0.4659		0.0913	
Male, Hispanic	0.4793		0.1694		0.9545	
Female, Hispanic	0.5608		0.2944		0.0941	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.41C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	133	139	89	90	48	47
GENDER						
Male	101	105	72	70	42	32
Female	82	85	52	50	22	28
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	118	122	81	79	46	35
White	127	129	82	78	41	30
Black or African American	37	43	23	22	14	8
American Indian or Alaska Native	12	11	4	*	2	1
Native Hawaiian or Other Pacific Islander	9	*	2	*	*	*
Asian	27	28	20	24	15	11
Two or More Races	23	20	14	12	5	6
Hispanic or Latino	58	64	35	41	13	26
EDUCATION						
< High School	55	53	31	28	20	15
High School Graduate	81	80	47	49	25	21
Some College	89	96	57	61	30	32
College Graduate	59	64	42	39	19	18
CURRENT EMPLOYMENT						
Full-Time	100	99	54	57	27	25
Part-Time	72	77	51	51	27	23
Unemployed	54	48	35	31	13	17
Other ¹	62	60	40	35	23	18

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.41D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.38	0.40	0.25	0.26	0.14	0.13
GENDER						
Male	0.58	0.60	0.41	0.40	0.24	0.18
Female	0.47	0.49	0.30	0.29	0.13	0.16
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.43	0.44	0.29	0.29	0.17	0.13
White	0.53	0.56	0.37	0.38	0.20	0.15
Black or African American	0.68	0.80	0.44	0.43	0.27	0.16
American Indian or Alaska Native	4.17	4.58	1.52	*	0.78	0.72
Native Hawaiian or Other Pacific Islander	4.17	*	1.27	*	*	*
Asian	1.36	1.35	1.04	1.15	0.78	0.55
Two or More Races	2.68	2.31	1.89	1.44	0.69	0.79
Hispanic or Latino	0.80	0.87	0.48	0.56	0.18	0.35
EDUCATION						
< High School	0.88	0.99	0.55	0.56	0.36	0.30
High School Graduate	0.64	0.62	0.38	0.39	0.21	0.18
Some College	0.61	0.67	0.43	0.46	0.24	0.25
College Graduate	0.95	0.98	0.74	0.65	0.38	0.31
CURRENT EMPLOYMENT						
Full-Time	0.62	0.61	0.37	0.38	0.20	0.18
Part-Time	0.66	0.71	0.49	0.50	0.28	0.24
Unemployed	1.18	1.15	0.81	0.77	0.32	0.44
Other ¹	0.78	0.74	0.51	0.45	0.31	0.25

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.41P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0590		0.2608		0.0780	
GENDER						
Male	0.0628		0.1070		0.0050	
Female	0.4307		0.7879		0.3782	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0566		0.1353		0.0018	
White	0.0635		0.0877		0.0025	
Black or African American	0.8309		0.6831		0.0306	
American Indian or Alaska Native	0.5134		*		0.7298	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.5119		0.9285		0.8604	
Two or More Races	0.6255		0.7316		0.2947	
Hispanic or Latino	0.6719		0.5924		0.0320	
EDUCATION						
< High School	0.3562		0.8221		0.4487	
High School Graduate	0.2578		0.7238		0.2283	
Some College	0.3972		0.4313		0.2597	
College Graduate	0.2410		0.2112		0.6343	
CURRENT EMPLOYMENT						
Full-Time	0.7186		0.7474		0.4275	
Part-Time	0.3911		0.5324		0.0512	
Unemployed	0.1051		0.3254		0.3865	
Other ¹	0.0552		0.1254		0.1905	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.42C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	678	601	177	120	123	74
GENDER						
Male	506	439	139	101	111	59
Female	425	340	103	55	51	35
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	646	564	163	115	105	70
White	692	583	135	103	71	67
Black or African American	202	154	46	33	34	17
American Indian or Alaska Native	57	54	*	9	4	1
Native Hawaiian or Other Pacific Islander	*	29	*	*	*	*
Asian	83	90	10	21	8	9
Two or More Races	122	81	*	13	*	9
Hispanic or Latino	224	160	55	39	44	23
EDUCATION						
< High School	206	169	37	22	14	16
High School Graduate	353	292	69	52	50	37
Some College	416	325	93	65	29	40
College Graduate	490	373	131	78	108	45
CURRENT EMPLOYMENT						
Full-Time	586	475	94	96	59	63
Part-Time	316	224	110	40	79	22
Unemployed	156	120	44	28	21	18
Other ¹	333	280	77	38	56	16

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.42D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.33	0.29	0.09	0.06	0.06	0.04
GENDER						
Male	0.52	0.45	0.14	0.10	0.11	0.06
Female	0.40	0.32	0.10	0.05	0.05	0.03
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.37	0.32	0.09	0.07	0.06	0.04
White	0.44	0.37	0.10	0.07	0.05	0.05
Black or African American	0.84	0.62	0.20	0.14	0.15	0.07
American Indian or Alaska Native	4.74	3.81	*	0.82	0.37	0.11
Native Hawaiian or Other Pacific Islander	*	3.75	*	*	*	*
Asian	0.82	0.84	0.10	0.20	0.08	0.09
Two or More Races	3.73	2.55	*	0.47	*	0.30
Hispanic or Latino	0.79	0.54	0.19	0.13	0.16	0.08
EDUCATION						
< High School	0.74	0.61	0.14	0.08	0.05	0.06
High School Graduate	0.56	0.47	0.12	0.09	0.09	0.06
Some College	0.72	0.55	0.18	0.12	0.06	0.08
College Graduate	0.65	0.50	0.19	0.11	0.16	0.07
CURRENT EMPLOYMENT						
Full-Time	0.48	0.38	0.09	0.09	0.06	0.06
Part-Time	1.13	0.86	0.45	0.17	0.33	0.09
Unemployed	1.76	1.51	0.56	0.41	0.27	0.26
Other ¹	0.49	0.41	0.12	0.06	0.09	0.02

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.42P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9291		0.9728		0.8146	
GENDER						
Male	0.6888		0.3558		0.9902	
Female	0.5199		0.1572		0.5709	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7900		0.8948		1.0000	
White	0.8437		0.6678		0.3413	
Black or African American	0.8670		0.8645		0.4957	
American Indian or Alaska Native	0.7118		*		0.4839	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.6403		0.1507		0.9389	
Two or More Races	0.6943		*		*	
Hispanic or Latino	0.3495		0.6323		0.5017	
EDUCATION						
< High School	0.4864		0.4820		0.3544	
High School Graduate	0.0487		0.3483		0.3262	
Some College	0.2364		0.5936		0.0938	
College Graduate	0.6872		0.7973		0.5175	
CURRENT EMPLOYMENT						
Full-Time	0.5687		0.0294		0.0937	
Part-Time	0.0728		0.1613		0.3434	
Unemployed	0.4988		0.6243		0.8523	
Other ¹	0.4995		0.2318		0.1066	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.43C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	706	630	200	154	131	84
GENDER						
Male	524	456	162	125	119	64
Female	436	355	117	75	56	45
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	672	587	184	143	115	79
White	718	610	157	132	80	73
Black or African American	206	165	51	40	37	18
American Indian or Alaska Native	58	55	*	13	4	2
Native Hawaiian or Other Pacific Islander	*	30	2	11	1	2
Asian	88	94	23	31	17	15
Two or More Races	124	83	74	18	*	11
Hispanic or Latino	236	178	67	59	46	36
EDUCATION						
< High School	218	177	49	36	25	22
High School Graduate	358	301	84	71	56	44
Some College	428	345	106	92	42	51
College Graduate	498	384	136	86	109	48
CURRENT EMPLOYMENT						
Full-Time	602	494	106	113	64	66
Part-Time	325	237	120	63	83	32
Unemployed	166	127	56	40	25	21
Other ¹	336	284	87	51	60	24

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.43D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.30	0.26	0.08	0.06	0.06	0.04
GENDER						
Male	0.46	0.39	0.14	0.11	0.10	0.06
Female	0.35	0.28	0.09	0.06	0.05	0.04
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.33	0.29	0.09	0.07	0.06	0.04
White	0.40	0.33	0.10	0.08	0.05	0.05
Black or African American	0.70	0.53	0.18	0.14	0.13	0.07
American Indian or Alaska Native	3.99	3.30	*	1.00	0.34	0.15
Native Hawaiian or Other Pacific Islander	*	3.19	0.33	1.24	0.10	0.20
Asian	0.74	0.74	0.19	0.25	0.14	0.12
Two or More Races	3.03	2.07	2.03	0.50	*	0.29
Hispanic or Latino	0.66	0.48	0.19	0.16	0.13	0.10
EDUCATION						
< High School	0.64	0.54	0.15	0.11	0.08	0.07
High School Graduate	0.48	0.40	0.12	0.10	0.08	0.06
Some College	0.59	0.48	0.17	0.14	0.07	0.08
College Graduate	0.61	0.47	0.19	0.11	0.15	0.06
CURRENT EMPLOYMENT						
Full-Time	0.43	0.35	0.09	0.09	0.05	0.05
Part-Time	0.84	0.65	0.35	0.19	0.24	0.10
Unemployed	1.22	1.03	0.47	0.38	0.21	0.19
Other ¹	0.44	0.37	0.12	0.07	0.08	0.03

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.43P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.6374		0.5166		0.3199	
GENDER						
Male	1.0000		0.9506		0.2663	
Female	0.4286		0.2755		0.9453	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.9141		0.5076		0.1902	
White	0.8805		0.5676		0.5710	
Black or African American	0.9065		0.7198		0.1420	
American Indian or Alaska Native	0.7943		*		0.3944	
Native Hawaiian or Other Pacific Islander	*		0.3054		0.6638	
Asian	0.7856		0.3156		0.9457	
Two or More Races	0.6362		0.6239		*	
Hispanic or Latino	0.3105		0.9242		0.6436	
EDUCATION						
< High School	0.2918		0.2399		0.8222	
High School Graduate	0.0999		0.3932		0.1678	
Some College	0.1848		0.9418		0.6241	
College Graduate	0.5851		0.9945		0.4864	
CURRENT EMPLOYMENT						
Full-Time	0.6208		0.0325		0.1956	
Part-Time	0.0522		0.1285		0.0995	
Unemployed	0.2125		0.3504		0.6390	
Other ¹	0.7675		0.0644		0.0389	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.44C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	519	428	105	95	64	65
AGE						
12-17	52	56	29	36	15	19
18-25	85	93	41	37	18	16
26 or Older	505	412	90	85	59	59
GENDER						
Male	416	346	95	83	56	57
Female	289	227	44	48	29	30
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	489	402	96	82	60	49
White	458	376	93	73	55	45
Black or African American	99	89	19	32	14	12
American Indian or Alaska Native	25	44	2	3	1	2
Native Hawaiian or Other Pacific Islander	*	22	*	2	*	1
Asian	54	56	18	20	17	15
Two or More Races	108	62	12	11	8	8
Hispanic or Latino	185	142	34	48	23	39

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.44D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.20	0.16	0.04	0.04	0.02	0.02
AGE						
12-17	0.21	0.23	0.12	0.14	0.06	0.07
18-25	0.24	0.27	0.12	0.11	0.05	0.05
26 or Older	0.25	0.20	0.04	0.04	0.03	0.03
GENDER						
Male	0.33	0.27	0.07	0.06	0.04	0.04
Female	0.21	0.17	0.03	0.03	0.02	0.02
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.22	0.18	0.04	0.04	0.03	0.02
White	0.27	0.22	0.05	0.04	0.03	0.03
Black or African American	0.32	0.28	0.06	0.10	0.04	0.04
American Indian or Alaska Native	1.74	2.79	0.13	0.22	0.09	0.16
Native Hawaiian or Other Pacific Islander	*	2.07	*	0.22	*	0.12
Asian	0.40	0.40	0.14	0.15	0.12	0.11
Two or More Races	2.26	1.36	0.29	0.24	0.20	0.18
Hispanic or Latino	0.45	0.33	0.08	0.11	0.05	0.09

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.44P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9936		0.6320		0.6223	
AGE						
12-17	0.7986		0.4803		0.2319	
18-25	0.1541		0.0698		0.2937	
26 or Older	0.7675		0.2517		0.6026	
GENDER						
Male	0.6564		0.7158		0.5786	
Female	0.5238		0.7355		0.9927	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.9096		0.8795		0.6013	
White	0.9951		0.7468		0.5747	
Black or African American	0.2654		0.9524		0.6970	
American Indian or Alaska Native	0.5061		0.3127		0.6349	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.6796		0.8059		0.7858	
Two or More Races	0.5722		0.5809		0.4003	
Hispanic or Latino	0.8436		0.1397		0.0572	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.45C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	52	56	29	36	15	19
GENDER						
Male	40	37	21	23	10	13
Female	34	41	20	27	11	13
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	46	47	26	30	14	15
White	38	40	23	27	11	12
Black or African American	21	22	13	10	8	6
American Indian or Alaska Native	4	3	1	2	1	1
Native Hawaiian or Other Pacific Islander	1	*	*	*	*	*
Asian	20	11	6	8	1	2
Two or More Races	5	11	3	7	1	6
Hispanic or Latino	26	30	15	19	6	11
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	28	27	17	17	8	8
Female, White, Not Hispanic	24	29	15	21	8	9
Male, Black, Not Hispanic	15	14	8	7	5	4
Female, Black, Not Hispanic	15	18	11	7	7	5
Male, Hispanic	21	22	13	13	6	9
Female, Hispanic	18	22	8	15	3	5

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.45D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.21	0.23	0.12	0.14	0.06	0.07
GENDER						
Male	0.31	0.29	0.17	0.18	0.08	0.10
Female	0.28	0.34	0.17	0.22	0.09	0.11
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.24	0.24	0.13	0.16	0.07	0.08
White	0.26	0.29	0.16	0.20	0.08	0.09
Black or African American	0.57	0.59	0.37	0.27	0.23	0.17
American Indian or Alaska Native	2.31	1.58	0.67	1.12	0.50	0.57
Native Hawaiian or Other Pacific Islander	1.08	*	*	*	*	*
Asian	1.49	0.92	0.49	0.64	0.05	0.19
Two or More Races	0.73	1.37	0.39	0.86	0.20	0.70
Hispanic or Latino	0.46	0.54	0.27	0.34	0.11	0.19
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.39	0.38	0.23	0.24	0.12	0.12
Female, White, Not Hispanic	0.35	0.43	0.22	0.31	0.11	0.14
Male, Black, Not Hispanic	0.81	0.74	0.43	0.38	0.26	0.20
Female, Black, Not Hispanic	0.81	0.98	0.62	0.40	0.39	0.28
Male, Hispanic	0.68	0.73	0.44	0.43	0.20	0.33
Female, Hispanic	0.63	0.76	0.30	0.53	0.11	0.18

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.45P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.7986		0.4803		0.2319	
GENDER						
Male	0.1107		0.5242		0.4913	
Female	0.2061		0.1202		0.3155	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.5584		0.8968		0.5915	
White	0.5970		0.4595		0.5167	
Black or African American	0.7180		0.0055		0.3908	
American Indian or Alaska Native	0.3267		0.4159		0.9476	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.3416		0.8536		0.5143	
Two or More Races	0.0929		0.2382		0.3208	
Hispanic or Latino	0.6365		0.1215		0.1511	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.1683		0.4640		0.6971	
Female, White, Not Hispanic	0.5152		0.0917		0.6032	
Male, Black, Not Hispanic	0.2660		0.1649		0.6424	
Female, Black, Not Hispanic	0.6079		0.0156		0.4691	
Male, Hispanic	0.9638		0.9203		0.3567	
Female, Hispanic	0.4644		0.0310		0.1858	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.46C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	85	93	41	37	18	16
GENDER						
Male	64	71	34	29	17	13
Female	53	53	22	23	8	10
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	75	81	35	29	17	15
White	74	79	32	28	14	13
Black or African American	20	19	8	5	5	4
American Indian or Alaska Native	6	7	1	2	1	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	15	13	5	4	3	1
Two or More Races	15	12	12	6	*	4
Hispanic or Latino	40	41	20	20	6	7
EDUCATION						
< High School	36	40	12	13	5	8
High School Graduate	51	49	27	18	12	7
Some College	52	59	23	26	11	10
College Graduate	35	40	17	16	6	8
CURRENT EMPLOYMENT						
Full-Time	55	59	20	18	10	7
Part-Time	44	49	22	23	9	10
Unemployed	38	33	23	14	11	9
Other ¹	36	40	16	17	6	6

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.46D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.24	0.27	0.12	0.11	0.05	0.05
GENDER						
Male	0.37	0.40	0.20	0.16	0.10	0.07
Female	0.31	0.31	0.13	0.13	0.04	0.06
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.27	0.29	0.13	0.11	0.06	0.05
White	0.35	0.38	0.16	0.14	0.07	0.07
Black or African American	0.39	0.37	0.16	0.09	0.09	0.08
American Indian or Alaska Native	2.30	3.39	0.53	0.86	0.44	0.28
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	0.81	0.61	0.29	0.22	0.17	0.05
Two or More Races	1.94	1.51	1.58	0.76	*	0.57
Hispanic or Latino	0.56	0.56	0.28	0.28	0.09	0.09
EDUCATION						
< High School	0.61	0.77	0.21	0.27	0.09	0.15
High School Graduate	0.41	0.40	0.22	0.15	0.10	0.06
Some College	0.40	0.44	0.19	0.21	0.09	0.08
College Graduate	0.65	0.67	0.33	0.27	0.12	0.14
CURRENT EMPLOYMENT						
Full-Time	0.39	0.40	0.15	0.13	0.07	0.05
Part-Time	0.42	0.50	0.23	0.24	0.09	0.10
Unemployed	0.87	0.81	0.53	0.36	0.27	0.23
Other ¹	0.46	0.52	0.21	0.22	0.08	0.08

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.46P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1541		0.0698		0.2937	
GENDER						
Male	0.7486		0.0227		0.1729	
Female	0.0491		0.9609		0.8001	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2486		0.0454		0.2654	
White	0.5756		0.1397		0.3437	
Black or African American	0.0893		0.0335		0.5425	
American Indian or Alaska Native	0.6505		0.6330		0.6067	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1216		0.3954		0.3066	
Two or More Races	0.8199		0.6713		*	
Hispanic or Latino	0.3713		0.7205		0.9598	
EDUCATION						
< High School	0.3212		0.4936		0.8031	
High School Graduate	0.0595		0.0083		0.2775	
Some College	0.3943		0.6271		0.4280	
College Graduate	0.5893		0.5890		0.8727	
CURRENT EMPLOYMENT						
Full-Time	0.3190		0.2456		0.2462	
Part-Time	0.6854		0.4748		0.7857	
Unemployed	0.1195		0.0738		0.5925	
Other ¹	0.9234		0.8596		0.9515	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.47C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	505	412	90	85	59	59
GENDER						
Male	405	335	84	80	53	54
Female	280	216	34	29	26	25
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	480	392	87	74	55	44
White	487	390	85	65	51	40
Black or African American	99	90	11	30	10	9
American Indian or Alaska Native	22	43	1	2	*	2
Native Hawaiian or Other Pacific Islander	*	20	*	*	*	*
Asian	47	53	16	18	16	15
Two or More Races	107	60	2	5	*	3
Hispanic or Latino	174	131	23	39	21	37
EDUCATION						
< High School	136	131	20	26	16	10
High School Graduate	253	190	30	32	24	14
Some College	279	232	40	39	32	31
College Graduate	356	265	72	61	39	47
CURRENT EMPLOYMENT						
Full-Time	411	332	78	69	48	52
Part-Time	207	155	25	29	21	19
Unemployed	103	82	22	20	*	11
Other ¹	240	197	32	38	27	16

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.47D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.25	0.20	0.04	0.04	0.03	0.03
GENDER						
Male	0.42	0.34	0.09	0.08	0.05	0.05
Female	0.27	0.20	0.03	0.03	0.02	0.02
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.28	0.22	0.05	0.04	0.03	0.02
White	0.33	0.26	0.06	0.05	0.04	0.03
Black or African American	0.43	0.38	0.05	0.13	0.05	0.04
American Indian or Alaska Native	2.22	3.63	0.11	0.20	*	0.19
Native Hawaiian or Other Pacific Islander	*	2.70	*	*	*	*
Asian	0.46	0.50	0.16	0.17	0.16	0.14
Two or More Races	3.38	2.05	0.07	0.17	*	0.11
Hispanic or Latino	0.61	0.45	0.08	0.13	0.07	0.12
EDUCATION						
< High School	0.50	0.47	0.08	0.10	0.06	0.04
High School Graduate	0.43	0.32	0.05	0.06	0.04	0.02
Some College	0.51	0.42	0.08	0.07	0.06	0.06
College Graduate	0.50	0.36	0.11	0.09	0.06	0.07
CURRENT EMPLOYMENT						
Full-Time	0.36	0.29	0.07	0.06	0.05	0.05
Part-Time	0.81	0.62	0.10	0.12	0.09	0.08
Unemployed	1.29	1.12	0.28	0.29	*	0.15
Other ¹	0.36	0.29	0.05	0.06	0.04	0.02

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.47P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.7675		0.2517		0.6026	
GENDER						
Male	0.5129		0.1381		0.4073	
Female	0.6676		0.5130		0.5974	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7147		0.4744		0.7246	
White	0.8945		0.9802		0.6333	
Black or African American	0.1166		0.0507		0.7098	
American Indian or Alaska Native	0.4817		0.6086		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.2367		0.6485		0.8512	
Two or More Races	0.4326		0.2819		*	
Hispanic or Latino	0.9554		0.1770		0.1041	
EDUCATION						
< High School	0.5029		0.3884		0.9293	
High School Graduate	0.7306		0.8602		0.1573	
Some College	0.4085		0.6154		0.7694	
College Graduate	0.6098		0.3356		0.2522	
CURRENT EMPLOYMENT						
Full-Time	0.8034		0.5092		0.4533	
Part-Time	0.5523		0.2994		0.8785	
Unemployed	0.6088		0.6963		*	
Other ¹	0.3721		0.6375		0.3613	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.48C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	515	424	101	88	61	61
GENDER						
Male	412	341	92	80	55	55
Female	287	225	40	38	27	27
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	486	399	94	75	58	46
White	496	402	90	67	53	42
Black or African American	99	92	14	30	11	10
American Indian or Alaska Native	25	44	2	3	1	2
Native Hawaiian or Other Pacific Islander	*	21	*	2	*	*
Asian	50	54	17	19	17	15
Two or More Races	107	61	12	8	8	5
Hispanic or Latino	182	140	31	44	22	37
EDUCATION						
< High School	141	135	23	29	17	13
High School Graduate	257	198	40	36	27	15
Some College	283	241	46	45	34	32
College Graduate	359	267	74	60	40	48
CURRENT EMPLOYMENT						
Full-Time	417	338	81	68	49	53
Part-Time	213	164	33	37	23	21
Unemployed	110	89	31	24	11	14
Other ¹	243	201	35	41	27	17

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.48D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.22	0.18	0.04	0.04	0.03	0.03
GENDER						
Male	0.36	0.29	0.08	0.07	0.05	0.05
Female	0.23	0.18	0.03	0.03	0.02	0.02
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.24	0.20	0.05	0.04	0.03	0.02
White	0.29	0.24	0.06	0.04	0.03	0.03
Black or African American	0.35	0.31	0.05	0.11	0.04	0.04
American Indian or Alaska Native	1.93	3.12	0.13	0.21	0.08	0.17
Native Hawaiian or Other Pacific Islander	*	2.30	*	0.20	*	*
Asian	0.41	0.43	0.14	0.15	0.14	0.12
Two or More Races	2.72	1.63	0.34	0.22	0.24	0.15
Hispanic or Latino	0.51	0.38	0.09	0.12	0.06	0.10
EDUCATION						
< High School	0.42	0.41	0.07	0.09	0.05	0.04
High School Graduate	0.36	0.27	0.06	0.05	0.04	0.02
Some College	0.42	0.35	0.07	0.07	0.05	0.05
College Graduate	0.47	0.34	0.10	0.08	0.06	0.07
CURRENT EMPLOYMENT						
Full-Time	0.32	0.26	0.07	0.05	0.04	0.04
Part-Time	0.59	0.47	0.10	0.11	0.07	0.06
Unemployed	0.89	0.78	0.26	0.22	0.10	0.13
Other ¹	0.32	0.26	0.05	0.06	0.04	0.02

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.48P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9802		0.7694		0.8427	
GENDER						
Male	0.5529		0.5786		0.6713	
Female	0.4089		0.6018		0.6749	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.8709		0.9305		0.4964	
White	0.9691		0.5860		0.4646	
Black or African American	0.2257		0.2011		0.9210	
American Indian or Alaska Native	0.4417		0.5317		0.5737	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4648		0.8314		0.7292	
Two or More Races	0.4534		0.9558		0.6637	
Hispanic or Latino	0.7774		0.3552		0.1153	
EDUCATION						
< High School	0.4161		0.3563		0.9833	
High School Graduate	0.9397		0.0938		0.0820	
Some College	0.3333		0.8751		0.9792	
College Graduate	0.6779		0.3896		0.2678	
CURRENT EMPLOYMENT						
Full-Time	0.9380		0.7105		0.5855	
Part-Time	0.4995		0.7284		0.8021	
Unemployed	0.2945		0.3904		0.4160	
Other ¹	0.3905		0.6395		0.3831	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.49C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	785	658	418	321	256	222
AGE						
12-17	63	69	50	60	32	41
18-25	149	165	110	115	69	71
26 or Older	751	622	386	286	244	207
GENDER						
Male	529	460	276	228	180	159
Female	556	445	284	212	179	149
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	731	617	364	291	228	204
White	671	568	327	264	203	175
Black or African American	205	184	111	104	94	90
American Indian or Alaska Native	45	49	30	32	8	15
Native Hawaiian or Other Pacific Islander	49	36	11	25	5	24
Asian	144	121	56	54	29	34
Two or More Races	126	82	66	37	23	27
Hispanic or Latino	272	216	172	134	121	90

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.49D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.30	0.25	0.16	0.12	0.10	0.08
AGE						
12-17	0.25	0.28	0.20	0.24	0.13	0.16
18-25	0.43	0.47	0.32	0.33	0.20	0.20
26 or Older	0.37	0.30	0.19	0.14	0.12	0.10
GENDER						
Male	0.42	0.36	0.22	0.18	0.14	0.12
Female	0.41	0.33	0.21	0.16	0.13	0.11
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.33	0.28	0.16	0.13	0.10	0.09
White	0.39	0.33	0.19	0.16	0.12	0.10
Black or African American	0.66	0.58	0.35	0.33	0.30	0.28
American Indian or Alaska Native	2.97	2.68	2.04	1.98	0.53	0.99
Native Hawaiian or Other Pacific Islander	4.88	3.30	1.29	2.42	0.61	2.30
Asian	1.02	0.85	0.42	0.40	0.22	0.25
Two or More Races	2.55	1.67	1.48	0.80	0.54	0.59
Hispanic or Latino	0.66	0.51	0.42	0.32	0.29	0.21

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.49P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5162		0.3072		0.9662	
AGE						
12-17	0.2713		0.1489		0.0396	
18-25	0.7131		0.4133		0.2389	
26 or Older	0.5229		0.2962		0.9959	
GENDER						
Male	0.0877		1.0000		0.9761	
Female	0.4551		0.1298		0.9266	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2524		0.4469		0.6722	
White	0.4254		0.1196		0.6057	
Black or African American	0.1029		0.0594		0.3702	
American Indian or Alaska Native	0.7613		0.8184		0.0154	
Native Hawaiian or Other Pacific Islander	0.3814		0.2179		0.0331	
Asian	0.4035		0.3663		0.3943	
Two or More Races	0.4872		0.3551		0.3801	
Hispanic or Latino	0.2631		0.3941		0.3342	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.50C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	63	69	50	60	32	41
GENDER						
Male	47	48	39	40	23	26
Female	44	51	32	45	21	31
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	55	58	44	50	28	33
White	51	54	40	45	23	28
Black or African American	26	28	19	23	15	16
American Indian or Alaska Native	7	7	6	6	2	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	10	10	7	8	3	6
Two or More Races	14	12	8	8	4	5
Hispanic or Latino	34	39	25	33	15	22
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	37	38	29	31	16	18
Female, White, Not Hispanic	33	40	26	34	15	21
Male, Black, Not Hispanic	19	17	14	14	12	10
Female, Black, Not Hispanic	18	22	13	18	9	13
Male, Hispanic	24	28	20	22	10	14
Female, Hispanic	25	29	18	25	12	18

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.50D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.25	0.28	0.20	0.24	0.13	0.16
GENDER						
Male	0.37	0.38	0.30	0.32	0.18	0.21
Female	0.36	0.42	0.27	0.37	0.17	0.25
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.28	0.30	0.23	0.26	0.14	0.17
White	0.33	0.37	0.27	0.32	0.17	0.20
Black or African American	0.70	0.74	0.52	0.62	0.42	0.45
American Indian or Alaska Native	3.84	3.21	3.61	2.96	1.14	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	0.79	0.83	0.55	0.67	0.20	0.53
Two or More Races	1.69	1.44	1.05	1.00	0.55	0.64
Hispanic or Latino	0.61	0.69	0.45	0.58	0.28	0.39
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.48	0.52	0.39	0.44	0.23	0.27
Female, White, Not Hispanic	0.45	0.57	0.37	0.50	0.22	0.31
Male, Black, Not Hispanic	1.00	0.93	0.78	0.74	0.66	0.56
Female, Black, Not Hispanic	0.97	1.14	0.72	0.97	0.54	0.72
Male, Hispanic	0.81	0.93	0.67	0.75	0.36	0.49
Female, Hispanic	0.86	0.99	0.61	0.87	0.43	0.62

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.50P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2713		0.1489		0.0396	
GENDER						
Male	0.5446		0.4700		0.2751	
Female	0.0316		0.0058		0.0681	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.5519		0.4308		0.1055	
White	0.8788		0.7985		0.1933	
Black or African American	0.0587		0.0462		0.5404	
American Indian or Alaska Native	0.6766		0.6081		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.9074		0.5202		0.0747	
Two or More Races	0.4684		0.4040		0.4371	
Hispanic or Latino	0.2814		0.1413		0.1701	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.4596		0.2544		0.5191	
Female, White, Not Hispanic	0.3337		0.1373		0.2284	
Male, Black, Not Hispanic	0.6782		0.6058		0.8497	
Female, Black, Not Hispanic	0.0262		0.0266		0.2967	
Male, Hispanic	0.8609		0.7405		0.3861	
Female, Hispanic	0.1709		0.0815		0.2934	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.51C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	149	165	110	115	69	71
GENDER						
Male	106	112	82	85	51	52
Female	101	111	72	77	45	46
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	130	148	96	105	62	64
White	160	163	109	100	60	58
Black or African American	53	64	32	43	23	23
American Indian or Alaska Native	11	13	7	9	6	6
Native Hawaiian or Other Pacific Islander	10	*	6	*	1	*
Asian	29	36	18	28	10	14
Two or More Races	24	24	18	17	8	11
Hispanic or Latino	69	72	47	51	30	33
EDUCATION						
< High School	68	74	48	45	31	27
High School Graduate	98	105	68	71	39	45
Some College	108	120	74	79	42	45
College Graduate	76	84	48	51	26	29
CURRENT EMPLOYMENT						
Full-Time	119	121	78	79	45	50
Part-Time	100	102	68	67	40	35
Unemployed	63	58	44	40	26	23
Other ¹	72	78	49	51	29	33

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.51D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.43	0.47	0.32	0.33	0.20	0.20
GENDER						
Male	0.61	0.64	0.47	0.48	0.29	0.30
Female	0.58	0.64	0.41	0.45	0.26	0.26
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.47	0.54	0.35	0.38	0.22	0.23
White	0.58	0.65	0.44	0.46	0.28	0.28
Black or African American	0.88	1.04	0.59	0.79	0.45	0.44
American Indian or Alaska Native	3.64	4.90	2.66	3.97	2.41	2.98
Native Hawaiian or Other Pacific Islander	4.51	*	3.05	*	0.48	*
Asian	1.48	1.59	0.99	1.30	0.54	0.67
Two or More Races	2.83	2.61	2.31	2.03	1.11	1.40
Hispanic or Latino	0.95	0.98	0.65	0.69	0.42	0.45
EDUCATION						
< High School	1.03	1.21	0.81	0.83	0.53	0.54
High School Graduate	0.69	0.75	0.52	0.55	0.32	0.36
Some College	0.69	0.78	0.52	0.58	0.33	0.35
College Graduate	1.05	1.14	0.79	0.81	0.50	0.49
CURRENT EMPLOYMENT						
Full-Time	0.70	0.67	0.51	0.51	0.32	0.34
Part-Time	0.79	0.87	0.60	0.63	0.40	0.35
Unemployed	1.26	1.28	0.97	0.97	0.60	0.57
Other ¹	0.84	0.87	0.61	0.62	0.38	0.43

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.51P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.7131		0.4133		0.2389	
GENDER						
Male	0.2663		0.2913		0.3004	
Female	0.5490		0.9595		0.5290	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.9764		0.4450		0.1322	
White	0.5214		0.0875		0.1475	
Black or African American	0.1246		0.0722		0.2255	
American Indian or Alaska Native	0.2052		0.2683		0.7293	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.8704		0.8941		0.9728	
Two or More Races	0.7626		0.6098		0.7326	
Hispanic or Latino	0.4747		0.7843		0.6919	
EDUCATION						
< High School	0.8683		0.9777		0.2338	
High School Graduate	0.2404		0.3111		0.3901	
Some College	0.9626		0.8469		0.8570	
College Graduate	0.3518		0.3899		0.5583	
CURRENT EMPLOYMENT						
Full-Time	0.6108		0.5363		0.8214	
Part-Time	0.5740		0.4158		0.2812	
Unemployed	0.2877		0.1636		0.1225	
Other ¹	0.7294		0.4755		0.5421	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.52C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	751	622	386	286	244	207
GENDER						
Male	514	437	253	206	173	148
Female	539	421	273	187	173	136
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	699	580	334	262	216	190
White	774	615	318	239	199	163
Black or African American	215	196	106	100	90	88
American Indian or Alaska Native	42	43	28	29	5	13
Native Hawaiian or Other Pacific Islander	*	32	7	21	1	*
Asian	138	115	52	45	27	29
Two or More Races	122	75	63	32	21	23
Hispanic or Latino	253	203	159	121	114	81
EDUCATION						
< High School	256	221	145	128	115	95
High School Graduate	377	323	186	148	123	106
Some College	459	367	207	152	137	108
College Graduate	542	401	218	153	120	91
CURRENT EMPLOYMENT						
Full-Time	622	503	278	210	158	146
Part-Time	335	231	160	94	111	66
Unemployed	166	139	99	64	70	45
Other ¹	437	344	204	155	138	114

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.52D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.37	0.30	0.19	0.14	0.12	0.10
GENDER						
Male	0.53	0.44	0.26	0.21	0.18	0.15
Female	0.51	0.39	0.26	0.17	0.16	0.13
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.40	0.33	0.19	0.15	0.12	0.11
White	0.47	0.39	0.22	0.17	0.14	0.12
Black or African American	0.87	0.75	0.46	0.41	0.39	0.37
American Indian or Alaska Native	4.06	3.29	2.75	2.52	0.49	1.16
Native Hawaiian or Other Pacific Islander	*	4.10	1.18	2.78	0.22	*
Asian	1.28	1.06	0.51	0.44	0.27	0.28
Two or More Races	3.81	2.42	2.18	1.08	0.77	0.79
Hispanic or Latino	0.89	0.69	0.56	0.41	0.40	0.27
EDUCATION						
< High School	0.91	0.77	0.53	0.45	0.42	0.34
High School Graduate	0.59	0.52	0.31	0.25	0.21	0.18
Some College	0.77	0.61	0.39	0.28	0.26	0.20
College Graduate	0.71	0.50	0.31	0.22	0.18	0.13
CURRENT EMPLOYMENT						
Full-Time	0.50	0.40	0.25	0.18	0.15	0.13
Part-Time	1.16	0.89	0.65	0.39	0.46	0.28
Unemployed	1.93	1.71	1.25	0.89	0.91	0.63
Other ¹	0.62	0.49	0.31	0.23	0.21	0.17

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.52P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5229		0.2962		0.9959	
GENDER						
Male	0.0391		0.5937		0.8348	
Female	0.2605		0.0417		0.8347	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2554		0.5057		0.4911	
White	0.3479		0.2514		0.7917	
Black or African American	0.2887		0.3047		0.2493	
American Indian or Alaska Native	0.9115		0.7042		0.0164	
Native Hawaiian or Other Pacific Islander	*		0.1886		*	
Asian	0.3783		0.4134		0.5427	
Two or More Races	0.5668		0.4799		0.2073	
Hispanic or Latino	0.2512		0.2662		0.1556	
EDUCATION						
< High School	0.1038		0.2725		0.9635	
High School Graduate	0.1065		0.2008		0.3942	
Some College	0.7081		0.1529		0.6511	
College Graduate	0.3956		0.7816		0.6936	
CURRENT EMPLOYMENT						
Full-Time	0.0977		0.7581		0.2251	
Part-Time	0.0096		0.0712		0.3247	
Unemployed	0.8495		0.1604		0.3552	
Other ¹	0.4464		0.7382		0.8718	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.53C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	778	657	415	314	255	217
GENDER						
Male	525	457	273	225	179	156
Female	551	443	283	206	179	145
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	724	612	361	283	226	199
White	804	657	344	262	207	174
Black or African American	225	211	111	111	95	92
American Indian or Alaska Native	44	48	29	31	8	14
Native Hawaiian or Other Pacific Islander	*	35	9	24	2	23
Asian	143	120	55	54	29	34
Two or More Races	124	81	66	36	22	26
Hispanic or Latino	270	213	169	129	119	85
EDUCATION						
< High School	272	233	156	136	120	98
High School Graduate	390	340	195	166	127	116
Some College	470	392	225	168	142	116
College Graduate	552	415	221	159	122	95
CURRENT EMPLOYMENT						
Full-Time	640	526	289	225	163	156
Part-Time	351	253	174	112	117	75
Unemployed	176	152	110	75	76	51
Other ¹	442	350	210	162	141	118

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.53D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.33	0.27	0.17	0.13	0.11	0.09
GENDER						
Male	0.46	0.39	0.24	0.19	0.16	0.13
Female	0.45	0.36	0.23	0.17	0.15	0.12
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.36	0.30	0.18	0.14	0.11	0.10
White	0.43	0.36	0.21	0.16	0.13	0.11
Black or African American	0.73	0.64	0.39	0.37	0.34	0.31
American Indian or Alaska Native	3.36	2.98	2.25	2.18	0.59	1.06
Native Hawaiian or Other Pacific Islander	*	3.65	1.19	2.60	0.20	2.46
Asian	1.12	0.92	0.46	0.43	0.24	0.27
Two or More Races	3.07	2.00	1.80	0.95	0.64	0.69
Hispanic or Latino	0.76	0.58	0.47	0.35	0.33	0.23
EDUCATION						
< High School	0.79	0.69	0.47	0.40	0.36	0.30
High School Graduate	0.51	0.44	0.27	0.23	0.18	0.17
Some College	0.63	0.53	0.34	0.25	0.22	0.18
College Graduate	0.67	0.47	0.29	0.21	0.17	0.13
CURRENT EMPLOYMENT						
Full-Time	0.45	0.36	0.23	0.17	0.13	0.12
Part-Time	0.87	0.67	0.50	0.32	0.34	0.22
Unemployed	1.31	1.20	0.90	0.67	0.63	0.46
Other ¹	0.56	0.45	0.28	0.22	0.19	0.16

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.53P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.6047		0.2110		0.7175	
GENDER						
Male	0.0810		0.9129		0.8932	
Female	0.3310		0.0493		0.6981	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2840		0.3756		0.8428	
White	0.4463		0.1053		0.4842	
Black or African American	0.1601		0.1238		0.4223	
American Indian or Alaska Native	0.7120		0.8914		0.0324	
Native Hawaiian or Other Pacific Islander	*		0.0652		0.0119	
Asian	0.4078		0.4141		0.5622	
Two or More Races	0.5360		0.4200		0.2962	
Hispanic or Latino	0.1905		0.2475		0.2138	
EDUCATION						
< High School	0.1959		0.4192		0.6450	
High School Graduate	0.2029		0.1321		0.2870	
Some College	0.7229		0.1990		0.6371	
College Graduate	0.5057		0.7219		0.7540	
CURRENT EMPLOYMENT						
Full-Time	0.0827		0.5972		0.2064	
Part-Time	0.0087		0.0491		0.1894	
Unemployed	0.8387		0.0698		0.1564	
Other ¹	0.5052		0.5872		0.7514	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.54C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	640	553	343	253	206	174
AGE						
12-17	58	63	46	53	29	33
18-25	133	146	97	90	59	57
26 or Older	616	520	315	228	196	162
GENDER						
Male	434	401	227	184	146	133
Female	434	346	241	173	147	115
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	594	511	300	230	193	159
White	549	473	263	205	169	128
Black or African American	174	157	102	95	79	78
American Indian or Alaska Native	36	43	24	29	5	7
Native Hawaiian or Other Pacific Islander	47	32	10	24	5	23
Asian	106	91	53	38	25	23
Two or More Races	88	67	66	33	22	23
Hispanic or Latino	243	189	142	113	90	72

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.54D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.24	0.21	0.13	0.10	0.08	0.07
AGE						
12-17	0.23	0.25	0.18	0.21	0.11	0.13
18-25	0.38	0.42	0.28	0.26	0.17	0.16
26 or Older	0.30	0.25	0.16	0.11	0.10	0.08
GENDER						
Male	0.34	0.31	0.18	0.14	0.11	0.10
Female	0.32	0.25	0.18	0.13	0.11	0.08
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.27	0.23	0.14	0.10	0.09	0.07
White	0.32	0.28	0.15	0.12	0.10	0.08
Black or African American	0.56	0.50	0.33	0.30	0.25	0.25
American Indian or Alaska Native	2.41	2.41	1.65	1.85	0.38	0.48
Native Hawaiian or Other Pacific Islander	4.76	2.92	1.14	2.29	0.61	2.20
Asian	0.79	0.64	0.39	0.28	0.19	0.16
Two or More Races	1.95	1.37	1.47	0.71	0.53	0.51
Hispanic or Latino	0.59	0.44	0.35	0.27	0.22	0.17

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.54P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.7945		0.0451		0.3709	
AGE						
12-17	0.9936		0.9004		0.3296	
18-25	0.2409		0.0124		0.0545	
26 or Older	0.5751		0.1846		0.6333	
GENDER						
Male	0.0367		0.2940		0.7867	
Female	0.0681		0.0649		0.3135	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3514		0.0861		0.4485	
White	0.6126		0.0211		0.0910	
Black or African American	0.0954		0.0799		0.3308	
American Indian or Alaska Native	0.7397		0.5974		0.5026	
Native Hawaiian or Other Pacific Islander	0.3661		0.2549		0.0724	
Asian	0.4954		0.9127		0.7914	
Two or More Races	0.6396		0.0885		0.7943	
Hispanic or Latino	0.1124		0.2570		0.6446	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.55C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	58	63	46	53	29	33
GENDER						
Male	43	42	34	34	20	22
Female	39	44	30	38	19	25
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	50	52	40	44	25	27
White	44	47	35	38	20	22
Black or African American	25	26	18	20	14	14
American Indian or Alaska Native	6	6	6	6	2	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	9	9	7	6	2	4
Two or More Races	13	12	8	8	4	5
Hispanic or Latino	31	34	22	28	14	18
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	33	33	26	26	14	14
Female, White, Not Hispanic	28	33	23	29	12	17
Male, Black, Not Hispanic	18	16	13	11	11	8
Female, Black, Not Hispanic	17	20	12	16	9	11
Male, Hispanic	21	25	17	19	9	13
Female, Hispanic	23	25	16	21	11	13

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.55D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.23	0.25	0.18	0.21	0.11	0.13
GENDER						
Male	0.34	0.33	0.27	0.27	0.16	0.17
Female	0.32	0.36	0.25	0.31	0.15	0.20
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.26	0.27	0.21	0.23	0.13	0.14
White	0.29	0.33	0.24	0.27	0.14	0.16
Black or African American	0.68	0.69	0.48	0.54	0.39	0.39
American Indian or Alaska Native	3.75	3.16	3.51	2.91	1.14	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	0.70	0.69	0.52	0.50	0.17	0.31
Two or More Races	1.63	1.38	1.01	0.93	0.49	0.64
Hispanic or Latino	0.56	0.61	0.40	0.49	0.25	0.32
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.44	0.46	0.36	0.37	0.20	0.21
Female, White, Not Hispanic	0.39	0.47	0.33	0.42	0.18	0.26
Male, Black, Not Hispanic	0.96	0.84	0.71	0.62	0.60	0.47
Female, Black, Not Hispanic	0.93	1.08	0.69	0.86	0.50	0.62
Male, Hispanic	0.73	0.83	0.57	0.66	0.31	0.44
Female, Hispanic	0.80	0.85	0.56	0.72	0.39	0.47

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 1.55P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9936		0.9004		0.3296	
GENDER						
Male	0.4647		0.1775		0.5876	
Female	0.4613		0.1270		0.4038	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6308		0.6094		0.5144	
White	0.4092		0.3515		0.7092	
Black or African American	0.1388		0.1055		0.7414	
American Indian or Alaska Native	0.6906		0.6341		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.8414		0.8957		0.2141	
Two or More Races	0.2722		0.3412		0.8909	
Hispanic or Latino	0.4651		0.2696		0.4110	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.4079		0.0950		0.9757	
Female, White, Not Hispanic	0.7461		0.7061		0.5847	
Male, Black, Not Hispanic	0.8204		0.7531		0.8398	
Female, Black, Not Hispanic	0.0678		0.0595		0.5166	
Male, Hispanic	0.6227		0.6026		0.3821	
Female, Hispanic	0.5729		0.3011		0.7589	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.56C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	133	146	97	90	59	57
GENDER						
Male	100	103	74	70	46	45
Female	84	100	63	58	38	36
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	118	130	86	83	52	51
White	132	137	89	79	48	45
Black or African American	48	53	30	35	22	20
American Indian or Alaska Native	10	11	6	5	5	3
Native Hawaiian or Other Pacific Islander	9	*	6	*	1	*
Asian	22	27	14	18	8	10
Two or More Races	24	21	18	15	8	11
Hispanic or Latino	63	65	43	43	27	30
EDUCATION						
< High School	63	68	46	41	28	25
High School Graduate	89	93	62	59	37	37
Some College	91	101	61	60	34	35
College Graduate	57	64	37	32	18	17
CURRENT EMPLOYMENT						
Full-Time	104	103	68	62	39	43
Part-Time	83	85	58	49	34	24
Unemployed	56	52	40	34	21	19
Other ¹	59	70	39	43	23	28

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.56D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.38	0.42	0.28	0.26	0.17	0.16
GENDER						
Male	0.57	0.59	0.42	0.40	0.26	0.25
Female	0.49	0.57	0.36	0.33	0.22	0.21
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.43	0.47	0.31	0.30	0.19	0.18
White	0.53	0.58	0.40	0.38	0.24	0.22
Black or African American	0.80	0.93	0.55	0.66	0.42	0.39
American Indian or Alaska Native	3.49	4.42	2.40	2.36	2.07	1.28
Native Hawaiian or Other Pacific Islander	4.24	*	3.04	*	0.41	*
Asian	1.19	1.17	0.76	0.83	0.45	0.48
Two or More Races	2.81	2.35	2.33	1.77	1.04	1.36
Hispanic or Latino	0.87	0.88	0.59	0.58	0.38	0.40
EDUCATION						
< High School	0.98	1.14	0.77	0.76	0.49	0.50
High School Graduate	0.64	0.69	0.49	0.46	0.30	0.30
Some College	0.64	0.68	0.45	0.46	0.27	0.27
College Graduate	0.90	0.94	0.65	0.53	0.36	0.31
CURRENT EMPLOYMENT						
Full-Time	0.64	0.62	0.46	0.41	0.28	0.29
Part-Time	0.73	0.76	0.53	0.48	0.35	0.25
Unemployed	1.17	1.19	0.88	0.83	0.50	0.48
Other ¹	0.72	0.83	0.51	0.54	0.30	0.36

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 1.56P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2409		0.0124		0.0545	
GENDER						
Male	0.2796		0.0226		0.1024	
Female	0.5585		0.2412		0.3000	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4034		0.0320		0.0244	
White	0.2411		0.0133		0.0494	
Black or African American	0.2283		0.2856		0.1404	
American Indian or Alaska Native	0.6406		0.5861		0.4629	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7476		0.7996		0.5640	
Two or More Races	0.1632		0.1413		0.8881	
Hispanic or Latino	0.3645		0.2141		0.8605	
EDUCATION						
< High School	0.9720		0.8970		0.5398	
High School Graduate	0.0712		0.0114		0.0910	
Some College	0.9564		0.7462		0.8853	
College Graduate	0.9493		0.0556		0.2582	
CURRENT EMPLOYMENT						
Full-Time	0.8794		0.3066		0.8072	
Part-Time	0.2410		0.1262		0.0158	
Unemployed	0.3451		0.0365		0.2510	
Other ¹	0.5426		0.5056		0.6605	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.57C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	616	520	315	228	196	162
GENDER						
Male	419	380	209	167	139	124
Female	425	323	226	155	141	103
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	569	481	275	206	182	147
White	581	477	251	185	165	118
Black or African American	176	163	98	92	74	78
American Indian or Alaska Native	32	38	22	27	2	4
Native Hawaiian or Other Pacific Islander	*	27	4	21	1	*
Asian	102	87	50	33	24	20
Two or More Races	84	62	63	28	21	19
Hispanic or Latino	226	177	133	100	84	65
EDUCATION						
< High School	201	190	114	112	83	84
High School Graduate	322	250	165	125	104	82
Some College	350	288	172	122	117	85
College Graduate	393	283	166	114	100	67
CURRENT EMPLOYMENT						
Full-Time	494	410	234	175	129	121
Part-Time	253	168	133	72	88	51
Unemployed	134	115	83	56	66	41
Other ¹	321	261	158	122	112	83

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.57D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.30	0.25	0.16	0.11	0.10	0.08
GENDER						
Male	0.43	0.39	0.22	0.17	0.14	0.13
Female	0.40	0.30	0.21	0.14	0.13	0.10
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.33	0.27	0.16	0.12	0.10	0.08
White	0.39	0.33	0.18	0.13	0.12	0.09
Black or African American	0.74	0.64	0.42	0.38	0.32	0.33
American Indian or Alaska Native	3.19	2.98	2.16	2.39	0.18	0.35
Native Hawaiian or Other Pacific Islander	*	3.52	0.78	2.74	0.22	*
Asian	0.98	0.80	0.49	0.32	0.23	0.19
Two or More Races	2.91	2.04	2.16	0.96	0.76	0.66
Hispanic or Latino	0.79	0.60	0.47	0.34	0.30	0.22
EDUCATION						
< High School	0.73	0.66	0.42	0.40	0.31	0.31
High School Graduate	0.51	0.42	0.28	0.21	0.18	0.14
Some College	0.63	0.50	0.33	0.23	0.23	0.16
College Graduate	0.55	0.39	0.24	0.16	0.15	0.10
CURRENT EMPLOYMENT						
Full-Time	0.42	0.35	0.22	0.15	0.12	0.11
Part-Time	0.95	0.68	0.54	0.30	0.36	0.22
Unemployed	1.63	1.48	1.06	0.79	0.85	0.58
Other ¹	0.47	0.39	0.24	0.18	0.17	0.13

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.57P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5751		0.1846		0.6333	
GENDER						
Male	0.0117		0.9955		0.8455	
Female	0.0673		0.0612		0.3669	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2229		0.3063		0.8470	
White	0.3912		0.1383		0.2279	
Black or African American	0.2263		0.2224		0.1688	
American Indian or Alaska Native	0.7111		0.7727		0.2308	
Native Hawaiian or Other Pacific Islander	*		0.0985		*	
Asian	0.4356		0.9738		0.7406	
Two or More Races	0.9594		0.2329		0.7138	
Hispanic or Latino	0.1280		0.3189		0.4661	
EDUCATION						
< High School	0.0106		0.0711		0.5192	
High School Graduate	0.3666		0.1447		0.2594	
Some College	0.7437		0.1630		0.8402	
College Graduate	0.2994		0.2619		0.8327	
CURRENT EMPLOYMENT						
Full-Time	0.0786		0.8305		0.5666	
Part-Time	0.0133		0.2254		0.6545	
Unemployed	0.5291		0.2103		0.3508	
Other ¹	0.9108		0.5187		0.5267	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.58C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	636	554	339	246	205	170
GENDER						
Male	431	400	225	181	145	131
Female	431	345	238	169	145	111
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	590	510	298	223	191	155
White	602	509	272	204	171	127
Black or African American	184	176	102	100	79	80
American Indian or Alaska Native	35	41	23	27	5	5
Native Hawaiian or Other Pacific Islander	*	31	8	23	1	22
Asian	106	90	52	38	25	22
Two or More Races	86	66	65	31	22	22
Hispanic or Latino	239	189	141	109	88	69
EDUCATION						
< High School	215	203	125	119	88	88
High School Graduate	334	270	173	140	109	89
Some College	363	308	187	134	122	90
College Graduate	396	293	169	116	101	69
CURRENT EMPLOYMENT						
Full-Time	510	428	244	186	132	130
Part-Time	266	190	144	87	94	57
Unemployed	145	129	94	67	70	45
Other ¹	325	271	164	131	114	88

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.58D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.27	0.23	0.14	0.10	0.09	0.07
GENDER						
Male	0.38	0.35	0.20	0.16	0.13	0.11
Female	0.35	0.28	0.19	0.14	0.12	0.09
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.29	0.25	0.15	0.11	0.09	0.08
White	0.35	0.30	0.17	0.13	0.11	0.08
Black or African American	0.62	0.55	0.36	0.33	0.28	0.27
American Indian or Alaska Native	2.70	2.66	1.80	2.02	0.42	0.37
Native Hawaiian or Other Pacific Islander	*	3.21	0.99	2.46	0.19	2.34
Asian	0.86	0.70	0.43	0.30	0.21	0.18
Two or More Races	2.36	1.66	1.78	0.83	0.63	0.59
Hispanic or Latino	0.67	0.51	0.39	0.30	0.25	0.19
EDUCATION						
< High School	0.64	0.60	0.38	0.36	0.27	0.27
High School Graduate	0.44	0.37	0.24	0.20	0.16	0.13
Some College	0.53	0.43	0.29	0.20	0.19	0.14
College Graduate	0.51	0.37	0.23	0.15	0.14	0.09
CURRENT EMPLOYMENT						
Full-Time	0.39	0.31	0.20	0.14	0.11	0.10
Part-Time	0.71	0.54	0.42	0.26	0.28	0.17
Unemployed	1.13	1.06	0.78	0.59	0.58	0.41
Other ¹	0.43	0.36	0.22	0.17	0.16	0.12

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.58P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.8108		0.0403		0.2866	
GENDER						
Male	0.0317		0.4020		0.7197	
Female	0.0545		0.0339		0.2430	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3379		0.0968		0.3863	
White	0.5786		0.0273		0.0774	
Black or African American	0.1391		0.1378		0.3531	
American Indian or Alaska Native	0.6847		0.6561		0.9589	
Native Hawaiian or Other Pacific Islander	*		0.0782		0.0261	
Asian	0.4907		0.9227		0.9187	
Two or More Races	0.7591		0.1146		0.7694	
Hispanic or Latino	0.0849		0.1699		0.5224	
EDUCATION						
< High School	0.0307		0.1352		0.7348	
High School Graduate	0.6960		0.0236		0.0984	
Some College	0.7531		0.1548		0.8047	
College Graduate	0.3271		0.1431		0.6831	
CURRENT EMPLOYMENT						
Full-Time	0.0856		0.6523		0.6138	
Part-Time	0.0067		0.0849		0.1879	
Unemployed	0.8230		0.0479		0.2068	
Other ¹	0.9625		0.4083		0.4635	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.59C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,015	873	677	595	509	476
GEOGRAPHIC DIVISION						
Northeast	463	362	258	277	185	222
New England	225	183	141	151	100	122
Middle Atlantic	405	311	219	236	155	190
Midwest	395	398	247	245	190	183
East North Central	329	310	203	206	153	155
West North Central	225	223	137	118	106	86
South	654	481	412	322	306	259
South Atlantic	523	341	355	243	262	202
East South Central	211	187	119	112	101	84
West South Central	312	276	173	176	137	126
West	535	470	402	339	311	269
Mountain	248	186	170	144	128	119
Pacific	477	430	361	305	282	239
COUNTY TYPE						
Large Metro	1,317	1,066	589	536	405	416
Small Metro	961	776	446	354	329	272
250K – 1 Mil. Pop.	831	663	338	305	243	237
< 250K Pop.	570	466	300	199	228	140
Nonmetro	623	575	279	261	209	178
Urbanized	458	425	223	179	168	122
Less Urbanized	454	435	165	199	118	134
Completely Rural	196	157	83	41	66	30

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.59D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.33	0.26	0.22	0.19	0.18
GEOGRAPHIC DIVISION						
Northeast	0.98	0.76	0.54	0.58	0.39	0.47
New England	1.80	1.46	1.13	1.20	0.80	0.97
Middle Atlantic	1.16	0.89	0.63	0.67	0.44	0.54
Midwest	0.70	0.71	0.44	0.43	0.34	0.32
East North Central	0.84	0.79	0.52	0.53	0.39	0.40
West North Central	1.31	1.28	0.79	0.68	0.62	0.49
South	0.67	0.49	0.42	0.33	0.31	0.26
South Atlantic	1.02	0.65	0.69	0.47	0.51	0.39
East South Central	1.36	1.20	0.77	0.72	0.65	0.54
West South Central	1.02	0.89	0.56	0.57	0.45	0.41
West	0.87	0.76	0.66	0.54	0.51	0.43
Mountain	1.33	0.98	0.91	0.76	0.68	0.63
Pacific	1.12	1.00	0.85	0.71	0.66	0.55
COUNTY TYPE						
Large Metro	0.58	0.47	0.36	0.33	0.27	0.26
Small Metro	0.67	0.55	0.49	0.38	0.38	0.31
250K – 1 Mil. Pop.	0.87	0.68	0.58	0.47	0.45	0.39
< 250K Pop.	1.07	1.00	0.92	0.65	0.75	0.46
Nonmetro	0.90	0.68	0.58	0.52	0.45	0.37
Urbanized	1.35	1.16	0.95	0.85	0.77	0.61
Less Urbanized	1.37	1.03	0.78	0.76	0.57	0.54
Completely Rural	2.38	1.87	1.57	0.82	1.29	0.69

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.59P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2511		0.0189		0.0022	
GEOGRAPHIC DIVISION						
Northeast	0.0366		0.0022		0.0008	
New England	0.1034		0.0495		0.0346	
Middle Atlantic	0.1395		0.0185		0.0090	
Midwest	0.1133		0.3477		0.3020	
East North Central	0.0390		0.6355		0.4788	
West North Central	0.8901		0.2918		0.3853	
South	0.5613		0.2131		0.1166	
South Atlantic	0.9815		0.9100		0.2391	
East South Central	0.4719		0.3331		0.9158	
West South Central	0.5412		0.0715		0.2253	
West	0.1405		0.9615		0.5285	
Mountain	0.3941		0.7652		0.7704	
Pacific	0.2212		0.9300		0.5734	
COUNTY TYPE						
Large Metro	0.3754		0.0262		0.0083	
Small Metro	0.8581		0.9559		0.3621	
250K – 1 Mil. Pop.	0.9786		0.5102		0.1807	
< 250K Pop.	0.7121		0.4166		0.7562	
Nonmetro	0.1782		0.1422		0.1819	
Urbanized	0.0565		0.3445		0.4693	
Less Urbanized	0.6839		0.0860		0.0539	
Completely Rural	0.5990		0.3306		0.2039	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.60C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	97	108	86	95	62	73
GEOGRAPHIC DIVISION						
Northeast	35	38	32	34	23	25
New England	16	17	14	16	10	11
Middle Atlantic	31	33	28	30	21	23
Midwest	36	46	34	40	24	29
East North Central	29	39	28	34	18	27
West North Central	20	25	18	21	15	12
South	61	65	49	57	35	44
South Atlantic	48	47	37	41	27	31
East South Central	18	21	16	18	11	13
West South Central	33	41	27	36	19	28
West	56	59	53	52	39	43
Mountain	28	27	22	26	20	19
Pacific	48	53	47	46	33	38
COUNTY TYPE						
Large Metro	89	103	75	86	51	63
Small Metro	73	67	59	55	38	36
250K – 1 Mil. Pop.	61	52	49	44	32	31
< 250K Pop.	38	40	31	31	19	20
Nonmetro	47	44	32	39	23	25
Urbanized	28	30	23	24	15	17
Less Urbanized	39	33	24	30	17	19
Completely Rural	12	11	10	8	7	6

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.60D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.43	0.35	0.38	0.25	0.30
GEOGRAPHIC DIVISION						
Northeast	0.84	0.91	0.75	0.83	0.56	0.61
New England	1.50	1.54	1.28	1.45	0.95	0.98
Middle Atlantic	1.00	1.09	0.90	0.99	0.67	0.75
Midwest	0.67	0.87	0.63	0.74	0.45	0.55
East North Central	0.78	1.03	0.76	0.92	0.47	0.72
West North Central	1.23	1.51	1.08	1.25	0.91	0.76
South	0.65	0.69	0.53	0.61	0.38	0.47
South Atlantic	1.04	1.00	0.80	0.87	0.57	0.65
East South Central	1.23	1.42	1.11	1.25	0.78	0.90
West South Central	1.01	1.25	0.85	1.09	0.59	0.86
West	0.94	0.99	0.88	0.88	0.65	0.72
Mountain	1.50	1.45	1.19	1.35	1.05	1.02
Pacific	1.18	1.30	1.16	1.14	0.81	0.94
COUNTY TYPE						
Large Metro	0.52	0.60	0.48	0.53	0.35	0.42
Small Metro	0.71	0.74	0.65	0.64	0.44	0.47
250K – 1 Mil. Pop.	0.86	0.83	0.79	0.74	0.56	0.58
< 250K Pop.	1.22	1.47	1.10	1.16	0.70	0.83
Nonmetro	0.98	0.95	0.73	0.87	0.56	0.62
Urbanized	1.38	1.50	1.16	1.27	0.79	1.03
Less Urbanized	1.63	1.33	1.11	1.32	0.91	0.87
Completely Rural	2.23	2.31	2.00	1.88	1.49	1.40

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.60P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9952		0.6826		0.1397	
GEOGRAPHIC DIVISION						
Northeast	0.7220		0.6595		0.7344	
New England	0.5933		0.5226		0.2512	
Middle Atlantic	0.4982		0.4104		0.8651	
Midwest	0.7829		0.6846		0.8286	
East North Central	0.9190		0.9484		0.5517	
West North Central	0.7318		0.5056		0.1623	
South	0.8453		0.4387		0.0627	
South Atlantic	0.3420		0.9910		0.5945	
East South Central	0.4515		0.1741		0.9531	
West South Central	0.0503		0.0386		0.0150	
West	0.8284		0.6228		0.3227	
Mountain	0.4091		0.6874		0.5600	
Pacific	0.4843		0.4473		0.1477	
COUNTY TYPE						
Large Metro	0.8107		0.7985		0.3466	
Small Metro	0.4736		0.5104		0.3919	
250K – 1 Mil. Pop.	0.2349		0.2371		0.2407	
< 250K Pop.	0.6806		0.5717		0.7577	
Nonmetro	0.6714		0.4241		0.3603	
Urbanized	0.6861		0.5154		0.3725	
Less Urbanized	0.2323		0.7207		0.7757	
Completely Rural	0.5397		0.7105		0.6388	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.61C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	168	190	166	185	143	160
GEOGRAPHIC DIVISION						
Northeast	71	74	67	72	52	58
New England	32	36	34	34	25	29
Middle Atlantic	59	66	53	66	45	51
Midwest	61	80	65	83	55	67
East North Central	51	67	53	71	44	58
West North Central	33	38	37	40	33	30
South	105	111	98	109	82	94
South Atlantic	80	73	77	73	65	63
East South Central	38	47	35	42	27	35
West South Central	53	68	51	64	41	55
West	96	102	93	98	85	85
Mountain	39	42	41	44	38	38
Pacific	89	93	84	87	76	76
COUNTY TYPE						
Large Metro	261	252	196	195	139	154
Small Metro	188	183	134	148	90	106
250K – 1 Mil. Pop.	142	145	101	126	67	88
< 250K Pop.	132	111	93	79	63	61
Nonmetro	118	136	87	102	60	70
Urbanized	93	105	66	81	45	53
Less Urbanized	81	86	59	60	41	47
Completely Rural	32	25	20	19	12	11

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.61D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.48	0.54	0.48	0.53	0.41	0.46
GEOGRAPHIC DIVISION						
Northeast	1.15	1.20	1.09	1.16	0.85	0.94
New England	1.99	2.20	2.10	2.07	1.54	1.79
Middle Atlantic	1.31	1.47	1.17	1.46	0.99	1.12
Midwest	0.82	1.08	0.88	1.12	0.75	0.90
East North Central	1.00	1.31	1.04	1.39	0.86	1.14
West North Central	1.43	1.62	1.59	1.72	1.41	1.29
South	0.82	0.86	0.76	0.84	0.64	0.73
South Atlantic	1.22	1.11	1.17	1.11	1.00	0.96
East South Central	1.88	2.27	1.70	2.06	1.33	1.71
West South Central	1.25	1.58	1.19	1.48	0.97	1.28
West	1.15	1.21	1.11	1.16	1.01	1.01
Mountain	1.52	1.64	1.62	1.72	1.51	1.49
Pacific	1.52	1.59	1.43	1.49	1.30	1.30
COUNTY TYPE						
Large Metro	0.71	0.79	0.69	0.74	0.61	0.64
Small Metro	0.82	0.90	0.79	0.97	0.66	0.76
250K – 1 Mil. Pop.	1.02	1.14	0.95	1.25	0.79	0.97
< 250K Pop.	1.37	1.49	1.36	1.49	1.10	1.28
Nonmetro	1.08	1.28	1.06	1.33	0.84	1.05
Urbanized	1.57	2.05	1.44	2.24	1.20	1.73
Less Urbanized	1.77	1.57	1.79	1.56	1.36	1.42
Completely Rural	3.92	3.86	2.78	3.38	1.95	2.44

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.61P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2296		0.7006		0.4747	
GEOGRAPHIC DIVISION						
Northeast	0.8358		0.2816		0.2153	
New England	0.8086		0.6692		0.5606	
Middle Atlantic	0.7221		0.1412		0.2812	
Midwest	0.1649		0.3464		0.9545	
East North Central	0.1040		0.7238		0.5741	
West North Central	0.9531		0.2070		0.4165	
South	0.2170		0.3522		0.2321	
South Atlantic	0.3077		0.6834		0.2677	
East South Central	0.8781		0.8372		0.7348	
West South Central	0.4163		0.3123		0.3512	
West	0.7529		0.0706		0.4209	
Mountain	0.6084		0.8975		0.8187	
Pacific	0.9079		0.0508		0.3111	
COUNTY TYPE						
Large Metro	0.4289		0.8246		0.7639	
Small Metro	0.9250		0.7019		0.1801	
250K – 1 Mil. Pop.	0.8543		0.7257		0.1390	
< 250K Pop.	0.6818		0.8655		0.7932	
Nonmetro	0.1333		0.3880		0.4742	
Urbanized	0.0975		0.2510		0.8667	
Less Urbanized	0.7149		0.9147		0.4612	
Completely Rural	0.9286		0.8364		0.7487	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.62C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	978	818	622	515	472	415
GEOGRAPHIC DIVISION						
Northeast	451	343	233	251	172	204
New England	217	174	115	136	90	111
Middle Atlantic	396	295	206	215	146	176
Midwest	379	373	216	200	170	156
East North Central	319	290	176	172	136	133
West North Central	211	211	122	94	98	75
South	631	454	383	275	289	221
South Atlantic	502	330	331	211	247	175
East South Central	200	178	103	99	92	71
West South Central	299	254	153	155	129	110
West	505	435	367	294	287	231
Mountain	234	181	157	124	116	99
Pacific	448	394	328	265	261	205
COUNTY TYPE						
Large Metro	1,214	976	518	466	366	364
Small Metro	877	697	392	283	299	226
250K – 1 Mil. Pop.	751	603	296	243	221	195
< 250K Pop.	511	405	259	161	203	119
Nonmetro	569	498	247	208	185	144
Urbanized	419	366	197	134	144	96
Less Urbanized	406	371	135	161	103	107
Completely Rural	172	140	69	33	59	30

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.62D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.48	0.40	0.31	0.25	0.23	0.20
GEOGRAPHIC DIVISION						
Northeast	1.22	0.92	0.63	0.67	0.46	0.55
New England	2.23	1.77	1.17	1.38	0.92	1.13
Middle Atlantic	1.45	1.07	0.75	0.78	0.54	0.64
Midwest	0.87	0.85	0.50	0.46	0.39	0.36
East North Central	1.06	0.96	0.59	0.57	0.45	0.44
West North Central	1.58	1.58	0.92	0.70	0.73	0.56
South	0.84	0.59	0.51	0.36	0.38	0.29
South Atlantic	1.25	0.81	0.82	0.52	0.61	0.43
East South Central	1.67	1.48	0.86	0.82	0.77	0.58
West South Central	1.30	1.08	0.66	0.66	0.56	0.47
West	1.07	0.91	0.78	0.61	0.61	0.48
Mountain	1.64	1.24	1.10	0.85	0.82	0.68
Pacific	1.37	1.18	1.00	0.79	0.80	0.62
COUNTY TYPE						
Large Metro	0.72	0.57	0.43	0.38	0.32	0.30
Small Metro	0.83	0.66	0.58	0.42	0.46	0.34
250K – 1 Mil. Pop.	1.07	0.82	0.68	0.51	0.53	0.43
< 250K Pop.	1.36	1.20	1.13	0.75	0.93	0.56
Nonmetro	1.11	0.83	0.68	0.57	0.52	0.40
Urbanized	1.70	1.36	1.14	0.88	0.89	0.64
Less Urbanized	1.68	1.27	0.85	0.89	0.66	0.60
Completely Rural	2.80	2.20	1.73	0.95	1.50	0.88

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.62P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3587		0.0120		0.0029	
GEOGRAPHIC DIVISION						
Northeast	0.0324		0.0015		0.0008	
New England	0.0935		0.0207		0.0214	
Middle Atlantic	0.1316		0.0231		0.0130	
Midwest	0.1578		0.3893		0.2277	
East North Central	0.0670		0.6456		0.6121	
West North Central	0.8975		0.3619		0.1410	
South	0.7438		0.3094		0.2867	
South Atlantic	0.9487		0.9553		0.3621	
East South Central	0.4282		0.2031		0.8068	
West South Central	0.8397		0.2043		0.6235	
West	0.1159		0.6652		0.4002	
Mountain	0.4854		0.8363		0.7096	
Pacific	0.1584		0.5611		0.4460	
COUNTY TYPE						
Large Metro	0.4214		0.0073		0.0030	
Small Metro	0.6963		0.9084		0.6200	
250K – 1 Mil. Pop.	0.8820		0.6438		0.3818	
< 250K Pop.	0.6352		0.4212		0.7310	
Nonmetro	0.2501		0.2384		0.2842	
Urbanized	0.1061		0.6590		0.6009	
Less Urbanized	0.6118		0.0619		0.0647	
Completely Rural	0.5087		0.3087		0.1678	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.63C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	999	857	666	581	503	462
GEOGRAPHIC DIVISION						
Northeast	459	354	256	272	184	218
New England	224	181	140	150	101	122
Middle Atlantic	402	305	216	230	155	185
Midwest	391	396	239	239	189	178
East North Central	327	307	196	200	152	151
West North Central	223	219	135	114	107	85
South	646	469	412	315	306	251
South Atlantic	519	338	357	235	263	193
East South Central	206	183	116	111	99	81
West South Central	303	264	169	176	136	125
West	516	463	389	328	301	263
Mountain	240	186	169	144	127	119
Pacific	460	421	347	294	272	231
COUNTY TYPE						
Large Metro	1,292	1,041	573	518	399	404
Small Metro	933	758	432	340	322	264
250K – 1 Mil. Pop.	805	649	325	292	237	230
< 250K Pop.	558	453	294	191	225	137
Nonmetro	607	563	274	249	207	170
Urbanized	451	416	219	171	165	116
Less Urbanized	437	420	158	187	114	127
Completely Rural	190	152	79	37	63	29

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.63D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.42	0.36	0.28	0.24	0.21	0.19
GEOGRAPHIC DIVISION						
Northeast	1.06	0.82	0.59	0.62	0.43	0.50
New England	1.96	1.58	1.23	1.31	0.88	1.06
Middle Atlantic	1.26	0.95	0.68	0.72	0.49	0.58
Midwest	0.77	0.78	0.47	0.47	0.37	0.35
East North Central	0.93	0.87	0.56	0.56	0.43	0.43
West North Central	1.43	1.39	0.86	0.73	0.69	0.54
South	0.73	0.52	0.47	0.35	0.35	0.28
South Atlantic	1.11	0.71	0.76	0.49	0.56	0.41
East South Central	1.47	1.30	0.83	0.78	0.71	0.58
West South Central	1.11	0.95	0.62	0.63	0.50	0.45
West	0.93	0.82	0.70	0.58	0.54	0.47
Mountain	1.43	1.09	1.00	0.84	0.76	0.70
Pacific	1.19	1.08	0.90	0.75	0.71	0.59
COUNTY TYPE						
Large Metro	0.63	0.51	0.39	0.36	0.29	0.28
Small Metro	0.72	0.59	0.53	0.41	0.42	0.33
250K – 1 Mil. Pop.	0.94	0.73	0.62	0.51	0.48	0.42
< 250K Pop.	1.18	1.06	1.01	0.70	0.82	0.50
Nonmetro	0.98	0.74	0.63	0.55	0.49	0.39
Urbanized	1.49	1.24	1.03	0.90	0.85	0.63
Less Urbanized	1.48	1.13	0.84	0.82	0.61	0.58
Completely Rural	2.60	1.98	1.69	0.84	1.38	0.74

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.63P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2681		0.0193		0.0036	
GEOGRAPHIC DIVISION						
Northeast	0.0339		0.0015		0.0006	
New England	0.1134		0.0565		0.0265	
Middle Atlantic	0.1248		0.0122		0.0085	
Midwest	0.1105		0.3028		0.2803	
East North Central	0.0393		0.6160		0.5274	
West North Central	0.9092		0.2378		0.2866	
South	0.5850		0.2523		0.1851	
South Atlantic	0.9132		0.9066		0.2601	
East South Central	0.4223		0.2308		0.9208	
West South Central	0.7146		0.1481		0.4373	
West	0.1136		0.9038		0.6159	
Mountain	0.4371		0.8120		0.7014	
Pacific	0.1670		0.9850		0.7082	
COUNTY TYPE						
Large Metro	0.3627		0.0209		0.0106	
Small Metro	0.7171		0.9787		0.4152	
250K – 1 Mil. Pop.	0.8595		0.6073		0.2269	
< 250K Pop.	0.7032		0.4548		0.7765	
Nonmetro	0.1576		0.1645		0.2150	
Urbanized	0.0564		0.3775		0.5268	
Less Urbanized	0.5675		0.0862		0.0518	
Completely Rural	0.4932		0.2958		0.1702	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.64C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,017	872	577	538	442	424
GEOGRAPHIC DIVISION						
Northeast	449	364	231	233	164	192
New England	223	180	128	133	95	111
Middle Atlantic	391	314	193	193	133	160
Midwest	394	386	224	209	168	161
East North Central	318	311	183	173	135	135
West North Central	233	210	128	106	100	78
South	648	489	336	280	262	214
South Atlantic	515	349	292	215	230	159
East South Central	209	181	104	98	80	74
West South Central	311	268	145	148	109	112
West	537	477	353	306	278	249
Mountain	253	178	151	129	110	111
Pacific	479	444	315	276	255	223
COUNTY TYPE						
Large Metro	1,234	1,009	497	470	352	371
Small Metro	891	730	375	285	274	221
250K – 1 Mil. Pop.	778	627	296	249	206	192
< 250K Pop.	527	423	241	153	187	115
Nonmetro	571	538	218	233	192	160
Urbanized	414	392	180	155	162	107
Less Urbanized	413	404	120	179	98	122
Completely Rural	183	145	64	34	55	30

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.64D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.33	0.22	0.20	0.17	0.16
GEOGRAPHIC DIVISION						
Northeast	0.95	0.76	0.49	0.49	0.35	0.40
New England	1.79	1.43	1.03	1.06	0.76	0.89
Middle Atlantic	1.12	0.89	0.55	0.55	0.38	0.46
Midwest	0.70	0.68	0.40	0.37	0.30	0.28
East North Central	0.82	0.80	0.47	0.44	0.35	0.35
West North Central	1.35	1.21	0.74	0.61	0.58	0.45
South	0.66	0.49	0.34	0.28	0.27	0.22
South Atlantic	1.00	0.67	0.57	0.41	0.45	0.30
East South Central	1.35	1.16	0.67	0.63	0.52	0.48
West South Central	1.02	0.86	0.47	0.48	0.36	0.36
West	0.88	0.77	0.58	0.49	0.45	0.40
Mountain	1.35	0.94	0.81	0.68	0.59	0.59
Pacific	1.12	1.03	0.74	0.64	0.60	0.52
COUNTY TYPE						
Large Metro	0.57	0.47	0.31	0.29	0.23	0.24
Small Metro	0.67	0.56	0.43	0.32	0.33	0.26
250K – 1 Mil. Pop.	0.85	0.68	0.51	0.40	0.38	0.32
< 250K Pop.	1.16	0.98	0.77	0.56	0.64	0.41
Nonmetro	0.87	0.68	0.47	0.48	0.42	0.34
Urbanized	1.27	1.12	0.81	0.78	0.76	0.55
Less Urbanized	1.38	1.00	0.60	0.71	0.49	0.50
Completely Rural	2.27	1.85	1.24	0.72	1.12	0.70

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.64P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3327		0.0214		0.0004	
GEOGRAPHIC DIVISION						
Northeast	0.0456		0.0095		0.0005	
New England	0.1148		0.0797		0.0633	
Middle Atlantic	0.1605		0.0536		0.0033	
Midwest	0.1741		0.4841		0.1982	
East North Central	0.0968		0.8596		0.4314	
West North Central	0.9693		0.3067		0.2380	
South	0.7415		0.1473		0.0541	
South Atlantic	0.5579		0.6246		0.3341	
East South Central	0.5735		0.5899		0.4599	
West South Central	0.8851		0.0669		0.0529	
West	0.2278		0.8057		0.3944	
Mountain	0.2508		0.9604		0.6668	
Pacific	0.4329		0.8012		0.4594	
COUNTY TYPE						
Large Metro	0.2399		0.0473		0.0075	
Small Metro	0.5328		0.6700		0.2244	
250K – 1 Mil. Pop.	0.4117		0.7493		0.1016	
< 250K Pop.	0.8998		0.2323		0.8583	
Nonmetro	0.3879		0.0104		0.0413	
Urbanized	0.1780		0.1856		0.5562	
Less Urbanized	0.6800		0.0042		0.0027	
Completely Rural	0.3973		0.5115		0.5708	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.65C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	86	95	78	82	56	67
GEOGRAPHIC DIVISION						
Northeast	31	33	28	32	21	24
New England	16	16	14	16	10	10
Middle Atlantic	27	28	25	27	19	21
Midwest	34	41	31	37	22	25
East North Central	28	35	26	32	17	23
West North Central	18	21	16	18	13	10
South	51	56	44	47	31	38
South Atlantic	39	38	34	33	24	26
East South Central	15	18	14	15	10	12
West South Central	28	39	24	30	16	25
West	51	55	49	48	36	42
Mountain	26	25	21	24	19	17
Pacific	44	49	43	43	30	38
COUNTY TYPE						
Large Metro	71	84	62	72	45	58
Small Metro	58	56	53	45	33	33
250K – 1 Mil. Pop.	49	44	45	37	29	27
< 250K Pop.	31	34	28	26	16	19
Nonmetro	35	36	28	32	19	22
Urbanized	22	24	20	21	13	14
Less Urbanized	26	26	20	24	14	16
Completely Rural	10	9	8	8	6	5

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.65D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.35	0.38	0.31	0.33	0.23	0.27
GEOGRAPHIC DIVISION						
Northeast	0.75	0.80	0.68	0.76	0.51	0.57
New England	1.43	1.47	1.26	1.45	0.92	0.89
Middle Atlantic	0.87	0.92	0.80	0.88	0.61	0.69
Midwest	0.63	0.76	0.58	0.68	0.40	0.47
East North Central	0.74	0.93	0.69	0.86	0.44	0.62
West North Central	1.11	1.31	1.00	1.07	0.79	0.61
South	0.55	0.60	0.47	0.50	0.33	0.40
South Atlantic	0.83	0.81	0.73	0.72	0.52	0.55
East South Central	1.02	1.22	0.94	1.00	0.68	0.79
West South Central	0.86	1.18	0.74	0.93	0.49	0.76
West	0.86	0.92	0.82	0.82	0.60	0.71
Mountain	1.38	1.30	1.13	1.26	0.99	0.92
Pacific	1.07	1.22	1.06	1.05	0.74	0.94
COUNTY TYPE						
Large Metro	0.47	0.53	0.42	0.47	0.32	0.39
Small Metro	0.65	0.68	0.62	0.56	0.40	0.44
250K – 1 Mil. Pop.	0.80	0.76	0.76	0.66	0.51	0.52
< 250K Pop.	1.12	1.33	1.05	1.04	0.64	0.81
Nonmetro	0.77	0.83	0.65	0.75	0.48	0.54
Urbanized	1.20	1.38	1.07	1.19	0.73	0.85
Less Urbanized	1.18	1.13	0.98	1.09	0.75	0.78
Completely Rural	1.76	1.99	1.69	1.71	1.34	1.34

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.65P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9930		0.4690		0.4269	
GEOGRAPHIC DIVISION						
Northeast	0.2177		0.3909		0.5893	
New England	0.6606		0.4674		0.2955	
Middle Atlantic	0.0918		0.1568		0.9266	
Midwest	0.5357		0.1892		0.3035	
East North Central	0.4454		0.2920		0.6979	
West North Central	0.9630		0.4143		0.1584	
South	0.7146		0.6740		0.1804	
South Atlantic	0.6340		0.5546		0.6689	
East South Central	0.4454		0.0543		0.5984	
West South Central	0.1331		0.2991		0.0464	
West	0.3992		0.5018		0.3074	
Mountain	0.3479		0.3410		0.4323	
Pacific	0.1431		0.2091		0.1165	
COUNTY TYPE						
Large Metro	0.7385		0.5184		0.6500	
Small Metro	0.3329		0.7189		0.7051	
250K – 1 Mil. Pop.	0.2176		0.7111		0.8037	
< 250K Pop.	0.9517		0.2269		0.7582	
Nonmetro	0.5009		0.9079		0.4739	
Urbanized	0.9038		0.8095		0.6761	
Less Urbanized	0.2937		0.6020		0.7027	
Completely Rural	0.6971		0.7523		0.4839	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.66C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	170	197	163	186	135	157
GEOGRAPHIC DIVISION						
Northeast	72	72	65	71	52	55
New England	33	36	31	35	24	30
Middle Atlantic	60	63	54	64	44	47
Midwest	63	85	62	82	54	63
East North Central	54	72	52	68	43	55
West North Central	33	37	34	41	32	28
South	104	115	95	110	76	91
South Atlantic	80	75	76	74	62	60
East South Central	37	48	32	42	24	32
West South Central	54	68	47	65	36	55
West	98	108	92	99	80	86
Mountain	39	43	40	43	38	38
Pacific	90	98	84	89	71	76
COUNTY TYPE						
Large Metro	241	239	184	185	130	145
Small Metro	174	173	124	135	86	96
250K – 1 Mil. Pop.	130	138	92	114	62	78
< 250K Pop.	123	103	87	72	61	58
Nonmetro	111	128	78	96	58	66
Urbanized	84	99	57	78	43	51
Less Urbanized	80	80	55	55	39	43
Completely Rural	31	25	19	15	12	9

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.66D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.49	0.56	0.47	0.53	0.39	0.45
GEOGRAPHIC DIVISION						
Northeast	1.17	1.17	1.06	1.16	0.84	0.90
New England	2.04	2.21	1.88	2.12	1.50	1.83
Middle Atlantic	1.32	1.40	1.19	1.42	0.97	1.04
Midwest	0.86	1.15	0.84	1.10	0.73	0.85
East North Central	1.06	1.41	1.01	1.33	0.83	1.07
West North Central	1.44	1.60	1.49	1.75	1.40	1.22
South	0.81	0.89	0.74	0.85	0.59	0.71
South Atlantic	1.22	1.13	1.15	1.12	0.95	0.92
East South Central	1.82	2.32	1.57	2.06	1.16	1.58
West South Central	1.26	1.59	1.11	1.50	0.85	1.29
West	1.17	1.29	1.10	1.17	0.96	1.02
Mountain	1.54	1.69	1.57	1.67	1.49	1.48
Pacific	1.55	1.68	1.44	1.51	1.22	1.30
COUNTY TYPE						
Large Metro	0.72	0.82	0.68	0.74	0.58	0.64
Small Metro	0.83	0.93	0.80	0.94	0.66	0.72
250K – 1 Mil. Pop.	1.05	1.17	0.97	1.19	0.76	0.89
< 250K Pop.	1.43	1.52	1.36	1.43	1.15	1.25
Nonmetro	1.10	1.28	1.00	1.29	0.84	1.00
Urbanized	1.58	1.99	1.34	2.23	1.21	1.67
Less Urbanized	1.84	1.65	1.72	1.48	1.33	1.32
Completely Rural	3.90	3.85	2.65	2.92	1.89	2.09

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.66P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3159		0.6654		0.4081	
GEOGRAPHIC DIVISION						
Northeast	0.9775		0.2584		0.1531	
New England	0.5161		0.5761		0.8540	
Middle Atlantic	0.7511		0.1107		0.1197	
Midwest	0.5649		0.4121		0.9172	
East North Central	0.4963		0.8388		0.8377	
West North Central	0.9954		0.2073		0.5961	
South	0.1638		0.4911		0.1105	
South Atlantic	0.3544		0.8200		0.1800	
East South Central	0.8451		0.8691		0.8716	
West South Central	0.2343		0.2869		0.2270	
West	0.9362		0.1722		0.3196	
Mountain	0.7639		0.8464		0.7593	
Pacific	0.9651		0.1560		0.2014	
COUNTY TYPE						
Large Metro	0.5559		0.6143		0.6538	
Small Metro	0.9519		0.6448		0.0836	
250K – 1 Mil. Pop.	0.4727		0.5728		0.0746	
< 250K Pop.	0.3808		0.9893		0.5720	
Nonmetro	0.1009		0.1179		0.4056	
Urbanized	0.0146		0.1136		0.8368	
Less Urbanized	0.6767		0.6033		0.3636	
Completely Rural	0.7275		0.9393		0.7334	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.67C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	986	811	519	451	407	361
GEOGRAPHIC DIVISION						
Northeast	437	348	205	205	148	174
New England	217	172	104	119	84	100
Middle Atlantic	381	301	177	172	122	147
Midwest	380	357	192	166	147	137
East North Central	310	286	156	142	117	118
West North Central	218	200	113	83	91	66
South	631	458	307	231	246	177
South Atlantic	505	332	270	182	215	135
East South Central	198	172	90	81	73	63
West South Central	298	246	121	124	98	95
West	507	435	317	262	254	215
Mountain	241	167	136	108	100	91
Pacific	448	401	281	234	232	192
COUNTY TYPE						
Large Metro	1,147	936	432	400	316	322
Small Metro	821	655	320	225	243	179
250K – 1 Mil. Pop.	713	569	256	194	186	154
< 250K Pop.	473	373	198	123	158	96
Nonmetro	526	463	188	182	168	130
Urbanized	380	334	152	109	138	81
Less Urbanized	374	349	96	147	84	99
Completely Rural	162	128	53	28	49	29

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.67D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.49	0.39	0.26	0.22	0.20	0.18
GEOGRAPHIC DIVISION						
Northeast	1.18	0.93	0.55	0.55	0.40	0.47
New England	2.22	1.75	1.07	1.21	0.86	1.01
Middle Atlantic	1.40	1.10	0.65	0.62	0.45	0.53
Midwest	0.87	0.82	0.44	0.38	0.34	0.31
East North Central	1.03	0.94	0.52	0.47	0.39	0.39
West North Central	1.64	1.50	0.85	0.62	0.69	0.50
South	0.84	0.60	0.41	0.30	0.33	0.23
South Atlantic	1.26	0.81	0.67	0.44	0.53	0.33
East South Central	1.65	1.43	0.75	0.67	0.61	0.52
West South Central	1.29	1.04	0.53	0.53	0.42	0.40
West	1.08	0.91	0.67	0.55	0.54	0.45
Mountain	1.69	1.15	0.95	0.74	0.70	0.63
Pacific	1.37	1.21	0.86	0.70	0.71	0.58
COUNTY TYPE						
Large Metro	0.72	0.57	0.36	0.33	0.28	0.27
Small Metro	0.83	0.66	0.49	0.34	0.38	0.28
250K – 1 Mil. Pop.	1.04	0.81	0.59	0.42	0.44	0.34
< 250K Pop.	1.48	1.18	0.89	0.62	0.74	0.48
Nonmetro	1.10	0.80	0.54	0.52	0.48	0.37
Urbanized	1.60	1.27	0.94	0.75	0.87	0.56
Less Urbanized	1.73	1.23	0.63	0.82	0.55	0.56
Completely Rural	2.71	2.07	1.38	0.83	1.29	0.86

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.67P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.4483		0.0073		0.0004	
GEOGRAPHIC DIVISION						
Northeast	0.0316		0.0066		0.0005	
New England	0.0898		0.0342		0.0323	
Middle Atlantic	0.1323		0.0655		0.0065	
Midwest	0.1750		0.4366		0.0876	
East North Central	0.0977		0.7224		0.3693	
West North Central	0.9781		0.3807		0.0842	
South	0.9892		0.1297		0.1771	
South Atlantic	0.6416		0.5464		0.5634	
East South Central	0.5605		0.2996		0.3197	
West South Central	0.7592		0.1330		0.2197	
West	0.1576		0.5048		0.2397	
Mountain	0.2949		0.7501		0.5884	
Pacific	0.2902		0.5499		0.2927	
COUNTY TYPE						
Large Metro	0.2527		0.0080		0.0015	
Small Metro	0.3892		0.5739		0.4765	
250K – 1 Mil. Pop.	0.3418		0.9090		0.2311	
< 250K Pop.	0.9202		0.2604		0.6734	
Nonmetro	0.5309		0.0193		0.0622	
Urbanized	0.3549		0.4502		0.6702	
Less Urbanized	0.5491		0.0018		0.0028	
Completely Rural	0.2793		0.5412		0.5002	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.68C Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,008	860	565	528	437	414
GEOGRAPHIC DIVISION						
Northeast	447	358	229	228	162	190
New England	223	179	127	132	95	111
Middle Atlantic	388	309	191	189	132	157
Midwest	392	383	217	205	167	158
East North Central	319	307	177	169	133	133
West North Central	230	208	125	103	100	77
South	646	481	336	276	262	209
South Atlantic	517	347	293	210	230	153
East South Central	205	179	102	98	78	72
West South Central	303	258	142	148	107	113
West	519	469	339	296	270	243
Mountain	246	176	149	130	111	112
Pacific	461	434	300	264	246	215
COUNTY TYPE						
Large Metro	1,220	994	485	458	346	362
Small Metro	871	716	361	276	267	216
250K – 1 Mil. Pop.	760	616	286	239	201	186
< 250K Pop.	517	416	233	150	183	114
Nonmetro	562	528	214	223	191	154
Urbanized	410	386	176	148	160	102
Less Urbanized	403	393	115	170	95	118
Completely Rural	179	141	59	31	53	29

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.68D Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.42	0.36	0.24	0.22	0.18	0.17
GEOGRAPHIC DIVISION						
Northeast	1.03	0.82	0.53	0.52	0.38	0.44
New England	1.96	1.56	1.12	1.15	0.83	0.97
Middle Atlantic	1.22	0.96	0.60	0.59	0.42	0.49
Midwest	0.77	0.75	0.43	0.40	0.33	0.31
East North Central	0.90	0.87	0.50	0.48	0.38	0.38
West North Central	1.48	1.32	0.80	0.65	0.64	0.49
South	0.73	0.54	0.38	0.31	0.30	0.23
South Atlantic	1.11	0.73	0.63	0.44	0.49	0.32
East South Central	1.46	1.27	0.73	0.70	0.56	0.51
West South Central	1.11	0.93	0.52	0.53	0.39	0.40
West	0.94	0.83	0.61	0.53	0.49	0.43
Mountain	1.46	1.03	0.88	0.76	0.66	0.65
Pacific	1.20	1.11	0.78	0.68	0.64	0.55
COUNTY TYPE						
Large Metro	0.63	0.52	0.33	0.32	0.25	0.25
Small Metro	0.73	0.60	0.46	0.35	0.35	0.28
250K – 1 Mil. Pop.	0.92	0.73	0.55	0.43	0.41	0.34
< 250K Pop.	1.28	1.06	0.82	0.60	0.69	0.45
Nonmetro	0.97	0.73	0.51	0.51	0.46	0.36
Urbanized	1.41	1.18	0.88	0.82	0.83	0.57
Less Urbanized	1.52	1.10	0.64	0.77	0.52	0.54
Completely Rural	2.51	1.94	1.29	0.74	1.19	0.76

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.68P Marijuana Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3598		0.0141		0.0006	
GEOGRAPHIC DIVISION						
Northeast	0.0368		0.0060		0.0003	
New England	0.1258		0.0922		0.0507	
Middle Atlantic	0.1270		0.0311		0.0027	
Midwest	0.1621		0.3521		0.1470	
East North Central	0.0859		0.7188		0.3922	
West North Central	0.9801		0.2419		0.1723	
South	0.7854		0.1284		0.0804	
South Atlantic	0.5438		0.5684		0.3578	
East South Central	0.6043		0.4128		0.4052	
West South Central	0.9645		0.0966		0.1180	
West	0.1694		0.8784		0.4720	
Mountain	0.2811		0.8380		0.5788	
Pacific	0.3156		0.9428		0.5946	
COUNTY TYPE						
Large Metro	0.2299		0.0345		0.0079	
Small Metro	0.4007		0.7056		0.2380	
250K – 1 Mil. Pop.	0.2946		0.7813		0.1011	
< 250K Pop.	0.9402		0.2936		0.8217	
Nonmetro	0.3506		0.0086		0.0476	
Urbanized	0.1599		0.1852		0.5774	
Less Urbanized	0.5980		0.0026		0.0025	
Completely Rural	0.3201		0.4654		0.5011	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.69C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	693	632	189	186	133	105
GEOGRAPHIC DIVISION						
Northeast	292	287	77	79	41	52
New England	144	147	51	36	22	21
Middle Atlantic	253	250	58	70	35	48
Midwest	263	244	68	77	47	41
East North Central	208	209	62	70	43	37
West North Central	157	117	27	28	18	19
South	414	347	123	115	93	59
South Atlantic	346	266	107	85	86	55
East South Central	147	112	28	30	20	14
West South Central	205	192	58	73	30	22
West	398	384	113	98	76	59
Mountain	177	153	49	52	32	35
Pacific	359	352	101	83	68	48
COUNTY TYPE						
Large Metro	639	573	158	158	115	93
Small Metro	419	402	96	103	52	54
250K – 1 Mil. Pop.	364	338	84	89	45	48
< 250K Pop.	257	222	46	58	25	24
Nonmetro	302	227	56	38	48	18
Urbanized	213	154	27	28	15	12
Less Urbanized	198	169	42	24	39	12
Completely Rural	98	69	25	10	24	6

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.69D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.26	0.24	0.07	0.07	0.05	0.04
GEOGRAPHIC DIVISION						
Northeast	0.62	0.60	0.16	0.17	0.09	0.11
New England	1.15	1.17	0.41	0.29	0.17	0.17
Middle Atlantic	0.73	0.71	0.17	0.20	0.10	0.14
Midwest	0.47	0.43	0.12	0.14	0.08	0.07
East North Central	0.53	0.53	0.16	0.18	0.11	0.09
West North Central	0.91	0.67	0.16	0.16	0.10	0.11
South	0.42	0.35	0.13	0.12	0.10	0.06
South Atlantic	0.67	0.51	0.21	0.16	0.17	0.10
East South Central	0.95	0.72	0.18	0.19	0.13	0.09
West South Central	0.67	0.62	0.19	0.24	0.10	0.07
West	0.65	0.62	0.18	0.16	0.12	0.10
Mountain	0.94	0.81	0.26	0.28	0.17	0.18
Pacific	0.84	0.82	0.24	0.19	0.16	0.11
COUNTY TYPE						
Large Metro	0.39	0.35	0.11	0.11	0.08	0.06
Small Metro	0.45	0.44	0.12	0.13	0.07	0.07
250K – 1 Mil. Pop.	0.57	0.53	0.16	0.16	0.08	0.09
< 250K Pop.	0.83	0.75	0.18	0.22	0.10	0.10
Nonmetro	0.62	0.47	0.13	0.09	0.11	0.04
Urbanized	0.91	0.75	0.14	0.16	0.08	0.07
Less Urbanized	0.94	0.68	0.22	0.12	0.20	0.06
Completely Rural	1.81	1.37	0.52	0.22	0.50	0.15

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.69P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2134		0.2191		0.8370	
GEOGRAPHIC DIVISION						
Northeast	0.1691		0.0705		0.0500	
New England	0.6243		0.8049		0.5755	
Middle Atlantic	0.1941		0.0179		0.0548	
Midwest	0.0671		0.9826		0.8348	
East North Central	0.0714		0.6521		0.8476	
West North Central	0.5177		0.3044		0.9376	
South	0.7317		0.6795		0.1559	
South Atlantic	0.9332		0.9704		0.2733	
East South Central	0.3752		0.3377		0.7673	
West South Central	0.9620		0.7904		0.3176	
West	0.8238		0.6992		0.9652	
Mountain	0.8257		0.8532		0.6938	
Pacific	0.8862		0.7336		0.8631	
COUNTY TYPE						
Large Metro	0.1038		0.2880		0.9862	
Small Metro	0.4963		0.4005		0.5771	
250K – 1 Mil. Pop.	0.3504		0.6208		0.7615	
< 250K Pop.	0.8654		0.4261		0.5299	
Nonmetro	0.1977		0.4992		0.0971	
Urbanized	0.3777		0.9141		0.7606	
Less Urbanized	0.6971		0.9290		0.3168	
Completely Rural	0.2301		0.1814		0.1021	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.70C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	19	26	15	23	9	10
GEOGRAPHIC DIVISION						
Northeast	6	8	5	7	3	3
New England	2	3	2	1	0	*
Middle Atlantic	5	7	4	7	3	3
Midwest	6	11	4	9	2	4
East North Central	6	10	3	7	1	3
West North Central	3	4	3	4	2	3
South	12	13	10	10	6	8
South Atlantic	10	8	8	6	5	3
East South Central	3	3	4	2	3	*
West South Central	7	10	4	8	1	8
West	12	18	10	18	6	4
Mountain	8	7	6	6	3	3
Pacific	9	16	8	16	5	1
COUNTY TYPE						
Large Metro	14	21	10	20	6	7
Small Metro	12	9	10	7	7	6
250K – 1 Mil. Pop.	10	7	8	6	4	4
< 250K Pop.	8	6	7	4	5	4
Nonmetro	7	13	4	10	2	4
Urbanized	5	9	3	9	*	1
Less Urbanized	3	9	2	5	2	3
Completely Rural	1	3	*	3	*	*

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.70D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.08	0.10	0.06	0.09	0.04	0.04
GEOGRAPHIC DIVISION						
Northeast	0.14	0.19	0.11	0.16	0.07	0.08
New England	0.17	0.25	0.16	0.07	0.02	*
Middle Atlantic	0.18	0.24	0.14	0.22	0.10	0.11
Midwest	0.12	0.20	0.07	0.16	0.04	0.07
East North Central	0.15	0.26	0.08	0.20	0.03	0.08
West North Central	0.17	0.27	0.16	0.27	0.10	0.16
South	0.13	0.14	0.10	0.11	0.06	0.08
South Atlantic	0.21	0.17	0.16	0.13	0.10	0.06
East South Central	0.20	0.23	0.27	0.16	0.20	*
West South Central	0.21	0.30	0.14	0.25	0.04	0.23
West	0.20	0.30	0.16	0.30	0.10	0.06
Mountain	0.43	0.38	0.34	0.34	0.16	0.18
Pacific	0.21	0.41	0.19	0.41	0.13	0.03
COUNTY TYPE						
Large Metro	0.10	0.15	0.08	0.14	0.04	0.05
Small Metro	0.16	0.13	0.14	0.10	0.09	0.08
250K – 1 Mil. Pop.	0.18	0.14	0.15	0.12	0.08	0.09
< 250K Pop.	0.34	0.27	0.28	0.18	0.22	0.17
Nonmetro	0.17	0.33	0.09	0.27	0.05	0.11
Urbanized	0.33	0.58	0.16	0.56	*	0.09
Less Urbanized	0.20	0.45	0.14	0.26	0.12	0.14
Completely Rural	0.30	0.76	*	0.76	*	*

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.70P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.8252		0.1860		0.7663	
GEOGRAPHIC DIVISION						
Northeast	0.4201		0.4850		0.9305	
New England	0.7951		0.3812		*	
Middle Atlantic	0.4441		0.3563		0.8678	
Midwest	0.7137		0.2925		0.3995	
East North Central	0.4840		0.1708		0.3597	
West North Central	0.5267		0.8554		0.7927	
South	0.2951		0.7463		0.6533	
South Atlantic	0.0544		0.1923		0.1815	
East South Central	0.7488		0.7768		*	
West South Central	0.5536		0.3132		0.0560	
West	0.5394		0.2114		0.0782	
Mountain	0.2362		0.5640		0.8755	
Pacific	0.1645		0.0931		0.0314	
COUNTY TYPE						
Large Metro	0.2018		0.0425		0.9612	
Small Metro	0.0091		0.0167		0.4050	
250K – 1 Mil. Pop.	0.1444		0.1666		0.6993	
< 250K Pop.	0.0269		0.0348		0.4349	
Nonmetro	0.3241		0.0559		0.5187	
Urbanized	0.6037		0.1781		*	
Less Urbanized	0.5171		0.3970		0.6177	
Completely Rural	0.3603		*		*	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.71C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	108	120	72	81	34	40
GEOGRAPHIC DIVISION						
Northeast	46	41	30	32	12	13
New England	22	21	21	16	6	6
Middle Atlantic	41	34	22	28	10	12
Midwest	40	45	24	28	10	16
East North Central	32	38	21	24	8	14
West North Central	23	22	11	14	5	7
South	62	76	37	48	19	21
South Atlantic	48	51	29	36	15	18
East South Central	18	26	10	14	5	7
West South Central	32	49	21	27	10	8
West	64	67	46	50	23	28
Mountain	31	30	17	21	10	14
Pacific	55	61	42	45	21	24
COUNTY TYPE						
Large Metro	96	99	59	64	27	33
Small Metro	59	73	40	53	18	25
250K – 1 Mil. Pop.	47	60	30	42	15	21
< 250K Pop.	36	41	25	31	11	13
Nonmetro	41	41	20	21	7	7
Urbanized	33	27	18	12	5	4
Less Urbanized	25	29	10	16	5	6
Completely Rural	11	11	4	*	2	0

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.71D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.31	0.34	0.21	0.23	0.10	0.11
GEOGRAPHIC DIVISION						
Northeast	0.75	0.66	0.49	0.52	0.19	0.22
New England	1.37	1.25	1.29	0.97	0.36	0.38
Middle Atlantic	0.90	0.77	0.49	0.62	0.23	0.26
Midwest	0.54	0.61	0.32	0.38	0.13	0.21
East North Central	0.62	0.74	0.40	0.48	0.16	0.28
West North Central	1.00	0.97	0.46	0.61	0.23	0.29
South	0.48	0.59	0.28	0.37	0.15	0.17
South Atlantic	0.73	0.77	0.45	0.54	0.23	0.28
East South Central	0.90	1.27	0.49	0.70	0.25	0.36
West South Central	0.75	1.13	0.49	0.63	0.23	0.20
West	0.76	0.80	0.55	0.60	0.27	0.33
Mountain	1.23	1.17	0.68	0.83	0.38	0.53
Pacific	0.95	1.04	0.73	0.77	0.36	0.42
COUNTY TYPE						
Large Metro	0.45	0.48	0.30	0.32	0.15	0.17
Small Metro	0.51	0.60	0.35	0.46	0.17	0.23
250K – 1 Mil. Pop.	0.63	0.74	0.43	0.56	0.21	0.29
< 250K Pop.	0.82	0.99	0.59	0.77	0.28	0.35
Nonmetro	0.72	0.67	0.37	0.39	0.13	0.14
Urbanized	1.15	0.94	0.64	0.51	0.19	0.16
Less Urbanized	1.02	1.00	0.46	0.61	0.20	0.25
Completely Rural	1.96	2.42	0.78	*	0.34	0.06

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.71P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2725		0.6188		0.1629	
GEOGRAPHIC DIVISION						
Northeast	0.1373		0.5886		0.9580	
New England	0.5124		0.7343		0.4190	
Middle Atlantic	0.1751		0.6668		0.7009	
Midwest	0.1256		0.4224		0.2378	
East North Central	0.4552		0.9509		0.3841	
West North Central	0.0854		0.1146		0.3878	
South	0.4719		0.3811		0.3770	
South Atlantic	0.2782		0.1590		0.1146	
East South Central	0.6240		0.3432		0.8253	
West South Central	0.7144		0.3955		0.2848	
West	0.4923		0.9513		0.4810	
Mountain	0.5899		0.5148		0.4863	
Pacific	0.6213		0.7252		0.6637	
COUNTY TYPE						
Large Metro	0.2290		0.9513		0.4551	
Small Metro	0.6285		0.2951		0.1071	
250K – 1 Mil. Pop.	0.8977		0.6305		0.1950	
< 250K Pop.	0.5170		0.2618		0.3342	
Nonmetro	0.1720		0.8046		0.4421	
Urbanized	0.0238		0.1334		0.3289	
Less Urbanized	0.9734		0.2636		0.9774	
Completely Rural	0.4217		*		0.2509	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.72C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	674	611	172	165	129	98
GEOGRAPHIC DIVISION						
Northeast	289	282	73	75	39	52
New England	144	144	50	32	21	20
Middle Atlantic	250	246	54	69	33	48
Midwest	261	238	64	71	45	38
East North Central	208	202	59	65	42	34
West North Central	153	114	25	24	17	17
South	409	336	114	99	92	52
South Atlantic	340	255	100	76	85	50
East South Central	142	108	26	24	19	12
West South Central	203	189	53	62	29	19
West	384	372	100	83	73	55
Mountain	169	146	45	45	29	31
Pacific	346	343	89	70	65	46
COUNTY TYPE						
Large Metro	618	558	147	143	112	88
Small Metro	410	387	84	84	47	47
250K – 1 Mil. Pop.	355	324	75	77	42	43
< 250K Pop.	250	214	38	41	21	19
Nonmetro	289	218	51	28	47	16
Urbanized	203	148	19	23	13	12
Less Urbanized	189	159	41	16	38	10
Completely Rural	92	66	24	6	23	6

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.72D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.33	0.30	0.08	0.08	0.06	0.05
GEOGRAPHIC DIVISION						
Northeast	0.78	0.75	0.20	0.20	0.10	0.14
New England	1.47	1.47	0.51	0.32	0.22	0.21
Middle Atlantic	0.92	0.90	0.20	0.25	0.12	0.17
Midwest	0.60	0.55	0.15	0.16	0.10	0.09
East North Central	0.69	0.67	0.20	0.22	0.14	0.11
West North Central	1.15	0.85	0.19	0.18	0.13	0.13
South	0.54	0.44	0.15	0.13	0.12	0.07
South Atlantic	0.84	0.62	0.25	0.18	0.21	0.12
East South Central	1.18	0.90	0.21	0.20	0.16	0.10
West South Central	0.88	0.80	0.23	0.26	0.13	0.08
West	0.82	0.78	0.21	0.17	0.15	0.12
Mountain	1.18	1.00	0.31	0.31	0.20	0.21
Pacific	1.06	1.03	0.27	0.21	0.20	0.14
COUNTY TYPE						
Large Metro	0.49	0.43	0.13	0.12	0.10	0.08
Small Metro	0.58	0.55	0.14	0.14	0.08	0.08
250K – 1 Mil. Pop.	0.71	0.66	0.18	0.18	0.10	0.10
< 250K Pop.	1.09	0.97	0.19	0.21	0.11	0.10
Nonmetro	0.76	0.59	0.15	0.09	0.14	0.05
Urbanized	1.12	0.95	0.13	0.17	0.09	0.09
Less Urbanized	1.14	0.87	0.27	0.10	0.25	0.06
Completely Rural	2.17	1.62	0.62	0.16	0.62	0.16

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.72P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1591		0.3133		0.5229	
GEOGRAPHIC DIVISION						
Northeast	0.1151		0.0379		0.0467	
New England	0.5639		0.9283		0.7232	
Middle Atlantic	0.1398		0.0104		0.0402	
Midwest	0.0370		0.6619		0.5156	
East North Central	0.0559		0.5322		0.6023	
West North Central	0.3360		0.7024		0.6697	
South	0.6354		0.8784		0.0793	
South Atlantic	0.8186		0.7135		0.1514	
East South Central	0.3154		0.4901		0.7828	
West South Central	0.9324		0.5814		0.2783	
West	0.7818		0.7580		0.9300	
Mountain	0.7054		0.9842		0.8544	
Pacific	0.8961		0.7321		0.8429	
COUNTY TYPE						
Large Metro	0.0742		0.3177		0.8170	
Small Metro	0.5416		0.4954		0.9965	
250K – 1 Mil. Pop.	0.3570		0.6559		0.8571	
< 250K Pop.	0.7944		0.5428		0.7112	
Nonmetro	0.2564		0.2791		0.1045	
Urbanized	0.6253		0.7876		0.8949	
Less Urbanized	0.6764		0.5133		0.3278	
Completely Rural	0.1414		0.0752		0.0892	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.73C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	692	627	189	184	132	105
GEOGRAPHIC DIVISION						
Northeast	291	287	77	79	41	52
New England	143	147	51	36	22	21
Middle Atlantic	253	250	58	70	34	48
Midwest	263	244	68	76	47	41
East North Central	208	208	62	69	43	37
West North Central	157	117	27	28	18	18
South	416	343	122	114	93	58
South Atlantic	348	264	107	85	86	55
East South Central	144	112	28	30	20	14
West South Central	204	188	57	73	30	20
West	394	382	113	96	75	59
Mountain	175	152	49	52	31	35
Pacific	355	350	101	80	67	48
COUNTY TYPE						
Large Metro	638	572	157	157	114	92
Small Metro	418	402	95	102	51	54
250K – 1 Mil. Pop.	363	338	83	88	45	48
< 250K Pop.	256	221	46	57	24	23
Nonmetro	302	227	56	35	48	18
Urbanized	213	156	27	25	15	12
Less Urbanized	198	168	42	23	39	12
Completely Rural	98	68	25	9	24	6

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.73D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.29	0.26	0.08	0.08	0.06	0.04
GEOGRAPHIC DIVISION						
Northeast	0.67	0.66	0.18	0.18	0.09	0.12
New England	1.26	1.28	0.45	0.32	0.19	0.19
Middle Atlantic	0.79	0.78	0.18	0.22	0.11	0.15
Midwest	0.52	0.48	0.13	0.15	0.09	0.08
East North Central	0.59	0.59	0.18	0.20	0.12	0.10
West North Central	1.00	0.74	0.17	0.18	0.12	0.12
South	0.47	0.38	0.14	0.13	0.11	0.07
South Atlantic	0.74	0.56	0.23	0.18	0.18	0.12
East South Central	1.03	0.79	0.20	0.21	0.14	0.10
West South Central	0.75	0.68	0.21	0.26	0.11	0.07
West	0.71	0.68	0.20	0.17	0.14	0.11
Mountain	1.04	0.89	0.29	0.31	0.19	0.20
Pacific	0.92	0.89	0.26	0.21	0.17	0.12
COUNTY TYPE						
Large Metro	0.43	0.38	0.12	0.12	0.09	0.07
Small Metro	0.50	0.48	0.13	0.14	0.07	0.08
250K – 1 Mil. Pop.	0.62	0.58	0.17	0.18	0.09	0.10
< 250K Pop.	0.92	0.82	0.19	0.24	0.10	0.10
Nonmetro	0.68	0.52	0.14	0.09	0.12	0.05
Urbanized	0.99	0.84	0.16	0.16	0.09	0.08
Less Urbanized	1.03	0.76	0.24	0.12	0.22	0.06
Completely Rural	2.02	1.49	0.58	0.23	0.55	0.15

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.73P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2311		0.2783		0.8494	
GEOGRAPHIC DIVISION						
Northeast	0.1810		0.0807		0.0510	
New England	0.6344		0.8210		0.5707	
Middle Atlantic	0.2065		0.0227		0.0562	
Midwest	0.0717		0.9008		0.7861	
East North Central	0.0781		0.5658		0.8051	
West North Central	0.5105		0.2903		0.9120	
South	0.7436		0.6615		0.1428	
South Atlantic	0.9667		0.8998		0.3042	
East South Central	0.3709		0.3198		0.8610	
West South Central	0.9902		0.8689		0.1551	
West	0.8680		0.8315		0.8658	
Mountain	0.7896		0.7974		0.6829	
Pacific	0.9530		0.9189		0.9713	
COUNTY TYPE						
Large Metro	0.1146		0.3899		0.9817	
Small Metro	0.4917		0.2981		0.5161	
250K – 1 Mil. Pop.	0.3587		0.5526		0.7429	
< 250K Pop.	0.8952		0.3116		0.4300	
Nonmetro	0.1869		0.3088		0.0846	
Urbanized	0.3769		0.6516		0.6989	
Less Urbanized	0.6872		0.8543		0.3352	
Completely Rural	0.1973		0.1286		0.0763	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.74C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	364	317	77	81	64	55
GEOGRAPHIC DIVISION						
Northeast	134	128	25	39	23	26
New England	62	72	16	18	16	10
Middle Atlantic	117	107	20	35	17	23
Midwest	161	130	47	30	40	24
East North Central	127	98	46	23	38	17
West North Central	99	75	12	19	11	17
South	222	192	49	61	35	37
South Atlantic	179	159	39	42	27	38
East South Central	91	62	20	10	18	8
West South Central	99	101	21	45	14	10
West	200	167	30	24	28	21
Mountain	91	81	21	16	21	14
Pacific	180	150	21	18	19	15
COUNTY TYPE						
Large Metro	274	246	51	67	48	50
Small Metro	204	182	47	49	33	25
250K – 1 Mil. Pop.	176	144	44	37	30	25
< 250K Pop.	101	99	17	33	15	5
Nonmetro	158	121	34	15	26	13
Urbanized	85	72	13	9	9	5
Less Urbanized	120	84	21	10	18	10
Completely Rural	57	51	23	6	17	6

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.74D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.14	0.12	0.03	0.03	0.02	0.02
GEOGRAPHIC DIVISION						
Northeast	0.28	0.27	0.05	0.08	0.05	0.05
New England	0.49	0.57	0.13	0.15	0.12	0.08
Middle Atlantic	0.34	0.31	0.06	0.10	0.05	0.07
Midwest	0.29	0.23	0.08	0.05	0.07	0.04
East North Central	0.33	0.25	0.12	0.06	0.10	0.04
West North Central	0.57	0.43	0.07	0.11	0.07	0.10
South	0.23	0.19	0.05	0.06	0.04	0.04
South Atlantic	0.35	0.31	0.08	0.08	0.05	0.07
East South Central	0.59	0.39	0.13	0.07	0.11	0.05
West South Central	0.32	0.33	0.07	0.14	0.05	0.03
West	0.33	0.27	0.05	0.04	0.05	0.03
Mountain	0.48	0.42	0.11	0.08	0.11	0.07
Pacific	0.42	0.35	0.05	0.04	0.04	0.04
COUNTY TYPE						
Large Metro	0.19	0.16	0.04	0.05	0.03	0.03
Small Metro	0.25	0.22	0.06	0.06	0.04	0.03
250K – 1 Mil. Pop.	0.32	0.26	0.08	0.07	0.06	0.05
< 250K Pop.	0.37	0.37	0.07	0.13	0.06	0.02
Nonmetro	0.36	0.28	0.08	0.04	0.06	0.03
Urbanized	0.43	0.40	0.07	0.05	0.05	0.03
Less Urbanized	0.62	0.40	0.11	0.05	0.10	0.05
Completely Rural	1.17	1.12	0.49	0.13	0.36	0.13

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.74P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3420		0.2313		0.7509	
GEOGRAPHIC DIVISION						
Northeast	0.0808		0.0023		0.3579	
New England	0.3951		0.0810		0.9533	
Middle Atlantic	0.1261		0.0122		0.2564	
Midwest	0.6252		0.4528		0.5891	
East North Central	0.6416		0.2566		0.4617	
West North Central	0.8311		0.4700		0.7793	
South	0.7697		0.4720		0.8468	
South Atlantic	0.8653		0.8485		0.9437	
East South Central	0.9927		0.8712		0.7825	
West South Central	0.3638		0.1537		0.6560	
West	0.8304		0.5655		0.5089	
Mountain	0.5974		0.5756		0.5231	
Pacific	0.6077		0.7920		0.7706	
COUNTY TYPE						
Large Metro	0.4738		0.0058		0.5466	
Small Metro	0.1613		0.6764		0.3288	
250K – 1 Mil. Pop.	0.4672		0.3515		0.4363	
< 250K Pop.	0.1201		0.5014		0.4942	
Nonmetro	0.4483		0.0654		0.3193	
Urbanized	0.5887		0.3588		0.5987	
Less Urbanized	0.5431		0.4473		0.8095	
Completely Rural	0.9251		0.1225		0.2630	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.75C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	707	636	204	159	132	88
GEOGRAPHIC DIVISION						
Northeast	281	277	92	64	73	30
New England	146	132	30	36	9	10
Middle Atlantic	240	244	86	51	71	28
Midwest	265	258	65	60	24	30
East North Central	204	213	55	54	16	25
West North Central	170	151	35	26	18	15
South	430	347	101	82	52	49
South Atlantic	364	274	86	62	45	41
East South Central	130	109	29	29	17	22
West South Central	210	178	45	44	22	18
West	430	361	138	104	96	60
Mountain	199	135	62	40	22	19
Pacific	382	334	123	95	94	56
COUNTY TYPE						
Large Metro	651	570	162	138	94	77
Small Metro	458	376	96	74	57	42
250K – 1 Mil. Pop.	392	331	84	64	52	39
< 250K Pop.	264	189	46	40	24	19
Nonmetro	292	233	91	42	76	23
Urbanized	217	170	86	30	74	17
Less Urbanized	187	164	25	29	14	16
Completely Rural	81	64	15	6	13	2

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.75D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.27	0.24	0.08	0.06	0.05	0.03
GEOGRAPHIC DIVISION						
Northeast	0.59	0.58	0.19	0.13	0.15	0.06
New England	1.17	1.05	0.24	0.29	0.07	0.08
Middle Atlantic	0.69	0.69	0.25	0.15	0.20	0.08
Midwest	0.47	0.46	0.12	0.11	0.04	0.05
East North Central	0.52	0.54	0.14	0.14	0.04	0.07
West North Central	0.98	0.87	0.20	0.15	0.10	0.09
South	0.44	0.35	0.10	0.08	0.05	0.05
South Atlantic	0.71	0.52	0.17	0.12	0.09	0.08
East South Central	0.84	0.70	0.19	0.19	0.11	0.14
West South Central	0.69	0.57	0.15	0.14	0.07	0.06
West	0.70	0.58	0.22	0.17	0.16	0.10
Mountain	1.06	0.71	0.33	0.21	0.12	0.10
Pacific	0.90	0.77	0.29	0.22	0.22	0.13
COUNTY TYPE						
Large Metro	0.39	0.34	0.11	0.09	0.07	0.05
Small Metro	0.48	0.42	0.12	0.09	0.07	0.05
250K – 1 Mil. Pop.	0.60	0.52	0.16	0.12	0.10	0.07
< 250K Pop.	0.83	0.63	0.17	0.15	0.09	0.07
Nonmetro	0.61	0.49	0.21	0.10	0.18	0.06
Urbanized	0.94	0.86	0.44	0.17	0.38	0.10
Less Urbanized	0.89	0.67	0.13	0.14	0.07	0.08
Completely Rural	1.50	1.34	0.32	0.13	0.28	0.04

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.75P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5990		0.3865		0.2769	
GEOGRAPHIC DIVISION						
Northeast	0.8669		0.4940		0.4635	
New England	0.7268		0.8270		0.9342	
Middle Atlantic	0.6913		0.5090		0.4421	
Midwest	0.2304		0.6406		0.5639	
East North Central	0.3423		0.7867		0.4132	
West North Central	0.4709		0.6411		0.9172	
South	0.4727		0.1044		0.2485	
South Atlantic	0.8206		0.1532		0.3288	
East South Central	0.4885		0.7940		0.9588	
West South Central	0.5195		0.4371		0.4333	
West	0.3886		0.7749		0.6248	
Mountain	0.4916		0.3548		0.7628	
Pacific	0.5298		0.9035		0.5608	
COUNTY TYPE						
Large Metro	0.8587		0.7301		0.8179	
Small Metro	0.6981		0.4101		0.1202	
250K – 1 Mil. Pop.	0.9884		0.2574		0.1415	
< 250K Pop.	0.4484		0.7310		0.6239	
Nonmetro	0.4777		0.4727		0.2202	
Urbanized	0.8970		0.4578		0.2816	
Less Urbanized	0.6343		0.5134		0.6923	
Completely Rural	0.4340		0.1159		0.0531	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.76C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	519	428	105	95	64	65
GEOGRAPHIC DIVISION						
Northeast	240	194	32	47	21	41
New England	109	98	17	17	7	8
Middle Atlantic	215	168	27	44	20	40
Midwest	188	175	31	42	26	17
East North Central	141	138	29	31	26	16
West North Central	126	101	11	28	4	6
South	319	245	65	49	25	35
South Atlantic	257	193	59	31	19	22
East South Central	104	89	20	31	14	26
West South Central	162	117	18	22	7	8
West	301	248	68	56	48	30
Mountain	153	109	38	21	23	10
Pacific	261	220	56	52	42	29
COUNTY TYPE						
Large Metro	458	371	78	78	52	56
Small Metro	293	240	58	46	23	18
250K – 1 Mil. Pop.	267	207	54	39	15	17
< 250K Pop.	149	129	23	24	18	7
Nonmetro	199	160	37	29	27	26
Urbanized	151	107	30	24	24	23
Less Urbanized	121	108	18	17	7	12
Completely Rural	53	57	10	4	10	3

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.76D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.20	0.16	0.04	0.04	0.02	0.02
GEOGRAPHIC DIVISION						
Northeast	0.51	0.41	0.07	0.10	0.04	0.09
New England	0.87	0.78	0.13	0.13	0.06	0.06
Middle Atlantic	0.61	0.48	0.08	0.13	0.06	0.11
Midwest	0.34	0.31	0.06	0.07	0.05	0.03
East North Central	0.36	0.35	0.07	0.08	0.07	0.04
West North Central	0.73	0.58	0.06	0.16	0.03	0.03
South	0.33	0.25	0.07	0.05	0.03	0.04
South Atlantic	0.50	0.37	0.12	0.06	0.04	0.04
East South Central	0.67	0.57	0.13	0.20	0.09	0.17
West South Central	0.53	0.38	0.06	0.07	0.02	0.02
West	0.49	0.40	0.11	0.09	0.08	0.05
Mountain	0.82	0.58	0.20	0.11	0.12	0.05
Pacific	0.61	0.51	0.13	0.12	0.10	0.07
COUNTY TYPE						
Large Metro	0.30	0.23	0.06	0.05	0.04	0.04
Small Metro	0.33	0.28	0.07	0.06	0.03	0.02
250K – 1 Mil. Pop.	0.44	0.34	0.10	0.07	0.03	0.03
< 250K Pop.	0.53	0.47	0.09	0.10	0.07	0.03
Nonmetro	0.44	0.35	0.09	0.07	0.06	0.06
Urbanized	0.71	0.55	0.16	0.14	0.13	0.13
Less Urbanized	0.62	0.50	0.10	0.08	0.04	0.06
Completely Rural	1.08	1.19	0.22	0.09	0.21	0.06

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.76P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9936		0.6320		0.6223	
GEOGRAPHIC DIVISION						
Northeast	0.6156		0.1592		0.3441	
New England	0.8652		0.9590		0.8052	
Middle Atlantic	0.5099		0.1252		0.3024	
Midwest	0.1992		0.2425		0.9671	
East North Central	0.0972		0.3845		0.9357	
West North Central	0.9890		0.4244		0.8687	
South	0.9022		0.7263		0.1889	
South Atlantic	0.5612		0.4441		0.3428	
East South Central	0.8544		0.2626		0.2168	
West South Central	0.1924		0.5304		0.4736	
West	0.2713		0.6058		0.3421	
Mountain	0.2552		0.0391		0.1520	
Pacific	0.5285		0.5574		0.7264	
COUNTY TYPE						
Large Metro	0.8349		0.3201		0.3546	
Small Metro	0.5763		0.9793		0.3093	
250K – 1 Mil. Pop.	0.4986		0.9239		0.8794	
< 250K Pop.	0.9532		0.8947		0.0867	
Nonmetro	0.6711		0.3248		0.9587	
Urbanized	0.4679		0.3576		0.8310	
Less Urbanized	0.9387		0.8046		0.2295	
Completely Rural	0.5448		0.7355		0.5416	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.77C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	52	56	29	36	15	19
GEOGRAPHIC DIVISION						
Northeast	15	20	11	12	5	7
New England	6	9	3	4	2	3
Middle Atlantic	13	18	10	11	5	6
Midwest	18	21	11	15	5	8
East North Central	15	19	11	14	5	8
West North Central	11	9	4	6	3	0
South	38	36	20	22	12	12
South Atlantic	32	25	16	14	10	9
East South Central	11	11	6	8	2	5
West South Central	17	24	11	14	6	7
West	26	31	14	21	5	9
Mountain	13	12	7	6	3	4
Pacific	23	28	12	20	4	8
COUNTY TYPE						
Large Metro	39	48	23	29	11	14
Small Metro	34	29	16	19	9	10
250K – 1 Mil. Pop.	29	23	13	15	6	9
< 250K Pop.	17	17	10	11	6	4
Nonmetro	17	18	11	14	7	7
Urbanized	10	12	5	8	3	5
Less Urbanized	13	13	9	10	6	5
Completely Rural	5	5	2	3	0	3

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.77D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.21	0.23	0.12	0.14	0.06	0.07
GEOGRAPHIC DIVISION						
Northeast	0.35	0.47	0.25	0.29	0.13	0.16
New England	0.51	0.79	0.32	0.41	0.20	0.26
Middle Atlantic	0.43	0.58	0.32	0.37	0.16	0.20
Midwest	0.33	0.38	0.21	0.29	0.10	0.15
East North Central	0.41	0.50	0.28	0.37	0.12	0.21
West North Central	0.65	0.55	0.26	0.37	0.16	0.01
South	0.40	0.38	0.21	0.23	0.13	0.13
South Atlantic	0.68	0.53	0.34	0.31	0.21	0.19
East South Central	0.76	0.76	0.41	0.55	0.14	0.31
West South Central	0.54	0.72	0.33	0.42	0.19	0.22
West	0.45	0.52	0.23	0.35	0.08	0.15
Mountain	0.71	0.64	0.39	0.34	0.16	0.20
Pacific	0.56	0.70	0.29	0.49	0.09	0.20
COUNTY TYPE						
Large Metro	0.27	0.32	0.17	0.20	0.08	0.10
Small Metro	0.42	0.38	0.21	0.26	0.11	0.14
250K – 1 Mil. Pop.	0.53	0.45	0.25	0.31	0.12	0.18
< 250K Pop.	0.67	0.72	0.39	0.48	0.26	0.19
Nonmetro	0.42	0.47	0.27	0.35	0.18	0.19
Urbanized	0.59	0.74	0.31	0.51	0.17	0.29
Less Urbanized	0.69	0.62	0.51	0.52	0.36	0.27
Completely Rural	1.06	1.26	0.54	0.85	0.04	0.68

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.77P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.7986		0.4803		0.2319	
GEOGRAPHIC DIVISION						
Northeast	0.7278		0.9780		0.5651	
New England	0.5245		0.7824		0.7736	
Middle Atlantic	0.9275		0.9423		0.6161	
Midwest	0.3304		0.3587		0.4194	
East North Central	0.0786		0.3854		0.1492	
West North Central	0.2640		0.7377		0.0366	
South	0.1191		0.4596		0.7828	
South Atlantic	0.0869		0.2294		0.9551	
East South Central	0.1371		0.5047		0.0911	
West South Central	0.6972		0.8758		0.7468	
West	0.5300		0.1474		0.2775	
Mountain	0.7003		0.2874		0.9818	
Pacific	0.3701		0.0438		0.2168	
COUNTY TYPE						
Large Metro	0.6947		0.6973		0.2853	
Small Metro	0.5036		0.6410		0.5834	
250K – 1 Mil. Pop.	0.6996		0.8093		0.2447	
< 250K Pop.	0.5264		0.6457		0.4460	
Nonmetro	0.5647		0.6331		0.6964	
Urbanized	0.9096		0.8667		0.8770	
Less Urbanized	0.1176		0.9199		0.8215	
Completely Rural	0.0947		0.0719		0.0994	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.78C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	785	658	418	321	256	222
GEOGRAPHIC DIVISION						
Northeast	273	283	138	145	77	101
New England	142	160	65	68	35	51
Middle Atlantic	232	237	122	129	69	88
Midwest	304	285	132	126	81	81
East North Central	252	228	111	110	70	71
West North Central	178	159	69	66	41	36
South	505	390	280	189	176	133
South Atlantic	409	294	232	130	141	103
East South Central	159	122	83	71	64	45
West South Central	233	217	117	117	84	73
West	431	361	238	174	144	117
Mountain	220	157	127	83	64	54
Pacific	374	326	202	155	128	107
COUNTY TYPE						
Large Metro	750	618	326	265	198	185
Small Metro	524	424	241	185	154	120
250K – 1 Mil. Pop.	427	332	183	150	128	100
< 250K Pop.	322	280	164	107	89	68
Nonmetro	362	279	176	118	75	64
Urbanized	280	206	135	90	45	46
Less Urbanized	233	209	111	79	59	45
Completely Rural	91	72	39	24	23	15

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.78D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.30	0.25	0.16	0.12	0.10	0.08
GEOGRAPHIC DIVISION						
Northeast	0.58	0.59	0.29	0.30	0.16	0.21
New England	1.14	1.27	0.52	0.54	0.28	0.41
Middle Atlantic	0.66	0.68	0.35	0.37	0.20	0.25
Midwest	0.54	0.50	0.23	0.22	0.14	0.14
East North Central	0.65	0.58	0.29	0.28	0.18	0.18
West North Central	1.03	0.92	0.40	0.38	0.24	0.21
South	0.52	0.39	0.29	0.19	0.18	0.13
South Atlantic	0.79	0.56	0.45	0.25	0.27	0.20
East South Central	1.03	0.78	0.53	0.46	0.41	0.29
West South Central	0.76	0.70	0.38	0.38	0.28	0.23
West	0.70	0.58	0.39	0.28	0.23	0.19
Mountain	1.18	0.82	0.68	0.44	0.34	0.28
Pacific	0.88	0.75	0.47	0.36	0.30	0.25
COUNTY TYPE						
Large Metro	0.43	0.35	0.22	0.17	0.14	0.12
Small Metro	0.53	0.45	0.29	0.22	0.19	0.15
250K – 1 Mil. Pop.	0.65	0.51	0.33	0.27	0.24	0.18
< 250K Pop.	0.96	0.85	0.58	0.36	0.32	0.25
Nonmetro	0.70	0.54	0.39	0.26	0.17	0.15
Urbanized	1.14	0.89	0.64	0.47	0.22	0.25
Less Urbanized	0.98	0.82	0.56	0.35	0.30	0.21
Completely Rural	1.57	1.23	0.77	0.52	0.49	0.34

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.78P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5162		0.3072		0.9662	
GEOGRAPHIC DIVISION						
Northeast	0.0291		0.1690		0.0460	
New England	0.1800		0.5042		0.0473	
Middle Atlantic	0.0849		0.2324		0.2431	
Midwest	0.2586		0.9435		0.9553	
East North Central	0.0591		0.7683		0.9599	
West North Central	0.4728		0.5318		0.9810	
South	0.8630		0.2424		0.2994	
South Atlantic	0.6694		0.1371		0.7058	
East South Central	0.7922		0.6974		0.3181	
West South Central	0.3932		0.8045		0.4470	
West	0.1607		0.1413		0.8513	
Mountain	0.4044		0.2747		0.6833	
Pacific	0.2565		0.2924		0.6806	
COUNTY TYPE						
Large Metro	0.5379		0.6339		0.6975	
Small Metro	0.5521		0.3293		0.4594	
250K – 1 Mil. Pop.	0.7722		0.7807		0.6559	
< 250K Pop.	0.5434		0.2169		0.5020	
Nonmetro	0.6326		0.6717		0.8877	
Urbanized	0.7531		0.6820		0.1938	
Less Urbanized	0.8592		0.4663		0.4190	
Completely Rural	0.8366		0.3632		0.9008	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.79C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	63	69	50	60	32	41
GEOGRAPHIC DIVISION						
Northeast	19	24	17	20	9	13
New England	8	9	7	8	4	4
Middle Atlantic	17	22	15	18	8	12
Midwest	24	29	19	25	13	16
East North Central	19	26	15	23	10	14
West North Central	14	15	11	12	8	9
South	41	44	31	39	20	26
South Atlantic	28	29	22	26	14	17
East South Central	13	16	11	12	7	9
West South Central	25	31	18	28	13	18
West	37	36	30	30	19	21
Mountain	18	16	14	14	7	9
Pacific	33	33	26	27	18	19
COUNTY TYPE						
Large Metro	52	58	41	49	25	32
Small Metro	40	38	31	31	19	19
250K – 1 Mil. Pop.	33	30	25	26	16	16
< 250K Pop.	23	22	17	18	10	11
Nonmetro	26	30	17	27	11	16
Urbanized	17	22	13	19	9	10
Less Urbanized	21	20	10	18	7	12
Completely Rural	8	6	6	5	3	3

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.79D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.25	0.28	0.20	0.24	0.13	0.16
GEOGRAPHIC DIVISION						
Northeast	0.45	0.57	0.40	0.48	0.22	0.31
New England	0.76	0.86	0.67	0.76	0.37	0.41
Middle Atlantic	0.54	0.71	0.49	0.59	0.26	0.39
Midwest	0.44	0.54	0.35	0.47	0.24	0.30
East North Central	0.49	0.70	0.40	0.61	0.26	0.37
West North Central	0.84	0.89	0.67	0.75	0.48	0.56
South	0.44	0.46	0.33	0.42	0.21	0.27
South Atlantic	0.61	0.61	0.48	0.55	0.29	0.36
East South Central	0.91	1.09	0.77	0.78	0.45	0.60
West South Central	0.78	0.94	0.55	0.85	0.40	0.55
West	0.63	0.61	0.50	0.51	0.32	0.36
Mountain	0.93	0.83	0.74	0.75	0.39	0.50
Pacific	0.81	0.82	0.65	0.67	0.43	0.46
COUNTY TYPE						
Large Metro	0.34	0.38	0.28	0.33	0.18	0.23
Small Metro	0.47	0.48	0.37	0.40	0.24	0.26
250K – 1 Mil. Pop.	0.56	0.55	0.44	0.47	0.30	0.32
< 250K Pop.	0.86	0.91	0.68	0.75	0.44	0.50
Nonmetro	0.63	0.72	0.42	0.66	0.28	0.40
Urbanized	0.94	1.24	0.69	1.14	0.49	0.65
Less Urbanized	1.06	0.95	0.56	0.89	0.37	0.60
Completely Rural	1.80	1.60	1.37	1.27	0.64	0.77

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.79P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2713		0.1489		0.0396	
GEOGRAPHIC DIVISION						
Northeast	0.4162		0.3780		0.3377	
New England	0.6484		0.6307		0.5982	
Middle Atlantic	0.2694		0.2294		0.2075	
Midwest	0.8812		0.5545		0.7384	
East North Central	0.4772		0.2886		0.6575	
West North Central	0.3928		0.5333		0.9640	
South	0.1355		0.0615		0.0175	
South Atlantic	0.2822		0.1607		0.0868	
East South Central	0.7423		0.3324		0.3655	
West South Central	0.1809		0.0610		0.1696	
West	0.7353		0.6006		0.8121	
Mountain	0.2191		0.1370		0.6418	
Pacific	0.8042		0.8454		0.9581	
COUNTY TYPE						
Large Metro	0.3845		0.4933		0.3158	
Small Metro	0.9115		0.8279		0.3137	
250K – 1 Mil. Pop.	0.3622		0.5489		0.1346	
< 250K Pop.	0.2826		0.2227		0.6516	
Nonmetro	0.2410		0.0051		0.0234	
Urbanized	0.2030		0.0458		0.0618	
Less Urbanized	0.6652		0.0374		0.2638	
Completely Rural	0.9573		0.7914		0.3623	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.80C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	149	165	110	115	69	71
GEOGRAPHIC DIVISION						
Northeast	60	69	43	52	24	33
New England	29	35	26	30	13	17
Middle Atlantic	52	60	33	44	20	29
Midwest	54	67	42	50	26	34
East North Central	46	57	34	44	21	29
West North Central	28	34	24	27	16	16
South	92	94	70	72	43	41
South Atlantic	73	64	54	47	31	24
East South Central	33	42	23	30	17	20
West South Central	46	57	38	45	24	27
West	82	87	59	53	40	35
Mountain	36	36	27	26	17	14
Pacific	74	80	52	48	36	32
COUNTY TYPE						
Large Metro	156	157	97	102	57	64
Small Metro	107	118	73	71	38	36
250K – 1 Mil. Pop.	83	90	56	59	29	30
< 250K Pop.	73	76	47	40	25	21
Nonmetro	70	78	50	42	27	23
Urbanized	55	60	39	30	20	15
Less Urbanized	46	49	29	28	19	17
Completely Rural	17	16	11	6	3	4

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.80D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.43	0.47	0.32	0.33	0.20	0.20
GEOGRAPHIC DIVISION						
Northeast	0.97	1.13	0.70	0.84	0.39	0.53
New England	1.79	2.15	1.61	1.81	0.77	1.01
Middle Atlantic	1.15	1.34	0.74	0.97	0.45	0.65
Midwest	0.74	0.90	0.57	0.68	0.35	0.46
East North Central	0.91	1.11	0.67	0.86	0.40	0.56
West North Central	1.24	1.49	1.05	1.18	0.69	0.70
South	0.72	0.73	0.54	0.56	0.34	0.32
South Atlantic	1.12	0.97	0.83	0.72	0.48	0.36
East South Central	1.63	2.05	1.12	1.47	0.82	0.99
West South Central	1.08	1.34	0.88	1.05	0.57	0.63
West	0.98	1.04	0.70	0.63	0.47	0.41
Mountain	1.40	1.42	1.07	1.00	0.67	0.56
Pacific	1.26	1.36	0.90	0.81	0.61	0.54
COUNTY TYPE						
Large Metro	0.61	0.65	0.43	0.47	0.29	0.31
Small Metro	0.72	0.84	0.54	0.57	0.32	0.31
250K – 1 Mil. Pop.	0.92	1.00	0.67	0.75	0.39	0.41
< 250K Pop.	1.10	1.54	0.88	0.90	0.54	0.52
Nonmetro	0.92	1.08	0.77	0.70	0.48	0.42
Urbanized	1.21	1.81	1.14	1.05	0.71	0.61
Less Urbanized	1.53	1.31	1.09	0.97	0.77	0.66
Completely Rural	2.68	3.03	1.92	1.52	0.61	0.97

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.80P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.7131		0.4133		0.2389	
GEOGRAPHIC DIVISION						
Northeast	0.8478		0.4750		0.8968	
New England	0.9770		0.1225		0.5869	
Middle Atlantic	0.8392		0.8112		0.6426	
Midwest	0.2145		0.2548		0.9277	
East North Central	0.0630		0.3677		0.6866	
West North Central	0.5150		0.4934		0.6489	
South	0.2864		0.4668		0.3380	
South Atlantic	0.1483		0.7116		0.2213	
East South Central	0.7789		0.1982		0.8343	
West South Central	0.8327		0.9749		0.9231	
West	0.0038		0.0039		0.2713	
Mountain	0.0582		0.1421		0.1537	
Pacific	0.0208		0.0133		0.5840	
COUNTY TYPE						
Large Metro	0.5371		0.8057		0.8105	
Small Metro	0.5312		0.6342		0.1398	
250K – 1 Mil. Pop.	0.3503		0.8641		0.5830	
< 250K Pop.	0.8552		0.5530		0.0863	
Nonmetro	0.4956		0.2608		0.4111	
Urbanized	0.4746		0.3466		0.6060	
Less Urbanized	0.6613		0.6082		0.3179	
Completely Rural	0.7315		0.3679		0.2242	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.81C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	751	622	386	286	244	207
GEOGRAPHIC DIVISION						
Northeast	265	269	127	133	71	94
New England	138	152	53	58	31	46
Middle Atlantic	226	225	115	120	64	83
Midwest	297	268	119	102	76	68
East North Central	245	212	103	90	66	61
West North Central	171	151	59	53	37	32
South	492	378	266	169	169	123
South Atlantic	393	288	221	119	137	100
East South Central	155	116	77	66	60	37
West South Central	230	207	108	105	79	64
West	406	331	216	157	139	112
Mountain	211	156	116	78	61	51
Pacific	352	291	183	138	125	103
COUNTY TYPE						
Large Metro	701	571	293	236	187	169
Small Metro	490	379	223	154	149	108
250K – 1 Mil. Pop.	394	307	163	125	121	90
< 250K Pop.	295	238	155	87	87	60
Nonmetro	341	247	166	102	68	56
Urbanized	266	177	127	76	37	39
Less Urbanized	212	185	105	67	54	38
Completely Rural	78	62	35	21	21	15

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.81D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.37	0.30	0.19	0.14	0.12	0.10
GEOGRAPHIC DIVISION						
Northeast	0.72	0.72	0.34	0.36	0.19	0.25
New England	1.41	1.54	0.55	0.59	0.32	0.47
Middle Atlantic	0.83	0.82	0.42	0.44	0.24	0.30
Midwest	0.68	0.61	0.27	0.23	0.18	0.16
East North Central	0.81	0.70	0.34	0.30	0.22	0.20
West North Central	1.29	1.13	0.44	0.40	0.28	0.24
South	0.65	0.49	0.35	0.22	0.22	0.16
South Atlantic	0.98	0.70	0.55	0.29	0.34	0.24
East South Central	1.29	0.96	0.64	0.54	0.50	0.30
West South Central	1.00	0.88	0.47	0.44	0.34	0.27
West	0.87	0.69	0.46	0.33	0.30	0.23
Mountain	1.47	1.07	0.81	0.54	0.43	0.35
Pacific	1.08	0.87	0.56	0.41	0.38	0.31
COUNTY TYPE						
Large Metro	0.53	0.43	0.26	0.20	0.17	0.15
Small Metro	0.66	0.54	0.35	0.25	0.24	0.18
250K – 1 Mil. Pop.	0.80	0.64	0.39	0.30	0.29	0.22
< 250K Pop.	1.21	0.99	0.74	0.42	0.42	0.30
Nonmetro	0.85	0.64	0.48	0.30	0.20	0.17
Urbanized	1.41	1.02	0.80	0.53	0.25	0.28
Less Urbanized	1.19	1.00	0.68	0.41	0.36	0.24
Completely Rural	1.75	1.38	0.93	0.58	0.58	0.42

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.81P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5229		0.2962		0.9959	
GEOGRAPHIC DIVISION						
Northeast	0.0315		0.0987		0.0385	
New England	0.1584		0.1014		0.0369	
Middle Atlantic	0.1010		0.2967		0.2168	
Midwest	0.3776		0.6142		0.8714	
East North Central	0.1394		0.3648		0.7708	
West North Central	0.5807		0.5709		0.8440	
South	0.8456		0.0798		0.2372	
South Atlantic	0.4119		0.0654		0.7507	
East South Central	0.8203		0.9223		0.2408	
West South Central	0.4496		0.5142		0.2787	
West	0.4038		0.5104		0.9205	
Mountain	0.6987		0.5624		0.4453	
Pacific	0.4618		0.6787		0.7918	
COUNTY TYPE						
Large Metro	0.4754		0.6212		0.7176	
Small Metro	0.6739		0.3965		0.6064	
250K – 1 Mil. Pop.	0.9667		0.7348		0.5848	
< 250K Pop.	0.4320		0.3526		0.9141	
Nonmetro	0.6086		0.5310		0.9445	
Urbanized	0.7311		0.7648		0.2466	
Less Urbanized	0.8771		0.3253		0.4432	
Completely Rural	0.7694		0.5111		0.6076	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.82C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	778	657	415	314	255	217
GEOGRAPHIC DIVISION						
Northeast	273	281	136	142	76	99
New England	142	160	65	68	35	51
Middle Atlantic	231	235	119	126	68	86
Midwest	302	284	130	121	81	78
East North Central	249	226	110	105	69	68
West North Central	178	159	69	65	41	35
South	502	387	280	185	176	129
South Atlantic	406	294	233	127	141	101
East South Central	158	120	81	70	64	43
West South Central	230	213	117	114	84	70
West	422	362	234	171	143	117
Mountain	217	159	125	82	65	52
Pacific	366	325	198	152	127	108
COUNTY TYPE						
Large Metro	743	612	322	257	198	180
Small Metro	514	418	236	177	153	117
250K – 1 Mil. Pop.	417	330	178	145	127	98
< 250K Pop.	316	272	162	101	88	66
Nonmetro	359	272	174	113	73	61
Urbanized	279	200	133	84	42	43
Less Urbanized	228	203	111	76	58	42
Completely Rural	87	68	36	23	21	15

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.82D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.33	0.27	0.17	0.13	0.11	0.09
GEOGRAPHIC DIVISION						
Northeast	0.63	0.65	0.31	0.33	0.18	0.23
New England	1.25	1.39	0.57	0.59	0.30	0.45
Middle Atlantic	0.73	0.73	0.38	0.39	0.21	0.27
Midwest	0.59	0.56	0.26	0.24	0.16	0.15
East North Central	0.71	0.64	0.31	0.30	0.20	0.19
West North Central	1.14	1.02	0.44	0.41	0.26	0.22
South	0.57	0.43	0.32	0.21	0.20	0.14
South Atlantic	0.87	0.62	0.50	0.27	0.30	0.21
East South Central	1.12	0.85	0.58	0.50	0.45	0.31
West South Central	0.84	0.77	0.43	0.41	0.31	0.25
West	0.76	0.64	0.42	0.30	0.26	0.21
Mountain	1.29	0.93	0.74	0.48	0.39	0.31
Pacific	0.95	0.83	0.52	0.39	0.33	0.27
COUNTY TYPE						
Large Metro	0.47	0.39	0.24	0.19	0.15	0.13
Small Metro	0.58	0.49	0.31	0.24	0.21	0.16
250K – 1 Mil. Pop.	0.71	0.57	0.36	0.29	0.26	0.20
< 250K Pop.	1.04	0.90	0.63	0.38	0.35	0.26
Nonmetro	0.77	0.58	0.43	0.28	0.19	0.16
Urbanized	1.24	0.96	0.70	0.49	0.23	0.26
Less Urbanized	1.08	0.90	0.61	0.39	0.33	0.22
Completely Rural	1.66	1.30	0.82	0.56	0.51	0.38

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.82P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.6047		0.2110		0.7175	
GEOGRAPHIC DIVISION						
Northeast	0.0347		0.2004		0.0564	
New England	0.1748		0.4696		0.0409	
Middle Atlantic	0.1036		0.2909		0.3045	
Midwest	0.2683		0.9745		0.9053	
East North Central	0.0679		0.6207		0.8978	
West North Central	0.5155		0.4622		0.9891	
South	0.9839		0.1465		0.1613	
South Atlantic	0.5998		0.0951		0.5533	
East South Central	0.7672		0.5882		0.2529	
West South Central	0.4905		0.5311		0.2922	
West	0.1574		0.1560		0.8210	
Mountain	0.4619		0.3611		0.7320	
Pacific	0.2297		0.2720		0.6730	
COUNTY TYPE						
Large Metro	0.5947		0.5555		0.8100	
Small Metro	0.5927		0.3374		0.3759	
250K – 1 Mil. Pop.	0.8720		0.7153		0.5124	
< 250K Pop.	0.4929		0.2729		0.5385	
Nonmetro	0.5496		0.3869		0.7477	
Urbanized	0.6686		0.9084		0.3445	
Less Urbanized	0.8287		0.2954		0.3017	
Completely Rural	0.7893		0.3301		0.7783	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.83C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	640	553	343	253	206	174
GEOGRAPHIC DIVISION						
Northeast	204	217	101	112	67	76
New England	115	121	51	49	26	35
Middle Atlantic	168	185	88	103	62	68
Midwest	226	227	112	108	68	72
East North Central	177	192	94	96	61	64
West North Central	138	116	55	52	31	29
South	401	316	234	156	148	107
South Atlantic	320	235	199	109	118	85
East South Central	130	114	78	55	60	30
West South Central	195	170	98	98	65	60
West	375	307	191	133	112	91
Mountain	189	131	113	66	45	41
Pacific	326	281	154	121	103	83
COUNTY TYPE						
Large Metro	570	472	258	208	166	139
Small Metro	376	328	191	148	122	103
250K – 1 Mil. Pop.	313	268	152	119	108	84
< 250K Pop.	221	200	113	88	60	60
Nonmetro	263	199	161	95	57	49
Urbanized	193	142	129	73	38	39
Less Urbanized	176	158	91	61	39	30
Completely Rural	71	61	34	19	21	11

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.83D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.24	0.21	0.13	0.10	0.08	0.07
GEOGRAPHIC DIVISION						
Northeast	0.43	0.46	0.21	0.24	0.14	0.16
New England	0.92	0.97	0.41	0.39	0.20	0.28
Middle Atlantic	0.48	0.53	0.25	0.29	0.18	0.19
Midwest	0.40	0.40	0.20	0.19	0.12	0.13
East North Central	0.45	0.49	0.24	0.25	0.16	0.16
West North Central	0.80	0.67	0.32	0.30	0.18	0.17
South	0.41	0.32	0.24	0.16	0.15	0.11
South Atlantic	0.62	0.45	0.39	0.21	0.23	0.16
East South Central	0.84	0.73	0.50	0.35	0.39	0.20
West South Central	0.64	0.55	0.32	0.31	0.21	0.19
West	0.61	0.49	0.31	0.21	0.18	0.15
Mountain	1.01	0.69	0.60	0.35	0.24	0.22
Pacific	0.77	0.65	0.36	0.28	0.24	0.19
COUNTY TYPE						
Large Metro	0.35	0.29	0.17	0.14	0.12	0.09
Small Metro	0.42	0.37	0.23	0.18	0.15	0.13
250K – 1 Mil. Pop.	0.53	0.45	0.27	0.22	0.20	0.16
< 250K Pop.	0.72	0.65	0.41	0.32	0.23	0.22
Nonmetro	0.54	0.40	0.36	0.21	0.13	0.11
Urbanized	0.84	0.62	0.63	0.38	0.20	0.21
Less Urbanized	0.78	0.65	0.46	0.29	0.20	0.14
Completely Rural	1.31	1.18	0.70	0.44	0.47	0.27

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.83P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.7945		0.0451		0.3709	
GEOGRAPHIC DIVISION						
Northeast	0.0011		0.3068		0.3015	
New England	0.0179		0.6110		0.0799	
Middle Atlantic	0.0200		0.3822		0.7599	
Midwest	0.2903		0.8403		0.9100	
East North Central	0.1338		0.8425		0.8015	
West North Central	0.7718		0.9490		0.7992	
South	0.4069		0.0925		0.1872	
South Atlantic	0.2521		0.1420		0.5897	
East South Central	0.8552		0.3484		0.1295	
West South Central	0.8060		0.7200		0.4952	
West	0.1594		0.0147		0.4157	
Mountain	0.0740		0.1072		0.5635	
Pacific	0.5297		0.0665		0.2492	
COUNTY TYPE						
Large Metro	0.9742		0.1089		0.4338	
Small Metro	0.4352		0.4043		0.5265	
250K – 1 Mil. Pop.	0.5959		0.5199		0.4834	
< 250K Pop.	0.5471		0.5917		0.9555	
Nonmetro	0.5609		0.3744		0.7709	
Urbanized	0.5596		0.8992		0.0692	
Less Urbanized	0.7725		0.4066		0.3482	
Completely Rural	0.7338		0.3058		0.3363	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.84C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	58	63	46	53	29	33
GEOGRAPHIC DIVISION						
Northeast	16	21	13	18	8	12
New England	7	6	6	5	4	4
Middle Atlantic	15	20	11	17	7	11
Midwest	22	27	18	24	11	12
East North Central	18	25	14	22	9	11
West North Central	12	12	10	9	7	6
South	38	39	29	33	18	21
South Atlantic	26	25	20	21	12	13
East South Central	12	14	11	11	6	8
West South Central	25	26	17	23	11	15
West	34	34	27	28	17	17
Mountain	15	15	12	13	6	8
Pacific	30	30	24	24	16	15
COUNTY TYPE						
Large Metro	46	50	37	42	22	26
Small Metro	36	35	27	27	17	17
250K – 1 Mil. Pop.	30	29	22	23	14	15
< 250K Pop.	20	19	15	15	9	9
Nonmetro	23	26	16	23	11	12
Urbanized	15	19	12	18	8	8
Less Urbanized	18	17	10	15	7	10
Completely Rural	8	6	5	4	2	2

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.84D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.23	0.25	0.18	0.21	0.11	0.13
GEOGRAPHIC DIVISION						
Northeast	0.39	0.51	0.31	0.43	0.19	0.29
New England	0.66	0.60	0.58	0.48	0.37	0.41
Middle Atlantic	0.47	0.66	0.37	0.55	0.22	0.36
Midwest	0.41	0.50	0.33	0.44	0.20	0.23
East North Central	0.48	0.66	0.38	0.58	0.23	0.30
West North Central	0.72	0.72	0.62	0.58	0.40	0.36
South	0.40	0.41	0.31	0.35	0.19	0.23
South Atlantic	0.55	0.53	0.44	0.45	0.26	0.29
East South Central	0.82	0.97	0.73	0.74	0.41	0.54
West South Central	0.76	0.80	0.54	0.70	0.36	0.47
West	0.57	0.57	0.45	0.46	0.29	0.29
Mountain	0.82	0.80	0.62	0.68	0.32	0.40
Pacific	0.74	0.75	0.59	0.60	0.40	0.38
COUNTY TYPE						
Large Metro	0.31	0.34	0.25	0.29	0.16	0.18
Small Metro	0.44	0.45	0.33	0.36	0.22	0.23
250K – 1 Mil. Pop.	0.53	0.54	0.39	0.44	0.27	0.30
< 250K Pop.	0.78	0.80	0.62	0.64	0.39	0.42
Nonmetro	0.57	0.64	0.39	0.59	0.27	0.32
Urbanized	0.87	1.16	0.64	1.11	0.47	0.51
Less Urbanized	0.92	0.82	0.53	0.73	0.36	0.49
Completely Rural	1.64	1.57	1.14	1.07	0.36	0.45

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 1.84P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.9936		0.9004		0.3296	
GEOGRAPHIC DIVISION						
Northeast	0.2350		0.1574		0.1175	
New England	0.6493		0.6256		0.6746	
Middle Atlantic	0.1466		0.0849		0.0560	
Midwest	0.9314		0.9612		0.7256	
East North Central	0.5183		0.4332		0.9496	
West North Central	0.3218		0.1187		0.4461	
South	0.8431		0.7540		0.2794	
South Atlantic	0.8095		0.6913		0.4683	
East South Central	0.7096		0.9282		0.5018	
West South Central	0.8890		0.9070		0.6047	
West	0.3255		0.3584		0.8543	
Mountain	0.3660		0.2017		0.4023	
Pacific	0.5210		0.7009		0.5780	
COUNTY TYPE						
Large Metro	0.7549		0.4703		0.9754	
Small Metro	0.9034		0.9784		0.3619	
250K – 1 Mil. Pop.	0.4080		0.3369		0.0987	
< 250K Pop.	0.3102		0.1506		0.3649	
Nonmetro	0.5268		0.0537		0.1254	
Urbanized	0.4449		0.0837		0.0759	
Less Urbanized	0.9924		0.3312		0.8883	
Completely Rural	0.6920		0.8379		0.2413	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.85C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	133	146	97	90	59	57
GEOGRAPHIC DIVISION						
Northeast	53	59	38	39	22	24
New England	28	30	25	20	10	12
Middle Atlantic	45	51	29	35	20	20
Midwest	52	62	36	42	20	28
East North Central	42	54	28	37	15	24
West North Central	29	30	23	20	13	13
South	82	88	61	58	34	34
South Atlantic	64	60	47	37	24	20
East South Central	30	36	20	24	14	16
West South Central	43	52	34	36	19	22
West	75	78	54	43	36	29
Mountain	32	31	25	20	14	12
Pacific	67	72	48	38	34	27
COUNTY TYPE						
Large Metro	127	132	82	77	49	49
Small Metro	91	101	60	54	31	26
250K – 1 Mil. Pop.	73	74	49	44	25	21
< 250K Pop.	60	70	35	33	19	16
Nonmetro	61	65	41	33	20	20
Urbanized	48	49	34	22	16	13
Less Urbanized	40	43	22	23	13	16
Completely Rural	13	15	9	5	3	2

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.85D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.38	0.42	0.28	0.26	0.17	0.16
GEOGRAPHIC DIVISION						
Northeast	0.86	0.95	0.62	0.64	0.36	0.39
New England	1.73	1.83	1.54	1.24	0.64	0.75
Middle Atlantic	0.99	1.14	0.64	0.78	0.44	0.45
Midwest	0.70	0.84	0.48	0.56	0.27	0.38
East North Central	0.82	1.05	0.54	0.73	0.30	0.48
West North Central	1.26	1.28	0.99	0.87	0.55	0.56
South	0.64	0.68	0.47	0.45	0.26	0.27
South Atlantic	0.98	0.91	0.71	0.56	0.37	0.30
East South Central	1.49	1.74	0.98	1.15	0.69	0.80
West South Central	1.02	1.21	0.79	0.84	0.44	0.51
West	0.89	0.93	0.64	0.51	0.43	0.35
Mountain	1.28	1.22	0.97	0.79	0.56	0.46
Pacific	1.15	1.24	0.82	0.65	0.57	0.47
COUNTY TYPE						
Large Metro	0.55	0.57	0.39	0.36	0.25	0.24
Small Metro	0.64	0.78	0.48	0.47	0.27	0.24
250K – 1 Mil. Pop.	0.84	0.89	0.62	0.60	0.35	0.30
< 250K Pop.	1.04	1.56	0.73	0.81	0.44	0.42
Nonmetro	0.86	0.94	0.67	0.57	0.37	0.37
Urbanized	1.09	1.51	1.05	0.80	0.57	0.51
Less Urbanized	1.42	1.20	0.87	0.88	0.54	0.63
Completely Rural	2.21	2.78	1.56	1.27	0.60	0.63

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 1.85P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2409		0.0124		0.0545	
GEOGRAPHIC DIVISION						
Northeast	0.4305		0.4500		0.2119	
New England	0.5535		0.3593		0.5421	
Middle Atlantic	0.5753		0.7974		0.0763	
Midwest	0.8119		0.8643		0.6802	
East North Central	0.2380		0.5611		0.3448	
West North Central	0.1412		0.5662		0.4810	
South	0.9065		0.6939		0.1948	
South Atlantic	0.6780		0.5036		0.1250	
East South Central	0.9444		0.2294		0.8899	
West South Central	0.4822		0.5414		0.5433	
West	0.0058		0.0001		0.1402	
Mountain	0.1455		0.0664		0.3519	
Pacific	0.0180		0.0006		0.2311	
COUNTY TYPE						
Large Metro	0.2490		0.0934		0.3838	
Small Metro	0.9265		0.1619		0.0252	
250K – 1 Mil. Pop.	0.9444		0.1646		0.0908	
< 250K Pop.	0.8209		0.6688		0.1600	
Nonmetro	0.3304		0.1838		0.5970	
Urbanized	0.2415		0.0650		0.4275	
Less Urbanized	0.4600		0.8597		0.8567	
Completely Rural	0.2376		0.4170		0.5077	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.86C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	616	520	315	228	196	162
GEOGRAPHIC DIVISION						
Northeast	199	206	92	100	62	71
New England	112	115	43	43	23	33
Middle Atlantic	163	174	82	92	58	63
Midwest	218	211	98	91	64	62
East North Central	169	177	84	80	58	56
West North Central	133	109	48	45	29	26
South	396	298	219	140	141	102
South Atlantic	313	225	190	104	114	82
East South Central	125	104	71	49	56	27
West South Central	194	161	87	87	62	53
West	358	284	178	124	108	85
Mountain	184	131	104	64	40	38
Pacific	309	254	143	111	100	78
COUNTY TYPE						
Large Metro	541	431	231	183	156	127
Small Metro	349	293	174	127	118	96
250K – 1 Mil. Pop.	284	244	136	102	102	78
< 250K Pop.	203	168	105	74	59	55
Nonmetro	245	175	154	83	51	42
Urbanized	180	121	123	65	32	34
Less Urbanized	159	139	88	51	36	23
Completely Rural	64	51	32	18	20	12

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.86D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.30	0.25	0.16	0.11	0.10	0.08
GEOGRAPHIC DIVISION						
Northeast	0.54	0.55	0.25	0.27	0.17	0.19
New England	1.15	1.17	0.44	0.44	0.23	0.34
Middle Atlantic	0.60	0.63	0.30	0.33	0.21	0.23
Midwest	0.50	0.48	0.23	0.21	0.15	0.14
East North Central	0.56	0.59	0.28	0.26	0.19	0.19
West North Central	1.00	0.81	0.36	0.34	0.22	0.20
South	0.53	0.39	0.29	0.18	0.19	0.13
South Atlantic	0.78	0.55	0.47	0.25	0.28	0.20
East South Central	1.05	0.86	0.59	0.40	0.47	0.22
West South Central	0.84	0.68	0.38	0.37	0.27	0.22
West	0.76	0.59	0.38	0.26	0.23	0.18
Mountain	1.29	0.90	0.73	0.44	0.28	0.26
Pacific	0.95	0.76	0.44	0.33	0.31	0.23
COUNTY TYPE						
Large Metro	0.44	0.35	0.20	0.16	0.14	0.11
Small Metro	0.52	0.45	0.28	0.21	0.19	0.16
250K – 1 Mil. Pop.	0.65	0.55	0.33	0.25	0.25	0.19
< 250K Pop.	0.92	0.77	0.51	0.37	0.30	0.27
Nonmetro	0.65	0.47	0.45	0.25	0.15	0.13
Urbanized	1.04	0.74	0.79	0.46	0.22	0.25
Less Urbanized	0.94	0.79	0.57	0.32	0.24	0.14
Completely Rural	1.56	1.24	0.86	0.50	0.56	0.34

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.86P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5751		0.1846		0.6333	
GEOGRAPHIC DIVISION						
Northeast	0.0021		0.2213		0.1911	
New England	0.0198		0.2544		0.0852	
Middle Atlantic	0.0336		0.4304		0.5274	
Midwest	0.2993		0.8535		0.9720	
East North Central	0.2364		0.8679		0.9463	
West North Central	0.8849		0.5004		0.8245	
South	0.4001		0.0854		0.2363	
South Atlantic	0.1960		0.1637		0.7601	
East South Central	0.8281		0.1458		0.0721	
West South Central	0.6429		0.8658		0.5249	
West	0.4591		0.2277		0.7575	
Mountain	0.1486		0.2892		0.4223	
Pacific	0.9762		0.4659		0.4946	
COUNTY TYPE						
Large Metro	0.7129		0.3182		0.6053	
Small Metro	0.4465		0.6970		0.8298	
250K – 1 Mil. Pop.	0.6474		0.7131		0.5668	
< 250K Pop.	0.4910		0.8730		0.5981	
Nonmetro	0.6217		0.3587		0.8947	
Urbanized	0.6260		1.0000		0.0606	
Less Urbanized	0.8771		0.2690		0.2946	
Completely Rural	0.9023		0.4205		0.2252	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.87C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	636	554	339	246	205	170
GEOGRAPHIC DIVISION						
Northeast	203	216	100	109	66	74
New England	115	121	51	49	25	35
Middle Atlantic	167	184	86	100	61	66
Midwest	223	228	109	103	67	69
East North Central	174	192	92	91	60	62
West North Central	139	115	56	51	31	29
South	401	313	233	153	148	104
South Atlantic	319	234	199	108	119	83
East South Central	128	111	76	54	59	29
West South Central	195	169	97	96	65	56
West	369	311	187	132	111	90
Mountain	188	134	111	65	45	40
Pacific	319	284	150	120	101	82
COUNTY TYPE						
Large Metro	564	469	253	200	164	135
Small Metro	370	323	187	142	122	101
250K – 1 Mil. Pop.	306	265	149	114	108	83
< 250K Pop.	218	195	110	84	59	59
Nonmetro	260	193	159	92	55	47
Urbanized	191	137	128	70	35	36
Less Urbanized	173	153	91	59	39	28
Completely Rural	67	59	33	18	20	12

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.87D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.27	0.23	0.14	0.10	0.09	0.07
GEOGRAPHIC DIVISION						
Northeast	0.47	0.50	0.23	0.25	0.15	0.17
New England	1.01	1.06	0.45	0.42	0.22	0.31
Middle Atlantic	0.52	0.57	0.27	0.31	0.19	0.21
Midwest	0.44	0.45	0.21	0.20	0.13	0.14
East North Central	0.49	0.54	0.26	0.26	0.17	0.17
West North Central	0.89	0.73	0.36	0.32	0.20	0.18
South	0.45	0.35	0.26	0.17	0.17	0.12
South Atlantic	0.68	0.49	0.43	0.23	0.25	0.18
East South Central	0.92	0.79	0.54	0.38	0.42	0.20
West South Central	0.71	0.61	0.35	0.34	0.24	0.20
West	0.67	0.55	0.34	0.23	0.20	0.16
Mountain	1.12	0.78	0.66	0.38	0.27	0.23
Pacific	0.83	0.73	0.39	0.31	0.26	0.21
COUNTY TYPE						
Large Metro	0.38	0.32	0.19	0.14	0.13	0.10
Small Metro	0.46	0.41	0.25	0.19	0.17	0.14
250K – 1 Mil. Pop.	0.58	0.50	0.30	0.23	0.22	0.17
< 250K Pop.	0.79	0.70	0.44	0.33	0.25	0.24
Nonmetro	0.59	0.43	0.39	0.23	0.14	0.12
Urbanized	0.91	0.67	0.68	0.41	0.20	0.22
Less Urbanized	0.86	0.71	0.51	0.31	0.22	0.15
Completely Rural	1.39	1.23	0.75	0.46	0.49	0.30

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.87P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.8108		0.0403		0.2866	
GEOGRAPHIC DIVISION						
Northeast	0.0015		0.4049		0.4093	
New England	0.0168		0.5723		0.0688	
Middle Atlantic	0.0286		0.5275		0.9732	
Midwest	0.2968		0.8260		0.8601	
East North Central	0.1541		0.9596		0.8047	
West North Central	0.8407		0.7244		0.9212	
South	0.3883		0.0819		0.1347	
South Atlantic	0.2383		0.1286		0.5272	
East South Central	0.8177		0.3457		0.0977	
West South Central	0.7949		0.6987		0.4153	
West	0.1825		0.0202		0.4294	
Mountain	0.0893		0.1432		0.6563	
Pacific	0.5579		0.0728		0.2843	
COUNTY TYPE						
Large Metro	0.9559		0.1279		0.4262	
Small Metro	0.4612		0.4043		0.4408	
250K – 1 Mil. Pop.	0.6758		0.4257		0.3444	
< 250K Pop.	0.4881		0.7509		0.9250	
Nonmetro	0.5067		0.2295		0.9627	
Urbanized	0.4990		0.6881		0.1261	
Less Urbanized	0.7737		0.3204		0.3180	
Completely Rural	0.7746		0.2768		0.2645	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.88C Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug or Method of Administration	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
HALLUCINOGENS	707	636	35	37	133	139	678	601
PCP (Angel Dust, Phencyclidine)	346	274	8	11	30	31	342	273
LSD (Acid)	580	508	19	27	81	92	567	491
Peyote	328	267	9	9	33	27	327	264
Mescaline	378	312	6	6	35	28	375	309
Psilocybin (Mushrooms)	561	488	22	24	100	106	542	464
Ecstasy (MDMA)	464	384	24	27	114	120	442	345
INHALANTS	519	428	52	56	85	93	505	412
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	364	294	18	20	40	40	357	291
Correction Fluid, Degreaser, or Cleaning Fluid	100	115	26	25	26	27	91	107
Gasoline or Lighter Fluid	164	143	28	31	29	34	156	136
Glue, Shoe Polish, or Toluene	175	154	31	35	30	31	168	149
Halothane, Ether, or Other Anesthetics	85	89	10	9	17	17	84	88
Lacquer Thinner or Other Paint Solvents	117	96	18	22	21	24	114	93
Lighter Gases (Butane, Propane)	80	60	15	16	17	18	77	56
Nitrous Oxide or Whippits	409	324	17	22	70	77	399	313
Spray Paints	121	106	24	32	28	24	115	100
Other Aerosol Sprays	130	110	21	26	39	38	121	102
NEEDLE USE^{1,2,3}	265	198	10	11	40	42	261	194
Heroin Needle Use ^{1,2}	211	146	6	7	33	34	209	141
Cocaine Needle Use ^{1,2}	193	154	1	6	21	22	192	152
Stimulant Needle Use ^{1,2,3}	199	141	5	9	26	28	196	137
Methamphetamine Needle Use ^{1,2,3}	185	124	5	9	22	25	182	121
HEROIN	281	213	9	9	44	48	278	207
Smoke Heroin ²	152	120	5	8	32	32	150	115
Sniff or Snort Heroin ²	215	184	5	5	35	42	212	179

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

³ Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.88D Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Percentages, 2013 and 2014

Drug or Method of Administration	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
HALLUCINOGENS	0.27	0.24	0.14	0.15	0.38	0.40	0.33	0.29
PCP (Angel Dust, Phencyclidine)	0.13	0.10	0.03	0.05	0.09	0.09	0.17	0.13
LSD (Acid)	0.22	0.19	0.08	0.11	0.23	0.26	0.28	0.24
Peyote	0.13	0.10	0.03	0.04	0.09	0.08	0.16	0.13
Mescaline	0.14	0.12	0.02	0.02	0.10	0.08	0.18	0.15
Psilocybin (Mushrooms)	0.21	0.18	0.09	0.10	0.29	0.30	0.27	0.23
Ecstasy (MDMA)	0.18	0.14	0.10	0.11	0.33	0.34	0.22	0.17
INHALANTS	0.20	0.16	0.21	0.23	0.24	0.27	0.25	0.20
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	0.14	0.11	0.07	0.08	0.12	0.12	0.18	0.14
Correction Fluid, Degreaser, or Cleaning Fluid	0.04	0.04	0.11	0.10	0.08	0.08	0.05	0.05
Gasoline or Lighter Fluid	0.06	0.05	0.11	0.13	0.08	0.10	0.08	0.07
Glue, Shoe Polish, or Toluene	0.07	0.06	0.12	0.14	0.09	0.09	0.08	0.07
Halothane, Ether, or Other Anesthetics	0.03	0.03	0.04	0.04	0.05	0.05	0.04	0.04
Lacquer Thinner or Other Paint Solvents	0.04	0.04	0.07	0.09	0.06	0.07	0.06	0.05
Lighter Gases (Butane, Propane)	0.03	0.02	0.06	0.07	0.05	0.05	0.04	0.03
Nitrous Oxide or Whippits	0.16	0.12	0.07	0.09	0.20	0.22	0.20	0.15
Spray Paints	0.05	0.04	0.10	0.13	0.08	0.07	0.06	0.05
Other Aerosol Sprays	0.05	0.04	0.08	0.10	0.11	0.11	0.06	0.05
NEEDLE USE^{1,2,3}	0.10	0.07	0.04	0.05	0.11	0.12	0.13	0.09
Heroin Needle Use ^{1,2}	0.08	0.06	0.02	0.03	0.09	0.10	0.10	0.07
Cocaine Needle Use ^{1,2}	0.07	0.06	0.00	0.02	0.06	0.06	0.09	0.07
Stimulant Needle Use ^{1,2,3}	0.08	0.05	0.02	0.03	0.07	0.08	0.10	0.07
Methamphetamine Needle Use ^{1,2,3}	0.07	0.05	0.02	0.03	0.06	0.07	0.09	0.06
HEROIN	0.11	0.08	0.04	0.03	0.13	0.14	0.14	0.10
Smoke Heroin ²	0.06	0.05	0.02	0.03	0.09	0.09	0.07	0.06
Sniff or Snort Heroin ²	0.08	0.07	0.02	0.02	0.10	0.12	0.10	0.09

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

³ Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.88P Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug or Method of Administration	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
HALLUCINOGENS	0.5990		0.0681		0.0590		0.9291	
PCP (Angel Dust, Phencyclidine)	0.7509		0.2942		0.5934		0.7424	
LSD (Acid)	0.9928		0.0332		0.1116		0.6946	
Peyote	0.7501		0.8960		0.0040		0.9601	
Mescaline	0.6186		0.6295		0.0294		0.7435	
Psilocybin (Mushrooms)	0.3915		0.2853		0.1281		0.5719	
Ecstasy (MDMA)	0.4555		0.0462		0.1242		0.8254	
INHALANTS	0.9936		0.7986		0.1541		0.7675	
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	0.8626		0.4645		0.1594		0.9449	
Correction Fluid, Degreaser, or Cleaning Fluid	0.3950		0.5588		0.9494		0.2832	
Gasoline or Lighter Fluid	0.7513		0.5024		0.2438		0.9669	
Glue, Shoe Polish, or Toluene	0.8693		0.3997		0.2672		0.9034	
Halothane, Ether, or Other Anesthetics	0.1732		0.3282		0.9696		0.1392	
Lacquer Thinner or Other Paint Solvents	1.0000		0.3107		0.6054		0.7631	
Lighter Gases (Butane, Propane)	0.2617		0.7371		0.7821		0.1848	
Nitrous Oxide or Whippits	0.7393		0.7128		0.4104		0.6447	
Spray Paints	0.6559		0.0338		0.0463		0.5985	
Other Aerosol Sprays	0.9734		0.6157		0.3510		0.7886	
NEEDLE USE^{1,2,3}	0.4861		0.9400		0.8821		0.4817	
Heroin Needle Use ^{1,2}	0.3047		0.7031		0.6206		0.3269	
Cocaine Needle Use ^{1,2}	0.8901		0.0648		0.7808		0.8668	
Stimulant Needle Use ^{1,2,3}	0.4459		0.3534		0.8540		0.3903	
Methamphetamine Needle Use ^{1,2,3}	0.6156		0.3293		0.7097		0.5326	
HEROIN	0.8887		0.8141		0.3514		0.7454	
Smoke Heroin ²	0.8986		0.3744		0.5643		0.7517	
Sniff or Snort Heroin ²	0.4615		0.4737		0.1796		0.6500	

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

³ Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.89C Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Pain Reliever	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	453	392	40	41	79	84	436	380
Percocet [®] , Percodan [®] , or Tylox [®]	408	358	21	28	93	96	393	345
Vicodin [®] , Lortab [®] , or Lorcet [®]	481	429	36	41	118	126	461	399
Codeine	293	228	24	29	76	75	284	211
Demerol [®]	196	146	8	11	25	28	195	142
Dilaudid [®]	144	116	5	8	31	34	140	108
Fioricet [®]	94	56	2	7	14	13	93	54
Fiorinal [®]	85	41	4	7	10	8	84	40
Hydrocodone	359	274	29	28	89	96	344	249
Methadone	159	120	11	18	45	41	153	108
Morphine	200	155	17	21	57	56	192	141
OxyContin [®]	280	220	22	24	74	78	268	201
Phenaphen [®] with Codeine	81	69	7	9	23	23	78	65
Propoxyphene	42	48	4	6	13	11	40	46
SK-65 [®]	17	28	7	5	7	12	14	25
Stadol [®]	43	24	2	4	4	8	43	23
Talacen [®]	17	17	4	3	7	6	15	15
Talwin [®]	60	48	3	3	3	5	60	48
Talwin [®] NX	18	18	3	3	9	6	15	17
Tramadol	192	134	11	17	52	51	186	119
Ultram [®]	145	88	6	7	29	23	142	86
SELECTED GROUPS OF DRUGS¹								
Propoxyphene or Codeine Products ²	479	415	43	46	95	100	460	398
Oxycodone Products ³	443	379	28	33	100	105	427	361
Hydrocodone Products ⁴	508	445	41	46	123	131	488	414
Tramadol Products ⁵	210	144	12	18	52	52	204	131

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.89D Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Percentages, 2013 and 2014

Pain Reliever	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	0.17	0.15	0.16	0.16	0.23	0.24	0.21	0.18
Percocet [®] , Percodan [®] , or Tylox [®]	0.16	0.13	0.09	0.11	0.27	0.27	0.19	0.17
Vicodin [®] , Lortab [®] , or Lorcet [®]	0.18	0.16	0.14	0.17	0.34	0.36	0.23	0.19
Codeine	0.11	0.09	0.09	0.12	0.22	0.21	0.14	0.10
Demerol [®]	0.07	0.05	0.03	0.05	0.07	0.08	0.10	0.07
Dilaudid [®]	0.06	0.04	0.02	0.03	0.09	0.10	0.07	0.05
Fioricet [®]	0.04	0.02	0.01	0.03	0.04	0.04	0.05	0.03
Fiorinal [®]	0.03	0.02	0.02	0.03	0.03	0.02	0.04	0.02
Hydrocodone	0.14	0.10	0.12	0.11	0.26	0.27	0.17	0.12
Methadone	0.06	0.05	0.04	0.07	0.13	0.12	0.08	0.05
Morphine	0.08	0.06	0.07	0.08	0.16	0.16	0.09	0.07
OxyContin [®]	0.11	0.08	0.09	0.10	0.21	0.22	0.13	0.10
Phenaphen [®] with Codeine	0.03	0.03	0.03	0.04	0.07	0.06	0.04	0.03
Propoxyphene	0.02	0.02	0.02	0.03	0.04	0.03	0.02	0.02
SK-65 [®]	0.01	0.01	0.03	0.02	0.02	0.04	0.01	0.01
Stadol [®]	0.02	0.01	0.01	0.02	0.01	0.02	0.02	0.01
Talacen [®]	0.01	0.01	0.02	0.01	0.02	0.02	0.01	0.01
Talwin [®]	0.02	0.02	0.01	0.01	0.01	0.01	0.03	0.02
Talwin [®] NX	0.01	0.01	0.01	0.01	0.03	0.02	0.01	0.01
Tramadol	0.07	0.05	0.04	0.07	0.15	0.15	0.09	0.06
Ultram [®]	0.06	0.03	0.03	0.03	0.08	0.07	0.07	0.04
SELECTED GROUPS OF DRUGS¹								
Propoxyphene or Codeine Products ²	0.18	0.16	0.17	0.19	0.27	0.29	0.23	0.19
Oxycodone Products ³	0.17	0.14	0.11	0.13	0.29	0.30	0.21	0.18
Hydrocodone Products ⁴	0.19	0.17	0.17	0.18	0.35	0.38	0.24	0.20
Tramadol Products ⁵	0.08	0.05	0.05	0.07	0.15	0.15	0.10	0.06

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.89P Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Pain Reliever	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	0.8527		0.1273		0.5128		0.8769	
Percocet [®] , Percodan [®] , or Tylox [®]	0.0668		0.6275		0.3106		0.0348	
Vicodin [®] , Lortab [®] , or Lorcet [®]	0.4669		0.1940		0.0110		0.9511	
Codeine	0.2127		0.3476		0.2321		0.2973	
Demerol [®]	0.3676		0.3058		0.5303		0.2809	
Dilaudid [®]	0.5817		0.2397		0.9934		0.6110	
Fioricet [®]	0.3985		0.0763		0.7973		0.3501	
Fiorinal [®]	0.0836		0.6873		0.9704		0.0750	
Hydrocodone	0.2939		0.3434		0.4934		0.4153	
Methadone	0.1887		0.0158		0.0255		0.3498	
Morphine	0.5506		0.6362		0.6897		0.5847	
OxyContin [®]	0.9676		0.5099		0.2930		0.7152	
Phenaphen [®] with Codeine	0.2889		0.6700		0.3450		0.3881	
Propoxyphene	0.2558		0.4392		0.3599		0.1741	
SK-65 [®]	0.2432		0.4805		0.1286		0.4073	
Stadol [®]	0.3156		0.1461		0.2453		0.1730	
Talacen [®]	0.3808		0.7080		0.9073		0.3082	
Talwin [®]	0.9567		0.9476		0.4834		0.9071	
Talwin [®] NX	0.8025		0.6891		0.9602		0.7051	
Tramadol	0.4206		0.3082		0.3846		0.5122	
Ultram [®]	0.0441		0.6322		0.0906		0.0964	
SELECTED GROUPS OF DRUGS¹								
Propoxyphene or Codeine Products ²	0.9862		0.1566		0.2847		0.9464	
Oxycodone Products ³	0.1398		0.5910		0.3106		0.0666	
Hydrocodone Products ⁴	0.6204		0.2536		0.0348		0.8594	
Tramadol Products ⁵	0.2445		0.3848		0.2685		0.3360	

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.90C Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Tranquilizer	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Klonopin® or Clonazepam	293	226	16	18	68	74	284	205
Xanax®, Alprazolam, Ativan®, or Lorazepam	454	349	28	39	102	111	435	324
Valium® or Diazepam	443	346	14	20	70	71	433	337
Atarax®	64	40	4	6	10	9	63	38
BuSpar®	73	67	4	6	23	22	69	64
Equanil®	31	19	2	7	9	5	30	17
Flexeril®	189	146	8	10	37	36	186	139
Librium®	106	66	4	7	13	10	105	65
Limbitrol®	13	11	4	4	6	4	11	9
Meprobamate	46	45	3	7	6	8	45	44
Miltown®	23	35	3	2	8	5	22	35
Rohypnol®	71	35	4	3	15	20	70	29
Serax®	18	19	2	6	5	5	17	17
Soma®	196	130	12	11	46	45	187	119
Tranxene®	20	30	2	*	7	6	19	30
Vistaril®	56	46	4	7	17	14	53	42
SELECTED GROUPS OF DRUGS¹								
Benzodiazepines ²	564	435	30	41	107	121	546	418
Meprobamate Products ³	57	56	3	9	10	10	56	55
Muscle Relaxants ⁴	245	181	14	15	53	52	237	167

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan® or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

³ Includes Equanil®, meprobamate, and Miltown®.

⁴ Includes Flexeril® and Soma®.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.90D Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Percentages, 2013 and 2014

Tranquilizer	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Klonopin® or Clonazepam	0.11	0.09	0.06	0.07	0.20	0.21	0.14	0.10
Xanax®, Alprazolam, Ativan®, or Lorazepam	0.17	0.13	0.11	0.16	0.29	0.32	0.21	0.16
Valium® or Diazepam	0.17	0.13	0.06	0.08	0.20	0.20	0.21	0.16
Atarax®	0.02	0.01	0.02	0.02	0.03	0.03	0.03	0.02
BuSpar®	0.03	0.03	0.02	0.02	0.07	0.06	0.03	0.03
Equanil®	0.01	0.01	0.01	0.03	0.03	0.01	0.01	0.01
Flexeril®	0.07	0.06	0.03	0.04	0.11	0.10	0.09	0.07
Librium®	0.04	0.03	0.02	0.03	0.04	0.03	0.05	0.03
Limbitrol®	0.00	0.00	0.02	0.02	0.02	0.01	0.01	0.00
Meprobamate	0.02	0.02	0.01	0.03	0.02	0.02	0.02	0.02
Miltown®	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.02
Rohypnol®	0.03	0.01	0.02	0.01	0.04	0.06	0.03	0.01
Serax®	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.01
Soma®	0.07	0.05	0.05	0.04	0.13	0.13	0.09	0.06
Tranxene®	0.01	0.01	0.01	*	0.02	0.02	0.01	0.01
Vistaril®	0.02	0.02	0.02	0.03	0.05	0.04	0.03	0.02
SELECTED GROUPS OF DRUGS¹								
Benzodiazepines ²	0.22	0.16	0.12	0.16	0.31	0.35	0.27	0.20
Meprobamate Products ³	0.02	0.02	0.01	0.04	0.03	0.03	0.03	0.03
Muscle Relaxants ⁴	0.09	0.07	0.06	0.06	0.15	0.15	0.12	0.08

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan® or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

³ Includes Equanil®, meprobamate, and Miltown®.

⁴ Includes Flexeril® and Soma®.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.90P Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Tranquilizer	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Klonopin® or Clonazepam	0.3057		0.9767		0.9924		0.2810	
Xanax®, Alprazolam, Ativan®, or Lorazepam	0.3689		0.0107		0.6127		0.5610	
Valium® or Diazepam	0.1192		0.4256		0.9431		0.1223	
Atarax®	0.8419		0.9657		0.5539		0.7521	
BuSpar®	0.0852		0.7986		0.7824		0.0874	
Equanil®	0.5330		0.2043		0.1856		0.6039	
Flexeril®	0.5117		0.3153		0.4569		0.4403	
Librium®	0.0993		0.5135		0.1592		0.1250	
Limbitrol®	0.5610		0.3998		0.9241		0.7626	
Meprobamate	0.2123		0.2413		0.2284		0.3493	
Miltown®	0.6588		0.5671		0.3857		0.4907	
Rohypnol®	0.0685		0.2154		0.8384		0.0582	
Serax®	0.9328		0.0669		0.9197		0.7267	
Soma®	0.1208		0.0832		0.0163		0.4075	
Tranxene®	0.1691		*		0.8214		0.1378	
Vistaril®	0.6462		0.2524		0.6524		0.6117	
SELECTED GROUPS OF DRUGS¹								
Benzodiazepines ²	0.2057		0.0250		0.4950		0.3322	
Meprobamate Products ³	0.3159		0.1257		0.9001		0.3885	
Muscle Relaxants ⁴	0.3887		0.7024		0.0489		0.6966	

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan® or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

³ Includes Equanil®, meprobamate, and Miltown®.

⁴ Includes Flexeril® and Soma®.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.91C Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Stimulant	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Methamphetamine, Desoxyn®, or Methedrine ¹	393	363	14	22	57	58	386	356
Prescription Diet Pills ²	326	270	12	18	41	48	326	264
Ritalin® or Methylphenidate	265	219	19	22	74	74	255	200
Cylert®	50	56	4	8	5	9	50	55
Dexedrine®	189	155	4	6	16	18	188	154
Dextroamphetamine	128	111	10	8	23	28	125	106
Didrex®	22	40	2	4	9	4	20	40
Eskatrol®	28	42	1	3	5	4	28	42
Ionamin®	69	50	6	6	4	5	69	49
Mazanor®	8	21	2	9	2	2	8	19
Obedrin-LA®	6	8	2	3	5	2	*	7
Plegine®	12	9	3	9	2	*	11	*
Preludin®	51	59	*	6	7	3	50	59
Sanorex®	*	29	1	8	2	5	*	27
Tenuate®	34	37	2	8	5	6	34	36
SELECTED GROUPS OF DRUGS³								
Amphetamine, Dextroamphetamine, and Phentermine Products ⁴	363	294	20	23	71	70	357	285
Mazindol Products ⁵	9	30	3	10	3	6	8	28
Methylphenidate or Dexmethylphenidate Products ⁶	267	220	20	22	74	74	257	201

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn®, or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine®, Biphedamine®, Fastin®, or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine®, dextroamphetamine, and Ionamin®.

⁵ Includes Mazanor® and Sanorex®.

⁶ Includes Ritalin®, other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.91D Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Percentages, 2013 and 2014

Stimulant	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Methamphetamine, Desoxyn®, or Methedrine ¹	0.15	0.14	0.06	0.09	0.17	0.17	0.19	0.17
Prescription Diet Pills ²	0.12	0.10	0.05	0.07	0.12	0.14	0.16	0.13
Ritalin® or Methylphenidate	0.10	0.08	0.08	0.09	0.21	0.21	0.13	0.10
Cylert®	0.02	0.02	0.01	0.03	0.01	0.03	0.02	0.03
Dexedrine®	0.07	0.06	0.02	0.02	0.05	0.05	0.09	0.07
Dextroamphetamine	0.05	0.04	0.04	0.03	0.07	0.08	0.06	0.05
Didrex®	0.01	0.02	0.01	0.02	0.03	0.01	0.01	0.02
Eskatrol®	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.02
Ionamin®	0.03	0.02	0.02	0.02	0.01	0.01	0.03	0.02
Mazanor®	0.00	0.01	0.01	0.04	0.00	0.01	0.00	0.01
Obedrin-LA®	0.00	0.00	0.01	0.01	0.01	0.01	*	0.00
Plegine®	0.00	0.00	0.01	0.04	0.00	*	0.01	*
Preludin®	0.02	0.02	*	0.02	0.02	0.01	0.02	0.03
Sanorex®	*	0.01	0.01	0.03	0.01	0.02	*	0.01
Tenuate®	0.01	0.01	0.01	0.03	0.01	0.02	0.02	0.02
SELECTED GROUPS OF DRUGS³								
Amphetamine, Dextroamphetamine, and Phentermine Products ⁴	0.14	0.11	0.08	0.09	0.20	0.20	0.18	0.14
Mazindol Products ⁵	0.00	0.01	0.01	0.04	0.01	0.02	0.00	0.01
Methylphenidate or Dexmethylphenidate Products ⁶	0.10	0.08	0.08	0.09	0.21	0.21	0.13	0.10

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn®, or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine®, Biphedamine®, Fastin®, or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine®, dextroamphetamine, and Ionamin®.

⁵ Includes Mazanor® and Sanorex®.

⁶ Includes Ritalin®, other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.91P Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Stimulant	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Methamphetamine, Desoxyn®, or Methedrine ¹	0.2996		0.0428		0.9414		0.3517	
Prescription Diet Pills ²	0.3008		0.1609		0.0472		0.5167	
Ritalin® or Methylphenidate	0.5993		0.8323		0.0704		0.2439	
Cylert®	0.9320		0.2053		0.1622		0.8072	
Dexedrine®	0.9551		0.5744		0.1636		0.9076	
Dextroamphetamine	0.1998		0.9470		0.0254		0.4107	
Didrex®	0.0569		0.2296		0.2001		0.0340	
Eskatrol®	0.2284		0.3708		0.4713		0.2104	
Ionamin®	0.2983		0.8638		0.6522		0.3020	
Mazanor®	0.1653		0.0674		1.0000		0.4750	
Obedrin-LA®	0.7682		0.6200		0.2058		*	
Plegine®	0.9952		0.0195		*		*	
Preludin®	0.1993		*		0.2807		0.2067	
Sanorex®	*		0.0066		0.2719		*	
Tenuate®	0.3523		0.0457		0.9085		0.5504	
SELECTED GROUPS OF DRUGS³								
Amphetamine, Dextroamphetamine, and Phentermine Products ⁴	0.3374		0.4224		0.4780		0.4513	
Mazindol Products ⁵	0.0004		0.0077		0.2351		0.0086	
Methylphenidate or Dexmethylphenidate Products ⁶	0.6465		0.6973		0.0437		0.2411	

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn®, or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine®, Biphedamine®, Fastin®, or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine®, dextroamphetamine, and Ionamin®.

⁵ Includes Mazanor® and Sanorex®.

⁶ Includes Ritalin®, other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.92C Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Sedative	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Methaqualone, Sopor [®] , or Quaalude [®]	312	271	6	10	14	14	312	269
Barbiturates ¹	198	175	4	10	17	12	198	173
Restoril [®] or Temazepam	125	94	8	10	22	20	121	91
Amytal [®]	36	42	4	5	8	4	35	42
Butisol [®]	9	26	2	5	5	4	7	25
Chloral Hydrate	39	41	3	5	6	*	38	40
Dalmane [®]	64	45	2	5	8	2	64	44
Halcion [®]	93	54	3	5	9	12	92	52
Phenobarbital	103	102	5	7	17	12	101	100
Placidyl [®]	109	83	2	5	5	2	109	83
Tuinal [®]	121	89	2	4	4	*	121	89
SELECTED GROUPS OF DRUGS²								
Temazepam, Flurazepam, or Triazolam ³	162	113	9	12	24	23	158	109
Any Barbiturates ⁴	217	193	7	13	21	16	216	190

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.92D Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Percentages, 2013 and 2014

Sedative	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Methaqualone, Sopor [®] , or Quaalude [®]	0.12	0.10	0.02	0.04	0.04	0.04	0.15	0.13
Barbiturates ¹	0.08	0.07	0.01	0.04	0.05	0.04	0.10	0.08
Restoril [®] or Temazepam	0.05	0.04	0.03	0.04	0.06	0.06	0.06	0.04
Amytal [®]	0.01	0.02	0.01	0.02	0.02	0.01	0.02	0.02
Butisol [®]	0.00	0.01	0.01	0.02	0.01	0.01	0.00	0.01
Chloral Hydrate	0.01	0.02	0.01	0.02	0.02	*	0.02	0.02
Dalmane [®]	0.02	0.02	0.01	0.02	0.02	0.01	0.03	0.02
Halcion [®]	0.04	0.02	0.01	0.02	0.03	0.03	0.05	0.03
Phenobarbital	0.04	0.04	0.02	0.03	0.05	0.03	0.05	0.05
Placidyl [®]	0.04	0.03	0.01	0.02	0.01	0.01	0.05	0.04
Tuinal [®]	0.05	0.03	0.01	0.01	0.01	*	0.06	0.04
SELECTED GROUPS OF DRUGS²								
Temazepam, Flurazepam, or Triazolam ³	0.06	0.04	0.04	0.05	0.07	0.07	0.08	0.05
Any Barbiturates ⁴	0.08	0.07	0.03	0.05	0.06	0.05	0.11	0.09

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 1.92P Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Sedative	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
Methaqualone, Sopor [®] , or Quaalude [®]	0.9134		0.2017		0.3834		0.9293	
Barbiturates ¹	0.7864		0.0255		0.3599		0.8094	
Restoril [®] or Temazepam	0.4504		0.6438		0.7674		0.4983	
Amytal [®]	0.2577		0.2375		0.0712		0.1871	
Butisol [®]	0.0122		0.2384		0.7883		0.0139	
Chloral Hydrate	0.5691		0.4452		*		0.4989	
Dalmane [®]	0.2886		0.1212		0.0976		0.3104	
Halcion [®]	0.2627		0.2718		0.5459		0.1961	
Phenobarbital	0.6593		0.4401		0.5884		0.6380	
Placidyl [®]	0.8816		0.1639		0.4121		0.8610	
Tuinal [®]	0.6968		0.7674		*		0.6984	
SELECTED GROUPS OF DRUGS²								
Temazepam, Flurazepam, or Triazolam ³	0.8547		0.3167		0.7568		0.7440	
Any Barbiturates ⁴	0.4453		0.0398		0.3454		0.4616	

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.1A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	175,260	175,416	80,846	81,141	66,879	66,899
Cigarettes	162,138	161,789	66,256	65,858	55,778	55,241
Smokeless Tobacco	46,139	45,290	11,866	11,559	8,813	8,662
Cigars	89,117	89,267	24,933	23,734	12,438	11,964
Pipe Tobacco ²	34,173	33,695	--	--	2,271	2,247
ALCOHOL	213,794 ^b	217,765	173,966 ^a	176,556	136,868 ^a	139,677
Binge Alcohol Use ³	--	--	--	--	60,064	60,941
Heavy Alcohol Use ³	--	--	--	--	16,541	16,342

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.1B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	66.8	66.2	30.8	30.6	25.5	25.2
Cigarettes	61.8	61.0	25.3	24.8	21.3	20.8
Smokeless Tobacco	17.6	17.1	4.5	4.4	3.4	3.3
Cigars	34.0	33.7	9.5 ^a	9.0	4.7	4.5
Pipe Tobacco ²	13.0	12.7	--	--	0.9	0.8
ALCOHOL	81.5	82.1	66.3	66.6	52.2	52.7
Binge Alcohol Use ³	--	--	--	--	22.9	23.0
Heavy Alcohol Use ³	--	--	--	--	6.3	6.2

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.2A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	454	430	263	217	107	88
Cigarettes	317	291	184	136	71	58
Smokeless Tobacco	136	118	75	57	25	25
Cigars	130	120	83	77	18	20
Pipe Tobacco ²	50	47	--	--	17	15
ALCOHOL	730	649	470	443	167	168
Binge Alcohol Use ³	--	--	--	--	67	64
Heavy Alcohol Use ³	--	--	--	--	10	8

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.2B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	5.7	5.4	3.3	2.7	1.3	1.1
Cigarettes	4.0	3.7	2.3	1.7	0.9	0.7
Smokeless Tobacco	1.7	1.5	0.9	0.7	0.3	0.3
Cigars	1.6	1.5	1.0	1.0	0.2	0.3
Pipe Tobacco ²	0.6	0.6	--	--	0.2	0.2
ALCOHOL	9.2	8.2	5.9	5.6	2.1	2.1
Binge Alcohol Use ³	--	--	--	--	0.8	0.8
Heavy Alcohol Use ³	--	--	--	--	0.1	0.1

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.3A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	1,577 ^b	1,338	1,067 ^b	893	544 ^a	433
Cigarettes	1,224 ^b	1,015	774 ^b	607	367 ^a	288
Smokeless Tobacco	456	374	319	275	154	131
Cigars	540	522	360	364	122	124
Pipe Tobacco ²	165	131	--	--	47	43
ALCOHOL	2,427	2,320	1,896	1,775	806	722
Binge Alcohol Use ³	--	--	--	--	385	330
Heavy Alcohol Use ³	--	--	--	--	57	44

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.3B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	18.5 ^b	15.8	12.5 ^b	10.6	6.4 ^a	5.1
Cigarettes	14.4 ^b	12.0	9.1 ^b	7.2	4.3 ^a	3.4
Smokeless Tobacco	5.4	4.4	3.7	3.3	1.8	1.6
Cigars	6.3	6.2	4.2	4.3	1.4	1.5
Pipe Tobacco ²	1.9	1.6	--	--	0.6	0.5
ALCOHOL	28.5	27.4	22.3	21.0	9.5	8.5
Binge Alcohol Use ³	--	--	--	--	4.5	3.9
Heavy Alcohol Use ³	--	--	--	--	0.7	0.5

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.4A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	2,891	2,823	2,131	2,052	1,300	1,222
Cigarettes	2,362	2,217	1,603	1,466	959	867
Smokeless Tobacco	908	922	617	609	309	334
Cigars	1,400	1,343	981	984	436	376
Pipe Tobacco ²	303	376	--	--	96	121
ALCOHOL	4,512	4,406	3,766	3,761	1,903	1,983
Binge Alcohol Use ³	--	--	--	--	1,104	1,118
Heavy Alcohol Use ³	--	--	--	--	226	205

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.4B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	34.4	33.2	25.4	24.1	15.5	14.4
Cigarettes	28.1 ^a	26.0	19.1 ^a	17.2	11.4	10.2
Smokeless Tobacco	10.8	10.8	7.3	7.2	3.7	3.9
Cigars	16.7	15.8	11.7	11.6	5.2	4.4
Pipe Tobacco ²	3.6	4.4	--	--	1.1	1.4
ALCOHOL	53.7	51.8	44.8	44.2	22.7	23.3
Binge Alcohol Use ³	--	--	--	--	13.1	13.1
Heavy Alcohol Use ³	--	--	--	--	2.7	2.4

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.5A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	7,478	7,239	5,952	5,677	4,479 ^a	4,062
Cigarettes	6,420	6,085	4,792	4,460	3,571 ^b	3,145
Smokeless Tobacco	2,394	2,158	1,393	1,248	849 ^a	706
Cigars	4,273	4,176	2,999	2,870	1,462	1,381
Pipe Tobacco ²	1,204	1,058	--	--	400 ^a	286
ALCOHOL	9,581	9,374	8,559	8,326	5,774	5,788
Binge Alcohol Use ³	--	--	--	--	3,843	3,739
Heavy Alcohol Use ³	--	--	--	--	1,120	1,043

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.5B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	56.7	55.2	45.1	43.3	33.9 ^b	31.0
Cigarettes	48.7	46.4	36.3 ^a	34.0	27.1 ^b	24.0
Smokeless Tobacco	18.1 ^a	16.5	10.6	9.5	6.4 ^a	5.4
Cigars	32.4	31.9	22.7	21.9	11.1	10.5
Pipe Tobacco ²	9.1	8.1	--	--	3.0 ^a	2.2
ALCOHOL	72.6	71.5	64.9	63.5	43.8	44.2
Binge Alcohol Use ³	--	--	--	--	29.1	28.5
Heavy Alcohol Use ³	--	--	--	--	8.5	8.0

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.6A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	15,172	15,051	10,645	10,406	8,403	8,182
Cigarettes	13,721	13,523	8,958	8,698	7,085	6,759
Smokeless Tobacco	4,671	4,617	1,839	1,842	1,170	1,257
Cigars	9,106	8,994	4,215	4,106	2,010	2,011
Pipe Tobacco ²	2,393	2,416	--	--	357	379
ALCOHOL	19,567	19,752	18,148	18,387	14,964	15,026
Binge Alcohol Use ³	--	--	--	--	9,343	9,446
Heavy Alcohol Use ³	--	--	--	--	2,819	2,739

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.6B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	70.3	69.0	49.3	47.7	38.9	37.5
Cigarettes	63.6	62.0	41.5	39.8	32.8 ^a	31.0
Smokeless Tobacco	21.6	21.2	8.5	8.4	5.4	5.8
Cigars	42.2	41.2	19.5	18.8	9.3	9.2
Pipe Tobacco ²	11.1	11.1	--	--	1.7	1.7
ALCOHOL	90.6	90.5	84.1	84.2	69.3	68.8
Binge Alcohol Use ³	--	--	--	--	43.3	43.3
Heavy Alcohol Use ³	--	--	--	--	13.1	12.5

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.7A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	27,926	27,558	17,361 ^b	16,166	14,316 ^b	13,176
Cigarettes	26,160	25,487	14,760 ^b	13,434	12,312 ^b	11,128
Smokeless Tobacco	7,987	7,969	2,374	2,366	1,736	1,759
Cigars	16,641	16,140	5,951 ^a	5,367	2,895	2,562
Pipe Tobacco ²	3,304	3,595	--	--	349	363
ALCOHOL	33,703 ^b	34,538	30,065	30,415	24,686	24,987
Binge Alcohol Use ³	--	--	--	--	13,983	13,463
Heavy Alcohol Use ³	--	--	--	--	4,058 ^a	3,367

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.7B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	74.8 ^a	72.7	46.5 ^b	42.7	38.3 ^b	34.8
Cigarettes	70.0 ^b	67.3	39.5 ^b	35.5	33.0 ^b	29.4
Smokeless Tobacco	21.4	21.0	6.4	6.2	4.6	4.6
Cigars	44.6	42.6	15.9 ^a	14.2	7.8	6.8
Pipe Tobacco ²	8.8	9.5	--	--	0.9	1.0
ALCOHOL	90.2	91.2	80.5	80.3	66.1	66.0
Binge Alcohol Use ³	--	--	--	--	37.4	35.5
Heavy Alcohol Use ³	--	--	--	--	10.9 ^b	8.9

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.8A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	119,762	120,977	43,427 ^a	45,730	37,730 ^a	39,737
Cigarettes	111,934	113,171	35,185 ^a	37,057	31,412	32,995
Smokeless Tobacco	29,587	29,132	5,251	5,163	4,572	4,450
Cigars	57,027	57,972	10,345	9,966	5,494	5,489
Pipe Tobacco ²	26,754	26,071	--	--	1,004	1,039
ALCOHOL	143,274 ^b	146,726	111,061 ^a	113,449	88,567 ^a	91,003
Binge Alcohol Use ³	--	--	--	--	31,339	32,782
Heavy Alcohol Use ³	--	--	--	--	8,252	8,936

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.8B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	72.4	72.3	26.3	27.3	22.8	23.7
Cigarettes	67.7	67.6	21.3	22.1	19.0	19.7
Smokeless Tobacco	17.9	17.4	3.2	3.1	2.8	2.7
Cigars	34.5	34.6	6.3	6.0	3.3	3.3
Pipe Tobacco ²	16.2	15.6	--	--	0.6	0.6
ALCOHOL	86.6	87.6	67.2	67.8	53.6	54.4
Binge Alcohol Use ³	--	--	--	--	19.0	19.6
Heavy Alcohol Use ³	--	--	--	--	5.0	5.3

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.9A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	94,449	94,796	47,773	48,527	39,549	39,952
Cigarettes	84,457	84,510	35,858	35,638	30,041	29,822
Smokeless Tobacco	38,921	38,512	10,806	10,718	8,211	8,205
Cigars	64,644	65,196	18,848	18,424	9,755	9,580
Pipe Tobacco ²	29,642	28,851	--	--	1,878	1,774
ALCOHOL	107,186 ^a	108,611	88,185	88,879	72,605	73,559
Binge Alcohol Use ³	--	--	--	--	38,379	38,586
Heavy Alcohol Use ³	--	--	--	--	12,093	11,938

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.9B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	74.3	73.8	37.6	37.8	31.1	31.1
Cigarettes	66.4	65.8	28.2	27.8	23.6	23.2
Smokeless Tobacco	30.6	30.0	8.5	8.3	6.5	6.4
Cigars	50.9	50.8	14.8	14.3	7.7	7.5
Pipe Tobacco ²	23.3	22.5	--	--	1.5	1.4
ALCOHOL	84.3	84.6	69.4	69.2	57.1	57.3
Binge Alcohol Use ³	--	--	--	--	30.2	30.0
Heavy Alcohol Use ³	--	--	--	--	9.5	9.3

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.10A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	80,810	80,620	33,073	32,614	27,331	26,947
Cigarettes	77,682	77,279	30,398	30,220	25,737	25,418
Smokeless Tobacco	7,218	6,777	1,060	842	603	457
Cigars	24,473	24,071	6,085 ^b	5,310	2,683	2,384
Pipe Tobacco ²	4,532	4,845	--	--	392	473
ALCOHOL	106,608 ^b	109,155	85,781 ^a	87,676	64,263	66,118
Binge Alcohol Use ³	--	--	--	--	21,685	22,355
Heavy Alcohol Use ³	--	--	--	--	4,448	4,404

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.10B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	59.7	59.0	24.4	23.9	20.2	19.7
Cigarettes	57.4	56.5	22.5	22.1	19.0	18.6
Smokeless Tobacco	5.3	5.0	0.8	0.6	0.4	0.3
Cigars	18.1	17.6	4.5 ^b	3.9	2.0	1.7
Pipe Tobacco ²	3.4	3.5	--	--	0.3	0.3
ALCOHOL	78.8	79.8	63.4	64.1	47.5	48.4
Binge Alcohol Use ³	--	--	--	--	16.0	16.4
Heavy Alcohol Use ³	--	--	--	--	3.3	3.2

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.11A Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	175,260	175,416	80,846	81,141	66,879	66,899
12	130	147	78	68	26	24
13	324	282	184	149	81	64
14	643 ^b	506	391	339	189	144
15	934	832	676 ^a	554	355	289
16	1,275	1,226	942	872	538	519
17	1,617	1,597	1,189	1,179	762	703
18	2,297	2,125	1,874	1,721	1,333	1,215
19	2,450	2,390	1,941	1,886	1,468 ^a	1,274
20	2,731	2,725	2,137	2,070	1,678	1,573
21	2,897	2,841	2,145	2,104	1,654	1,613
22	3,071	3,018	2,138	2,204	1,681	1,739
23	3,092	3,066	2,188	2,062	1,801	1,631
24	3,154	3,046	2,205	2,020	1,700	1,584
25	2,957	3,080	1,969	2,015	1,566	1,614
26-29	12,541	12,144	8,156	7,438	6,502	6,086
30-34	15,385	15,414	9,205	8,728	7,814 ^a	7,090
35-39	13,465	14,119	6,768	7,365	5,701	6,201
40-44	15,693 ^a	14,445	7,262	6,991	6,258	5,860
45-49	14,977	15,081	6,869	6,626	6,029	5,805
50-54	17,810	17,832	7,118	7,677	6,141	6,823
55-59	14,664	15,190	5,475	6,266	4,718 ^a	5,630
60-64	13,072	13,065	4,034	4,417	3,490	3,882
65 or Older	30,080	31,246	5,899	6,389	5,392	5,535

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.11B Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	66.8	66.2	30.8	30.6	25.5	25.2
12	3.5	3.9	2.1	1.8	0.7	0.6
13	7.7	6.9	4.4	3.6	1.9	1.6
14	14.7 ^a	12.1	8.9	8.1	4.3	3.5
15	22.6 ^a	19.4	16.4 ^b	12.9	8.6 ^a	6.7
16	29.8	28.3	22.0	20.1	12.6	12.0
17	39.2	38.3	28.8	28.3	18.5	16.8
18	48.5	46.1	39.6	37.4	28.2	26.4
19	58.8	57.7	46.6	45.6	35.2 ^a	30.8
20	63.6	62.5	49.8	47.5	39.1	36.1
21	68.4	65.4	50.6	48.4	39.0	37.1
22	68.6	66.9	47.8	48.8	37.6	38.5
23	70.6	69.8	50.0	46.9	41.1 ^a	37.1
24	71.8	70.4	50.2	46.7	38.7	36.6
25	71.9	72.5	47.9	47.4	38.1	38.0
26-29	72.9	71.3	47.4 ^a	43.7	37.8	35.7
30-34	76.4	73.9	45.7 ^b	41.9	38.8 ^b	34.0
35-39	70.7	72.2	35.5	37.7	29.9	31.7
40-44	73.1	71.6	33.8	34.6	29.1	29.0
45-49	74.9	73.4	34.4	32.3	30.2	28.3
50-54	77.2	75.1	30.8	32.3	26.6	28.7
55-59	71.4	72.8	26.6	30.0	23.0 ^a	27.0
60-64	73.5	74.5	22.7	25.2	19.6	22.1
65 or Older	69.2	69.4	13.6	14.2	12.4	12.3

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.12A Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	162,138	161,789	66,256	65,858	55,778	55,241
12	80	98	54	39	16	12
13	237	194	130	97	55	45
14	491 ^b	365	274	219	115	86
15	733	650	500 ^b	389	252	202
16	1,020	981	681	654	372	374
17	1,342	1,236	922	813	587 ^a	493
18	1,868	1,699	1,409	1,292	992	892
19	2,118	1,999	1,594	1,451	1,188 ^a	993
20	2,434	2,387	1,789	1,717	1,391	1,261
21	2,547	2,467	1,772	1,697	1,336	1,285
22	2,789	2,697	1,844	1,864	1,468	1,442
23	2,799	2,797	1,829	1,755	1,489	1,358
24	2,868	2,759	1,887	1,681	1,463	1,305
25	2,719	2,801	1,626	1,701	1,329	1,369
26-29	11,707	11,156	6,897 ^a	6,069	5,623	5,089
30-34	14,453	14,332	7,863	7,365	6,689 ^a	6,039
35-39	12,467	13,150	5,720	6,060	4,876	5,244
40-44	14,461 ^a	13,334	5,795	5,574	5,018	4,797
45-49	13,874	13,819	5,442	5,340	4,882	4,813
50-54	16,726	16,863	5,720	6,306	5,323	5,764
55-59	13,975	14,152	4,456	5,206	4,036 ^a	4,784
60-64	12,237	12,397	3,234	3,682	2,915	3,272
65 or Older	28,193	29,457	4,817	4,888	4,363	4,321

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.12B Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	61.8	61.0	25.3	24.8	21.3	20.8
12	2.1	2.6	1.4	1.0	0.4	0.3
13	5.6	4.7	3.1	2.4	1.3	1.1
14	11.2 ^b	8.8	6.2	5.2	2.6	2.1
15	17.7 ^a	15.2	12.1 ^b	9.1	6.1 ^a	4.7
16	23.9	22.6	15.9	15.1	8.7	8.6
17	32.5	29.6	22.3 ^a	19.5	14.2 ^a	11.8
18	39.5	36.9	29.8	28.0	21.0	19.4
19	50.8	48.3	38.3	35.0	28.5 ^b	24.0
20	56.7	54.7	41.7	39.4	32.4	28.9
21	60.1	56.8	41.8	39.0	31.5	29.6
22	62.3	59.8	41.2	41.3	32.8	32.0
23	63.9	63.7	41.8	39.9	34.0	30.9
24	65.3	63.8	43.0 ^a	38.9	33.3	30.2
25	66.1	65.9	39.5	40.0	32.3	32.2
26-29	68.0	65.5	40.1 ^b	35.6	32.7	29.9
30-34	71.8 ^a	68.7	39.0 ^b	35.3	33.2 ^b	29.0
35-39	65.5	67.2	30.0	31.0	25.6	26.8
40-44	67.3	66.1	27.0	27.6	23.4	23.8
45-49	69.4	67.3	27.2	26.0	24.4	23.4
50-54	72.5	71.0	24.8	26.5	23.1	24.3
55-59	68.0	67.8	21.7	24.9	19.6 ^a	22.9
60-64	68.8	70.7	18.2	21.0	16.4	18.6
65 or Older	64.9	65.5	11.1	10.9	10.0	9.6

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.13A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	46,139	45,290	11,866	11,559	8,813	8,662
12	45	47	30	20	10	4
13	91	71	45	37	15	21
14	187	141	126	103	63	48
15	269	233	193	172	91	84
16	416	380	292	259	138	143
17	493	542	325	350	171	191
18	726	633	489	408	306 ^a	224
19	834 ^b	646	476	379	281	216
20	834	879	428	461	262	266
21	835	951	428	458	266	285
22	1,047	1,009	394	429	237	293
23	967	859	353	345	234	233
24	999	929	335	302	232	218
25	823	870	328	308	201	229
26-29	3,623	3,565	1,131	1,138	774	776
30-34	4,364	4,404	1,243	1,228	962	983
35-39	4,294	4,559	982	1,089	761	868
40-44	5,314	4,817	1,114	1,035	1,037	885
45-49	4,806	4,754	1,021	797	902	738
50-54	4,930	4,430	777	793	603	672
55-59	2,858	3,144	535	434	500	363
60-64	2,521	2,458	219	261	179	224
65 or Older	4,865	4,970	603	755	590	700

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.13B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	17.6	17.1	4.5	4.4	3.4	3.3
12	1.2	1.2	0.8	0.5	0.3	0.1
13	2.2	1.7	1.1	0.9	0.3	0.5
14	4.3	3.4	2.9	2.5	1.4	1.1
15	6.5	5.4	4.7	4.0	2.2	1.9
16	9.7	8.8	6.8	6.0	3.2	3.3
17	11.9	13.0	7.9	8.4	4.1	4.6
18	15.3	13.7	10.3	8.9	6.5	4.9
19	20.0 ^b	15.6	11.4	9.2	6.8	5.2
20	19.4	20.1	10.0	10.6	6.1	6.1
21	19.7	21.9	10.1	10.5	6.3	6.6
22	23.4	22.4	8.8	9.5	5.3	6.5
23	22.1	19.5	8.1	7.8	5.3	5.3
24	22.7	21.5	7.6	7.0	5.3	5.0
25	20.0	20.5	8.0	7.3	4.9	5.4
26-29	21.1	20.9	6.6	6.7	4.5	4.6
30-34	21.7	21.1	6.2	5.9	4.8	4.7
35-39	22.5	23.3	5.2	5.6	4.0	4.4
40-44	24.7	23.9	5.2	5.1	4.8	4.4
45-49	24.0	23.2	5.1 ^a	3.9	4.5	3.6
50-54	21.4	18.6	3.4	3.3	2.6	2.8
55-59	13.9	15.1	2.6	2.1	2.4	1.7
60-64	14.2	14.0	1.2	1.5	1.0	1.3
65 or Older	11.2	11.0	1.4	1.7	1.4	1.6

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.14A Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	89,117	89,267	24,933	23,734	12,438	11,964
12	33	38	18	24	5	10
13	97	82	65	53	13	10
14	202	189	118	126	29	43
15	338	333	242	238	93	82
16	586	515	415	376	184	150
17	813	828	566	608	252	226
18	1,258	1,145	979	915	495	445
19	1,404	1,377	996	1,008	455	479
20	1,611	1,655	1,024	946	513	456
21	1,754	1,729	960	938	477	463
22	1,873	1,876	869	908	386	452
23	1,899	1,768	876	759	463 ^a	362
24	1,876	1,812	820	802	384	399
25	1,704	1,808	690	698	300	336
26-29	7,724	7,103	3,116	2,710	1,465	1,269
30-34	8,916	9,037	2,835	2,657	1,430	1,294
35-39	7,255 ^a	8,064	1,713	1,928	812	908
40-44	8,012	7,486	1,789	1,786	972	960
45-49	7,302	7,451	1,714	1,491	927	727
50-54	9,140	8,188	1,926	1,552	939	880
55-59	6,789	7,114	1,232	1,295	582	861
60-64	6,342	5,993	1,061	833	640	542
65 or Older	12,187 ^a	13,676	909	1,080	621	611

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.14B Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	34.0	33.7	9.5 ^a	9.0	4.7	4.5
12	0.9	1.0	0.5	0.6	0.1	0.3
13	2.3	2.0	1.5	1.3	0.3	0.3
14	4.6	4.5	2.7	3.0	0.7	1.0
15	8.2	7.8	5.8	5.5	2.3	1.9
16	13.7	11.9	9.7	8.7	4.3	3.5
17	19.7	19.8	13.7	14.6	6.1	5.4
18	26.6	24.9	20.7	19.9	10.5	9.7
19	33.7	33.3	23.9	24.4	10.9	11.6
20	37.5	37.9	23.8	21.7	11.9	10.5
21	41.4	39.8	22.6	21.6	11.3	10.6
22	41.9	41.6	19.4	20.1	8.6	10.0
23	43.4	40.2	20.0	17.3	10.6 ^a	8.2
24	42.7	41.9	18.7	18.6	8.7	9.2
25	41.5	42.5	16.8	16.4	7.3	7.9
26-29	44.9 ^a	41.7	18.1	15.9	8.5	7.5
30-34	44.3	43.3	14.1	12.7	7.1	6.2
35-39	38.1 ^a	41.2	9.0	9.9	4.3	4.6
40-44	37.3	37.1	8.3	8.9	4.5	4.8
45-49	36.5	36.3	8.6	7.3	4.6	3.5
50-54	39.6 ^b	34.5	8.3	6.5	4.1	3.7
55-59	33.0	34.1	6.0	6.2	2.8	4.1
60-64	35.7	34.2	6.0	4.7	3.6	3.1
65 or Older	28.0	30.4	2.1	2.4	1.4	1.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.15A Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	213,794 ^b	217,765	173,966 ^a	176,556	136,868 ^a	139,677
12	227	173	129	90	41	36
13	503	476	341	353	126	132
14	977 ^a	833	717 ^a	607	306 ^a	220
15	1,450	1,488	1,179	1,168	501	502
16	2,061	2,002	1,678	1,696	763	817
17	2,451	2,403	2,089	2,065	1,140	1,165
18	3,097	2,972	2,751	2,585	1,661	1,678
19	3,096	2,964	2,762	2,620	1,894	1,783
20	3,388	3,438	3,046	3,121	2,219	2,327
21	3,746	3,816	3,597	3,682	2,941	3,095
22	4,025	4,068	3,757	3,819	3,132	3,150
23	4,021	4,021	3,694	3,674	3,052	2,988
24	4,011	3,924	3,674	3,616	3,095	2,924
25	3,763	3,923	3,427	3,596	2,745	2,869
26-29	15,662	15,468	14,205	13,937	11,878	11,649
30-34	18,041	19,071	15,860	16,478	12,808	13,338
35-39	16,897	17,507	14,258	14,931	11,473	11,955
40-44	19,313	18,257	16,345	15,519	13,043	12,548
45-49	17,858	18,386	14,693	15,191	11,784	12,367
50-54	21,234	21,462	17,259	16,880	13,827	13,687
55-59	18,135	18,364	13,481	14,240	10,792	11,430
60-64	15,390	15,706	11,355	11,457	9,524	9,192
65 or Older	34,448 ^b	37,044	23,671 ^a	25,230	18,123 ^a	19,824

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.15B Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	81.5	82.1	66.3	66.6	52.2	52.7
12	6.1 ^a	4.5	3.4 ^a	2.4	1.1	0.9
13	11.9	11.7	8.1	8.6	3.0	3.2
14	22.3	20.0	16.3	14.5	7.0 ^a	5.3
15	35.1	34.7	28.5	27.2	12.1	11.7
16	48.2	46.2	39.2	39.1	17.9	18.8
17	59.4	57.6	50.6	49.5	27.6	27.9
18	65.4	64.5	58.1	56.1	35.1	36.4
19	74.3	71.6	66.3	63.3	45.5	43.1
20	78.9	78.8	70.9	71.6	51.7	53.4
21	88.4	87.8	84.9	84.7	69.4	71.2
22	89.9	90.2	83.9	84.6	70.0	69.8
23	91.9	91.5	84.4	83.6	69.7	68.0
24	91.3	90.7	83.6	83.6	70.5	67.6
25	91.6	92.3	83.4	84.6	66.8	67.5
26-29	91.0	90.8	82.5	81.9	69.0	68.4
30-34	89.6 ^a	91.4	78.8	79.0	63.6	64.0
35-39	88.7	89.5	74.9	76.3	60.2	61.1
40-44	89.9	90.5	76.1	76.9	60.7	62.2
45-49	89.3	89.5	73.5	74.0	58.9	60.2
50-54	92.0	90.3	74.8 ^a	71.1	59.9	57.6
55-59	88.3	88.0	65.6	68.2	52.5	54.8
60-64	86.6	89.5	63.9	65.3	53.6	52.4
65 or Older	79.3 ^a	82.3	54.5	56.1	41.7	44.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.16A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	136,868 ^a	139,677	60,064	60,941	16,541	16,342
12	41	36	12	11	4	*
13	126	132	55	53	5	8
14	306 ^a	220	136	95	15	14
15	501	502	249	235	43	30
16	763	817	419	469	73	72
17	1,140	1,165	685	649	152	133
18	1,661	1,678	1,053	1,098	284	270
19	1,894	1,783	1,312	1,131	356	325
20	2,219	2,327	1,478	1,510	480	449
21	2,941	3,095	1,914	2,046	585	675
22	3,132	3,150	2,042	1,965	625	599
23	3,052	2,988	1,895	1,888	591	536
24	3,095	2,924	1,883	1,795	577 ^b	420
25	2,745	2,869	1,609	1,752	440	509
26-29	11,878	11,649	6,890	6,424	1,934 ^a	1,567
30-34	12,808	13,338	7,092	7,039	2,124	1,800
35-39	11,473	11,955	5,636	5,856	1,425	1,573
40-44	13,043	12,548	5,512	5,567	1,567	1,552
45-49	11,784	12,367	5,151	5,217	1,387	1,462
50-54	13,827	13,687	5,314	5,516	1,294	1,597
55-59	10,792	11,430	3,275 ^b	4,237	810	1,083
60-64	9,524	9,192	2,513	2,373	844	694
65 or Older	18,123 ^a	19,824	3,938	4,016	925	975

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.16B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	52.2	52.7	22.9	23.0	6.3	6.2
12	1.1	0.9	0.3	0.3	0.1	*
13	3.0	3.2	1.3	1.3	0.1	0.2
14	7.0 ^a	5.3	3.1	2.3	0.3	0.3
15	12.1	11.7	6.0	5.5	1.0	0.7
16	17.9	18.8	9.8	10.8	1.7	1.7
17	27.6	27.9	16.6	15.5	3.7	3.2
18	35.1	36.4	22.3	23.8	6.0	5.9
19	45.5	43.1	31.5 ^a	27.3	8.5	7.8
20	51.7	53.4	34.4	34.6	11.2	10.3
21	69.4	71.2	45.2	47.1	13.8	15.5
22	70.0	69.8	45.6	43.6	14.0	13.3
23	69.7	68.0	43.3	43.0	13.5	12.2
24	70.5	67.6	42.9	41.5	13.1 ^b	9.7
25	66.8	67.5	39.2	41.2	10.7	12.0
26-29	69.0	68.4	40.0	37.7	11.2 ^a	9.2
30-34	63.6	64.0	35.2	33.8	10.5	8.6
35-39	60.2	61.1	29.6	29.9	7.5	8.0
40-44	60.7	62.2	25.7	27.6	7.3	7.7
45-49	58.9	60.2	25.8	25.4	6.9	7.1
50-54	59.9	57.6	23.0	23.2	5.6	6.7
55-59	52.5	54.8	15.9 ^b	20.3	3.9	5.2
60-64	53.6	52.4	14.1	13.5	4.7	4.0
65 or Older	41.7	44.1	9.1	8.9	2.1	2.2

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.17A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics:
Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	175,260	175,416	80,846	81,141	66,879	66,899
AGE						
12-17	4,922 ^a	4,591	3,461 ^a	3,161	1,951 ^a	1,742
18-25	22,650	22,290	16,597	16,083	12,882 ^a	12,244
26 or Older	147,688	148,534	60,788	61,896	52,046	52,913
GENDER						
Male	94,449	94,796	47,773	48,527	39,549	39,952
Female	80,810	80,620	33,073	32,614	27,331	26,947
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	153,298	153,315	70,628	70,735	59,108	58,933
White	126,277	126,091	56,518	56,442	47,170	46,927
Black or African American	17,942	17,526	9,675	9,683	8,461	8,425
American Indian or Alaska Native	1,049	1,031	663	616	562	545
Native Hawaiian or Other Pacific Islander	526	652	312	365	225	312
Asian	4,494	4,960	1,819	2,019	1,353	1,418
Two or More Races	3,010	3,054	1,640	1,609	1,337	1,305
Hispanic or Latino	21,961	22,102	10,218	10,406	7,771	7,966

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.17B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	66.8	66.2	30.8	30.6	25.5	25.2
AGE						
12-17	19.8 ^a	18.5	13.9 ^a	12.7	7.8 ^a	7.0
18-25	65.1	63.8	47.7 ^a	46.0	37.0 ^b	35.0
26 or Older	72.9	72.3	30.0	30.1	25.7	25.8
GENDER						
Male	74.3	73.8	37.6	37.8	31.1	31.1
Female	59.7	59.0	24.4	23.9	20.2	19.7
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	69.3	68.9	31.9	31.8	26.7	26.5
White	74.3	74.1	33.2	33.2	27.7	27.6
Black or African American	57.4	55.3	31.0	30.6	27.1	26.6
American Indian or Alaska Native	74.8	71.5	47.3	42.7	40.1	37.8
Native Hawaiian or Other Pacific Islander	60.4	63.8	35.9	35.8	25.8	30.6
Asian	33.7	35.8	13.6	14.6	10.1	10.2
Two or More Races	70.2	69.1	38.2	36.4	31.2	29.5
Hispanic or Latino	53.2	52.0	24.8	24.5	18.8	18.8

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.18A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	4,922 ^a	4,591	3,461 ^a	3,161	1,951 ^a	1,742
GENDER						
Male	2,814 ^a	2,563	1,986	1,820	1,158	1,036
Female	2,108	2,028	1,475	1,342	793	706
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4,007 ^b	3,643	2,842 ^b	2,557	1,648 ^a	1,461
White	3,126	2,912	2,295	2,103	1,350	1,221
Black or African American	545 ^a	442	337 ^a	254	170	140
American Indian or Alaska Native	59	45	32	35	18	*
Native Hawaiian or Other Pacific Islander	*	*	8	*	7	*
Asian	99	85	69	56	36	23
Two or More Races	163	140	100	95	67	50
Hispanic or Latino	916	948	619	605	303	281
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	1,810	1,681	1,329	1,252	801	751
Female, White, Not Hispanic	1,316	1,231	966	851	549	470
Male, Black, Not Hispanic	318 ^a	242	211 ^a	144	106	85
Female, Black, Not Hispanic	228	199	126	109	64	55
Male, Hispanic	508	488	332	318	176	145
Female, Hispanic	408	460	287	286	127	136

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.18B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	19.8 ^a	18.5	13.9 ^a	12.7	7.8 ^a	7.0
GENDER						
Male	22.1 ^a	20.2	15.6	14.3	9.1	8.2
Female	17.3	16.6	12.1	11.0	6.5	5.8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	20.7 ^b	18.9	14.7 ^b	13.3	8.5 ^a	7.6
White	22.9	21.6	16.8	15.6	9.9	9.1
Black or African American	15.8 ^b	12.7	9.8 ^b	7.3	4.9	4.0
American Indian or Alaska Native	36.9	27.1	20.3	21.1	11.4	*
Native Hawaiian or Other Pacific Islander	*	*	7.9	*	7.4	*
Asian	8.0	6.9	5.6	4.6	2.9	1.9
Two or More Races	21.6	18.3	13.3	12.4	8.9	6.5
Hispanic or Latino	16.5	16.8	11.1	10.7	5.5	5.0
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	26.0	24.4	19.1	18.2	11.5	10.9
Female, White, Not Hispanic	19.7	18.7	14.5	12.9	8.2	7.1
Male, Black, Not Hispanic	18.1 ^a	13.7	12.0 ^b	8.2	6.1	4.8
Female, Black, Not Hispanic	13.5	11.6	7.5	6.4	3.8	3.2
Male, Hispanic	17.9	17.0	11.7	11.1	6.2	5.0
Female, Hispanic	15.0	16.6	10.5	10.3	4.7	4.9

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.19A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	22,650	22,290	16,597	16,083	12,882 ^a	12,244
GENDER						
Male	12,579	12,314	10,103 ^a	9,726	8,080 ^a	7,666
Female	10,070	9,977	6,493	6,357	4,802	4,578
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	18,345	18,080	13,766	13,386	10,881 ^a	10,324
White	14,188	13,914	10,766	10,336	8,562 ^a	7,988
Black or African American	2,539	2,543	1,887	1,900	1,495	1,564
American Indian or Alaska Native	180	165	130	125	115	109
Native Hawaiian or Other Pacific Islander	*	*	*	*	57	*
Asian	788	799	516	544	355	346
Two or More Races	517	557	374	395	298	256
Hispanic or Latino	4,305	4,210	2,831	2,697	2,002	1,920
EDUCATION						
< High School	3,677 ^a	3,267	2,971	2,724	2,525	2,292
High School Graduate	7,643	7,449	5,868	5,673	4,707	4,549
Some College	7,931	7,980	5,677	5,613	4,304	4,114
College Graduate	3,399	3,595	2,080	2,073	1,346	1,289
CURRENT EMPLOYMENT						
Full-Time	9,819	10,158	7,065	7,266	5,676	5,759
Part-Time	6,015	5,851	4,366	4,069	3,197 ^a	2,829
Unemployed	2,828 ^b	2,478	2,274	2,044	1,894 ^a	1,645
Other ¹	3,988	3,803	2,892	2,704	2,116	2,011

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.19B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	65.1	63.8	47.7 ^a	46.0	37.0 ^b	35.0
GENDER						
Male	72.0	70.2	57.9 ^a	55.5	46.3 ^a	43.7
Female	58.1	57.3	37.5	36.5	27.7	26.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	66.6	65.6	50.0	48.6	39.5 ^b	37.4
White	72.8	71.9	55.2 ^a	53.4	43.9 ^b	41.3
Black or African American	50.7	50.5	37.7	37.7	29.9	31.1
American Indian or Alaska Native	78.2	81.5	56.7	61.7	49.9	53.5
Native Hawaiian or Other Pacific Islander	*	*	*	*	28.4	*
Asian	42.1	39.8	27.6	27.1	19.0	17.2
Two or More Races	70.5	71.0	51.1	50.4	40.6 ^a	32.7
Hispanic or Latino	59.4	57.2	39.0	36.6	27.6	26.1
EDUCATION						
< High School	65.8	66.1	53.2	55.1	45.2	46.3
High School Graduate	63.8	62.2	49.0	47.4	39.3	38.0
Some College	65.1	64.1	46.6	45.1	35.3 ^a	33.1
College Graduate	67.3	64.5	41.2 ^a	37.2	26.7 ^a	23.1
CURRENT EMPLOYMENT						
Full-Time	72.7	71.3	52.3	51.0	42.0	40.4
Part-Time	62.2	61.7	45.2	42.9	33.1 ^b	29.8
Unemployed	68.0 ^a	63.7	54.7	52.6	45.6	42.3
Other ¹	53.5	52.0	38.8	37.0	28.4	27.5

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.20A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics:
Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	147,688	148,534	60,788	61,896	52,046	52,913
GENDER						
Male	79,056	79,919	35,683	36,981	30,310	31,250
Female	68,632	68,615	25,105	24,915	21,736	21,663
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	130,947	131,591	54,019	54,792	46,580	47,148
White	108,964	109,266	43,456	44,003	37,258	37,719
Black or African American	14,857	14,542	7,451	7,530	6,797	6,721
American Indian or Alaska Native	810	821	501	456	429	412
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	3,607	4,076	1,234	1,419	963	1,049
Two or More Races	2,330	2,356	1,165	1,120	972	999
Hispanic or Latino	16,741	16,943	6,769	7,104	5,466	5,765
EDUCATION						
< High School	18,302	18,142	10,185	10,189	9,386	9,237
High School Graduate	42,408	42,075	19,947	20,350	17,903	18,487
Some College	39,068	40,081	17,118	17,648	14,852	15,372
College Graduate	47,910	48,236	13,537	13,709	9,905	9,817
CURRENT EMPLOYMENT						
Full-Time	79,065	80,517	34,667	35,631	28,362	29,453
Part-Time	17,770	16,667	6,787	6,028	5,832	5,184
Unemployed	5,753	5,339	3,716	3,396	3,493	3,058
Other ¹	45,100	46,011	15,619	16,842	14,359	15,217

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.20B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	72.9	72.3	30.0	30.1	25.7	25.8
GENDER						
Male	81.5	81.4	36.8	37.7	31.3	31.8
Female	64.9	64.1	23.7	23.3	20.6	20.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	75.1	74.8	31.0	31.2	26.7	26.8
White	79.6	79.5	31.8	32.0	27.2	27.5
Black or African American	65.2	62.8	32.7	32.5	29.8	29.0
American Indian or Alaska Native	80.1	76.4	49.5	42.5	42.4	38.4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	35.2	38.4	12.1	13.4	9.4	9.9
Two or More Races	83.1	82.2	41.6	39.1	34.7	34.8
Hispanic or Latino	58.9	57.5	23.8	24.1	19.2	19.6
EDUCATION						
< High School	68.9	66.5	38.3	37.4	35.3	33.9
High School Graduate	73.1	73.9	34.4	35.7	30.9	32.5
Some College	76.3	75.4	33.4	33.2	29.0	28.9
College Graduate	71.6	71.0	20.2	20.2	14.8	14.4
CURRENT EMPLOYMENT						
Full-Time	75.0	74.1	32.9	32.8	26.9	27.1
Part-Time	73.8 ^a	70.8	28.2	25.6	24.2	22.0
Unemployed	75.2	77.3	48.6	49.2	45.7	44.3
Other ¹	68.8	69.5	23.8	25.4	21.9	23.0

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.21A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics:
Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	170,337	170,825	77,385	77,979	64,928	65,157
GENDER						
Male	91,635	92,233	45,787	46,707	38,390	38,916
Female	78,702	78,592	31,598	31,272	26,538	26,241
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	149,292	149,672	67,785	68,178	57,461	57,472
White	123,151	123,180	54,222	54,339	45,820	45,706
Black or African American	17,396	17,085	9,338	9,430	8,291	8,285
American Indian or Alaska Native	990	986	631	581	543	521
Native Hawaiian or Other Pacific Islander	*	633	305	351	218	309
Asian	4,395	4,875	1,750	1,962	1,318	1,395
Two or More Races	2,847	2,913	1,540	1,514	1,270	1,255
Hispanic or Latino	21,046	21,153	9,599	9,801	7,468	7,685
EDUCATION						
< High School	21,979	21,409	13,156	12,913	11,911	11,529
High School Graduate	50,051	49,524	25,815	26,023	22,610	23,035
Some College	46,999	48,061	22,796	23,262	19,156	19,486
College Graduate	51,309	51,831	15,618	15,782	11,252	11,106
CURRENT EMPLOYMENT						
Full-Time	88,884	90,676	41,732	42,897	34,038	35,212
Part-Time	23,785	22,518	11,153 ^a	10,096	9,028 ^a	8,013
Unemployed	8,581	7,817	5,990	5,440	5,387 ^a	4,703
Other ¹	49,088	49,814	18,511	19,546	16,474	17,229

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.21B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	71.7	71.1	32.6	32.5	27.3	27.1
GENDER						
Male	80.1	79.7	40.0	40.4	33.6	33.6
Female	63.9	63.1	25.7	25.1	21.6	21.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	74.0	73.6	33.6	33.5	28.5	28.3
White	78.8	78.6	34.7	34.7	29.3	29.2
Black or African American	62.6	60.6	33.6	33.4	29.8	29.4
American Indian or Alaska Native	79.7	77.3	50.8	45.6	43.8	40.8
Native Hawaiian or Other Pacific Islander	*	69.0	39.4	38.2	28.1	33.7
Asian	36.3	38.6	14.5	15.5	10.9	11.0
Two or More Races	80.5	79.8	43.5	41.5	35.9	34.4
Hispanic or Latino	59.0	57.4	26.9	26.6	20.9	20.9
EDUCATION						
< High School	68.3	66.4	40.9	40.1	37.0	35.8
High School Graduate	71.5	71.9	36.9	37.8	32.3	33.4
Some College	74.1	73.3	36.0	35.5	30.2	29.7
College Graduate	71.3	70.5	21.7	21.5	15.6	15.1
CURRENT EMPLOYMENT						
Full-Time	74.7	73.8	35.1	34.9	28.6	28.6
Part-Time	70.5 ^a	68.2	33.0 ^a	30.6	26.8 ^a	24.3
Unemployed	72.7	72.4	50.7	50.4	45.6	43.6
Other ¹	67.3	67.8	25.4	26.6	22.6	23.4

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.22A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	162,138	161,789	66,256	65,858	55,778	55,241
AGE						
12-17	3,903 ^b	3,523	2,561 ^b	2,210	1,397 ^a	1,213
18-25	20,141 ^a	19,608	13,750 ^a	13,157	10,657 ^b	9,905
26 or Older	138,095	138,658	49,945	50,491	43,724	44,123
GENDER						
Male	84,457	84,510	35,858	35,638	30,041	29,822
Female	77,682	77,279	30,398	30,220	25,737	25,418
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	141,708	141,135	57,330	56,700	48,833	48,166
White	117,091	116,260	45,398	44,716	38,644	37,966
Black or African American	16,091	15,914	8,120	8,119	7,193	7,120
American Indian or Alaska Native	976	947	573	544	511	469
Native Hawaiian or Other Pacific Islander	506	622	269	297	184	259
Asian	4,167	4,547	1,552	1,739	1,139	1,272
Two or More Races	2,877	2,844	1,418	1,286	1,162	1,080
Hispanic or Latino	20,430	20,654	8,926	9,158	6,945	7,075

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.22B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	61.8	61.0	25.3	24.8	21.3	20.8
AGE						
12-17	15.7 ^b	14.2	10.3 ^b	8.9	5.6 ^a	4.9
18-25	57.9 ^a	56.1	39.5 ^a	37.7	30.6 ^b	28.4
26 or Older	68.1	67.5	24.6	24.6	21.6	21.5
GENDER						
Male	66.4	65.8	28.2	27.8	23.6	23.2
Female	57.4	56.5	22.5	22.1	19.0	18.6
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	64.1	63.4	25.9	25.5	22.1	21.6
White	68.9	68.3	26.7	26.3	22.7	22.3
Black or African American	51.5	50.2	26.0	25.6	23.0	22.5
American Indian or Alaska Native	69.6	65.7	40.9	37.7	36.5	32.5
Native Hawaiian or Other Pacific Islander	58.1	61.0	30.9	29.1	21.1	25.4
Asian	31.2	32.8	11.6	12.5	8.5	9.2
Two or More Races	67.1	64.4	33.0	29.1	27.1	24.4
Hispanic or Latino	49.5	48.6	21.6	21.6	16.8	16.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.23A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	3,903 ^b	3,523	2,561 ^b	2,210	1,397 ^a	1,213
GENDER						
Male	2,067 ^b	1,842	1,312 ^a	1,154	722	649
Female	1,835	1,682	1,249 ^b	1,056	675 ^a	564
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	3,134 ^b	2,741	2,063 ^b	1,737	1,189 ^b	996
White	2,453 ^b	2,168	1,669 ^b	1,428	979 ^a	854
Black or African American	396	330	220 ^a	154	110	77
American Indian or Alaska Native	45	38	26	29	12	14
Native Hawaiian or Other Pacific Islander	*	*	6	*	4	*
Asian	93	63	64	41	31	16
Two or More Races	134	124	78	72	53	33
Hispanic or Latino	768	783	498	473	208	217
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	1,294 ^a	1,141	832	753	484	473
Female, White, Not Hispanic	1,160	1,027	837 ^b	675	496 ^b	382
Male, Black, Not Hispanic	217	179	126 ^a	85	62	41
Female, Black, Not Hispanic	180	151	94	69	48	35
Male, Hispanic	417	394	269	237	125	103
Female, Hispanic	352	388	229	236	84	114

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.23B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	15.7 ^b	14.2	10.3 ^b	8.9	5.6 ^a	4.9
GENDER						
Male	16.3 ^a	14.5	10.3 ^a	9.1	5.7	5.1
Female	15.1	13.8	10.3 ^b	8.7	5.5 ^a	4.6
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	16.2 ^b	14.3	10.7 ^b	9.0	6.1 ^b	5.2
White	18.0 ^b	16.1	12.2 ^b	10.6	7.2	6.3
Black or African American	11.5 ^a	9.5	6.4 ^b	4.4	3.2 ^a	2.2
American Indian or Alaska Native	28.3	22.8	16.1	17.5	7.5	8.6
Native Hawaiian or Other Pacific Islander	*	*	6.0	*	3.6	*
Asian	7.5	5.1	5.1	3.4	2.5	1.3
Two or More Races	17.8	16.2	10.3	9.3	7.0	4.4
Hispanic or Latino	13.8	13.9	9.0	8.4	3.7	3.8
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	18.6 ^a	16.6	12.0	10.9	7.0	6.9
Female, White, Not Hispanic	17.4	15.6	12.5 ^b	10.2	7.4 ^b	5.8
Male, Black, Not Hispanic	12.4	10.2	7.2 ^a	4.8	3.5	2.3
Female, Black, Not Hispanic	10.6	8.8	5.6	4.0	2.8	2.0
Male, Hispanic	14.7	13.7	9.5	8.3	4.4	3.6
Female, Hispanic	12.9	14.0	8.4	8.5	3.1	4.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.24A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	20,141 ^a	19,608	13,750 ^a	13,157	10,657 ^b	9,905
GENDER						
Male	10,990 ^a	10,638	8,061 ^a	7,642	6,346 ^b	5,883
Female	9,151	8,970	5,690	5,515	4,310 ^a	4,021
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	16,149	15,729	11,315 ^a	10,797	8,878 ^b	8,271
White	12,535	12,138	8,855	8,426	6,986 ^a	6,440
Black or African American	2,161	2,182	1,513	1,478	1,198	1,198
American Indian or Alaska Native	158	144	106	101	94	78
Native Hawaiian or Other Pacific Islander	*	*	*	*	52	*
Asian	713	704	432	441	301	307
Two or More Races	463	474	320	291	246	206
Hispanic or Latino	3,992	3,879	2,435	2,360	1,779	1,633
EDUCATION						
< High School	3,408 ^a	3,035	2,651	2,400	2,241 ^a	1,958
High School Graduate	6,865	6,691	4,957	4,765	3,995	3,785
Some College	7,025	6,923	4,659	4,470	3,464	3,226
College Graduate	2,843	2,959	1,484	1,522	956	935
CURRENT EMPLOYMENT						
Full-Time	8,828	9,065	5,858	6,008	4,664	4,574
Part-Time	5,176	4,986	3,500	3,207	2,583 ^b	2,232
Unemployed	2,581 ^b	2,247	1,970	1,756	1,613	1,432
Other ¹	3,556	3,310	2,423	2,186	1,796	1,667

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.24B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	57.9 ^a	56.1	39.5 ^a	37.7	30.6 ^b	28.4
GENDER						
Male	62.9 ^a	60.7	46.2 ^a	43.6	36.3 ^b	33.6
Female	52.8	51.5	32.8	31.7	24.9 ^a	23.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	58.6	57.1	41.1 ^a	39.2	32.2 ^b	30.0
White	64.3	62.7	45.4 ^a	43.5	35.8 ^b	33.3
Black or African American	43.2	43.4	30.2	29.4	23.9	23.8
American Indian or Alaska Native	68.8	70.8	46.2	49.8	41.0	38.7
Native Hawaiian or Other Pacific Islander	*	*	*	*	26.0	*
Asian	38.1	35.0	23.1	21.9	16.1	15.3
Two or More Races	63.2	60.5	43.7	37.1	33.6	26.3
Hispanic or Latino	55.1	52.7	33.6	32.0	24.5	22.2
EDUCATION						
< High School	61.0	61.4	47.4	48.5	40.1	39.6
High School Graduate	57.4	55.9	41.4	39.8	33.4	31.6
Some College	57.7	55.6	38.2 ^a	35.9	28.4 ^a	25.9
College Graduate	56.3	53.1	29.4	27.3	18.9	16.8
CURRENT EMPLOYMENT						
Full-Time	65.3	63.6	43.4	42.1	34.5 ^a	32.1
Part-Time	53.5	52.6	36.2	33.8	26.7 ^b	23.5
Unemployed	62.1 ^a	57.8	47.4	45.2	38.8	36.8
Other ¹	47.7	45.3	32.5	29.9	24.1	22.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.25A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	138,095	138,658	49,945	50,491	43,724	44,123
GENDER						
Male	71,400	72,030	26,486	26,842	22,973	23,290
Female	66,695	66,628	23,459	23,649	20,752	20,833
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	122,425	122,665	43,952	44,166	38,766	38,898
White	102,102	101,954	34,874	34,861	30,679	30,672
Black or African American	13,534	13,402	6,387	6,487	5,884	5,845
American Indian or Alaska Native	772	765	441	414	405	376
Native Hawaiian or Other Pacific Islander	*	*	*	223	*	216
Asian	3,360	3,780	1,056	1,257	807	948
Two or More Races	2,281	2,246	1,019	924	862	840
Hispanic or Latino	15,670	15,992	5,993	6,325	4,958	5,225
EDUCATION						
< High School	17,413	17,248	9,294	9,102	8,567	8,092
High School Graduate	40,210	39,882	16,880	17,474	15,361	15,915
Some College	36,978	38,029	14,484	14,739	12,694	13,058
College Graduate	43,494	43,498	9,286	9,176	7,103	7,059
CURRENT EMPLOYMENT						
Full-Time	72,715	73,930	26,706	27,486	22,514	23,459
Part-Time	17,032	15,906	5,913	5,206	5,303 ^a	4,568
Unemployed	5,423	5,039	3,344	3,101	3,117	2,786
Other ¹	42,924	43,782	13,982	14,698	12,790	13,311

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.25B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	68.1	67.5	24.6	24.6	21.6	21.5
GENDER						
Male	73.6	73.3	27.3	27.3	23.7	23.7
Female	63.1	62.2	22.2	22.1	19.6	19.5
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	70.3	69.8	25.2	25.1	22.2	22.1
White	74.6	74.2	25.5	25.4	22.4	22.3
Black or African American	59.4	57.9	28.0	28.0	25.8	25.2
American Indian or Alaska Native	76.3	71.3	43.6	38.5	40.0	35.0
Native Hawaiian or Other Pacific Islander	*	*	*	30.1	*	29.2
Asian	32.8	35.6	10.3	11.8	7.9	8.9
Two or More Races	81.3	78.3	36.4	32.2	30.8	29.3
Hispanic or Latino	55.1	54.3	21.1	21.5	17.4	17.7
EDUCATION						
< High School	65.5	63.2	35.0	33.4	32.2	29.7
High School Graduate	69.3	70.0	29.1	30.7	26.5	27.9
Some College	72.2	71.6	28.3	27.7	24.8	24.6
College Graduate	65.0	64.0	13.9	13.5	10.6	10.4
CURRENT EMPLOYMENT						
Full-Time	69.0	68.0	25.3	25.3	21.4	21.6
Part-Time	70.7 ^a	67.6	24.6	22.1	22.0 ^a	19.4
Unemployed	70.9	73.0	43.7	44.9	40.8	40.3
Other ¹	65.5	66.1	21.3	22.2	19.5	20.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.26A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	158,236	158,266	63,695	63,648	54,381	54,028
GENDER						
Male	82,389	82,668	34,546	34,484	29,319	29,173
Female	75,846	75,598	29,148	29,164	25,062	24,855
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	138,573	138,395	55,267	54,963	47,644	47,169
White	114,637	114,092	43,729	43,288	37,665	37,112
Black or African American	15,695	15,584	7,900	7,965	7,083	7,044
American Indian or Alaska Native	931	909	547	515	499	454
Native Hawaiian or Other Pacific Islander	*	605	263	284	180	258
Asian	4,074	4,484	1,488	1,697	1,108	1,255
Two or More Races	2,744	2,720	1,340	1,214	1,109	1,047
Hispanic or Latino	19,662	19,871	8,428	8,685	6,737	6,858
EDUCATION						
< High School	20,821	20,283	11,945	11,502	10,807	10,049
High School Graduate	47,075	46,573	21,837	22,240	19,356	19,700
Some College	44,002	44,952	19,143	19,209	16,158	16,284
College Graduate	46,337	46,458	10,770	10,698	8,059	7,994
CURRENT EMPLOYMENT						
Full-Time	81,543	82,995	32,564	33,494	27,178	28,032
Part-Time	22,209	20,892	9,413 ^a	8,414	7,886 ^b	6,799
Unemployed	8,004	7,286	5,314	4,856	4,730	4,218
Other ¹	46,480	47,092	16,404	16,884	14,586	14,978

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.26B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	66.6	65.9	26.8	26.5	22.9	22.5
GENDER						
Male	72.0	71.4	30.2	29.8	25.6	25.2
Female	61.6	60.7	23.7	23.4	20.4	20.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	68.7	68.0	27.4	27.0	23.6	23.2
White	73.3	72.8	28.0	27.6	24.1	23.7
Black or African American	56.5	55.3	28.4	28.2	25.5	25.0
American Indian or Alaska Native	74.9	71.2	44.1	40.3	40.2	35.6
Native Hawaiian or Other Pacific Islander	*	65.9	34.0	30.9	23.3	28.1
Asian	33.7	35.5	12.3	13.4	9.2	9.9
Two or More Races	77.6	74.5	37.9	33.3	31.3	28.7
Hispanic or Latino	55.1	54.0	23.6	23.6	18.9	18.6
EDUCATION						
< High School	64.7	62.9	37.1	35.7	33.6	31.2
High School Graduate	67.3	67.6	31.2	32.3	27.7	28.6
Some College	69.4	68.5	30.2	29.3	25.5	24.8
College Graduate	64.4	63.2	15.0	14.6	11.2	10.9
CURRENT EMPLOYMENT						
Full-Time	68.6	67.5	27.4	27.2	22.8	22.8
Part-Time	65.8 ^a	63.3	27.9 ^a	25.5	23.4 ^b	20.6
Unemployed	67.8	67.5	45.0	45.0	40.1	39.1
Other ¹	63.7	64.1	22.5	23.0	20.0	20.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.27A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics:
Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	46,139	45,290	11,866	11,559	8,813	8,662
AGE						
12-17	1,501	1,414	1,011	941	487	490
18-25	7,064	6,775	3,231	3,089	2,018	1,964
26 or Older	37,574	37,101	7,624	7,529	6,308	6,209
GENDER						
Male	38,921	38,512	10,806	10,718	8,211	8,205
Female	7,218	6,777	1,060	842	603	457
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	42,494	41,995	11,018	10,881	8,284	8,283
White	37,785	37,544	9,784	9,908	7,380	7,598
Black or African American	2,644	2,435	744 ^a	477	569	348
American Indian or Alaska Native	391	419	113	117	74	102
Native Hawaiian or Other Pacific Islander	156	116	41	38	34	31
Asian	628	591	139	127	92	73
Two or More Races	888	890	197	214	134	131
Hispanic or Latino	3,645	3,295	849	678	529	379

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.27B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	17.6	17.1	4.5	4.4	3.4	3.3
AGE						
12-17	6.0	5.7	4.1	3.8	2.0	2.0
18-25	20.3	19.4	9.3	8.8	5.8	5.6
26 or Older	18.5	18.1	3.8	3.7	3.1	3.0
GENDER						
Male	30.6	30.0	8.5	8.3	6.5	6.4
Female	5.3	5.0	0.8	0.6	0.4	0.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	19.2	18.9	5.0	4.9	3.7	3.7
White	22.2	22.1	5.8	5.8	4.3	4.5
Black or African American	8.5	7.7	2.4 ^a	1.5	1.8	1.1
American Indian or Alaska Native	27.9	29.0	8.1	8.1	5.3	7.0
Native Hawaiian or Other Pacific Islander	17.9	11.4	4.7	3.7	3.9	3.1
Asian	4.7	4.3	1.0	0.9	0.7	0.5
Two or More Races	20.7	20.1	4.6	4.8	3.1	3.0
Hispanic or Latino	8.8	7.8	2.1	1.6	1.3	0.9

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.28A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,501	1,414	1,011	941	487	490
GENDER						
Male	1,209	1,106	859	760	438	417
Female	291	308	152	181	49	73
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,364	1,264	928	855	463	455
White	1,201	1,128	823	773	405	423
Black or African American	72	47	47	27	21	6
American Indian or Alaska Native	24	*	12	*	7	*
Native Hawaiian or Other Pacific Islander	5	*	4	*	*	*
Asian	16	19	13	15	5	3
Two or More Races	46	40	29	23	20	10
Hispanic or Latino	137	150	83	86	25	35
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	960	871	702	613	364	361
Female, White, Not Hispanic	241	257	120	160	41	62
Male, Black, Not Hispanic	68	42	43	25	21	6
Female, Black, Not Hispanic	4	5	4	2	*	*
Male, Hispanic	109	128	67	81	24	30
Female, Hispanic	28	22	16	5	0	5

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.28B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	6.0	5.7	4.1	3.8	2.0	2.0
GENDER						
Male	9.5	8.7	6.8	6.0	3.4	3.3
Female	2.4	2.5	1.2	1.5	0.4	0.6
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	7.1	6.6	4.8	4.4	2.4	2.4
White	8.8	8.4	6.0	5.7	3.0	3.1
Black or African American	2.1	1.3	1.4	0.8	0.6	0.2
American Indian or Alaska Native	15.1	*	7.4	*	4.5	*
Native Hawaiian or Other Pacific Islander	5.5	*	4.4	*	*	*
Asian	1.3	1.6	1.0	1.2	0.4	0.3
Two or More Races	6.0	5.3	3.8	3.1	2.6	1.3
Hispanic or Latino	2.5	2.6	1.5	1.5	0.4	0.6
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	13.8	12.7	10.1	8.9	5.2	5.2
Female, White, Not Hispanic	3.6	3.9	1.8	2.4	0.6	0.9
Male, Black, Not Hispanic	3.9	2.4	2.5	1.4	1.2	0.4
Female, Black, Not Hispanic	0.3	0.3	0.2	0.1	*	*
Male, Hispanic	3.8	4.4	2.4	2.8	0.9	1.0
Female, Hispanic	1.0	0.8	0.6	0.2	0.0	0.2

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.29A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	7,064	6,775	3,231	3,089	2,018	1,964
GENDER						
Male	5,747 ^a	5,403	2,926	2,828	1,888	1,877
Female	1,318	1,372	305	262	130	87
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	6,249	5,943	2,922	2,803	1,891	1,837
White	5,628	5,283	2,707	2,537	1,764	1,695
Black or African American	267	308	73	111	42	48
American Indian or Alaska Native	82	80	32	33	23	28
Native Hawaiian or Other Pacific Islander	14	*	7	*	*	*
Asian	106	85	42	27	15	14
Two or More Races	153	144	60	78	42	43
Hispanic or Latino	815	832	309	286	128	127
EDUCATION						
< High School	1,150	1,092	598	565	357	355
High School Graduate	2,554	2,469	1,229	1,203	771	756
Some College	2,362	2,347	1,067	1,004	661	641
College Graduate	999	867	337	317	229	211
CURRENT EMPLOYMENT						
Full-Time	3,467	3,484	1,615	1,608	1,081	1,139
Part-Time	1,678	1,562	722	717	399	383
Unemployed	891 ^a	746	435 ^a	331	247	185
Other ¹	1,028	984	459	432	291	257

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.29B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	20.3	19.4	9.3	8.8	5.8	5.6
GENDER						
Male	32.9 ^a	30.8	16.8	16.1	10.8	10.7
Female	7.6	7.9	1.8	1.5	0.8	0.5
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	22.7	21.6	10.6	10.2	6.9	6.7
White	28.9 ^a	27.3	13.9	13.1	9.0	8.8
Black or African American	5.3	6.1	1.5	2.2	0.8	1.0
American Indian or Alaska Native	35.5	39.5	14.0	16.4	9.9	13.9
Native Hawaiian or Other Pacific Islander	7.0	*	3.7	*	*	*
Asian	5.6	4.2	2.2	1.4	0.8	0.7
Two or More Races	20.9	18.3	8.2	10.0	5.7	5.4
Hispanic or Latino	11.2	11.3	4.3	3.9	1.8	1.7
EDUCATION						
< High School	20.6	22.1	10.7	11.4	6.4	7.2
High School Graduate	21.3	20.6	10.3	10.0	6.4	6.3
Some College	19.4	18.9	8.8	8.1	5.4	5.2
College Graduate	19.8 ^b	15.6	6.7	5.7	4.5	3.8
CURRENT EMPLOYMENT						
Full-Time	25.7	24.4	11.9	11.3	8.0	8.0
Part-Time	17.4	16.5	7.5	7.6	4.1	4.0
Unemployed	21.4	19.2	10.5	8.5	5.9	4.8
Other ¹	13.8	13.5	6.2	5.9	3.9	3.5

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.30A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics:
Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	37,574	37,101	7,624	7,529	6,308	6,209
GENDER						
Male	31,965	32,003	7,021	7,130	5,885	5,911
Female	5,609	5,097	604	399	423	297
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	34,881	34,787	7,168	7,223	5,931	5,991
White	30,956	31,133	6,255	6,598	5,211	5,480
Black or African American	2,306	2,081	624 ^a	339	506	294
American Indian or Alaska Native	286	312	69	69	44	63
Native Hawaiian or Other Pacific Islander	*	69	29	20	25	20
Asian	507	487	84	85	72	56
Two or More Races	689	706	108	112	72	79
Hispanic or Latino	2,693	2,314	456	306	377	217
EDUCATION						
< High School	4,200	4,278	1,216	1,226	946	1,043
High School Graduate	10,748	10,844	2,809	2,598	2,370	2,220
Some College	10,016	9,913	1,974	1,920	1,668	1,599
College Graduate	12,609	12,065	1,625	1,785	1,324	1,347
CURRENT EMPLOYMENT						
Full-Time	24,383	24,626	5,542	5,471	4,581	4,496
Part-Time	3,667 ^a	3,012	414	422	330	324
Unemployed	1,530	1,294	444	277	334	184
Other ¹	7,995	8,168	1,224	1,359	1,062	1,204

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.30B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	18.5	18.1	3.8	3.7	3.1	3.0
GENDER						
Male	33.0	32.6	7.2	7.3	6.1	6.0
Female	5.3	4.8	0.6	0.4	0.4	0.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	20.0	19.8	4.1	4.1	3.4	3.4
White	22.6	22.7	4.6	4.8	3.8	4.0
Black or African American	10.1	9.0	2.7 ^a	1.5	2.2	1.3
American Indian or Alaska Native	28.2	29.1	6.9	6.5	4.4	5.8
Native Hawaiian or Other Pacific Islander	*	9.3	5.0	2.7	4.4	2.7
Asian	5.0	4.6	0.8	0.8	0.7	0.5
Two or More Races	24.6	24.6	3.9	3.9	2.6	2.7
Hispanic or Latino	9.5	7.9	1.6	1.0	1.3	0.7
EDUCATION						
< High School	15.8	15.7	4.6	4.5	3.6	3.8
High School Graduate	18.5	19.0	4.8	4.6	4.1	3.9
Some College	19.6	18.7	3.9	3.6	3.3	3.0
College Graduate	18.8	17.8	2.4	2.6	2.0	2.0
CURRENT EMPLOYMENT						
Full-Time	23.1	22.7	5.3	5.0	4.3	4.1
Part-Time	15.2 ^a	12.8	1.7	1.8	1.4	1.4
Unemployed	20.0	18.7	5.8	4.0	4.4	2.7
Other ¹	12.2	12.3	1.9	2.1	1.6	1.8

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.31A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics:
Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	44,638	43,876	10,856	10,619	8,326	8,173
GENDER						
Male	37,712	37,406	9,947	9,958	7,773	7,788
Female	6,927	6,470	909 ^a	661	553	384
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	41,130	40,730	10,090	10,026	7,822	7,828
White	36,584	36,416	8,962	9,135	6,975	7,175
Black or African American	2,572	2,389	697	450	548	342
American Indian or Alaska Native	367	393	102	102	67	91
Native Hawaiian or Other Pacific Islander	*	112	36	37	31	30
Asian	613	572	126	112	87	70
Two or More Races	843	849	168	190	114	121
Hispanic or Latino	3,508	3,145	765	592	504	344
EDUCATION						
< High School	5,350	5,370	1,815	1,792	1,303	1,398
High School Graduate	13,302	13,313	4,038	3,800	3,141	2,976
Some College	12,378	12,260	3,042	2,924	2,329	2,241
College Graduate	13,608	12,932	1,962	2,102	1,553	1,558
CURRENT EMPLOYMENT						
Full-Time	27,849	28,110	7,156	7,079	5,662	5,635
Part-Time	5,345 ^a	4,574	1,136	1,139	730	707
Unemployed	2,421	2,040	879 ^a	608	581 ^a	369
Other ¹	9,022	9,152	1,684	1,792	1,354	1,461

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.31B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	18.8	18.3	4.6	4.4	3.5	3.4
GENDER						
Male	33.0	32.3	8.7	8.6	6.8	6.7
Female	5.6	5.2	0.7 ^a	0.5	0.4	0.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	20.4	20.0	5.0	4.9	3.9	3.8
White	23.4	23.2	5.7	5.8	4.5	4.6
Black or African American	9.3	8.5	2.5	1.6	2.0	1.2
American Indian or Alaska Native	29.6	30.8	8.2	8.0	5.4	7.1
Native Hawaiian or Other Pacific Islander	*	12.2	4.7	4.0	4.0	3.3
Asian	5.1	4.5	1.0	0.9	0.7	0.6
Two or More Races	23.8	23.3	4.8	5.2	3.2	3.3
Hispanic or Latino	9.8	8.5	2.1	1.6	1.4	0.9
EDUCATION						
< High School	16.6	16.7	5.6	5.6	4.1	4.3
High School Graduate	19.0	19.3	5.8	5.5	4.5	4.3
Some College	19.5	18.7	4.8	4.5	3.7	3.4
College Graduate	18.9	17.6	2.7	2.9	2.2	2.1
CURRENT EMPLOYMENT						
Full-Time	23.4	22.9	6.0	5.8	4.8	4.6
Part-Time	15.8 ^a	13.9	3.4	3.5	2.2	2.1
Unemployed	20.5	18.9	7.4 ^a	5.6	4.9 ^a	3.4
Other ¹	12.4	12.5	2.3	2.4	1.9	2.0

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.32A Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	89,117	89,267	24,933	23,734	12,438	11,964
AGE						
12-17	2,070	1,985	1,424	1,426	576	521
18-25	13,379	13,170	7,214	6,975	3,472	3,392
26 or Older	73,668	74,112	16,296	15,333	8,389	8,051
GENDER						
Male	64,644	65,196	18,848	18,424	9,755	9,580
Female	24,473	24,071	6,085 ^b	5,310	2,683	2,384
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	79,189	79,416	21,656	20,801	10,894	10,412
White	67,891	67,355	17,216	16,615	8,140	7,811
Black or African American	7,513	7,527	3,212	2,922	2,144	2,062
American Indian or Alaska Native	434	531	178	131	86	60
Native Hawaiian or Other Pacific Islander	245	241	58	77	19	32
Asian	1,675	2,031	521	501	270	160
Two or More Races	1,431	1,730	471	556	236	287
Hispanic or Latino	9,928	9,852	3,277	2,933	1,544	1,551

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.32B Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	34.0	33.7	9.5 ^a	9.0	4.7	4.5
AGE						
12-17	8.3	8.0	5.7	5.7	2.3	2.1
18-25	38.5	37.7	20.7	20.0	10.0	9.7
26 or Older	36.3	36.1	8.0 ^a	7.5	4.1	3.9
GENDER						
Male	50.9	50.8	14.8	14.3	7.7	7.5
Female	18.1	17.6	4.5 ^b	3.9	2.0	1.7
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	35.8	35.7	9.8	9.3	4.9	4.7
White	39.9	39.6	10.1	9.8	4.8	4.6
Black or African American	24.1	23.8	10.3	9.2	6.9	6.5
American Indian or Alaska Native	31.0	36.8	12.7	9.1	6.1	4.2
Native Hawaiian or Other Pacific Islander	28.1	23.6	6.7	7.5	2.1	3.2
Asian	12.5	14.7	3.9	3.6	2.0	1.2
Two or More Races	33.3	39.2	11.0	12.6	5.5	6.5
Hispanic or Latino	24.1	23.2	7.9	6.9	3.7	3.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.33A Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	2,070	1,985	1,424	1,426	576	521
GENDER						
Male	1,363	1,242	950	907	407	337
Female	707	744	474	519	169	184
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,679	1,617	1,166	1,188	477	458
White	1,353	1,308	961	970	385	363
Black or African American	212	205	128	142	55	74
American Indian or Alaska Native	19	16	11	8	3	2
Native Hawaiian or Other Pacific Islander	3	*	3	*	2	*
Asian	30	19	27	17	13	5
Two or More Races	61	58	37	42	19	14
Hispanic or Latino	390	369	258	238	99	63
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	931	878	670	671	286	238
Female, White, Not Hispanic	422	430	291	299	99	125
Male, Black, Not Hispanic	132	126	80	89	31	49
Female, Black, Not Hispanic	80	79	48	53	25	24
Male, Hispanic	230	189	152	112	61	39
Female, Hispanic	160	180	106	125	38	24

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.33B Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8.3	8.0	5.7	5.7	2.3	2.1
GENDER						
Male	10.7	9.8	7.5	7.1	3.2	2.7
Female	5.8	6.1	3.9	4.3	1.4	1.5
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8.7	8.4	6.0	6.2	2.5	2.4
White	9.9	9.7	7.1	7.2	2.8	2.7
Black or African American	6.2	5.9	3.7	4.1	1.6	2.1
American Indian or Alaska Native	12.1	9.6	6.6	5.1	1.9	1.3
Native Hawaiian or Other Pacific Islander	3.4	*	2.7	*	1.9	*
Asian	2.4	1.5	2.2	1.4	1.0	0.4
Two or More Races	8.1	7.6	4.9	5.5	2.5	1.8
Hispanic or Latino	7.0	6.5	4.6	4.2	1.8	1.1
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	13.4	12.7	9.6	9.7	4.1	3.5
Female, White, Not Hispanic	6.3	6.5	4.4	4.5	1.5	1.9
Male, Black, Not Hispanic	7.5	7.1	4.5	5.0	1.7	2.8
Female, Black, Not Hispanic	4.8	4.6	2.8	3.1	1.5	1.4
Male, Hispanic	8.1	6.6	5.3	3.9	2.2	1.4
Female, Hispanic	5.9	6.5	3.9	4.5	1.4	0.9

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.34A Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	13,379	13,170	7,214	6,975	3,472	3,392
GENDER						
Male	8,726	8,498	5,229 ^a	4,880	2,602	2,471
Female	4,654	4,672	1,985	2,096	870	921
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	11,122	11,086	6,031	5,992	2,970	2,853
White	9,084	8,989	4,815	4,629	2,248	2,082
Black or African American	1,251	1,218	803	870	540	576
American Indian or Alaska Native	77	89	40	54	22	29
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	368	371	184	190	64	52
Two or More Races	289	352	153	205	87	96
Hispanic or Latino	2,258	2,084	1,183 ^a	983	503	539
EDUCATION						
< High School	1,829	1,643	1,126	1,072	635	684
High School Graduate	4,192	4,069	2,412	2,292	1,154	1,182
Some College	4,926	5,006	2,636	2,631	1,250	1,172
College Graduate	2,432	2,451	1,040	980	433	353
CURRENT EMPLOYMENT						
Full-Time	6,051	6,089	3,119	3,004	1,500	1,558
Part-Time	3,673	3,583	2,022	1,981	940	873
Unemployed	1,529	1,425	945	885	554	474
Other ¹	2,126	2,073	1,128	1,105	478	487

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.34B Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	38.5	37.7	20.7	20.0	10.0	9.7
GENDER						
Male	50.0	48.5	29.9 ^a	27.8	14.9	14.1
Female	26.9	26.8	11.5	12.0	5.0	5.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	40.4	40.2	21.9	21.7	10.8	10.3
White	46.6	46.4	24.7	23.9	11.5	10.8
Black or African American	25.0	24.2	16.1	17.3	10.8	11.4
American Indian or Alaska Native	33.6	44.1	17.5	26.6	9.7	14.3
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	19.7	18.4	9.8	9.4	3.4	2.6
Two or More Races	39.4	44.8	20.9	26.2	11.9	12.2
Hispanic or Latino	31.1	28.3	16.3 ^a	13.3	6.9	7.3
EDUCATION						
< High School	32.7	33.2	20.1	21.7	11.4 ^a	13.8
High School Graduate	35.0	34.0	20.1	19.1	9.6	9.9
Some College	40.4	40.2	21.6	21.1	10.3	9.4
College Graduate	48.2 ^a	44.0	20.6 ^a	17.6	8.6 ^a	6.3
CURRENT EMPLOYMENT						
Full-Time	44.8	42.7	23.1 ^a	21.1	11.1	10.9
Part-Time	38.0	37.8	20.9	20.9	9.7	9.2
Unemployed	36.8	36.6	22.7	22.8	13.3	12.2
Other ¹	28.5	28.3	15.1	15.1	6.4	6.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.35A Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	73,668	74,112	16,296	15,333	8,389	8,051
GENDER						
Male	54,556	55,457	12,669	12,638	6,745	6,772
Female	19,112	18,655	3,627 ^b	2,695	1,644 ^a	1,279
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	66,388	66,713	14,459	13,621	7,448	7,102
White	57,454	57,058	11,440	11,016	5,507	5,366
Black or African American	6,050	6,104	2,280	1,910	1,549	1,413
American Indian or Alaska Native	338	426	127	68	61	29
Native Hawaiian or Other Pacific Islander	*	163	19	24	*	13
Asian	1,277	1,641	310	295	193	103
Two or More Races	1,081	1,321	281	308	130	178
Hispanic or Latino	7,280	7,399	1,836	1,713	941	949
EDUCATION						
< High School	6,600	6,858	2,006	1,887	1,327	1,232
High School Graduate	18,353	18,364	4,188	3,656	2,193	2,142
Some College	19,671	20,090	4,523	4,306	2,327	2,406
College Graduate	29,044	28,800	5,579	5,484	2,543	2,272
CURRENT EMPLOYMENT						
Full-Time	45,040	45,231	10,857	10,425	5,140	5,026
Part-Time	8,168	7,910	1,618	1,335	742	681
Unemployed	2,992 ^a	2,457	990	745	731	512
Other ¹	17,469	18,515	2,831	2,828	1,776	1,832

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.35B Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	36.3	36.1	8.0 ^a	7.5	4.1	3.9
GENDER						
Male	56.3	56.5	13.1	12.9	7.0	6.9
Female	18.1	17.4	3.4 ^b	2.5	1.6 ^a	1.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	38.1	37.9	8.3	7.7	4.3	4.0
White	42.0	41.5	8.4	8.0	4.0	3.9
Black or African American	26.5	26.4	10.0	8.2	6.8	6.1
American Indian or Alaska Native	33.4	39.7	12.6	6.3	6.0	2.7
Native Hawaiian or Other Pacific Islander	*	22.0	3.4	3.2	*	1.8
Asian	12.5	15.5	3.0	2.8	1.9	1.0
Two or More Races	38.5	46.1	10.0	10.8	4.6	6.2
Hispanic or Latino	25.6	25.1	6.5	5.8	3.3	3.2
EDUCATION						
< High School	24.8	25.1	7.5	6.9	5.0	4.5
High School Graduate	31.6	32.3	7.2	6.4	3.8	3.8
Some College	38.4	37.8	8.8	8.1	4.5	4.5
College Graduate	43.4	42.4	8.3	8.1	3.8	3.3
CURRENT EMPLOYMENT						
Full-Time	42.7	41.6	10.3	9.6	4.9	4.6
Part-Time	33.9	33.6	6.7	5.7	3.1	2.9
Unemployed	39.1	35.6	12.9	10.8	9.6	7.4
Other ¹	26.7	28.0	4.3	4.3	2.7	2.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.36A Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	87,047	87,282	23,509	22,309	11,862	11,443
GENDER						
Male	63,281	63,955	17,898	17,518	9,348	9,243
Female	23,766	23,327	5,611 ^b	4,791	2,514	2,200
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	77,509	77,799	20,491	19,613	10,417	9,955
White	66,538	66,047	16,255	15,645	7,755	7,448
Black or African American	7,300	7,322	3,084	2,780	2,089	1,988
American Indian or Alaska Native	415	515	168	122	83	58
Native Hawaiian or Other Pacific Islander	*	230	55	68	17	32
Asian	1,645	2,012	494	485	258	155
Two or More Races	1,370	1,672	435	514	217	273
Hispanic or Latino	9,538	9,483	3,019	2,696	1,444	1,488
EDUCATION						
< High School	8,430	8,501	3,132	2,960	1,962	1,915
High School Graduate	22,545	22,434	6,600	5,948	3,347	3,324
Some College	24,596	25,096	7,159	6,937	3,576	3,578
College Graduate	31,476	31,251	6,619	6,464	2,976	2,625
CURRENT EMPLOYMENT						
Full-Time	51,091	51,320	13,977	13,430	6,640	6,584
Part-Time	11,841	11,493	3,640	3,316	1,682	1,554
Unemployed	4,521 ^a	3,882	1,934	1,630	1,285 ^a	985
Other ¹	19,594	20,587	3,959	3,933	2,254	2,319

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.36B Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	36.7	36.3	9.9 ^a	9.3	5.0	4.8
GENDER						
Male	55.3	55.3	15.6	15.1	8.2	8.0
Female	19.3	18.7	4.6 ^b	3.8	2.0	1.8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	38.4	38.2	10.2	9.6	5.2	4.9
White	42.6	42.1	10.4	10.0	5.0	4.8
Black or African American	26.3	26.0	11.1	9.9	7.5	7.1
American Indian or Alaska Native	33.4	40.4	13.5	9.6	6.7	4.5
Native Hawaiian or Other Pacific Islander	*	25.1	7.2	7.4	2.2	3.5
Asian	13.6	15.9	4.1	3.8	2.1	1.2
Two or More Races	38.7	45.8	12.3	14.1	6.1	7.5
Hispanic or Latino	26.7	25.8	8.5	7.3	4.0	4.0
EDUCATION						
< High School	26.2	26.4	9.7	9.2	6.1	5.9
High School Graduate	32.2	32.6	9.4	8.6	4.8	4.8
Some College	38.8	38.3	11.3	10.6	5.6	5.5
College Graduate	43.7	42.5	9.2	8.8	4.1	3.6
CURRENT EMPLOYMENT						
Full-Time	43.0	41.7	11.7 ^a	10.9	5.6	5.4
Part-Time	35.1	34.8	10.8	10.0	5.0	4.7
Unemployed	38.3	36.0	16.4	15.1	10.9	9.1
Other ¹	26.8	28.0	5.4	5.4	3.1	3.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.37A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	213,794 ^b	217,765	173,966 ^a	176,556	136,868 ^a	139,677
AGE						
12-17	7,669	7,375	6,133	5,979	2,877	2,872
18-25	29,148	29,126	26,707	26,713	20,738	20,815
26 or Older	176,977 ^b	181,265	141,126 ^a	143,864	113,253 ^a	115,990
GENDER						
Male	107,186 ^a	108,611	88,185	88,879	72,605	73,559
Female	106,608 ^b	109,155	85,781 ^a	87,676	64,263	66,118
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	183,706 ^b	186,429	150,028	151,107	119,112	120,828
White	147,073	148,157	121,197	120,942	98,001	98,288
Black or African American	23,363	23,976	18,320	18,623	13,612	13,989
American Indian or Alaska Native	1,097	1,131	747	789	522	611
Native Hawaiian or Other Pacific Islander	623	726	470	532	334	387
Asian	8,102	8,923	6,683	7,359	4,607	5,365
Two or More Races	3,448	3,516	2,612	2,862	2,035	2,188
Hispanic or Latino	30,088 ^b	31,336	23,938 ^b	25,449	17,755 ^a	18,849

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.37B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	81.5	82.1	66.3	66.6	52.2	52.7
AGE						
12-17	30.8	29.6	24.6	24.0	11.6	11.5
18-25	83.8	83.4	76.8	76.5	59.6	59.6
26 or Older	87.3 ^a	88.3	69.6	70.1	55.9	56.5
GENDER						
Male	84.3	84.6	69.4	69.2	57.1	57.3
Female	78.8	79.8	63.4	64.1	47.5	48.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	83.1	83.7	67.8	67.9	53.9	54.3
White	86.5	87.0	71.3	71.0	57.7	57.7
Black or African American	74.8	75.7	58.7	58.8	43.6	44.2
American Indian or Alaska Native	78.3	78.4	53.3	54.7	37.3	42.3
Native Hawaiian or Other Pacific Islander	71.5	71.1	54.0	52.1	38.4	37.9
Asian	60.7	64.4	50.1	53.1	34.5	38.7
Two or More Races	80.4	79.6	60.9	64.8	47.4	49.5
Hispanic or Latino	72.9	73.8	58.0	59.9	43.0	44.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.38A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	7,669	7,375	6,133	5,979	2,877	2,872
GENDER						
Male	3,857 ^a	3,584	2,985	2,794	1,429	1,369
Female	3,812	3,791	3,148	3,185	1,448	1,503
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5,958 ^a	5,669	4,843	4,627	2,284	2,238
White	4,399	4,234	3,689	3,598	1,758	1,774
Black or African American	1,022	940	743	652	335	300
American Indian or Alaska Native	47	48	32	37	15	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	8	*
Asian	238	206	197	151	100	56
Two or More Races	222	216	164	170	68	75
Hispanic or Latino	1,711	1,706	1,290	1,352	593	634
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	2,226	2,098	1,809	1,708	878	875
Female, White, Not Hispanic	2,173	2,136	1,880	1,889	880	899
Male, Black, Not Hispanic	500	449	350	296	153	122
Female, Black, Not Hispanic	523	490	393	355	182	177
Male, Hispanic	857	791	624	601	297	290
Female, Hispanic	854	915	666	751	296	344

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.38B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	30.8	29.6	24.6	24.0	11.6	11.5
GENDER						
Male	30.4 ^a	28.3	23.5	22.0	11.2	10.8
Female	31.3	31.1	25.8	26.1	11.9	12.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	30.8 ^a	29.5	25.1	24.1	11.8	11.6
White	32.3	31.4	27.1	26.7	12.9	13.2
Black or African American	29.7	27.0	21.6 ^a	18.7	9.7	8.6
American Indian or Alaska Native	29.3	28.7	20.3	22.3	9.3	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	8.2	*
Asian	19.1	16.8	15.8	12.3	8.0 ^a	4.5
Two or More Races	29.4	28.2	21.8	22.1	9.0	9.7
Hispanic or Latino	30.8	30.2	23.2	23.9	10.7	11.2
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	32.0	30.5	26.0	24.8	12.6	12.7
Female, White, Not Hispanic	32.5	32.4	28.2	28.7	13.2	13.6
Male, Black, Not Hispanic	28.5	25.4	19.9	16.8	8.7	6.9
Female, Black, Not Hispanic	31.0	28.6	23.3	20.7	10.8	10.3
Male, Hispanic	30.2	27.5	21.9	20.9	10.4	10.1
Female, Hispanic	31.4	33.0	24.5	27.1	10.9	12.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.39A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	29,148	29,126	26,707	26,713	20,738	20,815
GENDER						
Male	14,621	14,546	13,530	13,439	10,875	10,793
Female	14,527	14,580	13,178	13,274	9,864	10,021
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	23,267	23,256	21,513	21,509	16,907	16,917
White	17,125	16,982	15,991	15,821	12,903	12,789
Black or African American	3,873	3,862	3,477	3,480	2,531	2,454
American Indian or Alaska Native	181	161	154	139	116	108
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1,274	1,471	1,148	1,347	812	988
Two or More Races	658	655	604	610	433	497
Hispanic or Latino	5,881	5,870	5,195	5,204	3,831	3,898
EDUCATION						
< High School	4,181 ^b	3,604	3,586 ^b	3,025	2,408 ^b	2,066
High School Graduate	9,529	9,382	8,492	8,435	6,072	6,113
Some College	10,691	10,902	10,024	10,180	8,113	8,252
College Graduate	4,747 ^a	5,237	4,605	5,073	4,145	4,384
CURRENT EMPLOYMENT						
Full-Time	12,210	12,844	11,384 ^a	12,084	9,287 ^a	9,868
Part-Time	8,083	7,964	7,509	7,328	5,855	5,575
Unemployed	3,428 ^a	3,054	3,109 ^a	2,759	2,225	2,024
Other ¹	5,426	5,263	4,705	4,542	3,372	3,347

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.39B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	83.8	83.4	76.8	76.5	59.6	59.6
GENDER						
Male	83.7	83.0	77.5	76.7	62.3	61.6
Female	83.9	83.8	76.1	76.3	56.9	57.6
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	84.5	84.4	78.1	78.0	61.4	61.4
White	87.8	87.7	82.0	81.7	66.2	66.0
Black or African American	77.4	76.7	69.5	69.1	50.6	48.7
American Indian or Alaska Native	78.8	79.2	67.1	68.3	50.4	53.1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	68.1	73.2	61.4	67.0	43.4	49.2
Two or More Races	89.7 ^a	83.5	82.5	77.8	59.1	63.4
Hispanic or Latino	81.1	79.7	71.6	70.7	52.8	52.9
EDUCATION						
< High School	74.8	72.9	64.2	61.2	43.1	41.8
High School Graduate	79.6	78.3	70.9	70.4	50.7	51.1
Some College	87.8	87.6	82.3	81.8	66.6	66.3
College Graduate	94.1	94.0	91.3	91.0	82.1 ^a	78.7
CURRENT EMPLOYMENT						
Full-Time	90.4	90.1	84.2	84.8	68.7	69.2
Part-Time	83.6	84.0	77.7	77.3	60.6	58.8
Unemployed	82.5 ^a	78.5	74.8 ^a	71.0	53.5	52.1
Other ¹	72.8	72.0	63.2	62.1	45.3	45.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.40A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	176,977 ^b	181,265	141,126 ^a	143,864	113,253 ^a	115,990
GENDER						
Male	88,708 ^b	90,481	71,671	72,646	60,302	61,396
Female	88,269 ^b	90,783	69,455 ^a	71,218	52,951	54,594
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	154,482 ^b	157,504	123,673	124,971	99,921	101,673
White	125,549	126,941	101,517	101,523	83,340	83,725
Black or African American	18,468	19,174	14,100	14,492	10,746	11,236
American Indian or Alaska Native	869	923	560	613	392	477
Native Hawaiian or Other Pacific Islander	*	575	*	*	*	*
Asian	6,590	7,245	5,338	5,862	3,695	4,321
Two or More Races	2,569	2,645	1,843	2,083	1,534	1,616
Hispanic or Latino	22,495 ^b	23,761	17,453 ^b	18,893	13,332 ^a	14,316
EDUCATION						
< High School	19,097	20,193	12,841	13,301	9,345	9,440
High School Graduate	49,467	49,597	36,660	36,962	27,788	28,583
Some College	46,847	48,778	37,922	39,026	30,445	31,480
College Graduate	61,567	62,697	53,702	54,575	45,674	46,487
CURRENT EMPLOYMENT						
Full-Time	96,954	100,000	83,124	85,414	69,029	70,683
Part-Time	21,362	20,984	17,054	16,908	13,706	13,743
Unemployed	6,853	6,119	5,408	4,978	4,130	3,750
Other ¹	51,807	54,161	35,539	36,564	26,388	27,814

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.40B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	87.3 ^a	88.3	69.6	70.1	55.9	56.5
GENDER						
Male	91.5	92.1	73.9	74.0	62.2	62.5
Female	83.5	84.8	65.7	66.5	50.1	51.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	88.6 ^a	89.6	71.0	71.1	57.3	57.8
White	91.7	92.4	74.2	73.9	60.9	60.9
Black or African American	81.1	82.8	61.9	62.6	47.2	48.5
American Indian or Alaska Native	85.9	86.0	55.4	57.1	38.7	44.5
Native Hawaiian or Other Pacific Islander	*	77.6	*	*	*	*
Asian	64.4	68.2	52.2	55.2	36.1	40.7
Two or More Races	91.6	92.3	65.7	72.6	54.7	56.4
Hispanic or Latino	79.1	80.7	61.4	64.1	46.9	48.6
EDUCATION						
< High School	71.8	74.0	48.3	48.8	35.2	34.6
High School Graduate	85.3 ^a	87.1	63.2	64.9	47.9 ^a	50.2
Some College	91.5	91.8	74.0	73.4	59.4	59.2
College Graduate	92.0	92.3	80.3	80.3	68.3	68.4
CURRENT EMPLOYMENT						
Full-Time	92.0	92.0	78.8	78.6	65.5	65.0
Part-Time	88.7	89.2	70.8	71.8	56.9	58.4
Unemployed	89.6	88.6	70.7	72.1	54.0	54.3
Other ¹	79.0 ^b	81.8	54.2	55.2	40.3	42.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.41A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	206,125 ^b	210,390	167,833 ^a	170,577	133,991 ^a	136,804
GENDER						
Male	103,329 ^b	105,027	85,200	86,085	71,176	72,189
Female	102,796 ^b	105,363	82,633 ^a	84,492	62,815	64,615
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	177,748 ^b	180,760	145,185	146,480	116,829	118,590
White	142,674	143,923	117,508	117,344	96,243	96,514
Black or African American	22,341	23,036	17,577	17,971	13,277	13,689
American Indian or Alaska Native	1,050	1,084	714	752	508	585
Native Hawaiian or Other Pacific Islander	*	701	*	511	326	379
Asian	7,864	8,717	6,486	7,208	4,507	5,309
Two or More Races	3,226	3,300	2,448	2,693	1,967	2,113
Hispanic or Latino	28,376 ^b	29,630	22,648 ^b	24,097	17,162 ^a	18,214
EDUCATION						
< High School	23,278	23,797	16,428	16,327	11,753	11,506
High School Graduate	58,995	58,979	45,152	45,397	33,861	34,696
Some College	57,538	59,680	47,946	49,206	38,558	39,732
College Graduate	66,314	67,934	58,308	59,647	49,819	50,871
CURRENT EMPLOYMENT						
Full-Time	109,164	112,845	94,509	97,498	78,316	80,551
Part-Time	29,446	28,949	24,563	24,236	19,561	19,318
Unemployed	10,282 ^a	9,173	8,517 ^a	7,737	6,354	5,774
Other ¹	57,233	59,424	40,244	41,106	29,760	31,160

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.41B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	86.8	87.6	70.7	71.0	56.4	56.9
GENDER						
Male	90.3	90.7	74.5	74.4	62.2	62.4
Female	83.5	84.6	67.1	67.9	51.0	51.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	88.1	88.9	71.9	72.0	57.9	58.3
White	91.3	91.8	75.2	74.9	61.6	61.6
Black or African American	80.4	81.7	63.2	63.7	47.8	48.6
American Indian or Alaska Native	84.6	84.9	57.5	58.9	40.9	45.8
Native Hawaiian or Other Pacific Islander	*	76.3	*	55.7	42.2	41.3
Asian	65.0	69.0	53.6	57.1	37.2	42.0
Two or More Races	91.2	90.4	69.2	73.8	55.6	57.9
Hispanic or Latino	79.5	80.5	63.4	65.4	48.1	49.5
EDUCATION						
< High School	72.4	73.8	51.1	50.7	36.5	35.7
High School Graduate	84.3	85.6	64.5	65.9	48.4 ^a	50.3
Some College	90.8	91.0	75.6	75.0	60.8	60.6
College Graduate	92.2	92.4	81.0	81.1	69.2	69.2
CURRENT EMPLOYMENT						
Full-Time	91.8	91.8	79.5	79.3	65.8	65.5
Part-Time	87.2	87.7	72.8	73.4	58.0	58.5
Unemployed	87.1	85.0	72.1	71.7	53.8	53.5
Other ¹	78.4 ^b	80.8	55.1	55.9	40.8	42.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.42A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	136,868 ^a	139,677	60,064	60,941	16,541	16,342
AGE						
12-17	2,877	2,872	1,556	1,512	293	257
18-25	20,738	20,815	13,187	13,184	3,939	3,782
26 or Older	113,253 ^a	115,990	45,322	46,245	12,310	12,303
GENDER						
Male	72,605	73,559	38,379	38,586	12,093	11,938
Female	64,263	66,118	21,685	22,355	4,448	4,404
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	119,112	120,828	50,115	50,456	14,544	14,164
White	98,001	98,288	40,790	40,076	12,395	12,022
Black or African American	13,612	13,989	6,281	6,857	1,406	1,434
American Indian or Alaska Native	522	611	330	400	82	132
Native Hawaiian or Other Pacific Islander	334	387	215	187	77	47
Asian	4,607	5,365	1,657	2,010	264	272
Two or More Races	2,035	2,188	842	926	320	256
Hispanic or Latino	17,755 ^a	18,849	9,949	10,486	1,997	2,179

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.42B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	52.2	52.7	22.9	23.0	6.3	6.2
AGE						
12-17	11.6	11.5	6.2	6.1	1.2	1.0
18-25	59.6	59.6	37.9	37.7	11.3	10.8
26 or Older	55.9	56.5	22.4	22.5	6.1	6.0
GENDER						
Male	57.1	57.3	30.2	30.0	9.5	9.3
Female	47.5	48.4	16.0	16.4	3.3	3.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	53.9	54.3	22.7	22.7	6.6	6.4
White	57.7	57.7	24.0	23.5	7.3	7.1
Black or African American	43.6	44.2	20.1	21.6	4.5	4.5
American Indian or Alaska Native	37.3	42.3	23.5	27.7	5.8	9.2
Native Hawaiian or Other Pacific Islander	38.4	37.9	24.7	18.3	8.9	4.6
Asian	34.5	38.7	12.4	14.5	2.0	2.0
Two or More Races	47.4	49.5	19.6	21.0	7.5	5.8
Hispanic or Latino	43.0	44.4	24.1	24.7	4.8	5.1

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.43A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	2,877	2,872	1,556	1,512	293	257
GENDER						
Male	1,429	1,369	845	813	173	147
Female	1,448	1,503	711	699	119	111
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,284	2,238	1,207	1,158	242	197
White	1,758	1,774	990	952	212	178
Black or African American	335	300	133	126	16	8
American Indian or Alaska Native	15	*	9	*	2	*
Native Hawaiian or Other Pacific Islander	8	*	4	*	0	*
Asian	100	56	35	18	3	0
Two or More Races	68	75	36	39	9	5
Hispanic or Latino	593	634	348	354	51	60
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	878	875	554	541	128	106
Female, White, Not Hispanic	880	899	437	412	84	72
Male, Black, Not Hispanic	153	122	65	55	15	2
Female, Black, Not Hispanic	182	177	69	71	0	6
Male, Hispanic	297	290	187	178	26	31
Female, Hispanic	296	344	162	176	24	30

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.43B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	11.6	11.5	6.2	6.1	1.2	1.0
GENDER						
Male	11.2	10.8	6.6	6.4	1.4	1.2
Female	11.9	12.3	5.8	5.7	1.0	0.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	11.8	11.6	6.2	6.0	1.3	1.0
White	12.9	13.2	7.3	7.1	1.6	1.3
Black or African American	9.7	8.6	3.9	3.6	0.5	0.2
American Indian or Alaska Native	9.3	*	5.6	*	1.5	*
Native Hawaiian or Other Pacific Islander	8.2	*	4.5	*	0.2	*
Asian	8.0 ^a	4.5	2.8	1.5	0.2	0.0
Two or More Races	9.0	9.7	4.7	5.1	1.3	0.6
Hispanic or Latino	10.7	11.2	6.3	6.3	0.9	1.1
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	12.6	12.7	8.0	7.9	1.8	1.5
Female, White, Not Hispanic	13.2	13.6	6.5	6.2	1.3	1.1
Male, Black, Not Hispanic	8.7	6.9	3.7	3.1	0.9	0.1
Female, Black, Not Hispanic	10.8	10.3	4.1	4.1	0.0	0.3
Male, Hispanic	10.4	10.1	6.6	6.2	0.9	1.1
Female, Hispanic	10.9	12.4	5.9	6.3	0.9	1.1

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.44A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	20,738	20,815	13,187	13,184	3,939	3,782
GENDER						
Male	10,875	10,793	7,750	7,650	2,659	2,497
Female	9,864	10,021	5,437	5,534	1,280	1,286
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	16,907	16,917	10,676	10,652	3,318	3,148
White	12,903	12,789	8,483	8,382	2,877	2,647
Black or African American	2,531	2,454	1,331	1,359	256	277
American Indian or Alaska Native	116	108	76	71	19	23
Native Hawaiian or Other Pacific Islander	*	*	*	*	15	*
Asian	812	988	446	523	83	93
Two or More Races	433	497	272	272	67	101
Hispanic or Latino	3,831	3,898	2,510	2,532	621	634
EDUCATION						
< High School	2,408 ^b	2,066	1,685 ^a	1,445	450	418
High School Graduate	6,072	6,113	3,981	4,099	1,094	1,101
Some College	8,113	8,252	5,052	4,968	1,631	1,520
College Graduate	4,145	4,384	2,469	2,673	764	744
CURRENT EMPLOYMENT						
Full-Time	9,287 ^a	9,868	5,961 ^a	6,405	1,885	1,866
Part-Time	5,855	5,575	3,660	3,356	1,082	954
Unemployed	2,225	2,024	1,452	1,400	394	392
Other ¹	3,372	3,347	2,115	2,023	577	571

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.44B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	59.6	59.6	37.9	37.7	11.3	10.8
GENDER						
Male	62.3	61.6	44.4	43.6	15.2	14.2
Female	56.9	57.6	31.4	31.8	7.4	7.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	61.4	61.4	38.8	38.6	12.1	11.4
White	66.2	66.0	43.5	43.3	14.8	13.7
Black or African American	50.6	48.7	26.6	27.0	5.1	5.5
American Indian or Alaska Native	50.4	53.1	33.0	35.0	8.3	11.2
Native Hawaiian or Other Pacific Islander	*	*	*	*	7.6	*
Asian	43.4	49.2	23.8	26.0	4.5	4.6
Two or More Races	59.1	63.4	37.1	34.8	9.2	12.9
Hispanic or Latino	52.8	52.9	34.6	34.4	8.6	8.6
EDUCATION						
< High School	43.1	41.8	30.2	29.2	8.0	8.4
High School Graduate	50.7	51.1	33.3	34.2	9.1	9.2
Some College	66.6	66.3	41.5	39.9	13.4	12.2
College Graduate	82.1 ^a	78.7	48.9	48.0	15.1	13.4
CURRENT EMPLOYMENT						
Full-Time	68.7	69.2	44.1	44.9	13.9	13.1
Part-Time	60.6	58.8	37.9	35.4	11.2	10.1
Unemployed	53.5	52.1	34.9	36.0	9.5	10.1
Other ¹	45.3	45.8	28.4	27.7	7.8	7.8

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.45A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Demographic
Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	113,253 ^a	115,990	45,322	46,245	12,310	12,303
GENDER						
Male	60,302	61,396	29,784	30,123	9,261	9,294
Female	52,951	54,594	15,538	16,122	3,049	3,008
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	99,921	101,673	38,232	38,645	10,984	10,818
White	83,340	83,725	31,317	30,741	9,307	9,197
Black or African American	10,746	11,236	4,816	5,373	1,134	1,149
American Indian or Alaska Native	392	477	245	312	60	104
Native Hawaiian or Other Pacific Islander	*	*	*	137	*	39
Asian	3,695	4,321	1,177	1,469	178	179
Two or More Races	1,534	1,616	534	614	244	151
Hispanic or Latino	13,332 ^a	14,316	7,090	7,600	1,325	1,485
EDUCATION						
< High School	9,345	9,440	5,779	5,969	1,420	1,701
High School Graduate	27,788	28,583	13,683	14,084	4,182	3,999
Some College	30,445	31,480	11,714	12,563	3,182	3,337
College Graduate	45,674	46,487	14,146	13,629	3,526	3,265
CURRENT EMPLOYMENT						
Full-Time	69,029	70,683	30,273	30,951	8,152	7,985
Part-Time	13,706	13,743	4,605	4,722	1,232	1,238
Unemployed	4,130	3,750	2,238	1,990	816	627
Other ¹	26,388	27,814	8,205	8,582	2,110	2,452

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.45B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	55.9	56.5	22.4	22.5	6.1	6.0
GENDER						
Male	62.2	62.5	30.7	30.7	9.6	9.5
Female	50.1	51.0	14.7	15.1	2.9	2.8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	57.3	57.8	21.9	22.0	6.3	6.2
White	60.9	60.9	22.9	22.4	6.8	6.7
Black or African American	47.2	48.5	21.1	23.2	5.0	5.0
American Indian or Alaska Native	38.7	44.5	24.2	29.0	5.9	9.7
Native Hawaiian or Other Pacific Islander	*	*	*	18.5	*	5.2
Asian	36.1	40.7	11.5	13.8	1.7	1.7
Two or More Races	54.7	56.4	19.1	21.4	8.7	5.3
Hispanic or Latino	46.9	48.6	24.9	25.8	4.7	5.0
EDUCATION						
< High School	35.2	34.6	21.7	21.9	5.3	6.2
High School Graduate	47.9 ^a	50.2	23.6	24.7	7.2	7.0
Some College	59.4	59.2	22.9	23.6	6.2	6.3
College Graduate	68.3	68.4	21.1	20.1	5.3	4.8
CURRENT EMPLOYMENT						
Full-Time	65.5	65.0	28.7	28.5	7.7	7.3
Part-Time	56.9	58.4	19.1	20.1	5.1	5.3
Unemployed	54.0	54.3	29.3	28.8	10.7	9.1
Other ¹	40.3	42.0	12.5	13.0	3.2	3.7

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.46A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic
Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	133,991 ^a	136,804	58,508	59,429	16,248	16,085
GENDER						
Male	71,176	72,189	37,534	37,773	11,920	11,791
Female	62,815	64,615	20,975	21,656	4,328	4,294
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	116,829	118,590	48,908	49,297	14,302	13,966
White	96,243	96,514	39,800	39,123	12,184	11,843
Black or African American	13,277	13,689	6,147	6,732	1,390	1,426
American Indian or Alaska Native	508	585	321	383	79	127
Native Hawaiian or Other Pacific Islander	326	379	211	182	77	47
Asian	4,507	5,309	1,623	1,992	261	272
Two or More Races	1,967	2,113	806	886	311	252
Hispanic or Latino	17,162 ^a	18,214	9,600	10,132	1,946	2,119
EDUCATION						
< High School	11,753	11,506	7,464	7,414	1,870	2,118
High School Graduate	33,861	34,696	17,664	18,182	5,276	5,100
Some College	38,558	39,732	16,766	17,531	4,813	4,857
College Graduate	49,819	50,871	16,614	16,302	4,290	4,009
CURRENT EMPLOYMENT						
Full-Time	78,316	80,551	36,233	37,356	10,037	9,851
Part-Time	19,561	19,318	8,265	8,078	2,314	2,192
Unemployed	6,354	5,774	3,690	3,390	1,210	1,020
Other ¹	29,760	31,160	10,320	10,605	2,687	3,023

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.46B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	56.4	56.9	24.6	24.7	6.8	6.7
GENDER						
Male	62.2	62.4	32.8	32.6	10.4	10.2
Female	51.0	51.9	17.0	17.4	3.5	3.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	57.9	58.3	24.2	24.2	7.1	6.9
White	61.6	61.6	25.5	25.0	7.8	7.6
Black or African American	47.8	48.6	22.1	23.9	5.0	5.1
American Indian or Alaska Native	40.9	45.8	25.9	30.0	6.4	9.9
Native Hawaiian or Other Pacific Islander	42.2	41.3	27.2	19.8	10.0	5.1
Asian	37.2	42.0	13.4	15.8	2.2	2.2
Two or More Races	55.6	57.9	22.8	24.3	8.8	6.9
Hispanic or Latino	48.1	49.5	26.9	27.5	5.5	5.8
EDUCATION						
< High School	36.5	35.7	23.2	23.0	5.8	6.6
High School Graduate	48.4 ^a	50.3	25.2	26.4	7.5	7.4
Some College	60.8	60.6	26.4	26.7	7.6	7.4
College Graduate	69.2	69.2	23.1	22.2	6.0	5.5
CURRENT EMPLOYMENT						
Full-Time	65.8	65.5	30.5	30.4	8.4	8.0
Part-Time	58.0	58.5	24.5	24.5	6.9	6.6
Unemployed	53.8	53.5	31.3	31.4	10.3	9.4
Other ¹	40.8	42.4	14.1	14.4	3.7	4.1

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.47A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	175,260	175,416	80,846	81,141	66,879	66,899
GEOGRAPHIC DIVISION						
Northeast	31,470	31,378	13,625	13,940	11,040	11,244
New England	8,440	8,595	3,427	3,494	2,667	2,771
Middle Atlantic	23,030	22,783	10,198	10,446	8,373	8,474
Midwest	40,009	40,071	19,653	19,127	16,485	16,056
East North Central	27,657	27,687	13,485	13,310	11,240	11,191
West North Central	12,352	12,384	6,168	5,817	5,245	4,865
South	65,126	65,120	31,268	31,850	26,270	26,873
South Atlantic	34,573	34,712	15,889	16,479	13,196	13,710
East South Central	10,552	10,645	5,598	5,742	4,961	4,991
West South Central	20,000	19,764	9,781	9,629	8,113	8,172
West	38,655	38,846	16,299	16,223	13,085	12,726
Mountain	12,591	12,458	5,545	5,408	4,537	4,406
Pacific	26,064	26,389	10,755	10,815	8,547	8,320
COUNTY TYPE						
Large Metro	90,220	93,446	39,839	41,734	32,136	33,430
Small Metro	54,563	52,485	25,424	24,548	21,229	20,458
250K – 1 Mil. Pop.	36,579	35,151	16,420	16,219	13,689	13,325
< 250K Pop.	17,984	17,334	9,004	8,329	7,540	7,133
Nonmetro	30,477	29,485	15,583	14,859	13,514	13,011
Urbanized	13,428	12,351	6,669	6,109	5,787	5,230
Less Urbanized	13,855	14,099	7,257	7,280	6,339	6,483
Completely Rural	3,194	3,036	1,657	1,470	1,388	1,299

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.47B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	66.8	66.2	30.8	30.6	25.5	25.2
GEOGRAPHIC DIVISION						
Northeast	66.4	65.9	28.8	29.3	23.3	23.6
New England	67.6	68.4	27.5	27.8	21.4	22.1
Middle Atlantic	66.0	65.0	29.2	29.8	24.0	24.2
Midwest	71.2	71.0	35.0	33.9	29.3	28.4
East North Central	71.0	70.8	34.6	34.0	28.8	28.6
West North Central	71.7	71.4	35.8	33.5	30.4	28.0
South	66.8	65.9	32.1	32.2	26.9	27.2
South Atlantic	67.2	66.5	30.9	31.6	25.6	26.3
East South Central	68.1	68.3	36.1	36.8	32.0	32.0
West South Central	65.4	63.6	32.0	31.0	26.5	26.3
West	63.1	62.5	26.6	26.1	21.4	20.5
Mountain	67.3	65.5	29.7	28.5	24.3	23.2
Pacific	61.2	61.1	25.3	25.0	20.1	19.3
COUNTY TYPE						
Large Metro	64.0	63.8	28.2	28.5	22.8	22.8
Small Metro	69.2	67.8	32.2	31.7	26.9	26.4
250K – 1 Mil. Pop.	68.5	67.1	30.8	31.0	25.6	25.4
< 250K Pop.	70.5	69.2	35.3	33.3	29.6	28.5
Nonmetro	71.8	71.6	36.7	36.1	31.8	31.6
Urbanized	70.7	71.6	35.1	35.4	30.5	30.3
Less Urbanized	73.0	71.6	38.2	37.0	33.4	32.9
Completely Rural	71.5	72.1	37.1	34.9	31.1	30.8

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.48A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	4,922 ^a	4,591	3,461 ^a	3,161	1,951 ^a	1,742
GEOGRAPHIC DIVISION						
Northeast	764 ^a	657	567 ^a	481	342 ^b	249
New England	181	179	132	147	80	73
Middle Atlantic	583 ^b	478	435 ^b	334	263 ^b	175
Midwest	1,120	1,042	804	713	456	429
East North Central	772	722	561	494	317	283
West North Central	348	320	243	219	139	146
South	1,946	1,852	1,354	1,282	750	705
South Atlantic	982 ^a	830	692	604	364	321
East South Central	354	377	252	246	151	149
West South Central	609	644	410	433	236	234
West	1,092	1,040	736	685	403	360
Mountain	413	355	278	248	146	160
Pacific	680	686	458	438	257	200
COUNTY TYPE						
Large Metro	2,302	2,220	1,626	1,499	884	764
Small Metro	1,590	1,458	1,099	1,033	609	563
250K – 1 Mil. Pop.	1,041	970	734	673	405	360
< 250K Pop.	549	488	366	360	204	204
Nonmetro	1,030	913	736 ^a	629	458	415
Urbanized	394	340	287	226	173	129
Less Urbanized	497	480	350	326	214	232
Completely Rural	140	93	99	77	71	55

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.48B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	19.8 ^a	18.5	13.9 ^a	12.7	7.8 ^a	7.0
GEOGRAPHIC DIVISION						
Northeast	18.2 ^a	15.8	13.5 ^a	11.6	8.2 ^b	6.0
New England	16.6	16.5	12.1	13.6	7.3	6.7
Middle Atlantic	18.8 ^a	15.6	14.1 ^b	10.9	8.5 ^b	5.7
Midwest	20.7	19.4	14.9	13.3	8.4	8.0
East North Central	20.5	19.4	14.9	13.2	8.4	7.6
West North Central	21.2	19.5	14.8	13.3	8.4	8.9
South	20.8	19.7	14.5	13.6	8.0	7.5
South Atlantic	21.1 ^a	17.7	14.9	12.9	7.8	6.9
East South Central	24.0	25.6	17.1	16.7	10.2	10.1
West South Central	18.9	19.8	12.7	13.3	7.3	7.2
West	18.4	17.5	12.4	11.5	6.8	6.1
Mountain	22.0	18.7	14.8	13.1	7.8	8.4
Pacific	16.7	17.0	11.2	10.8	6.3	5.0
COUNTY TYPE						
Large Metro	17.0	15.8	12.0 ^a	10.7	6.5 ^a	5.4
Small Metro	21.3	20.6	14.8	14.6	8.2	8.0
250K – 1 Mil. Pop.	20.3	19.9	14.3	13.8	7.9	7.4
< 250K Pop.	23.6	22.1	15.7	16.3	8.8	9.2
Nonmetro	26.5	24.2	18.9	16.7	11.8	11.0
Urbanized	23.7	22.1	17.3	14.7	10.4	8.4
Less Urbanized	27.9	25.7	19.6	17.4	12.0	12.4
Completely Rural	31.5	25.4	22.3	21.0	15.9	15.0

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.49A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	22,650	22,290	16,597	16,083	12,882 ^a	12,244
GEOGRAPHIC DIVISION						
Northeast	3,910	3,850	2,849	2,754	2,166	2,040
New England	1,099	1,093	790	795	590	619
Middle Atlantic	2,811	2,757	2,059	1,959	1,576	1,422
Midwest	5,125	5,196	3,870	3,850	3,086	3,002
East North Central	3,473	3,522	2,625	2,604	2,043	2,048
West North Central	1,652	1,673	1,244	1,246	1,043	954
South	8,284	8,164	6,129	6,030	4,817	4,761
South Atlantic	4,265	4,107	3,105	2,953	2,404	2,244
East South Central	1,349	1,385	1,051	1,091	872	908
West South Central	2,670	2,672	1,973	1,986	1,541	1,610
West	5,331	5,080	3,749 ^a	3,449	2,813 ^b	2,440
Mountain	1,723	1,650	1,242	1,134	975 ^a	833
Pacific	3,608	3,430	2,508	2,315	1,839 ^a	1,607
COUNTY TYPE						
Large Metro	11,651	11,609	8,390	8,193	6,382	6,119
Small Metro	7,296	7,085	5,314	5,187	4,114	3,974
250K – 1 Mil. Pop.	4,650	4,454	3,315	3,229	2,560	2,435
< 250K Pop.	2,645	2,631	1,999	1,958	1,553	1,538
Nonmetro	3,703	3,597	2,893	2,703	2,387	2,151
Urbanized	1,731	1,658	1,368	1,165	1,096	912
Less Urbanized	1,647	1,667	1,282	1,324	1,103	1,067
Completely Rural	325	272	243	213	189	172

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.49B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	65.1	63.8	47.7 ^a	46.0	37.0 ^b	35.0
GEOGRAPHIC DIVISION						
Northeast	63.6	62.6	46.3	44.8	35.2	33.2
New England	67.4	66.5	48.5	48.3	36.2	37.6
Middle Atlantic	62.2	61.2	45.6	43.5	34.9	31.5
Midwest	69.2	70.0	52.2	51.8	41.7	40.4
East North Central	68.0	68.9	51.4	51.0	40.0	40.1
West North Central	71.7	72.2	54.0	53.8	45.3	41.1
South	64.4	63.1	47.7	46.6	37.5	36.8
South Atlantic	65.0	62.3	47.3	44.8	36.6	34.0
East South Central	66.1	67.6	51.5	53.2	42.8	44.3
West South Central	62.8	62.1	46.4	46.2	36.2	37.4
West	63.7 ^a	60.4	44.8 ^a	41.0	33.6 ^b	29.0
Mountain	68.0	64.5	49.0 ^a	44.3	38.5 ^b	32.5
Pacific	61.8	58.6	42.9	39.5	31.5 ^a	27.4
COUNTY TYPE						
Large Metro	62.2	60.5	44.8	42.7	34.0 ^a	31.9
Small Metro	67.1	66.6	48.9	48.8	37.8	37.4
250K – 1 Mil. Pop.	66.5	64.8	47.4	47.0	36.6	35.4
< 250K Pop.	68.2	70.0	51.5	52.1	40.0	40.9
Nonmetro	71.6	70.2	56.0 ^a	52.7	46.2 ^a	42.0
Urbanized	70.0	69.3	55.3 ^b	48.7	44.3 ^a	38.1
Less Urbanized	74.1	71.2	57.7	56.6	49.6	45.6
Completely Rural	68.6	70.0	51.4	54.9	39.8	44.2

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.50A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	147,688	148,534	60,788	61,896	52,046	52,913
GEOGRAPHIC DIVISION						
Northeast	26,796	26,871	10,209	10,705	8,531	8,956
New England	7,160	7,323	2,505	2,552	1,997	2,079
Middle Atlantic	19,636	19,548	7,704	8,153	6,534	6,877
Midwest	33,765	33,834	14,979	14,564	12,943	12,624
East North Central	23,412	23,443	10,299	10,213	8,880	8,859
West North Central	10,352	10,391	4,680	4,351	4,063	3,765
South	54,896	55,104	23,785	24,538	20,703	21,407
South Atlantic	29,327	29,774	12,092	12,923	10,428	11,145
East South Central	8,848	8,883	4,295	4,406	3,938	3,934
West South Central	16,721	16,447	7,398	7,210	6,337	6,328
West	32,232	32,726	11,814	12,088	9,869	9,927
Mountain	10,455	10,453	4,025	4,027	3,416	3,414
Pacific	21,776	22,272	7,789	8,062	6,452	6,513
COUNTY TYPE						
Large Metro	76,266	79,618	29,823 ^a	32,041	24,870	26,547
Small Metro	45,678	43,942	19,011	18,328	16,506	15,921
250K – 1 Mil. Pop.	30,888	29,728	12,372	12,317	10,724	10,530
< 250K Pop.	14,790	14,214	6,639	6,011	5,782	5,392
Nonmetro	25,744	24,974	11,954	11,527	10,669	10,445
Urbanized	11,304	10,353	5,015	4,718	4,518	4,189
Less Urbanized	11,711	11,951	5,624	5,630	5,022	5,184
Completely Rural	2,730	2,670	1,315	1,180	1,129	1,072

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.50B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	72.9	72.3	30.0	30.1	25.7	25.8
GEOGRAPHIC DIVISION						
Northeast	72.3	72.0	27.6	28.7	23.0	24.0
New England	73.4	74.5	25.7	25.9	20.5	21.1
Middle Atlantic	71.9	71.1	28.2	29.7	23.9	25.0
Midwest	77.8	77.5	34.5	33.4	29.8	28.9
East North Central	77.7	77.4	34.2	33.7	29.5	29.3
West North Central	77.9	77.6	35.2	32.5	30.6	28.1
South	72.9	72.0	31.6	32.1	27.5	28.0
South Atlantic	72.9	72.8	30.1	31.6	25.9	27.3
East South Central	73.9	73.6	35.9	36.5	32.9	32.6
West South Central	72.4	69.9	32.0	30.6	27.5	26.9
West	68.6	68.4	25.2	25.3	21.0	20.8
Mountain	73.2	71.8	28.2	27.7	23.9	23.5
Pacific	66.7	66.9	23.8	24.2	19.8	19.6
COUNTY TYPE						
Large Metro	70.1	70.2	27.4	28.3	22.9	23.4
Small Metro	75.4 ^a	73.6	31.4	30.7	27.2	26.7
250K – 1 Mil. Pop.	74.8	73.2	30.0	30.3	26.0	25.9
< 250K Pop.	76.6	74.5	34.4	31.5	30.0	28.3
Nonmetro	77.1	77.4	35.8	35.7	31.9	32.4
Urbanized	76.0	77.8	33.7	35.4	30.4	31.5
Less Urbanized	78.2	77.2	37.5	36.4	33.5	33.5
Completely Rural	76.9	77.3	37.0	34.1	31.8	31.0

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.51A Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	170,337	170,825	77,385	77,979	64,928	65,157
GEOGRAPHIC DIVISION						
Northeast	30,706	30,721	13,058	13,459	10,698	10,996
New England	8,259	8,416	3,295	3,347	2,588	2,698
Middle Atlantic	22,447	22,305	9,763	10,112	8,110	8,298
Midwest	38,889	39,030	18,849	18,414	16,029	15,626
East North Central	26,885	26,965	12,925	12,817	10,922	10,908
West North Central	12,004	12,064	5,925	5,598	5,107	4,719
South	63,179	63,268	29,914	30,568	25,520	26,168
South Atlantic	33,591	33,881	15,197	15,876	12,832	13,389
East South Central	10,197	10,268	5,346	5,496	4,810	4,842
West South Central	19,391	19,119	9,371	9,196	7,878	7,938
West	37,563	37,806	15,564	15,537	12,682	12,366
Mountain	12,178	12,103	5,267	5,161	4,391	4,247
Pacific	25,385	25,703	10,297	10,377	8,291	8,120
COUNTY TYPE						
Large Metro	87,917	91,227	38,212	40,234	31,252	32,666
Small Metro	52,973	51,027	24,325	23,515	20,620	19,895
250K – 1 Mil. Pop.	35,538	34,181	15,687	15,546	13,285	12,965
< 250K Pop.	17,435	16,845	8,638	7,969	7,336	6,930
Nonmetro	29,447	28,572	14,847	14,230	13,057	12,596
Urbanized	13,035	12,011	6,382	5,883	5,614	5,101
Less Urbanized	13,358	13,618	6,907	6,954	6,125	6,251
Completely Rural	3,054	2,942	1,558	1,393	1,318	1,244

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.51B Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	71.7	71.1	32.6	32.5	27.3	27.1
GEOGRAPHIC DIVISION						
Northeast	71.1	70.7	30.2	31.0	24.8	25.3
New England	72.5	73.3	28.9	29.2	22.7	23.5
Middle Atlantic	70.6	69.7	30.7	31.6	25.5	25.9
Midwest	76.5	76.4	37.1	36.0	31.5	30.6
East North Central	76.3	76.2	36.7	36.2	31.0	30.8
West North Central	77.0	76.8	38.0	35.6	32.7	30.0
South	71.7	70.7	33.9	34.2	28.9	29.3
South Atlantic	71.8	71.4	32.5	33.4	27.4	28.2
East South Central	72.7	72.7	38.1	38.9	34.3	34.3
West South Central	70.9	68.7	34.3	33.0	28.8	28.5
West	67.9	67.2	28.1	27.6	22.9	22.0
Mountain	72.4	70.7	31.3	30.2	26.1	24.8
Pacific	65.9	65.7	26.7	26.5	21.5	20.7
COUNTY TYPE						
Large Metro	69.0	68.8	30.0	30.4	24.5	24.6
Small Metro	74.1 ^a	72.5	34.0	33.4	28.9	28.3
250K – 1 Mil. Pop.	73.6	72.0	32.5	32.7	27.5	27.3
< 250K Pop.	75.2	73.8	37.3	34.9	31.6	30.3
Nonmetro	76.4	76.4	38.5	38.1	33.9	33.7
Urbanized	75.2	76.5	36.8	37.5	32.4	32.5
Less Urbanized	77.6	76.4	40.1	39.0	35.6	35.1
Completely Rural	75.9	76.5	38.7	36.2	32.8	32.4

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.52A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	162,138	161,789	66,256	65,858	55,778	55,241
GEOGRAPHIC DIVISION						
Northeast	29,241	29,004	11,066	11,156	9,276	9,434
New England	7,865	7,856	2,759	2,670	2,266	2,231
Middle Atlantic	21,376	21,148	8,308	8,486	7,010	7,202
Midwest	37,217	37,029	16,243	15,565	13,845	13,270
East North Central	25,744	25,580	11,203	10,926	9,514	9,398
West North Central	11,473	11,449	5,040	4,639	4,331	3,871
South	60,097	59,917	25,633	25,967	21,848	22,159
South Atlantic	31,766	32,076	12,950	13,392	11,098	11,404
East South Central	9,892	9,800	4,640	4,463	4,084	3,897
West South Central	18,439	18,042	8,043	8,112	6,666	6,857
West	35,583	35,839	13,314	13,170	10,809	10,378
Mountain	11,559	11,512	4,425	4,461	3,631	3,667
Pacific	24,024	24,328	8,889	8,709	7,178	6,711
COUNTY TYPE						
Large Metro	83,080	85,837	32,601	33,539	26,853	27,660
Small Metro	50,444	48,565	20,710	20,239	17,641	17,124
250K – 1 Mil. Pop.	33,803	32,549	13,343	13,164	11,410	11,143
< 250K Pop.	16,642	16,016	7,368	7,075	6,231	5,981
Nonmetro	28,614	27,388	12,945	12,080	11,283	10,456
Urbanized	12,715	11,424	5,569	4,990	4,899	4,245
Less Urbanized	12,919	13,151	6,110	5,971	5,318	5,270
Completely Rural	2,980	2,812	1,265	1,119	1,067	941

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.52B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	61.8	61.0	25.3	24.8	21.3	20.8
GEOGRAPHIC DIVISION						
Northeast	61.7	60.9	23.4	23.4	19.6	19.8
New England	63.0	62.5	22.1	21.2	18.2	17.8
Middle Atlantic	61.2	60.3	23.8	24.2	20.1	20.5
Midwest	66.2	65.6	28.9	27.6	24.6	23.5
East North Central	66.0	65.4	28.7	27.9	24.4	24.0
West North Central	66.6	66.0	29.2	26.7	25.1	22.3
South	61.6	60.6	26.3	26.3	22.4	22.4
South Atlantic	61.7	61.5	25.2	25.7	21.6	21.9
East South Central	63.8	62.8	29.9	28.6	26.4	25.0
West South Central	60.3	58.0	26.3	26.1	21.8	22.1
West	58.1	57.6	21.7	21.2	17.6	16.7
Mountain	61.8	60.6	23.7	23.5	19.4	19.3
Pacific	56.4	56.3	20.9	20.2	16.9	15.5
COUNTY TYPE						
Large Metro	58.9	58.6	23.1	22.9	19.0	18.9
Small Metro	63.9	62.7	26.2	26.1	22.4	22.1
250K – 1 Mil. Pop.	63.3	62.1	25.0	25.1	21.4	21.3
< 250K Pop.	65.2	64.0	28.9	28.3	24.4	23.9
Nonmetro	67.4	66.5	30.5	29.4	26.6	25.4
Urbanized	66.9	66.3	29.3	28.9	25.8	24.6
Less Urbanized	68.0	66.8	32.2	30.3	28.0	26.8
Completely Rural	66.7	66.8	28.3	26.6	23.9	22.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.53A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	3,903 ^b	3,523	2,561 ^b	2,210	1,397 ^a	1,213
GEOGRAPHIC DIVISION						
Northeast	604 ^b	482	425 ^b	300	241 ^a	175
New England	129	115	95	78	54	45
Middle Atlantic	475 ^b	367	330 ^b	223	186 ^a	130
Midwest	911 ^a	785	597 ^b	488	338	309
East North Central	635	549	408 ^a	340	229	212
West North Central	276	237	189	148	109	98
South	1,471	1,430	969	888	519	473
South Atlantic	710	629	467	402	244	208
East South Central	270	292	176	159	100	94
West South Central	492	509	326	326	175	170
West	917	826	570	534	299	255
Mountain	346	291	221	199	112	113
Pacific	571	535	348	335	187	143
COUNTY TYPE						
Large Metro	1,820	1,659	1,210 ^b	992	622	543
Small Metro	1,285	1,145	814	764	449	374
250K – 1 Mil. Pop.	842	766	545	499	293	246
< 250K Pop.	443	379	269	265	156	127
Nonmetro	797	720	537	453	327	297
Urbanized	312	256	218 ^a	164	134	101
Less Urbanized	383	391	251	236	145	162
Completely Rural	103	73	67	53	48	34

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.53B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	15.7 ^b	14.2	10.3 ^b	8.9	5.6 ^a	4.9
GEOGRAPHIC DIVISION						
Northeast	14.4 ^b	11.6	10.1 ^b	7.2	5.7 ^a	4.2
New England	11.8	10.6	8.7	7.1	5.0	4.2
Middle Atlantic	15.3 ^b	11.9	10.7 ^b	7.2	6.0 ^a	4.2
Midwest	16.9 ^a	14.6	11.1 ^b	9.1	6.3	5.8
East North Central	16.9	14.7	10.9	9.1	6.1	5.7
West North Central	16.8	14.4	11.5	9.0	6.6	6.0
South	15.7	15.2	10.4	9.4	5.6	5.0
South Atlantic	15.2	13.4	10.0	8.6	5.2	4.5
East South Central	18.3	19.8	11.9	10.8	6.8	6.4
West South Central	15.2	15.6	10.1	10.0	5.4	5.2
West	15.4	13.9	9.6	9.0	5.0	4.3
Mountain	18.4	15.4	11.8	10.5	6.0	6.0
Pacific	14.0	13.2	8.5	8.3	4.6	3.5
COUNTY TYPE						
Large Metro	13.4 ^a	11.8	8.9 ^b	7.1	4.6	3.9
Small Metro	17.3	16.2	10.9	10.8	6.0	5.3
250K – 1 Mil. Pop.	16.4	15.7	10.6	10.3	5.7	5.1
< 250K Pop.	19.0	17.2	11.6	12.0	6.7	5.8
Nonmetro	20.5	19.1	13.8	12.0	8.4	7.9
Urbanized	18.8	16.7	13.1	10.6	8.1	6.6
Less Urbanized	21.5	20.9	14.1	12.6	8.1	8.6
Completely Rural	23.3	19.8	15.2	14.6	10.9	9.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.54A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	20,141 ^a	19,608	13,750 ^a	13,157	10,657 ^b	9,905
GEOGRAPHIC DIVISION						
Northeast	3,406	3,288	2,295	2,218	1,758	1,658
New England	934	867	633	616	468	481
Middle Atlantic	2,472	2,421	1,663	1,601	1,290	1,176
Midwest	4,535	4,564	3,209	3,119	2,542	2,389
East North Central	3,078	3,109	2,164	2,139	1,681	1,664
West North Central	1,456	1,455	1,045	980	861 ^a	725
South	7,421	7,227	5,113	4,949	3,979	3,869
South Atlantic	3,782 ^a	3,561	2,532 ^a	2,312	1,947	1,780
East South Central	1,222	1,201	907	927	742	740
West South Central	2,417	2,465	1,674	1,710	1,289	1,349
West	4,780	4,529	3,132	2,872	2,378 ^b	1,989
Mountain	1,557	1,453	1,019	925	806 ^a	681
Pacific	3,223	3,076	2,114	1,947	1,572 ^a	1,308
COUNTY TYPE						
Large Metro	10,385	10,218	6,976	6,756	5,297	4,979
Small Metro	6,447	6,163	4,331	4,155	3,364	3,170
250K – 1 Mil. Pop.	4,070	3,803	2,678	2,543	2,088	1,956
< 250K Pop.	2,377	2,360	1,654	1,611	1,276	1,214
Nonmetro	3,309	3,228	2,443	2,246	1,996	1,756
Urbanized	1,528	1,461	1,128	958	890	757
Less Urbanized	1,503	1,523	1,117	1,103	954	867
Completely Rural	278	244	198	186	153	132

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.54B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	57.9 ^a	56.1	39.5 ^a	37.7	30.6 ^b	28.4
GEOGRAPHIC DIVISION						
Northeast	55.4	53.5	37.3	36.1	28.6	27.0
New England	57.3	52.7	38.8	37.5	28.7	29.3
Middle Atlantic	54.7	53.7	36.8	35.5	28.5	26.1
Midwest	61.2	61.5	43.3	42.0	34.3	32.2
East North Central	60.3	60.9	42.4	41.9	32.9	32.6
West North Central	63.2	62.8	45.4	42.3	37.4 ^a	31.3
South	57.7	55.8	39.8	38.2	30.9	29.9
South Atlantic	57.6 ^a	54.0	38.6 ^a	35.1	29.7	27.0
East South Central	59.9	58.6	44.4	45.2	36.4	36.1
West South Central	56.8	57.3	39.4	39.8	30.3	31.4
West	57.1	53.8	37.4 ^a	34.1	28.4 ^b	23.6
Mountain	61.5 ^a	56.8	40.2	36.1	31.8 ^a	26.6
Pacific	55.2	52.5	36.2	33.3	26.9 ^a	22.3
COUNTY TYPE						
Large Metro	55.4	53.3	37.2	35.2	28.3 ^a	26.0
Small Metro	59.3	57.9	39.8	39.1	30.9	29.8
250K – 1 Mil. Pop.	58.2	55.3	38.3	37.0	29.9	28.4
< 250K Pop.	61.2	62.8	42.6	42.9	32.9	32.3
Nonmetro	64.0	63.0	47.3 ^a	43.8	38.6 ^b	34.3
Urbanized	61.8	61.0	45.6 ^a	40.0	36.0	31.6
Less Urbanized	67.6	65.0	50.3	47.1	42.9 ^b	37.0
Completely Rural	58.7	62.7	41.9	47.8	32.2	34.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.55A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	138,095	138,658	49,945	50,491	43,724	44,123
GEOGRAPHIC DIVISION						
Northeast	25,231	25,234	8,346	8,638	7,277	7,601
New England	6,802	6,874	2,031	1,976	1,744	1,705
Middle Atlantic	18,429	18,360	6,315	6,662	5,533	5,896
Midwest	31,771	31,679	12,437	11,958	10,965	10,572
East North Central	22,030	21,922	8,631	8,446	7,604	7,523
West North Central	9,741	9,757	3,806	3,511	3,361	3,049
South	51,206	51,260	19,551	20,130	17,350	17,817
South Atlantic	27,275	27,886	9,951	10,678	8,906	9,416
East South Central	8,400	8,307	3,557	3,377	3,242	3,063
West South Central	15,531	15,068	6,043	6,075	5,202	5,338
West	29,887	30,484	9,612	9,765	8,132	8,134
Mountain	9,656	9,767	3,185	3,338	2,713	2,874
Pacific	20,231	20,717	6,427	6,427	5,419	5,260
COUNTY TYPE						
Large Metro	70,874	73,960	24,414	25,790	20,935	22,138
Small Metro	42,713	41,258	15,565	15,320	13,828	13,581
250K – 1 Mil. Pop.	28,891	27,980	10,120	10,121	9,030	8,941
< 250K Pop.	13,822	13,278	5,445	5,199	4,799	4,640
Nonmetro	24,508	23,440	9,965	9,381	8,961	8,404
Urbanized	10,875	9,707	4,224	3,869	3,875	3,387
Less Urbanized	11,033	11,237	4,741	4,632	4,220	4,242
Completely Rural	2,599	2,496	1,000	880	866	775

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.55B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	68.1	67.5	24.6	24.6	21.6	21.5
GEOGRAPHIC DIVISION						
Northeast	68.1	67.6	22.5	23.1	19.6	20.4
New England	69.7	69.9	20.8	20.1	17.9	17.3
Middle Atlantic	67.5	66.8	23.1	24.2	20.3	21.4
Midwest	73.2	72.6	28.6	27.4	25.3	24.2
East North Central	73.1	72.4	28.7	27.9	25.2	24.9
West North Central	73.3	72.9	28.6	26.2	25.3	22.8
South	68.0	67.0	26.0	26.3	23.0	23.3
South Atlantic	67.8	68.2	24.7	26.1	22.1	23.0
East South Central	70.1	68.8	29.7	28.0	27.1	25.4
West South Central	67.3 ^a	64.0	26.2	25.8	22.5	22.7
West	63.7	63.7	20.5	20.4	17.3	17.0
Mountain	67.6	67.1	22.3	22.9	19.0	19.7
Pacific	61.9	62.3	19.7	19.3	16.6	15.8
COUNTY TYPE						
Large Metro	65.2	65.2	22.5	22.8	19.3	19.5
Small Metro	70.5	69.1	25.7	25.7	22.8	22.7
250K – 1 Mil. Pop.	70.0	68.9	24.5	24.9	21.9	22.0
< 250K Pop.	71.6	69.6	28.2	27.3	24.9	24.3
Nonmetro	73.4	72.7	29.8	29.1	26.8	26.1
Urbanized	73.1	72.9	28.4	29.1	26.1	25.4
Less Urbanized	73.6	72.6	31.6	29.9	28.2	27.4
Completely Rural	73.2	72.2	28.2	25.5	24.4	22.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.56A Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	158,236	158,266	63,695	63,648	54,381	54,028
GEOGRAPHIC DIVISION						
Northeast	28,637	28,522	10,641	10,856	9,035	9,258
New England	7,736	7,741	2,664	2,592	2,212	2,186
Middle Atlantic	20,901	20,781	7,978	8,263	6,823	7,072
Midwest	36,306	36,244	15,646	15,077	13,507	12,960
East North Central	25,109	25,032	10,795	10,586	9,285	9,187
West North Central	11,197	11,212	4,851	4,491	4,222	3,773
South	58,626	58,487	24,664	25,079	21,329	21,686
South Atlantic	31,057	31,447	12,483	12,990	10,853	11,196
East South Central	9,622	9,508	4,464	4,304	3,985	3,803
West South Central	17,947	17,533	7,717	7,785	6,491	6,687
West	34,667	35,014	12,744	12,637	10,510	10,123
Mountain	11,213	11,220	4,203	4,263	3,519	3,555
Pacific	23,454	23,793	8,540	8,374	6,991	6,568
COUNTY TYPE						
Large Metro	81,260	84,178	31,391	32,546	26,232	27,117
Small Metro	49,159	47,420	19,897	19,475	17,192	16,751
250K – 1 Mil. Pop.	32,961	31,783	12,798	12,664	11,118	10,896
< 250K Pop.	16,199	15,638	7,099	6,811	6,075	5,854
Nonmetro	27,816	26,668	12,408	11,627	10,957	10,160
Urbanized	12,403	11,168	5,351	4,826	4,765	4,144
Less Urbanized	12,536	12,760	5,858	5,735	5,174	5,109
Completely Rural	2,877	2,740	1,198	1,065	1,018	907

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.56B Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	66.6	65.9	26.8	26.5	22.9	22.5
GEOGRAPHIC DIVISION						
Northeast	66.3	65.6	24.6	25.0	20.9	21.3
New England	67.9	67.4	23.4	22.6	19.4	19.0
Middle Atlantic	65.7	64.9	25.1	25.8	21.5	22.1
Midwest	71.4	70.9	30.8	29.5	26.6	25.4
East North Central	71.3	70.7	30.6	29.9	26.4	26.0
West North Central	71.8	71.4	31.1	28.6	27.1	24.0
South	66.5	65.4	28.0	28.0	24.2	24.2
South Atlantic	66.4	66.2	26.7	27.4	23.2	23.6
East South Central	68.6	67.3	31.8	30.5	28.4	26.9
West South Central	65.6	63.0	28.2	28.0	23.7	24.0
West	62.7	62.2	23.0	22.5	19.0	18.0
Mountain	66.7	65.6	25.0	24.9	20.9	20.8
Pacific	60.9	60.8	22.2	21.4	18.2	16.8
COUNTY TYPE						
Large Metro	63.7	63.5	24.6	24.6	20.6	20.5
Small Metro	68.8	67.4	27.8	27.7	24.1	23.8
250K – 1 Mil. Pop.	68.3	66.9	26.5	26.7	23.0	22.9
< 250K Pop.	69.9	68.5	30.6	29.8	26.2	25.6
Nonmetro	72.1	71.3	32.2	31.1	28.4	27.2
Urbanized	71.5	71.1	30.9	30.7	27.5	26.4
Less Urbanized	72.9	71.6	34.1	32.2	30.1	28.7
Completely Rural	71.5	71.3	29.8	27.7	25.3	23.6

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.57A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Numbers in Thousands, 2013 and 2014**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	46,139	45,290	11,866	11,559	8,813	8,662
GEOGRAPHIC DIVISION						
Northeast	6,172	6,245	1,342	1,360	958	941
New England	1,631	1,699	291	342	165	207
Middle Atlantic	4,541	4,546	1,051	1,017	793	734
Midwest	11,430	11,388	2,963	3,027	2,166	2,331
East North Central	7,313	7,259	1,858	1,869	1,373	1,425
West North Central	4,117	4,129	1,105	1,157	793	906
South	18,030	18,022	5,164	5,071	4,021	4,009
South Atlantic	8,688	8,539	2,135	2,138	1,642	1,635
East South Central	3,203	3,484	1,139	1,321	940	1,121
West South Central	6,139	5,998	1,890	1,612	1,440	1,254
West	10,507	9,634	2,398	2,102	1,668	1,381
Mountain	3,879	3,606	995	860	725	552
Pacific	6,627	6,028	1,403	1,242	944	829
COUNTY TYPE						
Large Metro	19,187	20,264	4,382	4,495	3,025	3,175
Small Metro	16,109 ^a	14,579	3,870	3,758	2,926	2,792
250K – 1 Mil. Pop.	10,421 ^a	9,209	2,282	2,253	1,712	1,669
< 250K Pop.	5,688	5,370	1,588	1,505	1,215	1,123
Nonmetro	10,843	10,447	3,615	3,306	2,862	2,696
Urbanized	4,640	4,102	1,383	1,181	1,042	942
Less Urbanized	4,891	5,143	1,739	1,636	1,419	1,329
Completely Rural	1,311	1,201	492	489	401	425

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.57B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	17.6	17.1	4.5	4.4	3.4	3.3
GEOGRAPHIC DIVISION						
Northeast	13.0	13.1	2.8	2.9	2.0	2.0
New England	13.1	13.5	2.3	2.7	1.3	1.7
Middle Atlantic	13.0	13.0	3.0	2.9	2.3	2.1
Midwest	20.3	20.2	5.3	5.4	3.9	4.1
East North Central	18.8	18.6	4.8	4.8	3.5	3.6
West North Central	23.9	23.8	6.4	6.7	4.6	5.2
South	18.5	18.2	5.3	5.1	4.1	4.1
South Atlantic	16.9	16.4	4.1	4.1	3.2	3.1
East South Central	20.7	22.3	7.4	8.5	6.1	7.2
West South Central	20.1	19.3	6.2	5.2	4.7	4.0
West	17.1 ^a	15.5	3.9	3.4	2.7	2.2
Mountain	20.7	19.0	5.3	4.5	3.9	2.9
Pacific	15.6	14.0	3.3	2.9	2.2	1.9
COUNTY TYPE						
Large Metro	13.6	13.8	3.1	3.1	2.1	2.2
Small Metro	20.4 ^a	18.8	4.9	4.9	3.7	3.6
250K – 1 Mil. Pop.	19.5 ^a	17.6	4.3	4.3	3.2	3.2
< 250K Pop.	22.3	21.4	6.2	6.0	4.8	4.5
Nonmetro	25.5	25.4	8.5	8.0	6.7	6.6
Urbanized	24.4	23.8	7.3	6.8	5.5	5.5
Less Urbanized	25.8	26.1	9.2	8.3	7.5	6.7
Completely Rural	29.4	28.5	11.0	11.6	9.0	10.1

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.58A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	1,501	1,414	1,011	941	487	490
GEOGRAPHIC DIVISION						
Northeast	215	205	145	150	84	75
New England	46	56	39	48	15	19
Middle Atlantic	169	148	106	102	68	56
Midwest	357	339	244	206	115	113
East North Central	240	209	166 ^a	118	78	55
West North Central	117	131	78	88	37	58
South	723	641	483	446	230	235
South Atlantic	341	264	235	189	102	99
East South Central	164	173	102	115	62	68
West South Central	218	204	146	142	66	68
West	206	229	139	139	59	66
Mountain	104	105	66	67	23	35
Pacific	102	124	73	72	36	31
COUNTY TYPE						
Large Metro	493	510	340	358	155	155
Small Metro	527	489	353	313	153	164
250K – 1 Mil. Pop.	329	315	219	182	96	96
< 250K Pop.	198	174	134	130	58	68
Nonmetro	480	415	318	271	179	170
Urbanized	172	146	107	92	55	47
Less Urbanized	229	220	164	144	92	97
Completely Rural	79	49	47	35	33	27

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.58B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	6.0	5.7	4.1	3.8	2.0	2.0
GEOGRAPHIC DIVISION						
Northeast	5.1	4.9	3.5	3.6	2.0	1.8
New England	4.2	5.2	3.6	4.4	1.4	1.8
Middle Atlantic	5.5	4.8	3.4	3.3	2.2	1.8
Midwest	6.6	6.3	4.5	3.8	2.1	2.1
East North Central	6.4	5.6	4.4 ^a	3.2	2.1	1.5
West North Central	7.1	7.9	4.8	5.3	2.2	3.5
South	7.7	6.8	5.2	4.7	2.5	2.5
South Atlantic	7.3	5.6	5.0	4.0	2.2	2.1
East South Central	11.1	11.8	6.9	7.8	4.2	4.6
West South Central	6.7	6.3	4.5	4.4	2.0	2.1
West	3.5	3.9	2.3	2.3	1.0	1.1
Mountain	5.6	5.5	3.5	3.6	1.2	1.8
Pacific	2.5	3.1	1.8	1.8	0.9	0.8
COUNTY TYPE						
Large Metro	3.6	3.6	2.5	2.6	1.1	1.1
Small Metro	7.1	6.9	4.7	4.4	2.1	2.3
250K – 1 Mil. Pop.	6.4	6.5	4.3	3.7	1.9	2.0
< 250K Pop.	8.5	7.9	5.8	5.9	2.5	3.1
Nonmetro	12.4	11.0	8.2	7.2	4.6	4.5
Urbanized	10.4	9.5	6.4	6.0	3.3	3.0
Less Urbanized	12.8	11.7	9.2	7.7	5.1	5.2
Completely Rural	17.9	13.4	10.6	9.6	7.4	7.4

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.59A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	7,064	6,775	3,231	3,089	2,018	1,964
GEOGRAPHIC DIVISION						
Northeast	1,133 ^b	943	449	383	267	226
New England	313	279	130	121	64	58
Middle Atlantic	820 ^b	663	320	262	203	168
Midwest	1,855	1,869	863	878	576	576
East North Central	1,194	1,166	537	536	344	358
West North Central	660	703	325	342	232	218
South	2,576	2,546	1,252	1,209	829	819
South Atlantic	1,247	1,217	558	540	329	345
East South Central	519	535	282	287	210	207
West South Central	810	794	412	382	291	266
West	1,501	1,417	667	619	345	342
Mountain	549	562	246	249	145	145
Pacific	951	855	421	371	200	197
COUNTY TYPE						
Large Metro	3,016	2,872	1,211	1,173	672	708
Small Metro	2,417	2,385	1,174	1,146	769	739
250K – 1 Mil. Pop.	1,471	1,393	694	616	452	376
< 250K Pop.	946	992	480	531	317	363
Nonmetro	1,632	1,518	846	770	577	517
Urbanized	771	611	360	299	238	184
Less Urbanized	690	772	379	400	269	279
Completely Rural	170	135	106	72	70	54

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.59B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	20.3	19.4	9.3	8.8	5.8	5.6
GEOGRAPHIC DIVISION						
Northeast	18.4 ^b	15.3	7.3	6.2	4.3	3.7
New England	19.2	17.0	7.9	7.3	3.9	3.5
Middle Atlantic	18.2 ^b	14.7	7.1	5.8	4.5	3.7
Midwest	25.0	25.2	11.7	11.8	7.8	7.8
East North Central	23.4	22.8	10.5	10.5	6.7	7.0
West North Central	28.7	30.3	14.1	14.7	10.1	9.4
South	20.0	19.7	9.7	9.3	6.5	6.3
South Atlantic	19.0	18.5	8.5	8.2	5.0	5.2
East South Central	25.4	26.1	13.8	14.0	10.3	10.1
West South Central	19.0	18.5	9.7	8.9	6.8	6.2
West	17.9	16.8	8.0	7.4	4.1	4.1
Mountain	21.7	22.0	9.7	9.7	5.7	5.7
Pacific	16.3	14.6	7.2	6.3	3.4	3.4
COUNTY TYPE						
Large Metro	16.1	15.0	6.5	6.1	3.6	3.7
Small Metro	22.2	22.4	10.8	10.8	7.1	6.9
250K – 1 Mil. Pop.	21.0	20.3	9.9	9.0	6.5	5.5
< 250K Pop.	24.4	26.4	12.4	14.1	8.2	9.7
Nonmetro	31.6	29.6	16.4	15.0	11.2	10.1
Urbanized	31.2 ^a	25.5	14.6	12.5	9.6	7.7
Less Urbanized	31.0	33.0	17.1	17.1	12.1	11.9
Completely Rural	36.0	34.8	22.5	18.5	14.9	13.8

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.60A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Numbers in Thousands, 2013 and 2014**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	37,574	37,101	7,624	7,529	6,308	6,209
GEOGRAPHIC DIVISION						
Northeast	4,824	5,098	748	827	607	639
New England	1,272	1,364	122	174	86	130
Middle Atlantic	3,552	3,734	626	654	521	509
Midwest	9,218	9,180	1,856	1,942	1,475	1,642
East North Central	5,879	5,884	1,155	1,214	951	1,011
West North Central	3,340	3,296	701	728	524	630
South	14,731	14,835	3,429	3,417	2,962	2,956
South Atlantic	7,100	7,058	1,343	1,409	1,211	1,191
East South Central	2,520	2,775	755	920	668	845
West South Central	5,111	5,001	1,331	1,087	1,083	919
West	8,800	7,988	1,592	1,343	1,264	972
Mountain	3,226	2,939	683	543	556	371
Pacific	5,574	5,049	909	800	708	601
COUNTY TYPE						
Large Metro	15,678	16,882	2,830	2,965	2,199	2,311
Small Metro	13,164 ^a	11,705	2,343	2,299	2,004	1,889
250K – 1 Mil. Pop.	8,621 ^a	7,501	1,369	1,455	1,164	1,197
< 250K Pop.	4,543	4,204	974	844	840	691
Nonmetro	8,731	8,514	2,451	2,266	2,105	2,009
Urbanized	3,697	3,346	916	791	750	712
Less Urbanized	3,973	4,151	1,196	1,093	1,058	953
Completely Rural	1,062	1,017	339	382	298	344

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.60B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	18.5	18.1	3.8	3.7	3.1	3.0
GEOGRAPHIC DIVISION						
Northeast	13.0	13.7	2.0	2.2	1.6	1.7
New England	13.0	13.9	1.3	1.8	0.9	1.3
Middle Atlantic	13.0	13.6	2.3	2.4	1.9	1.9
Midwest	21.2	21.0	4.3	4.4	3.4	3.8
East North Central	19.5	19.4	3.8	4.0	3.2	3.3
West North Central	25.1	24.6	5.3	5.4	3.9	4.7
South	19.6	19.4	4.6	4.5	3.9	3.9
South Atlantic	17.6	17.3	3.3	3.4	3.0	2.9
East South Central	21.0	23.0	6.3	7.6	5.6	7.0
West South Central	22.1	21.3	5.8	4.6	4.7	3.9
West	18.7 ^a	16.7	3.4	2.8	2.7	2.0
Mountain	22.6	20.2	4.8	3.7	3.9 ^a	2.6
Pacific	17.1	15.2	2.8	2.4	2.2	1.8
COUNTY TYPE						
Large Metro	14.4	14.9	2.6	2.6	2.0	2.0
Small Metro	21.7 ^b	19.6	3.9	3.9	3.3	3.2
250K – 1 Mil. Pop.	20.9 ^a	18.5	3.3	3.6	2.8	2.9
< 250K Pop.	23.5	22.0	5.0	4.4	4.4	3.6
Nonmetro	26.1	26.4	7.3	7.0	6.3	6.2
Urbanized	24.9	25.1	6.2	5.9	5.0	5.3
Less Urbanized	26.5	26.8	8.0	7.1	7.1	6.2
Completely Rural	29.9	29.4	9.5	11.1	8.4	10.0

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.61A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Numbers in Thousands, 2013 and 2014**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	44,638	43,876	10,856	10,619	8,326	8,173
GEOGRAPHIC DIVISION						
Northeast	5,957	6,041	1,197	1,210	874	866
New England	1,585	1,643	252	294	150	188
Middle Atlantic	4,372	4,398	946	916	725	678
Midwest	11,073	11,049	2,719	2,821	2,051	2,218
East North Central	7,073	7,051	1,692	1,751	1,295	1,370
West North Central	4,000	3,998	1,027	1,070	756	848
South	17,307	17,380	4,681	4,626	3,792	3,774
South Atlantic	8,347	8,275	1,900	1,949	1,540	1,536
East South Central	3,039	3,311	1,037	1,207	878	1,052
West South Central	5,921	5,795	1,743	1,470	1,374	1,186
West	10,301	9,406	2,259	1,962	1,609	1,315
Mountain	3,775	3,501	929	792	701	517
Pacific	6,526	5,904	1,330	1,170	908	798
COUNTY TYPE						
Large Metro	18,694	19,754	4,042	4,137	2,870	3,019
Small Metro	15,581 ^a	14,090	3,517	3,445	2,773	2,628
250K – 1 Mil. Pop.	10,092 ^a	8,894	2,062	2,071	1,616	1,573
< 250K Pop.	5,490	5,196	1,454	1,374	1,157	1,055
Nonmetro	10,363	10,032	3,297	3,036	2,682	2,525
Urbanized	4,468	3,957	1,277	1,089	987	896
Less Urbanized	4,662	4,924	1,575	1,493	1,327	1,232
Completely Rural	1,232	1,152	445	454	368	398

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.61B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	18.8	18.3	4.6	4.4	3.5	3.4
GEOGRAPHIC DIVISION						
Northeast	13.8	13.9	2.8	2.8	2.0	2.0
New England	13.9	14.3	2.2	2.6	1.3	1.6
Middle Atlantic	13.7	13.7	3.0	2.9	2.3	2.1
Midwest	21.8	21.6	5.4	5.5	4.0	4.3
East North Central	20.1	19.9	4.8	4.9	3.7	3.9
West North Central	25.7	25.5	6.6	6.8	4.8	5.4
South	19.6	19.4	5.3	5.2	4.3	4.2
South Atlantic	17.8	17.4	4.1	4.1	3.3	3.2
East South Central	21.7	23.4	7.4	8.5	6.3	7.5
West South Central	21.7	20.8	6.4	5.3	5.0	4.3
West	18.6 ^a	16.7	4.1	3.5	2.9	2.3
Mountain	22.4	20.5	5.5	4.6	4.2	3.0
Pacific	16.9	15.1	3.5	3.0	2.4	2.0
COUNTY TYPE						
Large Metro	14.7	14.9	3.2	3.1	2.3	2.3
Small Metro	21.8 ^a	20.0	4.9	4.9	3.9	3.7
250K – 1 Mil. Pop.	20.9 ^a	18.7	4.3	4.4	3.3	3.3
< 250K Pop.	23.7	22.8	6.3	6.0	5.0	4.6
Nonmetro	26.9	26.8	8.5	8.1	7.0	6.8
Urbanized	25.8	25.2	7.4	6.9	5.7	5.7
Less Urbanized	27.1	27.6	9.2	8.4	7.7	6.9
Completely Rural	30.6	30.0	11.1	11.8	9.2	10.3

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.62A Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	89,117	89,267	24,933	23,734	12,438	11,964
GEOGRAPHIC DIVISION						
Northeast	15,370	15,838	4,362	4,450	2,061	2,181
New England	4,596	4,456	1,212	1,254	550	645
Middle Atlantic	10,774	11,382	3,150	3,196	1,511	1,536
Midwest	20,945	21,166	5,909	5,622	2,940	2,673
East North Central	14,188	14,447	3,950	3,846	1,896	1,880
West North Central	6,757	6,718	1,959	1,776	1,044	793
South	32,317	31,629	9,515 ^a	8,627	4,890	4,583
South Atlantic	17,317	17,104	5,172	4,828	2,476	2,571
East South Central	4,734	4,972	1,338	1,434	765	705
West South Central	10,266	9,553	3,005 ^b	2,365	1,649 ^a	1,307
West	20,486	20,634	5,147	5,036	2,546	2,527
Mountain	6,831	6,700	1,738	1,595	798	853
Pacific	13,655	13,935	3,409	3,441	1,749	1,674
COUNTY TYPE						
Large Metro	46,914	47,903	13,830	13,334	6,944	6,712
Small Metro	28,020	27,173	7,582	6,909	3,810	3,408
250K – 1 Mil. Pop.	19,276	18,500	5,138	4,702	2,516	2,260
< 250K Pop.	8,743	8,673	2,444	2,207	1,293	1,148
Nonmetro	14,183	14,191	3,521	3,491	1,684	1,843
Urbanized	6,536	5,947	1,570	1,601	707	817
Less Urbanized	6,194	6,833	1,505	1,633	762	910
Completely Rural	1,453	1,411	446 ^a	257	215	117

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.62B Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	34.0	33.7	9.5 ^a	9.0	4.7	4.5
GEOGRAPHIC DIVISION						
Northeast	32.4	33.3	9.2	9.3	4.3	4.6
New England	36.8	35.5	9.7	10.0	4.4	5.1
Middle Atlantic	30.9	32.5	9.0	9.1	4.3	4.4
Midwest	37.3	37.5	10.5	10.0	5.2	4.7
East North Central	36.4	36.9	10.1	9.8	4.9	4.8
West North Central	39.2	38.7	11.4	10.2	6.1	4.6
South	33.1	32.0	9.8 ^a	8.7	5.0	4.6
South Atlantic	33.7	32.8	10.1	9.3	4.8	4.9
East South Central	30.6	31.9	8.6	9.2	4.9	4.5
West South Central	33.6 ^a	30.7	9.8 ^b	7.6	5.4 ^a	4.2
West	33.4	33.2	8.4	8.1	4.2	4.1
Mountain	36.5	35.2	9.3	8.4	4.3	4.5
Pacific	32.1	32.3	8.0	8.0	4.1	3.9
COUNTY TYPE						
Large Metro	33.3	32.7	9.8 ^a	9.1	4.9	4.6
Small Metro	35.5	35.1	9.6	8.9	4.8	4.4
250K – 1 Mil. Pop.	36.1	35.3	9.6	9.0	4.7	4.3
< 250K Pop.	34.3	34.6	9.6	8.8	5.1	4.6
Nonmetro	33.4	34.5	8.3	8.5	4.0	4.5
Urbanized	34.4	34.5	8.3	9.3	3.7	4.7
Less Urbanized	32.6	34.7	7.9	8.3	4.0	4.6
Completely Rural	32.5	33.5	10.0 ^a	6.1	4.8	2.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.63A Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	2,070	1,985	1,424	1,426	576	521
GEOGRAPHIC DIVISION						
Northeast	378	348	253	271	114	99
New England	96	106	64	91	27	30
Middle Atlantic	282	242	188	180	87	69
Midwest	491	457	365	333	158	135
East North Central	349	324	262	226	113	89
West North Central	142	133	102	106	45	47
South	769	775	521	555	206	214
South Atlantic	380	369	257	283	95	108
East South Central	137	145	96	96	44	35
West South Central	253	260	168	176	67	71
West	432	405	285	267	98	73
Mountain	188	151	122	106	43	37
Pacific	244	255	163	161	56	36
COUNTY TYPE						
Large Metro	1,006	1,013	709	738	280	256
Small Metro	663	611	435	431	174	164
250K – 1 Mil. Pop.	436	404	300	286	125	102
< 250K Pop.	228	207	135	145	49	62
Nonmetro	401	361	280	257	122	101
Urbanized	161	141	113	100	46	41
Less Urbanized	183	175	131	129	62	50
Completely Rural	57	45	36	28	14	10

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.63B Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	8.3	8.0	5.7	5.7	2.3	2.1
GEOGRAPHIC DIVISION						
Northeast	9.0	8.4	6.0	6.5	2.7	2.4
New England	8.8	9.8	5.9	8.4	2.5	2.8
Middle Atlantic	9.1	7.9	6.1	5.9	2.8	2.2
Midwest	9.1	8.5	6.8	6.2	2.9	2.5
East North Central	9.3	8.7	7.0	6.1	3.0	2.4
West North Central	8.6	8.1	6.2	6.5	2.7	2.8
South	8.2	8.2	5.6	5.9	2.2	2.3
South Atlantic	8.2	7.9	5.5	6.0	2.0	2.3
East South Central	9.3	9.8	6.5	6.5	3.0	2.3
West South Central	7.8	8.0	5.2	5.4	2.1	2.2
West	7.3	6.8	4.8	4.5	1.7	1.2
Mountain	10.0	8.0	6.5	5.6	2.3	1.9
Pacific	6.0	6.3	4.0	4.0	1.4	0.9
COUNTY TYPE						
Large Metro	7.4	7.2	5.2	5.3	2.1	1.8
Small Metro	8.9	8.6	5.8	6.1	2.3	2.3
250K – 1 Mil. Pop.	8.5	8.3	5.9	5.9	2.4	2.1
< 250K Pop.	9.8	9.4	5.8	6.6	2.1	2.8
Nonmetro	10.3	9.6	7.2	6.8	3.1	2.7
Urbanized	9.7	9.2	6.8	6.5	2.8	2.6
Less Urbanized	10.2	9.4	7.4	6.9	3.5	2.7
Completely Rural	12.8	12.3	8.0	7.6	3.1	2.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.64A Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	13,379	13,170	7,214	6,975	3,472	3,392
GEOGRAPHIC DIVISION						
Northeast	2,397	2,359	1,280	1,235	607	619
New England	716	687	400	379	186	217
Middle Atlantic	1,681	1,672	880	856	421	401
Midwest	3,205	3,276	1,748	1,779	839	856
East North Central	2,158	2,193	1,181	1,150	565	577
West North Central	1,047	1,083	567	630	274	279
South	4,641	4,530	2,568	2,510	1,321	1,261
South Atlantic	2,407	2,344	1,326	1,294	703	616
East South Central	819	762	421	436	222	206
West South Central	1,416	1,424	821	780	396	439
West	3,137	3,005	1,617	1,451	704	656
Mountain	1,031	1,007	511	469	200	227
Pacific	2,106	1,998	1,106	981	504	429
COUNTY TYPE						
Large Metro	6,782	6,833	3,678	3,637	1,799	1,803
Small Metro	4,491	4,259	2,379	2,254	1,110	1,083
250K – 1 Mil. Pop.	2,852	2,669	1,504	1,422	650	683
< 250K Pop.	1,639	1,590	875	831	460	400
Nonmetro	2,106	2,077	1,157	1,085	563	505
Urbanized	1,075	991	612	531	285	247
Less Urbanized	860	925	465	496	241	229
Completely Rural	171	162	79	58	37	29

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.64B Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	38.5	37.7	20.7	20.0	10.0	9.7
GEOGRAPHIC DIVISION						
Northeast	39.0	38.4	20.8	20.1	9.9	10.1
New England	43.9	41.8	24.5	23.1	11.4	13.2
Middle Atlantic	37.2	37.1	19.5	19.0	9.3	8.9
Midwest	43.3	44.1	23.6	24.0	11.3	11.5
East North Central	42.3	42.9	23.1	22.5	11.1	11.3
West North Central	45.5	46.7	24.6	27.2	11.9	12.0
South	36.1	35.0	20.0	19.4	10.3	9.7
South Atlantic	36.7	35.6	20.2	19.6	10.7	9.3
East South Central	40.1	37.2	20.6	21.3	10.9	10.0
West South Central	33.3	33.1	19.3	18.1	9.3	10.2
West	37.5	35.7	19.3	17.2	8.4	7.8
Mountain	40.7	39.3	20.2	18.4	7.9	8.9
Pacific	36.1	34.1	18.9	16.8	8.6	7.3
COUNTY TYPE						
Large Metro	36.2	35.6	19.6	19.0	9.6	9.4
Small Metro	41.3	40.0	21.9	21.2	10.2	10.2
250K – 1 Mil. Pop.	40.8	38.8	21.5	20.7	9.3	9.9
< 250K Pop.	42.2	42.3	22.6	22.1	11.8	10.6
Nonmetro	40.7	40.5	22.4	21.2	10.9	9.9
Urbanized	43.5	41.4	24.8	22.2	11.5	10.3
Less Urbanized	38.7	39.5	20.9	21.2	10.9	9.8
Completely Rural	36.1	41.6	16.7	14.9	7.7	7.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.65A Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	73,668	74,112	16,296	15,333	8,389	8,051
GEOGRAPHIC DIVISION						
Northeast	12,595	13,131	2,830	2,944	1,340	1,464
New England	3,784	3,663	748	783	337	398
Middle Atlantic	8,811	9,467	2,082	2,161	1,002	1,066
Midwest	17,249	17,433	3,796	3,510	1,943	1,682
East North Central	11,681	11,930	2,507	2,470	1,218	1,215
West North Central	5,568	5,502	1,289	1,040	725	467
South	26,907	26,325	6,426 ^a	5,562	3,363	3,108
South Atlantic	14,531	14,391	3,589	3,251	1,678	1,846
East South Central	3,779	4,065	821	902	499	465
West South Central	8,598	7,869	2,016 ^b	1,409	1,187 ^a	796
West	16,917	17,224	3,245	3,318	1,744	1,798
Mountain	5,612	5,542	1,105	1,020	554	589
Pacific	11,305	11,682	2,140	2,298	1,189	1,209
COUNTY TYPE						
Large Metro	39,126	40,057	9,444	8,960	4,864	4,654
Small Metro	22,865	22,303	4,767	4,224	2,526	2,160
250K – 1 Mil. Pop.	15,989	15,427	3,334	2,993	1,741	1,475
< 250K Pop.	6,877	6,876	1,433	1,231	785	686
Nonmetro	11,676	11,752	2,085	2,149	999	1,237
Urbanized	5,300	4,816	845	970	376	529
Less Urbanized	5,151	5,732	908	1,008	458	630
Completely Rural	1,226	1,204	332	171	165	78

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.65B Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	36.3	36.1	8.0 ^a	7.5	4.1	3.9
GEOGRAPHIC DIVISION						
Northeast	34.0	35.2	7.6	7.9	3.6	3.9
New England	38.8	37.2	7.7	8.0	3.5	4.0
Middle Atlantic	32.3	34.4	7.6	7.9	3.7	3.9
Midwest	39.7	39.9	8.7	8.0	4.5	3.9
East North Central	38.8	39.4	8.3	8.2	4.0	4.0
West North Central	41.9	41.1	9.7	7.8	5.5	3.5
South	35.7	34.4	8.5 ^a	7.3	4.5	4.1
South Atlantic	36.1	35.2	8.9	7.9	4.2	4.5
East South Central	31.5	33.7	6.9	7.5	4.2	3.9
West South Central	37.2 ^a	33.4	8.7 ^b	6.0	5.1 ^b	3.4
West	36.0	36.0	6.9	6.9	3.7	3.8
Mountain	39.3	38.1	7.7	7.0	3.9	4.0
Pacific	34.6	35.1	6.6	6.9	3.6	3.6
COUNTY TYPE						
Large Metro	36.0	35.3	8.7	7.9	4.5	4.1
Small Metro	37.7	37.4	7.9	7.1	4.2	3.6
250K – 1 Mil. Pop.	38.7	38.0	8.1	7.4	4.2	3.6
< 250K Pop.	35.6	36.0	7.4	6.5	4.1	3.6
Nonmetro	35.0	36.4	6.2	6.7	3.0	3.8
Urbanized	35.6	36.2	5.7	7.3	2.5 ^a	4.0
Less Urbanized	34.4	37.0	6.1	6.5	3.1	4.1
Completely Rural	34.5	34.8	9.4 ^a	5.0	4.6	2.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.66A Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	87,047	87,282	23,509	22,309	11,862	11,443
GEOGRAPHIC DIVISION						
Northeast	14,992	15,490	4,110	4,179	1,947	2,082
New England	4,500	4,350	1,148	1,163	523	615
Middle Atlantic	10,492	11,140	2,962	3,016	1,424	1,467
Midwest	20,454	20,709	5,544	5,289	2,782	2,538
East North Central	13,839	14,123	3,688	3,620	1,783	1,792
West North Central	6,615	6,586	1,856	1,670	999	746
South	31,548	30,855	8,994 ^a	8,072	4,684	4,369
South Atlantic	16,937	16,735	4,915	4,545	2,381	2,462
East South Central	4,597	4,827	1,241	1,338	721	671
West South Central	10,013	9,293	2,837 ^b	2,189	1,583 ^a	1,236
West	20,053	20,229	4,862	4,769	2,448	2,454
Mountain	6,643	6,549	1,616	1,489	755	816
Pacific	13,411	13,680	3,246	3,280	1,693	1,638
COUNTY TYPE						
Large Metro	45,909	46,891	13,121	12,596	6,664	6,456
Small Metro	27,356	26,562	7,146	6,478	3,636	3,244
250K – 1 Mil. Pop.	18,841	18,096	4,838	4,415	2,391	2,158
< 250K Pop.	8,516	8,466	2,309	2,062	1,245	1,086
Nonmetro	13,782	13,830	3,242	3,234	1,562	1,743
Urbanized	6,375	5,807	1,458	1,501	661	776
Less Urbanized	6,011	6,657	1,373	1,504	699	860
Completely Rural	1,396	1,366	411 ^a	229	201	107

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.66B Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	36.7	36.3	9.9 ^a	9.3	5.0	4.8
GEOGRAPHIC DIVISION						
Northeast	34.7	35.6	9.5	9.6	4.5	4.8
New England	39.5	37.9	10.1	10.1	4.6	5.4
Middle Atlantic	33.0	34.8	9.3	9.4	4.5	4.6
Midwest	40.3	40.5	10.9	10.4	5.5	5.0
East North Central	39.3	39.9	10.5	10.2	5.1	5.1
West North Central	42.4	41.9	11.9	10.6	6.4	4.7
South	35.8	34.5	10.2 ^a	9.0	5.3	4.9
South Atlantic	36.2	35.2	10.5	9.6	5.1	5.2
East South Central	32.8	34.2	8.9	9.5	5.1	4.8
West South Central	36.6 ^a	33.4	10.4 ^b	7.9	5.8 ^a	4.4
West	36.2	36.0	8.8	8.5	4.4	4.4
Mountain	39.5	38.3	9.6	8.7	4.5	4.8
Pacific	34.8	35.0	8.4	8.4	4.4	4.2
COUNTY TYPE						
Large Metro	36.0	35.4	10.3 ^a	9.5	5.2	4.9
Small Metro	38.3	37.8	10.0	9.2	5.1	4.6
250K – 1 Mil. Pop.	39.0	38.1	10.0	9.3	5.0	4.5
< 250K Pop.	36.7	37.1	10.0	9.0	5.4	4.8
Nonmetro	35.7	37.0	8.4	8.7	4.1	4.7
Urbanized	36.8	37.0	8.4	9.6	3.8	4.9
Less Urbanized	34.9	37.3	8.0	8.4	4.1	4.8
Completely Rural	34.7	35.5	10.2 ^a	6.0	5.0	2.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.67A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	213,794 ^b	217,765	173,966 ^a	176,556	136,868 ^a	139,677
GEOGRAPHIC DIVISION						
Northeast	39,777	40,398	33,909	34,543	27,501	27,788
New England	10,932	10,951	9,304	9,385	7,724	7,608
Middle Atlantic	28,846	29,448	24,605	25,159	19,777	20,180
Midwest	47,725	47,856	39,158	38,946	31,317	31,121
East North Central	33,065	33,139	27,266	26,978	21,661	21,565
West North Central	14,660	14,717	11,892	11,968	9,657	9,556
South	77,508 ^b	79,390	61,443	62,164	46,979	48,184
South Atlantic	41,599 ^a	42,572	33,444	34,139	26,098	26,824
East South Central	11,817	12,076	8,775	8,964	6,536	6,742
West South Central	24,093	24,742	19,223	19,061	14,345	14,617
West	48,784 ^a	50,121	39,456 ^a	40,903	31,071 ^a	32,584
Mountain	15,221	15,274	12,204	12,268	9,717	9,661
Pacific	33,563 ^a	34,847	27,252 ^a	28,635	21,354 ^a	22,922
COUNTY TYPE						
Large Metro	114,457 ^a	120,346	95,831 ^a	100,940	76,521 ^a	80,932
Small Metro	65,114	63,994	51,938	50,783	40,688	39,824
250K – 1 Mil. Pop.	44,130	43,186	35,691	34,385	28,317	27,297
< 250K Pop.	20,985	20,808	16,246	16,398	12,371	12,527
Nonmetro	34,223	33,425	26,197	24,833	19,659	18,921
Urbanized	15,542	14,355	12,384	11,073	9,463	8,652
Less Urbanized	15,224	15,778	11,367	11,615	8,376	8,745
Completely Rural	3,458	3,293	2,446	2,145	1,820	1,524

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.67B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	81.5	82.1	66.3	66.6	52.2	52.7
GEOGRAPHIC DIVISION						
Northeast	83.9	84.8	71.6	72.5	58.0	58.3
New England	87.6	87.2	74.5	74.7	61.9	60.5
Middle Atlantic	82.6	84.0	70.5	71.7	56.7	57.5
Midwest	84.9	84.8	69.7	69.0	55.7	55.1
East North Central	84.8	84.7	69.9	69.0	55.6	55.1
West North Central	85.1	84.8	69.0	69.0	56.0	55.1
South	79.5	80.3	63.0	62.9	48.2	48.7
South Atlantic	80.8	81.6	65.0	65.5	50.7	51.4
East South Central	76.3	77.4	56.6	57.5	42.2	43.2
West South Central	78.8	79.6	62.9	61.3	46.9	47.0
West	79.6	80.6	64.4	65.8	50.7	52.4
Mountain	81.4	80.4	65.3	64.5	52.0	50.8
Pacific	78.8	80.7	64.0	66.3	50.2	53.1
COUNTY TYPE						
Large Metro	81.2	82.1	67.9	68.9	54.3	55.2
Small Metro	82.5	82.7	65.8	65.6	51.6	51.4
250K – 1 Mil. Pop.	82.7	82.5	66.9	65.7	53.0	52.1
< 250K Pop.	82.3	83.1	63.7	65.5	48.5	50.0
Nonmetro	80.6	81.2	61.7	60.3	46.3	46.0
Urbanized	81.8	83.2	65.2	64.2	49.8	50.2
Less Urbanized	80.2	80.1	59.9	59.0	44.1	44.4
Completely Rural	77.4	78.2	54.8	50.9	40.7	36.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.68A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	7,669	7,375	6,133	5,979	2,877	2,872
GEOGRAPHIC DIVISION						
Northeast	1,333	1,352	1,122	1,160	564	564
New England	350	361	287	309	149	130
Middle Atlantic	982	991	835	851	415	434
Midwest	1,704 ^a	1,558	1,374 ^a	1,225	619	556
East North Central	1,225	1,118	983 ^a	877	458 ^a	385
West North Central	479	440	390	348	161	171
South	2,878 ^a	2,675	2,240	2,166	1,027	1,019
South Atlantic	1,476 ^b	1,292	1,133	1,072	540	508
East South Central	412	385	326	299	140	130
West South Central	990	997	781	795	346	381
West	1,754	1,791	1,397	1,428	667	733
Mountain	593	529	465	431	216	240
Pacific	1,162	1,262	931	997	451	493
COUNTY TYPE						
Large Metro	4,092	4,107	3,337	3,364	1,589	1,668
Small Metro	2,339 ^a	2,103	1,845	1,695	850	780
250K – 1 Mil. Pop.	1,555	1,437	1,227	1,163	582	550
< 250K Pop.	784	665	618	533	268	229
Nonmetro	1,238	1,165	950	919	437	425
Urbanized	523	456	400	365	189	157
Less Urbanized	576	599	443	470	199	221
Completely Rural	140	110	107	85	50	47

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.68B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	30.8	29.6	24.6	24.0	11.6	11.5
GEOGRAPHIC DIVISION						
Northeast	31.8	32.5	26.8	27.9	13.5	13.6
New England	32.1	33.3	26.3	28.5	13.6	12.0
Middle Atlantic	31.7	32.3	27.0	27.7	13.4	14.1
Midwest	31.6 ^a	29.0	25.4 ^a	22.8	11.5	10.4
East North Central	32.6	30.0	26.2	23.5	12.2	10.3
West North Central	29.2	26.8	23.8	21.2	9.8	10.4
South	30.8 ^a	28.4	23.9	23.0	11.0	10.8
South Atlantic	31.7 ^b	27.6	24.3	22.9	11.6	10.8
East South Central	27.9	26.1	22.1	20.3	9.5	8.9
West South Central	30.7	30.6	24.2	24.4	10.7	11.7
West	29.5	30.2	23.5	24.1	11.2	12.4
Mountain	31.6	27.9	24.8	22.8	11.5	12.7
Pacific	28.5	31.2	22.9	24.7	11.1	12.2
COUNTY TYPE						
Large Metro	30.2	29.3	24.6	24.0	11.7	11.9
Small Metro	31.4	29.7	24.8	24.0	11.4	11.0
250K – 1 Mil. Pop.	30.4	29.5	24.0	23.9	11.4	11.3
< 250K Pop.	33.7	30.2	26.6	24.2	11.5	10.4
Nonmetro	31.9	30.9	24.4	24.3	11.3	11.2
Urbanized	31.5	29.6	24.1	23.7	11.4	10.2
Less Urbanized	32.3	32.0	24.8	25.1	11.2	11.8
Completely Rural	31.5	30.1	24.1	23.0	11.3	12.9

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.69A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	29,148	29,126	26,707	26,713	20,738	20,815
GEOGRAPHIC DIVISION						
Northeast	5,296	5,299	4,924	4,921	3,933	3,962
New England	1,461	1,457	1,374	1,397	1,127	1,154
Middle Atlantic	3,836	3,842	3,551	3,523	2,806	2,808
Midwest	6,423	6,365	6,000	5,892	4,773	4,662
East North Central	4,437	4,370	4,159	4,038	3,309	3,166
West North Central	1,986	1,995	1,841	1,854	1,464	1,496
South	10,590	10,658	9,640	9,715	7,277	7,291
South Atlantic	5,436	5,495	4,987	5,036	3,826	3,793
East South Central	1,648	1,643	1,475	1,495	1,060	1,097
West South Central	3,507	3,520	3,178	3,184	2,390	2,401
West	6,838	6,804	6,144	6,185	4,756	4,900
Mountain	2,086	2,059	1,857	1,877	1,404	1,435
Pacific	4,753	4,745	4,287	4,309	3,352	3,465
COUNTY TYPE						
Large Metro	15,676	15,945	14,404	14,675	11,242	11,503
Small Metro	9,141	8,850	8,388	8,187	6,564	6,389
250K – 1 Mil. Pop.	5,848	5,640	5,355	5,236	4,237	4,163
< 250K Pop.	3,293	3,210	3,033	2,951	2,327	2,226
Nonmetro	4,332	4,331	3,915	3,851	2,932	2,923
Urbanized	2,097	2,071	1,924	1,836	1,442	1,429
Less Urbanized	1,878	1,941	1,673	1,752	1,260	1,299
Completely Rural	357	319	318	263	230	196

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.69B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	83.8	83.4	76.8	76.5	59.6	59.6
GEOGRAPHIC DIVISION						
Northeast	86.1	86.2	80.1	80.0	64.0	64.4
New England	89.6	88.6	84.3	85.0	69.1	70.2
Middle Atlantic	84.9	85.3	78.6	78.2	62.1	62.3
Midwest	86.7	85.7	81.0	79.3	64.4	62.8
East North Central	86.9	85.5	81.5	79.0	64.8	62.0
West North Central	86.3	86.0	80.0	80.0	63.6	64.5
South	82.4	82.3	75.0	75.1	56.6	56.3
South Atlantic	82.8	83.4	76.0	76.4	58.3	57.5
East South Central	80.7	80.1	72.3	72.9	51.9	53.5
West South Central	82.4	81.8	74.7	74.0	56.2	55.8
West	81.7	80.9	73.4	73.5	56.8	58.2
Mountain	82.4	80.5	73.3	73.4	55.4	56.1
Pacific	81.4	81.0	73.4	73.6	57.4	59.2
COUNTY TYPE						
Large Metro	83.6	83.2	76.8	76.5	60.0	60.0
Small Metro	84.1	83.2	77.2	77.0	60.4	60.1
250K – 1 Mil. Pop.	83.7	82.0	76.6	76.2	60.6	60.5
< 250K Pop.	84.8	85.4	78.1	78.5	59.9	59.2
Nonmetro	83.8	84.5	75.7	75.2	56.7	57.0
Urbanized	84.8	86.5	77.8	76.7	58.3	59.7
Less Urbanized	84.5	82.9	75.3	74.8	56.7	55.4
Completely Rural	75.4	82.0	67.1	67.6	48.5	50.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.70A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	176,977 ^b	181,265	141,126 ^a	143,864	113,253 ^a	115,990
GEOGRAPHIC DIVISION						
Northeast	33,148	33,747	27,862	28,463	23,004	23,262
New England	9,121	9,133	7,643	7,678	6,448	6,324
Middle Atlantic	24,027	24,614	20,219	20,784	16,556	16,938
Midwest	39,598	39,934	31,785	31,828	25,925	25,903
East North Central	27,404	27,651	22,124	22,062	17,894	18,014
West North Central	12,194	12,283	9,660	9,766	8,031	7,889
South	64,040 ^b	66,057	49,563	50,283	38,675	39,874
South Atlantic	34,687 ^a	35,784	27,324	28,032	21,732	22,524
East South Central	9,757	10,048	6,975	7,170	5,336	5,515
West South Central	19,596 ^a	20,224	15,265	15,082	11,608	11,835
West	40,191 ^b	41,527	31,915 ^a	33,289	25,648	26,951
Mountain	12,543	12,687	9,882	9,961	8,097	7,986
Pacific	27,649 ^b	28,840	22,034 ^a	23,329	17,551 ^a	18,965
COUNTY TYPE						
Large Metro	94,689 ^a	100,294	78,089 ^a	82,901	63,690 ^a	67,761
Small Metro	53,635	53,042	41,704	40,900	33,274	32,656
250K – 1 Mil. Pop.	36,726	36,109	29,109	27,986	23,498	22,584
< 250K Pop.	16,908	16,933	12,595	12,914	9,776	10,072
Nonmetro	28,653	27,928	21,332	20,062	16,289	15,573
Urbanized	12,921	11,828	10,060	8,873	7,832	7,067
Less Urbanized	12,770	13,237	9,251	9,392	6,917	7,225
Completely Rural	2,961	2,864	2,021	1,798	1,540	1,280

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.70B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	87.3 ^a	88.3	69.6	70.1	55.9	56.5
GEOGRAPHIC DIVISION						
Northeast	89.5	90.4	75.2	76.3	62.1	62.3
New England	93.4	92.9	78.3	78.1	66.1	64.3
Middle Atlantic	88.0	89.5	74.1	75.6	60.7	61.6
Midwest	91.2	91.5	73.2	72.9	59.7	59.3
East North Central	91.0	91.3	73.5	72.9	59.4	59.5
West North Central	91.8	91.7	72.7	72.9	60.4	58.9
South	85.0	86.4	65.8	65.7	51.4	52.1
South Atlantic	86.2	87.5	67.9	68.6	54.0	55.1
East South Central	81.4	83.3	58.2	59.4	44.5	45.7
West South Central	84.9	85.9	66.1	64.1	50.3	50.3
West	85.6	86.8	68.0	69.6	54.6	56.3
Mountain	87.8	87.2	69.2	68.4	56.7	54.9
Pacific	84.6	86.7	67.5	70.1	53.7	57.0
COUNTY TYPE						
Large Metro	87.1 ^a	88.5	71.8	73.1	58.6	59.8
Small Metro	88.5	88.8	68.8	68.5	54.9	54.7
250K – 1 Mil. Pop.	89.0	88.9	70.5	68.9	56.9	55.6
< 250K Pop.	87.6	88.8	65.3	67.7	50.6	52.8
Nonmetro	85.8	86.6	63.9	62.2	48.8	48.3
Urbanized	86.9	88.8	67.7	66.6	52.7	53.1
Less Urbanized	85.2	85.5	61.7	60.7	46.2	46.7
Completely Rural	83.4	82.9	57.0	52.0	43.4	37.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.71A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	206,125 ^b	210,390	167,833 ^a	170,577	133,991 ^a	136,804
GEOGRAPHIC DIVISION						
Northeast	38,444	39,046	32,787	33,383	26,937	27,224
New England	10,581	10,590	9,017	9,076	7,575	7,478
Middle Atlantic	27,863	28,457	23,770	24,308	19,362	19,746
Midwest	46,021	46,298	37,784	37,720	30,698	30,565
East North Central	31,841	32,021	26,283	26,101	21,203	21,180
West North Central	14,180	14,278	11,501	11,620	9,495	9,385
South	74,630 ^b	76,715	59,203	59,998	45,952	47,165
South Atlantic	40,123 ^a	41,280	32,311	33,068	25,558	26,317
East South Central	11,405	11,691	8,450	8,665	6,396	6,612
West South Central	23,102 ^a	23,744	18,442	18,266	13,998	14,236
West	47,029 ^a	48,330	38,059 ^a	39,475	30,404 ^a	31,850
Mountain	14,628	14,745	11,739	11,837	9,501	9,421
Pacific	32,401 ^a	33,585	26,320 ^a	27,637	20,903 ^a	22,429
COUNTY TYPE						
Large Metro	110,365 ^a	116,239	92,494 ^a	97,576	74,932 ^a	79,264
Small Metro	62,775	61,891	50,092	49,087	39,838	39,044
250K – 1 Mil. Pop.	42,574	41,749	34,464	33,223	27,735	26,747
< 250K Pop.	20,201	20,143	15,628	15,865	12,103	12,297
Nonmetro	32,985	32,260	25,247	23,913	19,221	18,496
Urbanized	15,019	13,899	11,984	10,709	9,274	8,496
Less Urbanized	14,648	15,179	10,924	11,144	8,177	8,524
Completely Rural	3,318	3,182	2,339	2,060	1,770	1,476

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.71B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	86.8	87.6	70.7	71.0	56.4	56.9
GEOGRAPHIC DIVISION						
Northeast	89.0	89.8	75.9	76.8	62.4	62.6
New England	92.9	92.2	79.2	79.1	66.5	65.1
Middle Atlantic	87.6	88.9	74.7	76.0	60.9	61.7
Midwest	90.6	90.6	74.4	73.8	60.4	59.8
East North Central	90.4	90.5	74.6	73.8	60.2	59.9
West North Central	90.9	90.9	73.8	74.0	60.9	59.7
South	84.7	85.8	67.2	67.1	52.1	52.7
South Atlantic	85.7	86.9	69.0	69.6	54.6	55.4
East South Central	81.3	82.8	60.3	61.4	45.6	46.8
West South Central	84.5	85.3	67.5	65.6	51.2	51.2
West	85.0	85.9	68.8	70.2	55.0	56.6
Mountain	87.0	86.2	69.8	69.2	56.5	55.0
Pacific	84.2	85.8	68.4	70.6	54.3	57.3
COUNTY TYPE						
Large Metro	86.6	87.7	72.6	73.6	58.8	59.8
Small Metro	87.9	88.0	70.1	69.8	55.8	55.5
250K – 1 Mil. Pop.	88.2	87.9	71.4	69.9	57.5	56.3
< 250K Pop.	87.1	88.2	67.4	69.5	52.2	53.9
Nonmetro	85.5	86.3	65.5	64.0	49.8	49.5
Urbanized	86.6	88.5	69.1	68.2	53.5	54.1
Less Urbanized	85.1	85.1	63.5	62.5	47.5	47.8
Completely Rural	82.5	82.8	58.2	53.6	44.0	38.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.72A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	136,868 ^a	139,677	60,064	60,941	16,541	16,342
GEOGRAPHIC DIVISION						
Northeast	27,501	27,788	11,260	11,401	2,736	3,141
New England	7,724	7,608	3,059	2,944	799	807
Middle Atlantic	19,777	20,180	8,201	8,457	1,937 ^a	2,334
Midwest	31,317	31,121	14,408	14,396	4,042	4,000
East North Central	21,661	21,565	9,946	10,207	2,807	2,676
West North Central	9,657	9,556	4,462	4,189	1,235	1,324
South	46,979	48,184	20,795	21,260	5,988	5,893
South Atlantic	26,098	26,824	10,928	11,226	3,168	3,101
East South Central	6,536	6,742	2,860	3,105	900	840
West South Central	14,345	14,617	7,008	6,929	1,920	1,952
West	31,071 ^a	32,584	13,601	13,884	3,775	3,308
Mountain	9,717	9,661	4,580	4,133	1,375	1,140
Pacific	21,354 ^a	22,922	9,021	9,751	2,400	2,168
COUNTY TYPE						
Large Metro	76,521 ^a	80,932	32,885	34,476	8,748	8,816
Small Metro	40,688	39,824	18,212	17,616	5,275	4,909
250K – 1 Mil. Pop.	28,317	27,297	12,447	11,838	3,597	3,297
< 250K Pop.	12,371	12,527	5,765	5,778	1,678	1,612
Nonmetro	19,659	18,921	8,967	8,849	2,518	2,618
Urbanized	9,463	8,652	4,251	3,872	1,154	1,112
Less Urbanized	8,376	8,745	3,862	4,291	1,019 ^a	1,310
Completely Rural	1,820	1,524	854	685	344	196

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.72B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	52.2	52.7	22.9	23.0	6.3	6.2
GEOGRAPHIC DIVISION						
Northeast	58.0	58.3	23.8	23.9	5.8	6.6
New England	61.9	60.5	24.5	23.4	6.4	6.4
Middle Atlantic	56.7	57.5	23.5	24.1	5.5	6.7
Midwest	55.7	55.1	25.6	25.5	7.2	7.1
East North Central	55.6	55.1	25.5	26.1	7.2	6.8
West North Central	56.0	55.1	25.9	24.1	7.2	7.6
South	48.2	48.7	21.3	21.5	6.1	6.0
South Atlantic	50.7	51.4	21.2	21.5	6.2	5.9
East South Central	42.2	43.2	18.5	19.9	5.8	5.4
West South Central	46.9	47.0	22.9	22.3	6.3	6.3
West	50.7	52.4	22.2	22.3	6.2	5.3
Mountain	52.0	50.8	24.5 ^a	21.7	7.4	6.0
Pacific	50.2	53.1	21.2	22.6	5.6	5.0
COUNTY TYPE						
Large Metro	54.3	55.2	23.3	23.5	6.2	6.0
Small Metro	51.6	51.4	23.1	22.8	6.7	6.3
250K – 1 Mil. Pop.	53.0	52.1	23.3	22.6	6.7	6.3
< 250K Pop.	48.5	50.0	22.6	23.1	6.6	6.4
Nonmetro	46.3	46.0	21.1	21.5	5.9	6.4
Urbanized	49.8	50.2	22.4	22.5	6.1	6.4
Less Urbanized	44.1	44.4	20.3	21.8	5.4	6.6
Completely Rural	40.7	36.2	19.1	16.3	7.7 ^a	4.7

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.73A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	2,877	2,872	1,556	1,512	293	257
GEOGRAPHIC DIVISION						
Northeast	564	564	318	320	71	55
New England	149	130	92 ^a	60	18	9
Middle Atlantic	415	434	226	260	54	46
Midwest	619	556	322	275	61	52
East North Central	458 ^a	385	234	191	49	37
West North Central	161	171	88	84	12	15
South	1,027	1,019	527	527	83	107
South Atlantic	540	508	264	264	39	38
East South Central	140	130	74	64	19	9
West South Central	346	381	190	199	25 ^a	61
West	667	733	388	390	77 ^a	43
Mountain	216	240	119	123	28	18
Pacific	451	493	269	267	48	25
COUNTY TYPE						
Large Metro	1,589	1,668	835	869	152	133
Small Metro	850	780	463	393	87 ^a	51
250K – 1 Mil. Pop.	582	550	309	283	61	36
< 250K Pop.	268	229	154 ^a	110	26	15
Nonmetro	437	425	258	249	53	73
Urbanized	189	157	113	103	16	25
Less Urbanized	199	221	115	123	30	40
Completely Rural	50	47	29	23	7	8

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.73B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	11.6	11.5	6.2	6.1	1.2	1.0
GEOGRAPHIC DIVISION						
Northeast	13.5	13.6	7.6	7.7	1.7	1.3
New England	13.6	12.0	8.4 ^a	5.5	1.6	0.9
Middle Atlantic	13.4	14.1	7.3	8.5	1.7	1.5
Midwest	11.5	10.4	6.0	5.1	1.1	1.0
East North Central	12.2	10.3	6.2	5.1	1.3	1.0
West North Central	9.8	10.4	5.4	5.1	0.8	0.9
South	11.0	10.8	5.6	5.6	0.9	1.1
South Atlantic	11.6	10.8	5.7	5.6	0.8	0.8
East South Central	9.5	8.9	5.0	4.4	1.3	0.6
West South Central	10.7	11.7	5.9	6.1	0.8 ^a	1.9
West	11.2	12.4	6.5	6.6	1.3 ^a	0.7
Mountain	11.5	12.7	6.4	6.5	1.5	0.9
Pacific	11.1	12.2	6.6	6.6	1.2	0.6
COUNTY TYPE						
Large Metro	11.7	11.9	6.2	6.2	1.1	0.9
Small Metro	11.4	11.0	6.2	5.6	1.2 ^a	0.7
250K – 1 Mil. Pop.	11.4	11.3	6.0	5.8	1.2	0.7
< 250K Pop.	11.5	10.4	6.6	5.0	1.1	0.7
Nonmetro	11.3	11.2	6.6	6.6	1.4	1.9
Urbanized	11.4	10.2	6.8	6.7	1.0	1.6
Less Urbanized	11.2	11.8	6.5	6.6	1.7	2.1
Completely Rural	11.3	12.9	6.6	6.3	1.6	2.2

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.74A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	20,738	20,815	13,187	13,184	3,939	3,782
GEOGRAPHIC DIVISION						
Northeast	3,933	3,962	2,517	2,508	751	719
New England	1,127	1,154	738	732	245	241
Middle Atlantic	2,806	2,808	1,779	1,776	506	478
Midwest	4,773	4,662	3,169	3,192	1,015	999
East North Central	3,309	3,166	2,191	2,158	674	638
West North Central	1,464	1,496	978	1,035	341	361
South	7,277	7,291	4,427	4,411	1,349	1,269
South Atlantic	3,826	3,793	2,273	2,300	723	676
East South Central	1,060	1,097	664	648	199	174
West South Central	2,390	2,401	1,490	1,463	427	419
West	4,756	4,900	3,074	3,073	823	795
Mountain	1,404	1,435	905	920	215	270
Pacific	3,352	3,465	2,168	2,154	608	525
COUNTY TYPE						
Large Metro	11,242	11,503	7,045	7,210	2,081	1,985
Small Metro	6,564	6,389	4,219	4,033	1,286	1,203
250K – 1 Mil. Pop.	4,237	4,163	2,697	2,582	823	772
< 250K Pop.	2,327	2,226	1,523	1,451	463	431
Nonmetro	2,932	2,923	1,923	1,941	572	594
Urbanized	1,442	1,429	934	900	293	252
Less Urbanized	1,260	1,299	841	904	236	297
Completely Rural	230	196	148	137	43	46

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.74B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	59.6	59.6	37.9	37.7	11.3	10.8
GEOGRAPHIC DIVISION						
Northeast	64.0	64.4	40.9	40.8	12.2	11.7
New England	69.1	70.2	45.3	44.6	15.0	14.6
Middle Atlantic	62.1	62.3	39.4	39.4	11.2	10.6
Midwest	64.4	62.8	42.8	43.0	13.7	13.5
East North Central	64.8	62.0	42.9	42.2	13.2	12.5
West North Central	63.6	64.5	42.5	44.6	14.8	15.6
South	56.6	56.3	34.4	34.1	10.5	9.8
South Atlantic	58.3	57.5	34.6	34.9	11.0	10.3
East South Central	51.9	53.5	32.6	31.6	9.8	8.5
West South Central	56.2	55.8	35.0	34.0	10.0	9.7
West	56.8	58.2	36.7	36.5	9.8	9.4
Mountain	55.4	56.1	35.8	35.9	8.5	10.5
Pacific	57.4	59.2	37.1	36.8	10.4	9.0
COUNTY TYPE						
Large Metro	60.0	60.0	37.6	37.6	11.1	10.4
Small Metro	60.4	60.1	38.8	37.9	11.8	11.3
250K – 1 Mil. Pop.	60.6	60.5	38.6	37.5	11.8	11.2
< 250K Pop.	59.9	59.2	39.2	38.6	11.9	11.5
Nonmetro	56.7	57.0	37.2	37.9	11.1	11.6
Urbanized	58.3	59.7	37.8	37.6	11.9	10.5
Less Urbanized	56.7	55.4	37.9	38.6	10.6	12.7
Completely Rural	48.5	50.4	31.3	35.3	9.1	11.8

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.75A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	113,253 ^a	115,990	45,322	46,245	12,310	12,303
GEOGRAPHIC DIVISION						
Northeast	23,004	23,262	8,425	8,573	1,914 ^a	2,367
New England	6,448	6,324	2,229	2,151	537	557
Middle Atlantic	16,556	16,938	6,195	6,422	1,377 ^a	1,809
Midwest	25,925	25,903	10,917	10,929	2,965	2,948
East North Central	17,894	18,014	7,521	7,858	2,083	2,001
West North Central	8,031	7,889	3,396	3,070	882	947
South	38,675	39,874	15,841	16,322	4,556	4,517
South Atlantic	21,732	22,524	8,391	8,663	2,406	2,388
East South Central	5,336	5,515	2,122	2,392	682	657
West South Central	11,608	11,835	5,328	5,267	1,468	1,472
West	25,648	26,951	10,139	10,421	2,875	2,471
Mountain	8,097	7,986	3,555 ^a	3,090	1,131	853
Pacific	17,551 ^a	18,965	6,584	7,330	1,744	1,617
COUNTY TYPE						
Large Metro	63,690 ^a	67,761	25,005	26,397	6,515	6,698
Small Metro	33,274	32,656	13,530	13,190	3,902	3,655
250K – 1 Mil. Pop.	23,498	22,584	9,441	8,973	2,714	2,489
< 250K Pop.	9,776	10,072	4,089	4,216	1,188	1,166
Nonmetro	16,289	15,573	6,786	6,658	1,892	1,950
Urbanized	7,832	7,067	3,204	2,869	845	835
Less Urbanized	6,917	7,225	2,905	3,264	753	973
Completely Rural	1,540	1,280	676	525	294 ^a	142

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.75B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	55.9	56.5	22.4	22.5	6.1	6.0
GEOGRAPHIC DIVISION						
Northeast	62.1	62.3	22.7	23.0	5.2 ^a	6.3
New England	66.1	64.3	22.8	21.9	5.5	5.7
Middle Atlantic	60.7	61.6	22.7	23.4	5.0 ^a	6.6
Midwest	59.7	59.3	25.1	25.0	6.8	6.8
East North Central	59.4	59.5	25.0	26.0	6.9	6.6
West North Central	60.4	58.9	25.6	22.9	6.6	7.1
South	51.4	52.1	21.0	21.3	6.1	5.9
South Atlantic	54.0	55.1	20.9	21.2	6.0	5.8
East South Central	44.5	45.7	17.7	19.8	5.7	5.4
West South Central	50.3	50.3	23.1	22.4	6.4	6.3
West	54.6	56.3	21.6	21.8	6.1	5.2
Mountain	56.7	54.9	24.9 ^a	21.2	7.9 ^a	5.9
Pacific	53.7	57.0	20.2	22.0	5.3	4.9
COUNTY TYPE						
Large Metro	58.6	59.8	23.0	23.3	6.0	5.9
Small Metro	54.9	54.7	22.3	22.1	6.4	6.1
250K – 1 Mil. Pop.	56.9	55.6	22.9	22.1	6.6	6.1
< 250K Pop.	50.6	52.8	21.2	22.1	6.2	6.1
Nonmetro	48.8	48.3	20.3	20.6	5.7	6.0
Urbanized	52.7	53.1	21.6	21.6	5.7	6.3
Less Urbanized	46.2	46.7	19.4	21.1	5.0	6.3
Completely Rural	43.4	37.1	19.1	15.2	8.3 ^a	4.1

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.76A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	133,991 ^a	136,804	58,508	59,429	16,248	16,085
GEOGRAPHIC DIVISION						
Northeast	26,937	27,224	10,942	11,081	2,665	3,086
New England	7,575	7,478	2,967	2,884	782	798
Middle Atlantic	19,362	19,746	7,975	8,198	1,883 ^a	2,288
Midwest	30,698	30,565	14,086	14,121	3,980	3,948
East North Central	21,203	21,180	9,712	10,016	2,758	2,639
West North Central	9,495	9,385	4,374	4,105	1,222	1,308
South	45,952	47,165	20,268	20,733	5,905	5,786
South Atlantic	25,558	26,317	10,664	10,962	3,129	3,064
East South Central	6,396	6,612	2,786	3,041	881	831
West South Central	13,998	14,236	6,818	6,730	1,895	1,891
West	30,404 ^a	31,850	13,213	13,494	3,698	3,266
Mountain	9,501	9,421	4,461	4,010	1,347	1,123
Pacific	20,903 ^a	22,429	8,752	9,484	2,352	2,143
COUNTY TYPE						
Large Metro	74,932 ^a	79,264	32,050	33,607	8,596	8,683
Small Metro	39,838	39,044	17,749	17,223	5,188	4,858
250K – 1 Mil. Pop.	27,735	26,747	12,137	11,555	3,537	3,261
< 250K Pop.	12,103	12,297	5,612	5,668	1,651	1,597
Nonmetro	19,221	18,496	8,709	8,599	2,464	2,545
Urbanized	9,274	8,496	4,138	3,769	1,138	1,087
Less Urbanized	8,177	8,524	3,747	4,168	989	1,270
Completely Rural	1,770	1,476	824	662	337 ^a	188

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.76B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	56.4	56.9	24.6	24.7	6.8	6.7
GEOGRAPHIC DIVISION						
Northeast	62.4	62.6	25.3	25.5	6.2	7.1
New England	66.5	65.1	26.0	25.1	6.9	7.0
Middle Atlantic	60.9	61.7	25.1	25.6	5.9 ^a	7.2
Midwest	60.4	59.8	27.7	27.6	7.8	7.7
East North Central	60.2	59.9	27.6	28.3	7.8	7.5
West North Central	60.9	59.7	28.1	26.1	7.8	8.3
South	52.1	52.7	23.0	23.2	6.7	6.5
South Atlantic	54.6	55.4	22.8	23.1	6.7	6.5
East South Central	45.6	46.8	19.9	21.5	6.3	5.9
West South Central	51.2	51.2	24.9	24.2	6.9	6.8
West	55.0	56.6	23.9	24.0	6.7	5.8
Mountain	56.5	55.0	26.5 ^a	23.4	8.0	6.6
Pacific	54.3	57.3	22.7	24.2	6.1	5.5
COUNTY TYPE						
Large Metro	58.8	59.8	25.1	25.4	6.7	6.6
Small Metro	55.8	55.5	24.8	24.5	7.3	6.9
250K – 1 Mil. Pop.	57.5	56.3	25.1	24.3	7.3	6.9
< 250K Pop.	52.2	53.9	24.2	24.8	7.1	7.0
Nonmetro	49.8	49.5	22.6	23.0	6.4	6.8
Urbanized	53.5	54.1	23.9	24.0	6.6	6.9
Less Urbanized	47.5	47.8	21.8	23.4	5.8	7.1
Completely Rural	44.0	38.4	20.5	17.2	8.4 ^a	4.9

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.77A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	17,251	16,749	14,692	14,305	8,651	8,661
GENDER						
Male	8,791	8,334	7,425	7,041	4,482	4,424
Female	8,459	8,415	7,267	7,263	4,170	4,236
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	13,512	12,927	11,621	11,144	6,913	6,819
White	10,003	9,588	8,797	8,477	5,390	5,316
Black or African American	2,235	2,119	1,769	1,661	953	921
American Indian or Alaska Native	98	95	75	77	44	54
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	635	612	548	498	303	265
Two or More Races	465	454	378	380	188	232
Hispanic or Latino	3,738	3,822	3,071	3,161	1,738	1,841
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	5,110	4,788	4,465	4,182	2,817	2,746
Female, White, Not Hispanic	4,893	4,801	4,333	4,295	2,574	2,571
Male, Black, Not Hispanic	1,138	1,034	896	803	466	469
Female, Black, Not Hispanic	1,097	1,085	873	858	486	452
Male, Hispanic	1,876	1,890	1,523	1,543	893	895
Female, Hispanic	1,862	1,932	1,548	1,618	845	946

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.77B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	45.3	44.1	38.6	37.7	22.7	22.8
GENDER						
Male	45.0 ^a	42.9	38.0 ^a	36.2	23.0	22.8
Female	45.6	45.4	39.2	39.2	22.5	22.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	45.6	44.2	39.2	38.1	23.3	23.3
White	48.0	46.8	42.2	41.4	25.8	26.0
Black or African American	41.9	39.8	33.1	31.2	17.8	17.3
American Indian or Alaska Native	39.8	38.6	30.7	31.3	17.8	21.9
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	31.8	31.1	27.4	25.4	15.2	13.5
Two or More Races	44.0	41.3	35.7	34.6	17.8	21.1
Hispanic or Latino	44.3	43.9	36.4	36.3	20.6	21.2
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	47.7	45.7	41.7	39.9	26.3	26.2
Female, White, Not Hispanic	48.2	48.0	42.7	42.9	25.3	25.7
Male, Black, Not Hispanic	41.7	38.3	32.8	29.8	17.1	17.4
Female, Black, Not Hispanic	42.1	41.3	33.5	32.7	18.7	17.2
Male, Hispanic	43.6	42.0	35.4	34.3	20.7	19.9
Female, Hispanic	45.1	45.9	37.5	38.4	20.5	22.5

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.78A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	196,543	201,017	159,274	162,251	128,217	131,016
GENDER						
Male	98,395	100,277	80,760	81,838	68,124	69,134
Female	98,149	100,740	78,514	80,413	60,093	61,881
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	170,194	173,502	138,407	139,963	112,199	114,009
White	137,070	138,569	112,400	112,465	92,611	92,972
Black or African American	21,129	21,857	16,551	16,962	12,660	13,069
American Indian or Alaska Native	999	1,036	671	712	479	557
Native Hawaiian or Other Pacific Islander	*	668	*	482	*	357
Asian	7,467	8,310	6,135	6,860	4,304	5,099
Two or More Races	2,983	3,062	2,234	2,482	1,847	1,955
Hispanic or Latino	26,349	27,515	20,867	22,288	16,017	17,007
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	67,716	68,287	56,103	56,052	48,519	48,623
Female, White, Not Hispanic	69,354	70,282	56,296	56,413	44,091	44,349
Male, Black, Not Hispanic	10,157	10,314	8,130	8,052	6,523	6,431
Female, Black, Not Hispanic	10,971	11,544	8,421	8,910	6,137	6,637
Male, Hispanic	14,492	15,014	11,652	12,328	9,403	9,834
Female, Hispanic	11,857	12,501	9,215	9,960	6,614	7,174

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.78B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	87.6 ^a	88.5	71.0	71.4	57.2	57.7
GENDER						
Male	91.5	92.0	75.1	75.1	63.3	63.4
Female	84.1	85.3	67.3	68.1	51.5	52.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	88.9 ^a	89.7	72.3	72.4	58.6	59.0
White	91.9	92.5	75.4	75.1	62.1	62.1
Black or African American	81.6	82.9	63.9	64.4	48.9	49.6
American Indian or Alaska Native	86.5	86.7	58.1	59.5	41.4	46.6
Native Hawaiian or Other Pacific Islander	*	77.5	*	56.0	*	41.4
Asian	65.8	69.9	54.0	57.7	37.9	42.9
Two or More Races	92.3	92.3	69.1	74.8	57.2	58.9
Hispanic or Latino	80.3	81.5	63.6	66.0	48.8	50.4
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	94.0	94.2	77.9	77.3	67.3	67.1
Female, White, Not Hispanic	90.0	90.9	73.1	73.0	57.2	57.4
Male, Black, Not Hispanic	87.9	87.5	70.3	68.3	56.4	54.6
Female, Black, Not Hispanic	76.5	79.2	58.7	61.2	42.8	45.6
Male, Hispanic	88.4	89.8	71.1	73.7	57.3	58.8
Female, Hispanic	72.2	73.3	56.1	58.4	40.3	42.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.79A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	8,651	8,661	5,399	5,251	1,412	1,301
GENDER						
Male	4,482	4,424	3,094	2,999	908	829
Female	4,170	4,236	2,305	2,252	505	471
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	6,913	6,819	4,264	4,111	1,162	1,036
White	5,390	5,316	3,494	3,341	1,041	906
Black or African American	953	921	447	454	70	69
American Indian or Alaska Native	44	54	34	35	5	8
Native Hawaiian or Other Pacific Islander	*	*	19	*	*	2
Asian	303	265	152	132	19	21
Two or More Races	188	232	117	132	24	30
Hispanic or Latino	1,738	1,841	1,135	1,139	250	265
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	2,817	2,746	2,051	1,955	674	598
Female, White, Not Hispanic	2,574	2,571	1,443	1,386	367	308
Male, Black, Not Hispanic	466	469	219	244	40	34
Female, Black, Not Hispanic	486	452	228	211	30	35
Male, Hispanic	893	895	648	600	166	149
Female, Hispanic	845	946	487	539	84	115

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.79B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	22.7	22.8	14.2	13.8	3.7	3.4
GENDER						
Male	23.0	22.8	15.8	15.4	4.6	4.3
Female	22.5	22.9	12.4	12.1	2.7	2.5
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	23.3	23.3	14.4	14.0	3.9	3.5
White	25.8	26.0	16.8	16.3	5.0	4.4
Black or African American	17.8	17.3	8.4	8.5	1.3	1.3
American Indian or Alaska Native	17.8	21.9	13.9	14.3	1.8	3.4
Native Hawaiian or Other Pacific Islander	*	*	12.1	*	*	1.0
Asian	15.2	13.5	7.6	6.7	1.0	1.1
Two or More Races	17.8	21.1	11.1	12.0	2.2	2.7
Hispanic or Latino	20.6	21.2	13.5	13.1	3.0	3.0
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	26.3	26.2	19.2	18.7	6.3	5.7
Female, White, Not Hispanic	25.3	25.7	14.2	13.9	3.6	3.1
Male, Black, Not Hispanic	17.1	17.4	8.0	9.0	1.5	1.3
Female, Black, Not Hispanic	18.7	17.2	8.7	8.0	1.1	1.3
Male, Hispanic	20.7	19.9	15.1	13.4	3.9	3.3
Female, Hispanic	20.5	22.5	11.8	12.8	2.0	2.7

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.80A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	128,217	131,016	54,665	55,691	15,129	15,042
GENDER						
Male	68,124	69,134	35,284	35,587	11,186	11,109
Female	60,093	61,881	19,381	20,104	3,943	3,933
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	112,199	114,009	45,852	46,344	13,382	13,128
White	92,611	92,972	37,296	36,734	11,354	11,116
Black or African American	12,660	13,069	5,833	6,403	1,335	1,365
American Indian or Alaska Native	479	557	296	365	77	124
Native Hawaiian or Other Pacific Islander	*	357	196	169	74	45
Asian	4,304	5,099	1,506	1,879	244	251
Two or More Races	1,847	1,955	725	794	297	226
Hispanic or Latino	16,017	17,007	8,813	9,346	1,747	1,914
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	48,519	48,623	23,916	23,713	8,312	8,193
Female, White, Not Hispanic	44,091	44,349	13,380	13,021	3,043	2,923
Male, Black, Not Hispanic	6,523	6,431	3,595	3,706	959	935
Female, Black, Not Hispanic	6,137	6,637	2,238	2,697	377	430
Male, Hispanic	9,403	9,834	5,945	6,252	1,382	1,507
Female, Hispanic	6,614	7,174	2,868	3,095	364	407

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.80B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	57.2	57.7	24.4	24.5	6.7	6.6
GENDER						
Male	63.3	63.4	32.8	32.7	10.4	10.2
Female	51.5	52.4	16.6	17.0	3.4	3.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	58.6	59.0	23.9	24.0	7.0	6.8
White	62.1	62.1	25.0	24.5	7.6	7.4
Black or African American	48.9	49.6	22.5	24.3	5.2	5.2
American Indian or Alaska Native	41.4	46.6	25.6	30.5	6.7	10.4
Native Hawaiian or Other Pacific Islander	*	41.4	27.6	19.7	10.4	5.3
Asian	37.9	42.9	13.3	15.8	2.2	2.1
Two or More Races	57.2	58.9	22.4	23.9	9.2	6.8
Hispanic or Latino	48.8	50.4	26.8	27.7	5.3	5.7
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	67.3	67.1	33.2	32.7	11.5	11.3
Female, White, Not Hispanic	57.2	57.4	17.4	16.8	3.9	3.8
Male, Black, Not Hispanic	56.4	54.6	31.1	31.4	8.3	7.9
Female, Black, Not Hispanic	42.8	45.6	15.6	18.5	2.6	3.0
Male, Hispanic	57.3	58.8	36.3	37.4	8.4	9.0
Female, Hispanic	40.3	42.1	17.5	18.2	2.2	2.4

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.81A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	17,251	16,749	14,692	14,305	8,651	8,661
GEOGRAPHIC DIVISION						
Northeast	3,026	3,134	2,663	2,793	1,654	1,791
New England	839	920	750	844	503	551
Middle Atlantic	2,187	2,214	1,913	1,949	1,152	1,240
Midwest	3,868	3,617	3,325	3,062	2,009	1,834
East North Central	2,752	2,514	2,370	2,110	1,458 ^a	1,225
West North Central	1,116	1,104	955	953	552	609
South	6,136	5,861	5,132	4,989	2,871	2,888
South Atlantic	3,071	2,874	2,540	2,470	1,427	1,406
East South Central	916	887	778	756	414	446
West South Central	2,149	2,101	1,814	1,762	1,029	1,035
West	4,220	4,136	3,571	3,460	2,117	2,148
Mountain	1,290	1,175	1,084	1,007	600	615
Pacific	2,930	2,961	2,487	2,453	1,516	1,532
COUNTY TYPE						
Large Metro	9,214	9,129	7,928	7,828	4,690	4,729
Small Metro	5,336	5,026	4,527	4,318	2,665	2,633
250K – 1 Mil. Pop.	3,406	3,272	2,860	2,821	1,690	1,765
< 250K Pop.	1,930	1,754	1,667	1,497	976	868
Nonmetro	2,701	2,594	2,237	2,158	1,296	1,298
Urbanized	1,202	1,144	998	955	579	601
Less Urbanized	1,225	1,254	1,008	1,054	576	605
Completely Rural	274	196	230 ^a	148	140	92

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.81B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	45.3	44.1	38.6	37.7	22.7	22.8
GEOGRAPHIC DIVISION						
Northeast	47.4	48.2	41.7	42.9	25.9	27.5
New England	50.1	51.6	44.8	47.3	30.0	30.9
Middle Atlantic	46.5	46.9	40.7	41.3	24.5	26.3
Midwest	47.1	44.6	40.5 ^a	37.7	24.5	22.6
East North Central	48.1 ^a	44.8	41.4 ^a	37.6	25.5 ^a	21.8
West North Central	44.7	44.1	38.3	38.0	22.1	24.3
South	43.6	41.6	36.5	35.4	20.4	20.5
South Atlantic	43.9	41.3	36.3	35.5	20.4	20.2
East South Central	41.4	39.4	35.1	33.6	18.7	19.8
West South Central	44.2	43.1	37.3	36.2	21.2	21.3
West	44.8	44.5	37.9	37.3	22.5	23.1
Mountain	45.5	41.5	38.3	35.6	21.2	21.8
Pacific	44.5	45.8	37.8	38.0	23.0	23.7
COUNTY TYPE						
Large Metro	44.7	43.3	38.4	37.1	22.7	22.4
Small Metro	46.2	45.2	39.2	38.8	23.1	23.7
250K – 1 Mil. Pop.	44.4	43.6	37.3	37.6	22.0	23.5
< 250K Pop.	49.8	48.3	43.0	41.2	25.2	23.9
Nonmetro	45.7	45.1	37.9	37.5	21.9	22.6
Urbanized	46.5	46.8	38.6	39.1	22.4	24.6
Less Urbanized	45.9	44.7	37.8	37.6	21.6	21.6
Completely Rural	41.9	39.3	35.2	29.7	21.5	18.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.82A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	196,543	201,017	159,274	162,251	128,217	131,016
GEOGRAPHIC DIVISION						
Northeast	36,751	37,264	31,246	31,751	25,847	25,997
New England	10,093	10,031	8,554	8,541	7,221	7,057
Middle Atlantic	26,659	27,233	22,692	23,210	18,626	18,940
Midwest	43,857	44,238	35,833	35,883	29,308	29,287
East North Central	30,313	30,625	24,897	24,868	20,203	20,340
West North Central	13,544	13,614	10,936	11,015	9,105	8,947
South	71,372	73,529	56,310	57,175	44,108	45,296
South Atlantic	38,528	39,698	30,904	31,669	24,671	25,418
East South Central	10,901	11,190	7,997	8,207	6,122	6,296
West South Central	21,943	22,641	17,409	17,298	13,315	13,582
West	44,563	45,986	35,885	37,443	28,954	30,436
Mountain	13,931	14,099	11,120	11,261	9,116	9,046
Pacific	30,632	31,886	24,765	26,182	19,837	21,390
COUNTY TYPE						
Large Metro	105,243 ^b	111,217	87,903 ^a	93,112	71,831 ^a	76,203
Small Metro	59,778	58,969	47,411	46,464	38,023	37,191
250K – 1 Mil. Pop.	40,724	39,914	32,832	31,564	26,627	25,532
< 250K Pop.	19,054	19,055	14,579	14,901	11,395	11,658
Nonmetro	31,522	30,831	23,960	22,675	18,363	17,623
Urbanized	14,339	13,211	11,386	10,118	8,884	8,052
Less Urbanized	13,999	14,524	10,358	10,560	7,800	8,139
Completely Rural	3,184	3,096	2,216	1,997	1,679	1,432

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.82B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	87.6 ^a	88.5	71.0	71.4	57.2	57.7
GEOGRAPHIC DIVISION						
Northeast	89.6	90.6	76.2	77.2	63.0	63.2
New England	93.4	93.0	79.2	79.2	66.8	65.5
Middle Atlantic	88.3	89.7	75.1	76.5	61.7	62.4
Midwest	91.4	91.5	74.7	74.2	61.1	60.6
East North Central	91.1	91.4	74.9	74.2	60.7	60.7
West North Central	91.9	91.7	74.2	74.2	61.8	60.3
South	85.5	86.7	67.5	67.4	52.9	53.4
South Atlantic	86.7	87.8	69.5	70.1	55.5	56.2
East South Central	82.1	83.8	60.2	61.5	46.1	47.2
West South Central	85.4	86.4	67.7	66.0	51.8	51.8
West	85.9	86.9	69.2	70.8	55.8	57.5
Mountain	87.8	87.1	70.1	69.6	57.5	55.9
Pacific	85.1	86.8	68.8	71.3	55.1	58.3
COUNTY TYPE						
Large Metro	87.4 ^a	88.7	73.0	74.2	59.7	60.7
Small Metro	88.8	89.0	70.4	70.1	56.5	56.1
250K – 1 Mil. Pop.	89.1	88.9	71.8	70.3	58.2	56.9
< 250K Pop.	88.1	89.0	67.4	69.6	52.7	54.5
Nonmetro	86.3	87.1	65.6	64.0	50.2	49.8
Urbanized	87.4	89.3	69.4	68.4	54.1	54.4
Less Urbanized	85.8	86.0	63.5	62.5	47.8	48.2
Completely Rural	83.5	83.4	58.1	53.8	44.1	38.6

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.83A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	8,651	8,661	5,399	5,251	1,412	1,301
GEOGRAPHIC DIVISION						
Northeast	1,654	1,791	1,040	1,054	294	306
New England	503	551	330	316	96	102
Middle Atlantic	1,152	1,240	710	739	198	204
Midwest	2,009	1,834	1,265	1,147	361	303
East North Central	1,458 ^a	1,225	913	752	246	196
West North Central	552	609	352	396	115	107
South	2,871	2,888	1,720	1,728	461	445
South Atlantic	1,427	1,406	825	834	214	195
East South Central	414	446	247	272	71	63
West South Central	1,029	1,035	648	622	177	187
West	2,117	2,148	1,374	1,321	295	247
Mountain	600	615	388	378	89	84
Pacific	1,516	1,532	986	942	206	162
COUNTY TYPE						
Large Metro	4,690	4,729	2,811	2,776	712	648
Small Metro	2,665	2,633	1,738	1,612	469	416
250K – 1 Mil. Pop.	1,690	1,765	1,089	1,038	296	242
< 250K Pop.	976	868	649	574	172	174
Nonmetro	1,296	1,298	849	862	231	237
Urbanized	579	601	383	395	109	91
Less Urbanized	576	605	380	408	102	131
Completely Rural	140	92	86	58	20	15

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.83B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	22.7	22.8	14.2	13.8	3.7	3.4
GEOGRAPHIC DIVISION						
Northeast	25.9	27.5	16.3	16.2	4.6	4.7
New England	30.0	30.9	19.7	17.7	5.7	5.7
Middle Atlantic	24.5	26.3	15.1	15.6	4.2	4.3
Midwest	24.5	22.6	15.4	14.1	4.4	3.7
East North Central	25.5 ^a	21.8	15.9 ^a	13.4	4.3	3.5
West North Central	22.1	24.3	14.1	15.8	4.6	4.3
South	20.4	20.5	12.2	12.3	3.3	3.2
South Atlantic	20.4	20.2	11.8	12.0	3.1	2.8
East South Central	18.7	19.8	11.1	12.1	3.2	2.8
West South Central	21.2	21.3	13.3	12.8	3.6	3.8
West	22.5	23.1	14.6	14.2	3.1	2.7
Mountain	21.2	21.8	13.7	13.4	3.1	3.0
Pacific	23.0	23.7	15.0	14.6	3.1	2.5
COUNTY TYPE						
Large Metro	22.7	22.4	13.6	13.2	3.5	3.1
Small Metro	23.1	23.7	15.1	14.5	4.1	3.7
250K – 1 Mil. Pop.	22.0	23.5	14.2	13.8	3.9	3.2
< 250K Pop.	25.2	23.9	16.7	15.8	4.4	4.8
Nonmetro	21.9	22.6	14.4	15.0	3.9	4.1
Urbanized	22.4	24.6	14.8	16.2	4.2	3.7
Less Urbanized	21.6	21.6	14.3	14.5	3.8	4.7
Completely Rural	21.5	18.4	13.2	11.7	3.0	2.9

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.84A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	128,217	131,016	54,665	55,691	15,129	15,042
GEOGRAPHIC DIVISION						
Northeast	25,847	25,997	10,220	10,347	2,442	2,835
New England	7,221	7,057	2,729	2,628	703	705
Middle Atlantic	18,626	18,940	7,491	7,719	1,739	2,130
Midwest	29,308	29,287	13,143	13,249	3,680	3,697
East North Central	20,203	20,340	9,033	9,456	2,560	2,480
West North Central	9,105	8,947	4,110	3,793	1,120	1,217
South	44,108	45,296	19,075	19,532	5,527	5,448
South Atlantic	24,671	25,418	10,102	10,392	2,954	2,907
East South Central	6,122	6,296	2,613	2,833	829	777
West South Central	13,315	13,582	6,360	6,307	1,743	1,765
West	28,954	30,436	12,227	12,564	3,480	3,062
Mountain	9,116	9,046	4,192	3,755	1,286	1,056
Pacific	19,837	21,390	8,035	8,809	2,194	2,005
COUNTY TYPE						
Large Metro	71,831 ^a	76,203	30,074	31,700	8,036	8,168
Small Metro	38,023	37,191	16,474	16,004	4,806	4,493
250K – 1 Mil. Pop.	26,627	25,532	11,357	10,800	3,301	3,055
< 250K Pop.	11,395	11,658	5,116	5,204	1,505	1,439
Nonmetro	18,363	17,623	8,118	7,987	2,286	2,381
Urbanized	8,884	8,052	3,868	3,477	1,045	1,021
Less Urbanized	7,800	8,139	3,482	3,883	917	1,179
Completely Rural	1,679	1,432	767	627	324	181

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.84B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	57.2	57.7	24.4	24.5	6.7	6.6
GEOGRAPHIC DIVISION						
Northeast	63.0	63.2	24.9	25.2	6.0	6.9
New England	66.8	65.5	25.3	24.4	6.5	6.5
Middle Atlantic	61.7	62.4	24.8	25.4	5.8 ^a	7.0
Midwest	61.1	60.6	27.4	27.4	7.7	7.6
East North Central	60.7	60.7	27.2	28.2	7.7	7.4
West North Central	61.8	60.3	27.9	25.5	7.6	8.2
South	52.9	53.4	22.9	23.0	6.6	6.4
South Atlantic	55.5	56.2	22.7	23.0	6.6	6.4
East South Central	46.1	47.2	19.7	21.2	6.2	5.8
West South Central	51.8	51.8	24.7	24.1	6.8	6.7
West	55.8	57.5	23.6	23.8	6.7	5.8
Mountain	57.5	55.9	26.4 ^a	23.2	8.1	6.5
Pacific	55.1	58.3	22.3	24.0	6.1	5.5
COUNTY TYPE						
Large Metro	59.7	60.7	25.0	25.3	6.7	6.5
Small Metro	56.5	56.1	24.5	24.1	7.1	6.8
250K – 1 Mil. Pop.	58.2	56.9	24.8	24.1	7.2	6.8
< 250K Pop.	52.7	54.5	23.6	24.3	7.0	6.7
Nonmetro	50.2	49.8	22.2	22.6	6.3	6.7
Urbanized	54.1	54.4	23.6	23.5	6.4	6.9
Less Urbanized	47.8	48.2	21.3	23.0	5.6	7.0
Completely Rural	44.1	38.6	20.1	16.9	8.5 ^a	4.9

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150331

Table 2.1C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	942	826	875	778	835	736
Cigarettes	966	845	830	750	798	700
Smokeless Tobacco	727	579	334	276	300	249
Cigars	922	755	493	394	354	282
Pipe Tobacco ²	768	582	--	--	151	139
ALCOHOL	801	635	1,007	833	1,076	869
Binge Alcohol Use ³	--	--	--	--	804	690
Heavy Alcohol Use ³	--	--	--	--	438	368

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.1D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.36	0.31	0.33	0.29	0.32	0.28
Cigarettes	0.37	0.32	0.32	0.28	0.30	0.26
Smokeless Tobacco	0.28	0.22	0.13	0.10	0.11	0.09
Cigars	0.35	0.28	0.19	0.15	0.14	0.11
Pipe Tobacco ²	0.29	0.22	--	--	0.06	0.05
ALCOHOL	0.31	0.24	0.38	0.31	0.41	0.33
Binge Alcohol Use ³	--	--	--	--	0.31	0.26
Heavy Alcohol Use ³	--	--	--	--	0.17	0.14

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150331

Table 2.1P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.1860		0.6427		0.5459	
Cigarettes	0.1151		0.3341		0.2954	
Smokeless Tobacco	0.1556		0.3238		0.5359	
Cigars	0.5170		0.0219		0.1856	
Pipe Tobacco ²	0.3906		--		0.8172	
ALCOHOL	0.0901		0.5535		0.3203	
Binge Alcohol Use ³	--		--		0.8131	
Heavy Alcohol Use ³	--		--		0.5200	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.2C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	32	32	23	23	16	15
Cigarettes	26	28	20	19	13	12
Smokeless Tobacco	17	15	13	11	6	6
Cigars	15	15	12	13	5	7
Pipe Tobacco ²	10	11	--	--	6	5
ALCOHOL	41	39	31	33	17	22
Binge Alcohol Use ³	--	--	--	--	12	14
Heavy Alcohol Use ³	--	--	--	--	4	3

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.2D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.38	0.39	0.28	0.29	0.20	0.18
Cigarettes	0.32	0.34	0.24	0.24	0.16	0.15
Smokeless Tobacco	0.21	0.19	0.16	0.14	0.07	0.08
Cigars	0.18	0.20	0.15	0.16	0.07	0.09
Pipe Tobacco ²	0.13	0.13	--	--	0.08	0.06
ALCOHOL	0.47	0.46	0.37	0.40	0.22	0.27
Binge Alcohol Use ³	--	--	--	--	0.15	0.17
Heavy Alcohol Use ³	--	--	--	--	0.05	0.04

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150331

Table 2.2P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.6400		0.1638		0.3933	
Cigarettes	0.5319		0.0826		0.4531	
Smokeless Tobacco	0.4594		0.2957		0.9827	
Cigars	0.6707		0.7759		0.8162	
Pipe Tobacco ²	0.8659		--		0.8343	
ALCOHOL	0.1568		0.5855		0.9320	
Binge Alcohol Use ³	--		--		0.9162	
Heavy Alcohol Use ³	--		--		0.7633	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.3C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	59	54	47	42	33	31
Cigarettes	49	47	38	34	26	25
Smokeless Tobacco	32	28	27	25	17	17
Cigars	31	34	25	27	14	16
Pipe Tobacco ²	17	17	--	--	10	8
ALCOHOL	74	72	59	62	39	40
Binge Alcohol Use ³	--	--	--	--	27	26
Heavy Alcohol Use ³	--	--	--	--	9	10

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.3D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.59	0.59	0.51	0.47	0.37	0.36
Cigarettes	0.52	0.52	0.43	0.39	0.30	0.29
Smokeless Tobacco	0.36	0.32	0.30	0.29	0.20	0.20
Cigars	0.36	0.38	0.29	0.31	0.16	0.19
Pipe Tobacco ²	0.20	0.20	--	--	0.12	0.10
ALCOHOL	0.74	0.72	0.63	0.65	0.44	0.45
Binge Alcohol Use ³	--	--	--	--	0.31	0.30
Heavy Alcohol Use ³	--	--	--	--	0.11	0.11

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.3P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.0013		0.0045		0.0135	
Cigarettes	0.0013		0.0010		0.0312	
Smokeless Tobacco	0.0525		0.2460		0.3604	
Cigars	0.7566		0.8346		0.8948	
Pipe Tobacco ²	0.1668		--		0.7606	
ALCOHOL	0.3021		0.1624		0.1382	
Binge Alcohol Use ³	--		--		0.1547	
Heavy Alcohol Use ³	--		--		0.3317	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150331

Table 2.4C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	77	87	67	73	51	52
Cigarettes	68	76	54	61	43	44
Smokeless Tobacco	42	48	35	38	24	27
Cigars	55	56	44	49	29	29
Pipe Tobacco ²	22	31	--	--	13	17
ALCOHOL	101	107	93	97	65	72
Binge Alcohol Use ³	--	--	--	--	48	51
Heavy Alcohol Use ³	--	--	--	--	22	22

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.4D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.74	0.80	0.67	0.73	0.55	0.56
Cigarettes	0.69	0.74	0.58	0.63	0.48	0.48
Smokeless Tobacco	0.47	0.52	0.39	0.43	0.28	0.31
Cigars	0.58	0.61	0.49	0.55	0.33	0.33
Pipe Tobacco ²	0.26	0.35	--	--	0.15	0.20
ALCOHOL	0.79	0.80	0.80	0.81	0.66	0.72
Binge Alcohol Use ³	--	--	--	--	0.52	0.55
Heavy Alcohol Use ³	--	--	--	--	0.25	0.26

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.4P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.2523		0.2047		0.1557	
Cigarettes	0.0428		0.0309		0.0716	
Smokeless Tobacco	0.9789		0.7404		0.5491	
Cigars	0.2956		0.8669		0.1049	
Pipe Tobacco ²	0.0614		--		0.2508	
ALCOHOL	0.0832		0.5716		0.5122	
Binge Alcohol Use ³	--		--		0.9899	
Heavy Alcohol Use ³	--		--		0.4486	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.5C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	182	182	155	159	126	129
Cigarettes	160	162	133	138	111	111
Smokeless Tobacco	85	86	64	63	48	46
Cigars	131	139	108	111	68	70
Pipe Tobacco ²	64	60	--	--	36	32
ALCOHOL	224	227	212	216	168	186
Binge Alcohol Use ³	--	--	--	--	136	140
Heavy Alcohol Use ³	--	--	--	--	63	66

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.5D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.77	0.90	0.75	0.88	0.74	0.80
Cigarettes	0.80	0.90	0.75	0.86	0.71	0.74
Smokeless Tobacco	0.56	0.61	0.44	0.47	0.34	0.35
Cigars	0.70	0.86	0.64	0.72	0.46	0.49
Pipe Tobacco ²	0.45	0.44	--	--	0.27	0.24
ALCOHOL	0.70	0.78	0.75	0.84	0.79	0.92
Binge Alcohol Use ³	--	--	--	--	0.76	0.82
Heavy Alcohol Use ³	--	--	--	--	0.42	0.48

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150331

Table 2.5P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.2234		0.1200		0.0068	
Cigarettes	0.0649		0.0449		0.0029	
Smokeless Tobacco	0.0434		0.1066		0.0347	
Cigars	0.6369		0.3875		0.4146	
Pipe Tobacco ²	0.0942		--		0.0183	
ALCOHOL	0.2945		0.2324		0.7424	
Binge Alcohol Use ³	--		--		0.5887	
Heavy Alcohol Use ³	--		--		0.4091	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150331

Table 2.6C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	286	280	210	220	175	195
Cigarettes	257	266	184	203	157	179
Smokeless Tobacco	135	133	73	86	55	70
Cigars	211	206	123	131	81	87
Pipe Tobacco ²	88	100	--	--	36	35
ALCOHOL	346	331	331	322	294	286
Binge Alcohol Use ³	--	--	--	--	218	220
Heavy Alcohol Use ³	--	--	--	--	101	113

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.6D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.55	0.62	0.66	0.67	0.61	0.66
Cigarettes	0.56	0.67	0.63	0.69	0.59	0.65
Smokeless Tobacco	0.50	0.51	0.33	0.37	0.25	0.31
Cigars	0.60	0.65	0.49	0.51	0.36	0.37
Pipe Tobacco ²	0.36	0.41	--	--	0.16	0.16
ALCOHOL	0.37	0.40	0.48	0.51	0.60	0.64
Binge Alcohol Use ³	--	--	--	--	0.63	0.69
Heavy Alcohol Use ³	--	--	--	--	0.39	0.46

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.6P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.1153		0.0831		0.1097	
Cigarettes	0.0665		0.0787		0.0351	
Smokeless Tobacco	0.5016		0.8728		0.3874	
Cigars	0.2699		0.3191		0.8526	
Pipe Tobacco ²	0.9808		--		0.7091	
ALCOHOL	0.8092		0.7954		0.5945	
Binge Alcohol Use ³	--		--		1.0000	
Heavy Alcohol Use ³	--		--		0.3992	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.7C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	302	231	334	263	311	240
Cigarettes	312	248	320	252	302	228
Smokeless Tobacco	253	184	143	108	122	94
Cigars	331	231	238	169	166	124
Pipe Tobacco ²	176	136	--	--	63	46
ALCOHOL	209	149	277	199	322	258
Binge Alcohol Use ³	--	--	--	--	318	239
Heavy Alcohol Use ³	--	--	--	--	221	152

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.7D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.81	0.61	0.89	0.69	0.83	0.63
Cigarettes	0.83	0.65	0.86	0.66	0.81	0.60
Smokeless Tobacco	0.68	0.48	0.38	0.28	0.33	0.25
Cigars	0.89	0.61	0.64	0.45	0.44	0.33
Pipe Tobacco ²	0.47	0.36	--	--	0.17	0.12
ALCOHOL	0.56	0.39	0.74	0.53	0.86	0.68
Binge Alcohol Use ³	--	--	--	--	0.85	0.63
Heavy Alcohol Use ³	--	--	--	--	0.59	0.40

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.7P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.0460		0.0008		0.0007	
Cigarettes	0.0093		0.0002		0.0004	
Smokeless Tobacco	0.6744		0.8188		0.9935	
Cigars	0.0705		0.0235		0.0740	
Pipe Tobacco ²	0.2768		--		0.9134	
ALCOHOL	0.1715		0.8176		0.9006	
Binge Alcohol Use ³	--		--		0.0735	
Heavy Alcohol Use ³	--		--		0.0059	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150331

Table 2.8C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	828	710	747	652	714	618
Cigarettes	854	730	693	623	681	588
Smokeless Tobacco	667	527	267	224	254	210
Cigars	832	662	411	298	291	234
Pipe Tobacco ²	726	527	--	--	127	121
ALCOHOL	684	523	890	723	956	750
Binge Alcohol Use ³	--	--	--	--	664	543
Heavy Alcohol Use ³	--	--	--	--	344	297

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.8D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.50	0.42	0.45	0.39	0.43	0.37
Cigarettes	0.52	0.44	0.42	0.37	0.41	0.35
Smokeless Tobacco	0.40	0.31	0.16	0.13	0.15	0.13
Cigars	0.50	0.40	0.25	0.18	0.18	0.14
Pipe Tobacco ²	0.44	0.31	--	--	0.08	0.07
ALCOHOL	0.41	0.31	0.54	0.43	0.58	0.45
Binge Alcohol Use ³	--	--	--	--	0.40	0.32
Heavy Alcohol Use ³	--	--	--	--	0.21	0.18

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.8P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.7977		0.0783		0.1068	
Cigarettes	0.8860		0.1271		0.1893	
Smokeless Tobacco	0.3363		0.6626		0.5889	
Cigars	0.8286		0.3221		0.8436	
Pipe Tobacco ²	0.2609		--		0.8995	
ALCOHOL	0.0561		0.3868		0.2780	
Binge Alcohol Use ³	--		--		0.2239	
Heavy Alcohol Use ³	--		--		0.2047	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.9C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	562	467	630	543	612	513
Cigarettes	613	513	571	503	549	477
Smokeless Tobacco	634	509	312	265	282	240
Cigars	704	569	416	344	312	256
Pipe Tobacco ²	683	520	--	--	146	131
ALCOHOL	458	351	641	504	693	553
Binge Alcohol Use ³	--	--	--	--	607	519
Heavy Alcohol Use ³	--	--	--	--	367	311

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.9D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.44	0.36	0.50	0.42	0.48	0.40
Cigarettes	0.48	0.40	0.45	0.39	0.43	0.37
Smokeless Tobacco	0.50	0.40	0.25	0.21	0.22	0.19
Cigars	0.55	0.44	0.33	0.27	0.25	0.20
Pipe Tobacco ²	0.54	0.40	--	--	0.12	0.10
ALCOHOL	0.36	0.27	0.50	0.39	0.55	0.43
Binge Alcohol Use ³	--	--	--	--	0.48	0.40
Heavy Alcohol Use ³	--	--	--	--	0.29	0.24

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.9P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.3979		0.7522		1.0000	
Cigarettes	0.3125		0.4426		0.4719	
Smokeless Tobacco	0.3239		0.6290		0.8091	
Cigars	0.9035		0.2563		0.4990	
Pipe Tobacco ²	0.2055		--		0.5303	
ALCOHOL	0.5753		0.7985		0.8148	
Binge Alcohol Use ³	--		--		0.8169	
Heavy Alcohol Use ³	--		--		0.5640	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150331

Table 2.10C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	698	594	568	472	524	444
Cigarettes	693	606	546	470	513	438
Smokeless Tobacco	283	194	108	63	94	51
Cigars	506	389	250	161	161	111
Pipe Tobacco ²	237	195	--	--	35	43
ALCOHOL	598	467	709	600	736	620
Binge Alcohol Use ³	--	--	--	--	463	380
Heavy Alcohol Use ³	--	--	--	--	232	165

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.10D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.52	0.43	0.42	0.35	0.39	0.32
Cigarettes	0.51	0.44	0.40	0.34	0.38	0.32
Smokeless Tobacco	0.21	0.14	0.08	0.05	0.07	0.04
Cigars	0.37	0.28	0.18	0.12	0.12	0.08
Pipe Tobacco ²	0.18	0.14	--	--	0.03	0.03
ALCOHOL	0.44	0.34	0.52	0.44	0.54	0.45
Binge Alcohol Use ³	--	--	--	--	0.34	0.28
Heavy Alcohol Use ³	--	--	--	--	0.17	0.12

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.10P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOBACCO PRODUCTS¹	0.2583		0.2772		0.3309	
Cigarettes	0.1873		0.4918		0.3843	
Smokeless Tobacco	0.1353		0.0695		0.1602	
Cigars	0.3045		0.0051		0.0969	
Pipe Tobacco ²	0.3907		--		0.1678	
ALCOHOL	0.0632		0.2900		0.2246	
Binge Alcohol Use ³	--		--		0.4647	
Heavy Alcohol Use ³	--		--		0.7538	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.11C Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	942	826	875	778	835	736
12	16	17	13	10	8	8
13	26	26	19	21	13	12
14	39	33	30	27	20	18
15	41	44	36	33	26	25
16	51	55	43	45	31	33
17	57	62	48	53	39	38
18	87	97	76	88	63	70
19	96	92	81	79	67	65
20	96	102	84	89	74	76
21	95	104	83	83	70	74
22	103	111	80	90	71	80
23	102	102	82	80	75	70
24	102	104	84	81	72	70
25	104	111	70	91	61	81
26-29	417	281	326	216	278	193
30-34	463	330	324	248	290	207
35-39	392	314	257	220	240	203
40-44	438	320	277	225	248	204
45-49	395	334	271	219	259	206
50-54	686	531	400	347	358	327
55-59	582	456	349	286	303	271
60-64	503	442	266	254	249	237
65 or Older	497	430	344	275	335	259

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.11D Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.36	0.31	0.33	0.29	0.32	0.28
12	0.42	0.43	0.34	0.27	0.20	0.21
13	0.58	0.62	0.43	0.49	0.31	0.30
14	0.79	0.76	0.62	0.62	0.44	0.43
15	0.88	0.93	0.82	0.74	0.61	0.58
16	0.97	1.04	0.88	0.89	0.69	0.72
17	1.13	1.14	0.99	1.07	0.84	0.84
18	1.28	1.56	1.23	1.51	1.17	1.29
19	1.29	1.39	1.28	1.39	1.22	1.29
20	1.18	1.41	1.25	1.44	1.23	1.36
21	1.19	1.27	1.35	1.36	1.24	1.36
22	1.20	1.35	1.28	1.36	1.24	1.34
23	1.10	1.26	1.24	1.34	1.23	1.28
24	1.23	1.32	1.38	1.40	1.32	1.29
25	1.25	1.20	1.31	1.45	1.25	1.38
26-29	1.18	0.91	1.28	0.95	1.19	0.90
30-34	1.04	0.77	1.17	0.91	1.12	0.82
35-39	1.13	0.88	1.12	0.89	1.09	0.86
40-44	1.07	0.85	1.08	0.87	1.02	0.83
45-49	1.06	0.80	1.09	0.85	1.09	0.85
50-54	1.35	1.11	1.47	1.18	1.36	1.14
55-59	1.60	1.23	1.46	1.18	1.32	1.13
60-64	1.67	1.29	1.38	1.30	1.31	1.23
65 or Older	1.14	0.96	0.79	0.61	0.77	0.58

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.11P Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1860		0.6427		0.5459	
12	0.5219		0.4583		0.8534	
13	0.3813		0.2713		0.4019	
14	0.0213		0.3648		0.1609	
15	0.0126		0.0019		0.0278	
16	0.2763		0.1233		0.5309	
17	0.5734		0.7086		0.1710	
18	0.2350		0.2545		0.3100	
19	0.5708		0.5792		0.0122	
20	0.5494		0.2294		0.1019	
21	0.0830		0.2493		0.2946	
22	0.3351		0.5690		0.5948	
23	0.6021		0.0944		0.0235	
24	0.4396		0.0761		0.2564	
25	0.7576		0.8079		0.9446	
26-29	0.3031		0.0204		0.1720	
30-34	0.0544		0.0091		0.0006	
35-39	0.3007		0.1393		0.2025	
40-44	0.2746		0.5510		0.9395	
45-49	0.2691		0.1303		0.1722	
50-54	0.2253		0.4354		0.2353	
55-59	0.4794		0.0718		0.0205	
60-64	0.6586		0.1898		0.1653	
65 or Older	0.8823		0.5343		0.9120	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.12C Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	966	845	830	750	798	700
12	11	14	10	8	7	4
13	22	22	16	17	10	11
14	31	28	22	21	14	14
15	35	39	31	28	21	20
16	45	48	36	39	26	28
17	51	54	41	43	33	32
18	75	84	63	73	54	60
19	87	83	74	67	62	57
20	89	97	77	83	69	68
21	88	95	74	75	60	66
22	93	105	75	84	67	74
23	98	98	75	75	66	65
24	97	99	80	73	67	64
25	97	105	64	83	58	73
26-29	405	270	299	195	260	175
30-34	439	312	293	227	263	192
35-39	375	303	235	206	222	190
40-44	411	303	250	200	230	183
45-49	371	319	238	199	232	190
50-54	661	524	350	314	338	300
55-59	555	439	292	258	279	248
60-64	480	432	232	233	224	218
65 or Older	516	434	311	245	303	236

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.12D Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.37	0.32	0.32	0.28	0.30	0.26
12	0.30	0.37	0.27	0.22	0.17	0.12
13	0.52	0.53	0.38	0.41	0.23	0.26
14	0.66	0.65	0.47	0.49	0.32	0.33
15	0.78	0.84	0.72	0.64	0.51	0.47
16	0.91	0.96	0.77	0.80	0.60	0.61
17	1.06	1.07	0.88	0.91	0.74	0.71
18	1.31	1.47	1.18	1.36	1.05	1.14
19	1.35	1.44	1.32	1.33	1.25	1.21
20	1.26	1.47	1.25	1.47	1.24	1.31
21	1.28	1.35	1.31	1.40	1.15	1.28
22	1.23	1.38	1.34	1.38	1.25	1.33
23	1.20	1.33	1.27	1.29	1.19	1.22
24	1.27	1.35	1.36	1.34	1.27	1.23
25	1.24	1.28	1.24	1.42	1.22	1.31
26-29	1.25	0.97	1.26	0.91	1.17	0.85
30-34	1.05	0.84	1.11	0.87	1.07	0.77
35-39	1.17	0.91	1.07	0.89	1.04	0.85
40-44	1.12	0.85	1.03	0.82	0.98	0.78
45-49	1.08	0.87	1.00	0.82	1.00	0.82
50-54	1.39	1.21	1.34	1.11	1.30	1.08
55-59	1.65	1.34	1.28	1.12	1.24	1.07
60-64	1.73	1.34	1.25	1.21	1.22	1.15
65 or Older	1.19	0.96	0.72	0.54	0.70	0.52

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.12P Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1151		0.3341		0.2954	
12	0.3709		0.2200		0.6144	
13	0.2410		0.2092		0.5854	
14	0.0084		0.1379		0.2258	
15	0.0256		0.0018		0.0486	
16	0.3487		0.4334		0.9254	
17	0.0548		0.0242		0.0189	
18	0.1932		0.3373		0.3032	
19	0.1959		0.0872		0.0093	
20	0.3216		0.2360		0.0550	
21	0.0736		0.1473		0.2517	
22	0.1702		0.9608		0.6462	
23	0.8720		0.3044		0.0663	
24	0.4191		0.0330		0.0776	
25	0.8976		0.8037		0.9440	
26-29	0.1151		0.0044		0.0538	
30-34	0.0232		0.0083		0.0013	
35-39	0.2333		0.4940		0.3668	
40-44	0.3690		0.6302		0.7451	
45-49	0.1312		0.3501		0.4476	
50-54	0.4143		0.3139		0.4800	
55-59	0.9289		0.0552		0.0452	
60-64	0.4051		0.1096		0.1789	
65 or Older	0.7016		0.8043		0.6160	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.13C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	727	579	334	276	300	249
12	10	9	8	5	4	3
13	13	12	9	9	4	6
14	22	17	19	16	12	10
15	22	21	18	19	12	13
16	27	30	24	25	15	19
17	32	36	25	29	18	20
18	44	46	35	37	27	28
19	50	45	38	36	27	26
20	46	54	35	39	28	28
21	48	55	36	40	26	30
22	59	61	33	39	24	32
23	55	50	32	33	24	28
24	54	55	29	29	23	26
25	46	52	28	32	21	27
26-29	202	135	108	74	88	59
30-34	223	160	108	83	92	74
35-39	222	167	106	78	93	71
40-44	250	176	111	89	107	85
45-49	224	187	103	73	96	71
50-54	355	262	108	104	100	94
55-59	241	209	103	79	102	73
60-64	227	192	51	62	46	57
65 or Older	367	250	125	105	124	102

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.13D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.28	0.22	0.13	0.10	0.11	0.09
12	0.26	0.22	0.22	0.14	0.11	0.07
13	0.31	0.30	0.21	0.23	0.10	0.14
14	0.49	0.41	0.42	0.37	0.27	0.24
15	0.51	0.49	0.43	0.44	0.29	0.30
16	0.60	0.65	0.53	0.55	0.35	0.43
17	0.73	0.81	0.58	0.66	0.42	0.46
18	0.85	0.98	0.69	0.79	0.55	0.61
19	1.01	1.01	0.85	0.85	0.62	0.62
20	0.97	1.13	0.77	0.85	0.63	0.63
21	1.02	1.17	0.79	0.87	0.60	0.69
22	1.09	1.13	0.69	0.81	0.53	0.68
23	1.06	1.02	0.70	0.71	0.54	0.61
24	1.04	1.13	0.63	0.66	0.52	0.60
25	1.00	1.09	0.66	0.71	0.51	0.60
26-29	1.01	0.69	0.59	0.41	0.49	0.34
30-34	0.94	0.68	0.52	0.39	0.44	0.35
35-39	1.04	0.75	0.55	0.39	0.48	0.35
40-44	0.98	0.74	0.51	0.43	0.49	0.41
45-49	0.96	0.80	0.50	0.35	0.47	0.34
50-54	1.35	0.96	0.47	0.43	0.43	0.39
55-59	1.07	0.95	0.50	0.37	0.49	0.34
60-64	1.15	1.01	0.29	0.35	0.26	0.32
65 or Older	0.85	0.56	0.29	0.23	0.29	0.23

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.13P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1556		0.3238		0.5359	
12	0.9020		0.3058		0.1811	
13	0.3327		0.6069		0.3380	
14	0.1618		0.4918		0.4125	
15	0.1266		0.2803		0.5544	
16	0.2799		0.2513		0.9134	
17	0.3403		0.5591		0.4730	
18	0.2192		0.1577		0.0543	
19	0.0020		0.0591		0.0807	
20	0.6232		0.5946		0.9824	
21	0.1610		0.7236		0.7554	
22	0.5089		0.5083		0.1638	
23	0.0838		0.8242		0.9494	
24	0.4098		0.4832		0.7596	
25	0.7675		0.4487		0.5303	
26-29	0.9249		0.8754		0.9153	
30-34	0.6327		0.6640		0.9103	
35-39	0.5506		0.5378		0.4626	
40-44	0.4784		0.9275		0.4860	
45-49	0.4782		0.0445		0.1138	
50-54	0.1022		0.9664		0.7102	
55-59	0.4190		0.3967		0.2444	
60-64	0.9107		0.5748		0.5117	
65 or Older	0.8828		0.4329		0.5860	

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.14C Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	922	755	493	394	354	282
12	8	8	6	6	3	6
13	12	13	11	11	4	4
14	20	21	14	17	7	10
15	24	28	21	23	12	13
16	35	34	28	28	17	18
17	39	43	32	39	23	22
18	64	67	56	61	37	38
19	69	69	56	57	36	39
20	72	80	57	59	39	39
21	76	79	58	56	38	39
22	81	88	53	58	34	41
23	78	77	50	49	38	33
24	75	80	48	52	34	35
25	75	80	43	51	29	34
26-29	312	211	184	123	122	85
30-34	345	233	180	123	122	86
35-39	296	234	128	112	91	75
40-44	315	222	147	115	101	84
45-49	279	235	131	95	98	69
50-54	488	349	238	150	155	116
55-59	394	305	185	135	98	110
60-64	351	303	144	107	117	89
65 or Older	456	358	136	126	115	96

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.14D Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.35	0.28	0.19	0.15	0.14	0.11
12	0.22	0.21	0.15	0.16	0.08	0.16
13	0.29	0.32	0.25	0.27	0.10	0.09
14	0.46	0.49	0.32	0.41	0.15	0.23
15	0.56	0.63	0.49	0.53	0.29	0.31
16	0.73	0.73	0.60	0.63	0.38	0.41
17	0.87	0.92	0.74	0.85	0.54	0.51
18	1.10	1.27	1.02	1.17	0.73	0.77
19	1.21	1.33	1.09	1.18	0.80	0.87
20	1.22	1.47	1.13	1.19	0.83	0.84
21	1.25	1.33	1.14	1.16	0.84	0.87
22	1.28	1.39	1.04	1.09	0.73	0.83
23	1.23	1.35	1.00	1.04	0.82	0.73
24	1.24	1.37	0.98	1.05	0.73	0.77
25	1.32	1.41	1.01	1.07	0.71	0.77
26-29	1.25	0.90	0.92	0.68	0.66	0.48
30-34	1.21	0.82	0.82	0.56	0.58	0.40
35-39	1.23	0.92	0.65	0.54	0.47	0.37
40-44	1.13	0.86	0.64	0.54	0.46	0.40
45-49	1.08	0.88	0.61	0.44	0.48	0.33
50-54	1.55	1.18	0.98	0.61	0.67	0.48
55-59	1.57	1.18	0.87	0.64	0.47	0.52
60-64	1.59	1.39	0.78	0.60	0.64	0.50
65 or Older	1.05	0.80	0.31	0.28	0.27	0.21

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.14P Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5170		0.0219		0.1856	
12	0.6775		0.5361		0.5103	
13	0.4850		0.5433		0.6919	
14	0.9243		0.5097		0.2067	
15	0.6255		0.6803		0.4151	
16	0.0750		0.2265		0.1302	
17	0.9139		0.4525		0.3661	
18	0.3028		0.5932		0.4580	
19	0.8032		0.7787		0.5785	
20	0.8186		0.1930		0.2093	
21	0.3738		0.5152		0.6041	
22	0.8827		0.6504		0.2056	
23	0.0866		0.0585		0.0336	
24	0.6559		0.9306		0.6578	
25	0.5760		0.8110		0.5630	
26-29	0.0408		0.0554		0.1947	
30-34	0.5172		0.1775		0.2019	
35-39	0.0415		0.3084		0.5279	
40-44	0.8810		0.5348		0.7088	
45-49	0.8634		0.0828		0.0579	
50-54	0.0085		0.1168		0.6562	
55-59	0.5909		0.8446		0.0661	
60-64	0.4737		0.2162		0.5273	
65 or Older	0.0750		0.4613		0.8373	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.15C Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	801	635	1,007	833	1,076	869
12	22	19	16	13	8	8
13	32	34	25	30	15	20
14	50	45	39	38	25	22
15	51	57	45	51	30	34
16	66	72	58	64	40	45
17	72	75	66	70	49	54
18	114	130	108	120	87	97
19	109	107	103	104	85	86
20	113	118	107	112	92	102
21	118	131	116	131	105	121
22	122	134	117	131	106	117
23	118	118	112	115	104	104
24	114	121	108	115	99	104
25	116	126	110	120	99	106
26-29	468	333	446	310	410	294
30-34	506	392	470	363	413	337
35-39	445	363	405	330	367	293
40-44	513	358	472	339	412	310
45-49	443	374	407	340	371	311
50-54	739	594	676	515	602	448
55-59	656	530	550	478	493	418
60-64	578	483	488	410	458	374
65 or Older	462	343	533	443	533	451

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.15D Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.31	0.24	0.38	0.31	0.41	0.33
12	0.57	0.48	0.43	0.33	0.21	0.22
13	0.73	0.76	0.58	0.69	0.36	0.48
14	0.98	0.96	0.82	0.84	0.56	0.51
15	1.02	1.06	0.96	1.01	0.68	0.75
16	1.12	1.14	1.04	1.10	0.84	0.91
17	1.13	1.13	1.14	1.18	0.98	1.11
18	1.17	1.37	1.14	1.46	1.22	1.50
19	1.25	1.29	1.33	1.44	1.38	1.47
20	1.08	1.25	1.21	1.31	1.41	1.52
21	0.87	0.98	0.96	1.06	1.22	1.34
22	0.77	0.86	0.96	1.06	1.20	1.34
23	0.69	0.81	0.98	1.12	1.22	1.34
24	0.82	0.79	1.02	1.01	1.18	1.34
25	0.70	0.81	0.98	1.06	1.31	1.30
26-29	0.76	0.60	1.03	0.73	1.22	0.92
30-34	0.79	0.49	1.02	0.72	1.16	0.88
35-39	0.81	0.62	1.11	0.78	1.22	0.93
40-44	0.77	0.59	1.09	0.80	1.16	0.92
45-49	0.80	0.56	1.12	0.81	1.21	0.95
50-54	0.85	0.74	1.41	1.14	1.58	1.20
55-59	1.17	0.89	1.66	1.28	1.67	1.30
60-64	1.39	0.89	1.74	1.32	1.75	1.43
65 or Older	1.06	0.76	1.23	0.99	1.23	1.00

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.15P Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0901		0.5535		0.3203	
12	0.0406		0.0433		0.5875	
13	0.8201		0.5329		0.6649	
14	0.0922		0.1241		0.0255	
15	0.7860		0.3600		0.6850	
16	0.1953		0.9182		0.4316	
17	0.2608		0.4911		0.8346	
18	0.6162		0.2811		0.4899	
19	0.1282		0.1253		0.2352	
20	0.9734		0.7178		0.4137	
21	0.6353		0.8985		0.3175	
22	0.8600		0.6297		0.9133	
23	0.7305		0.5966		0.3452	
24	0.6027		0.9719		0.1092	
25	0.4781		0.3854		0.6914	
26-29	0.8723		0.5907		0.6962	
30-34	0.0459		0.8403		0.8095	
35-39	0.4366		0.2765		0.5647	
40-44	0.5768		0.5556		0.3290	
45-49	0.8284		0.7288		0.4027	
50-54	0.1388		0.0392		0.2465	
55-59	0.8654		0.2080		0.2869	
60-64	0.0752		0.5117		0.6008	
65 or Older	0.0198		0.3091		0.1383	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.16C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	1,076	869	804	690	438	368
12	8	8	4	5	3	*
13	15	20	11	13	3	3
14	25	22	17	14	5	5
15	30	34	21	22	8	9
16	40	45	28	33	11	13
17	49	54	38	41	19	18
18	87	97	63	78	30	35
19	85	86	71	68	33	34
20	92	102	71	79	40	39
21	105	121	81	95	42	51
22	106	117	83	91	44	51
23	104	104	78	80	43	43
24	99	104	78	78	43	37
25	99	106	72	84	34	44
26-29	410	294	309	210	156	97
30-34	413	337	306	227	190	117
35-39	367	293	254	192	130	104
40-44	412	310	263	199	133	102
45-49	371	311	239	201	119	99
50-54	602	448	360	292	166	149
55-59	493	418	272	250	134	117
60-64	458	374	217	177	118	94
65 or Older	533	451	297	237	150	117

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.16D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.41	0.33	0.31	0.26	0.17	0.14
12	0.21	0.22	0.11	0.13	0.07	*
13	0.36	0.48	0.25	0.31	0.07	0.08
14	0.56	0.51	0.39	0.33	0.10	0.11
15	0.68	0.75	0.50	0.50	0.20	0.20
16	0.84	0.91	0.62	0.69	0.25	0.29
17	0.98	1.11	0.83	0.91	0.44	0.44
18	1.22	1.50	1.05	1.35	0.57	0.72
19	1.38	1.47	1.28	1.33	0.71	0.78
20	1.41	1.52	1.28	1.39	0.85	0.83
21	1.22	1.34	1.31	1.43	0.86	1.03
22	1.20	1.34	1.29	1.45	0.88	1.02
23	1.22	1.34	1.30	1.29	0.87	0.90
24	1.18	1.34	1.28	1.33	0.89	0.80
25	1.31	1.30	1.28	1.41	0.78	0.94
26-29	1.22	0.92	1.28	0.93	0.82	0.53
30-34	1.16	0.88	1.10	0.82	0.86	0.53
35-39	1.22	0.93	1.14	0.82	0.67	0.50
40-44	1.16	0.92	1.03	0.81	0.59	0.48
45-49	1.21	0.95	1.00	0.83	0.58	0.45
50-54	1.58	1.20	1.35	1.07	0.70	0.61
55-59	1.67	1.30	1.20	1.06	0.64	0.55
60-64	1.75	1.43	1.17	0.94	0.65	0.52
65 or Older	1.23	1.00	0.68	0.53	0.34	0.26

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.16P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.3203		0.8131		0.5200	
12	0.5875		0.8735		*	
13	0.6649		0.9922		0.4985	
14	0.0255		0.1125		0.9615	
15	0.6850		0.4334		0.2279	
16	0.4316		0.2767		0.8986	
17	0.8346		0.3920		0.4170	
18	0.4899		0.3563		0.8814	
19	0.2352		0.0248		0.5099	
20	0.4137		0.9181		0.4550	
21	0.3175		0.3286		0.1977	
22	0.9133		0.2840		0.6027	
23	0.3452		0.8602		0.2933	
24	0.1092		0.4534		0.0043	
25	0.6914		0.2771		0.3007	
26-29	0.6962		0.1449		0.0373	
30-34	0.8095		0.2852		0.0586	
35-39	0.5647		0.8029		0.5038	
40-44	0.3290		0.1430		0.6047	
45-49	0.4027		0.7819		0.8062	
50-54	0.2465		0.9103		0.2310	
55-59	0.2869		0.0064		0.1391	
60-64	0.6008		0.6838		0.3425	
65 or Older	0.1383		0.8729		0.9278	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.17C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	942	826	875	778	835	736
AGE						
12-17	92	102	80	84	59	61
18-25	165	189	178	194	170	189
26 or Older	889	767	842	718	803	679
GENDER						
Male	562	467	630	543	612	513
Female	698	594	568	472	524	444
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	860	754	830	743	785	694
White	679	568	736	644	693	606
Black or African American	319	278	303	256	292	242
American Indian or Alaska Native	101	96	76	60	66	56
Native Hawaiian or Other Pacific Islander	74	88	55	60	47	57
Asian	316	253	156	139	136	113
Two or More Races	221	154	151	101	145	97
Hispanic or Latino	392	302	334	262	294	242

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.17D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.36	0.31	0.33	0.29	0.32	0.28
AGE						
12-17	0.37	0.41	0.32	0.34	0.24	0.25
18-25	0.48	0.54	0.51	0.55	0.49	0.54
26 or Older	0.44	0.37	0.42	0.35	0.40	0.33
GENDER						
Male	0.44	0.36	0.50	0.42	0.48	0.40
Female	0.52	0.43	0.42	0.35	0.39	0.32
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.39	0.34	0.38	0.33	0.35	0.31
White	0.40	0.33	0.43	0.38	0.41	0.36
Black or African American	1.02	0.88	0.97	0.81	0.94	0.76
American Indian or Alaska Native	2.79	2.67	3.82	2.84	3.65	2.81
Native Hawaiian or Other Pacific Islander	5.42	4.52	5.15	4.86	4.65	4.72
Asian	1.88	1.55	1.11	0.99	0.96	0.81
Two or More Races	2.12	1.79	2.74	1.92	2.73	1.87
Hispanic or Latino	0.95	0.71	0.81	0.62	0.71	0.57

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.17P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1860		0.6427		0.5459	
AGE						
12-17	0.0170		0.0108		0.0151	
18-25	0.0703		0.0268		0.0067	
26 or Older	0.3762		0.7683		0.8507	
GENDER						
Male	0.3979		0.7522		1.0000	
Female	0.2583		0.2772		0.3309	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3665		0.7350		0.5805	
White	0.6752		0.8711		0.7351	
Black or African American	0.1160		0.7442		0.6802	
American Indian or Alaska Native	0.3857		0.3360		0.6231	
Native Hawaiian or Other Pacific Islander	0.6298		0.9870		0.4722	
Asian	0.3799		0.5267		0.9396	
Two or More Races	0.7094		0.5931		0.6263	
Hispanic or Latino	0.3172		0.7951		0.9321	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.18C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	92	102	80	84	59	61
GENDER						
Male	68	74	59	66	45	47
Female	62	65	54	52	40	37
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	83	87	72	72	54	53
White	86	89	72	71	52	51
Black or African American	33	32	27	22	19	17
American Indian or Alaska Native	11	11	6	10	4	*
Native Hawaiian or Other Pacific Islander	*	*	3	*	3	*
Asian	18	15	14	13	10	8
Two or More Races	18	13	12	12	10	8
Hispanic or Latino	43	47	37	39	26	28
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	64	66	53	54	39	39
Female, White, Not Hispanic	52	55	43	43	33	31
Male, Black, Not Hispanic	25	23	22	17	15	14
Female, Black, Not Hispanic	21	21	15	15	11	10
Male, Hispanic	38	36	29	31	22	19
Female, Hispanic	32	36	27	27	17	19

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.18D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.37	0.41	0.32	0.34	0.24	0.25
GENDER						
Male	0.53	0.58	0.47	0.52	0.35	0.37
Female	0.51	0.54	0.44	0.42	0.33	0.30
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.43	0.45	0.37	0.37	0.28	0.28
White	0.51	0.57	0.45	0.48	0.35	0.36
Black or African American	0.82	0.79	0.71	0.59	0.54	0.47
American Indian or Alaska Native	3.98	4.85	3.45	4.43	2.46	*
Native Hawaiian or Other Pacific Islander	*	*	2.76	*	2.73	*
Asian	1.37	1.14	1.15	0.95	0.75	0.62
Two or More Races	2.08	1.66	1.49	1.48	1.27	1.09
Hispanic or Latino	0.78	0.84	0.67	0.70	0.47	0.49
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.73	0.83	0.64	0.72	0.51	0.55
Female, White, Not Hispanic	0.67	0.73	0.60	0.61	0.48	0.46
Male, Black, Not Hispanic	1.27	1.15	1.15	0.89	0.84	0.74
Female, Black, Not Hispanic	1.08	1.12	0.84	0.81	0.61	0.59
Male, Hispanic	1.13	1.16	0.95	1.03	0.72	0.66
Female, Hispanic	1.10	1.18	0.96	0.93	0.61	0.67

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.18P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0170		0.0108		0.0151	
GENDER						
Male	0.0146		0.0681		0.0652	
Female	0.3641		0.0714		0.1099	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0043		0.0079		0.0187	
White	0.0807		0.0590		0.0956	
Black or African American	0.0055		0.0061		0.1882	
American Indian or Alaska Native	0.1196		0.8783		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.5693		0.5088		0.3196	
Two or More Races	0.2170		0.6608		0.1646	
Hispanic or Latino	0.7652		0.6668		0.4844	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.1424		0.3355		0.4159	
Female, White, Not Hispanic	0.2910		0.0656		0.1006	
Male, Black, Not Hispanic	0.0102		0.0079		0.2518	
Female, Black, Not Hispanic	0.2239		0.3370		0.5076	
Male, Hispanic	0.5931		0.6646		0.2341	
Female, Hispanic	0.3195		0.8728		0.7854	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.19C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	165	189	178	194	170	189
GENDER						
Male	112	120	122	134	121	137
Female	106	132	110	123	101	114
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	137	166	147	172	143	164
White	271	274	224	224	183	191
Black or African American	114	116	92	100	79	90
American Indian or Alaska Native	22	20	18	17	16	16
Native Hawaiian or Other Pacific Islander	*	*	*	*	12	*
Asian	59	66	46	48	35	35
Two or More Races	38	38	30	32	27	25
Hispanic or Latino	80	89	84	87	74	80
EDUCATION						
< High School	113	114	98	101	89	92
High School Graduate	167	173	141	153	121	133
Some College	194	200	159	158	126	134
College Graduate	151	136	101	101	75	82
CURRENT EMPLOYMENT						
Full-Time	202	206	151	172	134	161
Part-Time	165	172	134	136	109	111
Unemployed	96	91	85	82	75	72
Other ¹	131	136	109	112	84	94

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.19D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.48	0.54	0.51	0.55	0.49	0.54
GENDER						
Male	0.64	0.68	0.70	0.77	0.70	0.78
Female	0.61	0.76	0.64	0.70	0.58	0.66
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.50	0.60	0.53	0.62	0.52	0.60
White	0.52	0.60	0.60	0.67	0.59	0.68
Black or African American	1.23	1.39	1.16	1.38	1.10	1.32
American Indian or Alaska Native	3.53	3.39	4.43	4.98	4.61	5.14
Native Hawaiian or Other Pacific Islander	*	*	*	*	5.30	*
Asian	2.49	2.38	2.06	2.07	1.68	1.63
Two or More Races	2.73	2.45	3.00	2.80	2.91	2.66
Hispanic or Latino	1.11	1.21	1.16	1.18	1.02	1.09
EDUCATION						
< High School	1.08	1.34	1.20	1.36	1.15	1.35
High School Graduate	0.78	0.90	0.80	0.92	0.80	0.89
Some College	0.72	0.87	0.81	0.87	0.75	0.85
College Graduate	1.29	1.31	1.33	1.35	1.14	1.19
CURRENT EMPLOYMENT						
Full-Time	0.69	0.73	0.78	0.80	0.77	0.83
Part-Time	0.83	1.00	0.90	0.99	0.82	0.89
Unemployed	1.25	1.46	1.33	1.50	1.32	1.50
Other ¹	1.12	1.18	1.03	1.09	0.96	0.99

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.19P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0703		0.0268		0.0067	
GENDER						
Male	0.0547		0.0219		0.0149	
Female	0.4109		0.3138		0.1078	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1817		0.0798		0.0091	
White	0.2519		0.0415		0.0032	
Black or African American	0.9132		0.9770		0.4790	
American Indian or Alaska Native	0.4968		0.4523		0.5993	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4953		0.8588		0.4509	
Two or More Races	0.8956		0.8629		0.0435	
Hispanic or Latino	0.1779		0.1445		0.3042	
EDUCATION						
< High School	0.8781		0.2909		0.5139	
High School Graduate	0.1692		0.1777		0.2662	
Some College	0.3854		0.2062		0.0453	
College Graduate	0.1231		0.0340		0.0318	
CURRENT EMPLOYMENT						
Full-Time	0.1637		0.2430		0.1557	
Part-Time	0.7078		0.0941		0.0078	
Unemployed	0.0254		0.2896		0.1037	
Other ¹	0.3433		0.2177		0.5133	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.20C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	889	767	842	718	803	679
GENDER						
Male	529	421	606	515	590	491
Female	680	566	554	438	511	414
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	816	708	800	690	758	645
White	1,637	1,352	880	728	781	660
Black or African American	615	472	378	323	360	299
American Indian or Alaska Native	94	89	73	54	62	50
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	300	237	143	126	128	108
Two or More Races	217	145	146	95	141	91
Hispanic or Latino	364	281	313	244	282	228
EDUCATION						
< High School	606	479	420	351	391	334
High School Graduate	867	711	571	471	525	448
Some College	821	663	501	406	465	370
College Graduate	1,125	866	496	379	414	313
CURRENT EMPLOYMENT						
Full-Time	1,300	990	737	607	639	546
Part-Time	579	426	328	241	302	223
Unemployed	287	227	221	176	214	166
Other ¹	960	876	516	468	506	445

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.20D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.44	0.37	0.42	0.35	0.40	0.33
GENDER						
Male	0.55	0.43	0.63	0.52	0.61	0.50
Female	0.64	0.53	0.52	0.41	0.48	0.39
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.47	0.40	0.46	0.39	0.44	0.37
White	0.47	0.39	0.52	0.44	0.48	0.41
Black or African American	1.36	1.14	1.30	1.04	1.28	0.98
American Indian or Alaska Native	3.63	3.34	5.07	3.72	4.97	3.58
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.43	1.96	1.36	1.17	1.20	1.01
Two or More Races	2.61	2.45	4.06	2.83	4.03	2.75
Hispanic or Latino	1.28	0.95	1.10	0.83	0.99	0.77
EDUCATION						
< High School	1.23	1.04	1.26	1.04	1.21	1.05
High School Graduate	0.76	0.62	0.82	0.64	0.79	0.62
Some College	0.76	0.64	0.82	0.66	0.78	0.61
College Graduate	0.81	0.63	0.66	0.49	0.57	0.43
CURRENT EMPLOYMENT						
Full-Time	0.58	0.43	0.57	0.45	0.51	0.43
Part-Time	1.17	0.93	1.13	0.89	1.07	0.85
Unemployed	2.08	1.64	2.25	1.86	2.21	1.83
Other ¹	0.82	0.77	0.70	0.62	0.69	0.59

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.20P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3762		0.7683		0.8507	
GENDER						
Male	0.8096		0.2988		0.4805	
Female	0.3206		0.4764		0.5993	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6148		0.7913		0.8854	
White	0.8841		0.6866		0.7186	
Black or African American	0.1717		0.9083		0.6139	
American Indian or Alaska Native	0.4641		0.2693		0.5193	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.3151		0.4687		0.7663	
Two or More Races	0.7967		0.6115		0.9728	
Hispanic or Latino	0.4011		0.8166		0.7786	
EDUCATION						
< High School	0.1442		0.5533		0.3629	
High School Graduate	0.4306		0.1968		0.1111	
Some College	0.3935		0.8405		0.9444	
College Graduate	0.5503		0.9448		0.6195	
CURRENT EMPLOYMENT						
Full-Time	0.2137		0.8972		0.7619	
Part-Time	0.0471		0.0737		0.1094	
Unemployed	0.4273		0.8382		0.6285	
Other ¹	0.5335		0.0852		0.2368	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.21C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	909	803	873	771	834	729
GENDER						
Male	542	447	635	544	615	514
Female	685	583	565	465	522	440
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	827	733	829	734	785	687
White	1,690	1,420	929	792	824	715
Black or African American	653	501	407	351	381	323
American Indian or Alaska Native	99	94	75	58	65	54
Native Hawaiian or Other Pacific Islander	*	87	55	59	47	57
Asian	313	249	155	137	136	112
Two or More Races	219	152	151	100	145	96
Hispanic or Latino	379	301	332	261	293	241
EDUCATION						
< High School	627	502	443	371	412	353
High School Graduate	896	739	600	505	552	472
Some College	849	700	526	438	483	396
College Graduate	1,155	884	519	399	431	332
CURRENT EMPLOYMENT						
Full-Time	1,337	1,023	772	641	671	582
Part-Time	612	468	355	282	322	256
Unemployed	306	254	237	201	227	186
Other ¹	961	885	532	481	518	458

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.21D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.38	0.33	0.37	0.32	0.35	0.30
GENDER						
Male	0.47	0.39	0.55	0.47	0.54	0.44
Female	0.56	0.47	0.46	0.37	0.42	0.35
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.41	0.36	0.41	0.36	0.39	0.34
White	0.42	0.35	0.47	0.41	0.44	0.38
Black or African American	1.13	0.95	1.09	0.89	1.06	0.84
American Indian or Alaska Native	3.03	2.84	4.19	3.21	4.16	3.14
Native Hawaiian or Other Pacific Islander	*	4.72	5.76	5.26	5.16	5.15
Asian	2.09	1.70	1.23	1.08	1.06	0.89
Two or More Races	2.20	1.99	3.31	2.32	3.27	2.26
Hispanic or Latino	1.06	0.82	0.93	0.71	0.82	0.66
EDUCATION						
< High School	1.03	0.91	1.07	0.92	1.03	0.93
High School Graduate	0.65	0.55	0.71	0.56	0.68	0.53
Some College	0.63	0.56	0.68	0.56	0.63	0.52
College Graduate	0.76	0.59	0.64	0.47	0.55	0.42
CURRENT EMPLOYMENT						
Full-Time	0.52	0.40	0.52	0.41	0.47	0.39
Part-Time	0.88	0.74	0.84	0.73	0.80	0.68
Unemployed	1.42	1.22	1.55	1.34	1.52	1.31
Other ¹	0.75	0.70	0.65	0.57	0.64	0.55

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.21P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2243		0.7973		0.6390	
GENDER						
Male	0.5094		0.6444		0.9215	
Female	0.2613		0.3492		0.3803	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4611		0.8915		0.6676	
White	0.7376		0.9821		0.8013	
Black or African American	0.1770		0.9120		0.7396	
American Indian or Alaska Native	0.5539		0.3208		0.5724	
Native Hawaiian or Other Pacific Islander	*		0.8786		0.4445	
Asian	0.3938		0.5076		0.9099	
Two or More Races	0.8097		0.6119		0.6995	
Hispanic or Latino	0.2581		0.8136		0.9613	
EDUCATION						
< High School	0.1708		0.5577		0.3661	
High School Graduate	0.6951		0.3350		0.1976	
Some College	0.3148		0.5810		0.5406	
College Graduate	0.4041		0.7632		0.4422	
CURRENT EMPLOYMENT						
Full-Time	0.1414		0.7756		0.9638	
Part-Time	0.0499		0.0267		0.0185	
Unemployed	0.8888		0.8703		0.3043	
Other ¹	0.6125		0.1548		0.3021	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.22C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	966	845	830	750	798	700
AGE						
12-17	83	89	67	71	51	52
18-25	170	201	171	196	162	185
26 or Older	915	788	797	691	770	651
GENDER						
Male	613	513	571	503	549	477
Female	693	606	546	470	513	438
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	885	789	781	704	743	654
White	722	605	689	610	650	562
Black or African American	330	278	285	251	280	239
American Indian or Alaska Native	97	94	68	58	64	52
Native Hawaiian or Other Pacific Islander	74	87	53	49	44	47
Asian	302	246	140	134	121	113
Two or More Races	219	149	142	87	137	84
Hispanic or Latino	390	300	308	249	284	233

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.22D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.37	0.32	0.32	0.28	0.30	0.26
AGE						
12-17	0.33	0.36	0.27	0.29	0.20	0.21
18-25	0.49	0.57	0.49	0.56	0.46	0.53
26 or Older	0.45	0.38	0.39	0.34	0.38	0.32
GENDER						
Male	0.48	0.40	0.45	0.39	0.43	0.37
Female	0.51	0.44	0.40	0.34	0.38	0.32
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.40	0.35	0.35	0.32	0.34	0.29
White	0.42	0.36	0.41	0.36	0.38	0.33
Black or African American	1.06	0.88	0.91	0.79	0.90	0.75
American Indian or Alaska Native	3.06	2.95	3.68	2.87	3.61	2.80
Native Hawaiian or Other Pacific Islander	5.46	4.61	4.99	4.21	4.44	4.05
Asian	1.82	1.52	0.99	0.95	0.85	0.81
Two or More Races	2.22	1.83	2.69	1.80	2.65	1.74
Hispanic or Latino	0.95	0.71	0.75	0.59	0.69	0.55

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.22P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1151		0.3341		0.2954	
AGE						
12-17	0.0020		0.0004		0.0115	
18-25	0.0190		0.0126		0.0013	
26 or Older	0.3210		0.9293		0.8733	
GENDER						
Male	0.3125		0.4426		0.4719	
Female	0.1873		0.4918		0.3843	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1942		0.3323		0.3132	
White	0.2882		0.4163		0.3933	
Black or African American	0.3499		0.7594		0.6365	
American Indian or Alaska Native	0.3508		0.4962		0.3854	
Native Hawaiian or Other Pacific Islander	0.6937		0.7803		0.4749	
Asian	0.4979		0.5029		0.5820	
Two or More Races	0.3516		0.2246		0.4065	
Hispanic or Latino	0.4532		0.9412		0.8423	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.23C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	83	89	67	71	51	52
GENDER						
Male	59	63	47	52	36	36
Female	59	60	49	47	37	34
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	74	74	59	59	45	44
White	73	74	58	57	44	43
Black or African American	28	26	21	16	14	11
American Indian or Alaska Native	9	10	6	10	3	6
Native Hawaiian or Other Pacific Islander	*	*	2	*	1	*
Asian	18	14	14	12	9	7
Two or More Races	17	12	11	10	9	7
Hispanic or Latino	40	44	33	35	20	24
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	51	52	40	40	30	31
Female, White, Not Hispanic	49	49	39	38	32	28
Male, Black, Not Hispanic	20	19	16	12	10	8
Female, Black, Not Hispanic	18	18	13	11	10	9
Male, Hispanic	34	32	25	27	16	16
Female, Hispanic	30	34	24	25	13	17

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.23D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.33	0.36	0.27	0.29	0.20	0.21
GENDER						
Male	0.46	0.49	0.37	0.41	0.28	0.28
Female	0.48	0.49	0.40	0.38	0.30	0.28
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.38	0.38	0.30	0.30	0.23	0.23
White	0.46	0.49	0.38	0.39	0.31	0.31
Black or African American	0.74	0.67	0.57	0.44	0.39	0.31
American Indian or Alaska Native	4.16	4.42	3.13	4.55	1.95	2.83
Native Hawaiian or Other Pacific Islander	*	*	2.52	*	1.48	*
Asian	1.35	1.06	1.12	0.88	0.71	0.55
Two or More Races	1.99	1.58	1.36	1.31	1.18	0.87
Hispanic or Latino	0.72	0.78	0.60	0.63	0.37	0.42
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.63	0.69	0.51	0.55	0.40	0.44
Female, White, Not Hispanic	0.65	0.66	0.55	0.55	0.46	0.42
Male, Black, Not Hispanic	1.07	0.99	0.89	0.66	0.57	0.46
Female, Black, Not Hispanic	0.97	0.96	0.73	0.63	0.56	0.48
Male, Hispanic	1.04	1.07	0.83	0.90	0.55	0.55
Female, Hispanic	1.05	1.10	0.87	0.86	0.48	0.61

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.23P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0020		0.0004		0.0115	
GENDER						
Male	0.0101		0.0266		0.1600	
Female	0.0647		0.0043		0.0259	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0003		0.0002		0.0032	
White	0.0047		0.0025		0.0509	
Black or African American	0.0405		0.0057		0.0491	
American Indian or Alaska Native	0.3725		0.8004		0.7602	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1713		0.2218		0.1959	
Two or More Races	0.5442		0.6005		0.0671	
Hispanic or Latino	0.9627		0.5057		0.8687	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.0298		0.1665		0.8777	
Female, White, Not Hispanic	0.0530		0.0031		0.0085	
Male, Black, Not Hispanic	0.1311		0.0305		0.1015	
Female, Black, Not Hispanic	0.1698		0.1063		0.2979	
Male, Hispanic	0.5316		0.3235		0.3048	
Female, Hispanic	0.4789		0.9484		0.1913	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.24C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	170	201	171	196	162	185
GENDER						
Male	118	132	124	139	117	132
Female	108	136	104	119	97	109
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	142	177	145	175	138	163
White	247	250	193	202	160	172
Black or African American	101	108	79	89	69	80
American Indian or Alaska Native	20	18	14	14	14	11
Native Hawaiian or Other Pacific Islander	*	*	*	*	11	*
Asian	55	60	41	42	33	34
Two or More Races	36	35	28	28	24	22
Hispanic or Latino	79	90	77	85	69	78
EDUCATION						
< High School	110	109	94	95	83	83
High School Graduate	153	160	126	140	109	123
Some College	175	182	136	137	112	118
College Graduate	132	122	80	88	59	70
CURRENT EMPLOYMENT						
Full-Time	185	198	135	162	122	144
Part-Time	147	156	113	122	93	96
Unemployed	90	90	78	77	69	69
Other ¹	119	124	95	99	76	84

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.24D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.49	0.57	0.49	0.56	0.46	0.53
GENDER						
Male	0.67	0.75	0.71	0.79	0.67	0.75
Female	0.62	0.78	0.60	0.69	0.56	0.62
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.52	0.64	0.53	0.63	0.50	0.59
White	0.56	0.68	0.60	0.71	0.58	0.69
Black or African American	1.19	1.45	1.11	1.39	1.07	1.31
American Indian or Alaska Native	4.14	4.44	4.26	5.02	4.80	4.61
Native Hawaiian or Other Pacific Islander	*	*	*	*	5.06	*
Asian	2.45	2.20	1.97	1.89	1.62	1.58
Two or More Races	2.96	2.60	2.94	2.80	2.80	2.52
Hispanic or Latino	1.09	1.22	1.07	1.15	0.95	1.06
EDUCATION						
< High School	1.11	1.33	1.19	1.34	1.11	1.29
High School Graduate	0.86	0.93	0.82	0.93	0.77	0.87
Some College	0.75	0.91	0.76	0.84	0.72	0.80
College Graduate	1.30	1.36	1.22	1.25	0.97	1.08
CURRENT EMPLOYMENT						
Full-Time	0.69	0.78	0.76	0.84	0.73	0.81
Part-Time	0.83	1.03	0.84	0.96	0.76	0.83
Unemployed	1.30	1.56	1.35	1.49	1.30	1.43
Other ¹	1.16	1.18	1.02	1.06	0.90	0.93

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.24P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0190		0.0126		0.0013	
GENDER						
Male	0.0257		0.0160		0.0059	
Female	0.1976		0.2067		0.0345	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0531		0.0193		0.0040	
White	0.0682		0.0412		0.0046	
Black or African American	0.9214		0.6288		0.9351	
American Indian or Alaska Native	0.7379		0.5825		0.7224	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.3458		0.6664		0.7214	
Two or More Races	0.4986		0.1022		0.0524	
Hispanic or Latino	0.1444		0.3276		0.0982	
EDUCATION						
< High School	0.8252		0.5440		0.7656	
High School Graduate	0.2457		0.1935		0.1305	
Some College	0.0857		0.0405		0.0199	
College Graduate	0.0855		0.2340		0.1368	
CURRENT EMPLOYMENT						
Full-Time	0.0959		0.2906		0.0263	
Part-Time	0.4810		0.0635		0.0048	
Unemployed	0.0350		0.2681		0.3091	
Other ¹	0.1351		0.0729		0.3064	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.25C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	915	788	797	691	770	651
GENDER						
Male	585	467	548	473	530	453
Female	671	573	533	439	500	412
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	842	738	752	656	720	616
White	1,540	1,303	772	647	707	591
Black or African American	580	453	344	306	331	286
American Indian or Alaska Native	93	88	62	53	60	49
Native Hawaiian or Other Pacific Islander	*	*	*	46	*	46
Asian	288	232	129	123	114	107
Two or More Races	215	141	136	83	133	81
Hispanic or Latino	363	280	292	230	275	217
EDUCATION						
< High School	577	458	381	328	363	310
High School Graduate	838	688	509	445	487	424
Some College	801	650	464	366	437	342
College Graduate	1,063	818	396	309	339	257
CURRENT EMPLOYMENT						
Full-Time	1,221	945	648	543	587	496
Part-Time	565	420	297	222	284	210
Unemployed	277	221	204	171	197	160
Other ¹	933	852	484	430	471	409

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.25D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.45	0.38	0.39	0.34	0.38	0.32
GENDER						
Male	0.60	0.48	0.57	0.48	0.55	0.46
Female	0.63	0.53	0.50	0.41	0.47	0.38
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.48	0.42	0.43	0.37	0.41	0.35
White	0.50	0.41	0.48	0.41	0.45	0.39
Black or African American	1.40	1.15	1.24	1.02	1.23	0.97
American Indian or Alaska Native	3.81	3.63	4.91	3.74	4.88	3.58
Native Hawaiian or Other Pacific Islander	*	*	*	5.39	*	5.36
Asian	2.36	1.94	1.23	1.14	1.08	0.99
Two or More Races	2.68	2.52	3.93	2.65	3.93	2.59
Hispanic or Latino	1.28	0.95	1.02	0.78	0.97	0.74
EDUCATION						
< High School	1.28	1.04	1.16	1.01	1.12	0.98
High School Graduate	0.82	0.64	0.77	0.63	0.75	0.60
Some College	0.79	0.65	0.78	0.60	0.75	0.57
College Graduate	0.84	0.66	0.55	0.43	0.48	0.37
CURRENT EMPLOYMENT						
Full-Time	0.59	0.46	0.53	0.43	0.49	0.41
Part-Time	1.21	0.98	1.04	0.84	1.01	0.81
Unemployed	2.13	1.73	2.18	1.87	2.12	1.81
Other ¹	0.85	0.78	0.67	0.58	0.66	0.56

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.25P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3210		0.9293		0.8733	
GENDER						
Male	0.6938		0.9872		0.9789	
Female	0.3048		0.8779		0.7813	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4381		0.8529		0.8168	
White	0.5399		0.8657		0.8768	
Black or African American	0.3988		0.9892		0.7076	
American Indian or Alaska Native	0.3390		0.4151		0.4125	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.3647		0.3666		0.4777	
Two or More Races	0.4122		0.3832		0.7584	
Hispanic or Latino	0.6145		0.7544		0.8020	
EDUCATION						
< High School	0.1652		0.2977		0.0859	
High School Graduate	0.4909		0.1109		0.1272	
Some College	0.5389		0.5790		0.8215	
College Graduate	0.3542		0.5923		0.7077	
CURRENT EMPLOYMENT						
Full-Time	0.2084		0.9564		0.7152	
Part-Time	0.0428		0.0687		0.0445	
Unemployed	0.4484		0.6803		0.8847	
Other ¹	0.5714		0.3250		0.4902	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.26C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	935	823	824	741	796	691
GENDER						
Male	596	496	571	502	549	476
Female	680	591	543	462	511	433
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	856	768	776	694	742	646
White	1,585	1,367	813	703	744	635
Black or African American	613	477	366	328	347	304
American Indian or Alaska Native	96	92	67	56	63	51
Native Hawaiian or Other Pacific Islander	*	86	53	48	44	47
Asian	300	243	139	133	120	112
Two or More Races	218	148	141	86	136	84
Hispanic or Latino	378	298	306	248	283	231
EDUCATION						
< High School	597	480	404	345	383	326
High School Graduate	861	710	534	471	507	445
Some College	823	685	482	393	450	366
College Graduate	1,089	834	417	328	352	275
CURRENT EMPLOYMENT						
Full-Time	1,255	980	676	578	610	526
Part-Time	591	461	319	262	302	232
Unemployed	293	246	217	195	209	181
Other ¹	935	860	497	441	482	418

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.26D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.34	0.35	0.31	0.34	0.29
GENDER						
Male	0.52	0.43	0.50	0.43	0.48	0.41
Female	0.55	0.47	0.44	0.37	0.42	0.35
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.42	0.38	0.38	0.34	0.37	0.32
White	0.45	0.37	0.44	0.38	0.41	0.36
Black or African American	1.18	0.96	1.03	0.87	1.01	0.83
American Indian or Alaska Native	3.22	3.13	4.19	3.21	4.10	3.10
Native Hawaiian or Other Pacific Islander	*	4.79	5.60	4.54	4.94	4.44
Asian	2.02	1.68	1.10	1.04	0.94	0.88
Two or More Races	2.30	2.06	3.23	2.15	3.18	2.09
Hispanic or Latino	1.06	0.81	0.86	0.67	0.79	0.63
EDUCATION						
< High School	1.08	0.91	1.00	0.88	0.96	0.87
High School Graduate	0.69	0.56	0.66	0.54	0.64	0.52
Some College	0.66	0.57	0.64	0.51	0.61	0.49
College Graduate	0.79	0.62	0.53	0.42	0.46	0.36
CURRENT EMPLOYMENT						
Full-Time	0.53	0.42	0.48	0.40	0.44	0.37
Part-Time	0.91	0.78	0.78	0.69	0.76	0.63
Unemployed	1.47	1.28	1.50	1.35	1.46	1.31
Other ¹	0.78	0.71	0.62	0.53	0.61	0.51

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.26P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1512		0.4823		0.3546	
GENDER						
Male	0.3852		0.5445		0.5064	
Female	0.2156		0.6553		0.4616	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2637		0.4745		0.3858	
White	0.3600		0.5445		0.4472	
Black or African American	0.4287		0.8972		0.7008	
American Indian or Alaska Native	0.4083		0.4802		0.3740	
Native Hawaiian or Other Pacific Islander	*		0.6637		0.4729	
Asian	0.4797		0.4490		0.5426	
Two or More Races	0.3184		0.2340		0.4804	
Hispanic or Latino	0.4012		0.9823		0.8054	
EDUCATION						
< High School	0.2065		0.2792		0.0636	
High School Graduate	0.7320		0.2143		0.2620	
Some College	0.3221		0.2681		0.3986	
College Graduate	0.2302		0.5389		0.5757	
CURRENT EMPLOYMENT						
Full-Time	0.1235		0.8347		0.9380	
Part-Time	0.0359		0.0210		0.0051	
Unemployed	0.8815		0.9931		0.6152	
Other ¹	0.7119		0.5435		0.6193	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.27C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	727	579	334	276	300	249
AGE						
12-17	59	58	49	47	31	34
18-25	137	141	94	101	71	82
26 or Older	705	563	313	255	288	234
GENDER						
Male	634	509	312	265	282	240
Female	283	194	108	63	94	51
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	685	556	319	269	286	245
White	618	504	295	249	265	227
Black or African American	212	157	109	74	100	69
American Indian or Alaska Native	54	47	22	19	16	18
Native Hawaiian or Other Pacific Islander	45	23	15	13	14	12
Asian	94	73	33	27	29	20
Two or More Races	117	75	45	27	42	20
Hispanic or Latino	224	165	96	68	83	52

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.27D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.28	0.22	0.13	0.10	0.11	0.09
AGE						
12-17	0.24	0.23	0.20	0.19	0.12	0.14
18-25	0.39	0.40	0.27	0.29	0.20	0.23
26 or Older	0.35	0.27	0.15	0.12	0.14	0.11
GENDER						
Male	0.50	0.40	0.25	0.21	0.22	0.19
Female	0.21	0.14	0.08	0.05	0.07	0.04
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.31	0.25	0.14	0.12	0.13	0.11
White	0.36	0.30	0.17	0.15	0.16	0.13
Black or African American	0.68	0.50	0.35	0.23	0.32	0.22
American Indian or Alaska Native	3.21	2.40	1.44	1.22	1.08	1.19
Native Hawaiian or Other Pacific Islander	4.59	2.32	1.67	1.28	1.61	1.18
Asian	0.70	0.53	0.25	0.20	0.22	0.15
Two or More Races	2.38	1.58	1.04	0.60	0.97	0.45
Hispanic or Latino	0.54	0.39	0.23	0.16	0.20	0.12

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.27P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1556		0.3238		0.5359	
AGE						
12-17	0.2986		0.3099		0.9525	
18-25	0.1043		0.2600		0.5588	
26 or Older	0.2938		0.6357		0.6310	
GENDER						
Male	0.3239		0.6290		0.8091	
Female	0.1353		0.0695		0.1602	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3721		0.6122		0.8782	
White	0.7106		0.7775		0.5538	
Black or African American	0.3545		0.0365		0.0624	
American Indian or Alaska Native	0.7846		0.9885		0.2783	
Native Hawaiian or Other Pacific Islander	0.2017		0.6597		0.6654	
Asian	0.6135		0.7043		0.5219	
Two or More Races	0.8421		0.8377		0.8893	
Hispanic or Latino	0.1080		0.1037		0.0968	

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.28C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	59	58	49	47	31	34
GENDER						
Male	51	51	44	43	28	32
Female	23	25	16	20	9	12
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	55	54	45	44	30	32
White	55	53	44	43	28	31
Black or African American	14	9	12	7	9	3
American Indian or Alaska Native	6	*	3	*	3	*
Native Hawaiian or Other Pacific Islander	2	*	2	*	*	*
Asian	6	6	6	6	4	3
Two or More Races	8	7	7	5	6	4
Hispanic or Latino	17	20	13	14	8	8
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	47	46	41	38	25	29
Female, White, Not Hispanic	22	24	14	19	8	12
Male, Black, Not Hispanic	13	9	12	7	9	3
Female, Black, Not Hispanic	3	2	3	1	*	*
Male, Hispanic	16	19	12	14	9	8
Female, Hispanic	7	7	6	3	0	3

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.28D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.24	0.23	0.20	0.19	0.12	0.14
GENDER						
Male	0.40	0.40	0.35	0.34	0.22	0.25
Female	0.19	0.20	0.13	0.16	0.07	0.10
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.28	0.28	0.24	0.23	0.15	0.17
White	0.37	0.37	0.30	0.31	0.20	0.23
Black or African American	0.40	0.25	0.34	0.19	0.27	0.10
American Indian or Alaska Native	3.11	*	2.04	*	1.66	*
Native Hawaiian or Other Pacific Islander	2.51	*	2.38	*	*	*
Asian	0.49	0.51	0.44	0.46	0.31	0.25
Two or More Races	1.05	0.91	0.88	0.71	0.79	0.47
Hispanic or Latino	0.31	0.35	0.24	0.25	0.15	0.15
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.62	0.64	0.54	0.53	0.35	0.41
Female, White, Not Hispanic	0.32	0.35	0.21	0.29	0.12	0.18
Male, Black, Not Hispanic	0.75	0.47	0.65	0.37	0.52	0.19
Female, Black, Not Hispanic	0.17	0.13	0.16	0.09	*	*
Male, Hispanic	0.54	0.64	0.42	0.48	0.30	0.28
Female, Hispanic	0.26	0.24	0.22	0.09	0.01	0.09

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.28P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2986		0.3099		0.9525	
GENDER						
Male	0.1644		0.1132		0.6309	
Female	0.6367		0.2550		0.1203	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2291		0.2845		0.9029	
White	0.3991		0.4885		0.5829	
Black or African American	0.1075		0.1322		0.1183	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.6530		0.7600		0.6683	
Two or More Races	0.5723		0.4946		0.1427	
Hispanic or Latino	0.6814		0.9336		0.4176	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.1948		0.1153		0.9872	
Female, White, Not Hispanic	0.5523		0.0800		0.1368	
Male, Black, Not Hispanic	0.0944		0.1571		0.1179	
Female, Black, Not Hispanic	0.9461		0.5782		*	
Male, Hispanic	0.4694		0.4754		0.6480	
Female, Hispanic	0.5175		0.0894		0.0770	

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.29C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	137	141	94	101	71	82
GENDER						
Male	113	117	87	92	67	78
Female	58	59	27	28	18	16
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	123	126	84	95	67	77
White	145	141	90	97	68	78
Black or African American	28	32	12	21	9	15
American Indian or Alaska Native	14	14	9	9	7	9
Native Hawaiian or Other Pacific Islander	6	*	4	*	*	*
Asian	17	17	11	8	5	5
Two or More Races	19	18	12	14	10	9
Hispanic or Latino	54	55	37	35	19	23
EDUCATION						
< High School	57	62	41	46	30	36
High School Graduate	86	90	57	59	43	46
Some College	88	91	59	63	43	51
College Graduate	69	57	33	36	23	27
CURRENT EMPLOYMENT						
Full-Time	106	104	68	72	52	60
Part-Time	78	77	47	55	32	39
Unemployed	52	52	36	35	25	26
Other ¹	56	62	38	43	27	36

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.29D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.40	0.27	0.29	0.20	0.23
GENDER						
Male	0.65	0.67	0.50	0.53	0.38	0.44
Female	0.33	0.34	0.15	0.16	0.11	0.09
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.45	0.46	0.31	0.34	0.24	0.28
White	0.55	0.55	0.40	0.45	0.32	0.37
Black or African American	0.54	0.59	0.24	0.41	0.17	0.29
American Indian or Alaska Native	4.20	5.04	3.55	4.01	2.72	3.81
Native Hawaiian or Other Pacific Islander	2.73	*	2.03	*	*	*
Asian	0.91	0.84	0.59	0.41	0.28	0.26
Two or More Races	2.40	2.11	1.54	1.70	1.40	1.10
Hispanic or Latino	0.75	0.75	0.52	0.48	0.26	0.32
EDUCATION						
< High School	0.93	1.08	0.69	0.86	0.52	0.69
High School Graduate	0.64	0.69	0.46	0.47	0.35	0.37
Some College	0.62	0.64	0.44	0.47	0.33	0.39
College Graduate	1.04	0.88	0.63	0.62	0.44	0.47
CURRENT EMPLOYMENT						
Full-Time	0.68	0.64	0.48	0.48	0.37	0.41
Part-Time	0.68	0.71	0.46	0.53	0.33	0.39
Unemployed	1.14	1.23	0.83	0.87	0.60	0.65
Other ¹	0.68	0.75	0.48	0.56	0.36	0.47

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.29P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1043		0.2600		0.5588	
GENDER						
Male	0.0249		0.3860		0.8554	
Female	0.5597		0.2477		0.0697	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0754		0.3360		0.5838	
White	0.0427		0.1892		0.5523	
Black or African American	0.3216		0.1126		0.7315	
American Indian or Alaska Native	0.5440		0.6604		0.3896	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.2530		0.2207		0.7431	
Two or More Races	0.4094		0.4475		0.8858	
Hispanic or Latino	0.9655		0.5868		0.9222	
EDUCATION						
< High School	0.2932		0.5085		0.3573	
High School Graduate	0.4478		0.7399		0.8042	
Some College	0.5534		0.2849		0.5923	
College Graduate	0.0020		0.2603		0.2476	
CURRENT EMPLOYMENT						
Full-Time	0.1947		0.3275		0.9858	
Part-Time	0.3694		0.8893		0.8575	
Unemployed	0.1756		0.1074		0.1805	
Other ¹	0.7338		0.7285		0.5045	

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.30C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	705	563	313	255	288	234
GENDER						
Male	615	496	293	248	270	228
Female	277	183	102	53	91	47
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	669	545	299	248	275	229
White	730	572	282	239	257	219
Black or African American	225	166	110	71	101	69
American Indian or Alaska Native	52	42	19	13	14	13
Native Hawaiian or Other Pacific Islander	*	19	14	10	13	10
Asian	92	71	30	25	28	19
Two or More Races	116	73	43	22	40	19
Hispanic or Latino	212	157	88	55	82	47
EDUCATION						
< High School	286	214	145	108	133	103
High School Graduate	396	329	184	147	173	139
Some College	384	295	155	119	149	111
College Graduate	504	361	142	119	130	105
CURRENT EMPLOYMENT						
Full-Time	612	486	256	206	235	190
Part-Time	247	177	65	53	60	48
Unemployed	151	103	83	41	74	33
Other ¹	444	322	144	138	138	131

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.30D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.35	0.27	0.15	0.12	0.14	0.11
GENDER						
Male	0.63	0.50	0.30	0.25	0.28	0.23
Female	0.26	0.17	0.10	0.05	0.09	0.04
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.38	0.31	0.17	0.14	0.16	0.13
White	0.45	0.36	0.20	0.17	0.18	0.15
Black or African American	0.91	0.66	0.47	0.30	0.43	0.29
American Indian or Alaska Native	4.36	3.15	1.83	1.25	1.34	1.21
Native Hawaiian or Other Pacific Islander	*	2.64	2.39	1.32	2.34	1.32
Asian	0.89	0.68	0.29	0.24	0.27	0.18
Two or More Races	3.58	2.41	1.52	0.77	1.42	0.64
Hispanic or Latino	0.75	0.53	0.31	0.19	0.29	0.16
EDUCATION						
< High School	0.98	0.71	0.54	0.38	0.49	0.36
High School Graduate	0.61	0.54	0.31	0.26	0.29	0.24
Some College	0.67	0.53	0.30	0.22	0.29	0.21
College Graduate	0.66	0.46	0.21	0.17	0.19	0.15
CURRENT EMPLOYMENT						
Full-Time	0.49	0.38	0.24	0.18	0.22	0.17
Part-Time	0.93	0.70	0.27	0.23	0.25	0.20
Unemployed	1.79	1.33	1.04	0.58	0.94	0.47
Other ¹	0.63	0.45	0.22	0.20	0.21	0.20

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.30P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2938		0.6357		0.6310	
GENDER						
Male	0.6358		0.9623		0.8887	
Female	0.0829		0.0676		0.2080	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6357		0.9789		0.9857	
White	0.9431		0.3771		0.4521	
Black or African American	0.3124		0.0231		0.0696	
American Indian or Alaska Native	0.8714		0.8568		0.4185	
Native Hawaiian or Other Pacific Islander	*		0.3909		0.5194	
Asian	0.7397		0.9599		0.5913	
Two or More Races	0.9944		0.9699		0.9132	
Hispanic or Latino	0.0787		0.1179		0.0751	
EDUCATION						
< High School	0.9206		0.9025		0.6666	
High School Graduate	0.5273		0.4879		0.6215	
Some College	0.2932		0.5172		0.4878	
College Graduate	0.1750		0.4654		0.9876	
CURRENT EMPLOYMENT						
Full-Time	0.4519		0.4568		0.4510	
Part-Time	0.0378		0.8324		0.9849	
Unemployed	0.5735		0.1323		0.1056	
Other ¹	0.8538		0.5354		0.4890	

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.31C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	716	576	331	271	299	246
GENDER						
Male	624	506	309	263	279	238
Female	282	195	106	61	94	50
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	678	554	316	265	285	242
White	753	596	299	256	267	234
Black or African American	228	170	110	75	101	70
American Indian or Alaska Native	55	45	21	16	15	16
Native Hawaiian or Other Pacific Islander	*	23	14	13	14	12
Asian	93	72	32	27	29	20
Two or More Races	117	75	45	26	42	20
Hispanic or Latino	223	163	95	65	82	51
EDUCATION						
< High School	295	228	152	119	136	111
High School Graduate	404	344	196	154	181	144
Some College	392	310	168	134	157	122
College Graduate	513	364	147	123	133	107
CURRENT EMPLOYMENT						
Full-Time	634	501	269	216	243	198
Part-Time	261	193	80	78	70	62
Unemployed	160	117	91	55	78	42
Other ¹	451	328	149	146	142	137

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.31D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.30	0.24	0.14	0.11	0.13	0.10
GENDER						
Male	0.55	0.44	0.27	0.23	0.24	0.21
Female	0.23	0.16	0.09	0.05	0.08	0.04
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.34	0.27	0.16	0.13	0.14	0.12
White	0.39	0.32	0.19	0.16	0.17	0.14
Black or African American	0.76	0.56	0.39	0.26	0.36	0.24
American Indian or Alaska Native	3.68	2.75	1.59	1.29	1.18	1.24
Native Hawaiian or Other Pacific Islander	*	2.57	1.84	1.42	1.80	1.30
Asian	0.77	0.57	0.27	0.22	0.24	0.16
Two or More Races	2.87	1.92	1.25	0.71	1.16	0.54
Hispanic or Latino	0.63	0.44	0.26	0.18	0.23	0.14
EDUCATION						
< High School	0.83	0.63	0.46	0.35	0.41	0.33
High School Graduate	0.52	0.47	0.28	0.22	0.25	0.21
Some College	0.55	0.45	0.26	0.20	0.25	0.18
College Graduate	0.61	0.42	0.20	0.16	0.18	0.14
CURRENT EMPLOYMENT						
Full-Time	0.44	0.35	0.22	0.17	0.20	0.15
Part-Time	0.69	0.54	0.24	0.23	0.21	0.19
Unemployed	1.25	0.95	0.74	0.49	0.64	0.38
Other ¹	0.57	0.42	0.20	0.20	0.19	0.18

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.31P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1675		0.4003		0.5218	
GENDER						
Male	0.3584		0.7974		0.8396	
Female	0.1207		0.0371		0.1015	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4069		0.7269		0.8814	
White	0.7419		0.6957		0.6002	
Black or African American	0.4043		0.0510		0.0805	
American Indian or Alaska Native	0.7979		0.9414		0.3138	
Native Hawaiian or Other Pacific Islander	*		0.7609		0.7299	
Asian	0.5764		0.6621		0.5549	
Two or More Races	0.8687		0.7501		0.9365	
Hispanic or Latino	0.0939		0.0925		0.0751	
EDUCATION						
< High School	0.9761		0.8890		0.5874	
High School Graduate	0.6587		0.4728		0.6032	
Some College	0.2444		0.3073		0.4014	
College Graduate	0.0769		0.6112		0.8669	
CURRENT EMPLOYMENT						
Full-Time	0.3312		0.3535		0.4863	
Part-Time	0.0242		0.7995		0.9432	
Unemployed	0.3047		0.0420		0.0447	
Other ¹	0.8987		0.6436		0.6172	

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.32C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	922	755	493	394	354	282
AGE						
12-17	64	68	50	57	32	34
18-25	171	189	142	151	100	105
26 or Older	885	705	469	345	335	256
GENDER						
Male	704	569	416	344	312	256
Female	506	389	250	161	161	111
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	860	696	463	365	334	261
White	764	611	413	320	290	221
Black or African American	283	244	184	139	149	121
American Indian or Alaska Native	63	66	46	21	22	15
Native Hawaiian or Other Pacific Islander	54	43	18	19	10	11
Asian	173	132	84	53	69	33
Two or More Races	137	118	57	60	37	48
Hispanic or Latino	334	270	190	152	131	107

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.32D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.35	0.28	0.19	0.15	0.14	0.11
AGE						
12-17	0.26	0.27	0.20	0.23	0.13	0.13
18-25	0.49	0.54	0.41	0.43	0.29	0.30
26 or Older	0.44	0.34	0.23	0.17	0.17	0.12
GENDER						
Male	0.55	0.44	0.33	0.27	0.25	0.20
Female	0.37	0.28	0.18	0.12	0.12	0.08
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.39	0.31	0.21	0.16	0.15	0.12
White	0.45	0.36	0.24	0.19	0.17	0.13
Black or African American	0.90	0.77	0.59	0.44	0.48	0.38
American Indian or Alaska Native	3.70	3.27	2.99	1.46	1.44	1.02
Native Hawaiian or Other Pacific Islander	5.18	3.74	2.06	1.86	1.17	1.11
Asian	1.23	0.96	0.62	0.40	0.51	0.24
Two or More Races	2.61	2.04	1.34	1.29	0.88	1.04
Hispanic or Latino	0.81	0.63	0.46	0.36	0.32	0.25

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.32P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5170		0.0219		0.1856	
AGE						
12-17	0.3729		0.9712		0.2399	
18-25	0.2966		0.1950		0.5116	
26 or Older	0.6611		0.0462		0.2945	
GENDER						
Male	0.9035		0.2563		0.4990	
Female	0.3045		0.0051		0.0969	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7750		0.0906		0.1913	
White	0.5161		0.2312		0.3518	
Black or African American	0.8055		0.1502		0.5596	
American Indian or Alaska Native	0.2386		0.2692		0.2640	
Native Hawaiian or Other Pacific Islander	0.4814		0.7586		0.5179	
Asian	0.1757		0.7001		0.1224	
Two or More Races	0.0791		0.3907		0.4647	
Hispanic or Latino	0.4000		0.0762		0.8277	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.33C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	64	68	50	57	32	34
GENDER						
Male	49	53	40	46	28	26
Female	36	39	29	31	16	20
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	57	61	44	52	28	31
White	55	56	42	49	26	28
Black or African American	20	22	16	17	11	13
American Indian or Alaska Native	5	6	3	2	1	1
Native Hawaiian or Other Pacific Islander	1	*	1	*	1	*
Asian	9	5	9	5	6	3
Two or More Races	10	9	8	8	6	4
Hispanic or Latino	30	30	25	24	17	12
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	43	46	34	41	23	22
Female, White, Not Hispanic	27	31	22	25	12	18
Male, Black, Not Hispanic	16	17	13	14	7	11
Female, Black, Not Hispanic	12	13	9	10	7	6
Male, Hispanic	26	22	20	16	14	10
Female, Hispanic	21	22	15	17	9	7

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.33D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.26	0.27	0.20	0.23	0.13	0.13
GENDER						
Male	0.39	0.42	0.31	0.36	0.22	0.21
Female	0.30	0.32	0.24	0.26	0.13	0.16
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.29	0.32	0.23	0.27	0.15	0.16
White	0.36	0.40	0.28	0.35	0.18	0.21
Black or African American	0.54	0.58	0.45	0.46	0.30	0.36
American Indian or Alaska Native	2.60	2.91	2.02	1.46	0.72	0.72
Native Hawaiian or Other Pacific Islander	1.46	*	1.23	*	1.19	*
Asian	0.72	0.43	0.69	0.40	0.45	0.21
Two or More Races	1.28	1.16	0.97	1.00	0.76	0.50
Hispanic or Latino	0.54	0.53	0.44	0.42	0.30	0.22
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.54	0.63	0.44	0.56	0.32	0.31
Female, White, Not Hispanic	0.39	0.45	0.32	0.37	0.17	0.26
Male, Black, Not Hispanic	0.87	0.88	0.72	0.73	0.42	0.60
Female, Black, Not Hispanic	0.67	0.71	0.49	0.55	0.39	0.37
Male, Hispanic	0.83	0.74	0.68	0.56	0.47	0.35
Female, Hispanic	0.73	0.77	0.54	0.63	0.32	0.24

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.33P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3729		0.9712		0.2399	
GENDER						
Male	0.1019		0.4923		0.0726	
Female	0.4941		0.2891		0.5698	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.5166		0.6781		0.6901	
White	0.6800		0.7478		0.6389	
Black or African American	0.7286		0.5764		0.2866	
American Indian or Alaska Native	0.5299		0.5427		0.5608	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.2797		0.3159		0.2287	
Two or More Races	0.7505		0.6648		0.4286	
Hispanic or Latino	0.5184		0.4837		0.0699	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.4405		0.8837		0.1462	
Female, White, Not Hispanic	0.7266		0.7133		0.1947	
Male, Black, Not Hispanic	0.7558		0.6336		0.1519	
Female, Black, Not Hispanic	0.8770		0.7465		0.9166	
Male, Hispanic	0.1725		0.1068		0.1827	
Female, Hispanic	0.5757		0.4656		0.1837	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.34C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	171	189	142	151	100	105
GENDER						
Male	125	126	114	115	86	85
Female	104	112	68	81	46	54
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	149	167	123	136	89	95
White	211	214	139	143	82	85
Black or African American	69	70	55	60	44	49
American Indian or Alaska Native	11	16	8	12	5	9
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	38	42	26	28	14	12
Two or More Races	27	29	18	22	14	15
Hispanic or Latino	75	79	61	58	41	45
EDUCATION						
< High School	78	73	58	57	45	49
High School Graduate	119	126	95	97	57	63
Some College	151	159	98	108	60	66
College Graduate	115	108	61	64	39	34
CURRENT EMPLOYMENT						
Full-Time	153	148	98	100	65	75
Part-Time	122	131	86	91	54	55
Unemployed	71	67	55	53	39	38
Other ¹	93	98	65	66	36	38

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.34D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.49	0.54	0.41	0.43	0.29	0.30
GENDER						
Male	0.71	0.72	0.65	0.65	0.50	0.49
Female	0.60	0.64	0.39	0.47	0.27	0.31
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.54	0.61	0.45	0.49	0.32	0.35
White	0.61	0.69	0.53	0.59	0.38	0.40
Black or African American	1.10	1.10	0.94	1.00	0.78	0.85
American Indian or Alaska Native	3.74	5.21	3.00	4.90	2.29	4.11
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.81	1.91	1.25	1.35	0.74	0.62
Two or More Races	3.05	2.76	2.34	2.38	1.81	1.77
Hispanic or Latino	1.04	1.07	0.84	0.79	0.57	0.62
EDUCATION						
< High School	1.06	1.18	0.90	1.01	0.74	0.90
High School Graduate	0.75	0.87	0.68	0.73	0.44	0.50
Some College	0.78	0.93	0.65	0.74	0.46	0.50
College Graduate	1.37	1.30	1.08	0.98	0.75	0.57
CURRENT EMPLOYMENT						
Full-Time	0.75	0.77	0.63	0.62	0.45	0.49
Part-Time	0.86	0.98	0.75	0.80	0.52	0.53
Unemployed	1.38	1.41	1.16	1.22	0.88	0.92
Other ¹	0.97	1.03	0.73	0.79	0.45	0.49

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.34P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2966		0.1950		0.5116	
GENDER						
Male	0.1393		0.0224		0.2462	
Female	0.9851		0.3408		0.5174	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.8235		0.8004		0.3582	
White	0.8500		0.3224		0.1581	
Black or African American	0.6119		0.3722		0.5738	
American Indian or Alaska Native	0.1004		0.1136		0.3347	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.6429		0.8358		0.3623	
Two or More Races	0.1871		0.1148		0.9028	
Hispanic or Latino	0.0572		0.0104		0.6525	
EDUCATION						
< High School	0.7580		0.2553		0.0357	
High School Graduate	0.3690		0.3092		0.7222	
Some College	0.8643		0.6137		0.2181	
College Graduate	0.0252		0.0393		0.0174	
CURRENT EMPLOYMENT						
Full-Time	0.0550		0.0235		0.8012	
Part-Time	0.8881		0.9890		0.4826	
Unemployed	0.9393		0.9776		0.3692	
Other ¹	0.8857		0.9705		0.7192	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.35C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	885	705	469	345	335	256
GENDER						
Male	682	538	397	320	293	242
Female	492	365	237	125	152	91
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	834	656	443	320	317	237
White	1,110	843	431	297	285	212
Black or African American	333	288	182	132	145	116
American Indian or Alaska Native	62	62	43	16	20	11
Native Hawaiian or Other Pacific Islander	*	40	11	12	*	8
Asian	168	126	79	45	67	30
Two or More Races	136	112	54	55	34	46
Hispanic or Latino	316	257	175	140	123	96
EDUCATION						
< High School	341	285	183	137	147	111
High School Graduate	549	419	263	175	181	141
Some College	556	432	250	196	177	145
College Graduate	814	604	315	223	203	153
CURRENT EMPLOYMENT						
Full-Time	911	672	401	293	258	204
Part-Time	388	299	167	116	122	80
Unemployed	213	146	127	70	111	60
Other ¹	588	498	207	174	159	147

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.35D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.44	0.34	0.23	0.17	0.17	0.12
GENDER						
Male	0.70	0.55	0.41	0.33	0.30	0.25
Female	0.47	0.34	0.22	0.12	0.14	0.08
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.48	0.37	0.25	0.18	0.18	0.13
White	0.55	0.42	0.29	0.21	0.20	0.15
Black or African American	1.21	1.01	0.76	0.54	0.62	0.48
American Indian or Alaska Native	4.97	4.11	3.84	1.54	1.94	1.06
Native Hawaiian or Other Pacific Islander	*	4.80	1.92	1.63	*	1.14
Asian	1.56	1.20	0.76	0.43	0.65	0.29
Two or More Races	3.95	3.02	1.93	1.84	1.20	1.53
Hispanic or Latino	1.11	0.87	0.62	0.48	0.43	0.33
EDUCATION						
< High School	1.10	0.86	0.66	0.49	0.54	0.40
High School Graduate	0.77	0.60	0.43	0.30	0.31	0.24
Some College	0.82	0.68	0.47	0.36	0.34	0.27
College Graduate	0.86	0.61	0.44	0.30	0.29	0.22
CURRENT EMPLOYMENT						
Full-Time	0.60	0.45	0.36	0.25	0.24	0.18
Part-Time	1.26	1.00	0.67	0.48	0.50	0.33
Unemployed	2.23	1.66	1.54	0.96	1.39	0.85
Other ¹	0.78	0.60	0.31	0.26	0.24	0.22

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.35P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.6611		0.0462		0.2945	
GENDER						
Male	0.8259		0.7032		0.8732	
Female	0.2576		0.0003		0.0317	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7940		0.0779		0.2986	
White	0.5180		0.3401		0.6401	
Black or African American	0.9016		0.0598		0.3725	
American Indian or Alaska Native	0.3324		0.1309		0.1378	
Native Hawaiian or Other Pacific Islander	*		0.9340		*	
Asian	0.1318		0.7712		0.1977	
Two or More Races	0.1305		0.7872		0.4221	
Hispanic or Latino	0.7361		0.4102		0.8737	
EDUCATION						
< High School	0.8265		0.4425		0.4778	
High School Graduate	0.5312		0.1290		0.9602	
Some College	0.5738		0.2213		0.9721	
College Graduate	0.3309		0.6165		0.2106	
CURRENT EMPLOYMENT						
Full-Time	0.1448		0.1104		0.4039	
Part-Time	0.8453		0.2048		0.7556	
Unemployed	0.2042		0.2367		0.1869	
Other ¹	0.1806		0.9088		0.8572	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.36C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	913	750	492	389	354	280
GENDER						
Male	698	558	417	342	312	255
Female	504	388	250	157	161	109
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	852	692	465	359	333	258
White	1,145	901	454	339	298	229
Black or African American	347	301	195	148	154	128
American Indian or Alaska Native	63	65	45	20	21	15
Native Hawaiian or Other Pacific Islander	*	42	18	18	10	11
Asian	172	132	83	53	69	33
Two or More Races	136	117	57	60	36	48
Hispanic or Latino	332	272	187	151	129	108
EDUCATION						
< High School	356	297	194	147	154	120
High School Graduate	565	442	279	202	189	152
Some College	578	470	266	232	187	163
College Graduate	835	621	325	230	209	157
CURRENT EMPLOYMENT						
Full-Time	941	690	416	312	269	218
Part-Time	409	335	189	143	132	96
Unemployed	222	164	139	86	120	70
Other ¹	595	505	217	186	164	152

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.36D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.38	0.31	0.21	0.16	0.15	0.12
GENDER						
Male	0.61	0.48	0.36	0.30	0.27	0.22
Female	0.41	0.31	0.20	0.13	0.13	0.09
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.42	0.34	0.23	0.18	0.17	0.13
White	0.49	0.39	0.26	0.20	0.18	0.14
Black or African American	1.02	0.85	0.66	0.49	0.53	0.43
American Indian or Alaska Native	4.14	3.56	3.30	1.65	1.62	1.15
Native Hawaiian or Other Pacific Islander	*	4.09	2.35	1.97	1.31	1.24
Asian	1.35	1.06	0.68	0.44	0.56	0.26
Two or More Races	3.17	2.45	1.63	1.55	1.04	1.27
Hispanic or Latino	0.93	0.74	0.52	0.41	0.36	0.29
EDUCATION						
< High School	0.93	0.75	0.57	0.45	0.46	0.37
High School Graduate	0.66	0.52	0.38	0.28	0.27	0.21
Some College	0.68	0.59	0.40	0.34	0.29	0.24
College Graduate	0.80	0.58	0.42	0.29	0.28	0.21
CURRENT EMPLOYMENT						
Full-Time	0.54	0.41	0.33	0.24	0.22	0.17
Part-Time	0.92	0.79	0.53	0.42	0.38	0.29
Unemployed	1.52	1.18	1.10	0.74	0.97	0.63
Other ¹	0.71	0.55	0.29	0.25	0.22	0.20

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.36P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5161		0.0199		0.2215	
GENDER						
Male	0.9452		0.2789		0.5997	
Female	0.2660		0.0030		0.0805	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7628		0.0778		0.1976	
White	0.4978		0.2104		0.3694	
Black or African American	0.8217		0.1329		0.4967	
American Indian or Alaska Native	0.2046		0.2849		0.2815	
Native Hawaiian or Other Pacific Islander	*		0.9350		0.4462	
Asian	0.1728		0.7631		0.1444	
Two or More Races	0.0777		0.4292		0.4116	
Hispanic or Latino	0.4154		0.0881		0.9920	
EDUCATION						
< High School	0.8851		0.4447		0.7936	
High School Graduate	0.6904		0.0900		0.9066	
Some College	0.5584		0.1758		0.6217	
College Graduate	0.2108		0.4225		0.1029	
CURRENT EMPLOYMENT						
Full-Time	0.0774		0.0423		0.4164	
Part-Time	0.8220		0.2736		0.5616	
Unemployed	0.2254		0.3328		0.1293	
Other ¹	0.1967		0.8487		0.8242	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.37C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	801	635	1,007	833	1,076	869
AGE						
12-17	109	107	100	103	73	81
18-25	136	141	158	165	183	197
26 or Older	736	553	954	764	1,021	802
GENDER						
Male	458	351	641	504	693	553
Female	598	467	709	600	736	620
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	699	532	893	739	979	789
White	531	382	752	604	848	663
Black or African American	276	225	329	256	328	261
American Indian or Alaska Native	107	109	77	84	65	72
Native Hawaiian or Other Pacific Islander	80	92	68	74	51	61
Asian	489	429	441	370	360	285
Two or More Races	226	161	181	151	174	130
Hispanic or Latino	340	278	395	327	395	332

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.37D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.31	0.24	0.38	0.31	0.41	0.33
AGE						
12-17	0.44	0.43	0.40	0.42	0.29	0.33
18-25	0.39	0.40	0.45	0.47	0.53	0.56
26 or Older	0.36	0.27	0.47	0.37	0.50	0.39
GENDER						
Male	0.36	0.27	0.50	0.39	0.55	0.43
Female	0.44	0.34	0.52	0.44	0.54	0.45
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.32	0.24	0.40	0.33	0.44	0.35
White	0.31	0.22	0.44	0.35	0.50	0.39
Black or African American	0.88	0.71	1.05	0.81	1.05	0.83
American Indian or Alaska Native	2.65	2.43	3.63	3.56	3.45	3.47
Native Hawaiian or Other Pacific Islander	5.24	4.24	5.46	5.04	4.95	4.85
Asian	2.07	1.45	2.09	1.46	2.06	1.41
Two or More Races	1.72	1.30	2.60	1.90	2.75	1.93
Hispanic or Latino	0.82	0.66	0.96	0.77	0.96	0.78

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.37P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0901		0.5535		0.3203	
AGE						
12-17	0.0585		0.2985		0.9818	
18-25	0.4537		0.6344		0.9625	
26 or Older	0.0301		0.4515		0.3269	
GENDER						
Male	0.5753		0.7985		0.8148	
Female	0.0632		0.2900		0.2246	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0984		0.9676		0.4780	
White	0.1847		0.6538		0.8944	
Black or African American	0.4391		0.9233		0.6678	
American Indian or Alaska Native	0.9685		0.7813		0.3005	
Native Hawaiian or Other Pacific Islander	0.9535		0.8033		0.9447	
Asian	0.1428		0.2320		0.0922	
Two or More Races	0.7164		0.2222		0.5338	
Hispanic or Latino	0.4146		0.1215		0.2763	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.38C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	109	107	100	103	73	81
GENDER						
Male	74	79	71	71	52	54
Female	75	77	71	74	52	57
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	95	91	87	88	64	70
White	106	101	94	94	62	70
Black or African American	49	54	40	41	27	27
American Indian or Alaska Native	9	10	8	10	4	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	3	*
Asian	31	26	31	21	20	12
Two or More Races	20	18	15	15	8	10
Hispanic or Latino	57	57	50	55	37	40
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	73	69	65	60	42	45
Female, White, Not Hispanic	65	73	61	70	43	49
Male, Black, Not Hispanic	32	35	27	26	18	18
Female, Black, Not Hispanic	34	38	30	30	20	20
Male, Hispanic	54	50	43	43	29	29
Female, Hispanic	47	52	40	47	26	32

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.38D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.44	0.43	0.40	0.42	0.29	0.33
GENDER						
Male	0.58	0.63	0.56	0.56	0.41	0.42
Female	0.61	0.63	0.58	0.61	0.43	0.47
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.49	0.47	0.45	0.46	0.33	0.37
White	0.56	0.58	0.52	0.57	0.40	0.47
Black or African American	1.08	1.17	0.97	0.97	0.71	0.71
American Indian or Alaska Native	3.99	4.68	3.90	5.06	2.39	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	2.90	*
Asian	2.21	1.89	2.25	1.57	1.48	0.97
Two or More Races	2.19	2.05	1.79	1.92	1.10	1.35
Hispanic or Latino	1.02	1.01	0.91	0.97	0.66	0.71
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.77	0.81	0.73	0.76	0.55	0.62
Female, White, Not Hispanic	0.76	0.86	0.75	0.85	0.59	0.67
Male, Black, Not Hispanic	1.49	1.63	1.36	1.30	0.98	0.93
Female, Black, Not Hispanic	1.50	1.68	1.47	1.43	1.08	1.08
Male, Hispanic	1.45	1.50	1.25	1.34	0.93	0.96
Female, Hispanic	1.46	1.46	1.30	1.37	0.93	1.05

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.38P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0585		0.2985		0.9818	
GENDER						
Male	0.0144		0.0663		0.4485	
Female	0.8298		0.7373		0.4865	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0495		0.1232		0.7206	
White	0.2840		0.6245		0.6694	
Black or African American	0.0864		0.0359		0.2540	
American Indian or Alaska Native	0.9145		0.7450		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4275		0.1979		0.0488	
Two or More Races	0.6840		0.8829		0.6706	
Hispanic or Latino	0.7049		0.5684		0.5537	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.1675		0.2506		0.9213	
Female, White, Not Hispanic	0.8934		0.6669		0.6088	
Male, Black, Not Hispanic	0.1651		0.0946		0.1809	
Female, Black, Not Hispanic	0.2854		0.1998		0.7602	
Male, Hispanic	0.2071		0.5702		0.7901	
Female, Hispanic	0.4337		0.1664		0.2742	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.39C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	136	141	158	165	183	197
GENDER						
Male	93	98	105	115	121	130
Female	86	94	99	111	117	129
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	116	119	132	138	154	164
White	325	313	314	307	277	274
Black or African American	142	150	133	141	112	113
American Indian or Alaska Native	21	20	18	18	16	16
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	90	104	86	99	60	81
Two or More Races	44	42	40	41	34	38
Hispanic or Latino	62	75	72	84	79	92
EDUCATION						
< High School	120	116	111	105	86	91
High School Graduate	200	208	186	199	149	171
Some College	246	248	238	238	210	213
College Graduate	179	163	176	162	168	149
CURRENT EMPLOYMENT						
Full-Time	231	236	219	232	200	210
Part-Time	205	200	199	195	171	173
Unemployed	107	103	100	99	85	82
Other ¹	174	177	163	165	135	142

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.39D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.40	0.45	0.47	0.53	0.56
GENDER						
Male	0.53	0.56	0.60	0.66	0.69	0.74
Female	0.49	0.54	0.57	0.64	0.68	0.74
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.42	0.43	0.48	0.50	0.56	0.60
White	0.43	0.45	0.51	0.54	0.62	0.65
Black or African American	0.98	1.14	1.07	1.26	1.22	1.38
American Indian or Alaska Native	3.96	3.88	4.40	4.63	4.96	4.97
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.56	2.21	2.69	2.33	2.27	2.58
Two or More Races	1.71	2.17	2.07	2.45	3.17	2.85
Hispanic or Latino	0.85	1.01	0.99	1.14	1.08	1.25
EDUCATION						
< High School	0.96	1.16	1.06	1.22	1.09	1.34
High School Graduate	0.68	0.77	0.73	0.85	0.77	0.92
Some College	0.56	0.60	0.65	0.68	0.79	0.81
College Graduate	0.57	0.56	0.72	0.72	1.01	1.07
CURRENT EMPLOYMENT						
Full-Time	0.48	0.46	0.58	0.57	0.75	0.74
Part-Time	0.68	0.68	0.77	0.80	0.90	0.98
Unemployed	0.99	1.31	1.14	1.45	1.32	1.53
Other ¹	1.00	1.09	1.07	1.15	1.08	1.22

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.39P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.4537		0.6344		0.9625	
GENDER						
Male	0.3259		0.3572		0.4838	
Female	0.9108		0.8131		0.5205	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.8086		0.8709		0.9565	
White	0.8317		0.6724		0.8799	
Black or African American	0.6682		0.8411		0.3245	
American Indian or Alaska Native	0.9396		0.8447		0.7066	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1338		0.1157		0.0938	
Two or More Races	0.0249		0.1457		0.3141	
Hispanic or Latino	0.2855		0.5145		0.9560	
EDUCATION						
< High School	0.1995		0.0649		0.4446	
High School Graduate	0.2252		0.6595		0.7881	
Some College	0.8497		0.6077		0.7989	
College Graduate	0.9217		0.8204		0.0191	
CURRENT EMPLOYMENT						
Full-Time	0.7035		0.5194		0.6343	
Part-Time	0.6680		0.7495		0.1928	
Unemployed	0.0168		0.0381		0.4728	
Other ¹	0.5507		0.4950		0.7657	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.40C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	736	553	954	764	1,021	802
GENDER						
Male	428	289	629	471	669	524
Female	571	427	673	556	701	581
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	646	471	844	694	935	736
White	1,803	1,505	1,612	1,317	1,435	1,175
Black or African American	689	558	554	438	460	374
American Indian or Alaska Native	102	102	74	77	59	66
Native Hawaiian or Other Pacific Islander	*	89	*	*	*	*
Asian	461	400	415	342	345	268
Two or More Races	221	154	176	142	171	122
Hispanic or Latino	312	239	376	293	377	302
EDUCATION						
< High School	604	514	470	406	411	331
High School Graduate	945	761	788	647	669	570
Some College	895	740	808	635	713	570
College Graduate	1,321	1,032	1,189	938	1,094	847
CURRENT EMPLOYMENT						
Full-Time	1,480	1,166	1,289	1,066	1,173	971
Part-Time	600	469	534	428	506	391
Unemployed	319	236	286	213	255	188
Other ¹	1,045	958	898	774	790	698

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.40D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.36	0.27	0.47	0.37	0.50	0.39
GENDER						
Male	0.44	0.29	0.65	0.48	0.69	0.53
Female	0.54	0.40	0.64	0.52	0.66	0.54
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.37	0.27	0.48	0.39	0.54	0.42
White	0.36	0.24	0.52	0.42	0.59	0.46
Black or African American	1.17	0.88	1.42	1.03	1.41	1.08
American Indian or Alaska Native	2.84	2.57	4.78	4.42	4.58	4.26
Native Hawaiian or Other Pacific Islander	*	5.04	*	*	*	*
Asian	2.64	1.82	2.65	1.83	2.61	1.79
Two or More Races	2.19	1.49	3.85	2.58	3.98	2.73
Hispanic or Latino	1.10	0.81	1.32	0.99	1.33	1.02
EDUCATION						
< High School	1.34	0.95	1.35	1.08	1.28	0.99
High School Graduate	0.65	0.51	0.87	0.67	0.86	0.68
Some College	0.56	0.41	0.82	0.62	0.89	0.67
College Graduate	0.55	0.37	0.70	0.55	0.82	0.63
CURRENT EMPLOYMENT						
Full-Time	0.37	0.28	0.53	0.43	0.61	0.48
Part-Time	0.91	0.64	1.22	0.95	1.36	1.06
Unemployed	1.46	1.22	2.18	1.76	2.31	1.89
Other ¹	0.78	0.58	0.93	0.75	0.93	0.76

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.40P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0301		0.4515		0.3269	
GENDER						
Male	0.2345		0.9561		0.7170	
Female	0.0523		0.3158		0.2897	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0447		0.8750		0.4813	
White	0.1279		0.6729		0.9531	
Black or African American	0.2376		0.6955		0.4475	
American Indian or Alaska Native	0.9822		0.7861		0.3580	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.2312		0.3441		0.1481	
Two or More Races	0.8129		0.1367		0.7323	
Hispanic or Latino	0.2496		0.0939		0.3023	
EDUCATION						
< High School	0.1859		0.7970		0.7306	
High School Graduate	0.0276		0.1210		0.0373	
Some College	0.6372		0.5615		0.8567	
College Graduate	0.7009		0.9495		0.8827	
CURRENT EMPLOYMENT						
Full-Time	0.8948		0.7213		0.5793	
Part-Time	0.6901		0.5113		0.3930	
Unemployed	0.6069		0.6221		0.9160	
Other ¹	0.0043		0.3936		0.1432	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.41C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	760	584	984	800	1,052	843
GENDER						
Male	442	310	641	486	687	540
Female	577	445	688	584	717	611
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	664	492	868	718	960	772
White	1,876	1,595	1,688	1,399	1,498	1,241
Black or African American	738	617	601	495	495	410
American Indian or Alaska Native	105	107	76	82	64	70
Native Hawaiian or Other Pacific Islander	*	91	*	73	51	61
Asian	484	422	437	366	357	282
Two or More Races	224	159	180	150	174	130
Hispanic or Latino	325	260	389	318	387	327
EDUCATION						
< High School	625	542	492	426	427	349
High School Graduate	978	804	813	692	687	605
Some College	940	785	856	684	750	615
College Graduate	1,356	1,059	1,223	965	1,122	874
CURRENT EMPLOYMENT						
Full-Time	1,529	1,208	1,334	1,111	1,210	1,003
Part-Time	641	518	579	475	539	432
Unemployed	340	263	306	240	275	210
Other ¹	1,053	968	914	788	801	703

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.41D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.32	0.24	0.41	0.33	0.44	0.35
GENDER						
Male	0.39	0.27	0.56	0.42	0.60	0.47
Female	0.47	0.36	0.56	0.47	0.58	0.49
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.33	0.24	0.43	0.35	0.48	0.38
White	0.32	0.22	0.47	0.38	0.53	0.42
Black or African American	0.97	0.75	1.19	0.88	1.19	0.92
American Indian or Alaska Native	2.56	2.28	3.99	3.78	3.84	3.68
Native Hawaiian or Other Pacific Islander	*	4.19	*	5.46	5.65	5.32
Asian	2.28	1.57	2.29	1.59	2.24	1.54
Two or More Races	1.78	1.26	3.13	2.14	3.25	2.28
Hispanic or Latino	0.91	0.71	1.09	0.86	1.08	0.89
EDUCATION						
< High School	1.13	0.84	1.15	0.94	1.08	0.87
High School Graduate	0.55	0.45	0.74	0.58	0.73	0.59
Some College	0.47	0.36	0.67	0.53	0.74	0.57
College Graduate	0.51	0.35	0.65	0.51	0.76	0.59
CURRENT EMPLOYMENT						
Full-Time	0.34	0.26	0.48	0.39	0.55	0.44
Part-Time	0.68	0.51	0.90	0.72	1.00	0.82
Unemployed	1.03	0.93	1.47	1.23	1.59	1.32
Other ¹	0.71	0.53	0.85	0.69	0.85	0.69

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.41P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0520		0.5307		0.3529	
GENDER						
Male	0.3608		0.8969		0.8328	
Female	0.0602		0.3192		0.2573	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0568		0.9097		0.5064	
White	0.1453		0.6219		0.9815	
Black or African American	0.2848		0.7404		0.6052	
American Indian or Alaska Native	0.9246		0.8023		0.3518	
Native Hawaiian or Other Pacific Islander	*		*		0.9082	
Asian	0.1427		0.2106		0.0773	
Two or More Races	0.6947		0.2312		0.5707	
Hispanic or Latino	0.4012		0.1523		0.3235	
EDUCATION						
< High School	0.2928		0.7852		0.5481	
High School Graduate	0.0749		0.1509		0.0370	
Some College	0.6790		0.4872		0.8012	
College Graduate	0.6973		0.9069		0.9639	
CURRENT EMPLOYMENT						
Full-Time	0.9603		0.8181		0.6532	
Part-Time	0.6088		0.5865		0.6683	
Unemployed	0.1288		0.8098		0.8745	
Other ¹	0.0064		0.4710		0.1392	

*Low precision; no estimate reported.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.42C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	1,076	869	804	690	438	368
AGE						
12-17	73	81	54	60	23	24
18-25	183	197	182	197	107	125
26 or Older	1,021	802	766	623	420	334
GENDER						
Male	693	553	607	519	367	311
Female	736	620	463	380	232	165
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	979	789	737	625	404	341
White	848	663	652	536	370	311
Black or African American	328	261	270	222	138	116
American Indian or Alaska Native	65	72	54	62	22	29
Native Hawaiian or Other Pacific Islander	51	61	39	31	27	19
Asian	360	285	177	155	53	45
Two or More Races	174	130	80	79	57	31
Hispanic or Latino	395	332	343	292	164	137

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.42D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.41	0.33	0.31	0.26	0.17	0.14
AGE						
12-17	0.29	0.33	0.22	0.24	0.09	0.10
18-25	0.53	0.56	0.52	0.57	0.31	0.36
26 or Older	0.50	0.39	0.38	0.30	0.21	0.16
GENDER						
Male	0.55	0.43	0.48	0.40	0.29	0.24
Female	0.54	0.45	0.34	0.28	0.17	0.12
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.44	0.35	0.33	0.28	0.18	0.15
White	0.50	0.39	0.38	0.31	0.22	0.18
Black or African American	1.05	0.83	0.86	0.70	0.44	0.36
American Indian or Alaska Native	3.45	3.47	3.01	3.12	1.46	1.79
Native Hawaiian or Other Pacific Islander	4.95	4.85	4.11	2.98	2.97	1.82
Asian	2.06	1.41	1.28	1.09	0.39	0.32
Two or More Races	2.75	1.93	1.82	1.64	1.31	0.67
Hispanic or Latino	0.96	0.78	0.83	0.69	0.40	0.32

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.42P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.3203		0.8131		0.5200	
AGE						
12-17	0.9818		0.5992		0.3006	
18-25	0.9625		0.8263		0.2930	
26 or Older	0.3269		0.7311		0.7608	
GENDER						
Male	0.8148		0.8169		0.5640	
Female	0.2246		0.4647		0.7538	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4780		0.9967		0.3654	
White	0.8944		0.3594		0.4184	
Black or African American	0.6678		0.1675		0.9632	
American Indian or Alaska Native	0.3005		0.3324		0.1478	
Native Hawaiian or Other Pacific Islander	0.9447		0.2066		0.2217	
Asian	0.0922		0.2125		0.9789	
Two or More Races	0.5338		0.5867		0.2583	
Hispanic or Latino	0.2763		0.5933		0.5702	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.43C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	73	81	54	60	23	24
GENDER						
Male	52	54	40	43	18	18
Female	52	57	37	41	16	16
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	64	70	47	52	22	21
White	62	70	46	50	20	20
Black or African American	27	27	16	17	7	3
American Indian or Alaska Native	4	*	3	*	2	*
Native Hawaiian or Other Pacific Islander	3	*	2	*	0	*
Asian	20	12	10	6	3	0
Two or More Races	8	10	6	7	3	2
Hispanic or Latino	37	40	29	29	10	12
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	42	45	34	36	14	15
Female, White, Not Hispanic	43	49	29	32	14	12
Male, Black, Not Hispanic	18	18	13	12	7	1
Female, Black, Not Hispanic	20	20	12	12	0	3
Male, Hispanic	29	29	23	22	8	9
Female, Hispanic	26	32	18	22	6	9

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.43D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.29	0.33	0.22	0.24	0.09	0.10
GENDER						
Male	0.41	0.42	0.32	0.34	0.14	0.14
Female	0.43	0.47	0.30	0.33	0.13	0.13
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.33	0.37	0.24	0.27	0.11	0.11
White	0.40	0.47	0.31	0.35	0.15	0.15
Black or African American	0.71	0.71	0.45	0.48	0.20	0.09
American Indian or Alaska Native	2.39	*	1.59	*	1.06	*
Native Hawaiian or Other Pacific Islander	2.90	*	1.76	*	0.17	*
Asian	1.48	0.97	0.75	0.52	0.20	0.02
Two or More Races	1.10	1.35	0.78	0.95	0.43	0.23
Hispanic or Latino	0.66	0.71	0.52	0.52	0.17	0.21
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.55	0.62	0.45	0.51	0.21	0.22
Female, White, Not Hispanic	0.59	0.67	0.42	0.46	0.21	0.18
Male, Black, Not Hispanic	0.98	0.93	0.72	0.66	0.40	0.08
Female, Black, Not Hispanic	1.08	1.08	0.69	0.68	0.02	0.18
Male, Hispanic	0.93	0.96	0.78	0.75	0.27	0.32
Female, Hispanic	0.93	1.05	0.67	0.79	0.22	0.34

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.43P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.9818		0.5992		0.3006	
GENDER						
Male	0.4485		0.6050		0.3055	
Female	0.4865		0.8255		0.6926	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7206		0.5425		0.1474	
White	0.6694		0.6675		0.2740	
Black or African American	0.2540		0.6833		0.3295	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0488		0.1657		0.2961	
Two or More Races	0.6706		0.7409		0.1741	
Hispanic or Latino	0.5537		1.0000		0.5637	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.9213		0.8720		0.3338	
Female, White, Not Hispanic	0.6088		0.6323		0.5621	
Male, Black, Not Hispanic	0.1809		0.5544		0.0737	
Female, Black, Not Hispanic	0.7602		0.9650		0.0825	
Male, Hispanic	0.7901		0.7296		0.7171	
Female, Hispanic	0.2742		0.7087		0.6801	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.44C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	183	197	182	197	107	125
GENDER						
Male	121	130	121	133	85	95
Female	117	129	112	126	65	66
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	154	164	155	174	97	111
White	277	274	212	213	107	113
Black or African American	112	113	74	82	28	33
American Indian or Alaska Native	16	16	13	12	5	6
Native Hawaiian or Other Pacific Islander	*	*	*	*	6	*
Asian	60	81	39	52	16	21
Two or More Races	34	38	26	28	12	17
Hispanic or Latino	79	92	80	89	45	50
EDUCATION						
< High School	86	91	73	73	38	41
High School Graduate	149	171	119	132	59	64
Some College	210	213	157	167	77	83
College Graduate	168	149	117	118	52	63
CURRENT EMPLOYMENT						
Full-Time	200	210	157	163	78	86
Part-Time	171	173	129	133	63	62
Unemployed	85	82	66	69	35	36
Other ¹	135	142	98	110	45	54

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.44D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.53	0.56	0.52	0.57	0.31	0.36
GENDER						
Male	0.69	0.74	0.69	0.76	0.49	0.54
Female	0.68	0.74	0.64	0.72	0.38	0.38
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.56	0.60	0.56	0.63	0.35	0.40
White	0.62	0.65	0.65	0.70	0.44	0.51
Black or African American	1.22	1.38	1.02	1.30	0.51	0.62
American Indian or Alaska Native	4.96	4.97	4.69	4.66	2.02	2.74
Native Hawaiian or Other Pacific Islander	*	*	*	*	3.01	*
Asian	2.27	2.58	1.92	2.03	0.86	1.02
Two or More Races	3.17	2.85	2.93	2.72	1.60	1.99
Hispanic or Latino	1.08	1.25	1.11	1.21	0.62	0.67
EDUCATION						
< High School	1.09	1.34	1.02	1.21	0.62	0.77
High School Graduate	0.77	0.92	0.78	0.84	0.45	0.50
Some College	0.79	0.81	0.79	0.90	0.54	0.58
College Graduate	1.01	1.07	1.28	1.39	0.83	0.99
CURRENT EMPLOYMENT						
Full-Time	0.75	0.74	0.80	0.77	0.50	0.55
Part-Time	0.90	0.98	0.91	1.02	0.57	0.59
Unemployed	1.32	1.53	1.26	1.47	0.80	0.86
Other ¹	1.08	1.22	0.97	1.11	0.54	0.67

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.44P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.9625		0.8263		0.2930	
GENDER						
Male	0.4838		0.4702		0.1759	
Female	0.5205		0.6690		1.0000	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.9565		0.8716		0.2398	
White	0.8799		0.8185		0.1055	
Black or African American	0.3245		0.7993		0.6285	
American Indian or Alaska Native	0.7066		0.7725		0.4025	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0938		0.4374		0.8966	
Two or More Races	0.3141		0.5609		0.1464	
Hispanic or Latino	0.9560		0.8818		0.9602	
EDUCATION						
< High School	0.4446		0.5532		0.6855	
High School Graduate	0.7881		0.3957		0.9347	
Some College	0.7989		0.1957		0.1352	
College Graduate	0.0191		0.6130		0.1668	
CURRENT EMPLOYMENT						
Full-Time	0.6343		0.4595		0.2496	
Part-Time	0.1928		0.0741		0.1681	
Unemployed	0.4728		0.5741		0.6108	
Other ¹	0.7657		0.6173		0.9529	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.45C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	1,021	802	766	623	420	334
GENDER						
Male	669	524	583	490	350	292
Female	701	581	435	338	219	151
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	935	736	700	567	388	314
White	1,435	1,175	751	563	371	290
Black or African American	460	374	292	258	139	116
American Indian or Alaska Native	59	66	50	59	21	28
Native Hawaiian or Other Pacific Islander	*	*	*	29	*	18
Asian	345	268	171	144	50	40
Two or More Races	171	122	75	72	55	26
Hispanic or Latino	377	302	330	268	154	125
EDUCATION						
< High School	411	331	311	257	132	126
High School Graduate	669	570	480	376	258	197
Some College	713	570	436	342	220	163
College Graduate	1,094	847	518	369	239	187
CURRENT EMPLOYMENT						
Full-Time	1,173	971	726	563	334	267
Part-Time	506	391	279	230	161	112
Unemployed	255	188	173	130	104	69
Other ¹	790	698	371	319	186	162

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.45D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.50	0.39	0.38	0.30	0.21	0.16
GENDER						
Male	0.69	0.53	0.60	0.50	0.36	0.30
Female	0.66	0.54	0.41	0.32	0.21	0.14
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.54	0.42	0.40	0.32	0.22	0.18
White	0.59	0.46	0.45	0.36	0.26	0.21
Black or African American	1.41	1.08	1.16	0.92	0.60	0.48
American Indian or Alaska Native	4.58	4.26	4.08	3.98	2.00	2.31
Native Hawaiian or Other Pacific Islander	*	*	*	3.85	*	2.46
Asian	2.61	1.79	1.62	1.34	0.49	0.38
Two or More Races	3.98	2.73	2.65	2.33	1.94	0.90
Hispanic or Latino	1.33	1.02	1.16	0.91	0.54	0.42
EDUCATION						
< High School	1.28	0.99	1.07	0.84	0.49	0.44
High School Graduate	0.86	0.68	0.71	0.57	0.44	0.34
Some College	0.89	0.67	0.75	0.55	0.42	0.30
College Graduate	0.82	0.63	0.66	0.50	0.34	0.27
CURRENT EMPLOYMENT						
Full-Time	0.61	0.48	0.57	0.42	0.30	0.23
Part-Time	1.36	1.06	1.05	0.87	0.65	0.46
Unemployed	2.31	1.89	1.96	1.57	1.29	0.93
Other ¹	0.93	0.76	0.54	0.45	0.28	0.24

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.45P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.3269		0.7311		0.7608	
GENDER						
Male	0.7170		0.9501		0.8500	
Female	0.2897		0.4855		0.7685	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4813		0.9424		0.5960	
White	0.9531		0.3802		0.7498	
Black or African American	0.4475		0.1653		0.9830	
American Indian or Alaska Native	0.3580		0.3981		0.2185	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1481		0.2664		0.9322	
Two or More Races	0.7323		0.5061		0.1080	
Hispanic or Latino	0.3023		0.5540		0.5804	
EDUCATION						
< High School	0.7306		0.9189		0.1794	
High School Graduate	0.0373		0.2125		0.7373	
Some College	0.8567		0.4091		0.8942	
College Graduate	0.8827		0.1902		0.2841	
CURRENT EMPLOYMENT						
Full-Time	0.5793		0.7419		0.3113	
Part-Time	0.3930		0.4906		0.8547	
Unemployed	0.9160		0.8597		0.3197	
Other ¹	0.1432		0.5256		0.1899	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.46C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	1,052	843	799	686	438	366
GENDER						
Male	687	540	606	521	367	311
Female	717	611	461	376	231	165
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	960	772	733	624	404	339
White	1,498	1,241	801	627	389	316
Black or African American	495	410	307	274	140	121
American Indian or Alaska Native	64	70	53	62	22	28
Native Hawaiian or Other Pacific Islander	51	61	39	31	27	19
Asian	357	282	177	156	53	45
Two or More Races	174	130	79	79	57	31
Hispanic or Latino	387	327	340	290	162	136
EDUCATION						
< High School	427	349	323	273	137	133
High School Graduate	687	605	498	409	265	206
Some College	750	615	461	383	231	180
College Graduate	1,122	874	542	402	247	199
CURRENT EMPLOYMENT						
Full-Time	1,210	1,003	758	597	346	286
Part-Time	539	432	306	271	172	129
Unemployed	275	210	189	153	112	78
Other ¹	801	703	383	335	192	171

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.46D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.44	0.35	0.34	0.29	0.18	0.15
GENDER						
Male	0.60	0.47	0.53	0.45	0.32	0.27
Female	0.58	0.49	0.37	0.30	0.19	0.13
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.48	0.38	0.36	0.31	0.20	0.17
White	0.53	0.42	0.42	0.34	0.24	0.20
Black or African American	1.19	0.92	0.98	0.78	0.50	0.41
American Indian or Alaska Native	3.84	3.68	3.35	3.40	1.66	1.98
Native Hawaiian or Other Pacific Islander	5.65	5.32	4.71	3.37	3.36	2.02
Asian	2.24	1.54	1.40	1.19	0.43	0.36
Two or More Races	3.25	2.28	2.28	1.98	1.61	0.81
Hispanic or Latino	1.08	0.89	0.95	0.79	0.45	0.37
EDUCATION						
< High School	1.08	0.87	0.90	0.76	0.42	0.40
High School Graduate	0.73	0.59	0.61	0.50	0.37	0.29
Some College	0.74	0.57	0.63	0.50	0.35	0.27
College Graduate	0.76	0.59	0.63	0.50	0.32	0.27
CURRENT EMPLOYMENT						
Full-Time	0.55	0.44	0.52	0.39	0.27	0.22
Part-Time	1.00	0.82	0.79	0.70	0.49	0.37
Unemployed	1.59	1.32	1.36	1.15	0.90	0.67
Other ¹	0.85	0.69	0.50	0.43	0.26	0.23

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.46P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.3529		0.8183		0.5408	
GENDER						
Male	0.8328		0.8071		0.5813	
Female	0.2573		0.4634		0.7684	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.5064		0.9971		0.3950	
White	0.9815		0.3561		0.4419	
Black or African American	0.6052		0.1608		0.9307	
American Indian or Alaska Native	0.3518		0.3883		0.1689	
Native Hawaiian or Other Pacific Islander	0.9082		0.1995		0.2128	
Asian	0.0773		0.1994		0.9946	
Two or More Races	0.5707		0.6247		0.2927	
Hispanic or Latino	0.3235		0.6165		0.6065	
EDUCATION						
< High School	0.5481		0.8670		0.1892	
High School Graduate	0.0370		0.1509		0.7686	
Some College	0.8012		0.7214		0.6759	
College Graduate	0.9639		0.2535		0.2211	
CURRENT EMPLOYMENT						
Full-Time	0.6532		0.9096		0.2245	
Part-Time	0.6683		0.9844		0.7249	
Unemployed	0.8745		0.9316		0.4717	
Other ¹	0.1392		0.6611		0.2155	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.47C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	942	826	875	778	835	736
GEOGRAPHIC DIVISION						
Northeast	373	343	340	345	332	327
New England	174	141	157	156	135	146
Middle Atlantic	330	311	303	308	303	291
Midwest	317	306	360	361	355	339
East North Central	261	262	289	299	281	285
West North Central	182	171	212	188	210	181
South	607	478	599	491	561	473
South Atlantic	480	347	472	373	440	353
East South Central	202	180	218	201	209	189
West South Central	335	267	287	244	268	246
West	508	457	433	364	417	335
Mountain	230	177	215	173	208	162
Pacific	449	415	374	315	365	292
COUNTY TYPE						
Large Metro	1,570	1,288	836	749	750	661
Small Metro	1,283	1,022	686	563	596	511
250K – 1 Mil. Pop.	1,069	835	521	445	454	404
< 250K Pop.	828	646	474	345	412	315
Nonmetro	901	859	509	482	459	441
Urbanized	617	623	339	334	310	302
Less Urbanized	703	652	388	361	344	335
Completely Rural	336	289	205	165	180	150

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.47D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.36	0.31	0.33	0.29	0.32	0.28
GEOGRAPHIC DIVISION						
Northeast	0.79	0.72	0.72	0.72	0.70	0.69
New England	1.39	1.12	1.26	1.24	1.08	1.16
Middle Atlantic	0.95	0.89	0.87	0.88	0.87	0.83
Midwest	0.56	0.54	0.64	0.64	0.63	0.60
East North Central	0.67	0.67	0.74	0.76	0.72	0.73
West North Central	1.06	0.98	1.23	1.09	1.22	1.04
South	0.62	0.48	0.61	0.50	0.58	0.48
South Atlantic	0.93	0.67	0.92	0.71	0.86	0.68
East South Central	1.30	1.15	1.41	1.29	1.35	1.21
West South Central	1.10	0.86	0.94	0.78	0.88	0.79
West	0.83	0.73	0.71	0.59	0.68	0.54
Mountain	1.23	0.93	1.15	0.91	1.11	0.85
Pacific	1.05	0.96	0.88	0.73	0.86	0.68
COUNTY TYPE						
Large Metro	0.54	0.46	0.46	0.40	0.45	0.37
Small Metro	0.58	0.48	0.62	0.54	0.57	0.51
250K – 1 Mil. Pop.	0.73	0.59	0.69	0.64	0.66	0.61
< 250K Pop.	0.91	0.86	1.15	0.99	1.06	0.94
Nonmetro	0.77	0.61	0.84	0.69	0.79	0.65
Urbanized	1.21	1.02	1.22	1.04	1.16	1.02
Less Urbanized	1.18	0.87	1.26	0.98	1.19	0.95
Completely Rural	2.10	1.81	2.59	2.46	2.45	2.29

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.47P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1860		0.6427		0.5459	
GEOGRAPHIC DIVISION						
Northeast	0.6176		0.6143		0.7521	
New England	0.6589		0.8406		0.6662	
Middle Atlantic	0.4373		0.6431		0.8830	
Midwest	0.7969		0.2312		0.3078	
East North Central	0.8661		0.5981		0.8288	
West North Central	0.8374		0.1680		0.1352	
South	0.2513		0.8423		0.7412	
South Atlantic	0.5744		0.5394		0.5585	
East South Central	0.9246		0.7161		0.9950	
West South Central	0.1799		0.4001		0.8256	
West	0.5793		0.5769		0.3066	
Mountain	0.2429		0.4124		0.4391	
Pacific	0.9464		0.8535		0.4615	
COUNTY TYPE						
Large Metro	0.7734		0.7085		0.9654	
Small Metro	0.0717		0.5339		0.5308	
250K – 1 Mil. Pop.	0.1378		0.8214		0.8249	
< 250K Pop.	0.3085		0.1797		0.4512	
Nonmetro	0.8808		0.5787		0.8316	
Urbanized	0.5500		0.8397		0.9333	
Less Urbanized	0.3372		0.4296		0.7526	
Completely Rural	0.8378		0.5395		0.9420	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.48C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	92	102	80	84	59	61
GEOGRAPHIC DIVISION						
Northeast	31	34	27	29	22	21
New England	16	15	13	14	10	10
Middle Atlantic	27	29	24	25	20	18
Midwest	37	47	30	39	24	29
East North Central	30	40	24	32	19	24
West North Central	21	23	17	20	14	16
South	60	61	53	51	38	39
South Atlantic	45	41	41	34	28	25
East South Central	20	25	19	20	15	17
West South Central	34	40	29	33	23	25
West	49	55	42	48	31	30
Mountain	23	25	18	22	15	17
Pacific	42	48	37	42	27	25
COUNTY TYPE						
Large Metro	77	84	63	66	46	46
Small Metro	67	62	54	51	35	33
250K – 1 Mil. Pop.	55	47	46	38	29	25
< 250K Pop.	37	40	27	34	19	22
Nonmetro	51	49	40	37	29	29
Urbanized	28	30	23	23	17	16
Less Urbanized	39	39	29	28	21	22
Completely Rural	23	13	19	12	13	11

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.48D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.37	0.41	0.32	0.34	0.24	0.25
GEOGRAPHIC DIVISION						
Northeast	0.75	0.81	0.65	0.69	0.53	0.51
New England	1.46	1.39	1.21	1.25	0.96	0.91
Middle Atlantic	0.88	0.96	0.77	0.81	0.64	0.60
Midwest	0.68	0.88	0.56	0.72	0.44	0.54
East North Central	0.80	1.08	0.65	0.86	0.50	0.63
West North Central	1.28	1.39	1.05	1.20	0.84	0.98
South	0.64	0.65	0.57	0.54	0.41	0.42
South Atlantic	0.97	0.88	0.89	0.73	0.59	0.54
East South Central	1.35	1.68	1.26	1.35	1.01	1.16
West South Central	1.06	1.22	0.89	1.00	0.70	0.77
West	0.82	0.92	0.70	0.80	0.52	0.50
Mountain	1.22	1.31	0.97	1.15	0.78	0.91
Pacific	1.04	1.20	0.90	1.04	0.65	0.61
COUNTY TYPE						
Large Metro	0.47	0.52	0.41	0.44	0.32	0.31
Small Metro	0.68	0.73	0.58	0.63	0.41	0.44
250K – 1 Mil. Pop.	0.80	0.81	0.72	0.67	0.49	0.49
< 250K Pop.	1.22	1.57	0.95	1.42	0.75	0.93
Nonmetro	1.00	1.06	0.84	0.85	0.65	0.70
Urbanized	1.32	1.61	1.14	1.31	0.93	0.97
Less Urbanized	1.55	1.56	1.31	1.24	1.01	1.03
Completely Rural	2.82	2.80	2.63	2.59	1.93	2.59

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.48P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0170		0.0108		0.0151	
GEOGRAPHIC DIVISION						
Northeast	0.0281		0.0383		0.0030	
New England	0.9680		0.3919		0.6765	
Middle Atlantic	0.0120		0.0042		0.0015	
Midwest	0.2268		0.0772		0.5096	
East North Central	0.3759		0.1189		0.2874	
West North Central	0.3628		0.3502		0.7313	
South	0.2178		0.2787		0.3650	
South Atlantic	0.0109		0.0884		0.2408	
East South Central	0.4673		0.8221		0.9463	
West South Central	0.5881		0.6726		0.9153	
West	0.5041		0.4434		0.3277	
Mountain	0.0679		0.2481		0.6004	
Pacific	0.8563		0.7684		0.1337	
COUNTY TYPE						
Large Metro	0.1014		0.0307		0.0155	
Small Metro	0.4669		0.8571		0.7206	
250K – 1 Mil. Pop.	0.7257		0.6161		0.4576	
< 250K Pop.	0.4686		0.7266		0.7103	
Nonmetro	0.1111		0.0563		0.4156	
Urbanized	0.4337		0.1364		0.1261	
Less Urbanized	0.3122		0.2198		0.7937	
Completely Rural	0.1256		0.7243		0.7753	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.49C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	165	189	178	194	170	189
GEOGRAPHIC DIVISION						
Northeast	66	79	70	76	68	71
New England	30	41	35	33	38	37
Middle Atlantic	58	66	61	67	56	60
Midwest	57	74	61	81	63	78
East North Central	47	64	51	65	53	64
West North Central	31	35	35	45	33	47
South	106	110	107	117	100	118
South Atlantic	79	72	82	70	76	75
East South Central	35	41	35	46	36	49
West South Central	55	71	56	74	53	72
West	94	100	97	100	93	93
Mountain	38	42	41	43	39	44
Pacific	86	88	88	90	84	82
COUNTY TYPE						
Large Metro	283	264	211	217	176	187
Small Metro	218	205	178	159	141	136
250K – 1 Mil. Pop.	159	156	130	123	105	104
< 250K Pop.	161	130	129	100	98	87
Nonmetro	144	152	119	119	101	101
Urbanized	112	115	94	87	75	71
Less Urbanized	97	99	78	84	71	73
Completely Rural	45	33	34	26	26	21

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.49D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.48	0.54	0.51	0.55	0.49	0.54
GEOGRAPHIC DIVISION						
Northeast	1.07	1.29	1.14	1.24	1.10	1.15
New England	1.83	2.52	2.14	2.01	2.31	2.28
Middle Atlantic	1.29	1.47	1.35	1.50	1.25	1.33
Midwest	0.77	1.00	0.83	1.09	0.85	1.05
East North Central	0.91	1.25	0.99	1.27	1.04	1.26
West North Central	1.37	1.52	1.53	1.94	1.45	2.03
South	0.82	0.85	0.83	0.90	0.78	0.91
South Atlantic	1.20	1.09	1.25	1.07	1.16	1.14
East South Central	1.73	2.00	1.70	2.24	1.76	2.41
West South Central	1.30	1.65	1.31	1.72	1.25	1.68
West	1.13	1.18	1.16	1.19	1.11	1.10
Mountain	1.52	1.66	1.61	1.68	1.53	1.71
Pacific	1.47	1.51	1.50	1.53	1.44	1.40
COUNTY TYPE						
Large Metro	0.71	0.80	0.74	0.81	0.71	0.76
Small Metro	0.83	0.86	0.88	0.87	0.80	0.88
250K – 1 Mil. Pop.	1.03	1.07	1.06	1.03	0.99	1.04
< 250K Pop.	1.40	1.38	1.50	1.54	1.39	1.53
Nonmetro	1.04	1.09	1.05	1.24	1.06	1.25
Urbanized	1.52	1.67	1.51	1.97	1.50	1.91
Less Urbanized	1.40	1.44	1.43	1.67	1.51	1.72
Completely Rural	4.19	4.02	4.19	3.24	3.83	3.70

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.49P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0703		0.0268		0.0067	
GEOGRAPHIC DIVISION						
Northeast	0.5531		0.3554		0.1965	
New England	0.7660		0.9627		0.6611	
Middle Atlantic	0.5987		0.3013		0.0674	
Midwest	0.5467		0.7632		0.3608	
East North Central	0.5604		0.7669		0.9646	
West North Central	0.8304		0.9097		0.0961	
South	0.2557		0.3813		0.5729	
South Atlantic	0.0998		0.1286		0.1130	
East South Central	0.5897		0.5476		0.6105	
West South Central	0.7621		0.9301		0.5646	
West	0.0442		0.0228		0.0034	
Mountain	0.1148		0.0427		0.0097	
Pacific	0.1293		0.1143		0.0448	
COUNTY TYPE						
Large Metro	0.1337		0.0653		0.0406	
Small Metro	0.6792		0.9296		0.6892	
250K – 1 Mil. Pop.	0.2366		0.7585		0.3994	
< 250K Pop.	0.3531		0.7916		0.6656	
Nonmetro	0.3360		0.0462		0.0102	
Urbanized	0.7445		0.0077		0.0108	
Less Urbanized	0.1437		0.5991		0.0769	
Completely Rural	0.8131		0.5121		0.4136	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.50C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	889	767	842	718	803	679
GEOGRAPHIC DIVISION						
Northeast	361	316	327	328	321	306
New England	160	136	144	142	131	130
Middle Atlantic	323	286	294	295	293	276
Midwest	296	280	343	333	342	318
East North Central	243	234	274	276	268	266
West North Central	171	158	201	167	202	160
South	575	450	580	460	542	443
South Atlantic	465	325	461	358	426	339
East South Central	188	162	209	181	204	171
West South Central	311	245	277	228	260	230
West	478	434	417	334	404	314
Mountain	219	168	206	160	199	146
Pacific	422	392	361	290	355	277
COUNTY TYPE						
Large Metro	1,465	1,201	757	682	683	607
Small Metro	1,191	943	614	494	549	455
250K – 1 Mil. Pop.	981	781	466	395	419	362
< 250K Pop.	769	581	416	296	371	276
Nonmetro	831	767	452	414	416	384
Urbanized	571	551	298	279	281	256
Less Urbanized	635	576	338	305	303	288
Completely Rural	299	261	180	142	161	131

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.50D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.44	0.37	0.42	0.35	0.40	0.33
GEOGRAPHIC DIVISION						
Northeast	0.97	0.85	0.88	0.88	0.87	0.82
New England	1.63	1.39	1.48	1.44	1.34	1.32
Middle Atlantic	1.18	1.04	1.08	1.07	1.07	1.00
Midwest	0.68	0.64	0.79	0.76	0.79	0.73
East North Central	0.81	0.77	0.91	0.91	0.89	0.88
West North Central	1.29	1.18	1.51	1.25	1.52	1.19
South	0.76	0.59	0.77	0.60	0.72	0.58
South Atlantic	1.16	0.80	1.15	0.88	1.06	0.83
East South Central	1.57	1.34	1.75	1.50	1.70	1.42
West South Central	1.35	1.04	1.20	0.97	1.13	0.98
West	1.02	0.91	0.89	0.70	0.86	0.66
Mountain	1.53	1.15	1.44	1.10	1.39	1.01
Pacific	1.29	1.18	1.11	0.87	1.09	0.83
COUNTY TYPE						
Large Metro	0.67	0.56	0.58	0.48	0.56	0.44
Small Metro	0.70	0.57	0.77	0.65	0.72	0.62
250K – 1 Mil. Pop.	0.86	0.70	0.87	0.77	0.84	0.72
< 250K Pop.	1.16	1.04	1.42	1.18	1.34	1.14
Nonmetro	0.94	0.72	1.03	0.82	0.98	0.79
Urbanized	1.42	1.18	1.50	1.23	1.43	1.20
Less Urbanized	1.46	1.03	1.55	1.20	1.45	1.16
Completely Rural	2.47	2.01	3.16	2.73	3.02	2.54

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.50P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3762		0.7683		0.8507	
GEOGRAPHIC DIVISION						
Northeast	0.7992		0.3651		0.4168	
New England	0.6085		0.8896		0.7199	
Middle Atlantic	0.5952		0.3478		0.4657	
Midwest	0.7553		0.2950		0.4001	
East North Central	0.7931		0.7226		0.8620	
West North Central	0.8686		0.1661		0.2033	
South	0.3710		0.6145		0.5946	
South Atlantic	0.9594		0.2825		0.3203	
East South Central	0.8966		0.7790		0.9001	
West South Central	0.1365		0.3613		0.7081	
West	0.8635		0.9237		0.8057	
Mountain	0.4784		0.7801		0.7912	
Pacific	0.8824		0.7888		0.8936	
COUNTY TYPE						
Large Metro	0.9098		0.2653		0.4439	
Small Metro	0.0472		0.4980		0.5386	
250K – 1 Mil. Pop.	0.1357		0.7683		0.9536	
< 250K Pop.	0.1770		0.1183		0.3373	
Nonmetro	0.7668		0.9679		0.7275	
Urbanized	0.3453		0.3790		0.5639	
Less Urbanized	0.5774		0.5446		0.9794	
Completely Rural	0.9059		0.4870		0.8411	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.51C Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	909	803	873	771	834	729
GEOGRAPHIC DIVISION						
Northeast	366	334	341	343	331	323
New England	164	139	157	154	134	144
Middle Atlantic	326	304	303	307	302	287
Midwest	305	294	355	358	350	339
East North Central	249	251	285	296	277	284
West North Central	178	163	209	182	208	176
South	586	462	604	485	562	468
South Atlantic	469	336	479	367	442	350
East South Central	191	168	217	195	208	185
West South Central	317	255	284	245	266	243
West	488	450	430	357	418	334
Mountain	226	176	215	174	209	162
Pacific	431	407	370	308	366	291
COUNTY TYPE						
Large Metro	1,553	1,270	827	738	744	655
Small Metro	1,255	1,005	669	552	588	506
250K – 1 Mil. Pop.	1,042	823	504	437	446	399
< 250K Pop.	817	636	465	337	408	311
Nonmetro	882	841	497	471	452	434
Urbanized	608	611	331	325	305	298
Less Urbanized	684	633	376	351	337	329
Completely Rural	322	283	197	160	175	144

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.51D Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.38	0.33	0.37	0.32	0.35	0.30
GEOGRAPHIC DIVISION						
Northeast	0.85	0.77	0.79	0.79	0.77	0.74
New England	1.44	1.22	1.38	1.34	1.18	1.25
Middle Atlantic	1.03	0.95	0.95	0.96	0.95	0.90
Midwest	0.60	0.58	0.70	0.70	0.69	0.66
East North Central	0.71	0.71	0.81	0.84	0.79	0.80
West North Central	1.14	1.04	1.34	1.16	1.33	1.12
South	0.67	0.52	0.69	0.54	0.64	0.52
South Atlantic	1.00	0.71	1.02	0.77	0.94	0.74
East South Central	1.36	1.19	1.55	1.38	1.48	1.31
West South Central	1.16	0.92	1.04	0.88	0.97	0.87
West	0.88	0.80	0.78	0.63	0.76	0.59
Mountain	1.34	1.03	1.28	1.02	1.24	0.95
Pacific	1.12	1.04	0.96	0.79	0.95	0.74
COUNTY TYPE						
Large Metro	0.58	0.50	0.51	0.44	0.50	0.40
Small Metro	0.61	0.52	0.68	0.59	0.63	0.56
250K – 1 Mil. Pop.	0.76	0.63	0.76	0.70	0.73	0.66
< 250K Pop.	0.98	0.94	1.25	1.07	1.17	1.03
Nonmetro	0.83	0.64	0.93	0.75	0.88	0.72
Urbanized	1.30	1.04	1.35	1.12	1.28	1.11
Less Urbanized	1.27	0.92	1.41	1.08	1.32	1.05
Completely Rural	2.30	1.83	2.93	2.58	2.76	2.37

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.51P Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2243		0.7973		0.6390	
GEOGRAPHIC DIVISION						
Northeast	0.7171		0.5119		0.6199	
New England	0.6683		0.9056		0.6489	
Middle Atlantic	0.5418		0.4993		0.7371	
Midwest	0.8706		0.2886		0.3173	
East North Central	0.9080		0.6873		0.8720	
West North Central	0.9064		0.1838		0.1206	
South	0.2733		0.7770		0.7057	
South Atlantic	0.7319		0.4528		0.5160	
East South Central	0.9915		0.7024		0.9919	
West South Central	0.1309		0.3645		0.8214	
West	0.5665		0.6121		0.3297	
Mountain	0.3194		0.4785		0.4086	
Pacific	0.8709		0.8556		0.5163	
COUNTY TYPE						
Large Metro	0.8655		0.5700		0.8353	
Small Metro	0.0469		0.4963		0.4938	
250K – 1 Mil. Pop.	0.0912		0.8277		0.8148	
< 250K Pop.	0.2908		0.1506		0.4066	
Nonmetro	0.9339		0.7210		0.8912	
Urbanized	0.4332		0.7117		0.9508	
Less Urbanized	0.4220		0.5214		0.7491	
Completely Rural	0.8344		0.5226		0.9117	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.52C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	966	845	830	750	798	700
GEOGRAPHIC DIVISION						
Northeast	379	353	329	341	315	317
New England	182	146	143	149	127	144
Middle Atlantic	332	323	297	305	289	280
Midwest	337	323	348	340	331	315
East North Central	278	280	278	279	264	266
West North Central	186	170	205	185	196	182
South	602	495	556	469	542	442
South Atlantic	486	365	433	343	427	324
East South Central	203	189	203	187	194	175
West South Central	323	247	273	235	252	226
West	526	441	411	333	394	300
Mountain	230	176	211	163	202	151
Pacific	472	398	353	287	340	260
COUNTY TYPE						
Large Metro	1,464	1,217	758	667	690	602
Small Metro	1,182	975	592	508	543	464
250K – 1 Mil. Pop.	966	793	447	399	408	368
< 250K Pop.	787	614	414	316	376	282
Nonmetro	865	814	433	430	397	392
Urbanized	604	585	284	288	267	258
Less Urbanized	662	624	330	329	295	304
Completely Rural	321	265	164	128	149	114

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.52D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.37	0.32	0.32	0.28	0.30	0.26
GEOGRAPHIC DIVISION						
Northeast	0.80	0.74	0.69	0.72	0.67	0.67
New England	1.46	1.16	1.15	1.19	1.02	1.14
Middle Atlantic	0.95	0.92	0.85	0.87	0.83	0.80
Midwest	0.60	0.57	0.62	0.60	0.59	0.56
East North Central	0.71	0.72	0.71	0.71	0.68	0.68
West North Central	1.08	0.98	1.19	1.07	1.14	1.05
South	0.62	0.50	0.57	0.47	0.56	0.45
South Atlantic	0.94	0.70	0.84	0.66	0.83	0.62
East South Central	1.31	1.21	1.31	1.20	1.25	1.12
West South Central	1.06	0.79	0.89	0.75	0.83	0.73
West	0.86	0.71	0.67	0.54	0.64	0.48
Mountain	1.23	0.92	1.13	0.86	1.08	0.79
Pacific	1.11	0.92	0.83	0.67	0.80	0.60
COUNTY TYPE						
Large Metro	0.53	0.46	0.44	0.39	0.42	0.36
Small Metro	0.62	0.50	0.58	0.50	0.56	0.48
250K – 1 Mil. Pop.	0.76	0.63	0.68	0.60	0.64	0.57
< 250K Pop.	1.03	0.86	1.06	0.91	1.02	0.86
Nonmetro	0.84	0.64	0.75	0.67	0.71	0.63
Urbanized	1.30	1.05	1.06	1.02	1.04	0.97
Less Urbanized	1.28	0.92	1.16	1.01	1.07	0.98
Completely Rural	2.12	1.82	2.30	2.01	2.21	1.87

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.52P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1151		0.3341		0.2954	
GEOGRAPHIC DIVISION						
Northeast	0.4559		0.9450		0.8060	
New England	0.7942		0.6068		0.7958	
Middle Atlantic	0.4825		0.7431		0.6913	
Midwest	0.4521		0.1246		0.1651	
East North Central	0.5279		0.4245		0.6939	
West North Central	0.6859		0.1168		0.0689	
South	0.2030		0.9827		0.9855	
South Atlantic	0.8363		0.6348		0.7748	
East South Central	0.5774		0.4552		0.4166	
West South Central	0.0826		0.8474		0.8237	
West	0.6945		0.5233		0.2374	
Mountain	0.4115		0.8908		0.9238	
Pacific	0.9567		0.5067		0.1887	
COUNTY TYPE						
Large Metro	0.6327		0.6969		0.7654	
Small Metro	0.1333		0.8910		0.7453	
250K – 1 Mil. Pop.	0.2396		0.8752		0.9117	
< 250K Pop.	0.3429		0.6540		0.6866	
Nonmetro	0.4186		0.2594		0.2195	
Urbanized	0.6898		0.7994		0.4115	
Less Urbanized	0.4151		0.2251		0.3858	
Completely Rural	0.9879		0.5641		0.5985	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.53C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	83	89	67	71	51	52
GEOGRAPHIC DIVISION						
Northeast	28	30	24	22	19	17
New England	13	12	12	9	9	7
Middle Atlantic	25	27	21	20	16	16
Midwest	34	41	27	31	20	24
East North Central	29	37	21	26	15	20
West North Central	19	19	15	16	12	13
South	53	54	43	44	32	33
South Atlantic	40	34	32	28	23	21
East South Central	17	23	16	18	13	14
West South Central	31	36	27	29	18	21
West	45	49	37	41	29	26
Mountain	22	22	17	19	13	15
Pacific	38	43	32	36	25	21
COUNTY TYPE						
Large Metro	65	71	51	54	37	40
Small Metro	56	54	43	44	30	27
250K – 1 Mil. Pop.	45	42	36	34	25	21
< 250K Pop.	33	34	23	28	17	16
Nonmetro	43	43	33	30	24	25
Urbanized	23	25	19	19	14	14
Less Urbanized	32	34	23	23	16	20
Completely Rural	19	12	15	9	11	9

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.53D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.33	0.36	0.27	0.29	0.20	0.21
GEOGRAPHIC DIVISION						
Northeast	0.67	0.72	0.57	0.54	0.44	0.42
New England	1.16	1.09	1.10	0.85	0.85	0.64
Middle Atlantic	0.81	0.87	0.67	0.65	0.52	0.52
Midwest	0.64	0.76	0.49	0.57	0.36	0.44
East North Central	0.76	0.98	0.56	0.71	0.41	0.54
West North Central	1.17	1.15	0.93	0.98	0.74	0.78
South	0.57	0.58	0.46	0.47	0.34	0.36
South Atlantic	0.85	0.73	0.68	0.60	0.50	0.45
East South Central	1.15	1.56	1.08	1.19	0.86	0.97
West South Central	0.96	1.09	0.82	0.89	0.56	0.65
West	0.75	0.82	0.63	0.68	0.49	0.43
Mountain	1.19	1.18	0.92	1.02	0.70	0.80
Pacific	0.93	1.07	0.79	0.89	0.62	0.53
COUNTY TYPE						
Large Metro	0.42	0.46	0.35	0.36	0.26	0.27
Small Metro	0.63	0.64	0.51	0.55	0.38	0.36
250K – 1 Mil. Pop.	0.73	0.72	0.61	0.60	0.46	0.42
< 250K Pop.	1.17	1.36	0.87	1.19	0.67	0.70
Nonmetro	0.89	0.98	0.73	0.72	0.55	0.61
Urbanized	1.15	1.41	0.98	1.08	0.78	0.84
Less Urbanized	1.40	1.42	1.15	1.07	0.84	0.93
Completely Rural	2.64	2.72	2.36	2.09	1.78	2.07

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.53P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0020		0.0004		0.0115	
GEOGRAPHIC DIVISION						
Northeast	0.0041		0.0002		0.0121	
New England	0.4573		0.2629		0.4491	
Middle Atlantic	0.0042		0.0003		0.0151	
Midwest	0.0224		0.0093		0.3803	
East North Central	0.0784		0.0528		0.5289	
West North Central	0.1383		0.0670		0.5311	
South	0.5186		0.1631		0.2861	
South Atlantic	0.1115		0.1147		0.2367	
East South Central	0.4254		0.4855		0.7911	
West South Central	0.7944		0.9432		0.8020	
West	0.1814		0.5294		0.2688	
Mountain	0.0687		0.3398		0.9785	
Pacific	0.5819		0.8263		0.1957	
COUNTY TYPE						
Large Metro	0.0103		0.0003		0.0590	
Small Metro	0.2352		0.8675		0.1564	
250K – 1 Mil. Pop.	0.4942		0.6576		0.2898	
< 250K Pop.	0.2946		0.7660		0.3338	
Nonmetro	0.2690		0.0766		0.4998	
Urbanized	0.2481		0.0862		0.1895	
Less Urbanized	0.7664		0.3450		0.6767	
Completely Rural	0.3578		0.8383		0.5747	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.54C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	170	201	171	196	162	185
GEOGRAPHIC DIVISION						
Northeast	65	83	64	78	61	71
New England	30	47	33	41	33	39
Middle Atlantic	57	68	54	68	51	60
Midwest	62	83	63	83	63	77
East North Central	54	71	53	64	54	64
West North Central	31	40	32	47	32	45
South	100	117	105	125	98	120
South Atlantic	75	76	79	75	72	72
East South Central	37	48	34	51	35	51
West South Central	55	73	56	77	53	75
West	103	100	96	92	91	83
Mountain	39	44	40	42	37	43
Pacific	94	89	86	82	82	72
COUNTY TYPE						
Large Metro	255	245	185	195	152	171
Small Metro	193	189	145	143	120	122
250K – 1 Mil. Pop.	140	143	108	110	90	96
< 250K Pop.	141	121	101	89	82	73
Nonmetro	131	140	102	110	88	89
Urbanized	99	104	75	78	61	64
Less Urbanized	92	92	72	74	65	61
Completely Rural	38	30	30	27	24	20

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.54D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.49	0.57	0.49	0.56	0.46	0.53
GEOGRAPHIC DIVISION						
Northeast	1.05	1.35	1.03	1.27	0.99	1.16
New England	1.85	2.84	2.03	2.51	2.02	2.40
Middle Atlantic	1.25	1.50	1.19	1.51	1.13	1.34
Midwest	0.84	1.12	0.84	1.12	0.85	1.04
East North Central	1.05	1.39	1.04	1.26	1.05	1.25
West North Central	1.35	1.74	1.40	2.01	1.41	1.95
South	0.78	0.91	0.81	0.96	0.76	0.93
South Atlantic	1.14	1.15	1.20	1.14	1.10	1.09
East South Central	1.81	2.35	1.68	2.47	1.70	2.47
West South Central	1.30	1.69	1.32	1.80	1.24	1.75
West	1.23	1.19	1.14	1.10	1.08	0.99
Mountain	1.55	1.71	1.58	1.64	1.45	1.67
Pacific	1.61	1.52	1.48	1.41	1.41	1.23
COUNTY TYPE						
Large Metro	0.71	0.82	0.72	0.77	0.66	0.74
Small Metro	0.85	0.97	0.79	0.93	0.77	0.88
250K – 1 Mil. Pop.	1.05	1.23	0.97	1.11	0.93	1.07
< 250K Pop.	1.38	1.48	1.35	1.61	1.37	1.49
Nonmetro	1.10	1.21	1.05	1.37	1.05	1.26
Urbanized	1.62	1.89	1.46	2.15	1.44	1.93
Less Urbanized	1.47	1.53	1.52	1.75	1.54	1.65
Completely Rural	4.00	4.06	4.16	4.19	3.53	3.89

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.54P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0190		0.0126		0.0013	
GEOGRAPHIC DIVISION						
Northeast	0.2584		0.4377		0.2830	
New England	0.1776		0.6852		0.8475	
Middle Atlantic	0.6184		0.5111		0.1615	
Midwest	0.8712		0.3393		0.1080	
East North Central	0.7555		0.7455		0.8177	
West North Central	0.8278		0.2020		0.0111	
South	0.1161		0.2243		0.3813	
South Atlantic	0.0259		0.0343		0.0863	
East South Central	0.6600		0.7993		0.9220	
West South Central	0.8148		0.8563		0.6216	
West	0.0569		0.0389		0.0012	
Mountain	0.0418		0.0736		0.0183	
Pacific	0.2312		0.1501		0.0147	
COUNTY TYPE						
Large Metro	0.0514		0.0615		0.0210	
Small Metro	0.2959		0.5237		0.3283	
250K – 1 Mil. Pop.	0.0718		0.3715		0.3152	
< 250K Pop.	0.4477		0.9026		0.7692	
Nonmetro	0.5264		0.0467		0.0080	
Urbanized	0.7592		0.0322		0.0714	
Less Urbanized	0.2214		0.1690		0.0092	
Completely Rural	0.4858		0.3194		0.7380	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.55C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	915	788	797	691	770	651
GEOGRAPHIC DIVISION						
Northeast	369	328	314	324	305	299
New England	168	138	136	138	126	134
Middle Atlantic	328	301	284	291	279	266
Midwest	316	291	332	307	318	293
East North Central	260	249	265	253	253	246
West North Central	175	158	194	161	188	159
South	574	464	535	438	525	415
South Atlantic	473	338	420	328	417	310
East South Central	190	170	196	161	188	155
West South Central	301	232	263	221	244	213
West	498	423	395	309	383	282
Mountain	226	164	203	147	195	135
Pacific	443	379	339	269	331	248
COUNTY TYPE						
Large Metro	1,373	1,147	685	614	634	560
Small Metro	1,113	906	544	445	510	416
250K – 1 Mil. Pop.	901	748	410	353	384	331
< 250K Pop.	740	554	375	272	345	252
Nonmetro	806	731	389	368	364	341
Urbanized	567	521	252	241	243	217
Less Urbanized	602	553	291	280	262	263
Completely Rural	291	239	145	110	135	102

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.55D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.45	0.38	0.39	0.34	0.38	0.32
GEOGRAPHIC DIVISION						
Northeast	1.00	0.88	0.85	0.87	0.82	0.80
New England	1.72	1.40	1.39	1.40	1.29	1.36
Middle Atlantic	1.20	1.10	1.04	1.06	1.02	0.97
Midwest	0.73	0.67	0.77	0.70	0.73	0.67
East North Central	0.86	0.82	0.88	0.84	0.84	0.81
West North Central	1.32	1.18	1.46	1.20	1.41	1.19
South	0.76	0.61	0.71	0.57	0.70	0.54
South Atlantic	1.18	0.83	1.04	0.80	1.04	0.76
East South Central	1.59	1.41	1.64	1.34	1.57	1.29
West South Central	1.31	0.98	1.14	0.94	1.06	0.90
West	1.06	0.88	0.84	0.65	0.82	0.59
Mountain	1.58	1.12	1.42	1.01	1.36	0.93
Pacific	1.36	1.14	1.04	0.81	1.01	0.75
COUNTY TYPE						
Large Metro	0.65	0.57	0.55	0.47	0.52	0.44
Small Metro	0.75	0.59	0.73	0.60	0.71	0.57
250K – 1 Mil. Pop.	0.91	0.75	0.86	0.71	0.82	0.68
< 250K Pop.	1.31	1.01	1.35	1.09	1.30	1.05
Nonmetro	1.02	0.75	0.92	0.79	0.88	0.75
Urbanized	1.52	1.20	1.30	1.18	1.28	1.12
Less Urbanized	1.59	1.10	1.41	1.22	1.31	1.18
Completely Rural	2.47	2.04	2.80	2.29	2.70	2.22

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.55P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3210		0.9293		0.8733	
GEOGRAPHIC DIVISION						
Northeast	0.7117		0.6104		0.5299	
New England	0.9286		0.7166		0.7738	
Middle Atlantic	0.6504		0.4604		0.4037	
Midwest	0.5209		0.2247		0.2926	
East North Central	0.5422		0.5353		0.7354	
West North Central	0.8111		0.2019		0.1719	
South	0.3106		0.6987		0.7758	
South Atlantic	0.7759		0.2937		0.4870	
East South Central	0.5403		0.4168		0.4062	
West South Central	0.0476		0.8068		0.9133	
West	0.9567		0.9568		0.7539	
Mountain	0.8055		0.7120		0.6469	
Pacific	0.8581		0.7823		0.5329	
COUNTY TYPE						
Large Metro	0.9392		0.6784		0.6846	
Small Metro	0.1442		0.9714		0.9302	
250K – 1 Mil. Pop.	0.3410		0.7245		0.9031	
< 250K Pop.	0.2270		0.5834		0.7487	
Nonmetro	0.5788		0.5361		0.5035	
Urbanized	0.9061		0.7113		0.7143	
Less Urbanized	0.5754		0.3517		0.6589	
Completely Rural	0.7539		0.4545		0.5741	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.56C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	935	823	824	741	796	691
GEOGRAPHIC DIVISION						
Northeast	374	344	326	337	314	313
New England	172	145	142	147	125	142
Middle Atlantic	331	316	294	302	288	277
Midwest	325	311	344	336	327	315
East North Central	267	269	275	275	261	264
West North Central	181	166	201	180	194	179
South	585	479	555	463	541	438
South Atlantic	475	354	434	339	427	321
East South Central	194	175	202	180	192	170
West South Central	308	239	271	236	250	224
West	509	432	408	328	395	297
Mountain	229	173	210	163	201	150
Pacific	456	388	351	283	341	257
COUNTY TYPE						
Large Metro	1,451	1,205	750	661	685	597
Small Metro	1,165	960	581	499	538	460
250K – 1 Mil. Pop.	950	783	435	391	403	365
< 250K Pop.	779	604	410	311	374	279
Nonmetro	852	799	424	421	393	386
Urbanized	596	576	278	281	263	254
Less Urbanized	649	608	323	320	291	298
Completely Rural	312	260	160	124	147	111

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.56D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.34	0.35	0.31	0.34	0.29
GEOGRAPHIC DIVISION						
Northeast	0.86	0.79	0.75	0.77	0.73	0.72
New England	1.51	1.26	1.25	1.28	1.10	1.24
Middle Atlantic	1.04	0.99	0.92	0.94	0.90	0.87
Midwest	0.64	0.61	0.68	0.66	0.64	0.62
East North Central	0.76	0.76	0.78	0.78	0.74	0.75
West North Central	1.16	1.06	1.29	1.15	1.24	1.14
South	0.66	0.54	0.63	0.52	0.61	0.49
South Atlantic	1.02	0.74	0.93	0.71	0.91	0.68
East South Central	1.38	1.24	1.44	1.28	1.37	1.21
West South Central	1.13	0.86	0.99	0.85	0.91	0.81
West	0.92	0.77	0.74	0.58	0.71	0.53
Mountain	1.36	1.01	1.25	0.95	1.19	0.88
Pacific	1.18	0.99	0.91	0.72	0.89	0.66
COUNTY TYPE						
Large Metro	0.57	0.50	0.49	0.42	0.46	0.39
Small Metro	0.66	0.53	0.63	0.55	0.61	0.52
250K – 1 Mil. Pop.	0.81	0.67	0.74	0.66	0.71	0.62
< 250K Pop.	1.12	0.91	1.16	0.99	1.13	0.94
Nonmetro	0.91	0.67	0.83	0.74	0.80	0.69
Urbanized	1.38	1.09	1.17	1.11	1.15	1.05
Less Urbanized	1.39	0.98	1.28	1.11	1.19	1.08
Completely Rural	2.27	1.88	2.60	2.16	2.49	2.01

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.56P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1512		0.4823		0.3546	
GEOGRAPHIC DIVISION						
Northeast	0.5601		0.7549		0.7093	
New England	0.8053		0.6538		0.8213	
Middle Atlantic	0.5987		0.5718		0.6019	
Midwest	0.5684		0.1752		0.1737	
East North Central	0.6171		0.5087		0.7066	
West North Central	0.7841		0.1451		0.0703	
South	0.1954		0.9366		0.9449	
South Atlantic	0.9157		0.5584		0.7321	
East South Central	0.4843		0.4790		0.4156	
West South Central	0.0612		0.8443		0.8160	
West	0.7306		0.5451		0.2600	
Mountain	0.5152		0.9584		0.9185	
Pacific	0.9416		0.5006		0.2139	
COUNTY TYPE						
Large Metro	0.7654		0.9188		0.8496	
Small Metro	0.1036		0.8488		0.7579	
250K – 1 Mil. Pop.	0.1874		0.8847		0.9194	
< 250K Pop.	0.3351		0.6041		0.6989	
Nonmetro	0.4899		0.3381		0.2443	
Urbanized	0.8111		0.9350		0.4824	
Less Urbanized	0.4467		0.2665		0.3771	
Completely Rural	0.9295		0.5410		0.5915	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.57C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	727	579	334	276	300	249
GEOGRAPHIC DIVISION						
Northeast	244	218	90	84	79	72
New England	112	112	37	45	31	33
Middle Atlantic	217	187	82	71	73	64
Midwest	285	260	140	129	127	113
East North Central	228	227	114	105	107	93
West North Central	174	155	82	70	70	62
South	498	368	231	202	211	189
South Atlantic	391	261	148	145	139	133
East South Central	178	137	111	97	106	95
West South Central	267	230	139	108	121	96
West	361	298	168	111	151	94
Mountain	196	139	96	62	90	50
Pacific	302	263	139	91	122	81
COUNTY TYPE						
Large Metro	603	504	221	203	185	171
Small Metro	542	404	194	158	170	138
250K – 1 Mil. Pop.	463	327	144	128	128	115
< 250K Pop.	326	252	124	93	112	77
Nonmetro	436	365	193	164	176	153
Urbanized	263	240	118	101	106	94
Less Urbanized	353	277	148	110	133	97
Completely Rural	155	136	73	86	66	83

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.57D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.28	0.22	0.13	0.10	0.11	0.09
GEOGRAPHIC DIVISION						
Northeast	0.51	0.46	0.19	0.18	0.17	0.15
New England	0.89	0.89	0.30	0.36	0.24	0.26
Middle Atlantic	0.62	0.53	0.23	0.20	0.21	0.18
Midwest	0.51	0.46	0.25	0.23	0.23	0.20
East North Central	0.59	0.58	0.29	0.27	0.27	0.24
West North Central	1.01	0.89	0.48	0.40	0.41	0.36
South	0.51	0.37	0.24	0.20	0.22	0.19
South Atlantic	0.76	0.50	0.29	0.28	0.27	0.25
East South Central	1.15	0.88	0.72	0.62	0.69	0.61
West South Central	0.87	0.74	0.45	0.35	0.40	0.31
West	0.59	0.48	0.27	0.18	0.25	0.15
Mountain	1.05	0.73	0.51	0.33	0.48	0.26
Pacific	0.71	0.61	0.33	0.21	0.29	0.19
COUNTY TYPE						
Large Metro	0.37	0.29	0.15	0.13	0.13	0.11
Small Metro	0.51	0.39	0.23	0.19	0.21	0.17
250K – 1 Mil. Pop.	0.62	0.49	0.25	0.23	0.23	0.21
< 250K Pop.	0.92	0.69	0.46	0.34	0.43	0.28
Nonmetro	0.74	0.60	0.42	0.34	0.39	0.33
Urbanized	0.94	0.91	0.57	0.47	0.51	0.47
Less Urbanized	1.30	0.84	0.68	0.45	0.62	0.42
Completely Rural	2.03	2.06	1.32	1.79	1.26	1.76

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.57P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1556		0.3238		0.5359	
GEOGRAPHIC DIVISION						
Northeast	0.8980		0.9308		0.8380	
New England	0.7184		0.3958		0.3607	
Middle Atlantic	0.9575		0.7210		0.5166	
Midwest	0.8113		0.7908		0.3604	
East North Central	0.8086		0.9746		0.7378	
West North Central	0.9445		0.6761		0.2490	
South	0.6839		0.5989		0.8152	
South Atlantic	0.5724		0.9003		0.8798	
East South Central	0.2474		0.2377		0.2208	
West South Central	0.4896		0.0824		0.1772	
West	0.0297		0.1039		0.0825	
Mountain	0.1640		0.1881		0.0765	
Pacific	0.0867		0.2841		0.3885	
COUNTY TYPE						
Large Metro	0.6370		0.8451		0.9014	
Small Metro	0.0136		0.8670		0.7037	
250K – 1 Mil. Pop.	0.0141		0.9330		0.9516	
< 250K Pop.	0.4599		0.7038		0.5883	
Nonmetro	0.8695		0.3768		0.7077	
Urbanized	0.6301		0.5608		0.9759	
Less Urbanized	0.8230		0.2935		0.3323	
Completely Rural	0.7713		0.7886		0.6070	

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.58C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	59	58	49	47	31	34
GEOGRAPHIC DIVISION						
Northeast	19	19	15	17	12	13
New England	8	8	7	8	4	5
Middle Atlantic	17	17	14	15	12	12
Midwest	23	27	18	21	12	15
East North Central	18	19	15	14	9	10
West North Central	13	18	10	15	7	11
South	45	38	38	32	23	23
South Atlantic	36	24	31	21	18	16
East South Central	16	18	13	14	10	11
West South Central	22	24	19	21	12	13
West	21	26	18	20	11	13
Mountain	12	15	10	12	5	8
Pacific	17	21	15	16	9	10
COUNTY TYPE						
Large Metro	39	38	32	32	19	20
Small Metro	38	35	31	28	17	20
250K – 1 Mil. Pop.	32	28	27	20	14	15
< 250K Pop.	20	22	16	19	10	14
Nonmetro	34	28	26	22	20	17
Urbanized	20	19	15	14	11	10
Less Urbanized	22	22	19	17	14	14
Completely Rural	17	9	10	7	8	6

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.58D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.24	0.23	0.20	0.19	0.12	0.14
GEOGRAPHIC DIVISION						
Northeast	0.45	0.46	0.37	0.40	0.29	0.31
New England	0.73	0.78	0.68	0.74	0.37	0.48
Middle Atlantic	0.55	0.57	0.44	0.49	0.37	0.39
Midwest	0.42	0.50	0.33	0.39	0.22	0.28
East North Central	0.48	0.52	0.39	0.38	0.25	0.27
West North Central	0.77	1.09	0.62	0.88	0.43	0.67
South	0.48	0.41	0.41	0.34	0.25	0.25
South Atlantic	0.77	0.52	0.67	0.45	0.38	0.33
East South Central	1.10	1.20	0.89	0.92	0.65	0.76
West South Central	0.69	0.75	0.58	0.64	0.37	0.41
West	0.35	0.43	0.30	0.34	0.18	0.22
Mountain	0.64	0.78	0.52	0.62	0.28	0.41
Pacific	0.41	0.52	0.36	0.41	0.23	0.26
COUNTY TYPE						
Large Metro	0.27	0.26	0.23	0.22	0.13	0.14
Small Metro	0.45	0.48	0.38	0.38	0.21	0.28
250K – 1 Mil. Pop.	0.56	0.55	0.48	0.40	0.25	0.30
< 250K Pop.	0.78	0.92	0.65	0.83	0.41	0.60
Nonmetro	0.78	0.68	0.62	0.56	0.49	0.45
Urbanized	1.13	1.14	0.85	0.89	0.65	0.65
Less Urbanized	1.10	1.03	0.99	0.84	0.76	0.69
Completely Rural	2.74	2.08	1.82	1.60	1.69	1.42

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.58P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2986		0.3099		0.9525	
GEOGRAPHIC DIVISION						
Northeast	0.7559		0.7882		0.6608	
New England	0.3580		0.4003		0.5372	
Middle Atlantic	0.4344		0.8749		0.4725	
Midwest	0.6466		0.1816		0.9737	
East North Central	0.2657		0.0222		0.1062	
West North Central	0.5438		0.5859		0.1015	
South	0.1479		0.4262		0.9052	
South Atlantic	0.0724		0.2128		0.8580	
East South Central	0.7007		0.5003		0.6411	
West South Central	0.6319		0.8467		0.9328	
West	0.4856		0.9847		0.6807	
Mountain	0.9802		0.9698		0.2164	
Pacific	0.3925		0.9837		0.7322	
COUNTY TYPE						
Large Metro	1.0000		0.8879		0.8635	
Small Metro	0.8048		0.5562		0.4597	
250K – 1 Mil. Pop.	0.9451		0.3967		0.7810	
< 250K Pop.	0.5977		0.8828		0.4118	
Nonmetro	0.1834		0.2236		0.8761	
Urbanized	0.5833		0.7046		0.7649	
Less Urbanized	0.4645		0.2416		0.9842	
Completely Rural	0.1917		0.6568		0.9952	

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.59C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	137	141	94	101	71	82
GEOGRAPHIC DIVISION						
Northeast	51	47	33	31	25	24
New England	26	24	16	19	12	11
Middle Atlantic	44	41	30	25	22	21
Midwest	56	67	40	55	33	45
East North Central	46	59	30	46	24	38
West North Central	33	36	25	29	22	24
South	86	88	66	62	52	53
South Atlantic	68	59	50	40	36	31
East South Central	30	38	22	29	20	26
West South Central	41	51	34	38	28	34
West	74	69	50	46	32	36
Mountain	31	36	23	26	17	20
Pacific	67	58	45	39	28	29
COUNTY TYPE						
Large Metro	121	115	71	73	46	58
Small Metro	105	98	66	68	50	55
250K – 1 Mil. Pop.	77	74	48	54	39	42
< 250K Pop.	72	65	45	43	33	36
Nonmetro	78	79	52	49	37	38
Urbanized	63	55	39	32	23	22
Less Urbanized	49	56	35	38	28	30
Completely Rural	28	20	20	12	13	10

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.59D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.40	0.27	0.29	0.20	0.23
GEOGRAPHIC DIVISION						
Northeast	0.83	0.76	0.54	0.51	0.41	0.39
New England	1.59	1.46	0.96	1.14	0.73	0.65
Middle Atlantic	0.98	0.90	0.65	0.56	0.50	0.48
Midwest	0.76	0.90	0.53	0.74	0.45	0.61
East North Central	0.90	1.15	0.59	0.91	0.48	0.75
West North Central	1.45	1.57	1.10	1.25	0.97	1.04
South	0.67	0.68	0.51	0.48	0.40	0.41
South Atlantic	1.04	0.90	0.77	0.60	0.55	0.47
East South Central	1.48	1.84	1.09	1.42	0.99	1.29
West South Central	0.96	1.18	0.79	0.89	0.65	0.79
West	0.89	0.82	0.60	0.55	0.39	0.42
Mountain	1.21	1.39	0.89	1.01	0.67	0.79
Pacific	1.15	0.99	0.77	0.66	0.48	0.50
COUNTY TYPE						
Large Metro	0.54	0.54	0.36	0.37	0.24	0.29
Small Metro	0.73	0.75	0.52	0.58	0.41	0.49
250K – 1 Mil. Pop.	0.91	0.88	0.63	0.71	0.50	0.57
< 250K Pop.	1.13	1.38	0.88	1.01	0.73	0.91
Nonmetro	1.06	1.04	0.83	0.80	0.63	0.66
Urbanized	1.67	1.59	1.27	1.11	0.81	0.84
Less Urbanized	1.56	1.58	1.25	1.33	1.08	1.09
Completely Rural	3.61	3.95	2.92	2.63	2.36	2.23

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.59P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1043		0.2600		0.5588	
GEOGRAPHIC DIVISION						
Northeast	0.0060		0.1441		0.2392	
New England	0.3064		0.6867		0.6946	
Middle Atlantic	0.0100		0.1413		0.2672	
Midwest	0.9170		0.8511		0.9729	
East North Central	0.6960		0.9766		0.7646	
West North Central	0.4454		0.7162		0.6321	
South	0.7001		0.5745		0.8268	
South Atlantic	0.6962		0.7546		0.7506	
East South Central	0.7747		0.9268		0.9223	
West South Central	0.6961		0.5022		0.5206	
West	0.3726		0.4550		0.9251	
Mountain	0.8800		0.9963		0.9488	
Pacific	0.2689		0.3856		0.9381	
COUNTY TYPE						
Large Metro	0.1467		0.5003		0.7675	
Small Metro	0.8571		0.9812		0.8373	
250K – 1 Mil. Pop.	0.5334		0.3075		0.1878	
< 250K Pop.	0.2601		0.1920		0.1998	
Nonmetro	0.1910		0.2465		0.2337	
Urbanized	0.0137		0.2137		0.0985	
Less Urbanized	0.3827		1.0000		0.8958	
Completely Rural	0.8202		0.3094		0.7551	

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.60C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	705	563	313	255	288	234
GEOGRAPHIC DIVISION						
Northeast	235	210	79	75	71	67
New England	107	106	30	36	25	30
Middle Atlantic	210	182	74	64	67	59
Midwest	276	248	132	114	121	101
East North Central	220	212	109	91	103	82
West North Central	171	145	75	64	64	55
South	480	355	218	188	208	177
South Atlantic	382	250	137	132	133	122
East South Central	171	133	111	93	107	91
West South Central	253	217	129	95	116	87
West	348	291	157	103	145	88
Mountain	191	133	89	57	85	47
Pacific	290	258	130	86	119	76
COUNTY TYPE						
Large Metro	565	473	203	180	178	155
Small Metro	509	375	174	135	161	123
250K – 1 Mil. Pop.	432	307	129	113	120	106
< 250K Pop.	307	227	113	73	107	61
Nonmetro	403	327	177	147	168	140
Urbanized	239	214	109	90	102	88
Less Urbanized	327	242	130	91	122	83
Completely Rural	134	122	64	80	62	78

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.60D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.35	0.27	0.15	0.12	0.14	0.11
GEOGRAPHIC DIVISION						
Northeast	0.64	0.56	0.21	0.20	0.19	0.18
New England	1.09	1.08	0.30	0.37	0.25	0.31
Middle Atlantic	0.77	0.66	0.27	0.23	0.25	0.22
Midwest	0.64	0.57	0.30	0.26	0.28	0.23
East North Central	0.73	0.70	0.36	0.30	0.34	0.27
West North Central	1.29	1.08	0.56	0.48	0.48	0.41
South	0.64	0.46	0.29	0.25	0.28	0.23
South Atlantic	0.95	0.61	0.34	0.32	0.33	0.30
East South Central	1.42	1.10	0.93	0.77	0.89	0.75
West South Central	1.10	0.92	0.56	0.41	0.50	0.37
West	0.74	0.61	0.33	0.21	0.31	0.18
Mountain	1.34	0.91	0.62	0.39	0.60	0.33
Pacific	0.89	0.78	0.40	0.26	0.36	0.23
COUNTY TYPE						
Large Metro	0.46	0.36	0.18	0.15	0.16	0.13
Small Metro	0.65	0.49	0.28	0.22	0.26	0.20
250K – 1 Mil. Pop.	0.77	0.60	0.30	0.27	0.28	0.26
< 250K Pop.	1.25	0.87	0.58	0.37	0.55	0.31
Nonmetro	0.91	0.74	0.51	0.41	0.48	0.40
Urbanized	1.16	1.13	0.70	0.58	0.65	0.58
Less Urbanized	1.59	1.00	0.80	0.53	0.75	0.49
Completely Rural	2.47	2.33	1.64	2.07	1.57	2.04

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.60P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2938		0.6357		0.6310	
GEOGRAPHIC DIVISION						
Northeast	0.4522		0.4986		0.7773	
New England	0.5902		0.2818		0.2676	
Middle Atlantic	0.5736		0.8108		0.8612	
Midwest	0.8043		0.6655		0.3189	
East North Central	0.9361		0.7046		0.6723	
West North Central	0.7593		0.8293		0.2284	
South	0.8300		0.8187		0.8463	
South Atlantic	0.7330		0.8161		0.8296	
East South Central	0.2767		0.2743		0.2227	
West South Central	0.5353		0.0976		0.2101	
West	0.0334		0.1424		0.0672	
Mountain	0.1411		0.1572		0.0488	
Pacific	0.1086		0.4225		0.4001	
COUNTY TYPE						
Large Metro	0.4174		0.9587		0.9366	
Small Metro	0.0093		0.9611		0.6615	
250K – 1 Mil. Pop.	0.0136		0.5107		0.7386	
< 250K Pop.	0.3246		0.3655		0.2504	
Nonmetro	0.8280		0.6337		0.9043	
Urbanized	0.8681		0.8049		0.7254	
Less Urbanized	0.8774		0.3330		0.3141	
Completely Rural	0.8863		0.5650		0.5449	

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.61C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	716	576	331	271	299	246
GEOGRAPHIC DIVISION						
Northeast	242	216	89	82	78	71
New England	112	112	37	43	30	32
Middle Atlantic	214	186	81	69	72	63
Midwest	283	259	140	128	126	112
East North Central	227	225	114	103	106	92
West North Central	173	153	82	70	70	61
South	489	363	230	197	212	186
South Atlantic	385	257	146	139	138	129
East South Central	175	136	112	95	107	94
West South Central	260	227	139	104	121	94
West	358	297	168	110	152	93
Mountain	195	140	95	62	90	50
Pacific	299	261	140	90	124	80
COUNTY TYPE						
Large Metro	595	497	217	200	184	169
Small Metro	531	399	190	151	168	136
250K – 1 Mil. Pop.	454	320	141	122	127	112
< 250K Pop.	322	251	121	88	111	76
Nonmetro	425	357	190	160	176	149
Urbanized	259	234	116	100	105	93
Less Urbanized	344	269	143	105	131	93
Completely Rural	146	132	70	83	65	80

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.61D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.30	0.24	0.14	0.11	0.13	0.10
GEOGRAPHIC DIVISION						
Northeast	0.56	0.50	0.21	0.19	0.18	0.16
New England	0.98	0.98	0.32	0.37	0.27	0.28
Middle Atlantic	0.67	0.58	0.25	0.22	0.23	0.20
Midwest	0.56	0.51	0.27	0.25	0.25	0.22
East North Central	0.64	0.64	0.32	0.29	0.30	0.26
West North Central	1.11	0.98	0.52	0.45	0.45	0.39
South	0.55	0.41	0.26	0.22	0.24	0.21
South Atlantic	0.82	0.54	0.31	0.29	0.30	0.27
East South Central	1.25	0.97	0.80	0.67	0.76	0.66
West South Central	0.95	0.81	0.51	0.37	0.44	0.34
West	0.65	0.53	0.30	0.19	0.27	0.16
Mountain	1.16	0.82	0.56	0.36	0.54	0.29
Pacific	0.78	0.67	0.36	0.23	0.32	0.20
COUNTY TYPE						
Large Metro	0.40	0.32	0.17	0.14	0.14	0.12
Small Metro	0.56	0.43	0.25	0.20	0.23	0.19
250K – 1 Mil. Pop.	0.68	0.53	0.28	0.25	0.25	0.23
< 250K Pop.	1.03	0.77	0.50	0.36	0.47	0.31
Nonmetro	0.81	0.65	0.47	0.37	0.43	0.36
Urbanized	1.03	0.98	0.63	0.52	0.57	0.51
Less Urbanized	1.41	0.91	0.74	0.49	0.68	0.45
Completely Rural	2.22	2.17	1.50	1.88	1.41	1.84

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.61P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.1675		0.4003		0.5218	
GEOGRAPHIC DIVISION						
Northeast	0.8883		0.9669		0.8938	
New England	0.7759		0.4738		0.3979	
Middle Atlantic	1.0000		0.7404		0.5936	
Midwest	0.8275		0.6468		0.3585	
East North Central	0.8652		0.7397		0.6236	
West North Central	0.8927		0.7442		0.3539	
South	0.7731		0.6877		0.7998	
South Atlantic	0.6787		0.9173		0.8932	
East South Central	0.2623		0.2707		0.2391	
West South Central	0.5027		0.0826		0.1703	
West	0.0234		0.1000		0.0749	
Mountain	0.1624		0.1825		0.0603	
Pacific	0.0699		0.2811		0.4028	
COUNTY TYPE						
Large Metro	0.6404		0.8256		0.8881	
Small Metro	0.0120		0.9413		0.6231	
250K – 1 Mil. Pop.	0.0109		0.8171		0.9145	
< 250K Pop.	0.4718		0.6793		0.5089	
Nonmetro	0.9768		0.4747		0.7219	
Urbanized	0.6851		0.6020		0.9910	
Less Urbanized	0.7580		0.3761		0.3264	
Completely Rural	0.8317		0.7551		0.6069	

*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.62C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	922	755	493	394	354	282
GEOGRAPHIC DIVISION						
Northeast	372	320	190	169	135	127
New England	176	161	95	89	59	70
Middle Atlantic	329	271	164	141	122	105
Midwest	346	325	224	175	171	125
East North Central	280	257	166	148	115	102
West North Central	199	202	144	98	121	74
South	630	443	360	253	235	186
South Atlantic	494	335	301	179	181	131
East South Central	212	172	113	103	85	72
West South Central	343	248	175	130	128	104
West	483	411	226	195	169	140
Mountain	248	181	111	90	88	72
Pacific	419	360	199	173	144	119
COUNTY TYPE						
Large Metro	1,044	821	433	350	300	245
Small Metro	787	628	319	218	192	142
250K – 1 Mil. Pop.	699	544	269	183	156	117
< 250K Pop.	439	353	181	122	118	87
Nonmetro	507	461	182	166	115	121
Urbanized	327	331	116	121	72	81
Less Urbanized	390	351	121	121	77	96
Completely Rural	176	154	81	37	48	25

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.62D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.35	0.28	0.19	0.15	0.14	0.11
GEOGRAPHIC DIVISION						
Northeast	0.79	0.67	0.40	0.35	0.29	0.27
New England	1.41	1.28	0.76	0.71	0.47	0.55
Middle Atlantic	0.94	0.77	0.47	0.40	0.35	0.30
Midwest	0.62	0.58	0.40	0.31	0.30	0.22
East North Central	0.72	0.66	0.43	0.38	0.29	0.26
West North Central	1.16	1.16	0.84	0.56	0.70	0.43
South	0.65	0.45	0.37	0.26	0.24	0.19
South Atlantic	0.96	0.64	0.58	0.34	0.35	0.25
East South Central	1.37	1.10	0.73	0.66	0.55	0.46
West South Central	1.12	0.80	0.57	0.42	0.42	0.33
West	0.79	0.66	0.37	0.31	0.28	0.22
Mountain	1.32	0.95	0.60	0.47	0.47	0.38
Pacific	0.98	0.83	0.47	0.40	0.34	0.28
COUNTY TYPE						
Large Metro	0.53	0.41	0.27	0.22	0.21	0.16
Small Metro	0.60	0.49	0.37	0.26	0.23	0.17
250K – 1 Mil. Pop.	0.76	0.62	0.44	0.32	0.28	0.21
< 250K Pop.	1.00	0.81	0.63	0.44	0.42	0.33
Nonmetro	0.79	0.64	0.39	0.34	0.26	0.26
Urbanized	1.08	1.13	0.56	0.57	0.38	0.41
Less Urbanized	1.27	0.90	0.57	0.49	0.38	0.41
Completely Rural	2.24	2.05	1.36	0.82	0.90	0.60

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.62P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5170		0.0219		0.1856	
GEOGRAPHIC DIVISION						
Northeast	0.4288		0.7980		0.5551	
New England	0.4765		0.7995		0.3141	
Middle Atlantic	0.1916		0.8833		0.9116	
Midwest	0.7866		0.2725		0.1886	
East North Central	0.5776		0.5978		0.8848	
West North Central	0.7667		0.2625		0.0702	
South	0.1465		0.0221		0.2156	
South Atlantic	0.4552		0.2404		0.7877	
East South Central	0.4483		0.5681		0.5650	
West South Central	0.0380		0.0018		0.0263	
West	0.8083		0.5328		0.7957	
Mountain	0.4299		0.2337		0.7144	
Pacific	0.8743		0.9509		0.5992	
COUNTY TYPE						
Large Metro	0.3917		0.0435		0.1843	
Small Metro	0.5938		0.1282		0.1355	
250K – 1 Mil. Pop.	0.4256		0.2381		0.2485	
< 250K Pop.	0.7784		0.3218		0.3648	
Nonmetro	0.2939		0.7155		0.1649	
Urbanized	0.9543		0.2035		0.0667	
Less Urbanized	0.1839		0.6271		0.2774	
Completely Rural	0.7511		0.0148		0.0591	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.63C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	64	68	50	57	32	34
GEOGRAPHIC DIVISION						
Northeast	24	28	18	24	14	13
New England	12	13	8	12	6	7
Middle Atlantic	20	24	16	20	12	11
Midwest	29	30	22	26	14	16
East North Central	24	26	18	21	11	13
West North Central	15	14	12	13	8	8
South	40	44	34	38	22	23
South Atlantic	29	30	24	27	15	15
East South Central	14	16	12	13	9	7
West South Central	24	28	19	22	14	15
West	30	31	25	25	14	12
Mountain	17	16	14	13	8	9
Pacific	25	26	20	22	11	9
COUNTY TYPE						
Large Metro	50	52	42	43	25	25
Small Metro	40	36	28	29	17	16
250K – 1 Mil. Pop.	32	29	25	24	15	12
< 250K Pop.	24	22	13	17	8	11
Nonmetro	29	26	23	23	14	14
Urbanized	17	17	14	15	9	9
Less Urbanized	21	19	16	17	11	11
Completely Rural	10	8	7	7	4	4

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.63D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.26	0.27	0.20	0.23	0.13	0.13
GEOGRAPHIC DIVISION						
Northeast	0.56	0.66	0.43	0.57	0.33	0.32
New England	1.14	1.18	0.73	1.09	0.57	0.63
Middle Atlantic	0.65	0.78	0.53	0.64	0.40	0.37
Midwest	0.53	0.56	0.40	0.48	0.26	0.29
East North Central	0.64	0.70	0.48	0.56	0.30	0.36
West North Central	0.90	0.84	0.73	0.81	0.51	0.47
South	0.43	0.47	0.36	0.40	0.24	0.24
South Atlantic	0.62	0.65	0.52	0.59	0.31	0.32
East South Central	0.93	1.10	0.85	0.90	0.60	0.48
West South Central	0.75	0.85	0.60	0.69	0.44	0.46
West	0.51	0.53	0.42	0.42	0.23	0.20
Mountain	0.91	0.87	0.73	0.69	0.45	0.47
Pacific	0.61	0.65	0.50	0.54	0.27	0.21
COUNTY TYPE						
Large Metro	0.34	0.36	0.29	0.30	0.18	0.18
Small Metro	0.46	0.48	0.35	0.39	0.22	0.23
250K – 1 Mil. Pop.	0.51	0.58	0.44	0.47	0.29	0.25
< 250K Pop.	0.96	0.89	0.54	0.71	0.32	0.47
Nonmetro	0.66	0.64	0.52	0.58	0.36	0.37
Urbanized	0.89	0.96	0.74	0.92	0.50	0.58
Less Urbanized	1.06	0.94	0.85	0.85	0.58	0.56
Completely Rural	1.91	1.95	1.43	1.64	0.93	0.97

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.63P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.3729		0.9712		0.2399	
GEOGRAPHIC DIVISION						
Northeast	0.4645		0.4944		0.4613	
New England	0.5561		0.0596		0.6766	
Middle Atlantic	0.2342		0.7986		0.2801	
Midwest	0.4511		0.3691		0.2930	
East North Central	0.5329		0.2144		0.1754	
West North Central	0.6519		0.8289		0.8925	
South	0.9854		0.5488		0.8300	
South Atlantic	0.7746		0.5022		0.5415	
East South Central	0.6817		0.9655		0.4208	
West South Central	0.9005		0.8228		0.8768	
West	0.5537		0.6197		0.1697	
Mountain	0.1008		0.3536		0.5916	
Pacific	0.7314		0.9878		0.1767	
COUNTY TYPE						
Large Metro	0.6902		0.9357		0.3394	
Small Metro	0.6953		0.6340		0.9697	
250K – 1 Mil. Pop.	0.7984		0.9793		0.3665	
< 250K Pop.	0.7639		0.3974		0.2044	
Nonmetro	0.4173		0.6065		0.3608	
Urbanized	0.6653		0.7936		0.8651	
Less Urbanized	0.5361		0.6899		0.3215	
Completely Rural	0.8525		0.8592		0.7665	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.64C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	171	189	142	151	100	105
GEOGRAPHIC DIVISION						
Northeast	69	74	56	57	40	41
New England	33	39	26	26	20	18
Middle Atlantic	60	62	50	49	35	37
Midwest	65	82	57	68	36	47
East North Central	53	70	48	55	30	40
West North Central	35	47	31	39	21	25
South	103	111	86	90	64	68
South Atlantic	81	77	68	58	52	40
East South Central	37	44	30	41	23	31
West South Central	47	63	42	53	31	43
West	91	101	73	78	50	50
Mountain	35	43	29	32	21	24
Pacific	85	90	69	71	46	44
COUNTY TYPE						
Large Metro	203	194	128	136	84	90
Small Metro	159	144	103	91	58	57
250K – 1 Mil. Pop.	116	113	78	72	42	45
< 250K Pop.	117	87	72	55	42	34
Nonmetro	98	104	68	69	40	41
Urbanized	78	82	54	55	30	31
Less Urbanized	59	65	41	43	26	26
Completely Rural	30	22	17	10	8	7

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.64D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.49	0.54	0.41	0.43	0.29	0.30
GEOGRAPHIC DIVISION						
Northeast	1.12	1.20	0.92	0.93	0.65	0.67
New England	2.01	2.40	1.60	1.56	1.25	1.12
Middle Atlantic	1.32	1.36	1.10	1.08	0.76	0.82
Midwest	0.87	1.11	0.76	0.91	0.49	0.63
East North Central	1.04	1.36	0.93	1.08	0.59	0.79
West North Central	1.51	2.01	1.35	1.69	0.91	1.07
South	0.80	0.86	0.67	0.70	0.50	0.52
South Atlantic	1.24	1.17	1.04	0.88	0.80	0.61
East South Central	1.82	2.14	1.45	2.00	1.13	1.49
West South Central	1.11	1.46	0.98	1.23	0.72	0.99
West	1.09	1.20	0.88	0.93	0.60	0.60
Mountain	1.40	1.69	1.16	1.25	0.82	0.95
Pacific	1.45	1.54	1.18	1.22	0.79	0.75
COUNTY TYPE						
Large Metro	0.70	0.77	0.58	0.62	0.42	0.44
Small Metro	0.87	0.85	0.70	0.70	0.46	0.49
250K – 1 Mil. Pop.	1.09	1.06	0.86	0.88	0.53	0.60
< 250K Pop.	1.48	1.46	1.24	1.16	0.88	0.85
Nonmetro	1.05	1.12	0.88	0.99	0.63	0.69
Urbanized	1.53	1.94	1.28	1.67	0.93	1.08
Less Urbanized	1.60	1.44	1.32	1.33	0.96	0.93
Completely Rural	3.46	3.64	2.56	2.33	1.57	1.76

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.64P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.2966		0.1950		0.5116	
GEOGRAPHIC DIVISION						
Northeast	0.6995		0.5705		0.8448	
New England	0.4920		0.5120		0.2793	
Middle Atlantic	0.9593		0.7510		0.7080	
Midwest	0.5529		0.7669		0.8163	
East North Central	0.7085		0.6564		0.8303	
West North Central	0.6174		0.2436		0.9288	
South	0.3498		0.5483		0.4631	
South Atlantic	0.5142		0.6726		0.1752	
East South Central	0.2939		0.7911		0.6510	
West South Central	0.9287		0.4568		0.4594	
West	0.2808		0.1044		0.4662	
Mountain	0.5338		0.2835		0.4380	
Pacific	0.3618		0.1973		0.2296	
COUNTY TYPE						
Large Metro	0.6038		0.4409		0.7469	
Small Metro	0.2983		0.4845		0.9740	
250K – 1 Mil. Pop.	0.1928		0.5002		0.4270	
< 250K Pop.	0.9777		0.7951		0.3277	
Nonmetro	0.8920		0.3641		0.2684	
Urbanized	0.3960		0.2183		0.3971	
Less Urbanized	0.7126		0.8927		0.4224	
Completely Rural	0.2709		0.6064		0.9013	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.65C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	885	705	469	345	335	256
GEOGRAPHIC DIVISION						
Northeast	359	302	180	153	128	113
New England	166	150	86	77	53	61
Middle Atlantic	320	257	158	132	117	95
Midwest	330	304	210	156	166	118
East North Central	265	242	154	134	111	95
West North Central	190	187	135	85	117	70
South	617	423	347	220	219	171
South Atlantic	494	319	294	162	166	126
East South Central	202	166	99	92	78	66
West South Central	327	238	164	114	123	93
West	455	380	211	172	163	129
Mountain	236	171	106	81	88	65
Pacific	394	329	184	150	138	110
COUNTY TYPE						
Large Metro	982	761	401	308	283	221
Small Metro	734	576	294	185	182	126
250K – 1 Mil. Pop.	651	506	246	159	149	105
< 250K Pop.	403	314	160	99	107	75
Nonmetro	477	406	156	135	103	109
Urbanized	305	282	95	91	63	67
Less Urbanized	363	314	103	101	67	86
Completely Rural	158	138	74	37	45	27

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.65D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.44	0.34	0.23	0.17	0.17	0.12
GEOGRAPHIC DIVISION						
Northeast	0.97	0.81	0.49	0.41	0.35	0.30
New England	1.70	1.52	0.88	0.79	0.54	0.62
Middle Atlantic	1.17	0.93	0.58	0.48	0.43	0.35
Midwest	0.76	0.70	0.48	0.36	0.38	0.27
East North Central	0.88	0.80	0.51	0.44	0.37	0.31
West North Central	1.43	1.40	1.02	0.64	0.88	0.53
South	0.82	0.55	0.46	0.29	0.29	0.22
South Atlantic	1.23	0.78	0.73	0.40	0.41	0.31
East South Central	1.69	1.38	0.82	0.76	0.65	0.55
West South Central	1.42	1.01	0.71	0.48	0.53	0.40
West	0.97	0.79	0.45	0.36	0.35	0.27
Mountain	1.65	1.18	0.74	0.56	0.61	0.45
Pacific	1.21	0.99	0.56	0.45	0.42	0.33
COUNTY TYPE						
Large Metro	0.67	0.49	0.34	0.25	0.26	0.18
Small Metro	0.76	0.61	0.46	0.29	0.29	0.20
250K – 1 Mil. Pop.	0.94	0.77	0.55	0.37	0.34	0.24
< 250K Pop.	1.31	1.01	0.77	0.50	0.52	0.38
Nonmetro	0.99	0.77	0.44	0.39	0.30	0.32
Urbanized	1.36	1.31	0.61	0.63	0.42	0.47
Less Urbanized	1.56	1.09	0.65	0.57	0.43	0.50
Completely Rural	2.68	2.29	1.69	1.05	1.12	0.79

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.65P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.6611		0.0462		0.2945	
GEOGRAPHIC DIVISION						
Northeast	0.3468		0.6934		0.5061	
New England	0.5051		0.7981		0.4748	
Middle Atlantic	0.1505		0.7573		0.7095	
Midwest	0.8535		0.2410		0.1843	
East North Central	0.5973		0.8084		0.9510	
West North Central	0.6873		0.1077		0.0560	
South	0.1829		0.0203		0.2717	
South Atlantic	0.5283		0.2428		0.5030	
East South Central	0.3271		0.5786		0.7134	
West South Central	0.0291		0.0015		0.0085	
West	0.9855		0.9652		0.9180	
Mountain	0.5539		0.4336		0.8276	
Pacific	0.7519		0.6241		0.9891	
COUNTY TYPE						
Large Metro	0.4365		0.0645		0.2427	
Small Metro	0.6901		0.1448		0.1159	
250K – 1 Mil. Pop.	0.5296		0.2838		0.1635	
< 250K Pop.	0.7985		0.2886		0.4681	
Nonmetro	0.2385		0.4736		0.0531	
Urbanized	0.7790		0.0694		0.0220	
Less Urbanized	0.1680		0.6028		0.1262	
Completely Rural	0.9300		0.0275		0.0828	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.66C Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	913	750	492	389	354	280
GEOGRAPHIC DIVISION						
Northeast	368	319	189	168	133	126
New England	173	160	95	87	58	69
Middle Atlantic	327	271	163	141	120	105
Midwest	341	322	223	172	170	124
East North Central	275	254	166	146	114	101
West North Central	197	200	143	96	120	73
South	623	437	359	247	234	183
South Atlantic	492	332	303	175	181	129
East South Central	209	172	111	103	85	73
West South Central	338	243	173	128	128	102
West	477	407	224	193	168	140
Mountain	243	179	111	90	88	72
Pacific	415	357	196	170	144	119
COUNTY TYPE						
Large Metro	1,034	813	425	342	296	242
Small Metro	775	621	317	216	191	140
250K – 1 Mil. Pop.	688	539	266	182	155	117
< 250K Pop.	433	348	179	118	117	85
Nonmetro	502	455	178	162	114	121
Urbanized	324	328	114	116	71	80
Less Urbanized	385	345	118	118	76	95
Completely Rural	171	152	80	36	47	25

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.66D Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.38	0.31	0.21	0.16	0.15	0.12
GEOGRAPHIC DIVISION						
Northeast	0.85	0.73	0.44	0.39	0.31	0.29
New England	1.52	1.39	0.83	0.76	0.51	0.60
Middle Atlantic	1.03	0.85	0.51	0.44	0.38	0.33
Midwest	0.67	0.63	0.44	0.34	0.33	0.24
East North Central	0.78	0.72	0.47	0.41	0.32	0.28
West North Central	1.26	1.28	0.92	0.61	0.77	0.46
South	0.71	0.49	0.41	0.28	0.27	0.20
South Atlantic	1.05	0.70	0.65	0.37	0.39	0.27
East South Central	1.49	1.22	0.79	0.73	0.60	0.52
West South Central	1.24	0.87	0.63	0.46	0.47	0.37
West	0.86	0.72	0.40	0.34	0.30	0.25
Mountain	1.44	1.05	0.66	0.53	0.53	0.42
Pacific	1.08	0.91	0.51	0.44	0.37	0.30
COUNTY TYPE						
Large Metro	0.58	0.45	0.30	0.23	0.23	0.17
Small Metro	0.65	0.54	0.41	0.28	0.25	0.19
250K – 1 Mil. Pop.	0.82	0.67	0.49	0.35	0.30	0.23
< 250K Pop.	1.08	0.90	0.69	0.48	0.47	0.35
Nonmetro	0.87	0.70	0.42	0.37	0.28	0.29
Urbanized	1.18	1.23	0.61	0.61	0.40	0.44
Less Urbanized	1.39	0.98	0.62	0.54	0.41	0.46
Completely Rural	2.46	2.22	1.52	0.89	1.00	0.65

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.66P Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.5161		0.0199		0.2215	
GEOGRAPHIC DIVISION						
Northeast	0.4108		0.8653		0.5050	
New England	0.4347		0.9641		0.3324	
Middle Atlantic	0.1688		0.8641		0.8260	
Midwest	0.7586		0.3141		0.2200	
East North Central	0.5545		0.7013		0.9969	
West North Central	0.7805		0.2468		0.0649	
South	0.1352		0.0172		0.2019	
South Atlantic	0.4534		0.2108		0.8350	
East South Central	0.4700		0.5627		0.6211	
West South Central	0.0326		0.0013		0.0235	
West	0.8011		0.5584		0.8744	
Mountain	0.4918		0.2848		0.6754	
Pacific	0.9281		0.9395		0.6590	
COUNTY TYPE						
Large Metro	0.3911		0.0392		0.2103	
Small Metro	0.5337		0.1102		0.1286	
250K – 1 Mil. Pop.	0.3757		0.2290		0.2782	
< 250K Pop.	0.8079		0.2684		0.2933	
Nonmetro	0.2597		0.6600		0.1327	
Urbanized	0.9029		0.1817		0.0595	
Less Urbanized	0.1590		0.5809		0.2207	
Completely Rural	0.8063		0.0163		0.0633	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.67C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	801	635	1,007	833	1,076	869
GEOGRAPHIC DIVISION						
Northeast	312	232	394	321	430	354
New England	109	105	175	161	211	172
Middle Atlantic	290	205	355	276	375	310
Midwest	240	228	359	330	388	346
East North Central	194	188	299	286	317	306
West North Central	139	118	212	157	233	178
South	513	401	664	533	683	565
South Atlantic	388	298	515	388	544	439
East South Central	204	148	254	195	246	200
West South Central	260	226	318	305	322	311
West	458	345	544	438	580	468
Mountain	195	140	249	179	278	215
Pacific	408	311	480	395	502	411
COUNTY TYPE						
Large Metro	1,811	1,528	1,590	1,350	1,396	1,202
Small Metro	1,454	1,184	1,245	999	1,053	837
250K – 1 Mil. Pop.	1,230	969	1,079	830	939	716
< 250K Pop.	909	754	747	617	599	488
Nonmetro	978	941	792	733	670	591
Urbanized	669	689	558	526	463	428
Less Urbanized	760	717	605	578	505	465
Completely Rural	353	309	249	206	195	154

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.67D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.31	0.24	0.38	0.31	0.41	0.33
GEOGRAPHIC DIVISION						
Northeast	0.66	0.49	0.83	0.67	0.91	0.74
New England	0.87	0.83	1.40	1.28	1.69	1.37
Middle Atlantic	0.83	0.58	1.02	0.79	1.07	0.89
Midwest	0.43	0.40	0.64	0.59	0.69	0.61
East North Central	0.50	0.48	0.77	0.73	0.81	0.78
West North Central	0.81	0.68	1.23	0.90	1.35	1.02
South	0.53	0.41	0.68	0.54	0.70	0.57
South Atlantic	0.75	0.57	1.00	0.74	1.06	0.84
East South Central	1.32	0.95	1.64	1.25	1.58	1.28
West South Central	0.85	0.73	1.04	0.98	1.06	1.00
West	0.75	0.56	0.89	0.70	0.95	0.75
Mountain	1.04	0.74	1.33	0.94	1.49	1.13
Pacific	0.96	0.72	1.13	0.91	1.18	0.95
COUNTY TYPE						
Large Metro	0.44	0.34	0.53	0.45	0.56	0.48
Small Metro	0.48	0.40	0.67	0.55	0.74	0.59
250K – 1 Mil. Pop.	0.58	0.53	0.81	0.70	0.88	0.74
< 250K Pop.	0.87	0.61	1.33	0.89	1.43	0.94
Nonmetro	0.74	0.56	0.92	0.74	0.94	0.75
Urbanized	1.09	0.80	1.30	1.01	1.30	1.06
Less Urbanized	1.10	0.89	1.40	1.14	1.47	1.15
Completely Rural	2.12	1.61	2.88	2.20	2.69	2.29

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.67P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0901		0.5535		0.3203	
GEOGRAPHIC DIVISION						
Northeast	0.2862		0.3673		0.7947	
New England	0.7271		0.9356		0.5414	
Middle Atlantic	0.1885		0.3295		0.5232	
Midwest	0.8113		0.4320		0.5215	
East North Central	0.8905		0.3601		0.7027	
West North Central	0.8185		0.9864		0.5724	
South	0.2102		0.8913		0.5282	
South Atlantic	0.4154		0.7161		0.6014	
East South Central	0.4672		0.6798		0.6048	
West South Central	0.5011		0.2673		0.9552	
West	0.2897		0.2220		0.1623	
Mountain	0.4098		0.6537		0.5405	
Pacific	0.1168		0.1108		0.0528	
COUNTY TYPE						
Large Metro	0.0819		0.1824		0.1881	
Small Metro	0.8259		0.7931		0.8941	
250K – 1 Mil. Pop.	0.8028		0.2633		0.4252	
< 250K Pop.	0.4328		0.2604		0.3722	
Nonmetro	0.5181		0.2460		0.7822	
Urbanized	0.2842		0.5584		0.8245	
Less Urbanized	0.9492		0.6146		0.8836	
Completely Rural	0.7757		0.2902		0.1970	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.68C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	109	107	100	103	73	81
GEOGRAPHIC DIVISION						
Northeast	40	45	36	46	30	34
New England	19	21	17	21	15	13
Middle Atlantic	35	40	32	41	26	31
Midwest	45	47	42	42	30	33
East North Central	36	42	35	37	24	27
West North Central	25	24	23	22	17	17
South	68	70	63	64	46	48
South Atlantic	51	47	48	43	36	33
East South Central	20	23	18	23	13	17
West South Central	42	47	38	42	26	30
West	59	56	52	58	39	47
Mountain	31	28	29	28	21	25
Pacific	50	47	44	49	33	39
COUNTY TYPE						
Large Metro	113	114	94	103	64	73
Small Metro	87	70	74	65	44	40
250K – 1 Mil. Pop.	72	57	63	51	37	35
< 250K Pop.	50	43	39	41	23	21
Nonmetro	54	54	42	47	27	31
Urbanized	31	37	26	31	18	19
Less Urbanized	40	41	32	36	19	23
Completely Rural	23	16	18	12	10	10

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.68D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.44	0.43	0.40	0.42	0.29	0.33
GEOGRAPHIC DIVISION						
Northeast	0.95	1.09	0.86	1.10	0.72	0.81
New England	1.73	1.94	1.57	1.96	1.37	1.21
Middle Atlantic	1.12	1.31	1.02	1.34	0.84	1.01
Midwest	0.83	0.87	0.77	0.78	0.55	0.61
East North Central	0.97	1.12	0.92	1.00	0.64	0.73
West North Central	1.54	1.44	1.38	1.33	1.06	1.06
South	0.73	0.74	0.67	0.68	0.49	0.51
South Atlantic	1.10	1.01	1.03	0.91	0.77	0.70
East South Central	1.32	1.58	1.21	1.58	0.86	1.16
West South Central	1.29	1.44	1.18	1.30	0.81	0.93
West	0.99	0.94	0.88	0.97	0.66	0.79
Mountain	1.65	1.51	1.52	1.48	1.12	1.31
Pacific	1.22	1.17	1.07	1.22	0.80	0.97
COUNTY TYPE						
Large Metro	0.62	0.61	0.55	0.59	0.42	0.48
Small Metro	0.78	0.78	0.73	0.76	0.51	0.52
250K – 1 Mil. Pop.	0.92	0.90	0.88	0.83	0.62	0.63
< 250K Pop.	1.45	1.57	1.22	1.61	0.88	0.90
Nonmetro	0.96	1.09	0.85	1.02	0.65	0.74
Urbanized	1.39	1.89	1.26	1.70	1.02	1.12
Less Urbanized	1.41	1.48	1.27	1.39	0.90	1.07
Completely Rural	2.74	3.25	2.57	2.59	1.86	2.28

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.68P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0585		0.2985		0.9818	
GEOGRAPHIC DIVISION						
Northeast	0.6297		0.4283		0.9264	
New England	0.6539		0.3968		0.3649	
Middle Atlantic	0.7578		0.6582		0.5835	
Midwest	0.0330		0.0162		0.1713	
East North Central	0.0773		0.0516		0.0547	
West North Central	0.2473		0.1765		0.6971	
South	0.0243		0.3353		0.8348	
South Atlantic	0.0064		0.2965		0.4710	
East South Central	0.3858		0.3706		0.6512	
West South Central	0.9645		0.9167		0.4446	
West	0.6113		0.6547		0.2635	
Mountain	0.1010		0.3324		0.4884	
Pacific	0.1081		0.2649		0.3726	
COUNTY TYPE						
Large Metro	0.3024		0.4367		0.7844	
Small Metro	0.1309		0.4463		0.5877	
250K – 1 Mil. Pop.	0.5177		0.9529		0.9434	
< 250K Pop.	0.0999		0.2335		0.3663	
Nonmetro	0.4864		0.9377		0.9909	
Urbanized	0.4284		0.8498		0.4337	
Less Urbanized	0.8911		0.8841		0.6446	
Completely Rural	0.7320		0.7703		0.5815	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.69C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	136	141	158	165	183	197
GEOGRAPHIC DIVISION						
Northeast	46	51	57	61	69	78
New England	20	24	23	25	33	34
Middle Atlantic	41	46	51	55	59	70
Midwest	42	59	49	70	65	83
East North Central	34	51	40	60	53	75
West North Central	25	30	29	38	35	42
South	91	86	105	100	122	118
South Atlantic	71	57	81	66	91	82
East South Central	33	35	40	40	45	45
West South Central	46	57	51	66	62	72
West	77	83	86	93	90	109
Mountain	34	39	39	43	42	49
Pacific	70	74	76	83	80	96
COUNTY TYPE						
Large Metro	351	320	340	308	294	273
Small Metro	263	248	246	237	204	207
250K – 1 Mil. Pop.	196	191	185	184	157	168
< 250K Pop.	191	154	181	144	144	121
Nonmetro	168	172	154	160	121	132
Urbanized	136	134	126	125	95	103
Less Urbanized	106	112	96	102	81	81
Completely Rural	46	39	43	35	33	27

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.69D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.39	0.40	0.45	0.47	0.53	0.56
GEOGRAPHIC DIVISION						
Northeast	0.75	0.83	0.92	0.99	1.12	1.27
New England	1.23	1.45	1.39	1.52	2.00	2.06
Middle Atlantic	0.90	1.01	1.13	1.22	1.31	1.54
Midwest	0.57	0.80	0.67	0.95	0.87	1.12
East North Central	0.66	0.99	0.79	1.17	1.04	1.47
West North Central	1.11	1.30	1.25	1.65	1.52	1.80
South	0.71	0.66	0.81	0.77	0.95	0.91
South Atlantic	1.08	0.86	1.23	1.00	1.39	1.25
East South Central	1.62	1.69	1.96	1.97	2.21	2.17
West South Central	1.09	1.31	1.21	1.52	1.45	1.68
West	0.92	0.99	1.03	1.11	1.08	1.29
Mountain	1.33	1.54	1.53	1.66	1.64	1.90
Pacific	1.19	1.26	1.31	1.42	1.37	1.65
COUNTY TYPE						
Large Metro	0.57	0.57	0.68	0.65	0.77	0.79
Small Metro	0.67	0.71	0.75	0.80	0.88	0.99
250K – 1 Mil. Pop.	0.85	0.92	0.96	0.99	1.12	1.25
< 250K Pop.	1.09	1.13	1.26	1.39	1.41	1.64
Nonmetro	0.87	0.93	0.94	1.16	1.06	1.23
Urbanized	1.16	1.39	1.34	1.89	1.49	1.96
Less Urbanized	1.28	1.30	1.42	1.53	1.77	1.69
Completely Rural	3.41	3.08	3.67	3.11	3.61	3.72

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.69P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.4537		0.6344		0.9625	
GEOGRAPHIC DIVISION						
Northeast	0.9823		0.9556		0.7844	
New England	0.6075		0.7115		0.7012	
Middle Atlantic	0.7812		0.8111		0.9176	
Midwest	0.2910		0.1478		0.2366	
East North Central	0.2399		0.0840		0.1108	
West North Central	0.8976		1.0000		0.6902	
South	0.9876		0.9374		0.8423	
South Atlantic	0.6889		0.7933		0.6843	
East South Central	0.7886		0.8242		0.6153	
West South Central	0.7291		0.7326		0.8701	
West	0.5462		0.9306		0.3955	
Mountain	0.3468		0.9963		0.7969	
Pacific	0.8439		0.9236		0.4066	
COUNTY TYPE						
Large Metro	0.5634		0.7437		0.9878	
Small Metro	0.3717		0.8705		0.8163	
250K – 1 Mil. Pop.	0.1892		0.7411		0.9649	
< 250K Pop.	0.7314		0.8552		0.7310	
Nonmetro	0.5781		0.6890		0.8484	
Urbanized	0.3478		0.6304		0.5818	
Less Urbanized	0.3832		0.8246		0.6094	
Completely Rural	0.1502		0.9223		0.7221	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.70C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	736	553	954	764	1,021	802
GEOGRAPHIC DIVISION						
Northeast	294	210	381	303	414	339
New England	93	89	167	154	196	162
Middle Atlantic	276	186	343	258	363	297
Midwest	217	204	345	310	379	318
East North Central	171	164	286	269	309	282
West North Central	128	112	206	148	227	167
South	480	354	626	498	647	527
South Atlantic	371	263	491	357	524	407
East South Central	194	133	243	185	232	189
West South Central	240	193	295	281	299	287
West	420	292	521	393	550	418
Mountain	174	113	232	168	258	201
Pacific	377	263	463	351	480	361
COUNTY TYPE						
Large Metro	1,682	1,420	1,463	1,245	1,298	1,129
Small Metro	1,319	1,086	1,119	905	974	768
250K – 1 Mil. Pop.	1,105	902	966	765	861	665
< 250K Pop.	817	670	661	542	549	434
Nonmetro	892	831	725	635	632	518
Urbanized	608	605	508	453	434	374
Less Urbanized	682	625	544	496	467	411
Completely Rural	314	276	216	180	173	136

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.70D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.36	0.27	0.47	0.37	0.50	0.39
GEOGRAPHIC DIVISION						
Northeast	0.79	0.56	1.03	0.81	1.12	0.91
New England	0.95	0.90	1.71	1.56	2.01	1.64
Middle Atlantic	1.01	0.68	1.26	0.94	1.33	1.08
Midwest	0.50	0.47	0.80	0.71	0.87	0.73
East North Central	0.57	0.54	0.95	0.89	1.03	0.93
West North Central	0.97	0.83	1.55	1.10	1.71	1.25
South	0.64	0.46	0.83	0.65	0.86	0.69
South Atlantic	0.92	0.64	1.22	0.87	1.30	1.00
East South Central	1.62	1.10	2.03	1.54	1.94	1.56
West South Central	1.04	0.82	1.28	1.20	1.30	1.22
West	0.89	0.61	1.11	0.82	1.17	0.87
Mountain	1.22	0.77	1.62	1.15	1.80	1.38
Pacific	1.15	0.79	1.42	1.06	1.47	1.09
COUNTY TYPE						
Large Metro	0.54	0.38	0.66	0.53	0.69	0.57
Small Metro	0.56	0.44	0.83	0.66	0.93	0.70
250K – 1 Mil. Pop.	0.66	0.58	0.99	0.84	1.08	0.88
< 250K Pop.	1.06	0.65	1.66	1.04	1.87	1.10
Nonmetro	0.91	0.65	1.13	0.88	1.14	0.90
Urbanized	1.35	0.87	1.63	1.19	1.60	1.27
Less Urbanized	1.36	1.05	1.71	1.38	1.79	1.40
Completely Rural	2.55	2.05	3.51	2.72	3.09	2.69

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.70P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0301		0.4515		0.3269	
GEOGRAPHIC DIVISION						
Northeast	0.3283		0.4202		0.8699	
New England	0.6549		0.9177		0.4953	
Middle Atlantic	0.2179		0.3331		0.5785	
Midwest	0.7252		0.7618		0.7280	
East North Central	0.6504		0.6575		0.9446	
West North Central	0.9871		0.8957		0.4760	
South	0.0960		0.9368		0.4862	
South Atlantic	0.2465		0.6678		0.5130	
East South Central	0.3559		0.6420		0.6442	
West South Central	0.4232		0.2456		0.9949	
West	0.2628		0.2415		0.2412	
Mountain	0.6668		0.7145		0.4262	
Pacific	0.1500		0.1349		0.0752	
COUNTY TYPE						
Large Metro	0.0347		0.1214		0.1791	
Small Metro	0.6683		0.7499		0.8433	
250K – 1 Mil. Pop.	0.9091		0.2077		0.3362	
< 250K Pop.	0.3461		0.2129		0.3224	
Nonmetro	0.4772		0.2454		0.7362	
Urbanized	0.2280		0.6162		0.8411	
Less Urbanized	0.8896		0.6173		0.8313	
Completely Rural	0.8622		0.2673		0.1230	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.71C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	760	584	984	800	1,052	843
GEOGRAPHIC DIVISION						
Northeast	301	217	390	311	425	348
New England	97	96	171	161	205	170
Middle Atlantic	282	191	351	263	372	303
Midwest	222	208	354	321	389	339
East North Central	174	171	294	279	318	302
West North Central	133	109	211	151	232	172
South	494	373	652	517	671	551
South Atlantic	380	277	509	372	539	428
East South Central	197	135	250	189	240	198
West South Central	248	209	308	296	313	302
West	432	314	528	418	560	450
Mountain	180	125	242	174	269	207
Pacific	388	284	467	376	485	394
COUNTY TYPE						
Large Metro	1,784	1,503	1,567	1,329	1,385	1,190
Small Metro	1,411	1,166	1,211	983	1,039	828
250K – 1 Mil. Pop.	1,193	956	1,051	819	926	708
< 250K Pop.	883	739	725	605	590	483
Nonmetro	955	919	778	717	665	583
Urbanized	656	676	550	516	460	424
Less Urbanized	738	693	588	557	498	454
Completely Rural	341	301	242	201	192	150

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.71D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.32	0.24	0.41	0.33	0.44	0.35
GEOGRAPHIC DIVISION						
Northeast	0.70	0.50	0.90	0.72	0.98	0.80
New England	0.85	0.83	1.50	1.40	1.80	1.48
Middle Atlantic	0.89	0.60	1.10	0.82	1.17	0.95
Midwest	0.44	0.41	0.70	0.63	0.76	0.66
East North Central	0.49	0.48	0.83	0.79	0.90	0.85
West North Central	0.85	0.69	1.35	0.96	1.49	1.10
South	0.56	0.42	0.74	0.58	0.76	0.62
South Atlantic	0.81	0.58	1.09	0.78	1.15	0.90
East South Central	1.40	0.95	1.78	1.34	1.71	1.40
West South Central	0.91	0.75	1.13	1.06	1.15	1.09
West	0.78	0.56	0.95	0.74	1.01	0.80
Mountain	1.07	0.73	1.44	1.02	1.60	1.21
Pacific	1.01	0.73	1.21	0.96	1.26	1.01
COUNTY TYPE						
Large Metro	0.47	0.34	0.58	0.47	0.61	0.51
Small Metro	0.49	0.40	0.73	0.59	0.81	0.63
250K – 1 Mil. Pop.	0.59	0.53	0.87	0.75	0.95	0.79
< 250K Pop.	0.91	0.61	1.45	0.93	1.57	1.00
Nonmetro	0.81	0.58	1.00	0.79	1.01	0.82
Urbanized	1.19	0.79	1.43	1.05	1.42	1.13
Less Urbanized	1.20	0.93	1.50	1.23	1.59	1.26
Completely Rural	2.30	1.76	3.19	2.47	2.89	2.51

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.71P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0520		0.5307		0.3529	
GEOGRAPHIC DIVISION						
Northeast	0.3368		0.4387		0.8337	
New England	0.5889		0.9630		0.5596	
Middle Atlantic	0.2087		0.3664		0.5738	
Midwest	0.9236		0.5770		0.5637	
East North Central	0.8805		0.4599		0.7877	
West North Central	0.9640		0.8981		0.5343	
South	0.1082		0.9413		0.5320	
South Atlantic	0.2291		0.6539		0.5784	
East South Central	0.3924		0.6231		0.5859	
West South Central	0.4933		0.2377		0.9733	
West	0.3407		0.2522		0.1961	
Mountain	0.5333		0.7238		0.4734	
Pacific	0.1792		0.1436		0.0608	
COUNTY TYPE						
Large Metro	0.0514		0.1533		0.1943	
Small Metro	0.8373		0.7310		0.8092	
250K – 1 Mil. Pop.	0.6829		0.1997		0.3473	
< 250K Pop.	0.3259		0.2303		0.3765	
Nonmetro	0.4332		0.2444		0.7851	
Urbanized	0.1877		0.6017		0.7372	
Less Urbanized	1.0000		0.6122		0.8904	
Completely Rural	0.9193		0.2597		0.1443	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.72C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	1,076	869	804	690	438	368
GEOGRAPHIC DIVISION						
Northeast	430	354	336	293	159	165
New England	211	172	154	149	84	76
Middle Atlantic	375	310	299	250	135	145
Midwest	388	346	332	282	180	159
East North Central	317	306	270	241	150	130
West North Central	233	178	198	152	93	86
South	683	565	523	412	291	233
South Atlantic	544	439	408	301	238	170
East South Central	246	200	190	148	103	74
West South Central	322	311	281	223	139	145
West	580	468	402	347	231	168
Mountain	278	215	199	152	122	88
Pacific	502	411	348	308	198	139
COUNTY TYPE						
Large Metro	1,396	1,202	782	655	375	311
Small Metro	1,053	837	559	449	238	204
250K – 1 Mil. Pop.	939	716	504	369	211	164
< 250K Pop.	599	488	303	271	118	121
Nonmetro	670	591	351	323	150	143
Urbanized	463	428	242	216	100	93
Less Urbanized	505	465	261	250	101	106
Completely Rural	195	154	117	81	65	39

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.72D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.41	0.33	0.31	0.26	0.17	0.14
GEOGRAPHIC DIVISION						
Northeast	0.91	0.74	0.71	0.61	0.34	0.35
New England	1.69	1.37	1.23	1.19	0.68	0.60
Middle Atlantic	1.07	0.89	0.86	0.71	0.39	0.41
Midwest	0.69	0.61	0.59	0.50	0.32	0.28
East North Central	0.81	0.78	0.69	0.62	0.39	0.33
West North Central	1.35	1.02	1.15	0.87	0.54	0.50
South	0.70	0.57	0.54	0.42	0.30	0.24
South Atlantic	1.06	0.84	0.79	0.58	0.46	0.33
East South Central	1.58	1.28	1.23	0.95	0.66	0.47
West South Central	1.06	1.00	0.92	0.72	0.45	0.47
West	0.95	0.75	0.66	0.56	0.38	0.27
Mountain	1.49	1.13	1.06	0.80	0.65	0.46
Pacific	1.18	0.95	0.82	0.71	0.47	0.32
COUNTY TYPE						
Large Metro	0.56	0.48	0.44	0.37	0.25	0.20
Small Metro	0.74	0.59	0.57	0.44	0.29	0.24
250K – 1 Mil. Pop.	0.88	0.74	0.70	0.53	0.38	0.29
< 250K Pop.	1.43	0.94	0.97	0.75	0.45	0.42
Nonmetro	0.94	0.75	0.67	0.55	0.33	0.30
Urbanized	1.30	1.06	0.99	0.85	0.47	0.47
Less Urbanized	1.47	1.15	1.06	0.80	0.48	0.44
Completely Rural	2.69	2.29	1.91	1.49	1.23	0.87

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.72P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.3203		0.8131		0.5200	
GEOGRAPHIC DIVISION						
Northeast	0.7947		0.8525		0.0891	
New England	0.5414		0.5283		0.9807	
Middle Atlantic	0.5232		0.5759		0.0509	
Midwest	0.5215		0.8629		0.8039	
East North Central	0.7027		0.5285		0.4815	
West North Central	0.5724		0.2257		0.5270	
South	0.5282		0.7877		0.6391	
South Atlantic	0.6014		0.7716		0.7103	
East South Central	0.6048		0.3482		0.6058	
West South Central	0.9552		0.5816		0.9941	
West	0.1623		0.8789		0.0704	
Mountain	0.5405		0.0391		0.0905	
Pacific	0.0528		0.1978		0.2765	
COUNTY TYPE						
Large Metro	0.1881		0.7195		0.5591	
Small Metro	0.8941		0.6493		0.3633	
250K – 1 Mil. Pop.	0.4252		0.4212		0.3536	
< 250K Pop.	0.3722		0.7006		0.8236	
Nonmetro	0.7822		0.6618		0.3338	
Urbanized	0.8245		0.9496		0.5773	
Less Urbanized	0.8836		0.2762		0.0504	
Completely Rural	0.1970		0.2401		0.0439	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.73C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	73	81	54	60	23	24
GEOGRAPHIC DIVISION						
Northeast	30	34	25	26	13	11
New England	15	13	12	8	6	3
Middle Atlantic	26	31	22	24	12	10
Midwest	30	33	21	25	9	11
East North Central	24	27	17	21	8	10
West North Central	17	17	12	13	4	4
South	46	48	33	36	12	16
South Atlantic	36	33	24	24	9	10
East South Central	13	17	9	13	5	4
West South Central	26	30	20	22	7	12
West	39	47	30	34	12	9
Mountain	21	25	14	16	6	5
Pacific	33	39	26	30	10	8
COUNTY TYPE						
Large Metro	64	73	45	50	19	18
Small Metro	44	40	31	28	12	9
250K – 1 Mil. Pop.	37	35	26	24	10	8
< 250K Pop.	23	21	17	14	6	5
Nonmetro	27	31	21	27	9	14
Urbanized	18	19	13	20	4	8
Less Urbanized	19	23	15	18	7	11
Completely Rural	10	10	7	7	3	3

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.73D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.29	0.33	0.22	0.24	0.09	0.10
GEOGRAPHIC DIVISION						
Northeast	0.72	0.81	0.60	0.61	0.32	0.27
New England	1.37	1.21	1.12	0.72	0.51	0.31
Middle Atlantic	0.84	1.01	0.71	0.79	0.39	0.34
Midwest	0.55	0.61	0.39	0.47	0.17	0.20
East North Central	0.64	0.73	0.44	0.57	0.22	0.26
West North Central	1.06	1.06	0.73	0.80	0.24	0.26
South	0.49	0.51	0.35	0.38	0.13	0.17
South Atlantic	0.77	0.70	0.51	0.50	0.20	0.20
East South Central	0.86	1.16	0.62	0.85	0.31	0.25
West South Central	0.81	0.93	0.63	0.69	0.22	0.38
West	0.66	0.79	0.50	0.57	0.20	0.16
Mountain	1.12	1.31	0.77	0.82	0.30	0.26
Pacific	0.80	0.97	0.63	0.74	0.26	0.20
COUNTY TYPE						
Large Metro	0.42	0.48	0.31	0.34	0.14	0.13
Small Metro	0.51	0.52	0.38	0.38	0.16	0.13
250K – 1 Mil. Pop.	0.62	0.63	0.45	0.47	0.19	0.17
< 250K Pop.	0.88	0.90	0.69	0.62	0.27	0.21
Nonmetro	0.65	0.74	0.52	0.68	0.23	0.37
Urbanized	1.02	1.12	0.74	1.23	0.27	0.51
Less Urbanized	0.90	1.07	0.77	0.86	0.39	0.58
Completely Rural	1.86	2.28	1.43	1.83	0.62	0.87

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.73P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.9818		0.5992		0.3006	
GEOGRAPHIC DIVISION						
Northeast	0.9264		0.9081		0.3658	
New England	0.3649		0.0300		0.2185	
Middle Atlantic	0.5835		0.2793		0.6332	
Midwest	0.1713		0.1665		0.5225	
East North Central	0.0547		0.1285		0.3500	
West North Central	0.6971		0.8114		0.6118	
South	0.8348		0.9391		0.2350	
South Atlantic	0.4710		0.9700		0.8931	
East South Central	0.6512		0.5511		0.0955	
West South Central	0.4446		0.8253		0.0135	
West	0.2635		0.9433		0.0240	
Mountain	0.4884		0.8894		0.1383	
Pacific	0.3726		0.9945		0.0813	
COUNTY TYPE						
Large Metro	0.7844		0.9377		0.3566	
Small Metro	0.5877		0.2295		0.0308	
250K – 1 Mil. Pop.	0.9434		0.7397		0.0778	
< 250K Pop.	0.3663		0.0841		0.2019	
Nonmetro	0.9909		0.9758		0.1956	
Urbanized	0.4337		0.9556		0.2683	
Less Urbanized	0.6446		0.9474		0.4992	
Completely Rural	0.5815		0.9194		0.5992	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.74C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	183	197	182	197	107	125
GEOGRAPHIC DIVISION						
Northeast	69	78	69	84	45	57
New England	33	34	33	39	27	25
Middle Atlantic	59	70	60	74	35	51
Midwest	65	83	69	86	45	53
East North Central	53	75	59	79	37	45
West North Central	35	42	34	37	25	29
South	122	118	109	110	64	69
South Atlantic	91	82	80	75	48	43
East South Central	45	45	38	39	23	22
West South Central	62	72	59	72	34	49
West	90	109	98	101	53	60
Mountain	42	49	38	41	22	25
Pacific	80	96	91	92	49	54
COUNTY TYPE						
Large Metro	294	273	217	215	99	108
Small Metro	204	207	152	156	65	73
250K – 1 Mil. Pop.	157	168	110	132	51	58
< 250K Pop.	144	121	110	87	41	43
Nonmetro	121	132	87	101	40	46
Urbanized	95	103	68	75	32	29
Less Urbanized	81	81	60	67	24	33
Completely Rural	33	27	22	22	11	11

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.74D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.53	0.56	0.52	0.57	0.31	0.36
GEOGRAPHIC DIVISION						
Northeast	1.12	1.27	1.13	1.37	0.74	0.93
New England	2.00	2.06	2.02	2.36	1.68	1.55
Middle Atlantic	1.31	1.54	1.33	1.64	0.77	1.14
Midwest	0.87	1.12	0.93	1.16	0.61	0.72
East North Central	1.04	1.47	1.16	1.54	0.72	0.87
West North Central	1.52	1.80	1.46	1.60	1.09	1.26
South	0.95	0.91	0.85	0.85	0.50	0.53
South Atlantic	1.39	1.25	1.22	1.14	0.73	0.65
East South Central	2.21	2.17	1.85	1.90	1.12	1.08
West South Central	1.45	1.68	1.39	1.67	0.81	1.13
West	1.08	1.29	1.17	1.20	0.64	0.71
Mountain	1.64	1.90	1.48	1.62	0.87	1.00
Pacific	1.37	1.65	1.56	1.57	0.84	0.92
COUNTY TYPE						
Large Metro	0.77	0.79	0.76	0.80	0.44	0.51
Small Metro	0.88	0.99	0.84	0.98	0.51	0.58
250K – 1 Mil. Pop.	1.12	1.25	1.01	1.29	0.64	0.72
< 250K Pop.	1.41	1.64	1.43	1.43	0.84	0.95
Nonmetro	1.06	1.23	1.01	1.19	0.65	0.77
Urbanized	1.49	1.96	1.55	1.85	1.03	1.05
Less Urbanized	1.77	1.69	1.59	1.68	0.87	1.20
Completely Rural	3.61	3.72	2.94	4.09	2.21	2.51

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.74P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.9625		0.8263		0.2930	
GEOGRAPHIC DIVISION						
Northeast	0.7844		0.9284		0.6602	
New England	0.7012		0.8205		0.8597	
Middle Atlantic	0.9176		0.9915		0.6762	
Midwest	0.2366		0.9007		0.7907	
East North Central	0.1108		0.7166		0.5233	
West North Central	0.6902		0.3237		0.6374	
South	0.8423		0.7715		0.3438	
South Atlantic	0.6843		0.8737		0.4395	
East South Central	0.6153		0.7234		0.4095	
West South Central	0.8701		0.6421		0.8283	
West	0.3955		0.9119		0.6832	
Mountain	0.7969		0.9293		0.1264	
Pacific	0.4066		0.8749		0.2477	
COUNTY TYPE						
Large Metro	0.9878		0.9841		0.2670	
Small Metro	0.8163		0.4889		0.5059	
250K – 1 Mil. Pop.	0.9649		0.5286		0.5700	
< 250K Pop.	0.7310		0.7608		0.7194	
Nonmetro	0.8484		0.6659		0.5999	
Urbanized	0.5818		0.9408		0.3600	
Less Urbanized	0.6094		0.7441		0.1650	
Completely Rural	0.7221		0.4187		0.4196	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.75C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	1,021	802	766	623	420	334
GEOGRAPHIC DIVISION						
Northeast	414	339	315	271	145	156
New England	196	162	139	131	72	72
Middle Atlantic	363	297	283	233	126	138
Midwest	379	318	311	256	173	148
East North Central	309	282	255	217	144	123
West North Central	227	167	181	138	90	79
South	647	527	500	380	282	214
South Atlantic	524	407	400	283	233	159
East South Central	232	189	175	137	96	69
West South Central	299	287	262	197	134	132
West	550	418	388	306	219	154
Mountain	258	201	190	141	119	79
Pacific	480	361	336	269	186	127
COUNTY TYPE						
Large Metro	1,298	1,129	714	593	353	287
Small Metro	974	768	512	388	227	182
250K – 1 Mil. Pop.	861	665	464	322	203	148
< 250K Pop.	549	434	259	236	107	107
Nonmetro	632	518	314	264	138	128
Urbanized	434	374	217	173	92	82
Less Urbanized	467	411	228	209	91	93
Completely Rural	173	136	104	69	61	36

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.75D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.50	0.39	0.38	0.30	0.21	0.16
GEOGRAPHIC DIVISION						
Northeast	1.12	0.91	0.85	0.73	0.39	0.42
New England	2.01	1.64	1.42	1.34	0.73	0.73
Middle Atlantic	1.33	1.08	1.04	0.85	0.46	0.50
Midwest	0.87	0.73	0.72	0.59	0.40	0.34
East North Central	1.03	0.93	0.85	0.72	0.48	0.40
West North Central	1.71	1.25	1.36	1.03	0.68	0.59
South	0.86	0.69	0.66	0.50	0.37	0.28
South Atlantic	1.30	1.00	0.99	0.69	0.58	0.39
East South Central	1.94	1.56	1.46	1.14	0.80	0.57
West South Central	1.30	1.22	1.14	0.84	0.58	0.56
West	1.17	0.87	0.83	0.64	0.47	0.32
Mountain	1.80	1.38	1.33	0.97	0.83	0.54
Pacific	1.47	1.09	1.03	0.81	0.57	0.38
COUNTY TYPE						
Large Metro	0.69	0.57	0.55	0.44	0.31	0.24
Small Metro	0.93	0.70	0.70	0.52	0.37	0.29
250K – 1 Mil. Pop.	1.08	0.88	0.87	0.63	0.48	0.35
< 250K Pop.	1.87	1.10	1.18	0.90	0.56	0.50
Nonmetro	1.14	0.90	0.80	0.61	0.39	0.36
Urbanized	1.60	1.27	1.21	0.93	0.57	0.56
Less Urbanized	1.79	1.40	1.25	0.94	0.57	0.52
Completely Rural	3.09	2.69	2.26	1.66	1.50	1.01

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.75P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.3269		0.7311		0.7608	
GEOGRAPHIC DIVISION						
Northeast	0.8699		0.8360		0.0403	
New England	0.4953		0.6199		0.8703	
Middle Atlantic	0.5785		0.6216		0.0248	
Midwest	0.7280		0.8989		0.8806	
East North Central	0.9446		0.3731		0.6244	
West North Central	0.4760		0.1245		0.6262	
South	0.4862		0.7163		0.7567	
South Atlantic	0.5130		0.7856		0.8410	
East South Central	0.6442		0.2550		0.8013	
West South Central	0.9949		0.6215		0.8975	
West	0.2412		0.8555		0.0917	
Mountain	0.4262		0.0268		0.0388	
Pacific	0.0752		0.1530		0.4854	
COUNTY TYPE						
Large Metro	0.1791		0.6819		0.8328	
Small Metro	0.8433		0.7805		0.4902	
250K – 1 Mil. Pop.	0.3362		0.4650		0.4449	
< 250K Pop.	0.3224		0.5381		0.9517	
Nonmetro	0.7362		0.7479		0.4723	
Urbanized	0.8411		1.0000		0.4598	
Less Urbanized	0.8313		0.2802		0.1049	
Completely Rural	0.1230		0.1670		0.0218	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.76C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	1,052	843	799	686	438	366
GEOGRAPHIC DIVISION						
Northeast	425	348	334	290	158	164
New England	205	170	153	147	83	75
Middle Atlantic	372	303	297	248	134	145
Midwest	389	339	331	284	180	159
East North Central	318	302	269	241	150	130
West North Central	232	172	196	150	93	86
South	671	551	522	408	291	230
South Atlantic	539	428	409	298	239	168
East South Central	240	198	190	147	102	73
West South Central	313	302	279	219	138	141
West	560	450	399	343	229	168
Mountain	269	207	199	153	121	88
Pacific	485	394	344	302	196	139
COUNTY TYPE						
Large Metro	1,385	1,190	777	649	375	310
Small Metro	1,039	828	552	443	237	203
250K – 1 Mil. Pop.	926	708	497	364	210	163
< 250K Pop.	590	483	298	269	117	121
Nonmetro	665	583	346	316	149	141
Urbanized	460	424	240	211	99	92
Less Urbanized	498	454	256	244	100	104
Completely Rural	192	150	116	78	65	38

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.76D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.44	0.35	0.34	0.29	0.18	0.15
GEOGRAPHIC DIVISION						
Northeast	0.98	0.80	0.77	0.67	0.37	0.38
New England	1.80	1.48	1.34	1.28	0.73	0.66
Middle Atlantic	1.17	0.95	0.94	0.77	0.42	0.45
Midwest	0.76	0.66	0.65	0.56	0.35	0.31
East North Central	0.90	0.85	0.76	0.68	0.43	0.37
West North Central	1.49	1.10	1.26	0.96	0.59	0.55
South	0.76	0.62	0.59	0.46	0.33	0.26
South Atlantic	1.15	0.90	0.87	0.63	0.51	0.35
East South Central	1.71	1.40	1.36	1.04	0.73	0.52
West South Central	1.15	1.09	1.02	0.79	0.51	0.51
West	1.01	0.80	0.72	0.61	0.41	0.30
Mountain	1.60	1.21	1.18	0.90	0.72	0.51
Pacific	1.26	1.01	0.89	0.77	0.51	0.35
COUNTY TYPE						
Large Metro	0.61	0.51	0.49	0.41	0.27	0.22
Small Metro	0.81	0.63	0.62	0.48	0.32	0.27
250K – 1 Mil. Pop.	0.95	0.79	0.77	0.58	0.42	0.32
< 250K Pop.	1.57	1.00	1.06	0.83	0.50	0.46
Nonmetro	1.01	0.82	0.74	0.59	0.36	0.33
Urbanized	1.42	1.13	1.10	0.90	0.51	0.51
Less Urbanized	1.59	1.26	1.16	0.87	0.53	0.48
Completely Rural	2.89	2.51	2.11	1.55	1.36	0.94

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.76P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.3529		0.8183		0.5408	
GEOGRAPHIC DIVISION						
Northeast	0.8337		0.8754		0.0772	
New England	0.5596		0.6167		0.9289	
Middle Atlantic	0.5738		0.6503		0.0468	
Midwest	0.5637		0.9253		0.8219	
East North Central	0.7877		0.4719		0.5106	
West North Central	0.5343		0.2239		0.5441	
South	0.5320		0.7972		0.5851	
South Atlantic	0.5784		0.7795		0.7087	
East South Central	0.5859		0.3315		0.6554	
West South Central	0.9733		0.5570		0.8469	
West	0.1961		0.9093		0.0855	
Mountain	0.4734		0.0377		0.1023	
Pacific	0.0608		0.2029		0.3087	
COUNTY TYPE						
Large Metro	0.1943		0.7328		0.5870	
Small Metro	0.8092		0.6510		0.3948	
250K – 1 Mil. Pop.	0.3473		0.3949		0.3787	
< 250K Pop.	0.3765		0.6482		0.8491	
Nonmetro	0.7851		0.6539		0.3908	
Urbanized	0.7372		0.9255		0.6241	
Less Urbanized	0.8904		0.2691		0.0562	
Completely Rural	0.1443		0.2120		0.0355	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.77C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	282	276	255	261	194	211
GENDER						
Male	177	181	162	169	124	138
Female	174	184	159	172	122	140
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	250	232	227	222	174	181
White	213	197	198	187	153	157
Black or African American	87	101	76	89	53	64
American Indian or Alaska Native	14	14	12	14	8	12
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	63	57	59	52	40	34
Two or More Races	33	31	29	28	19	24
Hispanic or Latino	123	131	107	120	78	89
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	131	130	123	121	96	102
Female, White, Not Hispanic	130	136	121	130	96	108
Male, Black, Not Hispanic	59	65	51	57	35	43
Female, Black, Not Hispanic	56	63	49	56	36	38
Male, Hispanic	87	94	77	86	58	66
Female, Hispanic	77	85	68	77	49	61

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.77D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.47	0.48	0.46	0.49	0.40	0.46
GENDER						
Male	0.58	0.66	0.58	0.66	0.51	0.60
Female	0.62	0.65	0.61	0.65	0.54	0.61
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.53	0.53	0.52	0.53	0.46	0.51
White	0.62	0.61	0.61	0.62	0.56	0.61
Black or African American	1.08	1.20	1.06	1.16	0.84	0.95
American Indian or Alaska Native	3.65	4.13	3.55	4.34	2.96	4.13
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.28	2.18	2.21	2.13	1.64	1.53
Two or More Races	2.18	2.09	2.11	2.13	1.58	1.85
Hispanic or Latino	0.95	1.05	0.90	1.05	0.79	0.87
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.77	0.85	0.76	0.86	0.71	0.81
Female, White, Not Hispanic	0.82	0.85	0.82	0.86	0.75	0.86
Male, Black, Not Hispanic	1.54	1.67	1.46	1.57	1.15	1.38
Female, Black, Not Hispanic	1.37	1.64	1.42	1.56	1.19	1.18
Male, Hispanic	1.38	1.49	1.32	1.48	1.13	1.25
Female, Hispanic	1.29	1.38	1.23	1.36	1.05	1.21

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.77P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0745		0.1733		0.8849	
GENDER						
Male	0.0133		0.0389		0.7981	
Female	0.8431		0.9697		0.6337	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0595		0.1324		0.9767	
White	0.1885		0.3605		0.8933	
Black or African American	0.2000		0.2169		0.6626	
American Indian or Alaska Native	0.8307		0.9076		0.4191	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.8281		0.4954		0.4575	
Two or More Races	0.3806		0.7137		0.1688	
Hispanic or Latino	0.7630		0.9408		0.6455	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.0766		0.1188		0.9230	
Female, White, Not Hispanic	0.8543		0.8293		0.7666	
Male, Black, Not Hispanic	0.1380		0.1559		0.8728	
Female, Black, Not Hispanic	0.7293		0.6977		0.3918	
Male, Hispanic	0.4464		0.6025		0.6227	
Female, Hispanic	0.6772		0.6159		0.2116	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.78C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	2,141	1,771	1,880	1,553	1,674	1,398
GENDER						
Male	1,351	1,143	1,184	1,022	1,088	938
Female	1,341	1,096	1,212	973	1,045	865
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,024	1,700	1,791	1,491	1,592	1,336
White	1,852	1,564	1,661	1,372	1,479	1,223
Black or African American	719	596	584	476	486	401
American Indian or Alaska Native	104	106	75	81	63	69
Native Hawaiian or Other Pacific Islander	*	90	*	72	*	60
Asian	476	417	428	359	352	278
Two or More Races	223	158	179	148	174	128
Hispanic or Latino	725	653	605	573	527	493
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	1,112	945	1,013	847	936	788
Female, White, Not Hispanic	1,189	989	1,075	874	928	765
Male, Black, Not Hispanic	476	374	401	305	343	266
Female, Black, Not Hispanic	445	388	369	313	304	274
Male, Hispanic	534	456	463	408	415	365
Female, Hispanic	430	391	370	340	317	276

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.78D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.33	0.25	0.43	0.35	0.46	0.36
GENDER						
Male	0.40	0.27	0.59	0.44	0.63	0.49
Female	0.49	0.37	0.59	0.48	0.61	0.51
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.34	0.25	0.45	0.37	0.50	0.39
White	0.33	0.23	0.48	0.39	0.55	0.43
Black or African American	1.03	0.79	1.27	0.93	1.27	0.98
American Indian or Alaska Native	2.54	2.33	4.19	4.06	4.07	3.89
Native Hawaiian or Other Pacific Islander	*	4.40	*	5.77	*	5.61
Asian	2.42	1.67	2.42	1.68	2.37	1.63
Two or More Races	1.91	1.29	3.40	2.25	3.52	2.42
Hispanic or Latino	0.98	0.75	1.17	0.91	1.17	0.94
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.41	0.28	0.67	0.51	0.73	0.57
Female, White, Not Hispanic	0.48	0.34	0.65	0.57	0.72	0.61
Male, Black, Not Hispanic	1.22	0.96	1.82	1.39	1.96	1.49
Female, Black, Not Hispanic	1.51	1.10	1.70	1.27	1.61	1.32
Male, Hispanic	1.19	0.81	1.66	1.20	1.69	1.33
Female, Hispanic	1.44	1.13	1.58	1.22	1.52	1.18

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.78P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0362		0.4444		0.3788	
GENDER						
Male	0.2469		0.9608		0.8772	
Female	0.0598		0.3051		0.2679	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0474		0.8690		0.5685	
White	0.1231		0.6639		0.9797	
Black or African American	0.3038		0.7791		0.6649	
American Indian or Alaska Native	0.9573		0.8061		0.3600	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1597		0.2137		0.0837	
Two or More Races	0.9907		0.1636		0.6759	
Hispanic or Latino	0.3227		0.1010		0.2960	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.5966		0.5482		0.8009	
Female, White, Not Hispanic	0.1084		0.9481		0.8564	
Male, Black, Not Hispanic	0.8035		0.3760		0.4475	
Female, Black, Not Hispanic	0.1473		0.2544		0.1868	
Male, Hispanic	0.3243		0.1901		0.4924	
Female, Hispanic	0.5271		0.2434		0.3449	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.79C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	194	211	153	159	68	73
GENDER						
Male	124	138	102	114	53	56
Female	122	140	88	97	40	38
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	174	181	133	134	61	59
White	153	157	123	120	58	57
Black or African American	53	64	34	40	14	13
American Indian or Alaska Native	8	12	7	10	2	5
Native Hawaiian or Other Pacific Islander	*	*	6	*	*	1
Asian	40	34	25	23	6	9
Two or More Races	19	24	15	17	5	8
Hispanic or Latino	78	89	68	72	30	37
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	96	102	84	86	44	46
Female, White, Not Hispanic	96	108	70	75	34	29
Male, Black, Not Hispanic	35	43	23	30	11	9
Female, Black, Not Hispanic	36	38	25	24	11	10
Male, Hispanic	58	66	50	53	26	28
Female, Hispanic	49	61	39	46	16	22

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.79D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.40	0.46	0.34	0.37	0.17	0.19
GENDER						
Male	0.51	0.60	0.45	0.53	0.25	0.28
Female	0.54	0.61	0.42	0.47	0.21	0.20
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.46	0.51	0.38	0.41	0.19	0.20
White	0.56	0.61	0.49	0.50	0.26	0.27
Black or African American	0.84	0.95	0.57	0.69	0.26	0.25
American Indian or Alaska Native	2.96	4.13	2.59	3.39	0.82	1.97
Native Hawaiian or Other Pacific Islander	*	*	3.69	*	*	0.62
Asian	1.64	1.53	1.13	1.09	0.30	0.43
Two or More Races	1.58	1.85	1.33	1.43	0.45	0.67
Hispanic or Latino	0.79	0.87	0.73	0.76	0.35	0.42
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.71	0.81	0.66	0.72	0.38	0.42
Female, White, Not Hispanic	0.75	0.86	0.60	0.66	0.32	0.28
Male, Black, Not Hispanic	1.15	1.38	0.82	1.05	0.38	0.32
Female, Black, Not Hispanic	1.19	1.18	0.87	0.82	0.41	0.38
Male, Hispanic	1.13	1.25	1.03	1.08	0.59	0.61
Female, Hispanic	1.05	1.21	0.89	1.00	0.37	0.51

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.79P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.8849		0.4869		0.2596	
GENDER						
Male	0.7981		0.5386		0.3108	
Female	0.6337		0.6693		0.5385	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.9767		0.5483		0.1713	
White	0.8933		0.5345		0.1290	
Black or African American	0.6626		0.8632		0.9535	
American Indian or Alaska Native	0.4191		0.9326		0.4571	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4575		0.5585		0.8515	
Two or More Races	0.1688		0.6490		0.5563	
Hispanic or Latino	0.6455		0.7205		0.8912	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.9230		0.6110		0.3032	
Female, White, Not Hispanic	0.7666		0.6917		0.2094	
Male, Black, Not Hispanic	0.8728		0.4557		0.6598	
Female, Black, Not Hispanic	0.3918		0.5495		0.7398	
Male, Hispanic	0.6227		0.2506		0.5247	
Female, Hispanic	0.2116		0.4508		0.2668	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.80C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	1,674	1,398	944	762	448	372
GENDER						
Male	1,088	938	717	605	372	323
Female	1,045	865	502	398	233	165
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,592	1,336	871	687	416	342
White	1,479	1,223	785	604	382	311
Black or African American	486	401	304	270	140	122
American Indian or Alaska Native	63	69	52	61	22	28
Native Hawaiian or Other Pacific Islander	*	60	38	30	27	19
Asian	352	278	176	152	53	43
Two or More Races	174	128	79	76	57	30
Hispanic or Latino	527	493	398	351	164	140
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	936	788	583	467	304	267
Female, White, Not Hispanic	928	765	422	315	212	143
Male, Black, Not Hispanic	343	266	244	201	114	103
Female, Black, Not Hispanic	304	274	172	174	79	61
Male, Hispanic	415	365	331	293	158	132
Female, Hispanic	317	276	206	161	53	51

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.80D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.46	0.36	0.35	0.29	0.19	0.16
GENDER						
Male	0.63	0.49	0.56	0.47	0.34	0.28
Female	0.61	0.51	0.39	0.31	0.20	0.14
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.50	0.39	0.38	0.31	0.21	0.17
White	0.55	0.43	0.43	0.35	0.24	0.20
Black or African American	1.27	0.98	1.05	0.83	0.53	0.44
American Indian or Alaska Native	4.07	3.89	3.56	3.61	1.78	2.09
Native Hawaiian or Other Pacific Islander	*	5.61	5.04	3.51	3.64	2.16
Asian	2.37	1.63	1.50	1.25	0.46	0.36
Two or More Races	3.52	2.42	2.48	2.13	1.76	0.88
Hispanic or Latino	1.17	0.94	1.03	0.83	0.48	0.39
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.73	0.57	0.67	0.54	0.40	0.36
Female, White, Not Hispanic	0.72	0.61	0.49	0.37	0.27	0.18
Male, Black, Not Hispanic	1.96	1.49	1.74	1.33	0.96	0.83
Female, Black, Not Hispanic	1.61	1.32	1.12	1.07	0.54	0.41
Male, Hispanic	1.69	1.33	1.63	1.36	0.91	0.74
Female, Hispanic	1.52	1.18	1.11	0.80	0.32	0.29

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.80P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.3788		0.7482		0.6236	
GENDER						
Male	0.8772		0.8493		0.6444	
Female	0.2679		0.4163		0.8361	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.5685		0.9677		0.4595	
White	0.9797		0.3859		0.5489	
Black or African American	0.6649		0.1856		0.9737	
American Indian or Alaska Native	0.3600		0.3302		0.1791	
Native Hawaiian or Other Pacific Islander	*		0.1980		0.2223	
Asian	0.0837		0.1928		0.9428	
Two or More Races	0.6759		0.6433		0.2323	
Hispanic or Latino	0.2960		0.5297		0.5756	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.8009		0.5886		0.6718	
Female, White, Not Hispanic	0.8564		0.4009		0.6099	
Male, Black, Not Hispanic	0.4475		0.8786		0.7727	
Female, Black, Not Hispanic	0.1868		0.0617		0.6295	
Male, Hispanic	0.4924		0.5909		0.6179	
Female, Hispanic	0.3449		0.6126		0.6939	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.81C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	282	276	255	261	194	211
GEOGRAPHIC DIVISION						
Northeast	105	133	99	130	81	111
New England	57	77	54	78	46	65
Middle Atlantic	88	111	83	106	66	92
Midwest	128	108	118	100	91	86
East North Central	114	92	105	86	83	70
West North Central	58	58	52	52	38	47
South	170	154	152	145	112	113
South Atlantic	127	102	114	96	87	80
East South Central	56	53	52	52	33	40
West South Central	96	102	86	95	64	67
West	156	157	139	138	105	113
Mountain	69	60	63	56	40	45
Pacific	139	144	124	126	97	103
COUNTY TYPE						
Large Metro	207	219	185	201	146	158
Small Metro	171	161	156	155	109	127
250K – 1 Mil. Pop.	127	127	113	125	81	107
< 250K Pop.	123	98	112	90	79	69
Nonmetro	124	114	107	101	79	79
Urbanized	93	92	83	79	60	64
Less Urbanized	80	74	66	66	48	47
Completely Rural	40	24	35	19	25	14

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.81D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.47	0.48	0.46	0.49	0.40	0.46
GEOGRAPHIC DIVISION						
Northeast	1.00	1.24	0.99	1.27	0.96	1.27
New England	1.96	2.17	1.96	2.39	2.10	2.43
Middle Atlantic	1.15	1.52	1.14	1.52	1.06	1.51
Midwest	0.93	0.96	0.93	0.96	0.82	0.95
East North Central	1.11	1.16	1.11	1.13	1.04	1.07
West North Central	1.65	1.68	1.66	1.69	1.29	1.69
South	0.75	0.79	0.75	0.79	0.65	0.69
South Atlantic	1.13	1.08	1.11	1.08	0.98	0.99
East South Central	1.59	1.69	1.63	1.78	1.30	1.57
West South Central	1.23	1.51	1.23	1.48	1.06	1.22
West	1.02	1.08	0.94	1.06	0.85	0.97
Mountain	1.47	1.41	1.44	1.40	1.15	1.30
Pacific	1.30	1.40	1.19	1.37	1.10	1.26
COUNTY TYPE						
Large Metro	0.61	0.65	0.60	0.65	0.55	0.61
Small Metro	0.86	0.89	0.86	0.94	0.73	0.90
250K – 1 Mil. Pop.	0.94	1.06	0.93	1.11	0.83	1.11
< 250K Pop.	1.65	1.62	1.66	1.70	1.37	1.49
Nonmetro	1.18	1.25	1.14	1.18	1.02	1.06
Urbanized	1.95	2.36	1.94	2.14	1.69	1.98
Less Urbanized	1.58	1.42	1.46	1.38	1.35	1.19
Completely Rural	2.86	2.75	3.22	2.62	2.71	2.24

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.81P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0745		0.1733		0.8849	
GEOGRAPHIC DIVISION						
Northeast	0.6312		0.4570		0.3140	
New England	0.6095		0.4042		0.7899	
Middle Atlantic	0.8216		0.7474		0.3300	
Midwest	0.0630		0.0424		0.1416	
East North Central	0.0420		0.0167		0.0152	
West North Central	0.7709		0.9093		0.3065	
South	0.0708		0.3448		0.9048	
South Atlantic	0.0968		0.6010		0.8927	
East South Central	0.4070		0.5411		0.5786	
West South Central	0.5827		0.5562		0.9583	
West	0.8504		0.6399		0.6135	
Mountain	0.0514		0.1894		0.7419	
Pacific	0.4854		0.9128		0.6784	
COUNTY TYPE						
Large Metro	0.1196		0.1318		0.6956	
Small Metro	0.3950		0.7560		0.6155	
250K – 1 Mil. Pop.	0.5994		0.8021		0.2742	
< 250K Pop.	0.5117		0.4494		0.5325	
Nonmetro	0.7134		0.8324		0.6650	
Urbanized	0.9268		0.8724		0.4047	
Less Urbanized	0.5512		0.9038		0.9836	
Completely Rural	0.5178		0.1859		0.3870	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.82C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	2,141	1,771	1,880	1,553	1,674	1,398
GEOGRAPHIC DIVISION						
Northeast	892	656	813	624	740	581
New England	463	293	404	288	388	261
Middle Atlantic	764	588	708	556	633	520
Midwest	827	711	718	622	640	539
East North Central	652	615	571	548	503	482
West North Central	495	328	423	284	392	261
South	1,470	1,084	1,276	914	1,118	808
South Atlantic	1,235	798	1,078	679	943	609
East South Central	481	356	400	297	359	266
West South Central	686	647	603	548	522	487
West	1,089	928	958	825	839	748
Mountain	500	403	439	351	397	339
Pacific	966	847	853	752	739	665
COUNTY TYPE						
Large Metro	1,756	1,485	1,537	1,311	1,361	1,178
Small Metro	1,380	1,136	1,180	953	1,023	806
250K – 1 Mil. Pop.	1,168	937	1,028	799	913	693
< 250K Pop.	855	714	695	583	573	468
Nonmetro	929	896	757	693	653	565
Urbanized	637	658	535	500	451	412
Less Urbanized	716	670	572	536	489	440
Completely Rural	330	296	231	197	184	147

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.82D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.33	0.25	0.43	0.35	0.46	0.36
GEOGRAPHIC DIVISION						
Northeast	0.72	0.51	0.94	0.74	1.03	0.83
New England	0.86	0.84	1.57	1.44	1.86	1.52
Middle Atlantic	0.92	0.62	1.15	0.85	1.22	0.98
Midwest	0.46	0.42	0.73	0.66	0.80	0.68
East North Central	0.52	0.49	0.87	0.82	0.94	0.87
West North Central	0.88	0.74	1.41	1.02	1.58	1.15
South	0.59	0.43	0.77	0.60	0.80	0.64
South Atlantic	0.85	0.59	1.13	0.81	1.21	0.93
East South Central	1.47	0.98	1.86	1.39	1.79	1.44
West South Central	0.95	0.77	1.18	1.11	1.20	1.12
West	0.82	0.58	1.02	0.78	1.07	0.82
Mountain	1.12	0.73	1.51	1.05	1.67	1.26
Pacific	1.06	0.75	1.30	1.00	1.33	1.03
COUNTY TYPE						
Large Metro	0.49	0.36	0.61	0.49	0.64	0.53
Small Metro	0.51	0.41	0.76	0.61	0.85	0.65
250K – 1 Mil. Pop.	0.60	0.54	0.90	0.77	0.99	0.81
< 250K Pop.	0.96	0.61	1.51	0.96	1.70	1.03
Nonmetro	0.84	0.60	1.04	0.82	1.05	0.85
Urbanized	1.25	0.79	1.51	1.10	1.46	1.18
Less Urbanized	1.24	0.97	1.57	1.29	1.67	1.31
Completely Rural	2.42	1.89	3.33	2.53	2.99	2.56

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.82P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Lifetime (2013)	Lifetime (2014)	Past Year (2013)	Past Year (2014)	Past Month (2013)	Past Month (2014)
TOTAL	0.0362		0.4444		0.3788	
GEOGRAPHIC DIVISION						
Northeast	0.2662		0.4023		0.8906	
New England	0.7720		0.9767		0.5717	
Middle Atlantic	0.1857		0.3476		0.6381	
Midwest	0.8408		0.6543		0.6432	
East North Central	0.7132		0.5961		0.9768	
West North Central	0.8709		1.0000		0.4422	
South	0.0963		0.9709		0.5733	
South Atlantic	0.2606		0.6925		0.6285	
East South Central	0.3175		0.5817		0.6372	
West South Central	0.4258		0.2765		0.9972	
West	0.3196		0.2163		0.2064	
Mountain	0.6107		0.7845		0.4536	
Pacific	0.1813		0.1275		0.0622	
COUNTY TYPE						
Large Metro	0.0404		0.1222		0.1896	
Small Metro	0.7517		0.7613		0.7459	
250K – 1 Mil. Pop.	0.8682		0.2130		0.2917	
< 250K Pop.	0.4167		0.2181		0.3696	
Nonmetro	0.4202		0.2547		0.7289	
Urbanized	0.1968		0.5933		0.8785	
Less Urbanized	0.8998		0.6361		0.8561	
Completely Rural	0.9720		0.3002		0.1642	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.83C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	194	211	153	159	68	73
GEOGRAPHIC DIVISION						
Northeast	81	111	65	81	32	36
New England	46	65	36	40	20	18
Middle Atlantic	66	92	53	71	26	30
Midwest	91	86	72	67	29	34
East North Central	83	70	65	56	24	29
West North Central	38	47	31	36	15	18
South	112	113	82	85	41	40
South Atlantic	87	80	57	60	29	25
East South Central	33	40	22	31	12	16
West South Central	64	67	54	50	26	28
West	105	113	83	85	35	30
Mountain	40	45	32	30	14	15
Pacific	97	103	76	80	32	26
COUNTY TYPE						
Large Metro	146	158	108	123	52	54
Small Metro	109	127	87	88	35	40
250K – 1 Mil. Pop.	81	107	65	71	26	27
< 250K Pop.	79	69	62	53	24	29
Nonmetro	79	79	60	61	27	28
Urbanized	60	64	46	46	22	15
Less Urbanized	48	47	37	39	15	22
Completely Rural	25	14	17	10	6	4

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.83D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.40	0.46	0.34	0.37	0.17	0.19
GEOGRAPHIC DIVISION						
Northeast	0.96	1.27	0.84	1.04	0.47	0.51
New England	2.10	2.43	1.74	1.68	1.03	0.93
Middle Atlantic	1.06	1.51	0.95	1.27	0.52	0.60
Midwest	0.82	0.95	0.73	0.78	0.31	0.41
East North Central	1.04	1.07	0.91	0.92	0.37	0.50
West North Central	1.29	1.69	1.13	1.32	0.58	0.71
South	0.65	0.69	0.50	0.57	0.28	0.28
South Atlantic	0.98	0.99	0.70	0.79	0.40	0.35
East South Central	1.30	1.57	0.93	1.32	0.53	0.72
West South Central	1.06	1.22	0.95	1.01	0.51	0.57
West	0.85	0.97	0.75	0.81	0.35	0.31
Mountain	1.15	1.30	1.00	0.94	0.51	0.52
Pacific	1.10	1.26	0.98	1.09	0.45	0.38
COUNTY TYPE						
Large Metro	0.55	0.61	0.44	0.52	0.24	0.25
Small Metro	0.73	0.90	0.63	0.68	0.28	0.34
250K – 1 Mil. Pop.	0.83	1.11	0.72	0.79	0.31	0.34
< 250K Pop.	1.37	1.49	1.20	1.24	0.56	0.75
Nonmetro	1.02	1.06	0.82	0.87	0.43	0.46
Urbanized	1.69	1.98	1.38	1.51	0.76	0.58
Less Urbanized	1.35	1.19	1.11	1.10	0.53	0.73
Completely Rural	2.71	2.24	2.09	1.72	0.91	0.86

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 2.83P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.8849		0.4869		0.2596	
GEOGRAPHIC DIVISION						
Northeast	0.3140		0.9496		0.8911	
New England	0.7899		0.4150		0.9948	
Middle Atlantic	0.3300		0.7272		0.8898	
Midwest	0.1416		0.2399		0.1981	
East North Central	0.0152		0.0494		0.1925	
West North Central	0.3065		0.3351		0.7189	
South	0.9048		0.9450		0.7708	
South Atlantic	0.8927		0.8588		0.6214	
East South Central	0.5786		0.5551		0.6802	
West South Central	0.9583		0.6910		0.7894	
West	0.6135		0.7363		0.3052	
Mountain	0.7419		0.8155		0.8179	
Pacific	0.6784		0.7915		0.2968	
COUNTY TYPE						
Large Metro	0.6956		0.4897		0.2696	
Small Metro	0.6155		0.5400		0.4664	
250K – 1 Mil. Pop.	0.2742		0.7453		0.1709	
< 250K Pop.	0.5325		0.5870		0.7141	
Nonmetro	0.6650		0.6114		0.7520	
Urbanized	0.4047		0.5028		0.6049	
Less Urbanized	0.9836		0.8571		0.3570	
Completely Rural	0.3870		0.5774		0.9173	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.84C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	1,674	1,398	944	762	448	372
GEOGRAPHIC DIVISION						
Northeast	740	581	393	323	159	171
New England	388	261	205	155	86	74
Middle Atlantic	633	520	336	285	133	153
Midwest	640	539	382	305	186	158
East North Central	503	482	303	263	152	131
West North Central	392	261	239	160	100	87
South	1,118	808	629	451	302	232
South Atlantic	943	609	499	322	248	168
East South Central	359	266	205	163	107	74
West South Central	522	487	345	264	140	145
West	839	748	463	378	241	173
Mountain	397	339	218	169	123	91
Pacific	739	665	409	340	210	143
COUNTY TYPE						
Large Metro	1,361	1,178	763	630	370	304
Small Metro	1,023	806	538	425	233	197
250K – 1 Mil. Pop.	913	693	487	352	208	159
< 250K Pop.	573	468	279	257	113	116
Nonmetro	653	565	334	298	145	139
Urbanized	451	412	231	194	97	90
Less Urbanized	489	440	246	232	96	100
Completely Rural	184	147	112	76	64	38

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.84D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.46	0.36	0.35	0.29	0.19	0.16
GEOGRAPHIC DIVISION						
Northeast	1.03	0.83	0.80	0.69	0.37	0.39
New England	1.86	1.52	1.39	1.35	0.74	0.69
Middle Atlantic	1.22	0.98	0.97	0.79	0.43	0.47
Midwest	0.80	0.68	0.67	0.57	0.37	0.33
East North Central	0.94	0.87	0.79	0.69	0.44	0.39
West North Central	1.58	1.15	1.30	0.98	0.62	0.57
South	0.80	0.64	0.62	0.47	0.35	0.27
South Atlantic	1.21	0.93	0.92	0.64	0.54	0.37
East South Central	1.79	1.44	1.40	1.07	0.76	0.52
West South Central	1.20	1.12	1.07	0.82	0.53	0.53
West	1.07	0.82	0.76	0.62	0.44	0.31
Mountain	1.67	1.26	1.24	0.92	0.76	0.53
Pacific	1.33	1.03	0.95	0.79	0.54	0.37
COUNTY TYPE						
Large Metro	0.64	0.53	0.52	0.42	0.29	0.23
Small Metro	0.85	0.65	0.65	0.50	0.34	0.28
250K – 1 Mil. Pop.	0.99	0.81	0.80	0.60	0.44	0.33
< 250K Pop.	1.70	1.03	1.11	0.85	0.52	0.48
Nonmetro	1.05	0.85	0.76	0.60	0.37	0.35
Urbanized	1.46	1.18	1.14	0.91	0.54	0.54
Less Urbanized	1.67	1.31	1.19	0.90	0.54	0.51
Completely Rural	2.99	2.56	2.21	1.57	1.45	0.96

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 2.84P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Alcohol Use (2013)	Alcohol Use (2014)	Binge Alcohol Use (2013)	Binge Alcohol Use (2014)	Heavy Alcohol Use (2013)	Heavy Alcohol Use (2014)
TOTAL	0.3788		0.7482		0.6236	
GEOGRAPHIC DIVISION						
Northeast	0.8906		0.8248		0.0833	
New England	0.5717		0.6488		0.9730	
Middle Atlantic	0.6381		0.6144		0.0479	
Midwest	0.6432		0.9822		0.9653	
East North Central	0.9768		0.3108		0.6159	
West North Central	0.4422		0.1500		0.4774	
South	0.5733		0.8160		0.6548	
South Atlantic	0.6285		0.8117		0.7442	
East South Central	0.6372		0.3791		0.6455	
West South Central	0.9972		0.6081		0.9441	
West	0.2064		0.8602		0.0872	
Mountain	0.4536		0.0377		0.0896	
Pacific	0.0622		0.1768		0.3341	
COUNTY TYPE						
Large Metro	0.1896		0.6612		0.6574	
Small Metro	0.7459		0.7000		0.4125	
250K – 1 Mil. Pop.	0.2917		0.4387		0.4531	
< 250K Pop.	0.3696		0.6386		0.7355	
Nonmetro	0.7289		0.7206		0.3571	
Urbanized	0.8785		0.9608		0.4882	
Less Urbanized	0.8561		0.2702		0.0667	
Completely Rural	0.1642		0.2318		0.0380	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.1A Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances, by Age Group: Numbers in Thousands, 2013 and 2014

Risk/Availability	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
PERCEPTIONS OF GREAT RISK¹								
Cigarettes								
Smoke One or More Packs Per Day	184,585 ^b	187,503	15,759 ^b	16,258	22,859	23,253	145,967	147,993
Marijuana								
Smoke Once a Month	72,547 ^b	68,922	5,899 ^a	5,572	5,147 ^b	4,689	61,502 ^a	58,661
Smoke Once or Twice a Week	95,410 ^b	89,132	9,645 ^b	9,139	7,257 ^b	6,357	78,509 ^b	73,636
Cocaine								
Use Once a Month	178,108	178,477	11,901	12,064	21,609	21,671	144,598	144,742
Use Once or Twice a Week	224,860	224,747	19,005	18,745	28,990	29,096	176,865	176,905
Heroin								
Try Once or Twice	211,257 ^b	216,026	13,984	14,282	27,206 ^a	27,640	170,067 ^b	174,105
Use Once or Twice a Week	240,078 ^b	242,829	19,185	19,304	31,930 ^b	32,251	188,963 ^b	191,273
LSD								
Try Once or Twice	174,818	175,144	10,928	10,753	19,690	19,207	144,200	145,185
Use Once or Twice a Week	216,427	216,174	16,199	15,929	26,380	26,322	173,848	173,923
Alcohol								
Have Four or Five Drinks Nearly Every Day	174,033	173,739	15,318	15,307	20,690	20,847	138,025	137,585
Have Five or More Drinks Once or Twice a Week	107,350	105,792	9,546	9,615	11,494	11,651	86,310	84,527
PERCEIVED AVAILABILITY²								
Fairly or Very Easy to Obtain ³								
Marijuana	146,638 ^b	152,532	11,659	11,453	25,751	25,703	109,228 ^b	115,376
Cocaine	58,931	60,741	3,397	3,385	9,324	9,290	46,210	48,066
Crack	51,867	53,766	3,517	3,372	6,571	6,371	41,779 ^a	44,023
Heroin	37,097 ^b	41,567	2,132	2,210	5,003	5,036	29,962 ^b	34,321
LSD	33,951	34,568	2,603	2,685	5,621	5,734	25,726	26,150
Approached in the Past Month by Someone Selling Drugs	17,650	17,335	3,087	3,013	5,656	5,644	8,906	8,678

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.1B Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances, by Age Group: Percentages, 2013 and 2014

Risk/Availability	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
PERCEPTIONS OF GREAT RISK¹								
Cigarettes								
Smoke One or More Packs Per Day	70.9	71.2	64.3 ^b	66.3	66.0	66.9	72.5	72.6
Marijuana								
Smoke Once a Month	28.2 ^b	26.5	24.2 ^a	22.9	14.9 ^b	13.5	31.0 ^b	29.2
Smoke Once or Twice a Week	37.1 ^b	34.3	39.5 ^b	37.4	21.0 ^b	18.3	39.6 ^b	36.6
Cocaine								
Use Once a Month	69.2	68.6	49.3	49.8	62.8	62.9	72.8	71.9
Use Once or Twice a Week	87.3 ^b	86.3	78.4 ^a	77.2	84.3	84.4	89.0 ^b	87.8
Heroin								
Try Once or Twice	82.2 ^a	83.1	58.2	59.2	79.2	80.2	85.7	86.5
Use Once or Twice a Week	93.3	93.3	79.8	79.9	92.9	93.5	95.1	94.8
LSD								
Try Once or Twice	68.7	68.0	47.1	46.2	57.9 ^a	56.4	73.1	72.5
Use Once or Twice a Week	85.0 ^b	83.9	69.7 ^a	68.4	77.5	77.3	88.1 ^b	86.8
Alcohol								
Have Four or Five Drinks Nearly Every Day	66.9	66.1	62.5	62.4	59.8	60.0	68.7 ^a	67.6
Have Five or More Drinks Once or Twice a Week	41.3 ^a	40.3	39.0	39.2	33.2	33.5	43.0 ^a	41.6
PERCEIVED AVAILABILITY²								
Fairly or Very Easy to Obtain ³								
Marijuana	58.6 ^b	60.2	48.6	47.8	75.3	74.9	56.9 ^b	59.2
Cocaine	24.0	24.4	14.4	14.4	27.9	27.5	24.5	25.2
Crack	21.2	21.7	14.9	14.3	19.7	19.0	22.2	23.1
Heroin	15.2 ^b	16.8	9.1	9.4	15.1	15.0	15.9 ^b	18.0
LSD	14.0	14.0	11.3	11.6	16.9	17.1	13.7	13.8
Approached in the Past Month by Someone Selling Drugs	6.7	6.5	12.4	12.1	16.3	16.2	4.4	4.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.2A Perceived Great Risk of Harm Associated with Cigarette and Alcohol Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	CIGARETTES		ALCOHOL			
	Smoke One or More Packs Per Day		Have Four or Five Drinks Nearly Every Day		Have Five or More Drinks Once or Twice a Week	
	2013	2014	2013	2014	2013	2014
TOTAL	15,759 ^b	16,258	15,318	15,307	9,546	9,615
AGE						
12-13	4,760	4,927	4,976	4,940	3,215	3,116
14-15	5,432	5,447	5,244	5,088	3,263	3,276
16-17	5,567	5,884	5,097	5,280	3,069	3,223
GENDER						
Male	7,732 ^a	7,994	7,150	7,159	4,554	4,542
Female	8,028 ^a	8,264	8,168	8,148	4,991	5,073
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	12,399 ^a	12,694	12,021	12,021	7,355	7,292
White	8,828	9,047	8,303	8,283	4,699	4,620
Black or African American	2,054	2,121	2,187	2,195	1,661	1,667
American Indian or Alaska Native	81	94	80	73	52	51
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	875	834	883	893	597	632
Two or More Races	507	525	503	506	298	287
Hispanic or Latino	3,361 ^a	3,564	3,296	3,287	2,191	2,323
COUNTY TYPE						
Large Metro	8,738 ^a	9,352	8,673	8,925	5,362	5,543
Small Metro	4,724	4,583	4,465	4,313	2,803	2,703
Nonmetro	2,298	2,323	2,180	2,069	1,381	1,369

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.2B Perceived Great Risk of Harm Associated with Cigarette and Alcohol Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	CIGARETTES		ALCOHOL			
	Smoke One or More Packs Per Day		Have Four or Five Drinks Nearly Every Day		Have Five or More Drinks Once or Twice a Week	
	2013	2014	2013	2014	2013	2014
TOTAL	64.3 ^b	66.3	62.5	62.4	39.0	39.2
AGE						
12-13	61.2 ^a	64.1	64.0	64.1	41.4	40.4
14-15	64.6	65.1	62.4	60.8	38.9	39.2
16-17	66.7 ^a	69.4	61.1	62.4	36.8	38.1
GENDER						
Male	61.8 ^a	63.9	57.3	57.2	36.5	36.3
Female	66.8 ^a	68.8	67.8	67.7	41.5	42.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	65.0 ^b	67.0	63.0	63.4	38.6	38.5
White	65.6 ^b	67.9	61.6	62.0	34.9	34.6
Black or African American	60.4	62.4	64.6	64.8	49.1	49.3
American Indian or Alaska Native	52.6	57.7	52.1	45.0	33.5	31.5
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	71.9	69.2	72.6	74.1	49.4	52.1
Two or More Races	68.3	69.6	67.3	66.8	39.9	38.0
Hispanic or Latino	61.6	64.0	60.6	59.0	40.3	41.7
COUNTY TYPE						
Large Metro	65.3 ^b	67.7	64.9	64.5	40.2	40.1
Small Metro	64.5	65.7	60.9	61.7	38.3	38.7
Nonmetro	60.0	62.4	56.9	55.6	36.0	36.9

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.3A Perceived Great Risk of Harm Associated with Marijuana and Cocaine Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	MARIJUANA				COCAINE			
	Smoke Once a Month		Smoke Once or Twice a Week		Use Once a Month		Use Once or Twice a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	5,899 ^a	5,572	9,645 ^b	9,139	11,901	12,064	19,005	18,745
AGE								
12-13	2,599 ^a	2,369	4,158	3,939	3,007	3,003	5,329	5,169
14-15	1,917	1,840	3,220	3,075	4,042	4,004	6,591	6,431
16-17	1,383	1,362	2,267	2,125	4,852	5,057	7,085	7,145
GENDER								
Male	2,809	2,657	4,659 ^b	4,345	6,209	6,318	9,822 ^a	9,595
Female	3,090	2,915	4,986	4,794	5,693	5,746	9,183	9,150
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	4,684 ^b	4,316	7,780 ^b	7,267	9,423	9,451	15,050 ^a	14,793
White	3,336 ^a	3,084	5,753 ^b	5,254	6,592	6,616	10,915	10,644
Black or African American	773	742	1,061	1,084	1,774	1,783	2,413	2,464
American Indian or Alaska Native	35	26	55	56	74	95	109	120
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	358	297	607	557	519	526	936	890
Two or More Races	162	147	263	271	411	380	603	596
Hispanic or Latino	1,215	1,256	1,865	1,872	2,478	2,613	3,955	3,953
COUNTY TYPE								
Large Metro	3,118	2,932	5,123	4,927	6,397	6,689	10,387	10,469
Small Metro	1,736	1,613	2,899	2,647	3,585	3,478	5,684	5,405
Nonmetro	1,044	1,028	1,623	1,565	1,920	1,896	2,934	2,872

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.3B Perceived Great Risk of Harm Associated with Marijuana and Cocaine Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	MARIJUANA				COCAINE			
	Smoke Once a Month		Smoke Once or Twice a Week		Use Once a Month		Use Once or Twice a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	24.2 ^a	22.9	39.5 ^b	37.4	49.3	49.8	78.4 ^a	77.2
AGE								
12-13	33.9 ^a	31.1	54.0 ^a	51.5	39.6	39.7	69.9	68.2
14-15	22.9	22.1	38.4	36.9	48.8	48.3	79.2	77.4
16-17	16.6	16.1	27.2 ^a	25.2	58.6	60.3	85.5	85.0
GENDER								
Male	22.6	21.4	37.4 ^b	34.8	50.5	51.1	79.6 ^b	77.5
Female	25.9	24.4	41.7	40.1	48.0	48.3	77.1	76.8
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	24.7 ^b	22.9	41.0 ^b	38.5	50.1	50.5	79.8	78.8
White	24.9 ^a	23.3	42.9 ^b	39.6	49.6	50.2	81.9	80.6
Black or African American	22.8	21.9	31.3	31.8	53.0	53.2	72.0	73.4
American Indian or Alaska Native	22.8	15.7	36.2	34.4	49.0	58.7	71.4	74.6
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	30.0	24.9	51.0	46.4	44.0	44.0	78.0	74.1
Two or More Races	21.7	19.6	35.4	36.1	56.3	51.0	82.6	79.1
Hispanic or Latino	22.4	22.7	34.3	33.7	46.2	47.3	73.5	71.7
COUNTY TYPE								
Large Metro	23.5 ^b	21.4	38.6 ^b	35.8	48.7	48.9	78.7 ^b	76.5
Small Metro	23.8	23.3	39.6	38.0	49.6	50.3	78.4	78.0
Nonmetro	27.4	27.8	42.7	42.2	50.8	51.8	77.4	78.2

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.4A Perceived Great Risk of Harm Associated with Heroin and LSD Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	HEROIN				LSD			
	Try Once or Twice		Use Once or Twice a Week		Try Once or Twice		Use Once or Twice a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	13,984	14,282	19,185	19,304	10,928	10,753	16,199	15,929
AGE								
12-13	3,355	3,426	5,167	5,142	2,817	2,788	4,348	4,225
14-15	4,799	4,799	6,647	6,629	3,773	3,649	5,570	5,508
16-17	5,830	6,057	7,372	7,533	4,338	4,315	6,281	6,196
GENDER								
Male	7,351	7,459	9,968	9,960	5,723	5,680	8,411 ^a	8,204
Female	6,634	6,823	9,218	9,344	5,205	5,073	7,788	7,725
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	10,915	10,976	15,180	15,117	8,463	8,254	12,871 ^b	12,521
White	7,705	7,700	11,069	10,927	5,974	5,727	9,441	9,065
Black or African American	2,039	2,068	2,451	2,483	1,599	1,630	2,035	2,040
American Indian or Alaska Native	84	104	101	131	67	70	83	91
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	590	599	888	891	460	450	788	779
Two or More Races	439	448	602	607	319	335	472	486
Hispanic or Latino	3,070 ^b	3,307	4,005 ^a	4,187	2,465	2,499	3,328	3,408
COUNTY TYPE								
Large Metro	7,545	7,981	10,464	10,891	5,813	5,897	8,834	8,906
Small Metro	4,185	4,107	5,726	5,501	3,299	3,091	4,770	4,540
Nonmetro	2,254	2,194	2,995	2,912	1,816	1,765	2,595	2,482

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.4B Perceived Great Risk of Harm Associated with Heroin and LSD Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	HEROIN				LSD			
	Try Once or Twice		Use Once or Twice a Week		Try Once or Twice		Use Once or Twice a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	58.2	59.2	79.8	79.9	47.1	46.2	69.7 ^a	68.4
AGE								
12-13	44.7	45.8	68.7	68.7	39.6	39.6	61.0	59.9
14-15	58.2	58.1	80.5	80.1	47.5	45.6	69.9	68.7
16-17	70.5	72.3	89.0	89.9	53.4	52.4	77.2 ^a	75.2
GENDER								
Male	60.2	60.7	81.5	80.9	48.3	47.7	70.8 ^a	68.9
Female	56.2	57.7	78.0	79.0	45.9	44.6	68.6	67.8
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	58.4	58.9	81.1	81.0	47.0	45.9	71.3 ^a	69.5
White	58.3	58.8	83.5	83.2	46.8	45.3	73.9 ^b	71.6
Black or African American	61.0	61.7	73.3	74.2	49.7	50.3	63.5	63.0
American Indian or Alaska Native	54.9	64.2	66.8 ^b	81.0	45.1	43.9	54.7	57.1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	50.9	50.8	76.8	75.4	40.9	39.4	69.3	68.3
Two or More Races	60.0	60.0	82.1	81.0	46.4	47.3	68.2	67.9
Hispanic or Latino	57.8	60.4	75.3	76.5	47.6	47.2	64.2	64.4
COUNTY TYPE								
Large Metro	57.7	58.7	79.9	80.1	46.3	45.0	70.3 ^b	67.8
Small Metro	58.2	59.9	79.5	80.0	47.2	46.5	68.2	68.4
Nonmetro	60.1	59.9	79.7	79.3	49.6	49.9	70.8	70.2

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.5A Perceived That Obtaining Specific Illicit Drugs Would Be Fairly or Very Easy among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/ Geographic Characteristic	Marijuana (2013)	Marijuana (2014)	Cocaine (2013)	Cocaine (2014)	Crack (2013)	Crack (2014)	Heroin (2013)	Heroin (2014)	LSD (2013)	LSD (2014)
TOTAL	11,659	11,453	3,397	3,385	3,517	3,372	2,132	2,210	2,603	2,685
AGE										
12-13	1,528	1,386	653	629	804	770	444	482	367	318
14-15	4,218	4,089	1,200	1,169	1,326	1,228	793	735	847	850
16-17	5,913	5,977	1,545	1,586	1,387	1,374	896	993	1,389	1,517
GENDER										
Male	5,772	5,569	1,357	1,349	1,269	1,243	861	910	1,086	1,134
Female	5,887	5,884	2,040	2,036	2,248	2,129	1,271	1,300	1,517	1,550
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	8,961	8,887	2,518	2,544	2,672	2,605	1,650	1,752	1,973	2,062
White	6,514	6,342	1,788	1,772	1,852	1,754	1,204	1,217	1,485	1,584
Black or African American	1,564	1,610	479	490	553	564	311	350	293	275
American Indian or Alaska Native	60	71	16	12	25	15	20	10	11	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*	*	*
Asian	415	425	120	132	122	136	53 ^a	93	90	84
Two or More Races	361	386	102	123	103	119	59	69	82	86
Hispanic or Latino	2,699	2,566	879	841	846	767	483	457	631	622
COUNTY TYPE										
Large Metro	6,545	6,540	1,840	1,909	1,884	1,841	1,160	1,231	1,494	1,567
Small Metro	3,461	3,314	1,056	985	1,106	1,035	649	659	771	795
Nonmetro	1,654	1,598	502	491	528	496	323	320	339	323

*Low precision; no estimate reported.

NOTE: Response categories for the Perceived Availability questions include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy." Respondents with unknown Perceived Availability data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.5B Perceived That Obtaining Specific Illicit Drugs Would Be Fairly or Very Easy among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/ Geographic Characteristic	Marijuana (2013)	Marijuana (2014)	Cocaine (2013)	Cocaine (2014)	Crack (2013)	Crack (2014)	Heroin (2013)	Heroin (2014)	LSD (2013)	LSD (2014)
TOTAL	48.6	47.8	14.4	14.4	14.9	14.3	9.1	9.4	11.3	11.6
AGE										
12-13	20.5	18.7	8.9	8.6	10.9	10.6	6.1	6.6	5.2	4.5
14-15	51.2	50.0	14.9	14.5	16.4	15.3	9.9	9.2	10.8	10.8
16-17	71.5	71.3	19.0	19.4	17.1	16.8	11.1	12.2	17.3	18.7
GENDER										
Male	47.2	45.5	11.3	11.2	10.6	10.3	7.2	7.6	9.2	9.6
Female	50.1	50.2	17.7	17.7	19.5	18.5	11.2	11.3	13.6	13.8
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	48.2	48.0	13.8	14.0	14.6	14.3	9.1	9.7	11.1	11.6
White	49.5	48.8	13.8	13.9	14.3	13.8	9.4	9.6	11.8	12.7
Black or African American	47.4	48.2	14.8	14.9	17.1	17.2	9.7	10.7	9.3	8.5
American Indian or Alaska Native	39.2	44.9	11.0	7.8	16.8	9.6	13.4 ^a	6.2	7.6	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*	*	*
Asian	35.2	35.8	10.3	11.2	10.5	11.5	4.6 ^a	8.0	8.0	7.3
Two or More Races	49.6	51.8	14.4	16.9	14.4	16.5	8.4	9.4	12.0	12.0
Hispanic or Latino	50.3 ^a	47.0	16.7	15.7	16.1	14.3	9.2	8.6	12.3	11.8
COUNTY TYPE										
Large Metro	50.2	48.3	14.4	14.3	14.7	13.9	9.1	9.3	12.0	12.0
Small Metro	48.2	48.7	15.0	14.7	15.7	15.5	9.3	9.9	11.2	12.1
Nonmetro	43.9	43.8	13.6	13.8	14.3	14.0	8.8	9.0	9.4	9.3

*Low precision; no estimate reported.

NOTE: Response categories for the Perceived Availability questions include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy." Respondents with unknown Perceived Availability data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.6A Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	22,920	22,965	21,677 ^a	21,423	22,209	22,028	22,232	22,210
AGE								
12-13	7,368	7,369	7,351	7,301	7,391	7,395	7,241	7,285
14-15	7,950	7,917	7,538	7,396	7,728	7,612	7,744	7,641
16-17	7,603	7,678	6,788	6,726	7,089	7,022	7,247	7,284
GENDER								
Male	11,587	11,681	10,920	10,928	11,180	11,224	11,166	11,194
Female	11,333	11,284	10,757 ^b	10,495	11,029 ^b	10,805	11,066	11,016
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	17,831	17,791	16,776 ^b	16,505	17,228 ^a	17,010	17,331	17,222
White	12,705	12,490	11,822	11,474	12,188	11,897	12,292	12,088
Black or African American	3,052	3,186	2,936	3,010	2,991	3,054	2,993	3,079
American Indian or Alaska Native	124	152	130	139	131	145	134	146
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1,175	1,156	1,166	1,138	1,172	1,153	1,149	1,120
Two or More Races	689	710	634	660	658	675	676	692
Hispanic or Latino	5,090 ^a	5,174	4,901	4,918	4,980	5,018	4,901	4,988
COUNTY TYPE								
Large Metro	12,590	13,078	11,795	12,119	12,056	12,484	12,196	12,667
Small Metro	6,808	6,471	6,460	6,032	6,659	6,205	6,617	6,255
Nonmetro	3,522	3,416	3,421	3,272	3,494	3,340	3,420	3,288

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Perceptions of Parents' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove." Respondents with unknown Perceptions of Parents' Feelings data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.6B Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	93.5	93.8	88.4 ^a	87.5	90.6	90.0	90.7	90.6
AGE								
12-13	95.0	95.9	94.4	94.8	95.1	95.9	93.2 ^a	94.4
14-15	94.4	94.7	89.7	88.5	92.1	91.1	92.2	91.4
16-17	91.3	90.9	81.3	79.8	85.0 ^a	83.4	86.9	86.3
GENDER								
Male	93.0	93.7	87.5	87.6	89.7	90.0	89.6	89.6
Female	94.1	93.8	89.3 ^b	87.3	91.6 ^b	89.9	91.9	91.5
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	93.4	93.7	87.8	87.0	90.3	89.7	90.8	90.7
White	94.1	93.5	87.6 ^b	85.9	90.3 ^a	89.1	91.1	90.4
Black or African American	90.6 ^b	93.9	87.0	89.0	88.8	90.5	89.0	91.0
American Indian or Alaska Native	82.5 ^a	93.8	86.4	86.5	87.1	90.1	88.9	91.0
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	96.1	96.0	95.0	94.0	95.7	95.4	93.5	92.8
Two or More Races	92.8	93.3	84.7	86.9	88.4	88.9	90.3	91.0
Hispanic or Latino	93.9	93.9	90.2	89.1	91.8	90.8	90.4	90.0
COUNTY TYPE								
Large Metro	94.3	94.8	88.2	87.8	90.3	90.5	91.4	91.6
Small Metro	92.8	93.0	88.1	86.7	90.8 ^a	89.1	90.2	89.7
Nonmetro	92.2	91.6	89.5	87.8	91.5	89.7	89.6	88.2

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Perceptions of Parents' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove." Respondents with unknown Perceptions of Parents' Feelings data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.7A Youths Felt That Close Friends Would Strongly Disapprove or Somewhat Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	21,788	21,978	19,450 ^a	19,164	19,680	19,511	21,123 ^a	21,371
AGE								
12-13	7,204	7,238	7,162	7,155	7,168	7,200	7,189	7,259
14-15	7,489	7,538	6,848	6,684	6,953	6,818	7,301	7,332
16-17	7,094	7,202	5,439	5,325	5,560	5,494	6,633	6,780
GENDER								
Male	10,855	10,959	9,618	9,584	9,721	9,708	10,466	10,617
Female	10,932	11,019	9,832 ^b	9,579	9,959	9,804	10,656	10,754
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	17,054	17,129	15,194 ^b	14,891	15,405 ^a	15,165	16,546	16,628
White	12,159	12,113	10,779	10,486	10,988	10,698	11,851	11,752
Black or African American	2,894	2,956	2,575	2,553	2,568	2,583	2,747	2,871
American Indian or Alaska Native	128	146	117	124	119	122	125	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1,142	1,131	1,070	1,080	1,084	1,094	1,110	1,133
Two or More Races	649	687	577	565	567	582	630	652
Hispanic or Latino	4,733 ^a	4,849	4,256	4,273	4,276	4,346	4,576 ^b	4,742
COUNTY TYPE								
Large Metro	11,944	12,545	10,506	10,763	10,659	10,980	11,573	12,217
Small Metro	6,512	6,179	5,794	5,390	5,849	5,491	6,316	5,975
Nonmetro	3,331	3,255	3,150	3,011	3,173	3,040	3,233	3,178

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Perceptions of Close Friends' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Perceptions of Close Friends' Feelings data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.7B Youths Felt That Close Friends Would Strongly Disapprove or Somewhat Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	89.1	89.9	79.6 ^a	78.4	80.6	79.9	86.6	87.4
AGE								
12-13	93.1	93.9	92.7	92.9	92.9	93.4	93.3	94.2
14-15	89.4	90.5	81.8	80.3	83.1	82.0	87.3	88.1
16-17	85.1	85.5	65.3	63.3	66.7	65.3	79.7	80.5
GENDER								
Male	87.5	88.3	77.6	77.3	78.4	78.3	84.6	85.6
Female	90.8	91.5	81.7 ^b	79.6	82.9	81.5	88.7	89.3
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	89.5	90.4	79.8	78.6	81.0	80.1	87.0	87.8
White	90.2	90.8	79.9	78.6	81.5	80.3	87.9	88.1
Black or African American	86.3	87.2	76.9	75.7	77.0	76.7	82.6	85.2
American Indian or Alaska Native	85.2	90.6	78.7	76.8	79.7	76.0	83.5	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	93.6	93.6	87.6	89.1	88.9	90.2	91.0	93.5
Two or More Races	87.7	91.1	77.9	75.4	76.5	77.5	85.2	86.4
Hispanic or Latino	87.7	88.2	79.0	77.6	79.3	79.0	85.1	86.0
COUNTY TYPE								
Large Metro	89.7 ^a	91.0	78.9	78.1	80.1	79.7	87.1 ^a	88.6
Small Metro	89.1	88.8	79.3	77.6	80.1	79.1	86.6	85.9
Nonmetro	87.4	87.7	82.6	81.1	83.3	82.0	85.0	85.8

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Perceptions of Close Friends' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Perceptions of Close Friends' Feelings data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.8A Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	22,424 ^b	22,736	19,509	19,516	19,436	19,425	21,778 ^a	22,015
AGE								
12-13	7,315	7,383	7,188	7,245	7,202	7,230	7,310	7,372
14-15	7,711	7,723	6,803	6,743	6,768	6,713	7,486	7,465
16-17	7,398	7,630	5,518	5,528	5,466	5,483	6,982	7,178
GENDER								
Male	11,278 ^b	11,527	9,754	9,909	9,672	9,826	10,869 ^b	11,120
Female	11,146	11,208	9,756	9,607	9,763	9,599	10,908	10,895
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	17,511	17,609	15,197	15,101	15,185	15,057	17,042	17,113
White	12,488	12,452	10,721	10,565	10,791	10,579	12,177	12,067
Black or African American	2,954	3,073	2,622	2,683	2,563	2,643	2,841	2,977
American Indian or Alaska Native	130	*	119	*	118	*	131	*
Native Hawaiian or Other Pacific Islander	92	*	*	*	*	*	*	*
Asian	1,153	1,142	1,071	1,054	1,066	1,052	1,141	1,149
Two or More Races	692	702	578	590	564	574	666	693
Hispanic or Latino	4,914 ^b	5,127	4,312	4,415	4,251	4,369	4,735 ^b	4,902
COUNTY TYPE								
Large Metro	12,284	12,944	10,528	10,879	10,494	10,840	11,862	12,498
Small Metro	6,714	6,346	5,851	5,536	5,813	5,506	6,546	6,206
Nonmetro	3,426	3,445	3,130	3,101	3,130	3,079	3,369	3,310

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Feelings about Peers include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Feelings about Peers data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.8B Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	91.4 ^b	92.5	79.5	79.5	79.2	79.2	88.7 ^a	89.7
AGE								
12-13	93.8 ^a	95.2	92.1 ^a	93.6	92.4	93.4	93.7 ^a	95.0
14-15	91.9	92.3	80.9	80.6	80.6	80.4	89.1	89.4
16-17	88.7 ^a	90.3	66.2	65.4	65.6	65.0	83.7	85.0
GENDER								
Male	90.4 ^b	92.2	78.1	79.4	77.6	78.8	87.1 ^b	89.1
Female	92.4	92.9	80.9	79.6	81.0	79.6	90.4	90.3
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	91.7 ^a	92.5	79.6	79.5	79.6	79.3	89.2	90.0
White	92.4	93.0	79.3	78.9	79.8	79.0	90.0	90.1
Black or African American	88.0 ^a	90.4	77.9	79.2	76.5	78.1	84.9 ^a	88.0
American Indian or Alaska Native	85.9	*	79.0	*	77.7	*	87.0	*
Native Hawaiian or Other Pacific Islander	95.3	*	*	*	*	*	*	*
Asian	93.8	94.2	87.8	87.2	87.4	87.3	93.3	95.1
Two or More Races	92.6	92.4	77.2	77.6	75.4	75.8	88.9	91.3
Hispanic or Latino	90.4 ^a	92.5	79.1	79.6	78.1	78.9	86.9	88.5
COUNTY TYPE								
Large Metro	91.8 ^b	93.5	78.7	78.6	78.5	78.4	88.7 ^b	90.3
Small Metro	91.7	91.0	79.7	79.3	79.3	78.9	89.2	89.0
Nonmetro	89.6 ^b	92.0	81.8	83.0	81.8	82.5	88.1	88.7

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Feelings about Peers include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Feelings about Peers data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.9A Participated in a Prevention Program or Group in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	PREVENTION PROGRAM OR GROUP					
	Problem-Solving, Communication Skills, or Self-Esteem Group		Violence Prevention Program		Pregnancy or Sexually Transmitted Disease Prevention Program	
	2013	2014	2013	2014	2013	2014
TOTAL	5,571	5,535	3,241	3,112	2,203	2,143
AGE						
12-13	1,892	1,941	1,441	1,326	629 ^a	515
14-15	1,972	1,925	1,133	1,009	910	967
16-17	1,707	1,669	667 ^a	776	663	661
GENDER						
Male	2,614	2,630	1,615	1,597	1,071	1,066
Female	2,957	2,905	1,626	1,515	1,132	1,078
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4,506	4,386	2,491	2,388	1,643	1,628
White	2,909	2,806	1,422	1,326	930	965
Black or African American	972	972	733	762	488	470
American Indian or Alaska Native	44	36	35	32	23	25
Native Hawaiian or Other Pacific Islander	*	*	10	*	*	*
Asian	355	356	152	161	107	90
Two or More Races	209	198	139 ^a	79	88	56
Hispanic or Latino	1,065	1,148	750	724	560	515
COUNTY TYPE						
Large Metro	3,065	3,139	1,761	1,720	1,105	1,232
Small Metro	1,690	1,592	973	890	713	600
Nonmetro	816	804	508	502	385	311

*Low precision; no estimate reported.

NOTE: Respondents with unknown Prevention Program or Group Participation data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.9B Participated in a Prevention Program or Group in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	PREVENTION PROGRAM OR GROUP					
	Problem-Solving, Communication Skills, or Self-Esteem Group		Violence Prevention Program		Pregnancy or Sexually Transmitted Disease Prevention Program	
	2013	2014	2013	2014	2013	2014
TOTAL	23.0	22.9	13.2	12.7	8.9	8.7
AGE						
12-13	24.9	25.7	18.5	17.1	8.0 ^a	6.6
14-15	23.6	23.3	13.5	12.0	10.8	11.5
16-17	20.5	19.8	8.0	9.2	7.9	7.8
GENDER						
Male	21.1	21.3	12.9	12.8	8.5	8.5
Female	24.9	24.5	13.5	12.6	9.4	8.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	23.9	23.4	13.1	12.6	8.6	8.5
White	21.9	21.3	10.5	9.9	6.9	7.2
Black or African American	28.8	28.7	21.6	22.3	14.3	13.7
American Indian or Alaska Native	28.0	22.1	22.1	19.6	14.5	15.0
Native Hawaiian or Other Pacific Islander	*	*	10.7	*	*	*
Asian	29.5	29.9	12.7	13.2	8.8	7.4
Two or More Races	28.5	26.4	18.7 ^b	10.5	11.9 ^a	7.4
Hispanic or Latino	19.7	21.0	13.7	13.0	10.2	9.2
COUNTY TYPE						
Large Metro	23.1	23.0	13.1	12.4	8.2	8.9
Small Metro	23.3	23.1	13.3	12.7	9.7	8.6
Nonmetro	21.7	22.1	13.2	13.5	10.0	8.3

*Low precision; no estimate reported.

NOTE: Respondents with unknown Prevention Program or Group Participation data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.10A Participated in a Substance Use Program or Exposure to Substance Use Prevention Message in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE PROGRAM OR PREVENTION MESSAGE							
	Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School		Participated in Program to Help Deal with Drug or Alcohol Use		Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School		Talked to Parents about Dangers of Tobacco, Alcohol, or Drug Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	2,840	2,735	892	1,003	17,792	17,849	13,565	13,728
AGE								
12-13	1,096	993	374	384	5,244	5,175	4,246	4,131
14-15	1,003	939	283	320	6,157	6,275	4,767	4,812
16-17	742	803	235	299	6,392	6,399	4,553	4,786
GENDER								
Male	1,470	1,461	483	550	8,820	8,874	6,720	6,903
Female	1,370	1,274	409	453	8,973	8,975	6,845	6,825
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	2,240	2,113	675	758	13,815	13,795	10,527	10,699
White	1,384	1,353	393	471	9,775	9,702	7,792	7,889
Black or African American	551	517	202	232	2,381	2,439	1,578	1,632
American Indian or Alaska Native	30	31	15	10	108	129	69	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	162	137	31	23	947	921	587	630
Two or More Races	98 ^b	50	32	18	533	541	439	424
Hispanic or Latino	600	621	217	245	3,977	4,054	3,038	3,029
COUNTY TYPE								
Large Metro	1,537	1,525	461 ^a	570	9,860	10,220	7,435	7,845
Small Metro	863	745	290	282	5,272	5,050	4,110	3,912
Nonmetro	440	465	141	151	2,660	2,579	2,020	1,971

*Low precision; no estimate reported.

NOTE: Respondents with unknown Substance Use Program Participation or Prevention Message Exposure data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.10B Participated in a Substance Use Program or Exposure to Substance Use Prevention Message in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE PROGRAM OR PREVENTION MESSAGE							
	Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School		Participated in Program to Help Deal with Drug or Alcohol Use		Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School		Talked to Parents about Dangers of Tobacco, Alcohol, or Drug Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	11.5	11.1	3.6	4.1	72.6	72.9	55.5	56.2
AGE								
12-13	14.0	12.8	4.8	5.0	67.6	67.3	54.9	53.7
14-15	11.9	11.2	3.4	3.8	73.2	75.1	56.9	57.7
16-17	8.9	9.5	2.8	3.5	76.6	75.8	54.7	56.9
GENDER								
Male	11.7	11.6	3.9	4.4	70.4	70.9	53.9	55.4
Female	11.4	10.5	3.4	3.8	74.8	74.9	57.2	57.0
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	11.7	11.1	3.5	4.0	72.5	72.8	55.5	56.5
White	10.3	10.1	2.9 ^a	3.5	72.7	72.9	58.1	59.3
Black or African American	16.2	15.1	6.0	6.8	70.3	71.5	46.6	47.9
American Indian or Alaska Native	19.4	18.6	9.5	5.9	69.1	79.5	44.3	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	13.2	11.2	2.6	1.9	77.7	75.7	48.8	52.6
Two or More Races	13.1 ^b	6.6	4.3	2.4	71.4	71.8	59.5	56.0
Hispanic or Latino	10.9	11.1	4.0	4.4	72.6	73.3	55.7	54.8
COUNTY TYPE								
Large Metro	11.5	11.0	3.4	4.1	73.8	73.9	55.8	56.9
Small Metro	11.7	10.6	4.0	4.0	72.0	72.5	56.4	56.2
Nonmetro	11.5	12.5	3.7	4.1	69.5	69.9	53.0	53.2

*Low precision; no estimate reported.

NOTE: Respondents with unknown Substance Use Program Participation or Prevention Message Exposure data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.11A Participated in a Physical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	DELINQUENT BEHAVIOR					
	Got into a Serious Fight at School or Work		Took Part in a Group-Against-Group Fight		Attacked Someone with the Intent to Seriously Hurt Them	
	2013	2014	2013	2014	2013	2014
TOTAL	4,381	4,164	2,714	2,615	1,260	1,195
AGE						
12-13	1,577	1,493	955	894	378	407
14-15	1,561	1,496	994	934	438	418
16-17	1,243	1,175	765	787	444	370
GENDER						
Male	2,637 ^a	2,435	1,502	1,422	759	687
Female	1,744	1,728	1,212	1,193	501	508
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	3,312	3,177	2,019	1,936	986	925
White	2,058	1,987	1,265	1,247	538	511
Black or African American	922	877	525	501	338	292
American Indian or Alaska Native	42	33	32	24	11	9
Native Hawaiian or Other Pacific Islander	*	*	*	*	1	*
Asian	97	114	81	67	37	52
Two or More Races	170	146	111	74	62	48
Hispanic or Latino	1,069	986	694	680	273	270
COUNTY TYPE						
Large Metro	2,403	2,312	1,485	1,490	647	666
Small Metro	1,278	1,199	798	702	409	335
Nonmetro	699	653	430	423	204	194

*Low precision; no estimate reported.

NOTE: Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time. Respondents with unknown Delinquent Behavior data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.11B Participated in a Physical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	DELINQUENT BEHAVIOR					
	Got into a Serious Fight at School or Work		Took Part in a Group-Against-Group Fight		Attacked Someone with the Intent to Seriously Hurt Them	
	2013	2014	2013	2014	2013	2014
TOTAL	17.7	16.9	11.0	10.6	5.1	4.8
AGE						
12-13	20.0	19.1	12.1	11.4	4.8	5.2
14-15	18.4	17.8	11.7	11.1	5.2	5.0
16-17	14.8	13.9	9.1	9.3	5.3	4.4
GENDER						
Male	20.9 ^a	19.4	11.9	11.3	6.0	5.4
Female	14.4	14.2	10.0	9.8	4.1	4.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	17.2	16.6	10.5	10.1	5.1	4.8
White	15.2	14.8	9.3	9.3	4.0	3.8
Black or African American	27.0	25.4	15.4	14.5	9.9	8.4
American Indian or Alaska Native	26.5	19.7	20.2	14.6	6.8	5.5
Native Hawaiian or Other Pacific Islander	*	*	*	*	0.7	*
Asian	7.9	9.3	6.5	5.5	3.0	4.3
Two or More Races	22.7	19.2	14.8 ^a	9.7	8.4	6.2
Hispanic or Latino	19.3	17.7	12.6	12.2	4.9	4.8
COUNTY TYPE						
Large Metro	17.8	16.6	11.0	10.7	4.8	4.8
Small Metro	17.3	17.0	10.8	10.0	5.5	4.8
Nonmetro	18.1	17.4	11.2	11.3	5.3	5.2

*Low precision; no estimate reported.

NOTE: Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time. Respondents with unknown Delinquent Behavior data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.12A Participated in a Nonphysical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	DELINQUENT BEHAVIOR					
	Carried a Handgun		Sold Illegal Drugs		Stole or Tried to Steal Anything Worth More Than \$50	
	2013	2014	2013	2014	2013	2014
TOTAL	851	945	595	502	697	649
AGE						
12-13	222 ^a	288	37	29	117	115
14-15	315	328	190	146	238	222
16-17	314	329	368	326	343	312
GENDER						
Male	682	741	430 ^a	341	453	437
Female	169	204	165	161	244	212
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	689	768	441	393	509	503
White	525	603	322	269	316	331
Black or African American	97	102	79	86	123	108
American Indian or Alaska Native	14	10	9	3	15	7
Native Hawaiian or Other Pacific Islander	*	*	4	*	3	*
Asian	16	16	12	5	25	30
Two or More Races	37	33	16	23	26	17
Hispanic or Latino	161	177	154	109	188	147
COUNTY TYPE						
Large Metro	352	425	317	286	398	365
Small Metro	280	308	203 ^b	136	213	212
Nonmetro	219	211	74	80	86	73

*Low precision; no estimate reported.

NOTE: Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time. Respondents with unknown Delinquent Behavior data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.12B Participated in a Nonphysical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	DELINQUENT BEHAVIOR					
	Carried a Handgun		Sold Illegal Drugs		Stole or Tried to Steal Anything Worth More Than \$50	
	2013	2014	2013	2014	2013	2014
TOTAL	3.4	3.8	2.4	2.0	2.8	2.6
AGE						
12-13	2.8 ^a	3.7	0.5	0.4	1.5	1.5
14-15	3.7	3.9	2.2	1.7	2.8	2.6
16-17	3.7	3.9	4.4	3.9	4.1	3.7
GENDER						
Male	5.4	5.9	3.4 ^a	2.7	3.6	3.5
Female	1.4	1.7	1.4	1.3	2.0	1.7
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	3.6	4.0	2.3	2.1	2.6	2.6
White	3.9	4.5	2.4	2.0	2.3	2.5
Black or African American	2.8	2.9	2.3	2.5	3.6	3.1
American Indian or Alaska Native	8.5	6.0	5.5	2.0	9.6	4.4
Native Hawaiian or Other Pacific Islander	*	*	4.2	*	3.4	*
Asian	1.3	1.3	1.0	0.4	2.1	2.4
Two or More Races	4.9	4.4	2.1	3.0	3.5	2.3
Hispanic or Latino	2.9	3.2	2.8	1.9	3.4	2.6
COUNTY TYPE						
Large Metro	2.6	3.0	2.4	2.1	3.0	2.6
Small Metro	3.8	4.4	2.7 ^a	1.9	2.9	3.0
Nonmetro	5.7	5.6	1.9	2.1	2.2	1.9

*Low precision; no estimate reported.

NOTE: Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time. Respondents with unknown Delinquent Behavior data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.13A Selected Risk and Protective Factors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	RISK FACTORS						PROTECTIVE FACTORS			
	Youth Moved 2 or More Times in the Past 5 Years		Youth Argued with at Least One Parent 10 or More Times in the Past Year		Youth Was Approached by Someone Selling Drugs in the Past Month		Youth Had Someone to Talk to about Problems ¹		Youth Attended 25 or More Religious Services in the Past Year ²	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	5,817	5,847	4,953	4,934	3,087	3,013	23,151	23,076	7,196	6,993
AGE										
12-13	1,994	2,020	1,193	1,232	361	319	7,386	7,284	2,155	2,190
14-15	2,022	1,972	1,912	1,768	1,087	1,098	7,907	7,855	2,560	2,352
16-17	1,800	1,855	1,847	1,934	1,639	1,596	7,858	7,936	2,481	2,451
GENDER										
Male	2,995	2,933	1,937	1,877	1,801	1,731	11,816	11,813	3,427	3,267
Female	2,822	2,914	3,016	3,056	1,287	1,281	11,336	11,263	3,769	3,726
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	4,248	4,348	3,936	3,913	2,276	2,181	18,006	17,938	5,878	5,763
White	2,540	2,610	3,054	3,034	1,503	1,457	12,822	12,714	4,399	4,252
Black or African American	1,221	1,209	397	405	526	496	3,150	3,196	880	937
American Indian or Alaska Native	48	48	29	*	16	*	145	155	33	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*	*	*
Asian	182	188	259	266	92	76	1,106	1,078	341	322
Two or More Races	230	263	185	167	124	104	698	702	194	172
Hispanic or Latino	1,569	1,500	1,017	1,020	811	832	5,145	5,138	1,318	1,231
COUNTY TYPE										
Large Metro	3,095	3,195	2,837	2,865	1,824	1,804	12,570	12,933	3,635	3,835
Small Metro	1,861	1,785	1,405	1,363	925	893	6,948	6,638	2,281 ^a	2,044
Nonmetro	860	868	711	705	338	317	3,634	3,505	1,280	1,114

*Low precision; no estimate reported.

NOTE: Respondents with unknown Risk or Protective Factors data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents were able to select multiple categories of persons that they would talk to about a serious problem, including the choice of "nobody." Respondents who reported both a person that they would talk to and that they had nobody to talk to were excluded from the analysis.

² Special occasions, such as weddings, funerals, or other special events, were not to have been considered in the number of times attended religious services.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.13B Selected Risk and Protective Factors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	RISK FACTORS						PROTECTIVE FACTORS			
	Youth Moved 2 or More Times in the Past 5 Years		Youth Argued with at Least One Parent 10 or More Times in the Past Year		Youth Was Approached by Someone Selling Drugs in the Past Month		Youth Had Someone to Talk to about Problems ¹		Youth Attended 25 or More Religious Services in the Past Year ²	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	24.0	24.1	20.3	20.2	12.4	12.1	95.5	95.3	29.8	29.0
AGE										
12-13	26.3	26.6	15.3	16.0	4.5	4.0	95.5	95.5	28.5	29.4
14-15	24.3	23.9	22.9	21.3	12.8	13.0	95.6	95.2	30.8 ^a	28.4
16-17	21.7	22.1	22.2	23.0	19.5	18.8	95.5	95.3	30.0	29.2
GENDER										
Male	24.3	23.8	15.5	15.1	14.2	13.7	95.3	95.6	27.8	26.5
Female	23.8	24.5	25.2	25.5	10.6	10.5	95.7	95.1	31.9	31.6
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	22.6	23.1	20.7	20.7	11.8	11.4	95.5	95.8	31.3	30.8
White	19.0	19.7	22.8	22.9	11.0	10.8	96.2	96.6	33.0	32.3
Black or African American	37.3	36.1	11.7	11.8	15.3	14.2	94.4	94.7	26.6	28.1
American Indian or Alaska Native	30.6	29.1	18.5	*	9.8	*	93.0	95.6	21.4	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*	*	*
Asian	15.0	15.5	21.8	22.2	7.4	6.2	92.2	91.5	29.2	27.7
Two or More Races	31.4	35.1	24.9	22.1	16.5	13.6	95.0	95.3	26.4	23.3
Hispanic or Latino	29.3	27.6	18.6	18.4	14.6	14.8	95.5 ^b	93.7	24.7	22.7
COUNTY TYPE										
Large Metro	23.5	23.3	21.3	20.8	13.5	12.9	95.2	94.8	27.7	28.2
Small Metro	25.7	25.9	19.2	19.6	12.5	12.7	96.0	96.1	31.6	29.9
Nonmetro	22.9	23.6	18.6	19.0	8.7	8.4	95.5	95.8	34.0 ^a	30.4

*Low precision; no estimate reported.

NOTE: Respondents with unknown Risk or Protective Factors data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents were able to select multiple categories of persons that they would talk to about a serious problem, including the choice of "nobody." Respondents who reported both a person that they would talk to and that they had nobody to talk to were excluded from the analysis.

² Special occasions, such as weddings, funerals, or other special events, were not to have been considered in the number of times attended religious services.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.14A Youths Agreed or Strongly Agreed with Statements about Religious Beliefs among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	STATEMENTS ABOUT RELIGIOUS BELIEFS					
	Religious Beliefs Are a Very Important Part of Youth's Life		Religious Beliefs Influence Youth's Decision Making in Life		It Is Important for Friends to Share Religious Beliefs	
	2013	2014	2013	2014	2013	2014
TOTAL	17,697 ^a	17,368	16,008 ^b	15,562	7,684 ^a	7,289
AGE						
12-13	6,085	5,935	5,534	5,344	2,800	2,650
14-15	6,087	5,882	5,486	5,267	2,571	2,453
16-17	5,525	5,551	4,988	4,950	2,312	2,186
GENDER						
Male	8,699	8,539	7,878	7,721	3,911 ^a	3,684
Female	8,998	8,829	8,131 ^a	7,841	3,772	3,604
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	13,610 ^a	13,308	12,393 ^a	12,040	6,019 ^b	5,626
White	9,330	9,025	8,486	8,154	4,195	3,912
Black or African American	2,696	2,751	2,516	2,492	1,236	1,172
American Indian or Alaska Native	125	130	106	120	66	73
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	888	829	786	758	304	253
Two or More Races	496	492	429	437	181	167
Hispanic or Latino	4,087	4,060	3,616	3,522	1,665	1,663
COUNTY TYPE						
Large Metro	9,458	9,642	8,472	8,578	3,719	3,642
Small Metro	5,381	5,003	4,915 ^a	4,491	2,441	2,210
Nonmetro	2,858	2,723	2,621	2,493	1,524	1,437

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Religious Beliefs include "Strongly disagree," "Disagree," "Agree," and "Strongly agree." The estimates in this table correspond to persons reporting "Agree" or "Strongly agree." Respondents with unknown Religious Beliefs data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.14B Youths Agreed or Strongly Agreed with Statements about Religious Beliefs among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	STATEMENTS ABOUT RELIGIOUS BELIEFS					
	Religious Beliefs Are a Very Important Part of Youth's Life		Religious Beliefs Influence Youth's Decision Making in Life		It Is Important for Friends to Share Religious Beliefs	
	2013	2014	2013	2014	2013	2014
TOTAL	73.3 ^a	71.9	66.2 ^b	64.4	32.0 ^a	30.4
AGE						
12-13	80.1	78.9	72.8	71.0	37.1	35.6
14-15	73.4 ^a	71.2	66.1 ^a	63.8	31.2	29.8
16-17	67.1	66.3	60.3	59.1	28.0	26.2
GENDER						
Male	71.0	69.5	64.1	62.9	32.0 ^a	30.2
Female	75.7	74.4	68.5 ^b	65.9	32.0	30.6
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	72.4	71.1	65.8	64.3	32.1 ^a	30.2
White	70.2 ^a	68.5	63.7 ^a	61.8	31.6 ^a	29.8
Black or African American	81.1	82.3	75.3	74.5	37.5	35.4
American Indian or Alaska Native	81.0	82.0	69.0	75.4	43.5	45.5
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	74.3	69.7	65.7	64.2	25.9	21.7
Two or More Races	68.2	66.3	58.6	58.8	24.8	22.5
Hispanic or Latino	76.5	74.4	67.7 ^a	64.5	31.4	30.8
COUNTY TYPE						
Large Metro	71.9	70.5	64.3	62.8	28.4	26.8
Small Metro	74.7	73.2	68.1 ^a	65.5	34.0	32.4
Nonmetro	75.8	74.4	69.5	68.2	40.6	39.7

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Religious Beliefs include "Strongly disagree," "Disagree," "Agree," and "Strongly agree." The estimates in this table correspond to persons reporting "Agree" or "Strongly agree." Respondents with unknown Religious Beliefs data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.15A Parents Always or Sometimes Took Part in Specific Behaviors in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	PARENTAL INVOLVEMENT							
	Parents Made Youth Do Chores around the House		Parents Limited Amount of Time Youth Watched TV		Parents Let Youth Know He or She Had Done a Good Job		Parents Let Youth Know They Were Proud of Something He or She Had Done	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	21,898 ^a	22,141	10,064	10,122	21,181	21,360	21,209	21,252
AGE								
12-13	7,086	7,049	4,102	4,085	7,148	7,106	7,174	7,157
14-15	7,500	7,590	3,386	3,383	7,215	7,224	7,183	7,130
16-17	7,313	7,502	2,575	2,654	6,818	7,030	6,852	6,964
GENDER								
Male	11,133 ^b	11,391	5,019	5,123	10,993 ^a	11,166	11,014	11,095
Female	10,765	10,751	5,044	4,998	10,188	10,194	10,195	10,156
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	16,964	17,086	7,693	7,608	16,597	16,670	16,645	16,566
White	11,888	11,949	5,279	5,271	11,894	11,844	11,928	11,782
Black or African American	3,139	3,199	1,337	1,351	2,863	2,948	2,918	2,962
American Indian or Alaska Native	144	153	61	70	133	*	127	131
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1,039	1,017	667	600	986	1,033	965	973
Two or More Races	678	680	311	265	642	621	628	635
Hispanic or Latino	4,934 ^a	5,055	2,371	2,514	4,584	4,690	4,564	4,685
COUNTY TYPE								
Large Metro	11,860	12,426	5,676	5,922	11,503	11,990	11,506	11,972
Small Metro	6,599	6,313	2,938	2,804	6,371	6,074	6,394	6,030
Nonmetro	3,439	3,402	1,450	1,397	3,307	3,297	3,308	3,250

*Low precision; no estimate reported.

NOTE: Response categories for the Parental Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes." Respondents with unknown Parental Involvement data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.15B Parents Always or Sometimes Took Part in Specific Behaviors in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	PARENTAL INVOLVEMENT							
	Parents Made Youth Do Chores around the House		Parents Limited Amount of Time Youth Watched TV		Parents Let Youth Know He or She Had Done a Good Job		Parents Let Youth Know They Were Proud of Something He or She Had Done	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	88.5 ^a	89.5	40.8	41.1	85.7	86.4	85.7	86.0
AGE								
12-13	89.5	89.8	52.2	52.4	90.6	90.5	90.7	91.3
14-15	88.6 ^a	90.2	40.2	40.4	85.1	85.9	84.8	84.8
16-17	87.4	88.5	30.8	31.3	81.5	83.0	81.8	82.3
GENDER								
Male	88.3 ^b	90.4	39.9	40.9	87.2 ^a	88.6	87.2	88.2
Female	88.7	88.6	41.7	41.3	84.0	84.0	84.1	83.8
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	88.2 ^a	89.3	40.1	40.0	86.4	87.2	86.6	86.7
White	87.6 ^b	89.0	39.0	39.4	87.7	88.2	87.9	87.9
Black or African American	92.0	92.7	39.3	39.4	83.8	85.6	85.4	85.8
American Indian or Alaska Native	90.3	92.4	38.2	42.5	83.3	*	81.1	79.7
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	84.3	83.5	54.1	49.6	80.9	84.7	78.4	79.6
Two or More Races	90.3	89.1	41.6	34.9	85.6	81.3	83.8	83.5
Hispanic or Latino	89.4	90.1	43.2	45.0	82.9	83.6	82.5	83.6
COUNTY TYPE								
Large Metro	88.1	89.1	42.3	42.6	85.4	86.0	85.3	85.9
Small Metro	89.1	89.7	39.8	40.0	86.1	86.4	86.4	85.8
Nonmetro	88.9	90.5	37.7	37.6	85.5 ^a	87.8	85.6	86.7

*Low precision; no estimate reported.

NOTE: Response categories for the Parental Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes." Respondents with unknown Parental Involvement data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.16A Measures of Student Characteristics in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	STUDENT CHARACTERISTICS							
	Participated in One or Fewer Activities ¹		Drug or Alcohol Prevention Messages Were Seen or Heard in School ²		Youth Liked a Lot or Kind of Liked Going to School ³		Grade Average for Last Completed Period Was "D" or Lower ⁴	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	2,924	2,858	16,909	16,857	18,810	18,531	967	971
AGE								
12-13	773	737	5,212	5,229	5,890	5,850	214	256
14-15	984	902	6,192	6,110	6,519	6,354	373	347
16-17	1,168	1,219	5,504	5,517	6,402	6,326	379	368
GENDER								
Male	1,727	1,591	8,410	8,410	9,696	9,597	604	547
Female	1,197	1,267	8,499	8,447	9,114	8,934	363	424
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	2,126	2,055	13,293	13,279	14,564	14,392	749	698
White	1,547	1,480	9,562	9,524	10,226	10,032	511	478
Black or African American	357	368	2,180	2,142	2,613	2,615	169	146
American Indian or Alaska Native	17	*	88	*	104	*	13	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	105	88	893	923	980	989	12	18
Two or More Races	77	79	506	541	568	575	41	39
Hispanic or Latino	798	803	3,616	3,578	4,246	4,139	217	273
COUNTY TYPE								
Large Metro	1,542	1,544	9,413	9,692	10,396	10,548	472	524
Small Metro	932	882	5,023	4,691	5,586	5,261	308	304
Nonmetro	451	432	2,473	2,474	2,829	2,722	187	143

*Low precision; no estimate reported.

NOTE: Respondents with unknown Student Characteristics data were excluded from the respective analyses.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The number of activities participated in is based on four questions: participation in school-based activities, community-based activities, church or faith-based activities, and other activities. For each question, the response categories include "None," "One," "Two," and "3 or more." Respondents were excluded from the analysis if they had unknown data for at least three of the questions and if their known responses did not place them in the recoded "Two or more" category.

² Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

³ Response categories for the School Satisfaction question include "You liked going to school a lot," "You kind of liked going to school," "You didn't like going to school very much," and "You hated going to school." The estimates in this table correspond to persons reporting "You liked going to school a lot" or "You kind of liked going to school."

⁴ Response categories for Grade Average question include "An 'A+', 'A,' or 'A-minus' average"; "A 'B+', 'B,' or 'B-minus' average"; "A 'C+', 'C,' or 'C-minus' average"; "A 'D' or less than a 'D' average"; and "My school does not give these grades." The estimates in this table correspond to persons reporting "A 'D' or less than a 'D' average." Respondents reporting that their school does not give these grades were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.16B Measures of Student Characteristics in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	STUDENT CHARACTERISTICS							
	Participated in One or Fewer Activities ¹		Drug or Alcohol Prevention Messages Were Seen or Heard in School ²		Youth Liked a Lot or Kind of Liked Going to School ³		Grade Average for Last Completed Period Was "D" or Lower ⁴	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	12.6	12.3	73.5	73.0	81.1 ^b	79.6	4.4	4.4
AGE								
12-13	10.9	10.4	74.3	74.7	82.9	82.7	3.3	3.9
14-15	12.3	11.3	77.8	76.5	81.2 ^a	79.3	4.8	4.5
16-17	14.5	14.9	68.6	67.9	79.4 ^a	77.3	4.8	4.6
GENDER								
Male	14.6	13.4	71.6	71.4	81.9	80.9	5.4	4.9
Female	10.6	11.1	75.5	74.6	80.2 ^a	78.3	3.3	3.9
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	11.7	11.3	73.6	73.4	80.0	78.9	4.3	4.0
White	11.9	11.4	73.9	73.8	78.4	77.3	4.1	3.9
Black or African American	11.6	11.8	71.1	69.0	84.6	83.5	5.8	4.9
American Indian or Alaska Native	13.2	*	69.3	*	80.7	*	10.2	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	9.1	7.6	77.7	80.7	84.8	85.6	1.1	1.7
Two or More Races	11.0	10.8	72.8	74.0	81.4	78.0	6.2	5.6
Hispanic or Latino	16.0	15.9	73.3	71.5	85.0 ^a	82.0	4.6	5.8
COUNTY TYPE								
Large Metro	12.2	11.7	74.9	74.1	82.0 ^a	80.1	3.9	4.2
Small Metro	13.4	13.3	72.8	71.3	80.4	79.3	4.6	4.8
Nonmetro	12.6	12.4	70.1	71.9	79.1	78.2	5.5	4.3

*Low precision; no estimate reported.

NOTE: Respondents with unknown Student Characteristics data were excluded from the respective analyses.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The number of activities participated in is based on four questions: participation in school-based activities, community-based activities, church or faith-based activities, and other activities. For each question, the response categories include "None," "One," "Two," and "3 or more." Respondents were excluded from the analysis if they had unknown data for at least three of the questions and if their known responses did not place them in the recoded "Two or more" category.

² Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

³ Response categories for the School Satisfaction question include "You liked going to school a lot," "You kind of liked going to school," "You didn't like going to school very much," and "You hated going to school." The estimates in this table correspond to persons reporting "You liked going to school a lot" or "You kind of liked going to school."

⁴ Response categories for Grade Average question include "An 'A+', 'A,' or 'A-minus' average"; "A 'B+', 'B,' or 'B-minus' average"; "A 'C+', 'C,' or 'C-minus' average"; "A 'D' or less than a 'D' average"; and "My school does not give these grades." The estimates in this table correspond to persons reporting "A 'D' or less than a 'D' average." Respondents reporting that their school does not give these grades were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.17A Adults Always or Sometimes Took Part in Specific School-Related Behaviors in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	ADULT INVOLVEMENT							
	Teacher Let Youth Know He or She Was Doing a Good Job with School Work		Parents Checked That Youth Did Homework		Parents Provided Help with Homework		Parents Limited Time Youth Went Out with Friends on a School Night	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	18,553	18,201	19,207	19,348	18,360	18,696	15,637	15,671
AGE								
12-13	6,049	5,952	6,451	6,369	6,319	6,332	4,934	4,998
14-15	6,345	6,096	6,777	6,773	6,462	6,521	5,554	5,447
16-17	6,159	6,153	5,979	6,206	5,580	5,843	5,148	5,227
GENDER								
Male	9,507	9,440	10,081	10,155	9,452	9,686	7,696	7,653
Female	9,046	8,761	9,126	9,193	8,908	9,010	7,941	8,018
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	14,498	14,177	15,059	15,111	14,710	14,865	12,211	12,130
White	10,170	9,858	10,808	10,775	10,794	10,747	8,789	8,509
Black or African American	2,633	2,639	2,539	2,651	2,386	2,477	2,102	2,192
American Indian or Alaska Native	110	*	106	115	103	113	85	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	972	951	943	915	799	848	746	759
Two or More Races	542	557	589	581	562	604	427	505
Hispanic or Latino	4,055	4,024	4,148	4,237	3,650	3,831	3,425	3,541
COUNTY TYPE								
Large Metro	10,237	10,369	10,460	10,938	9,866	10,399	8,553	8,838
Small Metro	5,519	5,147	5,794	5,490	5,582	5,395	4,703	4,449
Nonmetro	2,798	2,686	2,953	2,920	2,912	2,902	2,381	2,384

*Low precision; no estimate reported.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

NOTE: Response categories for the Adult Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes." Respondents with unknown Adult Involvement data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.17B Adults Always or Sometimes Took Part in Specific School-Related Behaviors in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	ADULT INVOLVEMENT							
	Teacher Let Youth Know He or She Was Doing a Good Job with School Work		Parents Checked That Youth Did Homework		Parents Provided Help with Homework		Parents Limited Time Youth Went Out with Friends on a School Night	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	80.1 ^b	78.3	83.0	83.3	79.4 ^a	80.6	68.4	68.4
AGE								
12-13	85.3	84.3	91.0	90.1	89.1	89.8	71.5	72.7
14-15	79.1 ^b	76.2	84.7	84.8	80.9	81.7	70.1	68.9
16-17	76.4	75.3	74.3	76.1	69.4 ^a	71.6	64.1	64.3
GENDER								
Male	80.4	79.8	85.5	85.9	80.3 ^a	82.0	66.2	65.8
Female	79.7 ^b	76.9	80.4	80.6	78.5	79.1	70.8	71.1
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	79.7 ^b	77.9	82.9	83.1	81.0	81.8	68.0	67.6
White	78.1 ^b	76.0	83.0	83.1	83.0	83.0	68.2 ^a	66.6
Black or African American	85.2	84.4	82.5 ^a	85.2	77.7	79.5	69.4	71.1
American Indian or Alaska Native	85.8	*	82.8	85.2	79.8	84.5	66.5	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	84.1	82.5	81.4	79.5	69.0	73.9	65.5	67.6
Two or More Races	77.4	76.3	84.3 ^a	78.9	80.6	81.9	62.5	69.1
Hispanic or Latino	81.4	79.9	83.4	84.1	73.5	76.1	69.9	71.3
COUNTY TYPE								
Large Metro	80.8 ^a	78.9	82.7	83.3	78.0	79.2	68.6	68.1
Small Metro	79.5	77.9	83.6	83.0	80.7	81.8	68.6	68.4
Nonmetro	78.3	77.2	82.9	84.0	81.8	83.7	67.7	69.8

*Low precision; no estimate reported.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

NOTE: Response categories for the Adult Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes." Respondents with unknown Adult Involvement data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.18A Youths Perceived That All or Most Students in Grade Level Use Specific Substances among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE OF STUDENTS IN GRADE LEVEL							
	Smoke Cigarettes		Use Marijuana or Hashish		Drink Alcoholic Beverages		Get Drunk at Least Once a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	4,094 ^b	3,620	6,286	6,215	7,846	7,529	2,964 ^a	2,697
AGE								
12-13	366	326	362	369	421	371	97	100
14-15	1,456	1,303	2,207	2,070	2,537	2,481	872 ^a	748
16-17	2,272 ^b	1,992	3,717	3,776	4,888	4,676	1,995	1,849
GENDER								
Male	1,629 ^a	1,413	2,834	2,750	3,404	3,159	1,174	1,038
Female	2,464 ^a	2,207	3,452	3,465	4,441	4,369	1,790	1,659
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	3,150 ^b	2,737	4,707	4,618	6,074	5,848	2,265 ^a	2,034
White	2,187 ^b	1,915	3,065	3,013	4,393	4,227	1,595 ^a	1,416
Black or African American	668	585	1,182	1,123	1,126	1,036	471	400
American Indian or Alaska Native	20	16	29	*	34	35	12	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	129	85	202	187	270	270	87	89
Two or More Races	121	112	199	224	217	241	86	99
Hispanic or Latino	944	883	1,579	1,597	1,772	1,681	699	663
COUNTY TYPE								
Large Metro	2,102 ^a	1,864	3,812	3,765	4,471	4,340	1,711	1,593
Small Metro	1,331 ^a	1,160	1,803	1,813	2,265	2,103	858	747
Nonmetro	660	596	671	637	1,110	1,085	395	357

*Low precision; no estimate reported.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months, or reported that they were "home-schooled" in the past 12 months were excluded from this analysis.

NOTE: Response categories for the Substance Use in Grade Level questions include "All of them," "Most of them," "A few of them," and "None of them." The estimates in this table correspond to persons reporting "All of them" or "Most of them." Respondents with unknown Substance Use in Grade Level data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.18B Youths Perceived That All or Most Students in Grade Level Use Specific Substances among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Percentages, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE OF STUDENTS IN GRADE LEVEL							
	Smoke Cigarettes		Use Marijuana or Hashish		Drink Alcoholic Beverages		Get Drunk at Least Once a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	18.4 ^b	16.3	28.4	28.1	35.4 ^a	34.0	13.6 ^a	12.3
AGE								
12-13	5.5	4.9	5.5	5.6	6.4	5.6	1.5	1.5
14-15	18.7	17.0	28.7	27.3	32.9	32.6	11.6 ^a	10.0
16-17	29.0 ^b	25.2	47.4	47.7	62.0 ^a	59.0	25.8 ^a	23.8
GENDER								
Male	14.4 ^b	12.5	25.1	24.4	30.1 ^a	28.0	10.5 ^a	9.3
Female	22.6 ^b	20.2	31.9	32.0	40.9	40.1	16.7	15.5
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	18.0 ^b	15.7	27.2	26.6	34.9	33.6	13.2 ^b	11.8
White	17.5 ^b	15.4	24.7	24.4	35.2	34.1	12.9 ^a	11.5
Black or African American	22.4	19.8	40.1	38.0	38.2	35.2	16.4	13.9
American Indian or Alaska Native	15.7	12.3	23.0	*	26.6	27.0	9.9	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	12.0 ^a	7.6	19.1	17.1	25.4	24.2	8.2	8.1
Two or More Races	17.8	15.6	29.5	31.3	32.0	33.7	12.9	14.0
Hispanic or Latino	19.6	18.4	32.9	33.5	37.0	35.2	14.8	14.1
COUNTY TYPE								
Large Metro	17.3 ^b	14.8	31.4	30.2	36.8 ^a	34.7	14.3 ^a	12.9
Small Metro	20.0	18.2	27.3	28.7	34.3	33.2	13.2	12.0
Nonmetro	19.1	17.9	19.7	19.2	32.4	32.6	11.7	10.9

*Low precision; no estimate reported.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months, or reported that they were "home-schooled" in the past 12 months were excluded from this analysis.

NOTE: Response categories for the Substance Use in Grade Level questions include "All of them," "Most of them," "A few of them," and "None of them." The estimates in this table correspond to persons reporting "All of them" or "Most of them." Respondents with unknown Substance Use in Grade Level data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.19A Past Month Use of Selected Substances, by Perceived Risk of Harm Associated with Substance Use Behaviors among Persons Aged 12 to 17: Numbers in Thousands, 2014

Perceptions of Risk	Used Illicit Drugs¹	Used Marijuana	Used Cigarettes	Binge Use of Alcohol²
CIGARETTES				
Smoke One or More Packs Per Day – Great Risk	1,392	1,103	552	935
Smoke One or More Packs Per Day – Moderate/Slight/No Risk	927	721	654	574
MARIJUANA				
Smoke Once a Month – Great Risk	127	47	77	108
Smoke Once a Month – Moderate/Slight/No Risk	2,198	1,781	1,132	1,400
Smoke Once or Twice a Week – Great Risk	195	63	105	143
Smoke Once or Twice a Week – Moderate/Slight/No Risk	2,131	1,763	1,103	1,364
COCAINE				
Use Once a Month – Great Risk	1,240	1,012	625	826
Use Once a Month – Moderate/Slight/No Risk	1,054	791	564	663
Use Once or Twice a Week – Great Risk	1,698	1,382	869	1,139
Use Once or Twice a Week – Moderate/Slight/No Risk	595	422	320	351
HEROIN				
Try Once or Twice – Great Risk	1,653	1,393	871	1,118
Try Once or Twice – Moderate/Slight/No Risk	646	412	319	371
Use Once or Twice a Week – Great Risk	1,962	1,625	1,028	1,316
Use Once or Twice a Week – Moderate/Slight/No Risk	331	178	160	168
LSD				
Try Once or Twice – Great Risk	829	658	479	598
Try Once or Twice – Moderate/Slight/No Risk	1,406	1,110	677	862
Use Once or Twice a Week – Great Risk	1,353	1,083	681	943
Use Once or Twice a Week – Moderate/Slight/No Risk	885	685	478	517
ALCOHOL				
Have Four or Five Drinks Nearly Every Day – Great Risk	1,116	838	463	613
Have Four or Five Drinks Nearly Every Day – Moderate/Slight/No Risk	1,200	982	736	884
Have Five or More Drinks Once or Twice a Week – Great Risk	707	528	310	340
Have Five or More Drinks Once or Twice a Week – Moderate/Slight/No Risk	1,608	1,291	888	1,163

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Risk data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.19B Past Month Use of Selected Substances, by Perceived Risk of Harm Associated with Substance Use Behaviors among Persons Aged 12 to 17: Percentages, 2014

Perceptions of Risk	Used Illicit Drugs ¹	Used Marijuana	Used Cigarettes	Binge Use of Alcohol ²
CIGARETTES				
Smoke One or More Packs Per Day – Great Risk	8.6	6.8	3.4	5.7
Smoke One or More Packs Per Day – Moderate/Slight/No Risk	11.2	8.7	7.9	6.9
MARIJUANA				
Smoke Once a Month – Great Risk	2.3	0.8	1.4	1.9
Smoke Once a Month – Moderate/Slight/No Risk	11.7	9.5	6.0	7.5
Smoke Once or Twice a Week – Great Risk	2.1	0.7	1.1	1.6
Smoke Once or Twice a Week – Moderate/Slight/No Risk	13.9	11.5	7.2	8.9
COCAINE				
Use Once a Month – Great Risk	10.3	8.4	5.2	6.8
Use Once a Month – Moderate/Slight/No Risk	8.7	6.5	4.6	5.4
Use Once or Twice a Week – Great Risk	9.1	7.4	4.6	6.1
Use Once or Twice a Week – Moderate/Slight/No Risk	10.7	7.6	5.8	6.3
HEROIN				
Try Once or Twice – Great Risk	11.6	9.8	6.1	7.8
Try Once or Twice – Moderate/Slight/No Risk	6.6	4.2	3.2	3.8
Use Once or Twice a Week – Great Risk	10.2	8.4	5.3	6.8
Use Once or Twice a Week – Moderate/Slight/No Risk	6.8	3.7	3.3	3.5
LSD				
Try Once or Twice – Great Risk	7.7	6.1	4.5	5.6
Try Once or Twice – Moderate/Slight/No Risk	11.2	8.9	5.4	6.9
Use Once or Twice a Week – Great Risk	8.5	6.8	4.3	5.9
Use Once or Twice a Week – Moderate/Slight/No Risk	12.0	9.3	6.5	7.0
ALCOHOL				
Have Four or Five Drinks Nearly Every Day – Great Risk	7.3	5.5	3.0	4.0
Have Four or Five Drinks Nearly Every Day – Moderate/Slight/No Risk	13.0	10.6	8.0	9.6
Have Five or More Drinks Once or Twice a Week – Great Risk	7.4	5.5	3.2	3.5
Have Five or More Drinks Once or Twice a Week – Moderate/Slight/No Risk	10.8	8.7	6.0	7.8

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Risk data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.20A Past Month Use of Selected Substances, by Perceived Availability of Selected Illicit Drugs among Persons Aged 12 to 17: Numbers in Thousands, 2014

Perceived Availability	Used Illicit Drugs ¹	Used Marijuana	Used Cigarettes	Binge Use of Alcohol ²
FAIRLY OR VERY EASY TO OBTAIN				
Marijuana	1,982	1,694	1,048	1,313
Cocaine	644	515	351	470
Crack	467	354	257	323
Heroin	347	252	214	240
LSD	607	511	354	431
FAIRLY OR VERY DIFFICULT, OR PROBABLY IMPOSSIBLE TO OBTAIN				
Marijuana	349	136	163	191
Cocaine	1,622	1,258	827	1,008
Crack	1,789	1,410	917	1,141
Heroin	1,906	1,514	962	1,233
LSD	1,643	1,255	811	1,038

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceived Availability data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.20B Past Month Use of Selected Substances, by Perceived Availability of Selected Illicit Drugs among Persons Aged 12 to 17: Percentages, 2014

Perceived Availability	Used Illicit Drugs¹	Used Marijuana	Used Cigarettes	Binge Use of Alcohol²
FAIRLY OR VERY EASY TO OBTAIN				
Marijuana	17.3	14.8	9.1	11.5
Cocaine	19.0	15.2	10.4	13.9
Crack	13.8	10.5	7.6	9.6
Heroin	15.7	11.4	9.7	10.9
LSD	22.6	19.0	13.2	16.1
FAIRLY OR VERY DIFFICULT, OR PROBABLY IMPOSSIBLE TO OBTAIN				
Marijuana	2.8	1.1	1.3	1.5
Cocaine	8.0	6.2	4.1	5.0
Crack	8.9	7.0	4.6	5.7
Heroin	9.0	7.1	4.5	5.8
LSD	8.1	6.2	4.0	5.1

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceived Availability data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.21A Past Month Use of Selected Substances, by Youth's Perceptions of Parents' Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Numbers in Thousands, 2014

Feelings about Substance Use Behaviors	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Strongly Disapprove	Somewhat Disapprove/ Neither Approve Nor Disapprove	Strongly Disapprove	Somewhat Disapprove/ Neither Approve Nor Disapprove	Strongly Disapprove	Somewhat Disapprove/ Neither Approve Nor Disapprove	Strongly Disapprove	Somewhat Disapprove/ Neither Approve Nor Disapprove
YOUTH'S PERCEPTIONS OF PARENTS' FEELINGS ABOUT YOUTH...								
Smoking One or More Packs of Cigarettes Per Day	1,886	390	1,461	328	751	427	1,218	263
Trying Marijuana or Hashish Once or Twice	1,315	953	918	863	664	510	942	535
Using Marijuana or Hashish Once a Month or More	1,382	887	982	800	698	478	981	494
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	1,736	547	1,326	467	798	379	1,073	411

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Parents' Feelings data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.21B Past Month Use of Selected Substances, by Youth's Perceptions of Parents' Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Percentages, 2014

Feelings about Substance Use Behaviors	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Strongly Disapprove	Somewhat Disapprove/ Approve Nor Neither Disapprove	Strongly Disapprove	Somewhat Disapprove/ Approve Nor Neither Disapprove	Strongly Disapprove	Somewhat Disapprove/ Approve Nor Neither Disapprove	Strongly Disapprove	Somewhat Disapprove/ Approve Nor Neither Disapprove
YOUTH'S PERCEPTIONS OF PARENTS' FEELINGS ABOUT YOUTH...								
Smoking One or More Packs of Cigarettes Per Day	8.2	25.6	6.4	21.5	3.3	28.1	5.3	17.3
Trying Marijuana or Hashish Once or Twice	6.1	31.1	4.3	28.1	3.1	16.6	4.4	17.4
Using Marijuana or Hashish Once a Month or More	6.3	36.1	4.5	32.5	3.2	19.4	4.5	20.1
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	7.8	23.6	6.0	20.1	3.6	16.4	4.8	17.8

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Parents' Feelings data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.22A Past Month Use of Selected Substances, by Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Numbers in Thousands, 2014

Feelings about Substance Use Behaviors	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove
YOUTH'S PERCEPTIONS OF CLOSE FRIENDS' FEELINGS ABOUT YOUTH...								
Smoking One or More Packs of Cigarettes Per Day	1,769	515	1,348	445	716	460	1,148	336
Trying Marijuana or Hashish Once or Twice	745	1,538	414	1,381	376	801	545	940
Using Marijuana or Hashish Once a Month or More	721	1,558	396	1,396	369	804	511	971
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	1,494	791	1,112	684	612	565	887	602
YOUTH'S FEELINGS ABOUT PEERS...								
Smoking One or More Packs of Cigarettes Per Day	1,897	388	1,468	327	784	393	1,209	278
Trying Marijuana or Hashish Once or Twice	798	1,479	453	1,340	432	743	601	885
Using Marijuana or Hashish Once a Month or More	716	1,556	388	1,403	423	751	537	941
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	1,587	687	1,204	587	702	473	980	504

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Close Friends' Feelings or Feelings about Peers data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.22B Past Month Use of Selected Substances, by Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Percentages, 2014

Feelings about Substance Use Behaviors	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove
YOUTH'S PERCEPTIONS OF CLOSE FRIENDS' FEELINGS ABOUT YOUTH...								
Smoking One or More Packs of Cigarettes Per Day	8.1	20.8	6.1	18.0	3.3	18.5	5.2	13.6
Trying Marijuana or Hashish Once or Twice	3.9	29.2	2.2	26.2	2.0	15.2	2.8	17.8
Using Marijuana or Hashish Once a Month or More	3.7	31.7	2.0	28.4	1.9	16.4	2.6	19.8
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	7.0	25.7	5.2	22.2	2.9	18.3	4.1	19.5
YOUTH'S FEELINGS ABOUT PEERS...								
Smoking One or More Packs of Cigarettes Per Day	8.3	21.2	6.5	17.9	3.4	21.4	5.3	15.2
Trying Marijuana or Hashish Once or Twice	4.1	29.4	2.3	26.6	2.2	14.8	3.1	17.6
Using Marijuana or Hashish Once a Month or More	3.7	30.5	2.0	27.5	2.2	14.7	2.8	18.4
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	7.2	27.1	5.5	23.2	3.2	18.7	4.5	19.9

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Close Friends' Feelings or Feelings about Peers data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.23A Past Month Use of Selected Substances, by Measures of Youth's Exposure to Prevention Programs and Messages and Youth's Delinquency in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2014

Prevention Programs/Messages and Delinquency	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior
EXPOSURE TO PREVENTION PROGRAMS AND MESSAGES IN PAST YEAR								
Participated in Problem-Solving, Communication Skills, or Self-Esteem Group	456	1,822	317	1,479	214	965	258	1,223
Participated in Violence Prevention Program	269	2,022	182	1,619	121	1,058	148	1,340
Participated in Pregnancy or Sexually Transmitted Disease Prevention Program	227	2,069	172	1,630	121	1,058	103	1,387
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School	257	2,037	193	1,608	131	1,049	147	1,341
Participated in Program to Help Deal with Drug or Alcohol Use	136	2,148	109	1,688	80	1,095	61	1,422
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School	1,624	660	1,273	520	799	379	1,059	424
Talked to Parents about Dangers of Tobacco, Alcohol, or Drug Use	1,262	1,018	1,013	783	622	555	811	671
DELINQUENT BEHAVIOR IN PAST YEAR³								
Got into a Serious Fight at School or Work	676	1,626	506	1,298	381	809	413	1,074
Took Part in a Group-Against-Group Fight	542	1,759	429	1,375	260	929	326	1,157
Attacked Someone with the Intent to Seriously Hurt Them	262	2,037	178	1,623	151	1,039	153	1,337
Carried a Handgun	151	2,152	132	1,668	103	1,087	120	1,371
Sold Illegal Drugs	323	1,977	310	1,488	173	1,014	214	1,267
Stole or Tried to Steal Anything Worth More Than \$50	265	2,036	228	1,571	138	1,047	173	1,317

*Low precision; no estimate reported.

NOTE: Respondents with unknown Prevention Programs and Messages Exposure or Delinquent Behavior data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.23B Past Month Use of Selected Substances, by Measures of Youth's Exposure to Prevention Programs and Messages and Youth's Delinquency in the Past Year among Persons Aged 12 to 17: Percentages, 2014

Prevention Programs/Messages and Delinquency	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior
EXPOSURE TO PREVENTION PROGRAMS AND MESSAGES IN PAST YEAR								
Participated in Problem-Solving, Communication Skills, or Self-Esteem Group	8.2	9.8	5.7	7.9	3.9	5.2	4.7	6.5
Participated in Violence Prevention Program	8.7	9.4	5.9	7.5	3.9	4.9	4.8	6.2
Participated in Pregnancy or Sexually Transmitted Disease Prevention Program	10.6	9.2	8.0	7.2	5.7	4.7	4.8	6.2
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School	9.4	9.3	7.0	7.3	4.8	4.8	5.4	6.1
Participated in Program to Help Deal with Drug or Alcohol Use	13.6	9.1	10.9	7.2	8.0	4.6	6.1	6.0
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School	9.1	9.9	7.1	7.8	4.5	5.7	5.9	6.4
Talked to Parents about Dangers of Tobacco, Alcohol, or Drug Use	9.2	9.5	7.4	7.3	4.5	5.2	5.9	6.3
DELINQUENT BEHAVIOR IN PAST YEAR³								
Got into a Serious Fight at School or Work	16.2	7.9	12.2	6.3	9.1	3.9	9.9	5.2
Took Part in a Group-Against-Group Fight	20.7	8.0	16.4	6.2	9.9	4.2	12.5	5.2
Attacked Someone with the Intent to Seriously Hurt Them	21.9	8.6	14.9	6.9	12.6	4.4	12.8	5.7
Carried a Handgun	16.0	9.0	13.9	7.0	10.9	4.6	12.7	5.8
Sold Illegal Drugs	64.3	8.1	61.8	6.1	34.5	4.2	42.6	5.2
Stole or Tried to Steal Anything Worth More Than \$50	40.8	8.4	35.1	6.5	21.2	4.3	26.6	5.5

*Low precision; no estimate reported.

NOTE: Respondents with unknown Prevention Programs and Messages Exposure or Delinquent Behavior data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.24A Past Month Use of Selected Substances, by Measures of Risk and Protective Factors, Religious Beliefs, and Past Year Parental Involvement among Persons Aged 12 to 17: Numbers in Thousands, 2014

Risk and Protective Factors, Religious Beliefs, and Parental Involvement	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement
RISK AND PROTECTIVE FACTORS								
Youth Moved 2 or More Times in the Past 5 Years	708	1,559	538	1,245	397	777	378	1,095
Youth Argued with at Least One Parent 10 or More Times in the Past Year	789	1,509	614	1,191	398	791	530	956
Youth Was Approached by Someone Selling Drugs in the Past Month	1,007	1,322	886	940	510	697	640	869
Youth Had Someone to Talk to about Problems ³	2,059	181	1,618	145	1,053	102	1,359	110
Youth Attended 25 or More Religious Services in the Past Year ⁴	448	1,827	314	1,473	191	978	261	1,210
YOUTH RELIGIOUS BELIEFS⁵								
Religious Beliefs Are a Very Important Part of Youth's Life	1,235	1,005	931	833	570	584	818	642
Religious Beliefs Influence Youth's Decision Making in Life	1,022	1,234	742	1,027	464	697	603	863
It Is Important for Friends to Share Religious Beliefs	462	1,790	298	1,479	199	957	247	1,212
PARENTAL INVOLVEMENT IN PAST YEAR⁶								
Parents Made Youth Do Chores around the House	1,981	326	1,544	265	1,047	148	1,283	209
Parents Limited Amount of Time Youth Watched TV	583	1,722	409	1,398	273	920	351	1,145
Parents Let Youth Know He or She Had Done a Good Job	1,771	535	1,361	446	893	300	1,146	347
Parents Let Youth Know They Were Proud of Something He or She Had Done	1,747	552	1,351	457	887	307	1,139	355

*Low precision; no estimate reported.

NOTE: Respondents with unknown Risk and Protective Factors, Religious Beliefs, or Parental Involvement data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Respondents were able to select multiple categories of persons that they would talk to about a serious problem, including the choice of "nobody." Respondents who reported both a person that they would talk to and that they had nobody to talk to were excluded from the analysis.

⁴ Special occasions, such as weddings, funerals, or other special events, were not to have been considered in the number of times attended religious services.

⁵ Response categories for the questions on Religious Beliefs include "Strongly disagree," "Disagree," "Agree," and "Strongly agree." The estimates in this table correspond to persons reporting "Agree" or "Strongly agree."

⁶ Response categories for the Parental Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.24B Past Month Use of Selected Substances, by Measures of Risk and Protective Factors, Religious Beliefs, and Past Year Parental Involvement among Persons Aged 12 to 17: Percentages, 2014

Risk and Protective Factors, Religious Beliefs, and Parental Involvement	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement
RISK AND PROTECTIVE FACTORS								
Youth Moved 2 or More Times in the Past 5 Years	12.1	8.5	9.2	6.8	6.8	4.2	6.5	5.9
Youth Argued with at Least One Parent 10 or More Times in the Past Year	16.0	7.7	12.4	6.1	8.1	4.1	10.7	4.9
Youth Was Approached by Someone Selling Drugs in the Past Month	33.4	6.1	29.4	4.3	16.9	3.2	21.2	4.0
Youth Had Someone to Talk to about Problems ³	8.9	16.0	7.0	12.8	4.6	9.0	5.9	9.7
Youth Attended 25 or More Religious Services in the Past Year ⁴	6.4	10.7	4.5	8.6	2.7	5.7	3.7	7.1
YOUTH RELIGIOUS BELIEFS⁵								
Religious Beliefs Are a Very Important Part of Youth's Life	7.1	14.8	5.4	12.3	3.3	8.6	4.7	9.4
Religious Beliefs Influence Youth's Decision Making in Life	6.6	14.3	4.8	11.9	3.0	8.1	3.9	10.0
It Is Important for Friends to Share Religious Beliefs	6.3	10.7	4.1	8.8	2.7	5.7	3.4	7.2
PARENTAL INVOLVEMENT IN PAST YEAR⁶								
Parents Made Youth Do Chores around the House	8.9	12.6	7.0	10.2	4.7	5.7	5.8	8.1
Parents Limited Amount of Time Youth Watched TV	5.8	11.9	4.0	9.6	2.7	6.3	3.5	7.9
Parents Let Youth Know He or She Had Done a Good Job	8.3	15.9	6.4	13.2	4.2	8.9	5.4	10.3
Parents Let Youth Know They Were Proud of Something He or She Had Done	8.2	16.0	6.4	13.2	4.2	8.9	5.4	10.3

*Low precision; no estimate reported.

NOTE: Respondents with unknown Risk and Protective Factors, Religious Beliefs, or Parental Involvement data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Respondents were able to select multiple categories of persons that they would talk to about a serious problem, including the choice of "nobody." Respondents who reported both a person that they would talk to and that they had nobody to talk to were excluded from the analysis.

⁴ Special occasions, such as weddings, funerals, or other special events, were not to have been considered in the number of times attended religious services.

⁵ Response categories for the questions on Religious Beliefs include "Strongly disagree," "Disagree," "Agree," and "Strongly agree." The estimates in this table correspond to persons reporting "Agree" or "Strongly agree."

⁶ Response categories for the Parental Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.25A Past Month Use of Selected Substances, by Measures of Scholastic Environment among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year: Numbers in Thousands, 2014

Scholastic Environment	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic
STUDENT CHARACTERISTICS IN PAST YEAR³								
Participated in One or Fewer Activities ⁴	424	1,784	373	1,372	289	847	272	1,154
Drug or Alcohol Prevention Messages Were Seen or Heard in School ⁵	1,426	768	1,117	619	660	465	916	496
Youth Liked a Lot or Kind of Liked Going to School ⁶	1,487	734	1,159	596	701	442	916	518
Grade Average for Last Completed Period Was "D" or Lower ⁷	181	1,961	154	1,538	157	949	138	1,261
ADULT INVOLVEMENT IN PAST YEAR^{3,8}								
Teacher Let Youth Know He or She Was Doing a Good Job with School Work	1,572	645	1,228	523	765	371	1,021	408
Parents Checked That Youth Did Homework	1,520	692	1,157	590	758	378	947	478
Parents Provided Help with Homework	1,465	749	1,116	633	792	350	924	503
Parents Limited Time Youth Went Out with Friends on a School Night	1,370	834	1,092	656	743	397	878	554
SUBSTANCE USE IN GRADE LEVEL⁹								
Students in Youth's Grade Smoke Cigarettes	629	1,537	505	1,205	480	639	418	987
Students in Youth's Grade Use Marijuana or Hashish	1,451	734	1,291	441	711	410	926	494
Students in Youth's Grade Drink Alcoholic Beverages	1,381	787	1,185	529	740	373	1,091	330
Students in Youth's Grade Get Drunk at Least Once a Week	602	1,545	523	1,178	359	752	463	943

*Low precision; no estimate reported.

NOTE: Respondents with unknown Student Characteristics, Adult Involvement, or Substance Use in Grade Level data were excluded from the respective analyses.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

⁴ The number of activities participated in is based on four questions: participation in school-based activities, community-based activities, church or faith-based activities, and other activities. For each question, the response categories include "None," "One," "Two," and "3 or more." Respondents were excluded from the analysis if they had unknown data for at least three of the questions and if their known responses did not place them in the recoded "Two or more" category.

⁵ Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

⁶ Response categories for the School Satisfaction question include "You liked going to school a lot," "You kind of liked going to school," "You didn't like going to school very much," and "You hated going to school." The estimates in this table correspond to persons reporting "You liked going to school a lot" or "You kind of liked going to school."

⁷ Response categories for Grade Average question include "An 'A+', 'A,' or 'A-minus' average"; "A 'B+', 'B,' or 'B-minus' average"; "A 'C+', 'C,' or 'C-minus' average"; "A 'D' or less than a 'D' average"; and "My school does not give these grades." The estimates in this table correspond to persons reporting "A 'D' or less than a 'D' average." Respondents reporting that their school does not give these grades were excluded from the analysis.

⁸ Response categories for the Adult Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes."

⁹ Response categories for the Substance Use in Grade Level questions include "All of them," "Most of them," "A few of them," and "None of them." The estimates in this table correspond to persons reporting "All of them" or "Most of them." Youths reporting that they were "home-schooled" in the past 12 months were not considered to be enrolled in school.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.25B Past Month Use of Selected Substances, by Measures of Scholastic Environment among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year: Percentages, 2014

Scholastic Environment	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic
STUDENT CHARACTERISTICS IN PAST YEAR³								
Participated in One or Fewer Activities ⁴	14.8	8.7	13.0	6.7	10.1	4.2	9.5	5.7
Drug or Alcohol Prevention Messages Were Seen or Heard in School ⁵	8.5	12.3	6.6	9.9	3.9	7.4	5.4	7.9
Youth Liked a Lot or Kind of Liked Going to School ⁶	8.0	15.5	6.3	12.6	3.8	9.3	4.9	10.9
Grade Average for Last Completed Period Was "D" or Lower ⁷	18.7	9.3	15.9	7.3	16.2	4.5	14.2	6.0
ADULT INVOLVEMENT IN PAST YEAR^{3,8}								
Teacher Let Youth Know He or She Was Doing a Good Job with School Work	8.6	12.8	6.7	10.4	4.2	7.4	5.6	8.1
Parents Checked That Youth Did Homework	7.9	17.9	6.0	15.2	3.9	9.8	4.9	12.3
Parents Provided Help with Homework	7.8	16.6	6.0	14.0	4.2	7.8	4.9	11.2
Parents Limited Time Youth Went Out with Friends on a School Night	8.7	11.5	7.0	9.1	4.7	5.5	5.6	7.6
SUBSTANCE USE IN GRADE LEVEL⁹								
Students in Youth's Grade Smoke Cigarettes	17.4	8.3	14.0	6.5	13.3	3.4	11.5	5.3
Students in Youth's Grade Use Marijuana or Hashish	23.3	4.6	20.8	2.8	11.4	2.6	14.9	3.1
Students in Youth's Grade Drink Alcoholic Beverages	18.3	5.4	15.7	3.6	9.8	2.5	14.5	2.3
Students in Youth's Grade Get Drunk at Least Once a Week	22.3	8.1	19.4	6.1	13.3	3.9	17.2	4.9

*Low precision; no estimate reported.

NOTE: Respondents with unknown Student Characteristics, Adult Involvement, or Substance Use in Grade Level data were excluded from the respective analyses.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

⁴ The number of activities participated in is based on four questions: participation in school-based activities, community-based activities, church or faith-based activities, and other activities. For each question, the response categories include "None," "One," "Two," and "3 or more." Respondents were excluded from the analysis if they had unknown data for at least three of the questions and if their known responses did not place them in the recoded "Two or more" category.

⁵ Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

⁶ Response categories for the School Satisfaction question include "You liked going to school a lot," "You kind of liked going to school," "You didn't like going to school very much," and "You hated going to school." The estimates in this table correspond to persons reporting "You liked going to school a lot" or "You kind of liked going to school."

⁷ Response categories for Grade Average question include "An 'A+', 'A,' or 'A-minus' average"; "A 'B+', 'B,' or 'B-minus' average"; "A 'C+', 'C,' or 'C-minus' average"; "A 'D' or less than a 'D' average"; and "My school does not give these grades." The estimates in this table correspond to persons reporting "A 'D' or less than a 'D' average." Respondents reporting that their school does not give these grades were excluded from the analysis.

⁸ Response categories for the Adult Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes."

⁹ Response categories for the Substance Use in Grade Level questions include "All of them," "Most of them," "A few of them," and "None of them." The estimates in this table correspond to persons reporting "All of them" or "Most of them." Youths reporting that they were "home-schooled" in the past 12 months were not considered to be enrolled in school.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.1C Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Risk/Availability	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
PERCEPTIONS OF GREAT RISK¹								
Cigarettes								
Smoke One or More Packs Per Day	857	671	109	113	149	160	827	634
Marijuana								
Smoke Once a Month	993	737	97	98	122	122	961	702
Smoke Once or Twice a Week	1,050	820	110	120	143	137	1,028	782
Cocaine								
Use Once a Month	845	762	108	115	160	180	816	707
Use Once or Twice a Week	593	514	97	95	112	128	570	480
Heroin								
Try Once or Twice	692	548	111	115	129	146	668	499
Use Once or Twice a Week	431	357	89	97	80	87	403	330
LSD								
Try Once or Twice	867	754	109	116	167	188	832	684
Use Once or Twice a Week	637	547	104	110	133	155	608	509
Alcohol								
Have Four or Five Drinks Nearly Every Day	889	733	106	121	156	166	859	702
Have Five or More Drinks Once or Twice a Week	967	810	109	116	151	165	939	764
PERCEIVED AVAILABILITY²								
Fairly or Very Easy to Obtain ³								
Marijuana	1,015	834	111	118	153	161	975	793
Cocaine	833	658	79	83	149	158	804	627
Crack	795	630	80	82	126	139	780	604
Heroin	703	575	64	68	113	120	691	552
LSD	636	514	70	76	122	131	626	488
Approached in the Past Month by Someone Selling Drugs	415	336	79	87	121	142	370	276

*Low precision; no estimate reported.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.1D Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances, by Age Group: Standard Errors of Percentages, 2013 and 2014

Risk/Availability	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
PERCEPTIONS OF GREAT RISK¹								
Cigarettes								
Smoke One or More Packs Per Day	0.33	0.25	0.44	0.46	0.43	0.46	0.41	0.31
Marijuana								
Smoke Once a Month	0.39	0.28	0.40	0.40	0.35	0.35	0.48	0.35
Smoke Once or Twice a Week	0.41	0.32	0.45	0.49	0.41	0.40	0.52	0.39
Cocaine								
Use Once a Month	0.33	0.29	0.45	0.47	0.47	0.52	0.41	0.35
Use Once or Twice a Week	0.23	0.20	0.40	0.39	0.33	0.37	0.29	0.24
Heroin								
Try Once or Twice	0.27	0.21	0.46	0.48	0.37	0.42	0.34	0.25
Use Once or Twice a Week	0.17	0.14	0.37	0.40	0.23	0.25	0.20	0.16
LSD								
Try Once or Twice	0.34	0.29	0.47	0.50	0.49	0.55	0.42	0.34
Use Once or Twice a Week	0.25	0.21	0.45	0.47	0.39	0.45	0.31	0.25
Alcohol								
Have Four or Five Drinks Nearly Every Day	0.34	0.28	0.43	0.49	0.45	0.48	0.43	0.34
Have Five or More Drinks Once or Twice a Week	0.37	0.31	0.45	0.47	0.44	0.48	0.47	0.38
PERCEIVED AVAILABILITY²								
Fairly or Very Easy to Obtain ³								
Marijuana	0.41	0.33	0.46	0.49	0.45	0.47	0.51	0.41
Cocaine	0.34	0.26	0.33	0.35	0.44	0.47	0.43	0.33
Crack	0.32	0.25	0.34	0.35	0.38	0.41	0.41	0.32
Heroin	0.29	0.23	0.28	0.29	0.34	0.36	0.37	0.29
LSD	0.26	0.21	0.30	0.33	0.37	0.39	0.33	0.26
Approached in the Past Month by Someone Selling Drugs	0.16	0.13	0.32	0.35	0.35	0.41	0.18	0.13

*Low precision; no estimate reported.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.1P Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Risk/Availability	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
PERCEPTIONS OF GREAT RISK¹								
Cigarettes								
Smoke One or More Packs Per Day	0.4131		0.0014		0.1784		0.9522	
Marijuana								
Smoke Once a Month	0.0004		0.0179		0.0066		0.0021	
Smoke Once or Twice a Week	0.0000		0.0016		0.0000		0.0000	
Cocaine								
Use Once a Month	0.1640		0.4543		0.9133		0.0961	
Use Once or Twice a Week	0.0008		0.0279		0.8420		0.0013	
Heroin								
Try Once or Twice	0.0109		0.1391		0.0877		0.0527	
Use Once or Twice a Week	0.7837		0.7477		0.0712		0.3953	
LSD								
Try Once or Twice	0.1181		0.1810		0.0406		0.2747	
Use Once or Twice a Week	0.0006		0.0337		0.6864		0.0014	
Alcohol								
Have Four or Five Drinks Nearly Every Day	0.0691		0.9015		0.7369		0.0491	
Have Five or More Drinks Once or Twice a Week	0.0327		0.7098		0.5966		0.0166	
PERCEIVED AVAILABILITY²								
Fairly or Very Easy to Obtain ³								
Marijuana	0.0020		0.2041		0.5722		0.0005	
Cocaine	0.2750		0.8839		0.5734		0.2111	
Crack	0.1915		0.2148		0.1821		0.0865	
Heroin	0.0000		0.4668		0.9254		0.0000	
LSD	0.8577		0.5077		0.7395		0.9809	
Approached in the Past Month by Someone Selling Drugs	0.3506		0.5288		0.8556		0.4586	

*Low precision; no estimate reported.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.2C Perceived Great Risk of Harm Associated with Cigarette and Alcohol Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	CIGARETTES		ALCOHOL			
	Smoke One or More Packs Per Day		Have Four or Five Drinks Nearly Every Day		Have Five or More Drinks Once or Twice a Week	
	2013	2014	2013	2014	2013	2014
TOTAL	109	113	106	121	109	116
AGE						
12-13	115	111	118	111	87	87
14-15	117	120	115	118	85	92
16-17	112	126	111	117	85	90
GENDER						
Male	78	83	76	88	78	87
Female	73	78	72	81	74	80
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	94	96	89	101	94	101
White	182	169	177	163	116	110
Black or African American	83	85	88	89	69	76
American Indian or Alaska Native	10	15	12	12	9	11
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	68	63	64	69	49	55
Two or More Races	34	36	35	37	25	23
Hispanic or Latino	56	60	60	61	56	59
COUNTY TYPE						
Large Metro	192	188	189	192	127	143
Small Metro	143	117	141	124	101	87
Nonmetro	82	90	93	82	66	60

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.2D Perceived Great Risk of Harm Associated with Cigarette and Alcohol Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	CIGARETTES		ALCOHOL			
	Smoke One or More Packs Per Day		Have Four or Five Drinks Nearly Every Day		Have Five or More Drinks Once or Twice a Week	
	2013	2014	2013	2014	2013	2014
TOTAL	0.44	0.46	0.43	0.49	0.45	0.47
AGE						
12-13	0.80	0.86	0.76	0.82	0.77	0.84
14-15	0.71	0.81	0.71	0.87	0.77	0.81
16-17	0.75	0.80	0.77	0.81	0.79	0.80
GENDER						
Male	0.62	0.66	0.61	0.70	0.62	0.69
Female	0.61	0.65	0.59	0.67	0.62	0.66
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.50	0.50	0.47	0.53	0.49	0.53
White	0.55	0.59	0.57	0.63	0.56	0.59
Black or African American	1.25	1.25	1.16	1.19	1.11	1.32
American Indian or Alaska Native	4.48	5.61	4.92	4.86	4.32	5.62
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.21	2.40	2.16	2.27	2.55	2.66
Two or More Races	2.23	2.16	2.24	2.21	2.56	2.26
Hispanic or Latino	1.02	1.07	1.11	1.09	1.03	1.06
COUNTY TYPE						
Large Metro	0.63	0.64	0.59	0.70	0.62	0.67
Small Metro	0.73	0.77	0.79	0.86	0.84	0.81
Nonmetro	1.01	1.11	1.23	1.12	1.04	1.04

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.2P Perceived Great Risk of Harm Associated with Cigarette and Alcohol Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	CIGARETTES		ALCOHOL			
	Smoke One or More Packs Per Day		Have Four or Five Drinks Nearly Every Day		Have Five or More Drinks Once or Twice a Week	
	2013	2014	2013	2014	2013	2014
TOTAL	0.0014		0.9015		0.7098	
AGE						
12-13	0.0136		0.9768		0.4037	
14-15	0.6164		0.1719		0.7888	
16-17	0.0144		0.2544		0.2395	
GENDER						
Male	0.0187		0.9534		0.8631	
Female	0.0296		0.9195		0.4644	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0058		0.5905		0.8717	
White	0.0047		0.5913		0.7451	
Black or African American	0.2608		0.9218		0.8806	
American Indian or Alaska Native	0.4729		0.3075		0.7717	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4113		0.6264		0.4656	
Two or More Races	0.6740		0.8774		0.5919	
Hispanic or Latino	0.1008		0.3037		0.3371	
COUNTY TYPE						
Large Metro	0.0096		0.6634		0.9038	
Small Metro	0.2706		0.4648		0.6911	
Nonmetro	0.1133		0.4408		0.5798	

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.3C Perceived Great Risk of Harm Associated with Marijuana and Cocaine Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	MARIJUANA				COCAINE			
	Smoke Once a Month		Smoke Once or Twice a Week		Use Once a Month		Use Once or Twice a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	97	98	110	120	108	115	97	95
AGE								
12-13	78	74	108	98	84	87	121	114
14-15	59	64	86	87	95	98	132	127
16-17	57	59	70	71	103	116	130	138
GENDER								
Male	67	68	80	83	77	84	65	68
Female	67	70	75	85	76	79	67	70
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	83	84	93	103	96	98	81	80
White	92	89	136	120	147	139	211	187
Black or African American	43	45	58	58	71	73	91	94
American Indian or Alaska Native	6	6	9	12	10	16	13	17
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	37	36	47	53	47	44	68	68
Two or More Races	17	20	22	27	29	30	37	37
Hispanic or Latino	51	51	61	55	56	59	53	56
COUNTY TYPE								
Large Metro	96	99	139	135	141	152	219	206
Small Metro	73	64	100	84	115	101	165	136
Nonmetro	48	53	65	67	73	75	98	100

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.3D Perceived Great Risk of Harm Associated with Marijuana and Cocaine Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	MARIJUANA				COCAINE			
	Smoke Once a Month		Smoke Once or Twice a Week		Use Once a Month		Use Once or Twice a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.40	0.40	0.45	0.49	0.45	0.47	0.40	0.39
AGE								
12-13	0.73	0.79	0.82	0.89	0.78	0.84	0.72	0.82
14-15	0.60	0.67	0.73	0.80	0.75	0.84	0.69	0.69
16-17	0.61	0.63	0.68	0.71	0.80	0.82	0.58	0.58
GENDER								
Male	0.54	0.55	0.64	0.67	0.63	0.68	0.52	0.55
Female	0.56	0.59	0.63	0.71	0.64	0.67	0.56	0.58
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.44	0.45	0.49	0.54	0.51	0.52	0.43	0.42
White	0.50	0.54	0.56	0.66	0.59	0.63	0.44	0.51
Black or African American	0.98	1.07	1.14	1.25	1.21	1.24	1.15	1.11
American Indian or Alaska Native	3.95	3.02	4.61	5.25	3.74	4.92	3.87	3.50
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	2.15	2.29	2.37	2.53	2.52	2.23	1.93	2.13
Two or More Races	2.05	2.29	2.48	2.56	2.49	2.50	1.76	1.96
Hispanic or Latino	0.95	0.92	1.12	1.00	1.04	1.06	0.98	1.01
COUNTY TYPE								
Large Metro	0.54	0.59	0.63	0.71	0.64	0.67	0.54	0.56
Small Metro	0.74	0.69	0.79	0.81	0.84	0.82	0.75	0.69
Nonmetro	0.90	1.05	1.00	1.10	1.12	1.14	0.94	0.98

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.3P Perceived Great Risk of Harm Associated with Marijuana and Cocaine Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	MARIJUANA				COCAINE			
	Smoke Once a Month		Smoke Once or Twice a Week		Use Once a Month		Use Once or Twice a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.0179		0.0016		0.4543		0.0279	
AGE								
12-13	0.0103		0.0377		0.9103		0.1244	
14-15	0.4089		0.1633		0.6387		0.0680	
16-17	0.5814		0.0331		0.1497		0.6057	
GENDER								
Male	0.1098		0.0051		0.5086		0.0062	
Female	0.0744		0.0882		0.7012		0.6749	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.0043		0.0006		0.6671		0.0941	
White	0.0296		0.0001		0.4735		0.0661	
Black or African American	0.5014		0.7684		0.8980		0.3900	
American Indian or Alaska Native	0.1537		0.8036		0.1131		0.5418	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.1050		0.1860		0.9847		0.1854	
Two or More Races	0.4785		0.8372		0.1317		0.1942	
Hispanic or Latino	0.8506		0.6939		0.4272		0.1887	
COUNTY TYPE								
Large Metro	0.0068		0.0035		0.7601		0.0044	
Small Metro	0.6011		0.1635		0.5413		0.7011	
Nonmetro	0.8028		0.7631		0.5379		0.5740	

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.4C Perceived Great Risk of Harm Associated with Heroin and LSD Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	HEROIN				LSD			
	Try Once or Twice		Use Once or Twice a Week		Try Once or Twice		Use Once or Twice a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	111	115	89	97	109	116	104	110
AGE								
12-13	90	93	120	116	85	81	110	103
14-15	102	108	127	131	90	94	119	115
16-17	117	129	135	145	94	105	122	126
GENDER								
Male	78	81	62	67	79	87	71	75
Female	78	79	64	68	76	78	72	77
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	97	100	76	80	91	100	88	93
White	156	151	214	192	131	127	191	169
Black or African American	79	82	90	95	70	70	84	83
American Indian or Alaska Native	11	19	13	20	9	13	10	14
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	51	52	67	68	42	41	59	64
Two or More Races	30	34	37	39	25	26	31	32
Hispanic or Latino	53	58	47	53	57	56	53	54
COUNTY TYPE								
Large Metro	161	172	217	218	136	137	196	186
Small Metro	131	115	166	137	110	95	141	122
Nonmetro	82	83	101	101	70	75	95	92

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.4D Perceived Great Risk of Harm Associated with Heroin and LSD Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	HEROIN				LSD			
	Try Once or Twice		Use Once or Twice a Week		Try Once or Twice		Use Once or Twice a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.46	0.48	0.37	0.40	0.47	0.50	0.45	0.47
AGE								
12-13	0.85	0.88	0.78	0.84	0.85	0.88	0.84	0.89
14-15	0.77	0.84	0.62	0.68	0.79	0.86	0.75	0.79
16-17	0.73	0.77	0.51	0.51	0.77	0.85	0.67	0.76
GENDER								
Male	0.64	0.66	0.51	0.55	0.67	0.73	0.60	0.63
Female	0.66	0.67	0.54	0.57	0.67	0.69	0.63	0.68
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.52	0.54	0.41	0.43	0.51	0.55	0.49	0.51
White	0.59	0.66	0.43	0.49	0.58	0.68	0.50	0.60
Black or African American	1.20	1.15	1.11	1.12	1.23	1.24	1.20	1.26
American Indian or Alaska Native	4.35	5.24	4.36	3.36	4.52	5.62	4.70	5.73
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	2.41	2.43	1.98	2.12	2.33	2.31	2.51	2.48
Two or More Races	2.59	2.65	2.05	2.02	2.62	2.69	2.56	2.33
Hispanic or Latino	1.00	1.07	0.89	0.96	1.09	1.06	1.02	1.03
COUNTY TYPE								
Large Metro	0.65	0.66	0.51	0.57	0.69	0.69	0.59	0.66
Small Metro	0.83	0.87	0.72	0.68	0.86	0.88	0.80	0.82
Nonmetro	1.11	1.06	0.85	0.93	1.07	1.33	0.96	1.12

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.4P Perceived Great Risk of Harm Associated with Heroin and LSD Use among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	HEROIN				LSD			
	Try Once or Twice		Use Once or Twice a Week		Try Once or Twice		Use Once or Twice a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.1391		0.7477		0.1810		0.0337	
AGE								
12-13	0.3791		0.9914		0.9458		0.3817	
14-15	0.9073		0.6037		0.1020		0.2743	
16-17	0.0861		0.2537		0.3977		0.0464	
GENDER								
Male	0.5593		0.4061		0.5675		0.0257	
Female	0.1264		0.2092		0.1794		0.3865	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.5016		0.8849		0.1453		0.0104	
White	0.5686		0.7006		0.0870		0.0043	
Black or African American	0.6663		0.5565		0.7304		0.7406	
American Indian or Alaska Native	0.1723		0.0099		0.8684		0.7486	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.9908		0.6224		0.6513		0.7646	
Two or More Races	0.9899		0.6925		0.7955		0.9423	
Hispanic or Latino	0.0720		0.3578		0.8280		0.8815	
COUNTY TYPE								
Large Metro	0.2922		0.8190		0.1763		0.0065	
Small Metro	0.1596		0.6459		0.5867		0.8703	
Nonmetro	0.8991		0.7556		0.8509		0.6550	

*Low precision; no estimate reported.

NOTE: Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.5C Perceived That Obtaining Specific Illicit Drugs Would Be Fairly or Very Easy among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/ Geographic Characteristic	Marijuana (2013)	Marijuana (2014)	Cocaine (2013)	Cocaine (2014)	Crack (2013)	Crack (2014)	Heroin (2013)	Heroin (2014)	LSD (2013)	LSD (2014)
TOTAL	111	118	79	83	80	82	64	68	70	76
AGE										
12-13	59	58	36	39	39	44	29	34	27	29
14-15	99	99	53	51	60	51	44	40	42	44
16-17	120	126	60	65	53	60	41	49	55	63
GENDER										
Male	80	87	49	51	51	50	42	45	46	47
Female	76	81	59	63	60	64	48	51	53	58
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	97	102	67	71	71	70	58	61	60	65
White	143	135	65	67	70	64	56	54	57	63
Black or African American	64	72	33	35	34	41	24	31	25	28
American Indian or Alaska Native	10	13	4	4	6	4	6	3	4	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*	*	*
Asian	41	40	18	21	18	21	11	17	15	15
Two or More Races	27	29	14	17	14	19	11	11	13	11
Hispanic or Latino	63	61	42	41	39	40	32	34	37	40
COUNTY TYPE										
Large Metro	154	149	73	74	78	71	59	58	64	68
Small Metro	111	96	50	52	49	51	35	41	41	45
Nonmetro	67	66	35	33	33	34	26	25	27	24

*Low precision; no estimate reported.

NOTE: Response categories for the Perceived Availability questions include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy." Respondents with unknown Perceived Availability data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.5D Perceived That Obtaining Specific Illicit Drugs Would Be Fairly or Very Easy among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/ Geographic Characteristic	Marijuana (2013)	Marijuana (2014)	Cocaine (2013)	Cocaine (2014)	Crack (2013)	Crack (2014)	Heroin (2013)	Heroin (2014)	LSD (2013)	LSD (2014)
TOTAL	0.46	0.49	0.33	0.35	0.34	0.35	0.28	0.29	0.30	0.33
AGE										
12-13	0.68	0.69	0.46	0.50	0.49	0.57	0.38	0.45	0.37	0.40
14-15	0.75	0.84	0.57	0.60	0.63	0.59	0.48	0.49	0.48	0.53
16-17	0.74	0.77	0.65	0.71	0.58	0.66	0.48	0.56	0.61	0.68
GENDER										
Male	0.65	0.71	0.41	0.42	0.42	0.42	0.35	0.38	0.39	0.40
Female	0.65	0.69	0.51	0.55	0.52	0.56	0.42	0.45	0.47	0.51
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	0.52	0.55	0.37	0.39	0.39	0.39	0.32	0.33	0.34	0.37
White	0.59	0.65	0.42	0.48	0.45	0.46	0.39	0.39	0.40	0.45
Black or African American	1.15	1.31	0.89	0.95	0.90	1.08	0.69	0.86	0.73	0.82
American Indian or Alaska Native	4.33	5.19	2.55	2.27	3.53	2.77	3.20	1.70	2.69	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*	*	*
Asian	2.40	2.50	1.44	1.60	1.41	1.58	0.94	1.36	1.18	1.31
Two or More Races	2.57	2.44	1.84	1.92	1.82	2.10	1.54	1.45	1.75	1.56
Hispanic or Latino	1.17	1.13	0.80	0.77	0.74	0.74	0.62	0.63	0.72	0.76
COUNTY TYPE										
Large Metro	0.66	0.72	0.47	0.49	0.50	0.48	0.40	0.41	0.43	0.47
Small Metro	0.82	0.80	0.57	0.63	0.57	0.63	0.44	0.54	0.51	0.61
Nonmetro	1.06	1.11	0.83	0.82	0.77	0.82	0.67	0.65	0.70	0.65

*Low precision; no estimate reported.

NOTE: Response categories for the Perceived Availability questions include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy." Respondents with unknown Perceived Availability data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.5P Perceived That Obtaining Specific Illicit Drugs Would Be Fairly or Very Easy among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/ Geographic Characteristic	Marijuana (2013)	Marijuana (2014)	Cocaine (2013)	Cocaine (2014)	Crack (2013)	Crack (2014)	Heroin (2013)	Heroin (2014)	LSD (2013)	LSD (2014)
TOTAL	0.2041		0.8839		0.2148		0.4668		0.5077	
AGE										
12-13	0.0640		0.6465		0.6000		0.4139		0.1862	
14-15	0.2897		0.6850		0.1778		0.3183		0.9667	
16-17	0.9099		0.7008		0.7518		0.1479		0.1251	
GENDER										
Male	0.0721		0.8864		0.7026		0.4513		0.5112	
Female	0.9514		0.9421		0.1945		0.7612		0.7515	
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	0.8248		0.7241		0.6234		0.2110		0.3078	
White	0.4648		0.9132		0.4223		0.6937		0.1178	
Black or African American	0.6617		0.9518		0.9372		0.3599		0.4742	
American Indian or Alaska Native	0.4009		0.3501		0.1091		0.0467		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*		*	
Asian	0.8740		0.6853		0.6404		0.0415		0.6926	
Two or More Races	0.5424		0.3358		0.4622		0.6167		0.9671	
Hispanic or Latino	0.0459		0.3697		0.0907		0.4420		0.6762	
COUNTY TYPE										
Large Metro	0.0542		0.9591		0.2140		0.8081		0.9594	
Small Metro	0.6508		0.7312		0.8090		0.3819		0.2413	
Nonmetro	0.9515		0.8399		0.7595		0.7787		0.9117	

*Low precision; no estimate reported.

NOTE: Response categories for the Perceived Availability questions include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy." Respondents with unknown Perceived Availability data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.6C Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	55	60	68	83	64	77	66	70
AGE								
12-13	148	138	149	136	149	138	147	136
14-15	149	137	145	133	149	138	150	139
16-17	143	146	132	140	134	141	135	143
GENDER								
Male	38	40	51	54	47	48	50	51
Female	38	42	46	57	44	54	41	47
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	50	52	60	73	57	68	56	60
White	236	208	228	197	232	205	231	204
Black or African American	110	115	105	112	108	113	107	111
American Indian or Alaska Native	16	21	16	22	16	21	15	21
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	81	80	81	80	82	81	80	78
Two or More Races	39	42	38	42	38	42	39	42
Hispanic or Latino	27	28	34	39	32	33	32	35
COUNTY TYPE								
Large Metro	257	238	248	234	252	239	251	235
Small Metro	193	152	182	149	188	150	184	150
Nonmetro	115	116	115	112	116	114	116	114

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Perceptions of Parents' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove." Respondents with unknown Perceptions of Parents' Feelings data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.6D Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.23	0.25	0.28	0.34	0.26	0.31	0.27	0.28
AGE								
12-13	0.37	0.35	0.37	0.40	0.36	0.35	0.42	0.37
14-15	0.33	0.38	0.45	0.51	0.42	0.47	0.40	0.47
16-17	0.43	0.49	0.58	0.69	0.52	0.64	0.52	0.58
GENDER								
Male	0.31	0.32	0.41	0.44	0.38	0.38	0.40	0.41
Female	0.32	0.35	0.38	0.48	0.36	0.45	0.34	0.39
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.26	0.28	0.31	0.38	0.30	0.36	0.29	0.32
White	0.27	0.33	0.38	0.48	0.35	0.45	0.33	0.39
Black or African American	0.80	0.57	0.88	0.78	0.80	0.75	0.84	0.70
American Indian or Alaska Native	4.21	1.52	3.53	4.06	3.58	2.52	2.91	2.28
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.09	1.00	1.01	1.20	0.97	1.14	1.10	1.27
Two or More Races	1.06	1.10	1.68	1.51	1.34	1.43	1.28	1.25
Hispanic or Latino	0.49	0.50	0.63	0.70	0.58	0.61	0.60	0.63
COUNTY TYPE								
Large Metro	0.30	0.29	0.40	0.47	0.38	0.43	0.37	0.37
Small Metro	0.45	0.40	0.52	0.60	0.46	0.52	0.49	0.57
Nonmetro	0.52	0.79	0.67	0.88	0.57	0.79	0.62	0.74

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Perceptions of Parents' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove." Respondents with unknown Perceptions of Parents' Feelings data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.6P Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.4371		0.0461		0.0977		0.6800	
AGE								
12-13	0.0598		0.4709		0.0949		0.0430	
14-15	0.5343		0.0681		0.1044		0.1622	
16-17	0.5681		0.0888		0.0500		0.4549	
GENDER								
Male	0.0811		0.7744		0.6293		0.8974	
Female	0.5592		0.0013		0.0046		0.4398	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.3744		0.0959		0.2010		0.7907	
White	0.2203		0.0069		0.0266		0.1758	
Black or African American	0.0008		0.0789		0.1110		0.0678	
American Indian or Alaska Native	0.0117		0.9875		0.4844		0.5591	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.9522		0.5434		0.8511		0.6837	
Two or More Races	0.7288		0.3355		0.8170		0.7042	
Hispanic or Latino	0.9769		0.2460		0.2473		0.7194	
COUNTY TYPE								
Large Metro	0.2740		0.5477		0.7501		0.6288	
Small Metro	0.7062		0.0758		0.0131		0.5707	
Nonmetro	0.5199		0.1181		0.0661		0.1538	

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Perceptions of Parents' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove." Respondents with unknown Perceptions of Parents' Feelings data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.7C Youths Felt That Close Friends Would Strongly Disapprove or Somewhat Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	72	75	88	103	89	96	76	81
AGE								
12-13	143	138	144	138	143	137	143	140
14-15	146	138	138	127	140	129	142	135
16-17	135	142	119	119	119	118	132	132
GENDER								
Male	54	56	63	68	66	67	57	62
Female	48	45	60	68	60	63	50	50
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	61	63	78	85	79	79	66	67
White	224	207	211	185	213	187	222	203
Black or African American	104	110	100	101	99	100	101	106
American Indian or Alaska Native	16	21	14	19	14	19	14	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	81	79	78	78	78	78	77	80
Two or More Races	38	42	35	38	34	39	37	40
Hispanic or Latino	41	37	46	53	48	51	40	43
COUNTY TYPE								
Large Metro	244	238	227	217	229	219	239	233
Small Metro	185	149	166	133	163	132	181	142
Nonmetro	111	113	109	108	111	108	109	114

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Perceptions of Close Friends' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Perceptions of Close Friends' Feelings data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.7D Youths Felt That Close Friends Would Strongly Disapprove or Somewhat Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.30	0.31	0.36	0.42	0.36	0.39	0.31	0.33
AGE								
12-13	0.46	0.45	0.43	0.47	0.44	0.45	0.42	0.42
14-15	0.49	0.47	0.59	0.66	0.58	0.64	0.52	0.55
16-17	0.58	0.59	0.76	0.82	0.76	0.75	0.64	0.65
GENDER								
Male	0.44	0.45	0.51	0.55	0.53	0.54	0.46	0.50
Female	0.40	0.37	0.50	0.57	0.50	0.53	0.42	0.42
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.32	0.33	0.41	0.45	0.42	0.42	0.35	0.35
White	0.36	0.38	0.48	0.56	0.48	0.51	0.39	0.41
Black or African American	0.91	0.83	1.10	1.12	1.09	1.08	1.00	0.88
American Indian or Alaska Native	3.48	2.79	3.36	4.99	3.36	4.87	3.23	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.19	1.28	1.56	1.55	1.51	1.45	1.36	1.28
Two or More Races	1.48	1.18	1.93	1.91	1.96	1.86	1.62	1.46
Hispanic or Latino	0.76	0.68	0.85	0.96	0.89	0.93	0.74	0.78
COUNTY TYPE								
Large Metro	0.43	0.41	0.53	0.60	0.52	0.54	0.45	0.45
Small Metro	0.52	0.52	0.66	0.71	0.63	0.69	0.55	0.61
Nonmetro	0.68	0.69	0.80	0.86	0.78	0.85	0.71	0.77

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Perceptions of Close Friends' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Perceptions of Close Friends' Feelings data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.7P Youths Felt That Close Friends Would Strongly Disapprove or Somewhat Disapprove of Youth's Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.0881		0.0300		0.1715		0.0878	
AGE								
12-13	0.2389		0.7263		0.3569		0.1347	
14-15	0.1064		0.0923		0.1966		0.3062	
16-17	0.6357		0.0699		0.1909		0.3978	
GENDER								
Male	0.2231		0.6665		0.8855		0.1624	
Female	0.2051		0.0054		0.0573		0.3115	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.0778		0.0603		0.1620		0.1190	
White	0.2389		0.0774		0.0793		0.7230	
Black or African American	0.4393		0.4454		0.8588		0.0512	
American Indian or Alaska Native	0.2238		0.7537		0.5295		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.9905		0.4963		0.5073		0.1818	
Two or More Races	0.0675		0.3616		0.6986		0.5677	
Hispanic or Latino	0.6394		0.2749		0.7654		0.4364	
COUNTY TYPE								
Large Metro	0.0311		0.3190		0.6050		0.0192	
Small Metro	0.7597		0.0718		0.2505		0.3971	
Nonmetro	0.7307		0.1925		0.2670		0.4427	

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Perceptions of Close Friends' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Perceptions of Close Friends' Feelings data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.8C Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	65	61	92	102	92	103	74	76
AGE								
12-13	147	139	145	137	147	137	148	139
14-15	146	137	137	126	138	126	144	134
16-17	139	147	121	121	118	117	134	140
GENDER								
Male	48	45	66	71	68	72	56	52
Female	44	43	61	68	61	68	47	51
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	57	53	79	85	81	85	64	63
White	235	210	210	189	213	188	226	206
Black or African American	106	113	100	105	99	104	104	111
American Indian or Alaska Native	15	*	15	*	16	*	17	*
Native Hawaiian or Other Pacific Islander	16	*	*	*	*	*	*	*
Asian	80	80	77	76	78	75	79	79
Two or More Races	39	42	35	39	35	39	38	41
Hispanic or Latino	35	33	50	52	48	54	40	40
COUNTY TYPE								
Large Metro	247	237	225	213	227	212	240	233
Small Metro	189	147	169	138	167	136	184	149
Nonmetro	112	119	108	112	107	112	112	116

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Feelings about Peers include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Feelings about Peers data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.8D Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.27	0.25	0.38	0.41	0.37	0.42	0.30	0.31
AGE								
12-13	0.42	0.39	0.44	0.47	0.42	0.47	0.40	0.40
14-15	0.43	0.43	0.63	0.64	0.61	0.65	0.50	0.50
16-17	0.50	0.48	0.77	0.79	0.75	0.78	0.61	0.59
GENDER								
Male	0.38	0.36	0.53	0.57	0.54	0.58	0.45	0.42
Female	0.36	0.36	0.51	0.56	0.51	0.56	0.39	0.42
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.30	0.28	0.42	0.45	0.42	0.45	0.34	0.33
White	0.33	0.32	0.49	0.56	0.51	0.55	0.38	0.40
Black or African American	0.89	0.73	1.09	1.04	1.08	1.02	0.96	0.80
American Indian or Alaska Native	3.19	*	4.09	*	4.14	*	2.99	*
Native Hawaiian or Other Pacific Islander	1.74	*	*	*	*	*	*	*
Asian	1.21	1.19	1.47	1.65	1.64	1.59	1.25	1.08
Two or More Races	1.23	1.15	2.05	1.94	2.17	1.98	1.32	1.19
Hispanic or Latino	0.65	0.59	0.92	0.94	0.88	0.97	0.73	0.73
COUNTY TYPE								
Large Metro	0.37	0.34	0.54	0.58	0.54	0.60	0.43	0.42
Small Metro	0.46	0.50	0.67	0.69	0.65	0.71	0.49	0.56
Nonmetro	0.72	0.60	0.83	1.00	0.80	0.92	0.76	0.73

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Feelings about Peers include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Feelings about Peers data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.8P Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	SUBSTANCE USE BEHAVIOR							
	Smoking One or More Packs of Cigarettes Per Day		Trying Marijuana or Hashish Once or Twice		Using Marijuana or Hashish Once a Month or More		Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.0021		0.9704		0.9141		0.0275	
AGE								
12-13	0.0142		0.0216		0.1199		0.0214	
14-15	0.4547		0.7596		0.8386		0.6472	
16-17	0.0212		0.4978		0.5579		0.1140	
GENDER								
Male	0.0010		0.0846		0.1397		0.0013	
Female	0.3268		0.0768		0.0779		0.8357	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.0390		0.8388		0.6233		0.0987	
White	0.2307		0.5881		0.2860		0.9020	
Black or African American	0.0333		0.3877		0.2795		0.0110	
American Indian or Alaska Native	*		*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.7909		0.7952		0.9779		0.2669	
Two or More Races	0.9238		0.8873		0.8835		0.1781	
Hispanic or Latino	0.0150		0.6900		0.5419		0.1233	
COUNTY TYPE								
Large Metro	0.0008		0.9699		0.9128		0.0072	
Small Metro	0.2888		0.6760		0.7275		0.8214	
Nonmetro	0.0081		0.3636		0.5584		0.5483	

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Youths' Feelings about Peers include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Feelings about Peers data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.9C Participated in a Prevention Program or Group in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	PREVENTION PROGRAM OR GROUP					
	Problem-Solving, Communication Skills, or Self-Esteem Group		Violence Prevention Program		Pregnancy or Sexually Transmitted Disease Prevention Program	
	2013	2014	2013	2014	2013	2014
TOTAL	92	101	79	79	67	67
AGE						
12-13	61	65	57	53	39	34
14-15	66	70	48	48	46	49
16-17	60	63	35	43	38	40
GENDER						
Male	64	71	52	54	48	46
Female	65	67	54	55	45	47
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	77	86	65	69	55	59
White	81	84	55	56	45	48
Black or African American	51	54	44	47	31	36
American Indian or Alaska Native	8	7	6	7	6	8
Native Hawaiian or Other Pacific Islander	*	*	4	*	*	*
Asian	38	38	19	23	22	16
Two or More Races	24	24	21	11	13	12
Hispanic or Latino	47	50	41	38	38	35
COUNTY TYPE						
Large Metro	91	101	70	67	54	57
Small Metro	68	61	43	47	44	40
Nonmetro	43	45	34	34	30	26

*Low precision; no estimate reported.

NOTE: Respondents with unknown Prevention Program or Group Participation data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.9D Participated in a Prevention Program or Group in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	PREVENTION PROGRAM OR GROUP					
	Problem-Solving, Communication Skills, or Self-Esteem Group		Violence Prevention Program		Pregnancy or Sexually Transmitted Disease Prevention Program	
	2013	2014	2013	2014	2013	2014
TOTAL	0.38	0.42	0.32	0.32	0.27	0.27
AGE						
12-13	0.69	0.74	0.65	0.64	0.47	0.42
14-15	0.65	0.71	0.53	0.54	0.52	0.55
16-17	0.63	0.66	0.40	0.48	0.43	0.44
GENDER						
Male	0.52	0.57	0.42	0.43	0.38	0.37
Female	0.55	0.57	0.45	0.46	0.38	0.39
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.41	0.46	0.34	0.37	0.29	0.31
White	0.46	0.52	0.35	0.39	0.30	0.34
Black or African American	1.03	1.19	1.03	1.07	0.79	0.92
American Indian or Alaska Native	3.67	3.84	2.95	3.87	3.33	4.14
Native Hawaiian or Other Pacific Islander	*	*	3.71	*	*	*
Asian	2.13	2.35	1.43	1.72	1.65	1.18
Two or More Races	2.51	2.36	2.39	1.34	1.61	1.51
Hispanic or Latino	0.87	0.90	0.75	0.68	0.70	0.62
COUNTY TYPE						
Large Metro	0.54	0.59	0.46	0.45	0.37	0.38
Small Metro	0.70	0.67	0.53	0.59	0.52	0.52
Nonmetro	0.84	0.99	0.76	0.79	0.70	0.61

*Low precision; no estimate reported.

NOTE: Respondents with unknown Prevention Program or Group Participation data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.9P Participated in a Prevention Program or Group in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	PREVENTION PROGRAM OR GROUP					
	Problem-Solving, Communication Skills, or Self-Esteem Group		Violence Prevention Program		Pregnancy or Sexually Transmitted Disease Prevention Program	
	2013	2014	2013	2014	2013	2014
TOTAL	0.8533		0.2495		0.5178	
AGE						
12-13	0.3915		0.1497		0.0239	
14-15	0.7133		0.0528		0.3397	
16-17	0.4578		0.0521		0.8358	
GENDER						
Male	0.7797		0.8151		0.9291	
Female	0.5724		0.1489		0.3944	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4340		0.3300		0.9224	
White	0.4332		0.2459		0.4432	
Black or African American	0.9757		0.6251		0.5703	
American Indian or Alaska Native	0.2612		0.6086		0.9348	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.9173		0.7961		0.4955	
Two or More Races	0.5549		0.0028		0.0425	
Hispanic or Latino	0.3196		0.5152		0.2913	
COUNTY TYPE						
Large Metro	0.8150		0.2600		0.2354	
Small Metro	0.8335		0.5139		0.1207	
Nonmetro	0.7784		0.8368		0.0768	

*Low precision; no estimate reported.

NOTE: Respondents with unknown Prevention Program or Group Participation data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.10C Participated in a Substance Use Program or Exposure to Substance Use Prevention Message in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE PROGRAM OR PREVENTION MESSAGE							
	Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School		Participated in Program to Help Deal with Drug or Alcohol Use		Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School		Talked to Parents about Dangers of Tobacco, Alcohol, or Drug Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	71	74	40	48	104	109	118	121
AGE								
12-13	47	47	26	29	119	114	104	98
14-15	48	46	24	26	122	127	106	108
16-17	38	43	20	27	123	130	101	116
GENDER								
Male	52	56	29	36	75	77	81	88
Female	48	49	28	31	66	75	80	79
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	60	64	31	40	88	92	101	103
White	54	54	26	33	187	171	165	156
Black or African American	35	37	19	23	91	97	70	75
American Indian or Alaska Native	7	7	5	3	13	20	10	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	26	19	8	6	67	68	43	53
Two or More Races	13	8	7	5	32	36	31	31
Hispanic or Latino	38	38	25	25	53	57	60	62
COUNTY TYPE								
Large Metro	63	63	32	39	208	207	170	177
Small Metro	44	41	22	26	156	131	123	106
Nonmetro	28	31	14	17	97	94	78	73

*Low precision; no estimate reported.

NOTE: Respondents with unknown Substance Use Program Participation or Prevention Message Exposure data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.10D Participated in a Substance Use Program or Exposure to Substance Use Prevention Message in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE PROGRAM OR PREVENTION MESSAGE							
	Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School		Participated in Program to Help Deal with Drug or Alcohol Use		Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School		Talked to Parents about Dangers of Tobacco, Alcohol, or Drug Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.29	0.30	0.16	0.19	0.42	0.44	0.48	0.49
AGE								
12-13	0.54	0.57	0.32	0.37	0.78	0.80	0.86	0.88
14-15	0.54	0.51	0.28	0.30	0.67	0.73	0.76	0.83
16-17	0.44	0.49	0.24	0.30	0.69	0.71	0.81	0.85
GENDER								
Male	0.41	0.45	0.23	0.29	0.60	0.61	0.65	0.70
Female	0.39	0.40	0.23	0.26	0.55	0.62	0.67	0.66
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.32	0.33	0.16	0.21	0.46	0.48	0.53	0.54
White	0.34	0.38	0.18	0.24	0.52	0.56	0.60	0.63
Black or African American	0.83	0.93	0.52	0.64	1.10	1.13	1.26	1.32
American Indian or Alaska Native	3.46	3.42	2.54	1.93	5.08	3.68	3.57	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.85	1.38	0.69	0.49	2.07	2.08	2.26	2.48
Two or More Races	1.55	0.99	0.96	0.65	2.46	2.25	2.25	2.42
Hispanic or Latino	0.69	0.69	0.46	0.45	0.97	1.02	1.10	1.12
COUNTY TYPE								
Large Metro	0.41	0.41	0.23	0.28	0.61	0.63	0.70	0.72
Small Metro	0.52	0.54	0.29	0.36	0.70	0.78	0.76	0.83
Nonmetro	0.65	0.72	0.36	0.43	1.01	1.11	1.07	1.06

*Low precision; no estimate reported.

NOTE: Respondents with unknown Substance Use Program Participation or Prevention Message Exposure data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.10P Participated in a Substance Use Program or Exposure to Substance Use Prevention Message in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	SUBSTANCE USE PROGRAM OR PREVENTION MESSAGE							
	Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School		Participated in Program to Help Deal with Drug or Alcohol Use		Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School		Talked to Parents about Dangers of Tobacco, Alcohol, or Drug Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.2917		0.0801		0.6126		0.3596	
AGE								
12-13	0.1259		0.7716		0.7878		0.3177	
14-15	0.3290		0.2616		0.0612		0.4649	
16-17	0.3608		0.0611		0.4516		0.0670	
GENDER								
Male	0.9008		0.1550		0.5775		0.1341	
Female	0.1496		0.2969		0.8677		0.8263	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.1825		0.0916		0.7427		0.1529	
White	0.7857		0.0438		0.8687		0.1643	
Black or African American	0.3696		0.3288		0.4366		0.4708	
American Indian or Alaska Native	0.8723		0.2596		0.0999		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.3913		0.3986		0.5146		0.2628	
Two or More Races	0.0004		0.0892		0.8934		0.2812	
Hispanic or Latino	0.8580		0.5033		0.6578		0.5550	
COUNTY TYPE								
Large Metro	0.4094		0.0683		0.9016		0.2515	
Small Metro	0.1410		0.8869		0.6180		0.8785	
Nonmetro	0.3061		0.4720		0.8139		0.8985	

*Low precision; no estimate reported.

NOTE: Respondents with unknown Substance Use Program Participation or Prevention Message Exposure data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.11C Participated in a Physical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	DELINQUENT BEHAVIOR					
	Got into a Serious Fight at School or Work		Took Part in a Group-Against-Group Fight		Attacked Someone with the Intent to Seriously Hurt Them	
	2013	2014	2013	2014	2013	2014
TOTAL	93	92	72	75	52	49
AGE						
12-13	61	59	47	43	27	29
14-15	58	59	46	44	30	28
16-17	51	54	39	44	31	30
GENDER						
Male	68	69	53	56	40	38
Female	58	61	45	49	30	33
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	75	81	59	63	44	45
White	65	70	50	50	30	34
Black or African American	49	48	31	35	26	27
American Indian or Alaska Native	9	6	9	6	4	3
Native Hawaiian or Other Pacific Islander	*	*	*	*	1	*
Asian	16	19	16	14	11	13
Two or More Races	22	17	18	10	12	14
Hispanic or Latino	50	47	40	41	25	26
COUNTY TYPE						
Large Metro	86	80	64	60	41	40
Small Metro	56	55	43	41	30	30
Nonmetro	39	39	29	30	19	21

*Low precision; no estimate reported.

NOTE: Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time. Respondents with unknown Delinquent Behavior data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.11D Participated in a Physical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	DELINQUENT BEHAVIOR					
	Got into a Serious Fight at School or Work		Took Part in a Group-Against-Group Fight		Attacked Someone with the Intent to Seriously Hurt Them	
	2013	2014	2013	2014	2013	2014
TOTAL	0.37	0.37	0.29	0.30	0.21	0.20
AGE						
12-13	0.65	0.65	0.53	0.53	0.33	0.36
14-15	0.61	0.67	0.51	0.51	0.34	0.32
16-17	0.57	0.58	0.45	0.49	0.36	0.34
GENDER						
Male	0.54	0.55	0.42	0.45	0.32	0.30
Female	0.48	0.50	0.37	0.41	0.25	0.27
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.39	0.42	0.31	0.33	0.23	0.23
White	0.41	0.49	0.33	0.36	0.22	0.24
Black or African American	1.08	1.05	0.80	0.85	0.69	0.68
American Indian or Alaska Native	4.04	3.67	4.62	3.63	2.21	1.69
Native Hawaiian or Other Pacific Islander	*	*	*	*	0.55	*
Asian	1.28	1.38	1.25	1.07	0.84	1.03
Two or More Races	2.43	1.92	2.16	1.27	1.51	1.73
Hispanic or Latino	0.91	0.84	0.72	0.73	0.45	0.47
COUNTY TYPE						
Large Metro	0.53	0.50	0.42	0.41	0.29	0.27
Small Metro	0.67	0.66	0.49	0.54	0.39	0.40
Nonmetro	0.84	0.91	0.67	0.70	0.48	0.50

*Low precision; no estimate reported.

NOTE: Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time. Respondents with unknown Delinquent Behavior data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.11P Participated in a Physical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	DELINQUENT BEHAVIOR					
	Got into a Serious Fight at School or Work		Took Part in a Group-Against-Group Fight		Attacked Someone with the Intent to Seriously Hurt Them	
	2013	2014	2013	2014	2013	2014
TOTAL	0.1075		0.3601		0.3699	
AGE						
12-13	0.3376		0.3921		0.4100	
14-15	0.4824		0.3711		0.6594	
16-17	0.2356		0.8192		0.0613	
GENDER						
Male	0.0480		0.3265		0.2036	
Female	0.8431		0.7810		0.8837	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2874		0.3922		0.3606	
White	0.5715		0.9500		0.6239	
Black or African American	0.2929		0.4689		0.1347	
American Indian or Alaska Native	0.2166		0.3427		0.6215	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4326		0.5283		0.3338	
Two or More Races	0.2529		0.0439		0.3571	
Hispanic or Latino	0.1721		0.6944		0.8526	
COUNTY TYPE						
Large Metro	0.0966		0.5873		0.9475	
Small Metro	0.7960		0.2789		0.1703	
Nonmetro	0.5847		0.8906		0.8860	

*Low precision; no estimate reported.

NOTE: Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time. Respondents with unknown Delinquent Behavior data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.12C Participated in a Nonphysical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	DELINQUENT BEHAVIOR					
	Carried a Handgun		Sold Illegal Drugs		Stole or Tried to Steal Anything Worth More Than \$50	
	2013	2014	2013	2014	2013	2014
TOTAL	40	44	35	33	36	38
AGE						
12-13	20	25	9	8	15	16
14-15	25	26	19	17	24	22
16-17	26	26	26	27	26	27
GENDER						
Male	36	39	28	27	30	31
Female	17	21	18	19	21	20
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	34	40	27	28	30	33
White	29	36	24	24	25	27
Black or African American	14	13	12	14	15	16
American Indian or Alaska Native	4	5	4	1	6	2
Native Hawaiian or Other Pacific Islander	*	*	2	*	2	*
Asian	7	6	5	3	10	7
Two or More Races	11	7	4	6	6	5
Hispanic or Latino	21	20	20	17	21	20
COUNTY TYPE						
Large Metro	29	31	26	28	30	31
Small Metro	23	24	20	15	18	21
Nonmetro	19	21	10	11	11	10

*Low precision; no estimate reported.

NOTE: Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time. Respondents with unknown Delinquent Behavior data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.12D Participated in a Nonphysical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	DELINQUENT BEHAVIOR					
	Carried a Handgun		Sold Illegal Drugs		Stole or Tried to Steal Anything Worth More Than \$50	
	2013	2014	2013	2014	2013	2014
TOTAL	0.16	0.18	0.14	0.13	0.15	0.15
AGE						
12-13	0.25	0.32	0.11	0.11	0.19	0.20
14-15	0.29	0.30	0.22	0.20	0.28	0.26
16-17	0.30	0.30	0.31	0.31	0.31	0.31
GENDER						
Male	0.29	0.31	0.23	0.21	0.24	0.25
Female	0.14	0.17	0.15	0.16	0.17	0.16
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.18	0.21	0.14	0.15	0.16	0.17
White	0.21	0.26	0.17	0.18	0.18	0.20
Black or African American	0.40	0.37	0.34	0.40	0.42	0.44
American Indian or Alaska Native	2.06	2.85	2.44	0.79	3.10	1.33
Native Hawaiian or Other Pacific Islander	*	*	2.28	*	1.77	*
Asian	0.53	0.49	0.39	0.26	0.79	0.60
Two or More Races	1.36	0.88	0.49	0.74	0.73	0.59
Hispanic or Latino	0.38	0.36	0.35	0.30	0.37	0.35
COUNTY TYPE						
Large Metro	0.21	0.22	0.19	0.20	0.22	0.22
Small Metro	0.29	0.33	0.27	0.21	0.25	0.29
Nonmetro	0.45	0.52	0.26	0.30	0.29	0.26

*Low precision; no estimate reported.

NOTE: Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time. Respondents with unknown Delinquent Behavior data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.12P Participated in a Nonphysical Delinquent Behavior in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	DELINQUENT BEHAVIOR					
	Carried a Handgun		Sold Illegal Drugs		Stole or Tried to Steal Anything Worth More Than \$50	
	2013	2014	2013	2014	2013	2014
TOTAL	0.1142		0.0543		0.3631	
AGE						
12-13	0.0328		0.5626		0.9512	
14-15	0.6722		0.0994		0.6636	
16-17	0.7408		0.2089		0.3519	
GENDER						
Male	0.2517		0.0233		0.7293	
Female	0.1990		0.8853		0.2574	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1189		0.2379		0.9221	
White	0.0642		0.1367		0.6176	
Black or African American	0.8363		0.7416		0.4303	
American Indian or Alaska Native	0.4900		0.1782		0.1203	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.9781		0.2272		0.7243	
Two or More Races	0.7181		0.3031		0.1950	
Hispanic or Latino	0.6466		0.0720		0.1256	
COUNTY TYPE						
Large Metro	0.1534		0.2753		0.2635	
Small Metro	0.1689		0.0191		0.7082	
Nonmetro	0.9563		0.6032		0.4439	

*Low precision; no estimate reported.

NOTE: Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time. Respondents with unknown Delinquent Behavior data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.13C Selected Risk and Protective Factors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	RISK FACTORS						PROTECTIVE FACTORS			
	Youth Moved 2 or More Times in the Past 5 Years		Youth Argued with at Least One Parent 10 or More Times in the Past Year		Youth Was Approached by Someone Selling Drugs in the Past Month		Youth Had Someone to Talk to about Problems ¹		Youth Attended 25 or More Religious Services in the Past Year ²	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	115	116	94	106	79	87	46	53	113	126
AGE										
12-13	66	72	49	52	29	26	149	136	72	76
14-15	73	68	69	66	47	53	147	140	76	74
16-17	67	68	61	75	60	66	142	150	78	80
GENDER										
Male	77	77	58	62	57	66	34	34	75	83
Female	74	74	71	75	52	49	29	37	73	83
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	93	97	81	90	66	72	40	41	100	114
White	83	85	86	92	57	65	236	212	119	121
Black or African American	60	62	30	32	34	36	111	116	54	55
American Indian or Alaska Native	9	10	7	*	4	*	18	21	6	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*	*	*
Asian	22	25	31	31	17	15	75	77	37	37
Two or More Races	23	26	19	17	16	13	40	41	21	19
Hispanic or Latino	62	58	45	51	42	48	22	30	52	51
COUNTY TYPE										
Large Metro	106	105	92	103	69	74	256	240	120	127
Small Metro	84	75	60	57	46	49	190	158	89	69
Nonmetro	46	49	38	42	25	23	121	116	61	59

*Low precision; no estimate reported.

NOTE: Respondents with unknown Risk or Protective Factors data were excluded from the respective analyses.

¹ Respondents were able to select multiple categories of persons that they would talk to about a serious problem, including the choice of "nobody." Respondents who reported both a person that they would talk to and that they had nobody to talk to were excluded from the analysis.

² Special occasions, such as weddings, funerals, or other special events, were not to have been considered in the number of times attended religious services.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.13D Selected Risk and Protective Factors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	RISK FACTORS						PROTECTIVE FACTORS			
	Youth Moved 2 or More Times in the Past 5 Years		Youth Argued with at Least One Parent 10 or More Times in the Past Year		Youth Was Approached by Someone Selling Drugs in the Past Month		Youth Had Someone to Talk to about Problems ¹		Youth Attended 25 or More Religious Services in the Past Year ²	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.48	0.48	0.38	0.43	0.32	0.35	0.19	0.22	0.47	0.52
AGE										
12-13	0.73	0.79	0.55	0.61	0.36	0.33	0.34	0.38	0.76	0.85
14-15	0.73	0.72	0.68	0.71	0.52	0.58	0.30	0.35	0.72	0.77
16-17	0.71	0.72	0.65	0.76	0.63	0.69	0.33	0.38	0.75	0.79
GENDER										
Male	0.62	0.62	0.46	0.50	0.45	0.52	0.28	0.28	0.61	0.67
Female	0.62	0.62	0.59	0.63	0.43	0.40	0.25	0.31	0.62	0.71
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	0.49	0.52	0.42	0.48	0.34	0.37	0.21	0.22	0.53	0.61
White	0.52	0.57	0.48	0.58	0.37	0.46	0.22	0.24	0.62	0.74
Black or African American	1.28	1.35	0.80	0.86	0.85	0.89	0.54	0.56	1.27	1.27
American Indian or Alaska Native	4.72	4.87	4.09	*	2.45	*	2.29	1.35	3.39	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*	*	*
Asian	1.64	1.87	2.20	2.13	1.36	1.11	1.43	1.38	2.45	2.54
Two or More Races	2.57	2.59	2.16	1.92	1.83	1.64	0.91	0.93	2.45	2.30
Hispanic or Latino	1.17	1.08	0.83	0.92	0.76	0.86	0.41	0.55	0.98	0.94
COUNTY TYPE										
Large Metro	0.66	0.66	0.55	0.63	0.46	0.49	0.28	0.33	0.65	0.71
Small Metro	0.90	0.87	0.61	0.66	0.54	0.61	0.28	0.32	0.83	0.80
Nonmetro	0.98	1.02	0.86	0.97	0.61	0.56	0.41	0.45	1.10	1.27

*Low precision; no estimate reported.

NOTE: Respondents with unknown Risk or Protective Factors data were excluded from the respective analyses.

¹ Respondents were able to select multiple categories of persons that they would talk to about a serious problem, including the choice of "nobody." Respondents who reported both a person that they would talk to and that they had nobody to talk to were excluded from the analysis.

² Special occasions, such as weddings, funerals, or other special events, were not to have been considered in the number of times attended religious services.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.13P Selected Risk and Protective Factors among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	RISK FACTORS						PROTECTIVE FACTORS			
	Youth Moved 2 or More Times in the Past 5 Years		Youth Argued with at Least One Parent 10 or More Times in the Past Year		Youth Was Approached by Someone Selling Drugs in the Past Month		Youth Had Someone to Talk to about Problems ¹		Youth Attended 25 or More Religious Services in the Past Year ²	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.9379		0.9184		0.5288		0.5417		0.2456	
AGE										
12-13	0.7752		0.4525		0.3034		0.9563		0.4433	
14-15	0.6800		0.1047		0.7894		0.4583		0.0241	
16-17	0.7475		0.4256		0.4098		0.6878		0.4758	
GENDER										
Male	0.5371		0.5268		0.4548		0.4779		0.1472	
Female	0.4437		0.7250		0.9121		0.1010		0.7624	
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	0.4581		0.9885		0.3819		0.3303		0.5738	
White	0.3801		0.8650		0.6975		0.2665		0.4433	
Black or African American	0.5089		0.9038		0.3832		0.6411		0.4263	
American Indian or Alaska Native	0.8283		*		*		0.3178		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*		*	
Asian	0.8386		0.9009		0.4897		0.7110		0.6806	
Two or More Races	0.3115		0.3294		0.2320		0.7998		0.3454	
Hispanic or Latino	0.2915		0.8721		0.8908		0.0093		0.1512	
COUNTY TYPE										
Large Metro	0.8971		0.5848		0.3558		0.3336		0.5883	
Small Metro	0.8773		0.6571		0.7998		0.8117		0.1405	
Nonmetro	0.6505		0.8144		0.7219		0.6476		0.0345	

*Low precision; no estimate reported.

NOTE: Respondents with unknown Risk or Protective Factors data were excluded from the respective analyses.

¹ Respondents were able to select multiple categories of persons that they would talk to about a serious problem, including the choice of "nobody." Respondents who reported both a person that they would talk to and that they had nobody to talk to were excluded from the analysis.

² Special occasions, such as weddings, funerals, or other special events, were not to have been considered in the number of times attended religious services.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.14C Youths Agreed or Strongly Agreed with Statements about Religious Beliefs among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	STATEMENTS ABOUT RELIGIOUS BELIEFS					
	Religious Beliefs Are a Very Important Part of Youth's Life		Religious Beliefs Influence Youth's Decision Making in Life		It Is Important for Friends to Share Religious Beliefs	
	2013	2014	2013	2014	2013	2014
TOTAL	103	113	116	120	117	113
AGE						
12-13	135	124	125	117	82	78
14-15	128	115	117	112	77	73
16-17	115	119	112	113	77	74
GENDER						
Male	71	79	79	81	78	80
Female	71	76	77	83	74	78
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	89	99	98	105	101	96
White	193	170	178	164	115	101
Black or African American	102	106	100	97	56	61
American Indian or Alaska Native	16	21	15	18	13	13
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	63	64	61	59	31	32
Two or More Races	32	36	30	31	18	17
Hispanic or Latino	51	57	58	60	57	58
COUNTY TYPE						
Large Metro	209	203	195	191	115	111
Small Metro	162	126	149	115	93	74
Nonmetro	104	100	98	94	67	66

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Religious Beliefs include "Strongly disagree," "Disagree," "Agree," and "Strongly agree." The estimates in this table correspond to persons reporting "Agree" or "Strongly agree." Respondents with unknown Religious Beliefs data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.14D Youths Agreed or Strongly Agreed with Statements about Religious Beliefs among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	STATEMENTS ABOUT RELIGIOUS BELIEFS					
	Religious Beliefs Are a Very Important Part of Youth's Life		Religious Beliefs Influence Youth's Decision Making in Life		It Is Important for Friends to Share Religious Beliefs	
	2013	2014	2013	2014	2013	2014
TOTAL	0.43	0.47	0.48	0.50	0.49	0.47
AGE						
12-13	0.65	0.72	0.73	0.81	0.83	0.81
14-15	0.70	0.74	0.75	0.80	0.75	0.76
16-17	0.74	0.79	0.80	0.82	0.76	0.74
GENDER						
Male	0.58	0.65	0.64	0.66	0.64	0.66
Female	0.60	0.64	0.65	0.70	0.63	0.67
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.47	0.53	0.52	0.56	0.54	0.52
White	0.56	0.63	0.57	0.68	0.62	0.61
Black or African American	0.95	0.97	1.14	1.10	1.22	1.31
American Indian or Alaska Native	3.38	3.80	4.01	4.16	5.77	5.34
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.36	2.27	2.67	2.37	2.36	2.29
Two or More Races	2.65	2.36	2.92	2.43	2.16	2.16
Hispanic or Latino	0.95	1.05	1.08	1.10	1.08	1.08
COUNTY TYPE						
Large Metro	0.62	0.68	0.72	0.71	0.68	0.63
Small Metro	0.71	0.77	0.76	0.83	0.85	0.85
Nonmetro	0.93	1.01	0.94	1.14	1.13	1.20

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Religious Beliefs include "Strongly disagree," "Disagree," "Agree," and "Strongly agree." The estimates in this table correspond to persons reporting "Agree" or "Strongly agree." Respondents with unknown Religious Beliefs data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.14P Youths Agreed or Strongly Agreed with Statements about Religious Beliefs among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	STATEMENTS ABOUT RELIGIOUS BELIEFS					
	Religious Beliefs Are a Very Important Part of Youth's Life		Religious Beliefs Influence Youth's Decision Making in Life		It Is Important for Friends to Share Religious Beliefs	
	2013	2014	2013	2014	2013	2014
TOTAL	0.0202		0.0075		0.0171	
AGE						
12-13	0.2122		0.0907		0.1994	
14-15	0.0305		0.0342		0.1908	
16-17	0.4477		0.2771		0.0830	
GENDER						
Male	0.0772		0.1882		0.0491	
Female	0.1129		0.0083		0.1205	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0667		0.0553		0.0103	
White	0.0493		0.0328		0.0398	
Black or African American	0.4042		0.6325		0.2410	
American Indian or Alaska Native	0.8484		0.2643		0.7957	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1597		0.6812		0.2025	
Two or More Races	0.5885		0.9596		0.4428	
Hispanic or Latino	0.1299		0.0383		0.7076	
COUNTY TYPE						
Large Metro	0.1369		0.1313		0.0900	
Small Metro	0.1457		0.0255		0.1958	
Nonmetro	0.3307		0.3913		0.5690	

*Low precision; no estimate reported.

NOTE: Response categories for the questions on Religious Beliefs include "Strongly disagree," "Disagree," "Agree," and "Strongly agree." The estimates in this table correspond to persons reporting "Agree" or "Strongly agree." Respondents with unknown Religious Beliefs data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.15C Parents Always or Sometimes Took Part in Specific Behaviors in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	PARENTAL INVOLVEMENT							
	Parents Made Youth Do Chores around the House		Parents Limited Amount of Time Youth Watched TV		Parents Let Youth Know He or She Had Done a Good Job		Parents Let Youth Know They Were Proud of Something He or She Had Done	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	73	77	112	123	80	83	81	92
AGE								
12-13	143	139	100	103	140	137	143	139
14-15	147	139	89	91	139	133	137	132
16-17	133	147	77	81	131	140	131	138
GENDER								
Male	52	53	76	82	53	55	54	57
Female	48	54	78	88	59	61	59	65
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	61	65	99	108	63	69	65	72
White	223	203	135	128	223	204	227	202
Black or African American	111	117	63	65	106	109	108	109
American Indian or Alaska Native	18	22	8	13	15	*	14	20
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	73	76	52	56	68	75	67	75
Two or More Races	39	41	27	23	38	38	37	38
Hispanic or Latino	36	40	57	63	46	45	49	49
COUNTY TYPE								
Large Metro	243	233	140	148	231	229	234	233
Small Metro	183	151	100	90	183	143	181	141
Nonmetro	116	116	60	64	115	114	116	112

*Low precision; no estimate reported.

NOTE: Response categories for the Parental Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes." Respondents with unknown Parental Involvement data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.15D Parents Always or Sometimes Took Part in Specific Behaviors in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	PARENTAL INVOLVEMENT							
	Parents Made Youth Do Chores around the House		Parents Limited Amount of Time Youth Watched TV		Parents Let Youth Know He or She Had Done a Good Job		Parents Let Youth Know They Were Proud of Something He or She Had Done	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.29	0.31	0.46	0.50	0.32	0.34	0.33	0.37
AGE								
12-13	0.48	0.52	0.77	0.89	0.46	0.47	0.46	0.48
14-15	0.49	0.51	0.76	0.84	0.53	0.56	0.55	0.63
16-17	0.51	0.53	0.71	0.76	0.62	0.66	0.62	0.68
GENDER								
Male	0.42	0.42	0.61	0.65	0.42	0.44	0.42	0.46
Female	0.40	0.44	0.65	0.73	0.48	0.51	0.49	0.53
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.32	0.34	0.52	0.57	0.33	0.36	0.34	0.37
White	0.38	0.39	0.60	0.68	0.36	0.40	0.37	0.43
Black or African American	0.70	0.70	1.24	1.27	0.89	0.95	0.83	0.92
American Indian or Alaska Native	2.35	2.43	3.51	5.18	2.81	*	3.66	4.82
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.69	1.87	2.46	2.77	1.88	1.63	2.02	1.96
Two or More Races	1.45	1.52	2.65	2.29	1.57	2.01	1.78	2.03
Hispanic or Latino	0.64	0.71	1.04	1.13	0.84	0.81	0.89	0.87
COUNTY TYPE								
Large Metro	0.42	0.43	0.65	0.71	0.46	0.50	0.48	0.55
Small Metro	0.48	0.56	0.84	0.89	0.55	0.60	0.53	0.59
Nonmetro	0.66	0.59	1.04	1.18	0.76	0.69	0.73	0.83

*Low precision; no estimate reported.

NOTE: Response categories for the Parental Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes." Respondents with unknown Parental Involvement data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.15P Parents Always or Sometimes Took Part in Specific Behaviors in the Past Year among Persons Aged 12 to 17, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	PARENTAL INVOLVEMENT							
	Parents Made Youth Do Chores around the House		Parents Limited Amount of Time Youth Watched TV		Parents Let Youth Know He or She Had Done a Good Job		Parents Let Youth Know They Were Proud of Something He or She Had Done	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.0181		0.6737		0.1208		0.5051	
AGE								
12-13	0.6195		0.8361		0.9179		0.3418	
14-15	0.0262		0.8032		0.3047		0.9932	
16-17	0.1431		0.5989		0.1033		0.6134	
GENDER								
Male	0.0004		0.2883		0.0205		0.1277	
Female	0.8590		0.6777		0.9752		0.6738	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.0160		0.8131		0.1216		0.8160	
White	0.0082		0.6708		0.3171		0.9650	
Black or African American	0.4557		0.9620		0.1797		0.7599	
American Indian or Alaska Native	0.5305		0.4912		*		0.8221	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.7590		0.2200		0.1321		0.6699	
Two or More Races	0.5814		0.0555		0.0953		0.9179	
Hispanic or Latino	0.5241		0.2353		0.5534		0.3664	
COUNTY TYPE								
Large Metro	0.0733		0.7391		0.4363		0.4091	
Small Metro	0.4175		0.8452		0.7143		0.4908	
Nonmetro	0.0636		0.9199		0.0219		0.3225	

*Low precision; no estimate reported.

NOTE: Response categories for the Parental Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes." Respondents with unknown Parental Involvement data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.16C Measures of Student Characteristics in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	STUDENT CHARACTERISTICS							
	Participated in One or Fewer Activities ¹		Drug or Alcohol Prevention Messages Were Seen or Heard in School ²		Youth Liked a Lot or Kind of Liked Going to School ³		Grade Average for Last Completed Period Was "D" or Lower ⁴	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	81	88	256	235	277	247	42	51
AGE								
12-13	41	41	123	117	132	121	18	27
14-15	46	47	127	122	132	126	26	27
16-17	47	55	112	121	128	131	26	30
GENDER								
Male	61	61	158	151	173	166	34	37
Female	48	58	154	156	166	156	26	31
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	66	70	217	208	234	216	37	42
White	56	61	190	175	200	184	31	34
Black or African American	26	30	84	85	97	98	17	18
American Indian or Alaska Native	4	*	12	*	12	*	4	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	18	16	64	68	67	73	6	8
Two or More Races	11	10	32	36	35	36	9	10
Hispanic or Latino	46	50	130	112	143	120	19	28
COUNTY TYPE								
Large Metro	60	68	203	196	221	205	29	38
Small Metro	47	47	151	126	160	130	25	28
Nonmetro	30	30	89	94	101	103	17	18

*Low precision; no estimate reported.

NOTE: Respondents with unknown Student Characteristics data were excluded from the respective analyses.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

¹ The number of activities participated in is based on four questions: participation in school-based activities, community-based activities, church or faith-based activities, and other activities. For each question, the response categories include "None," "One," "Two," and "3 or more." Respondents were excluded from the analysis if they had unknown data for at least three of the questions and if their known responses did not place them in the recoded "Two or more" category.

² Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

³ Response categories for the School Satisfaction question include "You liked going to school a lot," "You kind of liked going to school," "You didn't like going to school very much," and "You hated going to school." The estimates in this table correspond to persons reporting "You liked going to school a lot" or "You kind of liked going to school."

⁴ Response categories for Grade Average question include "An 'A+', 'A,' or 'A-minus' average"; "A 'B+', 'B,' or 'B-minus' average"; "A 'C+', 'C,' or 'C-minus' average"; "A 'D' or less than a 'D' average"; and "My school does not give these grades." The estimates in this table correspond to persons reporting "A 'D' or less than a 'D' average." Respondents reporting that their school does not give these grades were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.16D Measures of Student Characteristics in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	STUDENT CHARACTERISTICS							
	Participated in One or Fewer Activities ¹		Drug or Alcohol Prevention Messages Were Seen or Heard in School ²		Youth Liked a Lot or Kind of Liked Going to School ³		Grade Average for Last Completed Period Was "D" or Lower ⁴	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.32	0.35	0.44	0.47	0.37	0.41	0.19	0.22
AGE								
12-13	0.53	0.56	0.76	0.84	0.59	0.68	0.28	0.39
14-15	0.52	0.55	0.66	0.73	0.65	0.69	0.33	0.35
16-17	0.54	0.62	0.75	0.81	0.62	0.73	0.34	0.37
GENDER								
Male	0.47	0.49	0.60	0.66	0.49	0.56	0.30	0.32
Female	0.40	0.46	0.60	0.61	0.54	0.60	0.24	0.28
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.33	0.36	0.50	0.52	0.41	0.46	0.21	0.24
White	0.39	0.43	0.58	0.62	0.48	0.56	0.25	0.27
Black or African American	0.74	0.91	1.17	1.20	0.92	0.96	0.56	0.57
American Indian or Alaska Native	2.58	*	3.93	*	4.49	*	2.71	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.45	1.33	2.49	1.98	2.37	1.78	0.52	0.72
Two or More Races	1.53	1.35	2.43	2.31	1.72	2.10	1.29	1.37
Hispanic or Latino	0.78	0.87	1.05	1.10	0.78	0.89	0.40	0.56
COUNTY TYPE								
Large Metro	0.43	0.49	0.64	0.65	0.55	0.57	0.24	0.30
Small Metro	0.60	0.61	0.74	0.84	0.62	0.71	0.36	0.43
Nonmetro	0.73	0.77	1.05	1.09	0.82	0.97	0.49	0.53

*Low precision; no estimate reported.

NOTE: Respondents with unknown Student Characteristics data were excluded from the respective analyses.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

¹ The number of activities participated in is based on four questions: participation in school-based activities, community-based activities, church or faith-based activities, and other activities. For each question, the response categories include "None," "One," "Two," and "3 or more." Respondents were excluded from the analysis if they had unknown data for at least three of the questions and if their known responses did not place them in the recoded "Two or more" category.

² Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

³ Response categories for the School Satisfaction question include "You liked going to school a lot," "You kind of liked going to school," "You didn't like going to school very much," and "You hated going to school." The estimates in this table correspond to persons reporting "You liked going to school a lot" or "You kind of liked going to school."

⁴ Response categories for Grade Average question include "An 'A+', 'A,' or 'A-minus' average"; "A 'B+', 'B,' or 'B-minus' average"; "A 'C+', 'C,' or 'C-minus' average"; "A 'D' or less than a 'D' average"; and "My school does not give these grades." The estimates in this table correspond to persons reporting "A 'D' or less than a 'D' average." Respondents reporting that their school does not give these grades were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.16P Measures of Student Characteristics in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	STUDENT CHARACTERISTICS							
	Participated in One or Fewer Activities ¹		Drug or Alcohol Prevention Messages Were Seen or Heard in School ²		Youth Liked a Lot or Kind of Liked Going to School ³		Grade Average for Last Completed Period Was "D" or Lower ⁴	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.4663		0.3838		0.0065		0.9376	
AGE								
12-13	0.5097		0.7233		0.7836		0.1564	
14-15	0.1842		0.1890		0.0436		0.5121	
16-17	0.6079		0.5720		0.0252		0.6787	
GENDER								
Male	0.0765		0.7736		0.1396		0.2728	
Female	0.3657		0.3018		0.0170		0.1359	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.3848		0.7517		0.0780		0.3792	
White	0.4158		0.9205		0.1140		0.5470	
Black or African American	0.8935		0.2210		0.3851		0.2818	
American Indian or Alaska Native	*		*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.4581		0.3441		0.7974		0.5058	
Two or More Races	0.8985		0.7365		0.2147		0.7692	
Hispanic or Latino	0.9464		0.2429		0.0101		0.1016	
COUNTY TYPE								
Large Metro	0.4814		0.3741		0.0155		0.4679	
Small Metro	0.8834		0.1854		0.2455		0.7406	
Nonmetro	0.8524		0.2417		0.4584		0.1149	

*Low precision; no estimate reported.

NOTE: Respondents with unknown Student Characteristics data were excluded from the respective analyses.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

¹ The number of activities participated in is based on four questions: participation in school-based activities, community-based activities, church or faith-based activities, and other activities. For each question, the response categories include "None," "One," "Two," and "3 or more." Respondents were excluded from the analysis if they had unknown data for at least three of the questions and if their known responses did not place them in the recoded "Two or more" category.

² Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

³ Response categories for the School Satisfaction question include "You liked going to school a lot," "You kind of liked going to school," "You didn't like going to school very much," and "You hated going to school." The estimates in this table correspond to persons reporting "You liked going to school a lot" or "You kind of liked going to school."

⁴ Response categories for Grade Average question include "An 'A+,' 'A,' or 'A-minus' average"; "A 'B+,' 'B,' or 'B-minus' average"; "A 'C+,' 'C,' or 'C-minus' average"; "A 'D' or less than a 'D' average"; and "My school does not give these grades." The estimates in this table correspond to persons reporting "A 'D' or less than a 'D' average." Respondents reporting that their school does not give these grades were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.17C Adults Always or Sometimes Took Part in Specific School-Related Behaviors in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	ADULT INVOLVEMENT							
	Teacher Let Youth Know He or She Was Doing a Good Job with School Work		Parents Checked That Youth Did Homework		Parents Provided Help with Homework		Parents Limited Time Youth Went Out with Friends on a School Night	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	269	248	282	247	274	251	247	223
AGE								
12-13	132	123	137	127	135	130	116	114
14-15	130	122	137	126	133	124	123	113
16-17	122	131	120	129	118	126	115	118
GENDER								
Male	171	168	178	165	171	168	154	142
Female	161	152	165	156	159	159	155	150
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	228	217	239	219	239	224	212	198
White	194	181	211	188	210	193	187	162
Black or African American	96	99	96	101	94	95	82	89
American Indian or Alaska Native	13	*	12	20	12	19	10	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	66	67	67	66	61	69	60	60
Two or More Races	33	37	36	38	35	38	27	35
Hispanic or Latino	142	120	143	121	128	112	121	108
COUNTY TYPE								
Large Metro	213	210	219	206	217	208	195	183
Small Metro	164	131	174	133	165	133	144	117
Nonmetro	101	96	107	102	103	103	90	92

*Low precision; no estimate reported.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

NOTE: Response categories for the Adult Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes." Respondents with unknown Adult Involvement data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.17D Adults Always or Sometimes Took Part in Specific School-Related Behaviors in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	ADULT INVOLVEMENT							
	Teacher Let Youth Know He or She Was Doing a Good Job with School Work		Parents Checked That Youth Did Homework		Parents Provided Help with Homework		Parents Limited Time Youth Went Out with Friends on a School Night	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.39	0.41	0.34	0.36	0.39	0.41	0.44	0.48
AGE								
12-13	0.60	0.64	0.45	0.51	0.55	0.54	0.76	0.79
14-15	0.63	0.72	0.54	0.60	0.64	0.65	0.72	0.80
16-17	0.68	0.70	0.68	0.70	0.75	0.77	0.78	0.81
GENDER								
Male	0.50	0.57	0.45	0.46	0.56	0.57	0.62	0.69
Female	0.55	0.58	0.50	0.55	0.54	0.58	0.61	0.65
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.41	0.47	0.37	0.40	0.42	0.44	0.48	0.54
White	0.50	0.55	0.44	0.48	0.42	0.49	0.55	0.62
Black or African American	0.90	1.04	0.89	0.93	1.10	1.15	1.13	1.25
American Indian or Alaska Native	3.13	*	3.40	3.38	3.73	3.66	4.53	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.88	1.98	1.90	1.95	2.77	2.42	2.50	2.41
Two or More Races	1.90	2.29	1.59	1.97	1.85	1.95	2.77	2.22
Hispanic or Latino	0.95	0.85	0.82	0.84	1.01	0.99	1.04	1.06
COUNTY TYPE								
Large Metro	0.57	0.60	0.46	0.50	0.57	0.59	0.62	0.67
Small Metro	0.67	0.69	0.60	0.63	0.64	0.68	0.80	0.77
Nonmetro	0.88	0.97	0.83	0.81	0.81	0.85	1.02	1.00

*Low precision; no estimate reported.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

NOTE: Response categories for the Adult Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes." Respondents with unknown Adult Involvement data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.17P Adults Always or Sometimes Took Part in Specific School-Related Behaviors in the Past Year among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	ADULT INVOLVEMENT							
	Teacher Let Youth Know He or She Was Doing a Good Job with School Work		Parents Checked That Youth Did Homework		Parents Provided Help with Homework		Parents Limited Time Youth Went Out with Friends on a School Night	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.0025		0.4870		0.0410		0.9585	
AGE								
12-13	0.2298		0.1992		0.4085		0.2968	
14-15	0.0027		0.8889		0.3734		0.2856	
16-17	0.2604		0.0702		0.0455		0.9107	
GENDER								
Male	0.3756		0.4654		0.0316		0.6666	
Female	0.0005		0.7673		0.4685		0.7064	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.0040		0.6656		0.2101		0.5537	
White	0.0063		0.8588		0.9575		0.0435	
Black or African American	0.5744		0.0326		0.2711		0.3086	
American Indian or Alaska Native	*		0.6123		0.3666		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.5618		0.4944		0.1817		0.5467	
Two or More Races	0.6972		0.0330		0.6270		0.0653	
Hispanic or Latino	0.2488		0.5377		0.0593		0.3581	
COUNTY TYPE								
Large Metro	0.0177		0.3759		0.1630		0.5918	
Small Metro	0.0897		0.5261		0.2720		0.8158	
Nonmetro	0.3856		0.3263		0.1203		0.1427	

*Low precision; no estimate reported.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

NOTE: Response categories for the Adult Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes." Respondents with unknown Adult Involvement data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.18C Youths Perceived That All or Most Students in Grade Level Use Specific Substances among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE OF STUDENTS IN GRADE LEVEL							
	Smoke Cigarettes		Use Marijuana or Hashish		Drink Alcoholic Beverages		Get Drunk at Least Once a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	97	92	127	126	145	145	85	81
AGE								
12-13	27	27	27	31	29	28	13	16
14-15	57	55	73	71	76	78	44	42
16-17	69	67	86	98	102	112	66	67
GENDER								
Male	61	61	80	81	89	89	53	50
Female	72	71	87	92	103	107	62	64
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	81	75	107	106	124	127	72	67
White	67	65	84	86	106	110	60	56
Black or African American	40	38	52	58	51	55	32	31
American Indian or Alaska Native	5	4	7	*	7	9	4	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	20	16	28	25	30	31	16	17
Two or More Races	14	15	18	21	19	22	11	15
Hispanic or Latino	52	52	70	71	77	71	45	44
COUNTY TYPE								
Large Metro	75	71	107	112	117	123	70	69
Small Metro	60	52	73	64	83	74	45	41
Nonmetro	37	36	36	37	50	52	27	25

*Low precision; no estimate reported.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months, or reported that they were "home-schooled" in the past 12 months were excluded from this analysis.

NOTE: Response categories for the Substance Use in Grade Level questions include "All of them," "Most of them," "A few of them," and "None of them." The estimates in this table correspond to persons reporting "All of them" or "Most of them." Respondents with unknown Substance Use in Grade Level data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.18D Youths Perceived That All or Most Students in Grade Level Use Specific Substances among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic/Geographic Characteristic	SUBSTANCE USE OF STUDENTS IN GRADE LEVEL							
	Smoke Cigarettes		Use Marijuana or Hashish		Drink Alcoholic Beverages		Get Drunk at Least Once a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.39	0.38	0.44	0.47	0.45	0.49	0.34	0.34
AGE								
12-13	0.39	0.40	0.39	0.45	0.42	0.40	0.20	0.24
14-15	0.65	0.65	0.73	0.80	0.75	0.81	0.54	0.53
16-17	0.75	0.75	0.82	0.87	0.82	0.88	0.72	0.73
GENDER								
Male	0.48	0.49	0.58	0.63	0.59	0.65	0.43	0.41
Female	0.58	0.58	0.62	0.68	0.68	0.72	0.51	0.56
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.41	0.39	0.47	0.51	0.50	0.54	0.37	0.36
White	0.47	0.47	0.53	0.58	0.57	0.65	0.42	0.41
Black or African American	1.06	1.00	1.17	1.35	1.33	1.37	1.00	0.90
American Indian or Alaska Native	3.57	3.09	4.75	*	4.66	5.12	2.71	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.58	1.40	2.21	2.08	2.01	2.28	1.34	1.39
Two or More Races	1.92	1.85	2.22	2.43	2.37	2.40	1.57	1.84
Hispanic or Latino	0.95	0.93	1.13	1.10	1.12	1.09	0.81	0.83
COUNTY TYPE								
Large Metro	0.52	0.51	0.63	0.71	0.61	0.73	0.50	0.50
Small Metro	0.79	0.67	0.86	0.79	0.82	0.83	0.59	0.60
Nonmetro	0.88	0.93	0.87	0.94	1.04	1.10	0.70	0.74

*Low precision; no estimate reported.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months, or reported that they were "home-schooled" in the past 12 months were excluded from this analysis.

NOTE: Response categories for the Substance Use in Grade Level questions include "All of them," "Most of them," "A few of them," and "None of them." The estimates in this table correspond to persons reporting "All of them" or "Most of them." Respondents with unknown Substance Use in Grade Level data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.18P Youths Perceived That All or Most Students in Grade Level Use Specific Substances among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year, by Demographic and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic/Geographic Characteristic	SUBSTANCE USE OF STUDENTS IN GRADE LEVEL							
	Smoke Cigarettes		Use Marijuana or Hashish		Drink Alcoholic Beverages		Get Drunk at Least Once a Week	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.0001		0.6346		0.0346		0.0113	
AGE								
12-13	0.2841		0.9008		0.1774		0.8697	
14-15	0.0571		0.1966		0.7525		0.0368	
16-17	0.0004		0.7569		0.0146		0.0479	
GENDER								
Male	0.0062		0.4361		0.0190		0.0401	
Female	0.0050		0.9577		0.4474		0.1127	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.0000		0.4378		0.0778		0.0078	
White	0.0019		0.7312		0.1965		0.0176	
Black or African American	0.0667		0.2427		0.1169		0.0641	
American Indian or Alaska Native	0.4786		*		0.9523		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.0387		0.5121		0.7156		0.9457	
Two or More Races	0.3978		0.5868		0.6171		0.6553	
Hispanic or Latino	0.3712		0.7212		0.2629		0.5744	
COUNTY TYPE								
Large Metro	0.0009		0.1790		0.0269		0.0488	
Small Metro	0.0859		0.2351		0.3719		0.1500	
Nonmetro	0.3540		0.6957		0.8883		0.4516	

*Low precision; no estimate reported.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months, or reported that they were "home-schooled" in the past 12 months were excluded from this analysis.

NOTE: Response categories for the Substance Use in Grade Level questions include "All of them," "Most of them," "A few of them," and "None of them." The estimates in this table correspond to persons reporting "All of them" or "Most of them." Respondents with unknown Substance Use in Grade Level data were excluded from the respective analyses.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 3.19C Past Month Use of Selected Substances, by Perceived Risk of Harm Associated with Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014

Perceptions of Risk	Used Illicit Drugs¹	Used Marijuana	Used Cigarettes	Binge Use of Alcohol²
CIGARETTES				
Smoke One or More Packs Per Day – Great Risk	61	55	35	49
Smoke One or More Packs Per Day – Moderate/Slight/No Risk	46	41	39	38
MARIJUANA				
Smoke Once a Month – Great Risk	17	10	13	13
Smoke Once a Month – Moderate/Slight/No Risk	76	68	51	61
Smoke Once or Twice a Week – Great Risk	22	12	16	16
Smoke Once or Twice a Week – Moderate/Slight/No Risk	74	68	51	59
COCAINE				
Use Once a Month – Great Risk	55	50	34	43
Use Once a Month – Moderate/Slight/No Risk	51	46	37	41
Use Once or Twice a Week – Great Risk	65	58	42	54
Use Once or Twice a Week – Moderate/Slight/No Risk	39	34	28	30
HEROIN				
Try Once or Twice – Great Risk	64	59	42	53
Try Once or Twice – Moderate/Slight/No Risk	41	33	27	31
Use Once or Twice a Week – Great Risk	69	64	46	57
Use Once or Twice a Week – Moderate/Slight/No Risk	29	23	21	22
LSD				
Try Once or Twice – Great Risk	43	39	31	37
Try Once or Twice – Moderate/Slight/No Risk	61	52	40	47
Use Once or Twice a Week – Great Risk	57	51	37	47
Use Once or Twice a Week – Moderate/Slight/No Risk	48	42	34	36
ALCOHOL				
Have Four or Five Drinks Nearly Every Day – Great Risk	54	46	32	38
Have Four or Five Drinks Nearly Every Day – Moderate/Slight/No Risk	56	51	41	48
Have Five or More Drinks Once or Twice a Week – Great Risk	41	34	25	27
Have Five or More Drinks Once or Twice a Week – Moderate/Slight/No Risk	66	59	47	56

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Risk data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.19D Past Month Use of Selected Substances, by Perceived Risk of Harm Associated with Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Percentages, 2014

Perceptions of Risk	Used Illicit Drugs¹	Used Marijuana	Used Cigarettes	Binge Use of Alcohol²
CIGARETTES				
Smoke One or More Packs Per Day – Great Risk	0.36	0.33	0.21	0.29
Smoke One or More Packs Per Day – Moderate/Slight/No Risk	0.52	0.47	0.46	0.44
MARIJUANA				
Smoke Once a Month – Great Risk	0.29	0.18	0.23	0.24
Smoke Once a Month – Moderate/Slight/No Risk	0.38	0.34	0.26	0.31
Smoke Once or Twice a Week – Great Risk	0.23	0.14	0.18	0.18
Smoke Once or Twice a Week – Moderate/Slight/No Risk	0.45	0.41	0.32	0.36
COCAINE				
Use Once a Month – Great Risk	0.43	0.39	0.28	0.34
Use Once a Month – Moderate/Slight/No Risk	0.40	0.36	0.29	0.32
Use Once or Twice a Week – Great Risk	0.32	0.29	0.22	0.28
Use Once or Twice a Week – Moderate/Slight/No Risk	0.67	0.59	0.49	0.53
HEROIN				
Try Once or Twice – Great Risk	0.42	0.39	0.29	0.36
Try Once or Twice – Moderate/Slight/No Risk	0.39	0.32	0.27	0.30
Use Once or Twice a Week – Great Risk	0.34	0.31	0.23	0.28
Use Once or Twice a Week – Moderate/Slight/No Risk	0.57	0.46	0.43	0.44
LSD				
Try Once or Twice – Great Risk	0.38	0.35	0.28	0.34
Try Once or Twice – Moderate/Slight/No Risk	0.45	0.40	0.31	0.36
Use Once or Twice a Week – Great Risk	0.35	0.31	0.23	0.29
Use Once or Twice a Week – Moderate/Slight/No Risk	0.60	0.54	0.45	0.47
ALCOHOL				
Have Four or Five Drinks Nearly Every Day – Great Risk	0.34	0.29	0.21	0.25
Have Four or Five Drinks Nearly Every Day – Moderate/Slight/No Risk	0.57	0.51	0.42	0.50
Have Five or More Drinks Once or Twice a Week – Great Risk	0.41	0.35	0.26	0.28
Have Five or More Drinks Once or Twice a Week – Moderate/Slight/No Risk	0.41	0.37	0.31	0.36

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Risk data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.20C Past Month Use of Selected Substances, by Perceived Availability of Selected Illicit Drugs among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014

Perceived Availability	Used Illicit Drugs ¹	Used Marijuana	Used Cigarettes	Binge Use of Alcohol ²
FAIRLY OR VERY EASY TO OBTAIN				
Marijuana	72	67	51	59
Cocaine	42	37	29	35
Crack	35	30	25	28
Heroin	28	23	22	23
LSD	39	35	30	32
FAIRLY OR VERY DIFFICULT, OR PROBABLY IMPOSSIBLE TO OBTAIN				
Marijuana	29	18	17	22
Cocaine	65	57	43	48
Crack	70	62	44	54
Heroin	72	65	46	55
LSD	64	56	41	49

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceived Availability data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.20D Past Month Use of Selected Substances, by Perceived Availability of Selected Illicit Drugs among Persons Aged 12 to 17: Standard Errors of Percentages, 2014

Perceived Availability	Used Illicit Drugs ¹	Used Marijuana	Used Cigarettes	Binge Use of Alcohol ²
FAIRLY OR VERY EASY TO OBTAIN				
Marijuana	0.56	0.52	0.41	0.48
Cocaine	1.08	0.98	0.78	0.94
Crack	0.96	0.85	0.69	0.78
Heroin	1.16	1.01	0.93	1.00
LSD	1.23	1.16	1.02	1.07
FAIRLY OR VERY DIFFICULT, OR PROBABLY IMPOSSIBLE TO OBTAIN				
Marijuana	0.22	0.14	0.13	0.17
Cocaine	0.31	0.27	0.21	0.23
Crack	0.33	0.29	0.21	0.26
Heroin	0.32	0.29	0.21	0.25
LSD	0.30	0.27	0.20	0.23

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceived Availability data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.21C Past Month Use of Selected Substances, by Youth's Perceptions of Parents' Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014

Feelings about Substance Use Behaviors	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Strongly Disapprove	Somewhat Disapprove/ Approve Nor Neither Disapprove	Strongly Disapprove	Somewhat Disapprove/ Approve Nor Neither Disapprove	Strongly Disapprove	Somewhat Disapprove/ Approve Nor Neither Disapprove	Strongly Disapprove	Somewhat Disapprove/ Approve Nor Neither Disapprove
YOUTH'S PERCEPTIONS OF PARENTS' FEELINGS ABOUT YOUTH...								
Smoking One or More Packs of Cigarettes Per Day	68	31	61	28	40	31	55	25
Trying Marijuana or Hashish Once or Twice	58	51	49	48	37	34	48	37
Using Marijuana or Hashish Once a Month or More	58	48	49	44	38	33	50	35
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	67	36	59	33	41	28	53	31

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Parents' Feelings data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.21D Past Month Use of Selected Substances, by Youth's Perceptions of Parents' Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Percentages, 2014

Feelings about Substance Use Behaviors	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Strongly Disapprove	Somewhat Disapprove/ Neither Approve Nor Disapprove	Strongly Disapprove	Somewhat Disapprove/ Neither Approve Nor Disapprove	Strongly Disapprove	Somewhat Disapprove/ Neither Approve Nor Disapprove	Strongly Disapprove	Somewhat Disapprove/ Neither Approve Nor Disapprove
YOUTH'S PERCEPTIONS OF PARENTS' FEELINGS ABOUT YOUTH...								
Smoking One or More Packs of Cigarettes Per Day	0.29	1.68	0.26	1.55	0.17	1.64	0.23	1.40
Trying Marijuana or Hashish Once or Twice	0.26	1.32	0.22	1.27	0.17	1.01	0.22	1.07
Using Marijuana or Hashish Once a Month or More	0.25	1.46	0.22	1.39	0.17	1.22	0.22	1.18
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	0.29	1.31	0.26	1.24	0.18	1.11	0.23	1.20

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Parents' Feelings data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.22C Past Month Use of Selected Substances, by Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014

Feelings about Substance Use Behaviors	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove
YOUTH'S PERCEPTIONS OF CLOSE FRIENDS' FEELINGS ABOUT YOUTH...								
Smoking One or More Packs of Cigarettes Per Day	67	36	59	33	38	33	53	28
Trying Marijuana or Hashish Once or Twice	41	66	31	61	27	43	36	51
Using Marijuana or Hashish Once a Month or More	40	64	30	59	26	43	35	52
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	60	44	54	40	35	34	46	38
YOUTH'S FEELINGS ABOUT PEERS...								
Smoking One or More Packs of Cigarettes Per Day	69	30	61	28	41	29	54	26
Trying Marijuana or Hashish Once or Twice	45	63	35	59	31	42	39	49
Using Marijuana or Hashish Once a Month or More	40	64	31	60	30	41	35	49
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	63	42	56	38	38	33	49	37

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Close Friends' Feelings or Feelings about Peers data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.22D Past Month Use of Selected Substances, by Feelings about Substance Use Behaviors among Persons Aged 12 to 17: Standard Errors of Percentages, 2014

Feelings about Substance Use Behaviors	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove	Strongly/ Somewhat Disapprove	Neither Approve Nor Disapprove
YOUTH'S PERCEPTIONS OF CLOSE FRIENDS' FEELINGS ABOUT YOUTH...								
Smoking One or More Packs of Cigarettes Per Day	0.29	1.28	0.26	1.21	0.17	1.24	0.23	1.07
Trying Marijuana or Hashish Once or Twice	0.21	0.99	0.16	0.95	0.14	0.75	0.19	0.86
Using Marijuana or Hashish Once a Month or More	0.20	1.03	0.15	1.01	0.13	0.80	0.18	0.93
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	0.27	1.23	0.25	1.18	0.17	1.06	0.21	1.13
YOUTH'S FEELINGS ABOUT PEERS...								
Smoking One or More Packs of Cigarettes Per Day	0.29	1.44	0.26	1.37	0.18	1.41	0.23	1.31
Trying Marijuana or Hashish Once or Twice	0.22	1.00	0.17	0.97	0.16	0.77	0.20	0.87
Using Marijuana or Hashish Once a Month or More	0.20	0.99	0.16	0.96	0.16	0.75	0.18	0.86
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	0.28	1.36	0.25	1.29	0.17	1.15	0.22	1.26

*Low precision; no estimate reported.

NOTE: Respondents with unknown Perceptions of Close Friends' Feelings or Feelings about Peers data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.23C Past Month Use of Selected Substances, by Measures of Youth's Exposure to Prevention Programs and Messages and Youth's Delinquency in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014

Prevention Programs/Messages and Delinquency	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior
EXPOSURE TO PREVENTION PROGRAMS AND MESSAGES IN PAST YEAR								
Participated in Problem-Solving, Communication Skills, or Self-Esteem Group	34	70	26	63	20	47	23	57
Participated in Violence Prevention Program	25	73	20	66	16	50	19	58
Participated in Pregnancy or Sexually Transmitted Disease Prevention Program	21	75	19	65	16	49	15	59
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School	22	74	20	65	16	49	17	60
Participated in Program to Help Deal with Drug or Alcohol Use	17	75	15	67	12	51	11	61
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School	64	39	59	35	43	28	53	30
Talked to Parents about Dangers of Tobacco, Alcohol, or Drug Use	58	47	52	42	38	35	44	41
DELINQUENT BEHAVIOR IN PAST YEAR³								
Got into a Serious Fight at School or Work	38	67	32	60	29	42	29	54
Took Part in a Group-Against-Group Fight	36	71	33	61	22	47	28	55
Attacked Someone with the Intent to Seriously Hurt Them	25	74	19	66	18	49	20	59
Carried a Handgun	18	75	18	67	14	50	16	58
Sold Illegal Drugs	27	73	26	65	19	49	23	57
Stole or Tried to Steal Anything Worth More Than \$50	25	73	23	64	17	49	20	59

*Low precision; no estimate reported.

NOTE: Respondents with unknown Prevention Programs and Messages Exposure or Delinquent Behavior data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.23D Past Month Use of Selected Substances, by Measures of Youth's Exposure to Prevention Programs and Messages and Youth's Delinquency in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2014

Prevention Programs/Messages and Delinquency	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior	Had Exposure or Behavior	Did Not Have Exposure or Behavior
EXPOSURE TO PREVENTION PROGRAMS AND MESSAGES IN PAST YEAR								
Participated in Problem-Solving, Communication Skills, or Self-Esteem Group	0.57	0.35	0.46	0.32	0.35	0.24	0.40	0.29
Participated in Violence Prevention Program	0.76	0.32	0.62	0.29	0.49	0.23	0.59	0.26
Participated in Pregnancy or Sexually Transmitted Disease Prevention Program	0.93	0.31	0.84	0.28	0.72	0.21	0.68	0.25
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School	0.79	0.32	0.72	0.28	0.57	0.22	0.61	0.26
Participated in Program to Help Deal with Drug or Alcohol Use	1.61	0.30	1.48	0.27	1.18	0.21	1.13	0.25
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School	0.35	0.55	0.32	0.51	0.23	0.42	0.29	0.44
Talked to Parents about Dangers of Tobacco, Alcohol, or Drug Use	0.40	0.41	0.36	0.38	0.27	0.33	0.31	0.37
DELINQUENT BEHAVIOR IN PAST YEAR³								
Got into a Serious Fight at School or Work	0.83	0.31	0.74	0.28	0.66	0.20	0.68	0.25
Took Part in a Group-Against-Group Fight	1.21	0.30	1.14	0.27	0.80	0.21	0.99	0.24
Attacked Someone with the Intent to Seriously Hurt Them	1.79	0.30	1.47	0.27	1.41	0.20	1.48	0.24
Carried a Handgun	1.75	0.30	1.71	0.27	1.42	0.20	1.59	0.24
Sold Illegal Drugs	3.23	0.29	3.24	0.26	3.07	0.20	3.35	0.23
Stole or Tried to Steal Anything Worth More Than \$50	2.90	0.29	2.76	0.26	2.26	0.20	2.63	0.24

*Low precision; no estimate reported.

NOTE: Respondents with unknown Prevention Programs and Messages Exposure or Delinquent Behavior data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as having participated if they indicated taking part in the particular behavior at least one time.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.24C Past Month Use of Selected Substances, by Measures of Risk and Protective Factors, Religious Beliefs, and Past Year Parental Involvement among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2014

Risk and Protective Factors, Religious Beliefs, and Parental Involvement	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement
RISK AND PROTECTIVE FACTORS								
Youth Moved 2 or More Times in the Past 5 Years	39	64	35	58	28	41	27	55
Youth Argued with at Least One Parent 10 or More Times in the Past Year	49	61	42	56	31	42	38	48
Youth Was Approached by Someone Selling Drugs in the Past Month	53	57	49	50	35	39	39	47
Youth Had Someone to Talk to about Problems ³	73	21	66	19	48	14	59	17
Youth Attended 25 or More Religious Services in the Past Year ⁴	36	68	29	63	21	48	26	55
YOUTH RELIGIOUS BELIEFS⁵								
Religious Beliefs Are a Very Important Part of Youth's Life	58	53	49	50	33	37	45	40
Religious Beliefs Influence Youth's Decision Making in Life	54	58	46	55	32	40	39	48
It Is Important for Friends to Share Religious Beliefs	36	67	29	63	20	48	25	57
PARENTAL INVOLVEMENT IN PAST YEAR⁶								
Parents Made Youth Do Chores around the House	72	27	64	24	49	17	57	22
Parents Limited Amount of Time Youth Watched TV	38	65	32	59	24	46	29	53
Parents Let Youth Know He or She Had Done a Good Job	68	36	61	34	44	26	53	29
Parents Let Youth Know They Were Proud of Something He or She Had Done	67	36	60	34	44	27	53	30

*Low precision; no estimate reported.

NOTE: Respondents with unknown Risk and Protective Factors, Religious Beliefs, or Parental Involvement data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Respondents were able to select multiple categories of persons that they would talk to about a serious problem, including the choice of "nobody." Respondents who reported both a person that they would talk to and that they had nobody to talk to were excluded from the analysis.

⁴ Special occasions, such as weddings, funerals, or other special events, were not to have been considered in the number of times attended religious services.

⁵ Response categories for the questions on Religious Beliefs include "Strongly disagree," "Disagree," "Agree," and "Strongly agree." The estimates in this table correspond to persons reporting "Agree" or "Strongly agree."

⁶ Response categories for the Parental Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.24D Past Month Use of Selected Substances, by Measures of Risk and Protective Factors, Religious Beliefs, and Past Year Parental Involvement among Persons Aged 12 to 17: Standard Errors of Percentages, 2014

Risk and Protective Factors, Religious Beliefs, and Parental Involvement	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement	Had Factor, Belief, or Involvement	Did Not Have Factor, Belief, or Involvement
RISK AND PROTECTIVE FACTORS								
Youth Moved 2 or More Times in the Past 5 Years	0.63	0.33	0.58	0.30	0.47	0.22	0.46	0.28
Youth Argued with at Least One Parent 10 or More Times in the Past Year	0.87	0.30	0.77	0.28	0.60	0.21	0.71	0.24
Youth Was Approached by Someone Selling Drugs in the Past Month	1.35	0.25	1.31	0.22	1.06	0.18	1.15	0.21
Youth Had Someone to Talk to about Problems ³	0.30	1.72	0.27	1.59	0.21	1.22	0.24	1.44
Youth Attended 25 or More Religious Services in the Past Year ⁴	0.48	0.37	0.39	0.35	0.29	0.27	0.36	0.31
YOUTH RELIGIOUS BELIEFS⁵								
Religious Beliefs Are a Very Important Part of Youth's Life	0.32	0.69	0.28	0.66	0.19	0.52	0.25	0.55
Religious Beliefs Influence Youth's Decision Making in Life	0.33	0.61	0.29	0.59	0.20	0.44	0.25	0.52
It Is Important for Friends to Share Religious Beliefs	0.47	0.38	0.39	0.36	0.27	0.28	0.33	0.32
PARENTAL INVOLVEMENT IN PAST YEAR⁶								
Parents Made Youth Do Chores around the House	0.31	0.95	0.28	0.86	0.22	0.64	0.25	0.80
Parents Limited Amount of Time Youth Watched TV	0.37	0.41	0.32	0.38	0.24	0.31	0.29	0.35
Parents Let Youth Know He or She Had Done a Good Job	0.31	0.97	0.28	0.94	0.20	0.72	0.24	0.79
Parents Let Youth Know They Were Proud of Something He or She Had Done	0.30	0.96	0.28	0.91	0.21	0.75	0.25	0.80

*Low precision; no estimate reported.

NOTE: Respondents with unknown Risk and Protective Factors, Religious Beliefs, or Parental Involvement data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Respondents were able to select multiple categories of persons that they would talk to about a serious problem, including the choice of "nobody." Respondents who reported both a person that they would talk to and that they had nobody to talk to were excluded from the analysis.

⁴ Special occasions, such as weddings, funerals, or other special events, were not to have been considered in the number of times attended religious services.

⁵ Response categories for the questions on Religious Beliefs include "Strongly disagree," "Disagree," "Agree," and "Strongly agree." The estimates in this table correspond to persons reporting "Agree" or "Strongly agree."

⁶ Response categories for the Parental Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.25C Past Month Use of Selected Substances, by Measures of Scholastic Environment among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year: Standard Errors of Numbers in Thousands, 2014

Scholastic Environment	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic
STUDENT CHARACTERISTICS IN PAST YEAR³								
Participated in One or Fewer Activities ⁴	33	69	31	61	26	44	26	57
Drug or Alcohol Prevention Messages Were Seen or Heard in School ⁵	61	43	55	38	38	33	48	34
Youth Liked a Lot or Kind of Liked Going to School ⁶	61	41	54	39	39	32	46	37
Grade Average for Last Completed Period Was "D" or Lower ⁷	21	71	19	64	20	45	19	57
ADULT INVOLVEMENT IN PAST YEAR^{3,8}								
Teacher Let Youth Know He or She Was Doing a Good Job with School Work	66	37	59	34	40	30	52	30
Parents Checked That Youth Did Homework	62	42	53	38	42	30	48	33
Parents Provided Help with Homework	62	44	54	41	44	27	48	35
Parents Limited Time Youth Went Out with Friends on a School Night	57	46	51	41	42	30	48	37
SUBSTANCE USE IN GRADE LEVEL⁹								
Students in Youth's Grade Smoke Cigarettes	38	67	34	58	32	39	32	51
Students in Youth's Grade Use Marijuana or Hashish	62	43	58	34	41	29	49	34
Students in Youth's Grade Drink Alcoholic Beverages	60	43	55	36	42	28	52	27
Students in Youth's Grade Get Drunk at Least Once a Week	38	66	36	57	27	40	34	50

*Low precision; no estimate reported.

NOTE: Respondents with unknown Student Characteristics, Adult Involvement, or Substance Use in Grade Level data were excluded from the respective analyses.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

⁴ The number of activities participated in is based on four questions: participation in school-based activities, community-based activities, church or faith-based activities, and other activities. For each question, the response categories include "None," "One," "Two," and "3 or more." Respondents were excluded from the analysis if they had unknown data for at least three of the questions and if their known responses did not place them in the recoded "Two or more" category.

⁵ Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

⁶ Response categories for the School Satisfaction question include "You liked going to school a lot," "You kind of liked going to school," "You didn't like going to school very much," and "You hated going to school." The estimates in this table correspond to persons reporting "You liked going to school a lot" or "You kind of liked going to school."

⁷ Response categories for Grade Average question include "An 'A+', 'A,' or 'A-minus' average"; "A 'B+', 'B,' or 'B-minus' average"; "A 'C+', 'C,' or 'C-minus' average"; "A 'D' or less than a 'D' average"; and "My school does not give these grades." The estimates in this table correspond to persons reporting "A 'D' or less than a 'D' average." Respondents reporting that their school does not give these grades were excluded from the analysis.

⁸ Response categories for the Adult Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes."

⁹ Response categories for the Substance Use in Grade Level questions include "All of them," "Most of them," "A few of them," and "None of them." The estimates in this table correspond to persons reporting "All of them" or "Most of them." Youths reporting that they were "home-schooled" in the past 12 months were not considered to be enrolled in school.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 3.25D Past Month Use of Selected Substances, by Measures of Scholastic Environment among Persons Aged 12 to 17 Who Were Enrolled in School in the Past Year: Standard Errors of Percentages, 2014

Scholastic Environment	SUBSTANCE USE IN THE PAST MONTH							
	Used Illicit Drugs ¹		Used Marijuana		Used Cigarettes		Binge Use of Alcohol ²	
	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic	Had Scholastic Characteristic	Did Not Have Scholastic Characteristic
STUDENT CHARACTERISTICS IN PAST YEAR³								
Participated in One or Fewer Activities ⁴	1.07	0.32	1.03	0.29	0.85	0.21	0.86	0.27
Drug or Alcohol Prevention Messages Were Seen or Heard in School ⁵	0.34	0.64	0.31	0.58	0.22	0.51	0.27	0.52
Youth Liked a Lot or Kind of Liked Going to School ⁶	0.32	0.80	0.28	0.76	0.21	0.64	0.24	0.74
Grade Average for Last Completed Period Was "D" or Lower ⁷	1.94	0.32	1.85	0.29	1.90	0.21	1.83	0.26
ADULT INVOLVEMENT IN PAST YEAR^{3,8}								
Teacher Let Youth Know He or She Was Doing a Good Job with School Work	0.34	0.70	0.31	0.64	0.22	0.58	0.28	0.58
Parents Checked That Youth Did Homework	0.31	0.97	0.27	0.90	0.21	0.72	0.24	0.81
Parents Provided Help with Homework	0.31	0.90	0.27	0.85	0.23	0.58	0.25	0.73
Parents Limited Time Youth Went Out with Friends on a School Night	0.35	0.59	0.32	0.54	0.26	0.40	0.29	0.49
SUBSTANCE USE IN GRADE LEVEL⁹								
Students in Youth's Grade Smoke Cigarettes	0.94	0.34	0.88	0.30	0.84	0.20	0.80	0.27
Students in Youth's Grade Use Marijuana or Hashish	0.85	0.26	0.81	0.21	0.62	0.18	0.71	0.21
Students in Youth's Grade Drink Alcoholic Beverages	0.71	0.29	0.65	0.24	0.52	0.19	0.62	0.18
Students in Youth's Grade Get Drunk at Least Once a Week	1.27	0.33	1.19	0.29	0.96	0.21	1.16	0.26

*Low precision; no estimate reported.

NOTE: Respondents with unknown Student Characteristics, Adult Involvement, or Substance Use in Grade Level data were excluded from the respective analyses.

NOTE: Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

⁴ The number of activities participated in is based on four questions: participation in school-based activities, community-based activities, church or faith-based activities, and other activities. For each question, the response categories include "None," "One," "Two," and "3 or more." Respondents were excluded from the analysis if they had unknown data for at least three of the questions and if their known responses did not place them in the recoded "Two or more" category.

⁵ Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

⁶ Response categories for the School Satisfaction question include "You liked going to school a lot," "You kind of liked going to school," "You didn't like going to school very much," and "You hated going to school." The estimates in this table correspond to persons reporting "You liked going to school a lot" or "You kind of liked going to school."

⁷ Response categories for Grade Average question include "An 'A+', 'A,' or 'A-minus' average"; "A 'B+', 'B,' or 'B-minus' average"; "A 'C+', 'C,' or 'C-minus' average"; "A 'D' or less than a 'D' average"; and "My school does not give these grades." The estimates in this table correspond to persons reporting "A 'D' or less than a 'D' average." Respondents reporting that their school does not give these grades were excluded from the analysis.

⁸ Response categories for the Adult Involvement questions include "Always," "Sometimes," "Seldom," and "Never." The estimates in this table correspond to persons reporting "Always" or "Sometimes."

⁹ Response categories for the Substance Use in Grade Level questions include "All of them," "Most of them," "A few of them," and "None of them." The estimates in this table correspond to persons reporting "All of them" or "Most of them." Youths reporting that they were "home-schooled" in the past 12 months were not considered to be enrolled in school.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2014.

Table 4.1B Past Year Initiation of Illicit Drug Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Illicit Drug Use, and Past Year Illicit Drug Users Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	2,848	3,011	1.1	1.1	2.1	2.2	6.8	6.8
AGE								
12-25	2,459	2,541	4.1	4.2	6.7	7.0	14.7	15.0
12-17	1,353	1,410	5.4	5.7	6.6	6.9	31.6	32.5
12-13	210	234	2.6	3.0	2.8	3.2	45.1	42.0
14-15	510	529	6.0	6.3	7.1	7.4	37.9	38.7
16-17	633	648	7.5	7.6	10.8	10.7	25.5	26.8
18-25	1,106	1,131	3.2	3.2	6.9	7.1	8.9	9.0
18-20	755	741	5.7	5.7	10.6	10.5	14.9	15.0
21-25	351	391	1.6	1.8	3.9	4.4	4.7	5.1
26 or Older	389	469	0.2	0.2	0.4	0.5	1.6	1.7
GENDER								
Male	1,187 ^a	1,394	0.9 ^a	1.1	1.9 ^a	2.3	5.0	5.5
Female	1,661	1,617	1.2	1.2	2.2	2.1	9.3	8.7
GENDER/AGE								
Male 12-17	633	633	5.0	5.0	6.2	6.1	28.0	30.0
Female 12-17	720	777	5.9	6.4	7.1	7.7	35.6	34.9

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.2B Past Year Initiation of Marijuana Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Marijuana Use, and Past Year Marijuana Users Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	2,427	2,568	0.9	1.0	1.6	1.7	7.4	7.3
AGE								
12-25	2,217	2,298	3.7	3.8	5.6	5.8	15.5	15.9
12-17	1,200	1,203	4.8	4.8	5.5	5.5	35.9	36.9
12-13	129	133	1.6	1.7	1.6	1.7	65.3	71.6
14-15	451	461	5.3	5.4	5.8	5.9	46.3	48.2
16-17	620	609	7.4	7.2	9.7	9.5	28.5	28.7
18-25	1,017	1,094	2.9	3.1	5.7	6.2	9.2	9.8
18-20	696	755	5.3	5.8	9.0	9.7	15.4	16.8
21-25	320	340	1.5	1.6	3.2	3.4	4.9	5.1
26 or Older	210	271	0.1	0.1	0.2	0.2	1.1	1.3
GENDER								
Male	1,072	1,226	0.8	1.0	1.6	1.9	5.4	5.8
Female	1,355	1,342	1.0	1.0	1.6	1.6	10.2	9.6
GENDER/AGE								
Male 12-17	557	552	4.4	4.4	5.0	5.0	31.1	33.0
Female 12-17	643	651	5.3	5.3	5.9	6.0	41.4	41.0

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.3B Past Year Initiation of Cigarette Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Cigarette Use, and Past Year Cigarette Users Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	2,071	2,164	0.8	0.8	2.0	2.1	3.1	3.3
AGE								
12-25	1,963	2,019	3.3	3.4	5.2	5.2	12.0	13.1
12-17	932	838	3.7	3.4	4.3	3.8	36.4	37.9
12-13	92	71	1.2	0.9	1.2	0.9	49.9	*
14-15	326	268	3.8	3.2	4.3 ^a	3.5	42.2	44.1
16-17	514	500	6.1	5.9	7.8	7.4	32.0	34.1
18-25	1,031	1,181	3.0	3.4	6.6	7.2	7.5 ^a	9.0
18-20	766	807	5.8	6.2	10.2	10.3	16.0	18.1
21-25	265 ^a	374	1.2 ^a	1.7	3.3 ^a	4.3	3.0 ^b	4.3
26 or Older	108	144	0.1	0.1	0.2	0.2	0.2	0.3
GENDER								
Male	1,048	1,074	0.8	0.8	2.4	2.4	2.9	3.0
Female	1,023	1,089	0.8	0.8	1.7	1.8	3.4	3.6
GENDER/AGE								
Male 12-17	470 ^a	379	3.7 ^a	3.0	4.2 ^a	3.4	35.8	32.8
Female 12-17	462	459	3.8	3.8	4.3	4.2	37.0 ^a	43.5

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.4B Past Year Initiation of Alcohol Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Alcohol Use, and Past Year Alcohol Users Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	4,559	4,655	1.7	1.8	8.6	9.0	2.6	2.6
AGE								
12-25	4,473	4,561	7.5	7.6	16.4	16.4	13.6	14.0
12-17	2,417	2,335	9.7	9.4	12.3	11.8	39.4	39.1
12-13	230	242	2.9	3.1	3.1	3.2	49.0	54.7
14-15	929	902	10.9	10.7	13.2	12.8	49.0	50.8
16-17	1,258	1,191	15.0	14.0	24.4	22.5	33.4	31.7
18-25	2,056	2,225	5.9	6.4	26.7	27.7	7.7	8.3
18-20	1,308	1,365	9.9	10.4	26.6	26.8	15.3	16.4
21-25	748	861	3.5	3.9	27.0	29.3	4.1	4.7
26 or Older	85	95	0.0	0.0	0.3	0.4	0.1	0.1
GENDER								
Male	2,119	2,151	1.7	1.7	9.6	9.8	2.4	2.4
Female	2,439	2,504	1.8	1.8	7.8	8.3	2.8	2.9
GENDER/AGE								
Male 12-17	1,097	1,032	8.6	8.1	11.0	10.2	36.8	36.9
Female 12-17	1,320	1,303	10.8	10.7	13.6	13.4	41.9	40.9

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.5B Past Year Initiation of Substance Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 or Older: Numbers in Thousands and Percentages, 2013 and 2014

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS^{2,3}	2,848	3,011	1.1	1.1	2.1	2.2	6.8	6.8
Marijuana and Hashish	2,427	2,568	0.9	1.0	1.6	1.7	7.4	7.3
Cocaine	601 ^a	766	0.2 ^a	0.3	0.3 ^a	0.3	14.4	16.8
Crack	58	109	0.0	0.0	0.0	0.0	9.1	14.1
Heroin	169	212	0.1	0.1	0.1	0.1	24.9	23.1
Hallucinogens	1,065	936	0.4	0.4	0.5	0.4	24.0	22.0
LSD	482	586	0.2	0.2	0.2	0.2	43.3	45.4
PCP	32	41	0.0	0.0	0.0	0.0	*	*
Ecstasy	751	676	0.3	0.3	0.3	0.3	29.0	28.9
Inhalants	563	512	0.2	0.2	0.2	0.2	36.7	31.7
Nonmedical Use of Psychotherapeutics ^{3,4}	2,006	2,111	0.8	0.8	0.9	1.0	13.1	14.3
Pain Relievers	1,539	1,425	0.6	0.5	0.7	0.6	13.9	13.8
OxyContin [®]	436	300	0.2	0.1	0.2	0.1	30.2	24.3
Tranquilizers	1,180	1,133	0.4	0.4	0.5	0.5	22.4	21.8
Stimulants ³	603	690	0.2	0.3	0.2	0.3	18.5	20.2
Methamphetamine ³	144	183	0.1	0.1	0.1	0.1	14.7	18.1
Sedatives	128	173	0.0	0.1	0.1	0.1	20.1	22.3
ILLICIT DRUGS OTHER THAN MARIJUANA^{2,3}	2,220	2,413	0.8	0.9	1.2	1.3	11.2	12.2
CIGARETTES	2,071	2,164	0.8	0.8	2.0	2.1	3.1	3.3
Daily Cigarette Use ⁵	813	756	0.3	0.3	0.5	0.4	--	--
SMOKELESS TOBACCO⁶	1,116	1,016	0.4	0.4	0.5	0.5	9.4	8.8
CIGARS	2,770	2,597	1.1	1.0	1.6	1.5	11.1	10.9
ALCOHOL	4,559	4,655	1.7	1.8	8.6	9.0	2.6	2.6

*Low precision; no estimate reported.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.6B Past Year Initiation of Substance Use among Persons Aged 12 to 17, Persons Aged 12 to 17 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 to 17: Numbers in Thousands and Percentages, 2013 and 2014

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS^{2,3}	1,353	1,410	5.4	5.7	6.6	6.9	31.6	32.5
Marijuana and Hashish	1,200	1,203	4.8	4.8	5.5	5.5	35.9	36.9
Cocaine	94	117	0.4	0.5	0.4	0.5	71.5	*
Crack	10	11	0.0	0.0	0.0	0.0	*	*
Heroin	21	13	0.1	0.1	0.1	0.1	*	*
Hallucinogens	306	258	1.2	1.0	1.2	1.1	63.6	61.1
LSD	122	165	0.5	0.7	0.5	0.7	81.4	77.2
PCP	19	17	0.1	0.1	0.1	0.1	*	*
Ecstasy	182 ^a	125	0.7 ^a	0.5	0.7 ^a	0.5	78.6	68.8
Inhalants	254	271	1.0	1.1	1.1	1.1	52.9	52.9
Nonmedical Use of Psychotherapeutics ^{3,4}	601	641	2.4	2.6	2.6	2.8	41.8	41.4
Pain Relievers	508	489	2.0	2.0	2.2	2.1	43.9	42.0
OxyContin [®]	99	95	0.4	0.4	0.4	0.4	66.1	58.7
Tranquilizers	219	254	0.9	1.0	0.9	1.0	63.2	59.9
Stimulants ³	138	183	0.6	0.7	0.6	0.7	52.0	53.0
Methamphetamine ³	40	51	0.2	0.2	0.2	0.2	*	*
Sedatives	26 ^a	59	0.1 ^a	0.2	0.1 ^a	0.2	*	*
ILLCIT DRUGS OTHER THAN MARIJUANA^{2,3}	788	814	3.2	3.3	3.5	3.6	39.8	39.2
CIGARETTES	932	838	3.7	3.4	4.3	3.8	36.4	37.9
Daily Cigarette Use ⁵	209	165	0.8	0.7	0.9	0.7	--	--
SMOKELESS TOBACCO⁶	484	431	1.9	1.7	2.0	1.8	47.9	45.8
CIGARS	730	797	2.9	3.2	3.1	3.4	51.3	55.9
ALCOHOL	2,417	2,335	9.7	9.4	12.3	11.8	39.4	39.1

*Low precision; no estimate reported.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.7B Past Year Initiation of Substance Use among Persons Aged 18 to 25, Persons Aged 18 to 25 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 18 to 25: Numbers in Thousands and Percentages, 2013 and 2014

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS^{2,3}	1,106	1,131	3.2	3.2	6.9	7.1	8.9	9.0
Marijuana and Hashish	1,017	1,094	2.9	3.1	5.7	6.2	9.2	9.8
Cocaine	432	501	1.2	1.4	1.4	1.6	28.0	31.3
Crack	25	54	0.1	0.2	0.1	0.2	*	*
Heroin	66	75	0.2	0.2	0.2	0.2	27.1	27.9
Hallucinogens	611	608	1.8	1.7	2.1	2.0	26.1	27.6
LSD	312	371	0.9	1.1	1.0	1.1	44.1	46.3
PCP	13	24	0.0	0.1	0.0	0.1	*	*
Ecstasy	494	448	1.4	1.3	1.6	1.4	35.9	36.8
Inhalants	188	157	0.5	0.4	0.6	0.5	39.3	41.3
Nonmedical Use of Psychotherapeutics ^{3,4}	847	884	2.4	2.5	3.2	3.3	20.0	21.6
Pain Relievers	637	574	1.8	1.6	2.3	2.0	20.8	21.0
OxyContin [®]	144	139	0.4	0.4	0.4	0.4	30.5	32.5
Tranquilizers	453	440	1.3	1.3	1.5	1.4	28.9	30.4
Stimulants ³	345	356	1.0	1.0	1.1	1.1	27.3	27.5
Methamphetamine ³	93	67	0.3	0.2	0.3	0.2	34.9	23.6
Sedatives	35	64	0.1	0.2	0.1	0.2	*	*
ILLICIT DRUGS OTHER THAN MARIJUANA^{2,3}	952	1,044	2.7	3.0	4.0	4.3	16.3	18.1
CIGARETTES	1,031	1,181	3.0	3.4	6.6	7.2	7.5 ^a	9.0
Daily Cigarette Use ⁵	505	479	1.5	1.4	1.9	1.8	--	--
SMOKELESS TOBACCO⁶	514	468	1.5	1.3	1.8	1.6	15.9	15.1
CIGARS	1,334	1,311	3.8	3.8	5.9	5.7	18.5	18.8
ALCOHOL	2,056	2,225	5.9	6.4	26.7	27.7	7.7	8.3

*Low precision; no estimate reported.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.8B Past Year Initiation of Substance Use among Persons Aged 26 or Older, Persons Aged 26 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 26 or Older: Numbers in Thousands and Percentages, 2013 and 2014

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS^{2,3}	389	469	0.2	0.2	0.4	0.5	1.6	1.7
Marijuana and Hashish	210	271	0.1	0.1	0.2	0.2	1.1	1.3
Cocaine	75	148	0.0	0.1	0.0	0.1	3.0	5.3
Crack	23	44	0.0	0.0	0.0	0.0	*	7.0
Heroin	82	124	0.0	0.1	0.0	0.1	*	20.0
Hallucinogens	147	70	0.1	0.0	0.1	0.0	9.2	4.3
LSD	48	50	0.0	0.0	0.0	0.0	*	*
PCP	*	*	*	*	*	*	*	*
Ecstasy	75	104	0.0	0.1	0.0	0.1	7.6	11.0
Inhalants	121	84	0.1	0.0	0.1	0.0	*	*
Nonmedical Use of Psychotherapeutics ^{3,4}	558	585	0.3	0.3	0.3	0.4	5.8	6.4
Pain Relievers	394	362	0.2	0.2	0.2	0.2	5.7	5.6
OxyContin [®]	193	66	0.1	0.0	0.1	0.0	*	10.2
Tranquilizers	508	439	0.3	0.2	0.3	0.2	15.1	13.2
Stimulants ³	121	152	0.1	0.1	0.1	0.1	6.9	8.5
Methamphetamine ³	11 ^a	65	0.0 ^a	0.0	0.0 ^a	0.0	*	10.2
Sedatives	68	50	0.0	0.0	0.0	0.0	*	*
ILLCIT DRUGS OTHER THAN MARIJUANA^{2,3}	480	555	0.2	0.3	0.3	0.4	4.0	4.7
CIGARETTES	108	144	0.1	0.1	0.2	0.2	0.2	0.3
Daily Cigarette Use ⁵	99	113	0.0	0.1	0.1	0.1	--	--
SMOKELESS TOBACCO⁶	118	117	0.1	0.1	0.1	0.1	1.5	1.6
CIGARS	706	489	0.3	0.2	0.5	0.4	4.3	3.2
ALCOHOL	85	95	0.0	0.0	0.3	0.4	0.1	0.1

*Low precision; no estimate reported.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.9A Past Year Initiation of Substance Use among Persons Aged 12 or Older, by Gender: Numbers in Thousands, 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLICIT DRUGS^{1,2}	2,848	3,011	1,187 ^a	1,394	1,661	1,617
Marijuana and Hashish	2,427	2,568	1,072	1,226	1,355	1,342
Cocaine	601 ^a	766	344	442	256	324
Crack	58	109	33	56	25	53
Heroin	169	212	109	142	60	69
Hallucinogens	1,065	936	585	503	480	433
LSD	482	586	280	370	202	216
PCP	32	41	22	15	10	26
Ecstasy	751	676	398	395	354	281
Inhalants	563	512	332	275	231	237
Nonmedical Use of Psychotherapeutics ^{2,3}	2,006	2,111	827	893	1,179	1,218
Pain Relievers	1,539	1,425	700	653	839	773
OxyContin [®]	436	300	278	150	158	150
Tranquilizers	1,180	1,133	425	485	755	648
Stimulants ²	603	690	292	335	312	355
Methamphetamine ²	144	183	72	94	72	89
Sedatives	128	173	41	51	88	122
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	2,220	2,413	988	1,073	1,233	1,340
CIGARETTES	2,071	2,164	1,048	1,074	1,023	1,089
Daily Cigarette Use ⁴	813	756	460	392	354	365
SMOKELESS TOBACCO⁵	1,116	1,016	823	749	293	266
CIGARS	2,770	2,597	1,463	1,441	1,307	1,156
ALCOHOL	4,559	4,655	2,119	2,151	2,439	2,504

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.10A Past Year Initiation of Substance Use among Persons Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: Numbers in Thousands, 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLCIT DRUGS^{1,2}	1,542	1,550	733	717	809	833
Marijuana and Hashish	1,374	1,355	647	651	727	704
Cocaine	109	140	59	71	50	69
Crack	11	12	6	6	4	7
Heroin	21	17	10	9	*	*
Hallucinogens	374	303	230 ^a	162	144	140
LSD	140 ^a	201	87	108	53 ^a	93
PCP	19	17	12	6	7	12
Ecstasy	230 ^a	166	138	96	92	69
Inhalants	264	290	119	125	144	165
Nonmedical Use of Psychotherapeutics ^{2,3}	702	714	351	305	351	410
Pain Relievers	593	541	291	228	302	312
OxyContin [®]	110	102	51	56	60	46
Tranquilizers	255	288	128	132	127	156
Stimulants ²	161	210	86	96	75	114
Methamphetamine ²	48	59	24	29	24	31
Sedatives	32 ^a	65	9	25	23	40
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	896	894	458	401	439	492
CIGARETTES	1,046	955	541 ^a	433	505	522
Daily Cigarette Use ⁴	270	212	155 ^a	98	115	114
SMOKELESS TOBACCO⁵	535	477	434	362	101	115
CIGARS	876	938	555	560	320	378
ALCOHOL	2,693	2,615	1,247	1,146	1,446	1,469

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs

Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.11A Past Year Initiation of Substance Use among Persons Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: Numbers in Thousands, 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLCIT DRUGS^{1,2}	2,165	2,205	994	1,053	1,171	1,152
Marijuana and Hashish	1,944	2,009	899	995	1,045	1,013
Cocaine	351	339	210	175	141	164
Crack	25	40	11	26	14	14
Heroin	43	40	21	24	22	16
Hallucinogens	680	651	439	362	241	288
LSD	322	374	218	213	104 ^a	161
PCP	26	34	19	10	7 ^a	24
Ecstasy	458	384	278	222	180	162
Inhalants	364	355	193	166	171	190
Nonmedical Use of Psychotherapeutics ^{2,3}	1,135	1,167	573	519	562	648
Pain Relievers	879	848	456	386	423	462
OxyContin [®]	180	167	99	91	81	75
Tranquilizers	466	499	247	255	220	244
Stimulants ²	383	393	210	183	174	210
Methamphetamine ²	105	95	51	48	54	47
Sedatives	52 ^a	98	14 ^a	38	37	60
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	1,401	1,468	728	670	673 ^a	798
CIGARETTES	1,746	1,711	906	866	840	845
Daily Cigarette Use ⁴	617 ^a	500	369 ^a	265	248	236
SMOKELESS TOBACCO⁵	831	805	671	630	159	175
CIGARS	1,695	1,761	1,038	1,019	657	742
ALCOHOL	3,807	3,774	1,788	1,737	2,019	2,038

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs

Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.12A Past Year Initiation of Substance Use among Persons Aged 12 to 49, by Gender: Numbers in Thousands, 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLICIT DRUGS^{1,2}	2,784	2,940	1,155 ^a	1,370	1,629	1,570
Marijuana and Hashish	2,394	2,522	1,058	1,220	1,337	1,303
Cocaine	601 ^a	752	344	432	256	321
Crack	50 ^a	109	25	56	25	53
Heroin	151	198	91	129	60	69
Hallucinogens	1,032	936	585	503	447	433
LSD	482	586	280	370	202	216
PCP	32	41	22	15	10	26
Ecstasy	751	676	398	395	354	281
Inhalants	531	490	311	254	220	237
Nonmedical Use of Psychotherapeutics ^{2,3}	1,906	2,034	795	868	1,111	1,166
Pain Relievers	1,444	1,368	659	624	785	744
OxyContin [®]	340	300	189	150	151	150
Tranquilizers	1,107	1,043	425	466	682	578
Stimulants ²	603	682	292	326	312	355
Methamphetamine ²	144	183	72	94	72	89
Sedatives	122	160	41	51	81	109
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	2,099	2,339	970	1,048	1,128	1,290
CIGARETTES	2,059	2,164	1,036	1,074	1,023	1,089
Daily Cigarette Use ⁴	803	756	460	392	343	365
SMOKELESS TOBACCO⁵	1,096	1,016	823	749	273	266
CIGARS	2,593	2,486	1,432	1,396	1,161	1,089
ALCOHOL	4,551	4,617	2,119	2,151	2,432	2,466

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.13B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older, by Gender: 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLICIT DRUGS^{1,2}	19.8	20.0	18.4	19.5	20.9	20.5
Marijuana and Hashish	18.6	19.4	18.1	18.9	18.9	19.9
Cocaine	20.4 ^b	22.4	20.3 ^a	22.9	20.5	21.6
Crack	28.2	26.4	31.6	23.3	23.6	29.6
Heroin	27.4	29.6	29.3	31.0	24.0	26.7
Hallucinogens	21.6	19.3	19.0	19.3	24.7	19.3
LSD	19.7	19.7	18.9 ^b	20.3	20.9 ^b	18.5
PCP	17.1	17.7	17.1	*	16.9	17.1
Ecstasy	20.5	21.0	19.2 ^b	21.4	22.0	20.4
Inhalants	21.0	20.1	21.7	22.7	20.1	17.1
Nonmedical Use of Psychotherapeutics ^{2,3}	24.0	23.1	21.1	22.1	26.1	23.8
Pain Relievers	23.8	22.7	22.0	22.8	25.3	22.5
OxyContin [®]	31.3 ^a	21.2	34.0 ^a	19.8	26.6	22.5
Tranquilizers	27.2	26.1	22.3	23.1	30.0	28.3
Stimulants ²	21.6	22.0	20.9	23.0	22.2	21.0
Methamphetamine ²	18.9 ^a	22.0	19.2	23.0	18.5	21.0
Sedatives	26.3	23.9	23.4 ^a	18.4	27.6	26.2
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	22.7	21.7	19.4	20.9	25.4	22.4
CIGARETTES	18.0	18.6	17.9	18.7	18.2	18.5
Daily Cigarette Use ⁴	20.4	20.6	19.0	20.1	22.2	21.1
SMOKELESS TOBACCO⁵	19.0	19.0	18.1	18.3	21.7	20.7
CIGARS	23.6	21.8	21.2	21.0	26.3	22.8
ALCOHOL	17.3	17.6	17.2	17.4	17.5	17.7

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.14B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLCIT DRUGS^{1,2}	15.0	14.9	15.0	15.0	15.0	14.9
Marijuana and Hashish	15.2	15.2	15.2	15.2	15.3	15.1
Cocaine	15.9	15.9	15.8	16.2	16.1 ^a	15.5
Crack	15.3	16.0	*	*	*	*
Heroin	15.0	*	*	*	15.2	*
Hallucinogens	15.8	15.6	15.9	15.9	15.7	15.3
LSD	16.1	15.9	16.2	16.1	15.9	15.6
PCP	15.1	15.3	15.4	*	*	15.1
Ecstasy	15.9	15.9	16.0	16.1	15.8	15.5
Inhalants	14.1	14.2	14.1	14.5	14.1	14.0
Nonmedical Use of Psychotherapeutics ^{2,3}	15.3	15.2	15.3	15.3	15.2	15.1
Pain Relievers	15.3	15.2	15.3	15.4	15.3	15.1
OxyContin [®]	15.4	15.1	15.6	15.5	15.2	14.7
Tranquilizers	15.4	15.4	15.8	15.7	15.1	15.2
Stimulants ²	15.4	15.2	15.7	15.1	15.1	15.2
Methamphetamine ²	15.5	15.1	15.5	15.2	15.6	15.1
Sedatives	15.1	14.7	13.5	14.5	15.7	14.8
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	15.1	14.9	15.2	15.0	15.0	14.9
CIGARETTES	15.3	15.4	15.3	15.5	15.2	15.3
Daily Cigarette Use ⁴	15.8	15.7	15.9	16.0	15.6	15.5
SMOKELESS TOBACCO⁵	15.2	15.4	15.2	15.4	14.9	15.2
CIGARS	15.6	15.6	15.7	15.6	15.5	15.4
ALCOHOL	15.2	15.2	15.2	15.3	15.2	15.1

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs

Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.15B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLCIT DRUGS^{1,2}	16.1	16.0	15.9	16.2	16.2	15.9
Marijuana and Hashish	16.2	16.3	16.1 ^a	16.4	16.3	16.2
Cocaine	18.0	17.8	18.0	17.9	18.0	17.6
Crack	17.3	18.0	17.1	18.3	17.4	*
Heroin	17.0	17.7	17.1	17.9	16.9	*
Hallucinogens	17.2	17.3	17.3	17.5	17.0	17.1
LSD	17.6	17.3	17.8	17.6	17.2	17.0
PCP	15.9	16.9	16.4	*	*	16.8
Ecstasy	17.4	17.6	17.4	17.7	17.3	17.5
Inhalants	15.4	15.0	15.9	15.5	14.8	14.6
Nonmedical Use of Psychotherapeutics ^{2,3}	16.7	16.6	16.8	16.7	16.6	16.5
Pain Relievers	16.5	16.5	16.7	16.8	16.4	16.3
OxyContin [®]	16.9	16.5	17.4	16.7	16.2	16.3
Tranquilizers	17.0	16.9	17.3	17.3	16.7	16.5
Stimulants ²	17.5 ^a	16.9	17.7 ^a	16.7	17.2	16.9
Methamphetamine ²	17.2	16.5	17.1	16.7	17.3	16.3
Sedatives	16.5	16.1	15.8	16.1	16.8	16.1
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	16.5	16.4	16.6	16.5	16.4	16.4
CIGARETTES	16.6 ^a	16.8	16.6 ^b	17.0	16.6	16.6
Daily Cigarette Use ⁴	17.4	17.4	17.5	17.6	17.2	17.2
SMOKELESS TOBACCO⁵	16.4 ^a	16.7	16.4 ^a	16.8	16.3	16.5
CIGARS	17.1	17.0	17.0	16.9	17.2	17.1
ALCOHOL	16.2	16.2	16.2	16.4	16.1	16.1

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs

Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.16B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49, by Gender: 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLICIT DRUGS^{1,2}	19.0	19.0	17.3 ^b	18.9	20.2	19.1
Marijuana and Hashish	18.0	18.5	17.4 ^a	18.6	18.5	18.5
Cocaine	20.4 ^a	21.8	20.3 ^a	22.1	20.5	21.3
Crack	24.4	26.4	25.1	23.3	23.6	29.6
Heroin	24.5	28.0	24.8	28.7	24.0	26.7
Hallucinogens	19.9	19.3	19.0	19.3	21.1	19.3
LSD	19.7	19.7	18.9 ^b	20.3	20.9 ^b	18.5
PCP	17.1	17.7	17.1	*	16.9	17.1
Ecstasy	20.5	21.0	19.2 ^b	21.4	22.0	20.4
Inhalants	19.2	18.2	19.6	19.3	18.5	17.1
Nonmedical Use of Psychotherapeutics ^{2,3}	22.4	21.8	19.7	21.3	24.4 ^a	22.2
Pain Relievers	21.7	21.2	20.0	21.3	23.0	21.2
OxyContin [®]	23.6	21.2	22.1	19.8	25.4	22.5
Tranquilizers	25.4	23.4	22.3	21.9	27.3	24.6
Stimulants ²	21.6	21.6	20.9	22.1	22.2	21.0
Methamphetamine ²	18.9 ^a	22.0	19.2	23.0	18.5	21.0
Sedatives	25.0	21.4	23.4 ^a	18.4	25.8	22.8
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	20.6	20.6	18.8 ^a	20.2	22.1	21.0
CIGARETTES	17.8 ^a	18.6	17.5 ^b	18.7	18.2	18.5
Daily Cigarette Use ⁴	19.8	20.6	19.0	20.1	21.0	21.1
SMOKELESS TOBACCO⁵	18.4	19.0	18.1	18.3	19.6	20.7
CIGARS	21.6	20.4	20.3	20.1	23.2 ^b	20.8
ALCOHOL	17.3	17.3	17.2	17.4	17.4	17.2

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.1D Past Year Initiation of Illicit Drug Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Illicit Drug Use, and Past Year Illicit Drug Users Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	102	107	0.04	0.04	0.08	0.08	0.25	0.24
AGE								
12-25	75	84	0.13	0.14	0.20	0.23	0.43	0.46
12-17	50	58	0.20	0.23	0.25	0.28	0.96	1.08
12-13	22	27	0.27	0.34	0.29	0.36	3.24	3.46
14-15	32	34	0.37	0.39	0.43	0.46	1.86	1.92
16-17	34	40	0.40	0.45	0.56	0.61	1.18	1.39
18-25	57	63	0.16	0.18	0.34	0.39	0.45	0.47
18-20	51	55	0.36	0.39	0.65	0.72	0.87	0.95
21-25	31	36	0.14	0.16	0.33	0.40	0.40	0.46
26 or Older	63	64	0.03	0.03	0.06	0.06	0.25	0.24
GENDER								
Male	59	78	0.05	0.06	0.10	0.13	0.25	0.31
Female	82	74	0.06	0.05	0.11	0.09	0.47	0.40
GENDER/AGE								
Male 12-17	35	40	0.27	0.31	0.34	0.38	1.29	1.56
Female 12-17	37	42	0.30	0.35	0.36	0.41	1.45	1.60

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.1P Past Year Initiation of Illicit Drug Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Illicit Drug Use, and Past Year Illicit Drug Users Aged 12 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	N/A		0.3699		0.2745		0.9332	
AGE								
12-25	N/A		0.4926		0.3981		0.6168	
12-17	N/A		0.4479		0.4786		0.5073	
12-13	N/A		0.4592		0.4422		0.5110	
14-15	N/A		0.6139		0.6516		0.7822	
16-17	N/A		0.9008		0.9692		0.4786	
18-25	N/A		0.8066		0.6237		0.8810	
18-20	N/A		0.8930		0.9293		0.9608	
21-25	N/A		0.4390		0.3157		0.5592	
26 or Older	N/A		0.4062		0.3773		0.6464	
GENDER								
Male	N/A		0.0481		0.0265		0.2488	
Female	N/A		0.5786		0.6083		0.3148	
GENDER/AGE								
Male 12-17	N/A		0.9734		0.8840		0.3183	
Female 12-17	N/A		0.3161		0.2539		0.7578	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.2D Past Year Initiation of Marijuana Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Marijuana Use, and Past Year Marijuana Users Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	86	95	0.03	0.04	0.06	0.06	0.27	0.27
AGE								
12-25	71	81	0.12	0.13	0.18	0.20	0.47	0.53
12-17	46	52	0.18	0.21	0.21	0.24	1.09	1.28
12-13	16	20	0.20	0.25	0.20	0.25	4.48	4.64
14-15	29	30	0.34	0.35	0.37	0.38	2.15	2.35
16-17	34	40	0.40	0.45	0.53	0.58	1.33	1.57
18-25	54	62	0.16	0.18	0.30	0.35	0.48	0.53
18-20	48	57	0.33	0.40	0.57	0.67	0.90	1.06
21-25	31	33	0.14	0.15	0.29	0.32	0.45	0.48
26 or Older	44	45	0.02	0.02	0.04	0.04	0.24	0.22
GENDER								
Male	53	71	0.04	0.06	0.08	0.11	0.28	0.33
Female	67	63	0.05	0.05	0.08	0.07	0.54	0.45
GENDER/AGE								
Male 12-17	32	37	0.25	0.29	0.29	0.33	1.52	1.81
Female 12-17	35	37	0.29	0.30	0.32	0.34	1.71	1.87

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.2P Past Year Initiation of Marijuana Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Marijuana Use, and Past Year Marijuana Users Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	N/A		0.3679		0.2940		0.8894	
AGE								
12-25	N/A		0.4803		0.4107		0.4941	
12-17	N/A		0.9544		0.9564		0.5428	
12-13	N/A		0.8214		0.8359		0.3277	
14-15	N/A		0.7551		0.8066		0.5373	
16-17	N/A		0.7054		0.7729		0.9232	
18-25	N/A		0.3737		0.2969		0.4263	
18-20	N/A		0.3560		0.4205		0.3140	
21-25	N/A		0.7233		0.5756		0.8194	
26 or Older	N/A		0.3621		0.3458		0.5786	
GENDER								
Male	N/A		0.1062		0.0685		0.4243	
Female	N/A		0.7665		0.7763		0.3929	
GENDER/AGE								
Male 12-17	N/A		0.9355		0.8842		0.4265	
Female 12-17	N/A		0.8792		0.8237		0.8916	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.3D Past Year Initiation of Cigarette Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Cigarette Use, and Past Year Cigarette Users Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	81	90	0.03	0.03	0.08	0.09	0.12	0.14
AGE								
12-25	72	84	0.12	0.14	0.19	0.21	0.42	0.53
12-17	41	44	0.16	0.18	0.19	0.20	1.27	1.56
12-13	12	14	0.16	0.17	0.16	0.18	5.14	*
14-15	24	21	0.27	0.25	0.31	0.27	2.33	2.74
16-17	32	36	0.37	0.41	0.47	0.50	1.61	1.95
18-25	57	72	0.17	0.21	0.36	0.43	0.41	0.54
18-20	51	62	0.36	0.45	0.63	0.74	0.93	1.22
21-25	26	38	0.12	0.17	0.32	0.42	0.29	0.42
26 or Older	32	29	0.02	0.01	0.05	0.04	0.06	0.06
GENDER								
Male	56	68	0.04	0.05	0.13	0.15	0.16	0.19
Female	55	57	0.04	0.04	0.09	0.10	0.19	0.19
GENDER/AGE								
Male 12-17	30	30	0.23	0.24	0.27	0.27	1.81	2.09
Female 12-17	28	32	0.23	0.26	0.26	0.29	1.78	2.27

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.3P Past Year Initiation of Cigarette Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Cigarette Use, and Past Year Cigarette Users Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	N/A		0.5617		0.8214		0.3930	
AGE								
12-25	N/A		0.6388		0.9922		0.1041	
12-17	N/A		0.1219		0.0826		0.4429	
12-13	N/A		0.2604		0.2601		*	
14-15	N/A		0.0734		0.0493		0.5943	
16-17	N/A		0.6599		0.4848		0.4193	
18-25	N/A		0.1155		0.3030		0.0283	
18-20	N/A		0.5429		0.8776		0.1680	
21-25	N/A		0.0206		0.0471		0.0093	
26 or Older	N/A		0.4287		0.4621		0.4254	
GENDER								
Male	N/A		0.8628		0.9594		0.7129	
Female	N/A		0.4887		0.6837		0.3752	
GENDER/AGE								
Male 12-17	N/A		0.0335		0.0247		0.2822	
Female 12-17	N/A		0.9392		0.8204		0.0246	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.4D Past Year Initiation of Alcohol Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Alcohol Use, and Past Year Alcohol Users Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	113	127	0.04	0.05	0.23	0.24	0.07	0.07
AGE								
12-25	103	112	0.17	0.19	0.37	0.38	0.31	0.35
12-17	67	67	0.27	0.27	0.34	0.33	0.88	0.94
12-13	20	25	0.25	0.31	0.26	0.33	3.23	3.77
14-15	43	43	0.48	0.49	0.57	0.58	1.57	1.78
16-17	51	51	0.55	0.56	0.86	0.84	1.07	1.15
18-25	76	86	0.22	0.25	0.87	0.91	0.28	0.33
18-20	68	75	0.44	0.50	1.08	1.17	0.67	0.77
21-25	48	56	0.20	0.25	1.43	1.58	0.24	0.30
26 or Older	26	37	0.01	0.02	0.10	0.15	0.02	0.03
GENDER								
Male	76	84	0.06	0.07	0.37	0.37	0.09	0.10
Female	83	94	0.06	0.07	0.29	0.31	0.10	0.11
GENDER/AGE								
Male 12-17	45	45	0.35	0.35	0.45	0.44	1.24	1.34
Female 12-17	50	52	0.41	0.43	0.51	0.53	1.26	1.33

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.4P Past Year Initiation of Alcohol Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Alcohol Use, and Past Year Alcohol Users Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	N/A		0.7716		0.2602		0.8669	
AGE								
12-25	N/A		0.6107		1.0000		0.4810	
12-17	N/A		0.3988		0.2618		0.7856	
12-13	N/A		0.6600		0.7243		0.2488	
14-15	N/A		0.7253		0.6070		0.4415	
16-17	N/A		0.2136		0.1044		0.2734	
18-25	N/A		0.1639		0.4408		0.1460	
18-20	N/A		0.4558		0.9055		0.2767	
21-25	N/A		0.1404		0.2772		0.1478	
26 or Older	N/A		0.8555		0.7364		0.8654	
GENDER								
Male	N/A		0.9290		0.7235		0.8944	
Female	N/A		0.7556		0.2527		0.9313	
GENDER/AGE								
Male 12-17	N/A		0.3229		0.1801		0.9224	
Female 12-17	N/A		0.8101		0.7991		0.5856	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.5D Past Year Initiation of Substance Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 or Older: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS^{2,3}	102	107	0.04	0.04	0.08	0.08	0.25	0.24
Marijuana and Hashish	86	95	0.03	0.04	0.06	0.06	0.27	0.27
Cocaine	47	57	0.02	0.02	0.02	0.03	1.13	1.22
Crack	13	24	0.01	0.01	0.01	0.01	2.23	3.10
Heroin	36	35	0.01	0.01	0.01	0.01	4.47	3.31
Hallucinogens	72	58	0.03	0.02	0.03	0.03	1.56	1.28
LSD	40	48	0.02	0.02	0.02	0.02	3.49	2.73
PCP	7	10	0.00	0.00	0.00	0.00	*	*
Ecstasy	50	52	0.02	0.02	0.02	0.02	2.11	1.98
Inhalants	51	49	0.02	0.02	0.02	0.02	3.04	2.65
Nonmedical Use of Psychotherapeutics ^{3,4}	100	99	0.04	0.04	0.05	0.05	0.65	0.64
Pain Relievers	82	76	0.03	0.03	0.04	0.03	0.75	0.71
OxyContin [®]	75	31	0.03	0.01	0.03	0.01	4.10	2.40
Tranquilizers	89	73	0.03	0.03	0.04	0.03	1.64	1.27
Stimulants ³	48	58	0.02	0.02	0.02	0.02	1.53	1.55
Methamphetamine ³	21	31	0.01	0.01	0.01	0.01	2.36	3.00
Sedatives	27	25	0.01	0.01	0.01	0.01	4.23	3.27
ILLICIT DRUGS OTHER THAN MARIJUANA^{2,3}	108	103	0.04	0.04	0.06	0.06	0.54	0.52
CIGARETTES	81	90	0.03	0.03	0.08	0.09	0.12	0.14
Daily Cigarette Use ⁵	52	51	0.02	0.02	0.03	0.03	--	--
SMOKELESS TOBACCO⁶	65	59	0.02	0.02	0.03	0.03	0.57	0.51
CIGARS	144	104	0.06	0.04	0.08	0.06	0.55	0.42
ALCOHOL	113	127	0.04	0.05	0.23	0.24	0.07	0.07

*Low precision; no estimate reported.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.5P Past Year Initiation of Substance Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS^{2,3}	N/A		0.3699		0.2745		0.9332	
Marijuana and Hashish	N/A		0.3679		0.2940		0.8894	
Cocaine	N/A		0.0317		0.0289		0.1380	
Crack	N/A		0.0605		0.0600		0.1897	
Heroin	N/A		0.4241		0.4248		0.7552	
Hallucinogens	N/A		0.1303		0.1279		0.3156	
LSD	N/A		0.1129		0.1142		0.6361	
PCP	N/A		0.4850		0.4861		*	
Ecstasy	N/A		0.2557		0.2500		0.9589	
Inhalants	N/A		0.4202		0.4216		0.2085	
Nonmedical Use of Psychotherapeutics ^{3,4}	N/A		0.5508		0.5291		0.2127	
Pain Relievers	N/A		0.2508		0.2598		0.9270	
OxyContin [®]	N/A		0.0868		0.0872		0.2093	
Tranquilizers	N/A		0.6086		0.6428		0.7649	
Stimulants ³	N/A		0.2824		0.2752		0.4146	
Methamphetamine ³	N/A		0.3114		0.3079		0.3684	
Sedatives	N/A		0.2476		0.2462		0.6767	
ILLICIT DRUGS OTHER THAN MARIJUANA^{2,3}	N/A		0.2588		0.2250		0.1536	
CIGARETTES	N/A		0.5617		0.8214		0.3930	
Daily Cigarette Use ⁵	N/A		0.3732		0.3258		--	
SMOKELESS TOBACCO⁶	N/A		0.2074		0.1852		0.4243	
CIGARS	N/A		0.2601		0.2410		0.8104	
ALCOHOL	N/A		0.7716		0.2602		0.8669	

*Low precision; no estimate reported.

N/A: Not applicable.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.6D Past Year Initiation of Substance Use among Persons Aged 12 to 17, Persons Aged 12 to 17 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 to 17: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS^{2,3}	50	58	0.20	0.23	0.25	0.28	0.96	1.08
Marijuana and Hashish	46	52	0.18	0.21	0.21	0.24	1.09	1.28
Cocaine	13	20	0.05	0.08	0.05	0.08	4.83	*
Crack	4	5	0.01	0.02	0.01	0.02	*	*
Heroin	6	7	0.02	0.03	0.02	0.03	*	*
Hallucinogens	23	25	0.09	0.10	0.09	0.10	2.80	3.72
LSD	14	22	0.06	0.09	0.06	0.09	3.70	4.48
PCP	6	6	0.02	0.02	0.02	0.02	*	*
Ecstasy	18	18	0.07	0.07	0.07	0.07	3.81	5.60
Inhalants	22	28	0.09	0.11	0.09	0.12	3.13	3.54
Nonmedical Use of Psychotherapeutics ^{3,4}	34	40	0.13	0.16	0.14	0.17	1.84	1.96
Pain Relievers	31	34	0.12	0.14	0.13	0.15	2.15	2.20
OxyContin [®]	13	15	0.05	0.06	0.05	0.06	5.25	5.95
Tranquilizers	19	25	0.08	0.10	0.08	0.10	3.41	3.81
Stimulants ³	16	23	0.07	0.09	0.07	0.10	3.90	4.31
Methamphetamine ³	8	13	0.03	0.05	0.03	0.05	*	*
Sedatives	7	12	0.03	0.05	0.03	0.05	*	*
ILLICIT DRUGS OTHER THAN MARIJUANA^{2,3}	39	46	0.16	0.19	0.17	0.21	1.53	1.72
CIGARETTES	41	44	0.16	0.18	0.19	0.20	1.27	1.56
Daily Cigarette Use ⁵	19	19	0.08	0.08	0.08	0.08	--	--
SMOKELESS TOBACCO⁶	34	30	0.14	0.12	0.14	0.13	2.26	2.29
CIGARS	36	41	0.14	0.17	0.15	0.17	1.84	1.85
ALCOHOL	67	67	0.27	0.27	0.34	0.33	0.88	0.94

*Low precision; no estimate reported.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.6P Past Year Initiation of Substance Use among Persons Aged 12 to 17, Persons Aged 12 to 17 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS^{2,3}	N/A		0.4479		0.4786		0.5073	
Marijuana and Hashish	N/A		0.9544		0.9564		0.5428	
Cocaine	N/A		0.3438		0.3451		*	
Crack	N/A		0.8464		0.8458		*	
Heroin	N/A		0.4007		0.4009		*	
Hallucinogens	N/A		0.1579		0.1535		0.5965	
LSD	N/A		0.0970		0.0959		0.4663	
PCP	N/A		0.8243		0.8254		*	
Ecstasy	N/A		0.0236		0.0233		0.1494	
Inhalants	N/A		0.6196		0.6276		1.0000	
Nonmedical Use of Psychotherapeutics ^{3,4}	N/A		0.4359		0.4179		0.8994	
Pain Relievers	N/A		0.6845		0.6902		0.5283	
OxyContin [®]	N/A		0.8350		0.8323		0.3514	
Tranquilizers	N/A		0.2661		0.2588		0.5078	
Stimulants ³	N/A		0.1150		0.1134		0.8686	
Methamphetamine ³	N/A		0.4614		0.4584		*	
Sedatives	N/A		0.0157		0.0154		*	
ILLICIT DRUGS OTHER THAN MARIJUANA^{2,3}	N/A		0.6605		0.6520		0.7916	
CIGARETTES	N/A		0.1219		0.0826		0.4429	
Daily Cigarette Use ⁵	N/A		0.1008		0.1009		--	
SMOKELESS TOBACCO⁶	N/A		0.2490		0.2445		0.5270	
CIGARS	N/A		0.2174		0.2531		0.0773	
ALCOHOL	N/A		0.3988		0.2618		0.7856	

*Low precision; no estimate reported.

N/A: Not applicable.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.7D Past Year Initiation of Substance Use among Persons Aged 18 to 25, Persons Aged 18 to 25 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 18 to 25: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS^{2,3}	57	63	0.16	0.18	0.34	0.39	0.45	0.47
Marijuana and Hashish	54	62	0.16	0.18	0.30	0.35	0.48	0.53
Cocaine	37	40	0.11	0.11	0.12	0.13	1.97	2.05
Crack	6	14	0.02	0.04	0.02	0.04	*	*
Heroin	13	15	0.04	0.04	0.04	0.04	4.60	4.47
Hallucinogens	44	47	0.13	0.14	0.15	0.16	1.65	1.87
LSD	31	37	0.09	0.10	0.10	0.11	3.40	3.22
PCP	5	8	0.01	0.02	0.01	0.02	*	*
Ecstasy	39	42	0.11	0.12	0.12	0.13	2.22	2.81
Inhalants	24	22	0.07	0.06	0.07	0.07	4.12	4.68
Nonmedical Use of Psychotherapeutics ^{3,4}	53	55	0.15	0.16	0.20	0.21	1.13	1.20
Pain Relievers	48	45	0.14	0.13	0.17	0.16	1.42	1.48
OxyContin [®]	23	20	0.07	0.06	0.07	0.06	3.90	3.98
Tranquilizers	36	36	0.10	0.10	0.12	0.12	1.92	2.10
Stimulants ³	31	38	0.09	0.11	0.10	0.12	2.17	2.42
Methamphetamine ³	16	14	0.05	0.04	0.05	0.04	4.95	4.34
Sedatives	9	13	0.03	0.04	0.03	0.04	*	*
ILLICIT DRUGS OTHER THAN MARIJUANA^{2,3}	57	60	0.16	0.17	0.24	0.25	0.92	0.97
CIGARETTES	57	72	0.17	0.21	0.36	0.43	0.41	0.54
Daily Cigarette Use ⁵	36	40	0.10	0.12	0.14	0.15	--	--
SMOKELESS TOBACCO⁶	40	44	0.12	0.13	0.14	0.15	1.14	1.29
CIGARS	61	67	0.18	0.19	0.27	0.29	0.79	0.87
ALCOHOL	76	86	0.22	0.25	0.87	0.91	0.28	0.33

*Low precision; no estimate reported.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.7P Past Year Initiation of Substance Use among Persons Aged 18 to 25, Persons Aged 18 to 25 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 18 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS^{2,3}	N/A		0.8066		0.6237		0.8810	
Marijuana and Hashish	N/A		0.3737		0.2969		0.4263	
Cocaine	N/A		0.2130		0.2397		0.2468	
Crack	N/A		0.0598		0.0602		*	
Heroin	N/A		0.6850		0.6812		0.9021	
Hallucinogens	N/A		0.9327		0.8424		0.5629	
LSD	N/A		0.2396		0.2279		0.6330	
PCP	N/A		0.2419		0.2426		*	
Ecstasy	N/A		0.3917		0.3615		0.8035	
Inhalants	N/A		0.3363		0.3247		0.7390	
Nonmedical Use of Psychotherapeutics ^{3,4}	N/A		0.6601		0.6943		0.3464	
Pain Relievers	N/A		0.3189		0.2935		0.9425	
OxyContin [®]	N/A		0.8458		0.8341		0.7140	
Tranquilizers	N/A		0.7805		0.8186		0.5807	
Stimulants ³	N/A		0.8423		0.8700		0.9595	
Methamphetamine ³	N/A		0.2239		0.2251		0.0879	
Sedatives	N/A		0.0672		0.0674		*	
ILLICIT DRUGS OTHER THAN MARIJUANA^{2,3}	N/A		0.2924		0.3171		0.1801	
CIGARETTES	N/A		0.1155		0.3030		0.0283	
Daily Cigarette Use ⁵	N/A		0.5995		0.5204		--	
SMOKELESS TOBACCO⁶	N/A		0.4192		0.3755		0.6555	
CIGARS	N/A		0.7502		0.6356		0.8003	
ALCOHOL	N/A		0.1639		0.4408		0.1460	

*Low precision; no estimate reported.

N/A: Not applicable.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.8D Past Year Initiation of Substance Use among Persons Aged 26 or Older, Persons Aged 26 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 26 or Older: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS^{2,3}	63	64	0.03	0.03	0.06	0.06	0.25	0.24
Marijuana and Hashish	44	45	0.02	0.02	0.04	0.04	0.24	0.22
Cocaine	29	35	0.01	0.02	0.02	0.02	1.15	1.22
Crack	11	18	0.01	0.01	0.01	0.01	*	2.81
Heroin	32	31	0.02	0.02	0.02	0.02	*	4.45
Hallucinogens	49	19	0.02	0.01	0.03	0.01	2.91	1.12
LSD	25	19	0.01	0.01	0.01	0.01	*	*
PCP	*	*	*	*	*	*	*	*
Ecstasy	26	23	0.01	0.01	0.01	0.01	2.68	2.28
Inhalants	39	34	0.02	0.02	0.02	0.02	*	*
Nonmedical Use of Psychotherapeutics ^{3,4}	78	69	0.04	0.03	0.05	0.04	0.81	0.74
Pain Relievers	61	53	0.03	0.03	0.03	0.03	0.89	0.80
OxyContin [®]	68	17	0.03	0.01	0.03	0.01	*	2.68
Tranquilizers	80	59	0.04	0.03	0.04	0.03	2.25	1.64
Stimulants ³	31	35	0.02	0.02	0.02	0.02	1.79	1.90
Methamphetamine ³	11	22	0.01	0.01	0.01	0.01	*	3.40
Sedatives	25	18	0.01	0.01	0.01	0.01	*	*
ILLICIT DRUGS OTHER THAN MARIJUANA^{2,3}	78	67	0.04	0.03	0.06	0.05	0.64	0.56
CIGARETTES	32	29	0.02	0.01	0.05	0.04	0.06	0.06
Daily Cigarette Use ⁵	31	25	0.02	0.01	0.03	0.02	--	--
SMOKELESS TOBACCO⁶	31	26	0.02	0.01	0.02	0.02	0.41	0.35
CIGARS	126	62	0.06	0.03	0.10	0.05	0.75	0.40
ALCOHOL	26	37	0.01	0.02	0.10	0.15	0.02	0.03

*Low precision; no estimate reported.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.8P Past Year Initiation of Substance Use among Persons Aged 26 or Older, Persons Aged 26 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 26 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	NUMBER OF PAST YEAR INITIATES (1,000s)		PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION		PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ¹		PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS^{2,3}	N/A		0.4062		0.3773		0.6464	
Marijuana and Hashish	N/A		0.3621		0.3458		0.5786	
Cocaine	N/A		0.1165		0.1128		0.1674	
Crack	N/A		0.3166		0.3153		*	
Heroin	N/A		0.3772		0.3781		*	
Hallucinogens	N/A		0.1310		0.1314		0.1188	
LSD	N/A		0.9504		0.9525		*	
PCP	N/A		*		*		*	
Ecstasy	N/A		0.4234		0.4246		0.3402	
Inhalants	N/A		0.4543		0.4557		*	
Nonmedical Use of Psychotherapeutics ^{3,4}	N/A		0.8443		0.8308		0.5868	
Pain Relievers	N/A		0.6533		0.6625		0.9314	
OxyContin [®]	N/A		0.0702		0.0704		*	
Tranquilizers	N/A		0.4474		0.4613		0.4762	
Stimulants ³	N/A		0.5338		0.5279		0.5343	
Methamphetamine ³	N/A		0.0291		0.0290		*	
Sedatives	N/A		0.5503		0.5515		*	
ILLICIT DRUGS OTHER THAN MARIJUANA^{2,3}	N/A		0.5034		0.4785		0.4128	
CIGARETTES	N/A		0.4287		0.4621		0.4254	
Daily Cigarette Use ⁵	N/A		0.7594		0.7817		--	
SMOKELESS TOBACCO⁶	N/A		0.9516		0.9387		0.9906	
CIGARS	N/A		0.1123		0.1095		0.1800	
ALCOHOL	N/A		0.8555		0.7364		0.8654	

*Low precision; no estimate reported.

N/A: Not applicable.

-- Not available.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁶ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.9C Past Year Initiation of Substance Use among Persons Aged 12 or Older, by Gender: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLICIT DRUGS^{1,2}	102	107	59	78	82	74
Marijuana and Hashish	86	95	53	71	67	63
Cocaine	47	57	39	45	29	34
Crack	13	24	11	16	7	17
Heroin	36	35	33	28	13	21
Hallucinogens	72	58	43	42	55	36
LSD	40	48	28	39	32	25
PCP	7	10	6	5	4	8
Ecstasy	50	52	33	40	38	30
Inhalants	51	49	42	39	29	27
Nonmedical Use of Psychotherapeutics ^{2,3}	100	99	54	64	83	75
Pain Relievers	82	76	51	55	63	52
OxyContin [®]	75	31	68	22	25	21
Tranquilizers	89	73	40	46	78	56
Stimulants ²	48	58	33	40	35	39
Methamphetamine ²	21	31	16	20	14	21
Sedatives	27	25	14	11	23	23
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	108	103	60	66	84	77
CIGARETTES	81	90	56	68	55	57
Daily Cigarette Use ⁴	52	51	36	40	36	34
SMOKELESS TOBACCO⁵	65	59	50	50	36	28
CIGARS	144	104	89	76	106	66
ALCOHOL	113	127	76	84	83	94

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.10C Past Year Initiation of Substance Use among Persons Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLICIT DRUGS^{1,2}	59	65	41	46	42	46
Marijuana and Hashish	53	60	36	44	40	40
Cocaine	15	22	11	16	10	14
Crack	4	5	2	3	3	4
Heroin	6	7	4	5	*	*
Hallucinogens	29	28	25	19	15	19
LSD	16	24	13	16	10	17
PCP	6	6	5	3	3	5
Ecstasy	20	22	16	16	13	13
Inhalants	22	29	16	19	16	21
Nonmedical Use of Psychotherapeutics ^{2,3}	38	45	28	30	25	34
Pain Relievers	35	38	26	25	24	28
OxyContin [®]	13	15	8	12	10	9
Tranquilizers	22	27	17	18	13	20
Stimulants ²	18	26	14	18	11	18
Methamphetamine ²	9	14	7	10	7	10
Sedatives	7	12	4	8	6	10
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	44	50	34	32	27	37
CIGARETTES	46	49	34	33	31	35
Daily Cigarette Use ⁴	22	22	17	16	13	17
SMOKELESS TOBACCO⁵	36	32	32	29	13	15
CIGARS	43	49	35	41	24	28
ALCOHOL	79	83	52	54	56	61

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.11C Past Year Initiation of Substance Use among Persons Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLICIT DRUGS^{1,2}	76	82	49	58	57	56
Marijuana and Hashish	70	79	45	58	53	51
Cocaine	32	34	27	24	18	23
Crack	6	12	4	10	5	6
Heroin	9	11	7	9	6	8
Hallucinogens	43	45	36	35	22	27
LSD	28	35	24	26	16	22
PCP	7	9	6	4	3	8
Ecstasy	32	37	26	30	19	21
Inhalants	28	31	22	20	18	23
Nonmedical Use of Psychotherapeutics ^{2,3}	52	61	39	40	34	45
Pain Relievers	45	50	35	35	29	36
OxyContin [®]	22	20	18	16	13	12
Tranquilizers	33	38	25	28	20	25
Stimulants ²	32	37	24	25	22	26
Methamphetamine ²	16	17	11	11	12	13
Sedatives	10	16	5	10	9	13
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	58	68	45	44	37	50
CIGARETTES	70	77	49	60	46	49
Daily Cigarette Use ⁴	37	39	29	31	22	25
SMOKELESS TOBACCO⁵	49	49	44	45	18	20
CIGARS	71	74	54	58	40	45
ALCOHOL	105	107	69	71	73	78

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.12C Past Year Initiation of Substance Use among Persons Aged 12 to 49, by Gender: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLICIT DRUGS^{1,2}	96	100	54	74	79	70
Marijuana and Hashish	83	92	51	70	66	59
Cocaine	47	55	39	43	29	34
Crack	11	24	8	16	7	17
Heroin	31	35	28	28	13	21
Hallucinogens	63	58	43	42	44	36
LSD	40	48	28	39	32	25
PCP	7	10	6	5	4	8
Ecstasy	50	52	32	40	38	30
Inhalants	44	44	36	32	27	27
Nonmedical Use of Psychotherapeutics ^{2,3}	93	90	50	57	76	70
Pain Relievers	75	70	46	50	57	49
OxyContin [®]	41	31	33	22	24	21
Tranquilizers	83	65	40	43	71	48
Stimulants ²	47	57	33	39	35	39
Methamphetamine ²	21	31	16	20	14	21
Sedatives	26	21	14	11	22	19
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	94	95	57	60	69	72
CIGARETTES	80	90	55	68	55	57
Daily Cigarette Use ⁴	50	51	36	40	34	34
SMOKELESS TOBACCO⁵	63	59	50	50	32	28
CIGARS	119	96	84	70	76	61
ALCOHOL	112	118	75	83	82	85

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.13D Standard Errors of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older, by Gender: 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLICIT DRUGS^{1,2}	0.52	0.51	0.70	0.73	0.72	0.72
Marijuana and Hashish	0.37	0.48	0.57	0.50	0.47	0.78
Cocaine	0.36	0.66	0.55	0.97	0.43	0.77
Crack	3.92	2.78	5.71	1.78	3.05	4.79
Heroin	2.82	1.66	3.93	2.30	2.45	2.10
Hallucinogens	1.67	0.25	0.31	0.34	3.46	0.37
LSD	0.37	0.33	0.27	0.45	0.76	0.41
PCP	0.61	0.55	0.61	*	1.44	0.54
Ecstasy	0.58	0.47	0.33	0.74	1.09	0.43
Inhalants	1.53	1.91	2.09	3.26	2.26	0.77
Nonmedical Use of Psychotherapeutics ^{2,3}	0.80	0.67	0.93	1.00	1.13	0.91
Pain Relievers	0.96	0.81	1.13	1.23	1.45	1.07
OxyContin [®]	3.84	0.72	5.55	0.68	2.48	1.21
Tranquilizers	1.14	1.07	1.04	1.31	1.50	1.53
Stimulants ²	0.71	0.86	1.03	1.47	0.99	0.86
Methamphetamine ²	0.63	1.07	1.11	1.76	0.49	1.45
Sedatives	1.94	2.59	1.97	1.20	2.58	3.40
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	1.03	0.60	0.67	0.84	1.68	0.83
CIGARETTES	0.31	0.23	0.48	0.37	0.38	0.28
Daily Cigarette Use ⁴	0.83	0.49	0.37	0.56	1.74	0.80
SMOKELESS TOBACCO⁵	0.53	0.39	0.33	0.38	1.63	0.91
CIGARS	0.99	0.55	0.99	0.66	1.59	0.92
ALCOHOL	0.14	0.25	0.18	0.11	0.22	0.45

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.13P P Values from Tests of Differences of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older, by Gender: 2014 Versus 2013

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLCIT DRUGS^{1,2}	0.7734		0.2689		0.7231	
Marijuana and Hashish	0.1570		0.3095		0.2815	
Cocaine	0.0097		0.0233		0.2013	
Crack	0.7136		0.1797		0.3038	
Heroin	0.5136		0.7221		0.4015	
Hallucinogens	0.1816		0.5222		0.1238	
LSD	0.9288		0.0062		0.0078	
PCP	0.4184		*		0.9209	
Ecstasy	0.4891		0.0064		0.1924	
Inhalants	0.7049		0.7949		0.2097	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.3792		0.4279		0.1144	
Pain Relievers	0.3581		0.6518		0.1279	
OxyContin [®]	0.0103		0.0134		0.1427	
Tranquilizers	0.4805		0.5998		0.4368	
Stimulants ²	0.7167		0.2505		0.3764	
Methamphetamine ²	0.0124		0.0782		0.1069	
Sedatives	0.4678		0.0427		0.7437	
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.4021		0.1648		0.1090	
CIGARETTES	0.1474		0.1996		0.4970	
Daily Cigarette Use ⁴	0.8426		0.0855		0.5376	
SMOKELESS TOBACCO⁵	0.9700		0.5535		0.6218	
CIGARS	0.1170		0.8820		0.0578	
ALCOHOL	0.4261		0.3332		0.6170	

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.14D Standard Errors of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLCIT DRUGS^{1,2}	0.06	0.07	0.10	0.11	0.08	0.09
Marijuana and Hashish	0.06	0.07	0.10	0.10	0.08	0.08
Cocaine	0.20	0.14	0.32	0.14	0.17	0.21
Crack	0.40	0.41	*	*	*	*
Heroin	0.33	*	*	*	0.55	*
Hallucinogens	0.10	0.11	0.14	0.13	0.11	0.18
LSD	0.13	0.12	0.14	0.14	0.24	0.19
PCP	0.28	0.36	0.33	*	*	0.49
Ecstasy	0.13	0.14	0.18	0.17	0.17	0.21
Inhalants	0.16	0.14	0.23	0.25	0.22	0.17
Nonmedical Use of Psychotherapeutics ^{2,3}	0.08	0.10	0.13	0.17	0.11	0.12
Pain Relievers	0.09	0.10	0.14	0.18	0.11	0.13
OxyContin [®]	0.18	0.19	0.26	0.26	0.25	0.22
Tranquilizers	0.12	0.11	0.16	0.14	0.13	0.17
Stimulants ²	0.15	0.20	0.20	0.33	0.23	0.24
Methamphetamine ²	0.29	0.30	0.42	0.46	0.41	0.39
Sedatives	0.48	0.29	0.99	0.50	0.26	0.34
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.09	0.09	0.13	0.15	0.11	0.12
CIGARETTES	0.06	0.08	0.09	0.11	0.09	0.10
Daily Cigarette Use ⁴	0.12	0.14	0.15	0.19	0.17	0.21
SMOKELESS TOBACCO⁵	0.09	0.09	0.10	0.10	0.20	0.16
CIGARS	0.06	0.07	0.08	0.10	0.10	0.11
ALCOHOL	0.04	0.04	0.06	0.06	0.05	0.06

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.14P P Values from Tests of Differences of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: 2014 Versus 2013

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLCIT DRUGS^{1,2}	0.3303		0.9670		0.1729	
Marijuana and Hashish	0.5037		0.6215		0.1213	
Cocaine	0.9542		0.1684		0.0498	
Crack	0.2332		*		*	
Heroin	*		*		*	
Hallucinogens	0.1731		0.9445		0.0723	
LSD	0.2865		0.5924		0.4730	
PCP	0.6829		*		*	
Ecstasy	0.7522		0.5853		0.2405	
Inhalants	0.6082		0.2154		0.6344	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.5197		0.6987		0.7023	
Pain Relievers	0.6732		0.5956		0.2782	
OxyContin [®]	0.2932		0.7742		0.1056	
Tranquilizers	0.9565		0.8374		0.7403	
Stimulants ²	0.3306		0.1483		0.7628	
Methamphetamine ²	0.3739		0.7135		0.3735	
Sedatives	0.4878		0.4093		0.0519	
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.1680		0.2911		0.4835	
CIGARETTES	0.1906		0.1017		0.6519	
Daily Cigarette Use ⁴	0.7474		0.7161		0.6208	
SMOKELESS TOBACCO⁵	0.1528		0.2424		0.2440	
CIGARS	0.4313		0.6901		0.5168	
ALCOHOL	0.9525		0.6701		0.6821	

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.15D Standard Errors of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLCIT DRUGS^{1,2}	0.08	0.09	0.11	0.13	0.11	0.11
Marijuana and Hashish	0.07	0.08	0.10	0.12	0.10	0.11
Cocaine	0.14	0.17	0.20	0.22	0.19	0.26
Crack	0.45	0.35	0.69	0.32	0.59	*
Heroin	0.47	0.45	0.77	0.48	0.54	*
Hallucinogens	0.11	0.13	0.14	0.15	0.16	0.21
LSD	0.13	0.18	0.15	0.21	0.25	0.28
PCP	0.46	0.43	0.52	*	*	0.50
Ecstasy	0.13	0.16	0.17	0.19	0.20	0.26
Inhalants	0.20	0.19	0.30	0.28	0.25	0.24
Nonmedical Use of Psychotherapeutics ^{2,3}	0.11	0.11	0.17	0.17	0.15	0.14
Pain Relievers	0.13	0.13	0.19	0.19	0.16	0.17
OxyContin [®]	0.30	0.24	0.40	0.31	0.37	0.38
Tranquilizers	0.15	0.15	0.21	0.20	0.20	0.21
Stimulants ²	0.16	0.20	0.21	0.30	0.25	0.27
Methamphetamine ²	0.28	0.36	0.44	0.52	0.35	0.49
Sedatives	0.51	0.40	1.45	0.69	0.38	0.45
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.10	0.11	0.14	0.17	0.14	0.14
CIGARETTES	0.08	0.09	0.10	0.11	0.11	0.13
Daily Cigarette Use ⁴	0.10	0.13	0.13	0.15	0.16	0.22
SMOKELESS TOBACCO⁵	0.11	0.13	0.12	0.13	0.26	0.29
CIGARS	0.07	0.08	0.09	0.10	0.11	0.12
ALCOHOL	0.05	0.06	0.07	0.08	0.07	0.08

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.15P P Values from Tests of Differences of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: 2014 Versus 2013

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLICIT DRUGS^{1,2}	0.7522		0.1649		0.0763	
Marijuana and Hashish	0.4185		0.0378		0.3708	
Cocaine	0.3642		0.8956		0.2464	
Crack	0.2040		0.1386		*	
Heroin	0.2845		0.3858		*	
Hallucinogens	0.4131		0.3652		0.5873	
LSD	0.2674		0.5086		0.5715	
PCP	0.1414		*		*	
Ecstasy	0.2346		0.2876		0.5279	
Inhalants	0.1765		0.3601		0.4953	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.4630		0.8401		0.4481	
Pain Relievers	0.9540		0.5014		0.6588	
OxyContin [®]	0.3569		0.1913		0.9649	
Tranquilizers	0.6073		0.9772		0.4984	
Stimulants ²	0.0206		0.0135		0.4712	
Methamphetamine ²	0.1216		0.4868		0.1293	
Sedatives	0.5025		0.8354		0.2209	
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.8443		0.7337		0.8620	
CIGARETTES	0.0375		0.0022		0.8731	
Daily Cigarette Use ⁴	0.8751		0.5383		0.8730	
SMOKELESS TOBACCO⁵	0.0334		0.0265		0.6077	
CIGARS	0.3216		0.5018		0.4134	
ALCOHOL	0.4306		0.1258		0.7790	

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.16D Standard Errors of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49, by Gender: 2013 and 2014

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLICIT DRUGS^{1,2}	0.42	0.31	0.28	0.48	0.66	0.38
Marijuana and Hashish	0.26	0.28	0.28	0.46	0.40	0.31
Cocaine	0.36	0.48	0.55	0.63	0.43	0.71
Crack	2.70	2.78	4.33	1.78	3.05	4.79
Heroin	1.26	1.28	1.30	1.63	2.45	2.10
Hallucinogens	0.42	0.25	0.31	0.34	0.82	0.37
LSD	0.37	0.33	0.27	0.45	0.76	0.41
PCP	0.61	0.55	0.61	*	1.44	0.54
Ecstasy	0.58	0.47	0.33	0.74	1.09	0.43
Inhalants	0.95	0.80	0.94	1.31	1.87	0.77
Nonmedical Use of Psychotherapeutics ^{2,3}	0.64	0.40	0.54	0.62	0.95	0.53
Pain Relievers	0.62	0.48	0.61	0.75	0.96	0.63
OxyContin [®]	1.26	0.72	1.11	0.68	2.37	1.21
Tranquilizers	0.98	0.55	1.04	0.72	1.32	0.79
Stimulants ²	0.71	0.77	1.03	1.29	0.99	0.86
Methamphetamine ²	0.63	1.07	1.11	1.76	0.49	1.45
Sedatives	1.73	1.30	1.97	1.20	2.34	1.73
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.52	0.34	0.41	0.49	0.86	0.46
CIGARETTES	0.22	0.23	0.23	0.37	0.38	0.28
Daily Cigarette Use ⁴	0.65	0.49	0.37	0.56	1.37	0.80
SMOKELESS TOBACCO⁵	0.31	0.39	0.33	0.38	0.66	0.91
CIGARS	0.62	0.33	0.78	0.41	0.71	0.50
ALCOHOL	0.14	0.08	0.18	0.11	0.20	0.13

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 4.16P P Values from Tests of Differences of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49, by Gender: 2014 Versus 2013

Substance	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
ILLCIT DRUGS^{1,2}	0.9654		0.0042		0.1604	
Marijuana and Hashish	0.1891		0.0327		0.9673	
Cocaine	0.0234		0.0354		0.3294	
Crack	0.6075		0.6982		0.3038	
Heroin	0.0511		0.0666		0.4015	
Hallucinogens	0.2311		0.5222		0.0563	
LSD	0.9288		0.0062		0.0078	
PCP	0.4184		*		0.9209	
Ecstasy	0.4891		0.0064		0.1924	
Inhalants	0.4603		0.8637		0.4759	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.4268		0.0519		0.0484	
Pain Relievers	0.5873		0.1962		0.1115	
OxyContin [®]	0.0998		0.0807		0.2878	
Tranquilizers	0.0804		0.7668		0.0804	
Stimulants ²	0.9965		0.4521		0.3764	
Methamphetamine ²	0.0124		0.0782		0.1069	
Sedatives	0.1002		0.0427		0.3042	
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.9957		0.0353		0.2326	
CIGARETTES	0.0156		0.0056		0.4970	
Daily Cigarette Use ⁴	0.3601		0.0855		0.9746	
SMOKELESS TOBACCO⁵	0.2644		0.5535		0.2914	
CIGARS	0.0916		0.8092		0.0067	
ALCOHOL	0.9169		0.3332		0.5476	

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.1A Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2013 and 2014

Illicit Drug Dependence or Abuse Status ¹	Total		ALCOHOL DEPENDENCE OR ABUSE STATUS							
			No Dependence or Abuse		Dependence or Abuse					
	Total				Dependence		Abuse			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	262,391	265,123	245,093 ^b	248,128	17,298	16,994	7,998	8,063	9,301	8,931
NO DEPENDENCE OR ABUSE	255,540 ^b	258,045	240,831 ^b	243,643	14,709	14,403	6,588	6,651	8,121	7,752
DEPENDENCE OR ABUSE	6,852	7,077	4,262	4,486	2,589	2,592	1,410	1,412	1,179	1,179
Dependence	4,917	4,991	3,124	3,185	1,793	1,805	1,098	1,086	695	719
Abuse	1,935	2,087	1,138	1,301	797	786	312	326	485	460

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.1B Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older: Percentages, 2013 and 2014

Illicit Drug Dependence or Abuse Status ¹	Total		ALCOHOL DEPENDENCE OR ABUSE STATUS							
			No Dependence or Abuse		Dependence or Abuse					
	Total				Dependence		Abuse			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	100.0	100.0	93.4	93.6	6.6	6.4	3.0	3.0	3.5	3.4
NO DEPENDENCE OR ABUSE	97.4	97.3	91.8	91.9	5.6	5.4	2.5	2.5	3.1	2.9
DEPENDENCE OR ABUSE	2.6	2.7	1.6	1.7	1.0	1.0	0.5	0.5	0.4	0.4
Dependence	1.9	1.9	1.2	1.2	0.7	0.7	0.4	0.4	0.3	0.3
Abuse	0.7	0.8	0.4	0.5	0.3	0.3	0.1	0.1	0.2	0.2

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.2A Substance Dependence or Abuse for Specific Substances in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Past Year Dependence or Abuse	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	6,852	7,077	875	867	2,557	2,320	3,419	3,890
Marijuana and Hashish	4,206	4,176	715	667	1,878	1,706	1,612	1,804
Cocaine	855	913	22	27	232	185	601	702
Heroin	517	586	10	18	182	168	325	400
Hallucinogens	277	246	63	55	147	120	67	71
Inhalants	132	96	41	39	47 ^a	14	45	42
Nonmedical Use of Psychotherapeutics ^{2,3}	2,281	2,417	177 ^a	241	666	559	1,438	1,617
Pain Relievers	1,879	1,918	134	168	485	430	1,260	1,320
Tranquilizers	423	472	40	52	147	104	236	316
Stimulants ²	469	476	42	60	171	134	256	281
Sedatives	99	143	11	29	30	20	58	94
ALCOHOL	17,298	16,994	697	679	4,529	4,284	12,072	12,032
BOTH ILLICIT DRUGS AND ALCOHOL¹	2,589	2,592	270	296	1,073 ^a	908	1,246	1,388
ILLICIT DRUGS OR ALCOHOL¹	21,561	21,480	1,302	1,251	6,014	5,696	14,245	14,534

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.2B Substance Dependence or Abuse for Specific Substances in the Past Year, by Age Group: Percentages, 2013 and 2014

Past Year Dependence or Abuse	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	2.6	2.7	3.5	3.5	7.4	6.6	1.7	1.9
Marijuana and Hashish	1.6	1.6	2.9	2.7	5.4	4.9	0.8	0.9
Cocaine	0.3	0.3	0.1	0.1	0.7	0.5	0.3	0.3
Heroin	0.2	0.2	0.0	0.1	0.5	0.5	0.2	0.2
Hallucinogens	0.1	0.1	0.3	0.2	0.4	0.3	0.0	0.0
Inhalants	0.1	0.0	0.2	0.2	0.1 ^a	0.0	0.0	0.0
Nonmedical Use of Psychotherapeutics ^{2,3}	0.9	0.9	0.7 ^a	1.0	1.9	1.6	0.7	0.8
Pain Relievers	0.7	0.7	0.5	0.7	1.4	1.2	0.6	0.6
Tranquilizers	0.2	0.2	0.2	0.2	0.4	0.3	0.1	0.2
Stimulants ²	0.2	0.2	0.2	0.2	0.5	0.4	0.1	0.1
Sedatives	0.0	0.1	0.0	0.1	0.1	0.1	0.0	0.0
ALCOHOL	6.6	6.4	2.8	2.7	13.0	12.3	6.0	5.9
BOTH ILLICIT DRUGS AND ALCOHOL¹	1.0	1.0	1.1	1.2	3.1 ^a	2.6	0.6	0.7
ILLICIT DRUGS OR ALCOHOL¹	8.2	8.1	5.2	5.0	17.3	16.3	7.0	7.1

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.3A Substance Dependence or Abuse in the Past Year, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol ¹ (2013)	Illicit Drugs or Alcohol ¹ (2014)
TOTAL	6,852	7,077	17,298	16,994	21,561	21,480
12	21	35	9	7	27	39
13	45	42	17	26	58	60
14	107	75	71	48	137	104
15	187	180	143	139	269	235
16	252	267	177	194	355	381
17	264	269	279	264	455	430
18	375	312	440	368	660	597
19	328	284	460 ^a	357	642	547
20	339	295	549	549	746	708
21	370	306	609	686	832	839
22	322	365	673	628	855	842
23	302	313	668	600	824	789
24	295 ^a	204	624 ^a	505	809 ^a	645
25	227	241	507	592	646	729
26-29	690	877	2,197	1,849	2,669	2,408
30-34	916	689	2,015	1,899	2,515	2,323
35-39	421	469	1,451	1,475	1,706	1,760
40-44	348	390	1,415	1,398	1,657	1,668
45-49	258	358	1,367	1,354	1,521	1,579
50-54	376	484	1,149	1,441	1,427	1,721
55-59	151	285	817	1,038	901	1,234
60-64	94	177	771	600	820	726
65 or Older	163	161	891	978	1,028	1,114

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.3B Substance Dependence or Abuse in the Past Year, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol ¹ (2013)	Illicit Drugs or Alcohol ¹ (2014)
TOTAL	2.6	2.7	6.6	6.4	8.2	8.1
12	0.6	0.9	0.2	0.2	0.7	1.0
13	1.1	1.0	0.4	0.6	1.4	1.5
14	2.4	1.8	1.6	1.1	3.1	2.5
15	4.5	4.2	3.5	3.2	6.5	5.5
16	5.9	6.1	4.1	4.5	8.3	8.8
17	6.4	6.5	6.8	6.3	11.0	10.3
18	7.9	6.8	9.3	8.0	13.9	13.0
19	7.9	6.9	11.0 ^a	8.6	15.4	13.2
20	7.9	6.8	12.8	12.6	17.4	16.2
21	8.7	7.0	14.4	15.8	19.6	19.3
22	7.2	8.1	15.0	13.9	19.1	18.7
23	6.9	7.1	15.3	13.7	18.8	17.9
24	6.7 ^a	4.7	14.2 ^a	11.7	18.4 ^a	14.9
25	5.5	5.7	12.3	13.9	15.7	17.1
26-29	4.0	5.2	12.8	10.9	15.5	14.1
30-34	4.5 ^a	3.3	10.0	9.1	12.5	11.1
35-39	2.2	2.4	7.6	7.5	9.0	9.0
40-44	1.6	1.9	6.6	6.9	7.7	8.3
45-49	1.3	1.7	6.8	6.6	7.6	7.7
50-54	1.6	2.0	5.0	6.1	6.2	7.2
55-59	0.7	1.4	4.0	5.0	4.4	5.9
60-64	0.5	1.0	4.3	3.4	4.6	4.1
65 or Older	0.4	0.4	2.1	2.2	2.4	2.5

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.4A Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol ¹ (2013)	Illicit Drugs or Alcohol ¹ (2014)
TOTAL	6,852	7,077	17,298	16,994	21,561	21,480
AGE						
12-17	875	867	697	679	1,302	1,251
18-25	2,557	2,320	4,529	4,284	6,014	5,696
26 or Older	3,419	3,890	12,072	12,032	14,245	14,534
GENDER						
Male	4,278	4,422	11,080	10,945	13,733	13,731
Female	2,574	2,655	6,218	6,049	7,828	7,749
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5,800	5,898	14,306	14,080	18,006	17,867
White	4,284	4,133	11,539	11,059	14,291	13,721
Black or African American	1,045	1,313	1,661	1,889	2,324 ^a	2,713
American Indian or Alaska Native	69	40	180	211	208	231
Native Hawaiian or Other Pacific Islander	30	53	*	66	98	102
Asian	171	157	491	511	617	619
Two or More Races	202	200	361	344	468	481
Hispanic or Latino	1,051	1,180	2,992	2,915	3,554	3,614

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.4B Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	2.6	2.7	6.6	6.4	8.2	8.1
AGE						
12-17	3.5	3.5	2.8	2.7	5.2	5.0
18-25	7.4	6.6	13.0	12.3	17.3	16.3
26 or Older	1.7	1.9	6.0	5.9	7.0	7.1
GENDER						
Male	3.4	3.4	8.7	8.5	10.8	10.7
Female	1.9	1.9	4.6	4.4	5.8	5.7
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2.6	2.6	6.5	6.3	8.1	8.0
White	2.5	2.4	6.8	6.5	8.4	8.1
Black or African American	3.3	4.1	5.3	6.0	7.4	8.6
American Indian or Alaska Native	4.9	2.8	12.9	14.6	14.9	16.0
Native Hawaiian or Other Pacific Islander	3.5	5.2	*	6.4	11.3	10.0
Asian	1.3	1.1	3.7	3.7	4.6	4.5
Two or More Races	4.7	4.5	8.4	7.8	10.9	10.9
Hispanic or Latino	2.5	2.8	7.3	6.9	8.6	8.5

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.5A Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	875	867	697	679	1,302	1,251
GENDER						
Male	480	456	311	311	674	641
Female	395	412	385	367	628	610
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	670	628	554	511	1,009	932
White	485	469	476	430	782	727
Black or African American	138	121	44	55	161	148
American Indian or Alaska Native	7	5	7	5	11	8
Native Hawaiian or Other Pacific Islander	3	*	3	*	4	*
Asian	12	9	6	12	18	21
Two or More Races	24	21	18 ^a	7	32	26
Hispanic or Latino	205	239	143	168	292	318
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	246	240	204	196	377	369
Female, White, Not Hispanic	239	228	272	234	406	358
Male, Black, Not Hispanic	81	72	18	26	88	80
Female, Black, Not Hispanic	57	48	27	29	73	67
Male, Hispanic	132	125	75	77	179	163
Female, Hispanic	74	114	68	90	113	155

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.5B Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	3.5	3.5	2.8	2.7	5.2	5.0
GENDER						
Male	3.8	3.6	2.5	2.5	5.3	5.1
Female	3.2	3.4	3.2	3.0	5.2	5.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	3.5	3.3	2.9	2.7	5.2	4.8
White	3.6	3.5	3.5	3.2	5.7	5.4
Black or African American	4.0	3.5	1.3	1.6	4.7	4.2
American Indian or Alaska Native	4.6	3.3	4.2	3.0	7.1	4.7
Native Hawaiian or Other Pacific Islander	2.9	*	2.9	*	4.3	*
Asian	1.0	0.7	0.5	1.0	1.5	1.7
Two or More Races	3.2	2.7	2.3 ^a	0.9	4.3	3.4
Hispanic or Latino	3.7	4.2	2.6	3.0	5.3	5.6
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	3.5	3.5	2.9	2.8	5.4	5.4
Female, White, Not Hispanic	3.6	3.5	4.1	3.5	6.1	5.4
Male, Black, Not Hispanic	4.6	4.1	1.0	1.5	5.0	4.5
Female, Black, Not Hispanic	3.4	2.8	1.6	1.7	4.3	3.9
Male, Hispanic	4.6	4.3	2.7	2.7	6.3	5.7
Female, Hispanic	2.7	4.1	2.5	3.3	4.2	5.6

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.6A Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	2,557	2,320	4,529	4,284	6,014	5,696
GENDER						
Male	1,709 ^b	1,439	2,731	2,556	3,707	3,444
Female	849	881	1,798	1,729	2,307	2,252
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,143	1,940	3,598	3,385	4,850	4,567
White	1,546	1,379	2,874 ^a	2,565	3,755 ^a	3,386
Black or African American	395	393	422	439	675	701
American Indian or Alaska Native	25 ^a	10	37	38	51	42
Native Hawaiian or Other Pacific Islander	16	*	17	*	28	*
Asian	81	77	139	212	196	256
Two or More Races	80	71	109	109	145	156
Hispanic or Latino	414	380	932	900	1,164	1,129
EDUCATION						
< High School	550 ^a	410	632	537	979 ^a	790
High School Graduate	922	801	1,279	1,240	1,823	1,738
Some College	857	882	1,791	1,629	2,265	2,161
College Graduate	228	227	828	878	947	1,006
CURRENT EMPLOYMENT						
Full-Time	932	907	1,989	1,976	2,512	2,544
Part-Time	687	630	1,322	1,156	1,705	1,508
Unemployed	488	388	516	504	810	739
Other ²	450	395	702	649	987	905

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.6B Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	7.4	6.6	13.0	12.3	17.3	16.3
GENDER						
Male	9.8 ^b	8.2	15.6	14.6	21.2	19.6
Female	4.9	5.1	10.4	9.9	13.3	12.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	7.8	7.0	13.1	12.3	17.6	16.6
White	7.9	7.1	14.7 ^a	13.2	19.3 ^a	17.5
Black or African American	7.9	7.8	8.4	8.7	13.5	13.9
American Indian or Alaska Native	10.9 ^a	4.8	16.1	18.9	22.0	20.7
Native Hawaiian or Other Pacific Islander	7.9	*	8.3	*	14.2	*
Asian	4.3	3.8	7.4	10.5	10.5	12.8
Two or More Races	11.0	9.0	14.9	13.9	19.8	19.9
Hispanic or Latino	5.7	5.2	12.8	12.2	16.1	15.3
EDUCATION						
< High School	9.8	8.3	11.3	10.9	17.5	16.0
High School Graduate	7.7	6.7	10.7	10.4	15.2	14.5
Some College	7.0	7.1	14.7 ^a	13.1	18.6	17.4
College Graduate	4.5	4.1	16.4	15.8	18.8	18.1
CURRENT EMPLOYMENT						
Full-Time	6.9	6.4	14.7	13.9	18.6	17.8
Part-Time	7.1	6.6	13.7	12.2	17.6	15.9
Unemployed	11.7	10.0	12.4	13.0	19.5	19.0
Other ²	6.0	5.4	9.4	8.9	13.2	12.4

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.7A Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	3,419	3,890	12,072	12,032	14,245	14,534
GENDER						
Male	2,089 ^a	2,527	8,037	8,078	9,353	9,647
Female	1,330	1,363	4,035	3,954	4,892	4,887
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,987	3,330	10,155	10,184	12,147	12,367
White	2,252	2,286	8,189	8,064	9,754	9,608
Black or African American	511 ^a	799	1,195	1,395	1,489	1,865
American Indian or Alaska Native	37	25	136	167	147	181
Native Hawaiian or Other Pacific Islander	12	40	*	42	*	72
Asian	77	71	346	287	402	342
Two or More Races	97	109	234	228	291	299
Hispanic or Latino	432	560	1,918	1,848	2,098	2,167
EDUCATION						
< High School	665	773	1,526	1,589	1,950	2,095
High School Graduate	1,077	1,059	3,130	2,949	3,696	3,636
Some College	1,076	1,186	3,028	3,375	3,814	4,146
College Graduate	600 ^a	873	4,389	4,119	4,785	4,657
CURRENT EMPLOYMENT						
Full-Time	1,580	1,943	7,824	7,637	8,776	8,802
Part-Time	454	527	1,140	1,430	1,429	1,773
Unemployed	390	355	715	685	981	894
Other ²	996	1,065	2,394	2,278	3,059	3,066

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.7B Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	1.7	1.9	6.0	5.9	7.0	7.1
GENDER						
Male	2.2	2.6	8.3	8.2	9.6	9.8
Female	1.3	1.3	3.8	3.7	4.6	4.6
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1.7	1.9	5.8	5.8	7.0	7.0
White	1.6	1.7	6.0	5.9	7.1	7.0
Black or African American	2.2 ^a	3.4	5.2	6.0	6.5	8.0
American Indian or Alaska Native	3.6	2.4	13.5	15.6	14.5	16.9
Native Hawaiian or Other Pacific Islander	2.0	5.4	*	5.6	*	9.7
Asian	0.8	0.7	3.4	2.7	3.9	3.2
Two or More Races	3.5	3.8	8.4	8.0	10.4	10.4
Hispanic or Latino	1.5	1.9	6.7	6.3	7.4	7.4
EDUCATION						
< High School	2.5	2.8	5.7	5.8	7.3	7.7
High School Graduate	1.9	1.9	5.4	5.2	6.4	6.4
Some College	2.1	2.2	5.9	6.4	7.4	7.8
College Graduate	0.9 ^a	1.3	6.6	6.1	7.2	6.9
CURRENT EMPLOYMENT						
Full-Time	1.5	1.8	7.4	7.0	8.3	8.1
Part-Time	1.9	2.2	4.7	6.1	5.9 ^a	7.5
Unemployed	5.1	5.1	9.4	9.9	12.8	12.9
Other ²	1.5	1.6	3.7	3.4	4.7	4.6

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.8A Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	5,976	6,210	16,602	16,316	20,259	20,230
GENDER						
Male	3,798	3,966	10,769	10,633	13,059	13,091
Female	2,179	2,244	5,833	5,682	7,200	7,139
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5,130	5,269	13,752	13,568	16,997	16,934
White	3,799	3,665	11,063	10,629	13,508	12,994
Black or African American	906 ^a	1,192	1,617	1,834	2,163	2,566
American Indian or Alaska Native	62	35	173	206	197	223
Native Hawaiian or Other Pacific Islander	27	50	*	64	*	98
Asian	158	148	485	499	598	598
Two or More Races	178	179	343	338	435	455
Hispanic or Latino	846	941	2,850	2,747	3,262	3,296
EDUCATION						
< High School	1,215	1,183	2,158	2,126	2,930	2,885
High School Graduate	2,000	1,860	4,408	4,189	5,519	5,374
Some College	1,933	2,068	4,819	5,003	6,079	6,308
College Graduate	829 ^a	1,099	5,216	4,997	5,731	5,664
CURRENT EMPLOYMENT						
Full-Time	2,512	2,850	9,813	9,613	11,288	11,346
Part-Time	1,141	1,158	2,462	2,586	3,135	3,280
Unemployed	878	743	1,231	1,189	1,791	1,633
Other ²	1,446	1,459	3,096	2,927	4,045	3,971

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.8B Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	2.5	2.6	7.0	6.8	8.5	8.4
GENDER						
Male	3.3	3.4	9.4	9.2	11.4	11.3
Female	1.8	1.8	4.7	4.6	5.8	5.7
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2.5	2.6	6.8	6.7	8.4	8.3
White	2.4	2.3	7.1	6.8	8.6	8.3
Black or African American	3.3	4.2	5.8	6.5	7.8	9.1
American Indian or Alaska Native	5.0	2.7	14.0	16.1	15.9	17.5
Native Hawaiian or Other Pacific Islander	3.5	5.4	*	6.9	*	10.6
Asian	1.3	1.2	4.0	4.0	4.9	4.7
Two or More Races	5.0	4.9	9.7	9.2	12.3	12.5
Hispanic or Latino	2.4	2.6	8.0	7.5	9.1	8.9
EDUCATION						
< High School	3.8	3.7	6.7	6.6	9.1	9.0
High School Graduate	2.9	2.7	6.3	6.1	7.9	7.8
Some College	3.0	3.2	7.6	7.6	9.6	9.6
College Graduate	1.2 ^a	1.5	7.2	6.8	8.0	7.7
CURRENT EMPLOYMENT						
Full-Time	2.1	2.3	8.2	7.8	9.5	9.2
Part-Time	3.4	3.5	7.3	7.8	9.3	9.9
Unemployed	7.4	6.9	10.4	11.0	15.2	15.1
Other ²	2.0	2.0	4.2	4.0	5.5	5.4

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.9A Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	6,852	7,077	17,298	16,994	21,561	21,480
GEOGRAPHIC DIVISION						
Northeast	1,270	1,537	3,105	3,141	3,922	4,134
New England	409	393	816	828	1,098	1,087
Middle Atlantic	861 ^a	1,145	2,290	2,313	2,824	3,047
Midwest	1,356	1,336	3,755	3,756	4,601	4,588
East North Central	980	998	2,602	2,605	3,208	3,243
West North Central	375	338	1,153	1,151	1,393	1,346
South	2,420	2,399	6,051	6,055	7,602	7,584
South Atlantic	1,380	1,367	3,336	3,313	4,205	4,209
East South Central	428	328	797	765	1,092	980
West South Central	611	705	1,918	1,977	2,305	2,395
West	1,806	1,804	4,387	4,043	5,436	5,174
Mountain	603	552	1,328	1,475	1,709	1,804
Pacific	1,203	1,252	3,059	2,568	3,726	3,370
COUNTY TYPE						
Large Metro	3,798	4,137	9,819	9,726	12,119	12,312
Small Metro	2,198	2,128	5,175	4,958	6,633	6,344
250K – 1 Mil. Pop.	1,431	1,432	3,657	3,320	4,575	4,274
< 250K Pop.	768	697	1,518	1,639	2,058	2,071
Nonmetro	856	812	2,305	2,310	2,808	2,824
Urbanized	455	399	1,129	1,084	1,398	1,318
Less Urbanized	326	353	970	1,049	1,160	1,294
Completely Rural	74	60	205	177	250	211

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.9B Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	2.6	2.7	6.6	6.4	8.2	8.1
GEOGRAPHIC DIVISION						
Northeast	2.7	3.2	6.6	6.6	8.3	8.7
New England	3.3	3.1	6.5	6.6	8.8	8.6
Middle Atlantic	2.5 ^a	3.3	6.6	6.6	8.1	8.7
Midwest	2.4	2.4	6.7	6.7	8.2	8.1
East North Central	2.5	2.6	6.7	6.7	8.2	8.3
West North Central	2.2	1.9	6.7	6.6	8.1	7.8
South	2.5	2.4	6.2	6.1	7.8	7.7
South Atlantic	2.7	2.6	6.5	6.4	8.2	8.1
East South Central	2.8	2.1	5.1	4.9	7.0	6.3
West South Central	2.0	2.3	6.3	6.4	7.5	7.7
West	2.9	2.9	7.2	6.5	8.9	8.3
Mountain	3.2	2.9	7.1	7.8	9.1	9.5
Pacific	2.8	2.9	7.2 ^a	5.9	8.8	7.8
COUNTY TYPE						
Large Metro	2.7	2.8	7.0	6.6	8.6	8.4
Small Metro	2.8	2.7	6.6	6.4	8.4	8.2
250K – 1 Mil. Pop.	2.7	2.7	6.8	6.3	8.6	8.2
< 250K Pop.	3.0	2.8	6.0	6.5	8.1	8.3
Nonmetro	2.0	2.0	5.4	5.6	6.6	6.9
Urbanized	2.4	2.3	5.9	6.3	7.4	7.6
Less Urbanized	1.7	1.8	5.1	5.3	6.1	6.6
Completely Rural	1.7	1.4	4.6	4.2	5.6	5.0

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.10A Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	875	867	697	679	1,302	1,251
GEOGRAPHIC DIVISION						
Northeast	134	112	131	114	219	182
New England	38	37	36	28	57	53
Middle Atlantic	96	75	95	86	161	129
Midwest	161	178	141	132	253	258
East North Central	123	126	99	102	184	188
West North Central	38	51	42	30	69	70
South	329	303	254	239	500	440
South Atlantic	169	171	132	112	261	230
East South Central	47	33	45	27	80	52
West South Central	113	100	77	101	159	158
West	251	275	171	193	330	370
Mountain	97	81	68	67	122	113
Pacific	154	193	104	126	208	257
COUNTY TYPE						
Large Metro	477	499	349	366	694	703
Small Metro	291	233	229 ^a	171	426 ^a	331
250K – 1 Mil. Pop.	196	162	147	109	281	223
< 250K Pop.	95	71	82	61	145	108
Nonmetro	107	135	119	143	181	216
Urbanized	57	59	49	68	82	96
Less Urbanized	42	63	51	58	78	96
Completely Rural	8	13	19	16	21	24

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.10B Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	3.5	3.5	2.8	2.7	5.2	5.0
GEOGRAPHIC DIVISION						
Northeast	3.2	2.7	3.1	2.8	5.2	4.4
New England	3.5	3.4	3.3	2.6	5.2	4.9
Middle Atlantic	3.1	2.4	3.1	2.8	5.2	4.2
Midwest	3.0	3.3	2.6	2.5	4.7	4.8
East North Central	3.3	3.4	2.6	2.7	4.9	5.1
West North Central	2.3	3.1	2.6	1.8	4.2	4.3
South	3.5	3.2	2.7	2.5	5.3	4.7
South Atlantic	3.6	3.6	2.8	2.4	5.6	4.9
East South Central	3.2	2.2	3.1	1.8	5.4	3.5
West South Central	3.5	3.1	2.4	3.1	4.9	4.9
West	4.2	4.6	2.9	3.3	5.5	6.2
Mountain	5.2	4.3	3.6	3.5	6.5	6.0
Pacific	3.8	4.8	2.5	3.1	5.1	6.4
COUNTY TYPE						
Large Metro	3.5	3.6	2.6	2.6	5.1	5.0
Small Metro	3.9	3.3	3.1	2.4	5.7 ^a	4.7
250K – 1 Mil. Pop.	3.8	3.3	2.9	2.2	5.5	4.6
< 250K Pop.	4.1	3.2	3.5	2.8	6.3	4.9
Nonmetro	2.8	3.6	3.1	3.8	4.7	5.7
Urbanized	3.4	3.8	2.9	4.4	4.9	6.2
Less Urbanized	2.4	3.4	2.9	3.1	4.4	5.1
Completely Rural	1.9	3.7	4.3	4.3	4.7	6.6

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.11A Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	2,557	2,320	4,529	4,284	6,014	5,696
GEOGRAPHIC DIVISION						
Northeast	504	458	825	747	1,121	1,043
New England	145	165	252	231	326	333
Middle Atlantic	359	293	573	516	795	710
Midwest	527	506	1,028	1,030	1,334	1,300
East North Central	359	367	732	676	936	880
West North Central	168	139	296	354	398	420
South	859	776	1,550	1,472	2,066	1,941
South Atlantic	501	451	847	782	1,129	1,051
East South Central	106	114	241	215	316	287
West South Central	252	212	463	475	621	603
West	668	580	1,126	1,036	1,493	1,412
Mountain	195	161	314	320	440	407
Pacific	474	419	812	716	1,054	1,005
COUNTY TYPE						
Large Metro	1,477	1,350	2,475	2,362	3,334	3,224
Small Metro	737	709	1,388	1,274	1,826	1,694
250K – 1 Mil. Pop.	478	473	904	787	1,179	1,065
< 250K Pop.	258	236	484	487	647	629
Nonmetro	344 ^a	262	666	648	854	777
Urbanized	155	114	352	324	440	365
Less Urbanized	168	128	258	271	349	349
Completely Rural	20	20	56	53	66	63

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.11B Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	7.4	6.6	13.0	12.3	17.3	16.3
GEOGRAPHIC DIVISION						
Northeast	8.2	7.4	13.4	12.2	18.2	17.0
New England	8.9	10.0	15.4	14.1	20.0	20.2
Middle Atlantic	7.9	6.5	12.7	11.4	17.6	15.8
Midwest	7.1	6.8	13.9	13.9	18.0	17.5
East North Central	7.0	7.2	14.3	13.2	18.3	17.2
West North Central	7.3	6.0	12.9	15.3	17.3	18.1
South	6.7	6.0	12.1	11.4	16.1	15.0
South Atlantic	7.6	6.8	12.9	11.9	17.2	15.9
East South Central	5.2	5.6	11.8	10.5	15.5	14.0
West South Central	5.9	4.9	10.9	11.0	14.6	14.0
West	8.0	6.9	13.4	12.3	17.8	16.8
Mountain	7.7	6.3	12.4	12.5	17.4	15.9
Pacific	8.1	7.2	13.9	12.2	18.0	17.2
COUNTY TYPE						
Large Metro	7.9	7.0	13.2	12.3	17.8	16.8
Small Metro	6.8	6.7	12.8	12.0	16.8	15.9
250K – 1 Mil. Pop.	6.8	6.9	12.9	11.4	16.9	15.5
< 250K Pop.	6.6	6.3	12.5	13.0	16.7	16.7
Nonmetro	6.6 ^a	5.1	12.9	12.7	16.5	15.2
Urbanized	6.3	4.7	14.2	13.5	17.8	15.3
Less Urbanized	7.6	5.5	11.6	11.6	15.7	14.9
Completely Rural	4.2	5.2	11.8	13.6	13.9	16.2

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.12A Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	3,419	3,890	12,072	12,032	14,245	14,534
GEOGRAPHIC DIVISION						
Northeast	632 ^a	968	2,150	2,279	2,582	2,909
New England	226	191	529	569	715	701
Middle Atlantic	406 ^b	777	1,622	1,710	1,867	2,208
Midwest	668	652	2,586	2,595	3,014	3,030
East North Central	499	505	1,772	1,827	2,088	2,175
West North Central	170	148	814	767	926	856
South	1,232	1,320	4,247	4,343	5,036	5,203
South Atlantic	710	745	2,358	2,419	2,815	2,928
East South Central	275	181	511	523	696	642
West South Central	246	393	1,378	1,401	1,525	1,633
West	887	950	3,090	2,814	3,613	3,392
Mountain	311	311	946	1,088	1,148	1,283
Pacific	575	639	2,144	1,726	2,465	2,109
COUNTY TYPE						
Large Metro	1,843 ^a	2,288	6,994	6,999	8,091	8,385
Small Metro	1,171	1,186	3,559	3,514	4,381	4,319
250K – 1 Mil. Pop.	756	797	2,606	2,423	3,116	2,985
< 250K Pop.	415	390	953	1,091	1,265	1,334
Nonmetro	405	415	1,520	1,519	1,773	1,830
Urbanized	243	226	729	691	876	857
Less Urbanized	116	162	661	720	733	850
Completely Rural	46	27	130	108	164	124

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.12B Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	1.7	1.9	6.0	5.9	7.0	7.1
GEOGRAPHIC DIVISION						
Northeast	1.7 ^a	2.6	5.8	6.1	7.0	7.8
New England	2.3	1.9	5.4	5.8	7.3	7.1
Middle Atlantic	1.5 ^b	2.8	5.9	6.2	6.8	8.0
Midwest	1.5	1.5	6.0	5.9	6.9	6.9
East North Central	1.7	1.7	5.9	6.0	6.9	7.2
West North Central	1.3	1.1	6.1	5.7	7.0	6.4
South	1.6	1.7	5.6	5.7	6.7	6.8
South Atlantic	1.8	1.8	5.9	5.9	7.0	7.2
East South Central	2.3	1.5	4.3	4.3	5.8	5.3
West South Central	1.1	1.7	6.0	6.0	6.6	6.9
West	1.9	2.0	6.6	5.9	7.7	7.1
Mountain	2.2	2.1	6.6	7.5	8.0	8.8
Pacific	1.8	1.9	6.6	5.2	7.5	6.3
COUNTY TYPE						
Large Metro	1.7	2.0	6.4	6.2	7.4	7.4
Small Metro	1.9	2.0	5.9	5.9	7.2	7.2
250K – 1 Mil. Pop.	1.8	2.0	6.3	6.0	7.5	7.3
< 250K Pop.	2.1	2.0	4.9	5.7	6.6	7.0
Nonmetro	1.2	1.3	4.6	4.7	5.3	5.7
Urbanized	1.6	1.7	4.9	5.2	5.9	6.4
Less Urbanized	0.8	1.0	4.4	4.6	4.9	5.5
Completely Rural	1.3	0.8	3.7	3.1	4.6	3.6

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.13A Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	5,976	6,210	16,602	16,316	20,259	20,230
GEOGRAPHIC DIVISION						
Northeast	1,136 ^a	1,426	2,975	3,026	3,703	3,952
New England	371	356	780	800	1,041	1,034
Middle Atlantic	765 ^a	1,070	2,195	2,226	2,662	2,918
Midwest	1,195	1,159	3,614	3,624	4,348	4,330
East North Central	858	872	2,504	2,503	3,024	3,054
West North Central	337	287	1,110	1,121	1,324	1,276
South	2,090	2,096	5,797	5,815	7,102	7,144
South Atlantic	1,211	1,196	3,205	3,201	3,944	3,979
East South Central	381	295	752	738	1,012	929
West South Central	498	605	1,840	1,876	2,146	2,237
West	1,555	1,530	4,216	3,850	5,106	4,804
Mountain	506	471	1,260	1,408	1,587	1,690
Pacific	1,049	1,059	2,956 ^a	2,442	3,518	3,113
COUNTY TYPE						
Large Metro	3,320	3,638	9,470	9,361	11,425	11,609
Small Metro	1,907	1,895	4,947	4,788	6,206	6,013
250K – 1 Mil. Pop.	1,235	1,269	3,510	3,210	4,294	4,050
< 250K Pop.	673	626	1,437	1,578	1,912	1,963
Nonmetro	748	677	2,185	2,167	2,628	2,608
Urbanized	399	340	1,080	1,015	1,316	1,222
Less Urbanized	284	290	919	991	1,082	1,198
Completely Rural	66	47	186	161	230	187

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.13B Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	2.5	2.6	7.0	6.8	8.5	8.4
GEOGRAPHIC DIVISION						
Northeast	2.6	3.3	6.9	7.0	8.6	9.1
New England	3.3	3.1	6.9	7.0	9.1	9.0
Middle Atlantic	2.4 ^a	3.3	6.9	7.0	8.4	9.1
Midwest	2.4	2.3	7.1	7.1	8.6	8.5
East North Central	2.4	2.5	7.1	7.1	8.6	8.6
West North Central	2.2	1.8	7.1	7.1	8.5	8.1
South	2.4	2.3	6.6	6.5	8.1	8.0
South Atlantic	2.6	2.5	6.8	6.7	8.4	8.4
East South Central	2.7	2.1	5.4	5.2	7.2	6.6
West South Central	1.8	2.2	6.7	6.7	7.8	8.0
West	2.8	2.7	7.6	6.8	9.2	8.5
Mountain	3.0	2.8	7.5	8.2	9.4	9.9
Pacific	2.7	2.7	7.7 ^a	6.2	9.1	8.0
COUNTY TYPE						
Large Metro	2.6	2.7	7.4	7.1	9.0	8.8
Small Metro	2.7	2.7	6.9	6.8	8.7	8.5
250K – 1 Mil. Pop.	2.6	2.7	7.3	6.8	8.9	8.5
< 250K Pop.	2.9	2.7	6.2	6.9	8.2	8.6
Nonmetro	1.9	1.8	5.7	5.8	6.8	7.0
Urbanized	2.3	2.2	6.2	6.5	7.6	7.8
Less Urbanized	1.7	1.6	5.3	5.6	6.3	6.7
Completely Rural	1.6	1.2	4.6	4.2	5.7	4.9

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.14A Substance Dependence for Specific Substances in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Past Year Dependence	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	4,917	4,991	480	455	1,808	1,661	2,629	2,875
Marijuana and Hashish	2,804	2,656	383 ^a	304	1,251	1,148	1,171	1,204
Cocaine	624	571	11	11	141	114	472	446
Heroin	434	539	3	12	155	152	276	374
Hallucinogens	102	94	22	17	50	50	30	27
Inhalants	39	43	15	15	16	5	*	24
Nonmedical Use of Psychotherapeutics ^{2,3}	1,758	1,846	100 ^a	148	513	437	1,145	1,261
Pain Relievers	1,488	1,507	86	103	385	346	1,017	1,058
Tranquilizers	237	284	17	21	83	61	137	202
Stimulants ²	367	335	23	31	131	106	213	199
Sedatives	61	115	4	21	22	17	35	78
ALCOHOL	7,998	8,063	261	239	1,922	1,989	5,815	5,835
BOTH ILLICIT DRUGS AND ALCOHOL¹	1,098	1,086	91	74	444	377	562	635
ILLICIT DRUGS OR ALCOHOL¹	11,817	11,968	650	620	3,285	3,274	7,882	8,074

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.14B Substance Dependence for Specific Substances in the Past Year, by Age Group: Percentages, 2013 and 2014

Past Year Dependence	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	1.9	1.9	1.9	1.8	5.2	4.8	1.3	1.4
Marijuana and Hashish	1.1	1.0	1.5 ^a	1.2	3.6	3.3	0.6	0.6
Cocaine	0.2	0.2	0.0	0.0	0.4	0.3	0.2	0.2
Heroin	0.2	0.2	0.0	0.0	0.4	0.4	0.1	0.2
Hallucinogens	0.0	0.0	0.1	0.1	0.1	0.1	0.0	0.0
Inhalants	0.0	0.0	0.1	0.1	0.0	0.0	*	0.0
Nonmedical Use of Psychotherapeutics ^{2,3}	0.7	0.7	0.4 ^a	0.6	1.5	1.3	0.6	0.6
Pain Relievers	0.6	0.6	0.3	0.4	1.1	1.0	0.5	0.5
Tranquilizers	0.1	0.1	0.1	0.1	0.2	0.2	0.1	0.1
Stimulants ²	0.1	0.1	0.1	0.1	0.4	0.3	0.1	0.1
Sedatives	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0
ALCOHOL	3.0	3.0	1.0	1.0	5.5	5.7	2.9	2.8
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.4	0.4	0.4	0.3	1.3	1.1	0.3	0.3
ILLICIT DRUGS OR ALCOHOL¹	4.5	4.5	2.6	2.5	9.4	9.4	3.9	3.9

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.15A Received Substance Use Treatment in the Past Year, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	2,214	2,440	2,731	2,683	1,320	1,427	4,051	4,149
12	*	4	3	4	*	4	4	6
13	10	3	13	6	9	3	14	10
14	19	21	13	13	12	12	26	28
15	33	29	23	15	18	12	52	35
16	61	40	47 ^a	22	41 ^a	14	71	63
17	52	70	39	48	32	46	65	85
18	65	39	55	27	38	17	98 ^a	54
19	60	46	47	46	31	24	91	75
20	55	52	52	42	35	32	79	80
21	68	64	56	48	46	34	83	81
22	69	67	61	53	44	27	93	105
23	65	85	73	66	42	45	111	126
24	57	58	55	59	27	23	96	104
25	69	85	69	84	40	41	114	139
26-29	297	317	338	263	188	152	494	484
30-34	322	274	311	309	165	148	516	463
35-39	180	204	286	242	131	105	345	359
40-44	171	229	272	247	84	131	412	378
45-49	105	129	214	198	60	89	346	270
50-54	163	237	193	308	94	174	281	434
55-59	165	227	154	228	86	153	259	315
60-64	116	61	285	152	85	42	316	220
65 or Older	12 ^a	101	76 ^a	205	12 ^a	98	85 ^a	234

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.15B Received Substance Use Treatment in the Past Year, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.8	0.9	1.0	1.0	0.5	0.5	1.5	1.6
12	*	0.1	0.1	0.1	*	0.1	0.1	0.2
13	0.2	0.1	0.3	0.1	0.2	0.1	0.3	0.2
14	0.4	0.5	0.3	0.3	0.3	0.3	0.6	0.7
15	0.8	0.7	0.6	0.4	0.4	0.3	1.3	0.8
16	1.4	0.9	1.1 ^a	0.5	1.0 ^a	0.3	1.7	1.5
17	1.3	1.7	0.9	1.1	0.8	1.1	1.6	2.0
18	1.4	0.9	1.2	0.6	0.8	0.4	2.1 ^a	1.2
19	1.4	1.1	1.1	1.1	0.7	0.6	2.2	1.8
20	1.3	1.2	1.2	1.0	0.8	0.7	1.8	1.8
21	1.6	1.5	1.3	1.1	1.1	0.8	2.0	1.9
22	1.5	1.5	1.4	1.2	1.0	0.6	2.1	2.3
23	1.5	1.9	1.7	1.5	1.0	1.0	2.5	2.9
24	1.3	1.3	1.2	1.4	0.6	0.5	2.2	2.4
25	1.7	2.0	1.7	2.0	1.0	1.0	2.8	3.3
26-29	1.7	1.9	2.0	1.5	1.1	0.9	2.9	2.8
30-34	1.6	1.3	1.5	1.5	0.8	0.7	2.6	2.2
35-39	0.9	1.0	1.5	1.2	0.7	0.5	1.8	1.8
40-44	0.8	1.1	1.3	1.2	0.4	0.6	1.9	1.9
45-49	0.5	0.6	1.1	1.0	0.3	0.4	1.7	1.3
50-54	0.7	1.0	0.8	1.3	0.4	0.7	1.2	1.8
55-59	0.8	1.1	0.7	1.1	0.4	0.7	1.3	1.5
60-64	0.7	0.3	1.6	0.9	0.5	0.2	1.8	1.3
65 or Older	0.0 ^a	0.2	0.2	0.5	0.0 ^a	0.2	0.2 ^a	0.5

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.16A Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	2,214	2,440	2,731	2,683	1,320	1,427	4,051	4,149
AGE								
12-17	175	166	137	107	113	91	232	227
18-25	509	496	466	425	304	244	764	764
26 or Older	1,530	1,778	2,128	2,151	903	1,092	3,055	3,158
GENDER								
Male	1,281	1,462	1,747	1,768	774	867	2,569	2,703
Female	933	979	984	915	546	560	1,482	1,446
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	1,860	2,075	2,279	2,258	1,123	1,245	3,329	3,461
White	1,486	1,559	1,760	1,690	863	872	2,613	2,631
Black or African American	272 ^a	435	319	427	169 ^a	320	473	647
American Indian or Alaska Native	28	22	45	37	28	16	49	46
Native Hawaiian or Other Pacific Islander	9	11	9	7	9	4	17	16
Asian	27	19	68	35	25	14	86	44
Two or More Races	38	29	77	61	29	18	91	77
Hispanic or Latino	355	365	452	425	197	182	722	688

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.16B Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.8	0.9	1.0	1.0	0.5	0.5	1.5	1.6
AGE								
12-17	0.7	0.7	0.6	0.4	0.5	0.4	0.9	0.9
18-25	1.5	1.4	1.3	1.2	0.9	0.7	2.2	2.2
26 or Older	0.8	0.9	1.0	1.0	0.4	0.5	1.5	1.5
GENDER								
Male	1.0	1.1	1.4	1.4	0.6	0.7	2.0	2.1
Female	0.7	0.7	0.7	0.7	0.4	0.4	1.1	1.1
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.8	0.9	1.0	1.0	0.5	0.6	1.5	1.6
White	0.9	0.9	1.0	1.0	0.5	0.5	1.5	1.5
Black or African American	0.9 ^a	1.4	1.0	1.3	0.5 ^a	1.0	1.5	2.0
American Indian or Alaska Native	2.0	1.6	3.2	2.6	2.0	1.1	3.5	3.2
Native Hawaiian or Other Pacific Islander	1.0	1.1	1.0	0.7	1.0	0.4	2.0	1.6
Asian	0.2	0.1	0.5	0.3	0.2	0.1	0.6	0.3
Two or More Races	0.9	0.7	1.8	1.4	0.7	0.4	2.1	1.7
Hispanic or Latino	0.9	0.9	1.1	1.0	0.5	0.4	1.8	1.6

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.17A Received Substance Use Treatment in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	175	166	137	107	113	91	232	227
GENDER								
Male	111	78	76 ^a	41	66 ^a	37	142	111
Female	64	88	61	66	47	55	90	116
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	126	124	99	89	80	77	172	168
White	90	98	70	75	53	64	129	133
Black or African American	21	13	14	7	13	7	23	19
American Indian or Alaska Native	6	2	5	1	5	*	6	2
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	6	*	0	*	0	1	6
Two or More Races	8	6	7	5	5	5	9	7
Hispanic or Latino	49	42	38	18	33 ^a	14	61	59
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	55	52	39	29	30	26	77	74
Female, White, Not Hispanic	34	45	32	45	23	37	52	60
Male, Black, Not Hispanic	18	7	11	6	10	5	20	10
Female, Black, Not Hispanic	3	5	3	2	3	2	3	10
Male, Hispanic	28	15	18 ^a	5	18 ^a	5	33	22
Female, Hispanic	21	28	21	13	16	10	27	37

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.17B Received Substance Use Treatment in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.7	0.7	0.6	0.4	0.5	0.4	0.9	0.9
GENDER								
Male	0.9	0.6	0.6 ^a	0.3	0.5 ^a	0.3	1.1	0.9
Female	0.5	0.7	0.5	0.5	0.4	0.4	0.7	1.0
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.7	0.6	0.5	0.5	0.4	0.4	0.9	0.9
White	0.7	0.7	0.5	0.6	0.4	0.5	0.9	1.0
Black or African American	0.6	0.4	0.4	0.2	0.4	0.2	0.7	0.6
American Indian or Alaska Native	3.5	1.2	3.4	0.7	3.4	*	3.8	1.5
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	0.5	*	0.0	*	0.0	0.1	0.5
Two or More Races	1.0	0.8	0.9	0.7	0.7	0.7	1.2	0.9
Hispanic or Latino	0.9	0.7	0.7 ^a	0.3	0.6 ^a	0.3	1.1	1.0
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	0.8	0.8	0.6	0.4	0.4	0.4	1.1	1.1
Female, White, Not Hispanic	0.5	0.7	0.5	0.7	0.3	0.6	0.8	0.9
Male, Black, Not Hispanic	1.0	0.4	0.6	0.3	0.6	0.3	1.1	0.6
Female, Black, Not Hispanic	0.2	0.3	0.2	0.1	0.2	0.1	0.2	0.6
Male, Hispanic	1.0	0.5	0.6 ^a	0.2	0.6 ^a	0.2	1.2	0.8
Female, Hispanic	0.8	1.0	0.8	0.5	0.6	0.3	1.0	1.3

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.18A Received Substance Use Treatment in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	509	496	466	425	304	244	764	764
GENDER								
Male	337	310	314	280	211	153	510	497
Female	172	186	152	145	93	91	254	267
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	441	415	366	373	257	215	610	642
White	377	339	306	284	217	165	495	507
Black or African American	36	45	34	44	19	25	69	77
American Indian or Alaska Native	6	6	8	9	6	6	8	11
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	10	7	7	19	7	7	24	20
Two or More Races	12	17	11	14	9	11	15	25
Hispanic or Latino	69	81	100 ^a	52	46	29	154	122
EDUCATION								
< High School	139	121	112	84	73	63	203	167
High School Graduate	228	182	193	149	156 ^a	90	302	265
Some College	132	167	130	148	70	83	223	264
College Graduate	10	26	30	43	4	8	36	68
CURRENT EMPLOYMENT								
Full-Time	168	199	170	179	103	92	272	338
Part-Time	141	102	129	98	88	58	204	155
Unemployed	106	107	84	64	65	53	133	136
Other ³	95	88	83	84	48	41	155	135

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.18B Received Substance Use Treatment in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	1.5	1.4	1.3	1.2	0.9	0.7	2.2	2.2
GENDER								
Male	1.9	1.8	1.8	1.6	1.2	0.9	2.9	2.8
Female	1.0	1.1	0.9	0.8	0.5	0.5	1.5	1.5
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	1.6	1.5	1.3	1.4	0.9	0.8	2.2	2.3
White	1.9	1.7	1.6	1.5	1.1	0.8	2.5	2.6
Black or African American	0.7	0.9	0.7	0.9	0.4	0.5	1.4	1.5
American Indian or Alaska Native	2.7	3.1	3.3	4.4	2.7	3.0	3.3	5.2
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	0.5	0.3	0.4	1.0	0.4	0.3	1.3	1.0
Two or More Races	1.6	2.2	1.6	1.7	1.2	1.4	2.0	3.1
Hispanic or Latino	0.9	1.1	1.4 ^a	0.7	0.6	0.4	2.1	1.7
EDUCATION								
< High School	2.5	2.5	2.0	1.7	1.3	1.3	3.6	3.4
High School Graduate	1.9	1.5	1.6	1.2	1.3 ^a	0.7	2.5	2.2
Some College	1.1	1.3	1.1	1.2	0.6	0.7	1.8	2.1
College Graduate	0.2	0.5	0.6	0.8	0.1	0.1	0.7	1.2
CURRENT EMPLOYMENT								
Full-Time	1.2	1.4	1.3	1.3	0.8	0.6	2.0	2.4
Part-Time	1.5	1.1	1.3	1.0	0.9	0.6	2.1	1.6
Unemployed	2.6	2.8	2.0	1.6	1.6	1.4	3.2	3.5
Other ³	1.3	1.2	1.1	1.1	0.6	0.6	2.1	1.9

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.19A Received Substance Use Treatment in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	1,530	1,778	2,128	2,151	903	1,092	3,055	3,158
GENDER								
Male	833	1,074	1,358	1,447	497	677	1,917	2,095
Female	698	704	770	704	407	414	1,137	1,063
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	1,293	1,536	1,814	1,795	786	953	2,547	2,651
White	1,019	1,122	1,384	1,331	593	644	1,988	1,991
Black or African American	215 ^a	377	271	376	137 ^a	288	381	551
American Indian or Alaska Native	16	14	32	27	16	9	35	33
Native Hawaiian or Other Pacific Islander	7	10	7	*	7	*	14	12
Asian	17	7	61	16	17	7	61	18
Two or More Races	18	6	59	42	15	2	67	46
Hispanic or Latino	238	242	314	355	118	139	508	507
EDUCATION								
< High School	408	465	429	561	226	277	701	834
High School Graduate	465	615	570	665	262	409	874	981
Some College	435	496	572	607	275	291	834	922
College Graduate	223	202	557 ^a	318	141	114	646	421
CURRENT EMPLOYMENT								
Full-Time	324 ^b	561	701	878	160 ^a	291	961 ^a	1,268
Part-Time	254	199	238	207	155	129	373	304
Unemployed	327	192	393 ^a	188	259 ^a	93	502	325
Other ³	625	825	795	878	329 ^a	579	1,219	1,261

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.19B Received Substance Use Treatment in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.8	0.9	1.0	1.0	0.4	0.5	1.5	1.5
GENDER								
Male	0.9	1.1	1.4	1.5	0.5	0.7	2.0	2.1
Female	0.7	0.7	0.7	0.7	0.4	0.4	1.1	1.0
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.7	0.9	1.0	1.0	0.5	0.5	1.5	1.5
White	0.7	0.8	1.0	1.0	0.4	0.5	1.5	1.4
Black or African American	0.9 ^a	1.6	1.2	1.6	0.6 ^a	1.2	1.7	2.4
American Indian or Alaska Native	1.6	1.3	3.2	2.5	1.6	0.8	3.5	3.1
Native Hawaiian or Other Pacific Islander	1.2	1.4	1.2	*	1.2	*	2.4	1.6
Asian	0.2	0.1	0.6	0.1	0.2	0.1	0.6	0.2
Two or More Races	0.6	0.2	2.1	1.5	0.5	0.1	2.4	1.6
Hispanic or Latino	0.8	0.8	1.1	1.2	0.4	0.5	1.8	1.7
EDUCATION								
< High School	1.5	1.7	1.6	2.1	0.8	1.0	2.6	3.1
High School Graduate	0.8	1.1	1.0	1.2	0.5	0.7	1.5	1.7
Some College	0.8	0.9	1.1	1.1	0.5	0.5	1.6	1.7
College Graduate	0.3	0.3	0.8 ^a	0.5	0.2	0.2	1.0	0.6
CURRENT EMPLOYMENT								
Full-Time	0.3 ^b	0.5	0.7	0.8	0.2 ^a	0.3	0.9	1.2
Part-Time	1.1	0.8	1.0	0.9	0.6	0.5	1.5	1.3
Unemployed	4.3	2.8	5.1 ^a	2.7	3.4 ^a	1.3	6.6	4.7
Other ³	1.0	1.2	1.2	1.3	0.5 ^a	0.9	1.9	1.9

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.20A Received Substance Use Treatment in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	2,040	2,274	2,594	2,576	1,207	1,335	3,818	3,922
GENDER								
Male	1,170	1,384	1,671	1,727	708	830	2,427	2,593
Female	870	890	922	849	500	505	1,391	1,330
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	1,733	1,951	2,180	2,169	1,043	1,168	3,157	3,293
White	1,396	1,461	1,689	1,615	810	809	2,484	2,498
Black or African American	251 ^a	422	305	420	156 ^a	313	450	627
American Indian or Alaska Native	22	20	40	36	22	15	43	44
Native Hawaiian or Other Pacific Islander	7	11	7	7	7	4	14	16
Asian	27	14	68	35	25	14	85	38
Two or More Races	30	23	70	55	24	13	82	70
Hispanic or Latino	306	323	414	407	164	168	662	629
EDUCATION								
< High School	547	587	540	645	299	340	904	1,001
High School Graduate	693	797	763	815	418	499	1,176	1,245
Some College	567	663	702	755	345	374	1,056	1,187
College Graduate	233	228	588 ^a	361	145	122	682	489
CURRENT EMPLOYMENT								
Full-Time	492 ^b	760	871	1,057	264 ^a	383	1,234 ^a	1,606
Part-Time	395	301	367	306	242	186	577	459
Unemployed	434	300	477 ^a	252	324 ^a	145	635	461
Other ³	719	913	878	962	377 ^a	621	1,373	1,396

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.20B Received Substance Use Treatment in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.9	0.9	1.1	1.1	0.5	0.6	1.6	1.6
GENDER								
Male	1.0	1.2	1.5	1.5	0.6	0.7	2.1	2.2
Female	0.7	0.7	0.7	0.7	0.4	0.4	1.1	1.1
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.9	1.0	1.1	1.1	0.5	0.6	1.6	1.6
White	0.9	0.9	1.1	1.0	0.5	0.5	1.6	1.6
Black or African American	0.9 ^a	1.5	1.1	1.5	0.6 ^a	1.1	1.6	2.2
American Indian or Alaska Native	1.8	1.6	3.2	2.8	1.8	1.2	3.4	3.4
Native Hawaiian or Other Pacific Islander	0.9	1.2	0.9	0.8	0.9	0.5	1.8	1.7
Asian	0.2	0.1	0.6	0.3	0.2	0.1	0.7	0.3
Two or More Races	0.8	0.6	2.0	1.5	0.7	0.4	2.3	1.9
Hispanic or Latino	0.9	0.9	1.2	1.1	0.5	0.5	1.9	1.7
EDUCATION								
< High School	1.7	1.8	1.7	2.0	0.9	1.1	2.8	3.1
High School Graduate	1.0	1.2	1.1	1.2	0.6	0.7	1.7	1.8
Some College	0.9	1.0	1.1	1.2	0.5	0.6	1.7	1.8
College Graduate	0.3	0.3	0.8 ^a	0.5	0.2	0.2	0.9	0.7
CURRENT EMPLOYMENT								
Full-Time	0.4 ^b	0.6	0.7	0.9	0.2	0.3	1.0 ^a	1.3
Part-Time	1.2	0.9	1.1	0.9	0.7	0.6	1.7	1.4
Unemployed	3.7	2.8	4.0 ^a	2.3	2.7 ^a	1.3	5.4	4.3
Other ³	1.0	1.2	1.2	1.3	0.5 ^a	0.8	1.9	1.9

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.21A Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	2,214	2,440	2,731	2,683	1,320	1,427	4,051	4,149
GEOGRAPHIC DIVISION								
Northeast	471	582	508	515	221	337	822	839
New England	146	137	113	124	54	73	216	211
Middle Atlantic	325	445	395	391	167	264	606	628
Midwest	451	465	534	532	246	262	820	877
East North Central	292	335	354	386	149	181	557	631
West North Central	159	130	180	147	97	81	263	246
South	729	862	932	962	464	511	1,323	1,492
South Atlantic	377	556	465	589	243	323	655	907
East South Central	138	109	131	95	68	58	219	171
West South Central	214	197	335	279	154	130	449	413
West	564	532	757	673	389	317	1,086	942
Mountain	165	186	214	239	59 ^a	122	369	310
Pacific	399	346	543	434	329	196	717	632
COUNTY TYPE								
Large Metro	1,211	1,412	1,470	1,451	733	845	2,198	2,231
Small Metro	610	701	845	902	364	423	1,224	1,353
250K – 1 Mil. Pop.	405	475	635	596	248	269	889	899
< 250K Pop.	205	226	210	306	116	154	336	454
Nonmetro	393	328	417	329	222	159	628	565
Urbanized	147	149	172	152	87	72	263	268
Less Urbanized	232	150	204	135	131	67	315	235
Completely Rural	13	29	40	41	5	20	50	62

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.21B Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.8	0.9	1.0	1.0	0.5	0.5	1.5	1.6
GEOGRAPHIC DIVISION								
Northeast	1.0	1.2	1.1	1.1	0.5	0.7	1.7	1.8
New England	1.2	1.1	0.9	1.0	0.4	0.6	1.7	1.7
Middle Atlantic	0.9	1.3	1.1	1.1	0.5	0.8	1.7	1.8
Midwest	0.8	0.8	0.9	0.9	0.4	0.5	1.5	1.6
East North Central	0.7	0.9	0.9	1.0	0.4	0.5	1.4	1.6
West North Central	0.9	0.8	1.0	0.8	0.6	0.5	1.5	1.4
South	0.7	0.9	1.0	1.0	0.5	0.5	1.4	1.5
South Atlantic	0.7	1.1	0.9	1.1	0.5	0.6	1.3	1.7
East South Central	0.9	0.7	0.8	0.6	0.4	0.4	1.4	1.1
West South Central	0.7	0.6	1.1	0.9	0.5	0.4	1.5	1.3
West	0.9	0.9	1.2	1.1	0.6	0.5	1.8	1.5
Mountain	0.9	1.0	1.1	1.3	0.3 ^a	0.6	2.0	1.6
Pacific	0.9	0.8	1.3	1.0	0.8	0.5	1.7	1.5
COUNTY TYPE								
Large Metro	0.9	1.0	1.0	1.0	0.5	0.6	1.6	1.5
Small Metro	0.8	0.9	1.1	1.2	0.5	0.5	1.6	1.7
250K – 1 Mil. Pop.	0.8	0.9	1.2	1.1	0.5	0.5	1.7	1.7
< 250K Pop.	0.8	0.9	0.8	1.2	0.5	0.6	1.3	1.8
Nonmetro	0.9	0.8	1.0	0.8	0.5	0.4	1.5	1.4
Urbanized	0.8	0.9	0.9	0.9	0.5	0.4	1.4	1.6
Less Urbanized	1.2	0.8	1.1	0.7	0.7	0.3	1.7	1.2
Completely Rural	0.3	0.7	0.9	1.0	0.1	0.5	1.1	1.5

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.22A Substances for Which Last or Current Treatment Was Received among Persons Who Received Substance Use Treatment in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Substance for Which Last or Current Treatment Was Received ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	4,051	4,149	232	227	764	764	3,055	3,158
Marijuana	845	1,034	127	117	268	238	450 ^a	679
Cocaine	584	781	20	8	116	88	448 ^a	685
Heroin	526	618	*	12	128	133	392	473
Hallucinogens	303	339	25	13	90	72	189	254
Inhalants	187	189	12	11	46	35	129	143
Pain Relievers	746	772	29	22	190	183	527	567
Tranquilizers	376	480	19	23	104	98	253	359
Stimulants ²	461	425	21	19	94	97	346	308
Sedatives	154	152	*	11	25	21	123	121
Alcohol	2,513	2,406	105	79	398	377	2,010	1,950

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.22B Substances for Which Last or Current Treatment Was Received among Persons Who Received Substance Use Treatment in the Past Year, by Age Group: Percentages, 2013 and 2014

Substance for Which Last or Current Treatment Was Received ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marijuana	20.9	24.9	54.6	51.5	35.1	31.2	14.7 ^a	21.5
Cocaine	14.4	18.8	8.5	3.6	15.2	11.5	14.7 ^a	21.7
Heroin	13.0	14.9	*	5.3	16.8	17.4	12.8	15.0
Hallucinogens	7.5	8.2	10.6	5.8	11.8	9.4	6.2	8.0
Inhalants	4.6	4.6	5.2	4.7	6.0	4.6	4.2	4.5
Pain Relievers	18.4	18.6	12.5	9.9	24.9	23.9	17.2	18.0
Tranquilizers	9.3	11.6	8.3	10.3	13.6	12.8	8.3	11.4
Stimulants ²	11.4	10.2	9.0	8.5	12.3	12.7	11.3	9.8
Sedatives	3.8	3.7	*	4.7	3.3	2.7	4.0	3.8
Alcohol	62.0	58.0	45.0	34.9	52.1	49.3	65.8	61.8

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.23A Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	2,214	2,440	175	166	509	496	1,530	1,778
Hospital - Inpatient	475	560	26	25	104	114	345	421
Rehabilitation Facility - Inpatient	659	680	29	27	170	167	460	486
Rehabilitation Facility - Outpatient	1,126	1,121	60	43	283	236	783	843
Mental Health Center - Outpatient	625	751	42	35	137	145	446	571
Emergency Room	279	289	19	16	47	57	213	217
Private Doctor's Office	456	476	18	22	100	82	337	372
Self-Help Group	1,213	1,219	50	49	279	274	884	896
Prison/Jail	162	264	8	8	42	39	111 ^a	217

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment refers to treatment received in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.23B Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment in the Past Year, by Age Group: Percentages, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Hospital - Inpatient	21.4	22.9	14.7	15.1	20.5	23.0	22.5	23.7
Rehabilitation Facility - Inpatient	29.7	27.9	16.3	16.2	33.4	33.7	30.0	27.3
Rehabilitation Facility - Outpatient	50.8	46.0	34.4	25.8	55.5	47.5	51.2	47.4
Mental Health Center - Outpatient	28.2	30.8	23.9	21.0	26.8	29.2	29.1	32.1
Emergency Room	12.6	11.9	10.8	9.9	9.2	11.4	13.9	12.2
Private Doctor's Office	20.6	19.5	10.3	13.5	19.7	16.5	22.0	20.9
Self-Help Group	54.8	49.9	28.7	29.7	54.8	55.2	57.7	50.4
Prison/Jail	7.3	10.8	4.8	4.7	8.3	7.9	7.3	12.2

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment refers to treatment received in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.24A Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	2,731	2,683	137	*	466	425	2,128	2,151
Hospital - Inpatient	573	579	29	*	89	87	455	457
Rehabilitation Facility - Inpatient	619	601	31	*	97	87	492	488
Rehabilitation Facility - Outpatient	1,019	1,019	50	*	172	169	798	816
Mental Health Center - Outpatient	645	712	*	*	88	114	518	572
Emergency Room	318	330	*	*	37	27	259	276
Private Doctor's Office	417	460	20	*	55	42	342	387
Self-Help Group	1,649	1,517	51	*	210	217	1,387	1,264
Prison/Jail	156	187	14	*	32	25	109	146

*Low precision; no estimate reported.

NOTE: Received Alcohol Treatment refers to treatment received in order to reduce or stop alcohol use, or for medical problems associated with alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.24B Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: Percentages, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	100.0	100.0	100.0	*	100.0	100.0	100.0	100.0
Hospital - Inpatient	21.0	21.6	21.2	*	19.1	20.4	21.4	21.3
Rehabilitation Facility - Inpatient	22.7	22.4	22.3	*	20.8	20.6	23.1	22.7
Rehabilitation Facility - Outpatient	37.3	38.0	36.2	*	36.8	39.7	37.5	37.9
Mental Health Center - Outpatient	23.6	26.5	*	*	19.0	26.9	24.3	26.6
Emergency Room	11.6	12.3	*	*	8.0	6.4	12.2	12.8
Private Doctor's Office	15.3	17.1	14.6	*	11.8	9.9	16.1	18.0
Self-Help Group	60.4	56.5	37.3	*	45.1	51.0	65.2	58.7
Prison/Jail	5.7	7.0	10.3	*	6.9	6.0	5.1	6.8

*Low precision; no estimate reported.

NOTE: Received Alcohol Treatment refers to treatment received in order to reduce or stop alcohol use, or for medical problems associated with alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.25A Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	4,051	4,149	232	227	764	764	3,055	3,158
Hospital - Inpatient	879	921	43	51	148	154	688	716
Rehabilitation Facility - Inpatient	1,042	1,076	49	47	202	212	791	817
Rehabilitation Facility - Outpatient	1,753	1,731	89	59	349	326	1,314	1,346
Mental Health Center - Outpatient	1,176	1,157	62	51	181	206	933	900
Emergency Room	603	521	40	45	91	85	472	391
Private Doctor's Office	770	780	30	45	126	109	613	626
Self-Help Group	2,292	2,210	80	71	369	402	1,843	1,738
Prison/Jail	263	366	22	21	57	48	184	296

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.25B Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: Percentages, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Hospital - Inpatient	21.7	22.2	18.3	22.4	19.4	20.2	22.5	22.7
Rehabilitation Facility - Inpatient	25.7	25.9	21.1	20.5	26.4	27.8	25.9	25.9
Rehabilitation Facility - Outpatient	43.3	41.7	38.4 ^a	26.0	45.7	42.7	43.0	42.6
Mental Health Center - Outpatient	29.0	27.9	26.7	22.5	23.7	27.0	30.6	28.5
Emergency Room	14.9	12.6	17.3	20.0	11.9	11.1	15.4	12.4
Private Doctor's Office	19.0	18.8	13.0	19.8	16.5	14.3	20.1	19.8
Self-Help Group	56.6	53.3	34.6	31.1	48.3	52.5	60.3	55.0
Prison/Jail	6.5	8.8	9.3	9.3	7.5	6.3	6.0	9.4

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.26A Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Numbers in Thousands, 2013 and 2014

Dependence or Abuse Status	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol (2013)	Both Illicit Drugs and Alcohol (2014)	Illicit Drugs or Alcohol ¹ (2013)	Illicit Drugs or Alcohol ¹ (2014)
TOTAL	2,214	2,440	2,731	2,683	1,320	1,427	4,051	4,149
DEPENDENCE								
None	1,274	1,330	1,634	1,490	770	778	2,422	2,359
Both Illicit Drugs and Alcohol	204	288	182	300	163	276	246	338
Illicit Drugs or Alcohol	941	1,110	1,097	1,193	550	648	1,629	1,789
Illicit Drugs	799	911	486	548	439	475	931	1,093
Alcohol	346	487	794	945	274 ^a	449	944	1,034
DEPENDENCE OR ABUSE								
None	1,119	1,104	1,383	1,128	673	618	2,046	1,895
Both Illicit Drugs and Alcohol	352	403	323	473	265	353	462	578
Illicit Drugs or Alcohol	1,096	1,337	1,348	1,555	648	809	2,005	2,254
Illicit Drugs	917	1,030	583	731	502	562	1,103	1,333
Alcohol	531	709	1,088	1,297	411 ^a	599	1,365	1,500

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.26B Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Percentages, 2013 and 2014

Dependence or Abuse Status	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol (2013)	Both Illicit Drugs and Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.8	0.9	1.0	1.0	0.5	0.5	1.5	1.6
DEPENDENCE								
None	0.5	0.5	0.7	0.6	0.3	0.3	1.0	0.9
Both Illicit Drugs and Alcohol	18.6	26.5	16.6 ^a	27.6	14.9 ^a	25.5	22.4	31.1
Illicit Drugs or Alcohol	8.0	9.3	9.3	10.0	4.7	5.4	13.8	15.0
Illicit Drugs	16.2	18.3	9.9	11.0	8.9	9.5	18.9	21.9
Alcohol	4.3	6.0	9.9	11.7	3.4 ^a	5.6	11.8	12.8
DEPENDENCE OR ABUSE								
None	0.5	0.5	0.6	0.5	0.3	0.3	0.8	0.8
Both Illicit Drugs and Alcohol	13.6	15.5	12.5 ^a	18.3	10.2	13.6	17.8	22.3
Illicit Drugs or Alcohol	5.1	6.2	6.3	7.2	3.0	3.8	9.3	10.5
Illicit Drugs	13.4	14.6	8.5	10.3	7.3	7.9	16.1	18.8
Alcohol	3.1 ^a	4.2	6.3	7.6	2.4 ^a	3.5	7.9	8.8

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.27A Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	1,483	1,604	1,421	1,572	547	684	2,466	2,606
12	*	1	2	4	*	1	2	4
13	6	*	7	2	5	*	9	2
14	4	9	7	3	2	2	11	10
15	15	16	9	11	5	8	21	19
16	33	22	26	12	21 ^a	6	40	28
17	33	26	22	24	16	8	40	46
18	43 ^a	16	29	12	15	4	57 ^a	26
19	41	24	26	26	12	10	56	46
20	29	32	21	21	15	14	37	39
21	40	49	31	24	23	12	55	60
22	40	37	21	36	15	17	52	57
23	52	64	44	47	19	32	79	83
24	39	39	27	36	11	13	57	67
25	56	65	35	44	28	19	63	92
26-29	244	231	187	172	97	101	343	311
30-34	231	224	182	185	98	85	317	327
35-39	130	163	142	118	62	50	209	243
40-44	116	167	121	181	27	78	236	292
45-49	82	95	111	143	20	52	180	186
50-54	66	118	96	128	20	60	142	218
55-59	84	146	53	125	9	59	149	225
60-64	98	17	150	114	28	15	239	115
65 or Older	*	40	72	106	*	37	72	110

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.27B Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.6	0.6	0.5	0.6	0.2	0.3	0.9	1.0
12	*	0.0	0.1	0.1	*	0.0	0.1	0.1
13	0.1	*	0.2	0.0	0.1	*	0.2	0.0
14	0.1	0.2	0.2	0.1	0.0	0.0	0.2	0.2
15	0.4	0.4	0.2	0.3	0.1	0.2	0.5	0.4
16	0.8	0.5	0.6	0.3	0.5 ^a	0.1	0.9	0.7
17	0.8	0.6	0.5	0.6	0.4	0.2	1.0	1.1
18	0.9 ^a	0.4	0.6	0.3	0.3	0.1	1.2 ^a	0.6
19	1.0	0.6	0.6	0.6	0.3	0.3	1.4	1.1
20	0.7	0.7	0.5	0.5	0.3	0.3	0.9	0.9
21	0.9	1.1	0.7	0.5	0.5	0.3	1.3	1.4
22	0.9	0.8	0.5	0.8	0.3	0.4	1.2	1.3
23	1.2	1.5	1.0	1.1	0.4	0.7	1.8	1.9
24	0.9	0.9	0.6	0.8	0.3	0.3	1.3	1.5
25	1.4	1.5	0.8	1.0	0.7	0.4	1.5	2.2
26-29	1.4	1.4	1.1	1.0	0.6	0.6	2.0	1.8
30-34	1.1	1.1	0.9	0.9	0.5	0.4	1.6	1.6
35-39	0.7	0.8	0.7	0.6	0.3	0.3	1.1	1.2
40-44	0.5	0.8	0.6	0.9	0.1 ^a	0.4	1.1	1.4
45-49	0.4	0.5	0.6	0.7	0.1	0.3	0.9	0.9
50-54	0.3	0.5	0.4	0.5	0.1	0.3	0.6	0.9
55-59	0.4	0.7	0.3	0.6	0.0	0.3	0.7	1.1
60-64	0.6	0.1	0.8	0.6	0.2	0.1	1.3	0.7
65 or Older	*	0.1	0.2	0.2	*	0.1	0.2	0.2

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.28A Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	1,483	1,604	1,421	1,572	547	684	2,466	2,606
AGE								
12-17	90	74	73	55	49 ^a	25	122	109
18-25	341	328	233	244	137	122	458	470
26 or Older	1,051	1,201	1,115	1,272	361	536	1,886	2,028
GENDER								
Male	824	1,006	948	1,104	287 ^a	471	1,546	1,695
Female	659	597	473	468	259	213	920	911
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	1,258	1,350	1,203	1,332	478	600	2,069	2,193
White	1,011	1,024	880	999	349	404	1,615	1,688
Black or African American	185	269	202	254	77 ^a	163	310	400
American Indian or Alaska Native	20	20	30	24	18	13	33	32
Native Hawaiian or Other Pacific Islander	6	*	6	*	6	*	6	11
Asian	26	7	43	16	23	7	46	16
Two or More Races	11	23	42	35	5	13	58	45
Hispanic or Latino	224	254	218	240	69	83	397	413

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.28B Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.6	0.6	0.5	0.6	0.2	0.3	0.9	1.0
AGE								
12-17	0.4	0.3	0.3	0.2	0.2 ^a	0.1	0.5	0.4
18-25	1.0	0.9	0.7	0.7	0.4	0.3	1.3	1.3
26 or Older	0.5	0.6	0.5	0.6	0.2	0.3	0.9	1.0
GENDER								
Male	0.6	0.8	0.7	0.9	0.2 ^a	0.4	1.2	1.3
Female	0.5	0.4	0.3	0.3	0.2	0.2	0.7	0.7
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.6	0.6	0.5	0.6	0.2	0.3	0.9	1.0
White	0.6	0.6	0.5	0.6	0.2	0.2	1.0	1.0
Black or African American	0.6	0.8	0.6	0.8	0.2	0.5	1.0	1.3
American Indian or Alaska Native	1.4	1.4	2.2	1.7	1.3	0.9	2.4	2.2
Native Hawaiian or Other Pacific Islander	0.7	*	0.7	*	0.7	*	0.7	1.1
Asian	0.2	0.0	0.3	0.1	0.2	0.0	0.3	0.1
Two or More Races	0.2	0.5	1.0	0.8	0.1	0.3	1.4	1.0
Hispanic or Latino	0.5	0.6	0.5	0.6	0.2	0.2	1.0	1.0

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.29A Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	90	74	73	55	49 ^a	25	122	109
GENDER								
Male	47	30	44 ^a	18	27 ^a	9	68	42
Female	43	44	29	37	22	16	55	66
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	67	56	54	52	34	24	90	88
White	53	46	37	44	25	19	69	74
Black or African American	6	5	8	6	3	1	10	9
American Indian or Alaska Native	3	2	4	1	*	*	5	2
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	*	*	*	*	*	*	*
Two or More Races	3	3	3	2	2	2	4	3
Hispanic or Latino	23	18	19 ^b	3	15 ^a	2	33	21
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	30	22	27	10	18	6	40	30
Female, White, Not Hispanic	23	24	10 ^a	33	7	14	28	44
Male, Black, Not Hispanic	4	3	6	6	1	1	9	7
Female, Black, Not Hispanic	2	2	2	*	2	*	2	2
Male, Hispanic	9	4	6	2	5	2	13	4
Female, Hispanic	14	14	13 ^a	1	9	*	20	17

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.29B Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.4	0.3	0.3	0.2	0.2 ^a	0.1	0.5	0.4
GENDER								
Male	0.4	0.2	0.3 ^a	0.1	0.2 ^a	0.1	0.5	0.3
Female	0.4	0.4	0.2	0.3	0.2	0.1	0.4	0.5
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.3	0.3	0.3	0.3	0.2	0.1	0.5	0.5
White	0.4	0.3	0.3	0.3	0.2	0.1	0.5	0.5
Black or African American	0.2	0.1	0.2	0.2	0.1	0.0	0.3	0.2
American Indian or Alaska Native	2.1	1.2	2.8	0.7	*	*	3.2	1.4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	*	*	*	*	*	*	*
Two or More Races	0.4	0.4	0.3	0.3	0.2	0.3	0.5	0.4
Hispanic or Latino	0.4	0.3	0.3 ^b	0.0	0.3 ^a	0.0	0.6	0.4
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	0.4	0.3	0.4	0.2	0.3	0.1	0.6	0.4
Female, White, Not Hispanic	0.3	0.4	0.2 ^a	0.5	0.1	0.2	0.4	0.7
Male, Black, Not Hispanic	0.2	0.1	0.3	0.3	0.0	0.1	0.5	0.4
Female, Black, Not Hispanic	0.1	0.1	0.1	*	0.1	*	0.1	0.1
Male, Hispanic	0.3	0.1	0.2	0.1	0.2	0.1	0.5	0.1
Female, Hispanic	0.5	0.5	0.5 ^a	0.0	0.3	*	0.7	0.6

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.30A Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	341	328	233	244	137	122	458	470
GENDER								
Male	231	205	177	169	100	85	324	301
Female	110	123	57	75	37	37	134	169
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	298	280	190	226	117	120	388	406
White	265	232	155	172	103	89	334	333
Black or African American	19	28	15	29	4	17	30	41
American Indian or Alaska Native	1	5	5	5	1	*	6	6
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	8	*	6	9	6	*	8	9
Two or More Races	5	14	9	11	3	10	10	15
Hispanic or Latino	42	48	44	18	20 ^a	2	69	64
EDUCATION								
< High School	75	71	58	43	19	24	129	103
High School Graduate	166	126	108	90	79	50	198	172
Some College	94	114	55	81	39	43	113	153
College Graduate	5	17	12 ^a	31	0	5	17 ^a	43
CURRENT EMPLOYMENT								
Full-Time	99	125	83	86	44	47	145	169
Part-Time	89	69	56	67	33	31	122	105
Unemployed	95	78	60	44	44	32	111	101
Other ³	57	55	34	47	16	12	79	95

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.30B Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	1.0	0.9	0.7	0.7	0.4	0.3	1.3	1.3
GENDER								
Male	1.3	1.2	1.0	1.0	0.6	0.5	1.9	1.7
Female	0.6	0.7	0.3	0.4	0.2	0.2	0.8	1.0
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	1.1	1.0	0.7	0.8	0.4	0.4	1.4	1.5
White	1.4	1.2	0.8	0.9	0.5	0.5	1.7	1.7
Black or African American	0.4	0.6	0.3	0.6	0.1	0.3	0.6	0.8
American Indian or Alaska Native	0.6	2.3	2.4	2.7	0.6	*	2.4	3.0
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	0.4	*	0.3	0.5	0.3	*	0.4	0.5
Two or More Races	0.6	1.8	1.2	1.4	0.5	1.3	1.4	2.0
Hispanic or Latino	0.6	0.7	0.6	0.2	0.3 ^a	0.0	1.0	0.9
EDUCATION								
< High School	1.3	1.4	1.0	0.9	0.3	0.5	2.3	2.1
High School Graduate	1.4	1.1	0.9	0.8	0.7	0.4	1.7	1.4
Some College	0.8	0.9	0.5	0.6	0.3	0.3	0.9	1.2
College Graduate	0.1	0.3	0.2	0.5	0.0	0.1	0.3	0.8
CURRENT EMPLOYMENT								
Full-Time	0.7	0.9	0.6	0.6	0.3	0.3	1.1	1.2
Part-Time	0.9	0.7	0.6	0.7	0.3	0.3	1.3	1.1
Unemployed	2.3	2.0	1.5	1.1	1.1	0.8	2.7	2.6
Other ³	0.8	0.8	0.5	0.6	0.2	0.2	1.1	1.3

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.31A Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	1,051	1,201	1,115	1,272	361	536	1,886	2,028
GENDER								
Male	546	772	728	916	160 ^b	377	1,154	1,352
Female	506	430	387	356	201	160	732	675
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	893	1,014	959	1,053	327	456	1,591	1,700
White	694	746	688	783	222	296	1,213	1,282
Black or African American	159	236	180	220	70	145	269	351
American Indian or Alaska Native	*	13	20	18	*	8	22	24
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	10
Asian	17	7	38	7	17	7	38	7
Two or More Races	3	6	30	22	*	1	44	27
Hispanic or Latino	159	187	155	219	34	80	295	328
EDUCATION								
< High School	293	309	192	338	77	132	443	557
High School Graduate	327	431	315	467	128	239	514	663
Some College	315	331	376	321	133	123	603	570
College Graduate	115	130	232	147	22	42	325	237
CURRENT EMPLOYMENT								
Full-Time	211 ^b	386	289	422	63	119	438 ^b	719
Part-Time	167	153	132	123	76	82	229	194
Unemployed	260 ^a	105	215	126	128	25	358	232
Other ³	413	558	479	602	93 ^b	310	862	883

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.31B Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.5	0.6	0.5	0.6	0.2	0.3	0.9	1.0
GENDER								
Male	0.6	0.8	0.8	0.9	0.2 ^b	0.4	1.2	1.4
Female	0.5	0.4	0.4	0.3	0.2	0.1	0.7	0.6
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.5	0.6	0.6	0.6	0.2	0.3	0.9	1.0
White	0.5	0.5	0.5	0.6	0.2	0.2	0.9	0.9
Black or African American	0.7	1.0	0.8	0.9	0.3	0.6	1.2	1.5
American Indian or Alaska Native	*	1.2	2.0	1.7	*	0.8	2.2	2.2
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	1.4
Asian	0.2	0.1	0.4	0.1	0.2	0.1	0.4	0.1
Two or More Races	0.1	0.2	1.1	0.8	*	0.0	1.6	0.9
Hispanic or Latino	0.6	0.6	0.5	0.7	0.1	0.3	1.0	1.1
EDUCATION								
< High School	1.1	1.1	0.7	1.2	0.3	0.5	1.7	2.0
High School Graduate	0.6	0.8	0.5	0.8	0.2	0.4	0.9	1.2
Some College	0.6	0.6	0.7	0.6	0.3	0.2	1.2	1.1
College Graduate	0.2	0.2	0.3	0.2	0.0	0.1	0.5	0.3
CURRENT EMPLOYMENT								
Full-Time	0.2 ^a	0.4	0.3	0.4	0.1	0.1	0.4 ^b	0.7
Part-Time	0.7	0.6	0.5	0.5	0.3	0.3	0.9	0.8
Unemployed	3.4 ^a	1.5	2.8	1.8	1.7	0.4	4.7	3.4
Other ³	0.6	0.8	0.7	0.9	0.1 ^b	0.5	1.3	1.3

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.32A Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	1,392	1,529	1,348	1,517	498	658	2,343	2,498
GENDER								
Male	776	976	904	1,085	260 ^b	462	1,478	1,653
Female	616	553	444	431	238	196	866	845
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	1,191	1,294	1,149	1,280	444	577	1,979	2,105
White	959	978	842	955	325	385	1,547	1,615
Black or African American	179	264	195	249	74 ^a	161	300	392
American Indian or Alaska Native	17	18	26	23	*	12	28	30
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	11
Asian	26	7	43	16	23	7	46	16
Two or More Races	8	20	39	33	3	11	55	42
Hispanic or Latino	201	235	199	237	54	82	365	392
EDUCATION								
< High School	369	380	250	380	96	156	572	660
High School Graduate	493	557	423	557	207	289	713	835
Some College	410	445	431	401	173	167	716	723
College Graduate	121	147	245	178	23	47	343	279
CURRENT EMPLOYMENT								
Full-Time	310 ^b	512	372	508	107	167	583 ^b	888
Part-Time	256	222	188	189	110	113	351	299
Unemployed	355 ^a	183	275	170	172 ^a	57	469	333
Other ³	471	613	513	650	109 ^b	322	941	978

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.32B Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.6	0.6	0.6	0.6	0.2	0.3	1.0	1.0
GENDER								
Male	0.7	0.8	0.8	0.9	0.2 ^a	0.4	1.3	1.4
Female	0.5	0.4	0.4	0.3	0.2	0.2	0.7	0.7
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.6	0.6	0.6	0.6	0.2	0.3	1.0	1.0
White	0.6	0.6	0.5	0.6	0.2	0.2	1.0	1.0
Black or African American	0.6	0.9	0.7	0.9	0.3 ^a	0.6	1.1	1.4
American Indian or Alaska Native	1.3	1.4	2.1	1.8	*	1.0	2.3	2.3
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	1.2
Asian	0.2	0.1	0.4	0.1	0.2	0.1	0.4	0.1
Two or More Races	0.2	0.5	1.1	0.9	0.1	0.3	1.5	1.2
Hispanic or Latino	0.6	0.6	0.6	0.6	0.2	0.2	1.0	1.1
EDUCATION								
< High School	1.1	1.2	0.8	1.2	0.3	0.5	1.8	2.0
High School Graduate	0.7	0.8	0.6	0.8	0.3	0.4	1.0	1.2
Some College	0.6	0.7	0.7	0.6	0.3	0.3	1.1	1.1
College Graduate	0.2	0.2	0.3	0.2	0.0	0.1	0.5	0.4
CURRENT EMPLOYMENT								
Full-Time	0.3 ^b	0.4	0.3	0.4	0.1	0.1	0.5 ^b	0.7
Part-Time	0.8	0.7	0.6	0.6	0.3	0.3	1.0	0.9
Unemployed	3.0 ^a	1.7	2.3	1.6	1.5 ^a	0.5	4.0	3.1
Other ³	0.6	0.8	0.7	0.9	0.1 ^b	0.4	1.3	1.3

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.33A Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	1,483	1,604	1,421	1,572	547	684	2,466	2,606
GEOGRAPHIC DIVISION								
Northeast	345	400	298	390	74 ^a	171	576	647
New England	98	98	57	95	11	41	144	154
Middle Atlantic	248	302	241	295	64	131	432	493
Midwest	299	347	243	332	87 ^a	167	482	541
East North Central	195	247	163	219	56	105	302	381
West North Central	105	100	81	113	31	62	179	161
South	494	551	531	502	209	237	864	860
South Atlantic	275	352	278	338	119	157	463	560
East South Central	108	83	88	51	25	32	180	107
West South Central	110	116	165	113	65	48	221	194
West	344	306	350	347	176	109	545	558
Mountain	99	105	99	122	22	42	187	192
Pacific	245	201	251	225	154	67	358	366
COUNTY TYPE								
Large Metro	822	945	744	887	295	446	1,326	1,450
Small Metro	375	446	458	496	147	179	728	795
250K – 1 Mil. Pop.	231	322	345	348	99	128	495	571
< 250K Pop.	145	124	112	148	48	51	232	224
Nonmetro	285	213	220	188	104	58	412	361
Urbanized	103	100	78	85	31	25	153	165
Less Urbanized	171	96	111	75	72	23	218	161
Completely Rural	10	16	31	28	1	10	40	35

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.33B Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.6	0.6	0.5	0.6	0.2	0.3	0.9	1.0
GEOGRAPHIC DIVISION								
Northeast	0.7	0.8	0.6	0.8	0.2 ^a	0.4	1.2	1.4
New England	0.8	0.8	0.5	0.8	0.1	0.3	1.2	1.2
Middle Atlantic	0.7	0.9	0.7	0.8	0.2	0.4	1.2	1.4
Midwest	0.5	0.6	0.4	0.6	0.2 ^a	0.3	0.9	1.0
East North Central	0.5	0.6	0.4	0.6	0.1	0.3	0.8	1.0
West North Central	0.6	0.6	0.5	0.7	0.2	0.4	1.0	0.9
South	0.5	0.6	0.5	0.5	0.2	0.2	0.9	0.9
South Atlantic	0.5	0.7	0.5	0.6	0.2	0.3	0.9	1.1
East South Central	0.7	0.5	0.6	0.3	0.2	0.2	1.2	0.7
West South Central	0.4	0.4	0.5	0.4	0.2	0.2	0.7	0.6
West	0.6	0.5	0.6	0.6	0.3	0.2	0.9	0.9
Mountain	0.5	0.6	0.5	0.6	0.1	0.2	1.0	1.0
Pacific	0.6	0.5	0.6	0.5	0.4	0.2	0.8	0.8
COUNTY TYPE								
Large Metro	0.6	0.6	0.5	0.6	0.2	0.3	0.9	1.0
Small Metro	0.5	0.6	0.6	0.6	0.2	0.2	0.9	1.0
250K – 1 Mil. Pop.	0.4	0.6	0.6	0.7	0.2	0.2	0.9	1.1
< 250K Pop.	0.6	0.5	0.4	0.6	0.2	0.2	0.9	0.9
Nonmetro	0.7	0.5	0.5	0.5	0.2	0.1	1.0	0.9
Urbanized	0.5	0.6	0.4	0.5	0.2	0.1	0.8	1.0
Less Urbanized	0.9	0.5	0.6	0.4	0.4	0.1	1.1	0.8
Completely Rural	0.2	0.4	0.7	0.7	0.0	0.2	0.9	0.8

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.34A Substances for Which Last or Current Treatment Was Received among Persons Who Received Their Last or Current Substance Use Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Substance for Which Last or Current Treatment Was Received ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	1,680	1,848	*	*	330	337	1,260	1,435
Marijuana	485	574	*	*	152	118	277	401
Cocaine	359	503	*	*	71	52	277	443
Heroin	359	428	*	*	70	90	283	334
Hallucinogens	206	192	*	*	53	32	137	153
Inhalants	119	88	*	*	*	18	78	64
Pain Relievers	421	475	*	*	122	97	278	363
Tranquilizers	233	312	*	*	69	53	151	240
Stimulants ²	247	243	*	*	61	52	172	183
Sedatives	65	88	*	*	23	4	37	79
Alcohol	1,003	1,094	*	*	187	180	758	876

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.34B Substances for Which Last or Current Treatment Was Received among Persons Who Received Their Last or Current Substance Use Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014

Substance for Which Last or Current Treatment Was Received ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	100.0	100.0	*	*	100.0	100.0	100.0	100.0
Marijuana	28.9	31.0	*	*	46.0	35.0	22.0	27.9
Cocaine	21.4	27.2	*	*	21.5	15.6	22.0	30.9
Heroin	21.4	23.2	*	*	21.2	26.6	22.5	23.3
Hallucinogens	12.2	10.4	*	*	16.2	9.5	10.9	10.7
Inhalants	7.1	4.8	*	*	*	5.3	6.2	4.4
Pain Relievers	25.1	25.7	*	*	37.1	28.7	22.0	25.3
Tranquilizers	13.9	16.9	*	*	20.8	15.8	12.0	16.8
Stimulants ²	14.7	13.2	*	*	18.5	15.5	13.7	12.8
Sedatives	3.9	4.7	*	*	6.8 ^a	1.3	2.9	5.5
Alcohol	59.7	59.2	*	*	56.7	53.5	60.2	61.1

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.35A Source of Payment for Last or Current Illicit Drug Treatment among Persons Who Received Their Last or Current Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	1,075	1,180	*	*	245	229	770	896
Private Health Insurance	408	514	*	*	127	126	252	360
Medicare	301	337	*	*	47	45	233	278
Medicaid	303	391	*	*	45	50	240	325
Public Assistance Other Than Medicaid	332	294	*	*	67	50	254	236
Own Savings or Earnings	387	449	*	*	78	83	294	365
Family Members	310	297	*	*	84	*	203	172
Courts	112	153	*	*	40	22	69	119
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	42	40	*	*	10	8	31	31
Employer	16	33	*	*	6	9	10	25

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.35B Source of Payment for Last or Current Illicit Drug Treatment among Persons Who Received Their Last or Current Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	100.0	100.0	*	*	100.0	100.0	100.0	100.0
Private Health Insurance	37.9	43.6	*	*	51.7	55.1	32.7	40.2
Medicare	28.0	28.6	*	*	19.2	19.7	30.3	31.0
Medicaid	28.2	33.2	*	*	18.4	21.8	31.2	36.3
Public Assistance Other Than Medicaid	30.9	24.9	*	*	27.5	21.8	33.0	26.4
Own Savings or Earnings	36.0	38.1	*	*	31.8	36.0	38.2	40.8
Family Members	28.8	25.1	*	*	34.2	*	26.3	19.2
Courts	10.4	13.0	*	*	16.3	9.4	8.9	13.3
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	3.9	3.4	*	*	4.0	3.6	4.1	3.4
Employer	1.5	2.8	*	*	2.4	3.7	1.3	2.7

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.36A Source of Payment for Last or Current Alcohol Treatment among Persons Who Received Their Last or Current Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	864	1,049	*	*	156	*	656	836
Private Health Insurance	359	446	*	*	*	*	*	342
Medicare	186	227	*	*	17	*	*	186
Medicaid	237	261	*	*	16	*	*	218
Public Assistance Other Than Medicaid	176	232	*	*	*	*	137	202
Own Savings or Earnings	367	410	*	*	57	*	*	345
Family Members	179	207	*	*	*	*	106	127
Courts	75	125	*	*	*	*	50	101
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	*	105	*	*	*	*	*	93
Employer	27	52	*	*	7	*	20	45

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.36B Source of Payment for Last or Current Alcohol Treatment among Persons Who Received Their Last or Current Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	100.0	100.0	*	*	100.0	*	100.0	100.0
Private Health Insurance	41.5	42.6	*	*	*	*	*	40.9
Medicare	21.5	21.6	*	*	11.0	*	*	22.3
Medicaid	27.4	24.9	*	*	9.9	*	*	26.0
Public Assistance Other Than Medicaid	20.3	22.1	*	*	*	*	20.9	24.2
Own Savings or Earnings	42.5	39.1	*	*	36.4	*	*	41.3
Family Members	20.7	19.8	*	*	*	*	16.1	15.2
Courts	8.7	11.9	*	*	*	*	7.7	12.1
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	*	10.0	*	*	*	*	*	11.1
Employer	3.1	5.0	*	*	4.7	*	3.0	5.4

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.37A Source of Payment for Last or Current Illicit Drug or Alcohol Treatment among Persons Who Received Their Last or Current Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	1,680	1,848	*	*	330	337	1,260	1,435
Private Health Insurance	701	905	*	*	164	172	493	695
Medicare	450	561	*	*	63	68	363	473
Medicaid	487	626	*	*	56	75	407	527
Public Assistance Other Than Medicaid	487	496	*	*	81	68	389	411
Own Savings or Earnings	683	805	*	*	130	145	531	654
Family Members	387	456	*	*	117	151	237	274
Courts	174	226	*	*	44	30	122	178
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	133	159	*	*	16	15	117	138
Employer	42 ^a	112	*	*	12	12	30 ^a	91

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.37B Source of Payment for Last or Current Illicit Drug or Alcohol Treatment among Persons Who Received Their Last or Current Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	100.0	100.0	*	*	100.0	100.0	100.0	100.0
Private Health Insurance	41.7	49.0	*	*	49.7	51.1	39.1	48.4
Medicare	26.8	30.3	*	*	19.2	20.3	28.8	32.9
Medicaid	29.0	33.9	*	*	17.0	22.2	32.3	36.7
Public Assistance Other Than Medicaid	29.0	26.8	*	*	24.6	20.1	30.9	28.7
Own Savings or Earnings	40.6	43.6	*	*	39.3	43.0	42.2	45.6
Family Members	23.0	24.7	*	*	35.4	45.0	18.8	19.1
Courts	10.4	12.2	*	*	13.3	8.9	9.7	12.4
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	7.9	8.6	*	*	4.8	4.5	9.2	9.6
Employer	2.5 ^a	6.0	*	*	3.6	3.6	2.3 ^a	6.4

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.38A Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	1,483	1,604	*	*	341	328	1,051	1,201
Hospital - Inpatient	475	560	*	*	104	114	345	421
Rehabilitation Facility - Inpatient	659	680	*	*	170	167	460	486
Rehabilitation Facility - Outpatient	1,126	1,121	*	*	283	236	783	843
Mental Health Center - Outpatient	625	751	*	*	137	145	446	571
Emergency Room ²	267	273	*	*	41	57	210	201
Private Doctor's Office ²	329	350	*	*	71	57	247	279
Self-Help Group ²	905	913	*	*	213	221	667	662
Prison/Jail ²	130	203	*	*	35	31	90	169

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.38B Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	100.0	100.0	*	*	100.0	100.0	100.0	100.0
Hospital - Inpatient	32.0	34.9	*	*	30.6	34.7	32.8	35.0
Rehabilitation Facility - Inpatient	44.4	42.4	*	*	49.9	51.0	43.7	40.4
Rehabilitation Facility - Outpatient	75.9	69.9	*	*	83.0 ^a	71.9	74.5	70.2
Mental Health Center - Outpatient	42.1	46.8	*	*	40.1	44.2	42.4	47.5
Emergency Room ²	18.0	17.0	*	*	12.0	17.2	20.0	16.7
Private Doctor's Office ²	22.2	21.8	*	*	20.7	17.4	23.5	23.2
Self-Help Group ²	61.1	56.9	*	*	62.5	67.3	63.5	55.1
Prison/Jail ²	8.8	12.7	*	*	10.3	9.5	8.6	14.1

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.39A Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	1,421	1,572	*	*	233	244	1,115	1,272
Hospital - Inpatient	573	579	*	*	89	87	455	457
Rehabilitation Facility - Inpatient	619	601	*	*	97	87	492	488
Rehabilitation Facility - Outpatient	1,019	1,019	*	*	172	169	798	816
Mental Health Center - Outpatient	645	712	*	*	88	114	518	572
Emergency Room ²	304	314	*	*	27	26	259	262
Private Doctor's Office ²	276	315	*	*	*	27	227	265
Self-Help Group ²	907	940	*	*	122	150	756	766
Prison/Jail ²	92	134	*	*	27	*	52	103

*Low precision; no estimate reported.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.39B Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	100.0	100.0	*	*	100.0	100.0	100.0	100.0
Hospital - Inpatient	40.3	36.8	*	*	38.1	35.5	40.8	36.0
Rehabilitation Facility - Inpatient	43.6	38.2	*	*	41.6	35.8	44.1	38.4
Rehabilitation Facility - Outpatient	71.7	64.8	*	*	73.6	69.0	71.6	64.1
Mental Health Center - Outpatient	45.3	45.3	*	*	37.9	46.8	46.4	45.0
Emergency Room ²	21.4	20.0	*	*	11.5	10.6	23.3	20.6
Private Doctor's Office ²	19.4	20.0	*	*	*	11.1	20.4	20.8
Self-Help Group ²	63.8	59.8	*	*	52.4	61.2	67.8	60.2
Prison/Jail ²	6.4	8.5	*	*	11.6	*	4.7	8.1

*Low precision; no estimate reported.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.40A Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	2,466	2,606	122	*	458	470	1,886	2,028
Hospital - Inpatient	879	921	43	*	148	154	688	716
Rehabilitation Facility - Inpatient	1,042	1,076	49	*	202	212	791	817
Rehabilitation Facility - Outpatient	1,753	1,731	*	*	349	326	1,314	1,346
Mental Health Center - Outpatient	1,176	1,157	62	*	181	206	933	900
Emergency Room ²	574	499	*	*	73	84	468	375
Private Doctor's Office ²	522	561	21	*	82	75	419	456
Self-Help Group ²	1,505	1,554	46	*	269	302	1,190	1,204
Prison/Jail ²	189	280	*	*	47	35	127	228

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.40B Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	100.0	100.0	100.0	*	100.0	100.0	100.0	100.0
Hospital - Inpatient	35.6	35.3	34.7	*	32.3	32.8	36.5	35.3
Rehabilitation Facility - Inpatient	42.2	41.3	40.0	*	44.0	45.2	42.0	40.3
Rehabilitation Facility - Outpatient	71.1	66.4	*	*	76.3	69.4	69.7	66.4
Mental Health Center - Outpatient	47.7	44.4	50.7	*	39.5	43.9	49.5	44.4
Emergency Room ²	23.3	19.1	*	*	15.9	17.8	24.8	18.5
Private Doctor's Office ²	21.2	21.5	17.3	*	18.0	16.0	22.2	22.5
Self-Help Group ²	61.0	59.6	37.6	*	58.7	64.3	63.1	59.4
Prison/Jail ²	7.7	10.8	*	*	10.2	7.5	6.7	11.2

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.41A Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Numbers in Thousands, 2013 and 2014

Dependence or Abuse Status	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol (2013)	Both Illicit Drugs and Alcohol (2014)	Illicit Drugs or Alcohol ¹ (2013)	Illicit Drugs or Alcohol ¹ (2014)
TOTAL	1,483	1,604	1,421	1,572	547	684	2,466	2,606
DEPENDENCE								
None	768	782	734	762	261	318	1,301	1,274
Both Illicit Drugs and Alcohol	160	208	132	212	103	188	191	265
Illicit Drugs or Alcohol	715	821	687	810	286	366	1,165	1,332
Illicit Drugs	639	713	298	331	240	280	730	830
Alcohol	235	316	522	690	148 ^a	273	626	768
DEPENDENCE OR ABUSE								
None	673	643	626	567	224	251	1,126	998
Both Illicit Drugs and Alcohol	278	301	236	318	171	238	349	429
Illicit Drugs or Alcohol	809	961	795	1,005	323	433	1,340	1,609
Illicit Drugs	726	800	360	401	275	315	846	952
Alcohol	362	462	671 ^a	921	219 ^a	355	842	1,086

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.41B Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Percentages, 2013 and 2014

Dependence or Abuse Status	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol (2013)	Both Illicit Drugs and Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.6	0.6	0.5	0.6	0.2	0.3	0.9	1.0
DEPENDENCE								
None	0.3	0.3	0.3	0.3	0.1	0.1	0.5	0.5
Both Illicit Drugs and Alcohol	14.6	19.1	12.0	19.5	9.4	17.3	17.4	24.4
Illicit Drugs or Alcohol	6.0	6.9	5.8	6.8	2.4	3.1	9.9	11.1
Illicit Drugs	13.0	14.3	6.1	6.6	4.9	5.6	14.8	16.6
Alcohol	2.9	3.9	6.5	8.6	1.9 ^a	3.4	7.8	9.5
DEPENDENCE OR ABUSE								
None	0.3	0.3	0.3	0.2	0.1	0.1	0.5	0.4
Both Illicit Drugs and Alcohol	10.7	11.6	9.1	12.3	6.6	9.2	13.5	16.6
Illicit Drugs or Alcohol	3.8	4.5	3.7	4.7	1.5	2.0	6.2	7.5
Illicit Drugs	10.6	11.3	5.3	5.7	4.0	4.5	12.3	13.5
Alcohol	2.1	2.7	3.9 ^a	5.4	1.3 ^a	2.1	4.9 ^a	6.4

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.42A Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse of Specific Substances: Numbers in Thousands, 2013 and 2014

Past Year Dependence or Abuse ¹	Illicit Drugs ² (2013)	Illicit Drugs ² (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ² (2013)	Both Illicit Drugs and Alcohol ² (2014)	Illicit Drugs or Alcohol ^{2,3} (2013)	Illicit Drugs or Alcohol ^{2,3} (2014)
ILLCIT DRUGS²	726	800	360	401	275	315	846	952
Marijuana and Hashish	318	267	173	201	133	135	389	387
Cocaine	162	255	98	147	74	143	190	289
Heroin	*	270	58	80	53	76	*	276
Hallucinogens	*	18	*	15	*	15	*	18
Inhalants	*	*	*	*	*	*	*	*
Nonmedical Use of Psychotherapeutics ^{4,5}	389	413	167	174	136	146	422	478
Pain Relievers	345	348	138	148	125	125	360	392
Tranquilizers	*	126	*	31	*	23	*	155
Stimulants ⁴	78	81	45	34	35	33	88	101
Sedatives	*	*	*	*	*	*	*	*
ALCOHOL	362	462	671 ^a	921	219 ^a	355	842	1,086
BOTH ILLICIT DRUGS AND ALCOHOL²	278	301	236	318	171	238	349	429
ILLCIT DRUGS OR ALCOHOL²	809	961	795	1,005	323	433	1,340	1,609

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

³ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

⁴ Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

⁵ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.42B Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse of Specific Substances: Percentages, 2013 and 2014

Past Year Dependence or Abuse ¹	Illicit Drugs ² (2013)	Illicit Drugs ² (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ² (2013)	Both Illicit Drugs and Alcohol ² (2014)	Illicit Drugs or Alcohol ^{2,3} (2013)	Illicit Drugs or Alcohol ^{2,3} (2014)
ILLICIT DRUGS²	10.6	11.3	5.3	5.7	4.0	4.5	12.3	13.5
Marijuana and Hashish	7.6	6.4	4.1	4.8	3.2	3.2	9.2	9.3
Cocaine	18.9	28.0	11.5	16.2	8.7	15.6	22.2	31.6
Heroin	*	46.1	11.2	13.6	10.2	13.0	*	47.2
Hallucinogens	*	7.2	*	6.3	*	6.0	*	7.5
Inhalants	*	*	*	*	*	*	*	*
Nonmedical Use of Psychotherapeutics ^{4,5}	17.1	17.1	7.3	7.2	6.0	6.0	18.5	19.8
Pain Relievers	18.4	18.1	7.4	7.7	6.6	6.5	19.1	20.4
Tranquilizers	*	26.8	*	6.6	*	4.8	*	32.7
Stimulants ⁴	16.7	17.0	9.6	7.1	7.4	7.0	18.8	21.3
Sedatives	*	*	*	*	*	*	*	*
ALCOHOL	2.1	2.7	3.9 ^a	5.4	1.3 ^a	2.1	4.9 ^a	6.4
BOTH ILLICIT DRUGS AND ALCOHOL²	10.7	11.6	9.1	12.3	6.6	9.2	13.5	16.6
ILLICIT DRUGS OR ALCOHOL²	3.8	4.5	3.7	4.7	1.5	2.0	6.2	7.5

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

³ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

⁴ Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

⁵ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.43A Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year, by Detailed Age Category: Numbers in Thousands and Percentages, 2013 and 2014

Age Category	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	7,608	7,881	1,483	1,604	6,126	6,278	19.5	20.3
12	21	35	*	1	21	33	*	*
13	48	42	6	*	42	42	*	*
14	110	77	4	9	106 ^a	68	*	*
15	188	187	15	16	173	172	8.1	8.3
16	266	278	33	22	233	257	12.4	7.8
17	274	276	33	26	241	249	11.9	9.5
18	382	321	43 ^a	16	339	305	11.3	5.0
19	346	294	41	24	305	270	11.9	8.3
20	352	303	29	32	323	270	8.4	10.7
21	380	323	40	49	341	274	10.4	15.2
22	332	380	40	37	292	343	11.9	9.9
23	314	339	52	64	262	275	16.6	18.9
24	313 ^a	228	39	39	274 ^a	188	12.6	17.3
25	251	272	56	65	194	207	22.4	23.8
26-29	824	952	244	231	580	721	29.6	24.3
30-34	976	825	231	224	745	601	23.6	27.1
35-39	513	571	130	163	384	408	*	28.5
40-44	412	495	116	167	296	328	*	*
45-49	329	413	82	95	246	318	*	*
50-54	437	542	66	118	371	424	*	*
55-59	205	351	84	146	121	205	*	*
60-64	171	178	98	17	72	161	*	*
65 or Older	163	200	*	40	163	160	*	*

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.43B Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	2.9	3.0	0.6	0.6	2.3	2.4	19.5	20.3
12	0.6	0.9	*	0.0	0.6	0.9	*	*
13	1.1	1.0	0.1	*	1.0	1.0	*	*
14	2.5	1.8	0.1	0.2	2.4	1.6	*	*
15	4.6	4.4	0.4	0.4	4.2	4.0	8.1	8.3
16	6.2	6.4	0.8	0.5	5.5	5.9	12.4	7.8
17	6.6	6.6	0.8	0.6	5.8	6.0	11.9	9.5
18	8.1	7.0	0.9 ^a	0.4	7.2	6.6	11.3	5.0
19	8.3	7.1	1.0	0.6	7.3	6.5	11.9	8.3
20	8.2	6.9	0.7	0.7	7.5	6.2	8.4	10.7
21	9.0	7.4	0.9	1.1	8.0	6.3	10.4	15.2
22	7.4	8.4	0.9	0.8	6.5	7.6	11.9	9.9
23	7.2	7.7	1.2	1.5	6.0	6.3	16.6	18.9
24	7.1 ^a	5.3	0.9	0.9	6.2 ^a	4.3	12.6	17.3
25	6.1	6.4	1.4	1.5	4.7	4.9	22.4	23.8
26-29	4.8	5.6	1.4	1.4	3.4	4.2	29.6	24.3
30-34	4.8	4.0	1.1	1.1	3.7	2.9	23.6	27.1
35-39	2.7	2.9	0.7	0.8	2.0	2.1	*	28.5
40-44	1.9	2.5	0.5	0.8	1.4	1.6	*	*
45-49	1.6	2.0	0.4	0.5	1.2	1.5	*	*
50-54	1.9	2.3	0.3	0.5	1.6	1.8	*	*
55-59	1.0	1.7	0.4	0.7	0.6	1.0	*	*
60-64	1.0	1.0	0.6	0.1	0.4	0.9	*	*
65 or Older	0.4	0.4	*	0.1	0.4	0.4	*	*

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.44A Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	7,608	7,881	1,483	1,604	6,126	6,278	19.5	20.3
AGE								
12-17	908	894	90	74	817	820	10.0	8.3
18-25	2,671	2,460	341	328	2,330	2,132	12.8	13.3
26 or Older	4,030	4,527	1,051	1,201	2,978	3,326	26.1	26.5
GENDER								
Male	4,714	4,901	824	1,006	3,890	3,895	17.5	20.5
Female	2,895	2,980	659	597	2,236	2,383	22.8	20.0
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	6,418	6,569	1,258	1,350	5,160	5,219	19.6	20.6
White	4,766	4,677	1,011	1,024	3,755	3,653	21.2	21.9
Black or African American	1,151	1,409	185	269	966	1,140	16.0	19.1
American Indian or Alaska Native	86	49	20	20	66	29	*	*
Native Hawaiian or Other Pacific Islander	32	61	6	*	27	53	*	*
Asian	176	161	26	7	151	155	*	*
Two or More Races	206	211	11	23	195	189	5.2	10.7
Hispanic or Latino	1,190	1,312	224	254	966	1,059	18.9	19.3

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.44B Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	2.9	3.0	0.6	0.6	2.3	2.4	19.5	20.3
AGE								
12-17	3.6	3.6	0.4	0.3	3.3	3.3	10.0	8.3
18-25	7.7	7.0	1.0	0.9	6.7	6.1	12.8	13.3
26 or Older	2.0	2.2	0.5	0.6	1.5	1.6	26.1	26.5
GENDER								
Male	3.7	3.8	0.6	0.8	3.1	3.0	17.5	20.5
Female	2.1	2.2	0.5	0.4	1.7	1.7	22.8	20.0
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	2.9	3.0	0.6	0.6	2.3	2.3	19.6	20.6
White	2.8	2.7	0.6	0.6	2.2	2.1	21.2	21.9
Black or African American	3.7	4.4	0.6	0.8	3.1	3.6	16.0	19.1
American Indian or Alaska Native	6.2	3.4	1.4	1.4	4.7 ^a	2.0	*	*
Native Hawaiian or Other Pacific Islander	3.7	5.9	0.7	*	3.0	5.2	*	*
Asian	1.3	1.2	0.2	0.0	1.1	1.1	*	*
Two or More Races	4.8	4.8	0.2	0.5	4.6	4.3	5.2	10.7
Hispanic or Latino	2.9	3.1	0.5	0.6	2.3	2.5	18.9	19.3

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.45A Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	6,700	6,987	1,392	1,529	5,308	5,457	20.8	21.9
GENDER								
Male	4,214	4,437	776	976	3,438	3,460	18.4	22.0
Female	2,486	2,550	616	553	1,871	1,997	24.8	21.7
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	5,722	5,925	1,191	1,294	4,531	4,631	20.8	21.8
White	4,263	4,195	959	978	3,304	3,217	22.5	23.3
Black or African American	1,010	1,288	179	264	831	1,024	17.7	20.5
American Indian or Alaska Native	76	44	17	18	60	26	*	*
Native Hawaiian or Other Pacific Islander	27	57	*	*	24	50	*	*
Asian	164	153	26	7	138	146	*	*
Two or More Races	181	188	8	20	174	169	*	10.4
Hispanic or Latino	978	1,062	201	235	777	826	*	22.2
EDUCATION								
< High School	1,434	1,406	369	380	1,066	1,025	25.7	27.1
High School Graduate	2,245	2,181	493	557	1,752	1,624	22.0	25.5
Some College	2,147	2,269	410	445	1,738	1,824	19.1	19.6
College Graduate	874 ^a	1,132	121	147	753	985	13.8	13.0
CURRENT EMPLOYMENT								
Full-Time	2,667 ^a	3,141	310 ^b	512	2,357	2,629	11.6 ^a	16.3
Part-Time	1,257	1,236	256	222	1,001	1,014	20.4	18.0
Unemployed	1,074 ^a	820	355 ^a	183	718	637	33.1	22.3
Other ¹	1,702	1,790	471	613	1,232	1,177	27.7	34.3

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.45B Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	2.8	2.9	0.6	0.6	2.2	2.3	20.8	21.9
GENDER								
Male	3.7	3.8	0.7	0.8	3.0	3.0	18.4	22.0
Female	2.0	2.0	0.5	0.4	1.5	1.6	24.8	21.7
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	2.8	2.9	0.6	0.6	2.2	2.3	20.8	21.8
White	2.7	2.7	0.6	0.6	2.1	2.1	22.5	23.3
Black or African American	3.6	4.6	0.6	0.9	3.0	3.6	17.7	20.5
American Indian or Alaska Native	6.2	3.4	1.3	1.4	4.8 ^a	2.0	*	*
Native Hawaiian or Other Pacific Islander	3.5	6.2	*	*	3.1	5.4	*	*
Asian	1.4	1.2	0.2	0.1	1.1	1.2	*	*
Two or More Races	5.1	5.2	0.2	0.5	4.9	4.6	*	10.4
Hispanic or Latino	2.7	2.9	0.6	0.6	2.2	2.2	*	22.2
EDUCATION								
< High School	4.5	4.4	1.1	1.2	3.3	3.2	25.7	27.1
High School Graduate	3.2	3.2	0.7	0.8	2.5	2.4	22.0	25.5
Some College	3.4	3.5	0.6	0.7	2.7	2.8	19.1	19.6
College Graduate	1.2	1.5	0.2	0.2	1.0	1.3	13.8	13.0
CURRENT EMPLOYMENT								
Full-Time	2.2	2.6	0.3 ^b	0.4	2.0	2.1	11.6 ^a	16.3
Part-Time	3.7	3.7	0.8	0.7	3.0	3.1	20.4	18.0
Unemployed	9.1	7.6	3.0 ^a	1.7	6.1	5.9	33.1	22.3
Other ¹	2.3	2.4	0.6	0.8	1.7	1.6	27.7	34.3

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.46A Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

Geographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
							(2013)	(2014)
TOTAL	7,608	7,881	1,483	1,604	6,126	6,278	19.5	20.3
GEOGRAPHIC DIVISION								
Northeast	1,450	1,680	345	400	1,105	1,280	23.8	23.8
New England	456	450	98	98	359	351	*	21.9
Middle Atlantic	994	1,230	248	302	746	929	24.9	24.5
Midwest	1,482	1,495	299	347	1,182	1,148	20.2	23.2
East North Central	1,045	1,103	195	247	851	856	18.6	22.4
West North Central	436	392	105	100	331	293	*	*
South	2,692	2,692	494	551	2,198	2,142	18.3	20.5
South Atlantic	1,507	1,545	275	352	1,232	1,193	18.3	22.8
East South Central	510	383	108	83	402	301	*	21.6
West South Central	675	764	110	116	565	648	16.3	15.2
West	1,985	2,014	344	306	1,641	1,708	17.3	15.2
Mountain	666	617	99	105	567	512	*	17.0
Pacific	1,319	1,397	245	201	1,073	1,196	18.6	14.4
COUNTY TYPE								
Large Metro	4,160	4,590	822	945	3,338	3,645	19.8	20.6
Small Metro	2,398	2,343	375	446	2,023	1,897	15.6	19.0
250K – 1 Mil. Pop.	1,551	1,555	231	322	1,321	1,233	14.9	20.7
< 250K Pop.	847	788	145	124	702	664	17.1	15.7
Nonmetro	1,049	948	285	213	764	736	27.2	22.4
Urbanized	512	455	103	100	409	355	20.2	22.0
Less Urbanized	455	418	171	96	283	322	*	23.0
Completely Rural	83	75	10	16	72	59	*	*

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.46B Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	2.9	3.0	0.6	0.6	2.3	2.4	19.5	20.3
GEOGRAPHIC DIVISION								
Northeast	3.1	3.5	0.7	0.8	2.3	2.7	23.8	23.8
New England	3.7	3.6	0.8	0.8	2.9	2.8	*	21.9
Middle Atlantic	2.8	3.5	0.7	0.9	2.1	2.6	24.9	24.5
Midwest	2.6	2.6	0.5	0.6	2.1	2.0	20.2	23.2
East North Central	2.7	2.8	0.5	0.6	2.2	2.2	18.6	22.4
West North Central	2.5	2.3	0.6	0.6	1.9	1.7	*	*
South	2.8	2.7	0.5	0.6	2.3	2.2	18.3	20.5
South Atlantic	2.9	3.0	0.5	0.7	2.4	2.3	18.3	22.8
East South Central	3.3	2.5	0.7	0.5	2.6	1.9	*	21.6
West South Central	2.2	2.5	0.4	0.4	1.8	2.1	16.3	15.2
West	3.2	3.2	0.6	0.5	2.7	2.7	17.3	15.2
Mountain	3.6	3.2	0.5	0.6	3.0	2.7	*	17.0
Pacific	3.1	3.2	0.6	0.5	2.5	2.8	18.6	14.4
COUNTY TYPE								
Large Metro	2.9	3.1	0.6	0.6	2.4	2.5	19.8	20.6
Small Metro	3.0	3.0	0.5	0.6	2.6	2.5	15.6	19.0
250K – 1 Mil. Pop.	2.9	3.0	0.4	0.6	2.5	2.4	14.9	20.7
< 250K Pop.	3.3	3.1	0.6	0.5	2.8	2.7	17.1	15.7
Nonmetro	2.5	2.3	0.7	0.5	1.8	1.8	27.2	22.4
Urbanized	2.7	2.6	0.5	0.6	2.2	2.1	20.2	22.0
Less Urbanized	2.4	2.1	0.9	0.5	1.5	1.6	*	23.0
Completely Rural	1.8	1.8	0.2	0.4	1.6	1.4	*	*

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.47A Perceived Need for Illicit Drug Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug Problem, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment¹ (2013)	Felt Need for Treatment¹ (2014)	Felt Need and Made Effort to Get Treatment¹ (2013)	Felt Need and Made Effort to Get Treatment¹ (2014)	Felt Need and Made No Effort to Get Treatment¹ (2013)	Felt Need and Made No Effort to Get Treatment¹ (2014)	Did Not Feel Need for Treatment¹ (2013)	Did Not Feel Need for Treatment¹ (2014)
TOTAL	6,126	6,278	395	399	148	145	247	254	5,731	5,879
AGE										
12-17	817	820	24	29	7	20	17	9	793	791
18-25	2,330	2,132	88	85	33	28	55	58	2,242	2,046
26 or Older	2,978	3,326	283	284	108	97	175	187	2,695	3,042
GENDER										
Male	3,890	3,895	257	195	103	70	154	125	3,633	3,699
Female	2,236	2,383	138	203	46	74	93	129	2,097	2,180

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but have not received treatment for an illicit drug problem at a specialty facility.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.47B Perceived Need for Illicit Drug Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug Problem, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment¹ (2013)	Felt Need for Treatment¹ (2014)	Felt Need and Made Effort to Get Treatment¹ (2013)	Felt Need and Made Effort to Get Treatment¹ (2014)	Felt Need and Made No Effort to Get Treatment¹ (2013)	Felt Need and Made No Effort to Get Treatment¹ (2014)	Did Not Feel Need for Treatment¹ (2013)	Did Not Feel Need for Treatment¹ (2014)
TOTAL	100.0	100.0	6.4	6.4	2.4	2.3	4.0	4.0	93.6	93.6
AGE										
12-17	100.0	100.0	2.9	3.6	0.9	2.4	2.1	1.1	97.1	96.4
18-25	100.0	100.0	3.8	4.0	1.4	1.3	2.3	2.7	96.2	96.0
26 or Older	100.0	100.0	9.5	8.5	3.6	2.9	5.9	5.6	90.5	91.5
GENDER										
Male	100.0	100.0	6.6	5.0	2.6	1.8	4.0	3.2	93.4	95.0
Female	100.0	100.0	6.2	8.5	2.0	3.1	4.1	5.4	93.8	91.5

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but have not received treatment for an illicit drug problem at a specialty facility.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.48A Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	18,049	17,645	1,421	1,572	16,628	16,073	7.9	8.9
AGE								
12-17	735	699	73	55	662	644	10.0	7.9
18-25	4,633	4,361	233	244	4,400	4,117	5.0	5.6
26 or Older	12,681	12,584	1,115	1,272	11,567	11,312	8.8	10.1
GENDER								
Male	11,600	11,403	948	1,104	10,652	10,299	8.2	9.7
Female	6,449	6,242	473	468	5,976	5,774	7.3	7.5
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	14,926	14,605	1,203	1,332	13,723	13,273	8.1	9.1
White	11,970	11,477	880	999	11,090	10,478	7.4	8.7
Black or African American	1,781	1,977	202	254	1,579	1,723	11.4	12.9
American Indian or Alaska Native	207	215	30	24	177	191	*	11.4
Native Hawaiian or Other Pacific Islander	*	66	6	*	*	62	*	*
Asian	505	518	43	16	462	502	*	3.1
Two or More Races	382	353	42	35	341	318	10.9	*
Hispanic or Latino	3,124	3,039	218	240	2,905	2,800	7.0	7.9

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.48B Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	6.9	6.7	0.5	0.6	6.3	6.1	7.9	8.9
AGE								
12-17	3.0	2.8	0.3	0.2	2.7	2.6	10.0	7.9
18-25	13.3	12.5	0.7	0.7	12.6	11.8	5.0	5.6
26 or Older	6.3	6.1	0.5	0.6	5.7	5.5	8.8	10.1
GENDER								
Male	9.1	8.9	0.7	0.9	8.4	8.0	8.2	9.7
Female	4.8	4.6	0.3	0.3	4.4	4.2	7.3	7.5
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	6.7	6.6	0.5	0.6	6.2	6.0	8.1	9.1
White	7.0	6.7	0.5	0.6	6.5	6.2	7.4	8.7
Black or African American	5.7	6.2	0.6	0.8	5.1	5.4	11.4	12.9
American Indian or Alaska Native	14.8	14.9	2.2	1.7	12.6	13.2	*	11.4
Native Hawaiian or Other Pacific Islander	*	6.4	0.7	*	*	6.1	*	*
Asian	3.8	3.7	0.3	0.1	3.5	3.6	*	3.1
Two or More Races	8.9	8.0	1.0	0.8	7.9	7.2	10.9	*
Hispanic or Latino	7.6	7.2	0.5	0.6	7.0	6.6	7.0	7.9

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.49A Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	17,314	16,946	1,348	1,517	15,966	15,429	7.8	8.9
GENDER								
Male	11,267	11,085	904	1,085	10,363	10,000	8.0	9.8
Female	6,047	5,861	444	431	5,604	5,429	7.3	7.4
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	14,347	14,075	1,149	1,280	13,198	12,795	8.0	9.1
White	11,482	11,031	842	955	10,639	10,076	7.3	8.7
Black or African American	1,731	1,920	195	249	1,537	1,671	11.2	12.9
American Indian or Alaska Native	198	210	26	23	172	187	*	11.1
Native Hawaiian or Other Pacific Islander	*	64	*	*	*	60	*	*
Asian	499	506	43	16	456	489	*	3.2
Two or More Races	363	344	39	33	324	312	10.7	*
Hispanic or Latino	2,967	2,870	199	237	2,768	2,634	6.7	8.3
EDUCATION								
< High School	2,329	2,309	250	380	2,079	1,929	10.7	16.5
High School Graduate	4,613	4,422	423	557	4,190	3,865	9.2	12.6
Some College	5,087	5,165	431	401	4,656	4,763	8.5	7.8
College Graduate	5,285	5,050	245	178	5,041	4,872	4.6	3.5
CURRENT EMPLOYMENT								
Full-Time	10,019	9,875	372	508	9,647	9,367	3.7	5.1
Part-Time	2,580	2,631	188	189	2,392	2,442	7.3	7.2
Unemployed	1,340	1,216	275	170	1,064	1,047	20.5	13.9
Other ¹	3,376	3,223	513	650	2,862	2,574	15.2	20.2

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.49B Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	7.3	7.1	0.6	0.6	6.7	6.4	7.8	8.9
GENDER								
Male	9.8	9.6	0.8	0.9	9.1	8.6	8.0	9.8
Female	4.9	4.7	0.4	0.3	4.6	4.4	7.3	7.4
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	7.1	6.9	0.6	0.6	6.5	6.3	8.0	9.1
White	7.3	7.0	0.5	0.6	6.8	6.4	7.3	8.7
Black or African American	6.2	6.8	0.7	0.9	5.5	5.9	11.2	12.9
American Indian or Alaska Native	15.9	16.5	2.1	1.8	13.8	14.6	*	11.1
Native Hawaiian or Other Pacific Islander	*	6.9	*	*	*	6.5	*	*
Asian	4.1	4.0	0.4	0.1	3.8	3.9	*	3.2
Two or More Races	10.3	9.4	1.1	0.9	9.2	8.5	10.7	*
Hispanic or Latino	8.3	7.8	0.6	0.6	7.8	7.2	6.7	8.3
EDUCATION								
< High School	7.2	7.2	0.8	1.2	6.5	6.0	10.7	16.5
High School Graduate	6.6	6.4	0.6	0.8	6.0	5.6	9.2	12.6
Some College	8.0	7.9	0.7	0.6	7.3	7.3	8.5	7.8
College Graduate	7.3	6.9	0.3	0.2	7.0	6.6	4.6	3.5
CURRENT EMPLOYMENT								
Full-Time	8.4	8.0	0.3	0.4	8.1	7.6	3.7	5.1
Part-Time	7.6	8.0	0.6	0.6	7.1	7.4	7.3	7.2
Unemployed	11.3	11.3	2.3	1.6	9.0	9.7	20.5	13.9
Other ¹	4.6	4.4	0.7	0.9	3.9	3.5	15.2	20.2

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.50A Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment¹ (2013)	Felt Need for Treatment¹ (2014)	Felt Need and Made Effort to Get Treatment¹ (2013)	Felt Need and Made Effort to Get Treatment¹ (2014)	Felt Need and Made No Effort to Get Treatment¹ (2013)	Felt Need and Made No Effort to Get Treatment¹ (2014)	Did Not Feel Need for Treatment¹ (2013)	Did Not Feel Need for Treatment¹ (2014)
TOTAL	16,628	16,073	554	459	201	125	353	334	16,074	15,614
AGE										
12-17	662	644	13	8	7	*	6 ^a	1	649	636
18-25	4,400	4,117	77	72	38	30	39	42	4,322	4,045
26 or Older	11,567	11,312	464	380	156	88	308	291	11,103	10,933
GENDER										
Male	10,652	10,299	400 ^a	226	152	74	248	152	10,252	10,074
Female	5,976	5,774	154	234	49	52	105	182	5,822	5,540

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.50B Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment ¹ (2013)	Felt Need for Treatment ¹ (2014)	Felt Need and Made Effort to Get Treatment ¹ (2013)	Felt Need and Made Effort to Get Treatment ¹ (2014)	Felt Need and Made No Effort to Get Treatment ¹ (2013)	Felt Need and Made No Effort to Get Treatment ¹ (2014)	Did Not Feel Need for Treatment ¹ (2013)	Did Not Feel Need for Treatment ¹ (2014)
TOTAL	100.0	100.0	3.3	2.9	1.2	0.8	2.1	2.1	96.7	97.1
AGE										
12-17	100.0	100.0	1.9	1.2	1.0	*	0.9 ^a	0.2	98.1	98.8
18-25	100.0	100.0	1.8	1.8	0.9	0.7	0.9	1.0	98.2	98.2
26 or Older	100.0	100.0	4.0	3.4	1.4	0.8	2.7	2.6	96.0	96.6
GENDER										
Male	100.0	100.0	3.8 ^a	2.2	1.4	0.7	2.3	1.5	96.2 ^a	97.8
Female	100.0	100.0	2.6	4.0	0.8	0.9	1.8	3.1	97.4	96.0

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.51A Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	22,687	22,478	2,466	2,606	20,221	19,872	10.9	11.6
AGE								
12-17	1,340	1,284	122	109	1,218	1,175	9.1	8.5
18-25	6,146	5,845	458	470	5,689	5,375	7.4	8.0
26 or Older	15,200	15,349	1,886	2,028	13,314	13,321	12.4	13.2
GENDER								
Male	14,476	14,364	1,546	1,695	12,931	12,668	10.7	11.8
Female	8,210	8,115	920	911	7,290	7,204	11.2	11.2
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	18,909	18,681	2,069	2,193	16,841	16,488	10.9	11.7
White	14,967	14,407	1,615	1,688	13,351	12,719	10.8	11.7
Black or African American	2,478	2,814	310	400	2,168	2,413	12.5	14.2
American Indian or Alaska Native	232	236	33	32	199	204	*	13.6
Native Hawaiian or Other Pacific Islander	100	109	6	11	*	98	*	*
Asian	631	623	46	16	585	607	*	2.6
Two or More Races	501	492	58	45	442	447	11.7	9.2
Hispanic or Latino	3,777	3,797	397	413	3,380	3,384	10.5	10.9

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.51B Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	8.6	8.5	0.9	1.0	7.7	7.5	10.9	11.6
AGE								
12-17	5.4	5.2	0.5	0.4	4.9	4.7	9.1	8.5
18-25	17.7	16.7	1.3	1.3	16.4	15.4	7.4	8.0
26 or Older	7.5	7.5	0.9	1.0	6.6	6.5	12.4	13.2
GENDER								
Male	11.4	11.2	1.2	1.3	10.2	9.9	10.7	11.8
Female	6.1	5.9	0.7	0.7	5.4	5.3	11.2	11.2
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	8.6	8.4	0.9	1.0	7.6	7.4	10.9	11.7
White	8.8	8.5	1.0	1.0	7.9	7.5	10.8	11.7
Black or African American	7.9	8.9	1.0	1.3	6.9	7.6	12.5	14.2
American Indian or Alaska Native	16.6	16.4	2.4	2.2	14.2	14.2	*	13.6
Native Hawaiian or Other Pacific Islander	11.5	10.7	0.7	1.1	*	9.6	*	*
Asian	4.7	4.5	0.3	0.1	4.4	4.4	*	2.6
Two or More Races	11.7	11.1	1.4	1.0	10.3	10.1	11.7	9.2
Hispanic or Latino	9.2	8.9	1.0	1.0	8.2	8.0	10.5	10.9

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.52A Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands and Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	21,346	21,194	2,343	2,498	19,003	18,697	11.0	11.8
GENDER								
Male	13,785	13,710	1,478	1,653	12,307	12,056	10.7	12.1
Female	7,561	7,485	866	845	6,696	6,640	11.4	11.3
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	17,874	17,724	1,979	2,105	15,896	15,619	11.1	11.9
White	14,167	13,659	1,547	1,615	12,620	12,044	10.9	11.8
Black or African American	2,314	2,666	300	392	2,015	2,274	12.9	14.7
American Indian or Alaska Native	219	229	28	30	191	199	*	13.0
Native Hawaiian or Other Pacific Islander	*	105	*	11	*	94	*	*
Asian	613	602	46	16	567	586	*	2.7
Two or More Races	467	464	55	42	413	422	11.7	9.1
Hispanic or Latino	3,472	3,470	365	392	3,107	3,078	10.5	11.3
EDUCATION								
< High School	3,255	3,209	572	660	2,683	2,549	17.6	20.6
High School Graduate	5,822	5,700	713	835	5,110	4,865	12.2	14.6
Some College	6,424	6,550	716	723	5,707	5,827	11.2	11.0
College Graduate	5,845	5,735	343	279	5,503	5,456	5.9	4.9
CURRENT EMPLOYMENT								
Full-Time	11,561	11,765	583 ^b	888	10,978	10,877	5.0 ^b	7.5
Part-Time	3,291	3,345	351	299	2,940	3,046	10.7	8.9
Unemployed	1,998	1,749	469	333	1,529	1,416	23.5	19.0
Other ¹	4,497	4,336	941	978	3,556	3,357	20.9	22.6

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.52B Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	9.0	8.8	1.0	1.0	8.0	7.8	11.0	11.8
GENDER								
Male	12.0	11.8	1.3	1.4	10.8	10.4	10.7	12.1
Female	6.1	6.0	0.7	0.7	5.4	5.3	11.4	11.3
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	8.9	8.7	1.0	1.0	7.9	7.7	11.1	11.9
White	9.1	8.7	1.0	1.0	8.1	7.7	10.9	11.8
Black or African American	8.3	9.5	1.1	1.4	7.2	8.1	12.9	14.7
American Indian or Alaska Native	17.6	17.9	2.3	2.3	15.4	15.6	*	13.0
Native Hawaiian or Other Pacific Islander	*	11.4	*	1.2	*	10.3	*	*
Asian	5.1	4.8	0.4	0.1	4.7	4.6	*	2.7
Two or More Races	13.2	12.7	1.5	1.2	11.7	11.6	11.7	9.1
Hispanic or Latino	9.7	9.4	1.0	1.1	8.7	8.4	10.5	11.3
EDUCATION								
< High School	10.1	10.0	1.8	2.0	8.3	7.9	17.6	20.6
High School Graduate	8.3	8.3	1.0	1.2	7.3	7.1	12.2	14.6
Some College	10.1	10.0	1.1	1.1	9.0	8.9	11.2	11.0
College Graduate	8.1	7.8	0.5	0.4	7.6	7.4	5.9	4.9
CURRENT EMPLOYMENT								
Full-Time	9.7	9.6	0.5 ^b	0.7	9.2	8.8	5.0 ^b	7.5
Part-Time	9.8	10.1	1.0	0.9	8.7	9.2	10.7	8.9
Unemployed	16.9	16.2	4.0	3.1	13.0	13.1	23.5	19.0
Other ¹	6.2	5.9	1.3	1.3	4.9	4.6	20.9	22.6

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.53A Perceived Need for Illicit Drug or Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug or Alcohol Problem, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment¹ (2013)	Felt Need for Treatment¹ (2014)	Felt Need and Made Effort to Get Treatment¹ (2013)	Felt Need and Made Effort to Get Treatment¹ (2014)	Felt Need and Made No Effort to Get Treatment¹ (2013)	Felt Need and Made No Effort to Get Treatment¹ (2014)	Did Not Feel Need for Treatment¹ (2013)	Did Not Feel Need for Treatment¹ (2014)
TOTAL	20,221	19,872	908	798	316	231	592	567	19,313	19,073
AGE										
12-17	1,218	1,175	31	32	11	21	20	10	1,187	1,143
18-25	5,689	5,375	159	168	63	58	96	110	5,530	5,207
26 or Older	13,314	13,321	718	599	242	152	477	447	12,596	12,723
GENDER										
Male	12,931	12,668	622 ^a	392	224	119	399	274	12,308	12,276
Female	7,290	7,204	286	406	92	113	193	293	7,005	6,797

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs or alcohol, but have not received treatment for an illicit drug or alcohol problem at a specialty facility.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug or alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.53B Perceived Need for Illicit Drug or Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug or Alcohol Problem, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment¹ (2013)	Felt Need for Treatment¹ (2014)	Felt Need and Made Effort to Get Treatment¹ (2013)	Felt Need and Made Effort to Get Treatment¹ (2014)	Felt Need and Made No Effort to Get Treatment¹ (2013)	Felt Need and Made No Effort to Get Treatment¹ (2014)	Did Not Feel Need for Treatment¹ (2013)	Did Not Feel Need for Treatment¹ (2014)
TOTAL	100.0	100.0	4.5	4.0	1.6	1.2	2.9	2.9	95.5	96.0
AGE										
12-17	100.0	100.0	2.5	2.7	0.9	1.8	1.6	0.9	97.5	97.3
18-25	100.0	100.0	2.8	3.1	1.1	1.1	1.7	2.0	97.2	96.9
26 or Older	100.0	100.0	5.4	4.5	1.8	1.1	3.6	3.4	94.6	95.5
GENDER										
Male	100.0	100.0	4.8 ^a	3.1	1.7	0.9	3.1	2.2	95.2 ^a	96.9
Female	100.0	100.0	3.9	5.6	1.3	1.6	2.7	4.1	96.1	94.4

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs or alcohol, but have not received treatment for an illicit drug or alcohol problem at a specialty facility.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug or alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.54A Detailed Reasons for Not Receiving Illicit Drug Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Numbers in Thousands, Annual Averages Based on 2011-2014

Reason Did Not Receive Illicit Drug Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	467	171	297
No Health Coverage and Could Not Afford Cost	182	67	115
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	48	19	30
No Transportation/Inconvenient	37	20	17
No Program Having Type of Treatment	33	20	13
Not Ready to Stop Using	136	33	103
No Openings in a Program	30	*	8
Did Not Know Where to Go for Treatment	64	19	45
Might Cause Neighbors/Community to Have Negative Opinion	84	14	70
Might Have Negative Effect on Job	81	19	62
Did Not Feel Need for Treatment at the Time	12	6	7
Could Handle the Problem Without Treatment	37	14	23
Treatment Would Not Help	15	0	15
Did Not Have Time	32	10	22
Did Not Want Others to Find Out	25	4	20
Some Other Reason	12	6	7

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but have not received treatment for an illicit drug problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 5.54B Detailed Reasons for Not Receiving Illicit Drug Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Percentages, Annual Averages Based on 2011-2014

Reason Did Not Receive Illicit Drug Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	100.0	100.0	100.0
No Health Coverage and Could Not Afford Cost	39.0	39.2	38.8
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	10.4	11.0	10.0
No Transportation/Inconvenient	7.9	11.5	5.9
No Program Having Type of Treatment	7.1	11.9	4.3
Not Ready to Stop Using	29.1	19.4	34.7
No Openings in a Program	6.5	*	2.8
Did Not Know Where to Go for Treatment	13.8	11.4	15.2
Might Cause Neighbors/Community to Have Negative Opinion	18.0	8.3	23.6
Might Have Negative Effect on Job	17.3	11.1	20.9
Did Not Feel Need for Treatment at the Time	2.6	3.3	2.3
Could Handle the Problem Without Treatment	7.9	8.3	7.6
Treatment Would Not Help	3.3	0.1	5.1
Did Not Have Time	7.0	6.0	7.5
Did Not Want Others to Find Out	5.3	2.6	6.9
Some Other Reason	2.6	3.2	2.2

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but have not received treatment for an illicit drug problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 5.55A Detailed Reasons for Not Receiving Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Alcohol Treatment at a Specialty Facility and Who Felt a Need for Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Numbers in Thousands, Annual Averages Based on 2011-2014

Reason Did Not Receive Alcohol Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	546	159	386
No Health Coverage and Could Not Afford Cost	143	*	93
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	42	23	20
No Transportation/Inconvenient	35	*	21
No Program Having Type of Treatment	51	*	31
Not Ready to Stop Using	283	46	237
No Openings in a Program	10	*	2
Did Not Know Where to Go for Treatment	41	*	25
Might Cause Neighbors/Community to Have Negative Opinion	31	*	24
Might Have Negative Effect on Job	42	13	29
Did Not Feel Need for Treatment at the Time	20	10	11
Could Handle the Problem Without Treatment	37	*	24
Treatment Would Not Help	7	1	6
Did Not Have Time	18	*	14
Did Not Want Others to Find Out	13	1	13
Some Other Reason	18	7	12

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 5.55B Detailed Reasons for Not Receiving Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Alcohol Treatment at a Specialty Facility and Who Felt a Need for Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Percentages, Annual Averages Based on 2011-2014

Reason Did Not Receive Alcohol Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	100.0	100.0	100.0
No Health Coverage and Could Not Afford Cost	26.2	*	24.0
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	7.8	14.2	5.1
No Transportation/Inconvenient	6.5	*	5.4
No Program Having Type of Treatment	9.3	*	8.0
Not Ready to Stop Using	51.9	28.9	61.4
No Openings in a Program	1.8	*	0.5
Did Not Know Where to Go for Treatment	7.5	*	6.4
Might Cause Neighbors/Community to Have Negative Opinion	5.7	*	6.2
Might Have Negative Effect on Job	7.7	8.0	7.5
Did Not Feel Need for Treatment at the Time	3.7	6.1	2.8
Could Handle the Problem Without Treatment	6.7	*	6.3
Treatment Would Not Help	1.2	0.5	1.6
Did Not Have Time	3.3	*	3.6
Did Not Want Others to Find Out	2.4	0.4	3.3
Some Other Reason	3.4	4.1	3.1

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 5.56A Detailed Reasons for Not Receiving Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug or Alcohol Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug or Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Numbers in Thousands, Annual Averages Based on 2011-2014

Reason Did Not Receive Illicit Drug or Alcohol Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	931	294	638
No Health Coverage and Could Not Afford Cost	287	104	183
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	76	32	44
No Transportation/Inconvenient	69	29	40
No Program Having Type of Treatment	71	30	41
Not Ready to Stop Using	384	71	312
No Openings in a Program	35	25	9
Did Not Know Where to Go for Treatment	97	31	66
Might Cause Neighbors/Community to Have Negative Opinion	104	17	87
Might Have Negative Effect on Job	108	22	86
Did Not Feel Need for Treatment at the Time	34	13	21
Could Handle the Problem Without Treatment	67	19	48
Treatment Would Not Help	21	1	20
Did Not Have Time	49	11	38
Did Not Want Others to Find Out	38	5	33
Some Other Reason	25	11	14

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs or alcohol, but have not received treatment for an illicit drug or alcohol problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug or alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 5.56B Detailed Reasons for Not Receiving Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug or Alcohol Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug or Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Percentages, Annual Averages Based on 2011-2014

Reason Did Not Receive Illicit Drug or Alcohol Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	100.0	100.0	100.0
No Health Coverage and Could Not Afford Cost	30.8	35.3	28.7
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	8.2	11.0	6.9
No Transportation/Inconvenient	7.5	10.0	6.3
No Program Having Type of Treatment	7.6	10.3	6.4
Not Ready to Stop Using	41.2	24.2	49.0
No Openings in a Program	3.7	8.7	1.5
Did Not Know Where to Go for Treatment	10.4	10.6	10.3
Might Cause Neighbors/Community to Have Negative Opinion	11.1	5.7	13.6
Might Have Negative Effect on Job	11.6	7.6	13.4
Did Not Feel Need for Treatment at the Time	3.7	4.6	3.3
Could Handle the Problem Without Treatment	7.2	6.3	7.5
Treatment Would Not Help	2.3	0.5	3.1
Did Not Have Time	5.3	3.9	5.9
Did Not Want Others to Find Out	4.1	1.6	5.2
Some Other Reason	2.7	3.8	2.2

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs or alcohol, but have not received treatment for an illicit drug or alcohol problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug or alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 5.1C Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2013 and 2014

Illicit Drug Dependence or Abuse Status ¹	Total		ALCOHOL DEPENDENCE OR ABUSE STATUS							
			No Dependence or Abuse		Dependence or Abuse					
	Total				Dependence		Abuse			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0	0	415	362	415	362	286	246	307	250
NO DEPENDENCE OR ABUSE	239	220	460	389	401	336	270	227	298	232
DEPENDENCE OR ABUSE	239	220	199	160	140	138	101	106	98	86
Dependence	211	186	183	139	117	115	87	93	78	61
Abuse	113	126	80	94	78	76	51	45	58	61

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.1D Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2013 and 2014

Illicit Drug Dependence or Abuse Status ¹	Total		ALCOHOL DEPENDENCE OR ABUSE STATUS							
			No Dependence or Abuse		Dependence or Abuse					
	Total				Dependence		Abuse			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.00	0.00	0.16	0.14	0.16	0.14	0.11	0.09	0.12	0.09
NO DEPENDENCE OR ABUSE	0.09	0.08	0.18	0.15	0.15	0.13	0.10	0.09	0.11	0.09
DEPENDENCE OR ABUSE	0.09	0.08	0.08	0.06	0.05	0.05	0.04	0.04	0.04	0.03
Dependence	0.08	0.07	0.07	0.05	0.04	0.04	0.03	0.04	0.03	0.02
Abuse	0.04	0.05	0.03	0.04	0.03	0.03	0.02	0.02	0.02	0.02

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.1P Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Illicit Drug Dependence or Abuse Status ¹	ALCOHOL DEPENDENCE OR ABUSE STATUS									
	Total		No Dependence or Abuse		Dependence or Abuse					
					Total		Dependence		Abuse	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	N/A		0.3824		0.3824		0.9627		0.2422	
NO DEPENDENCE OR ABUSE	0.6367		0.6153		0.3828		0.9877		0.2331	
DEPENDENCE OR ABUSE	0.6367		0.4857		0.8997		0.9336		0.9231	
Dependence	0.9361		0.9024		0.9721		0.8575		0.8634	
Abuse	0.4377		0.2249		0.8615		0.8755		0.7245	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.2C Substance Dependence or Abuse for Specific Substances in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Past Year Dependence or Abuse	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	239	220	41	46	88	92	219	187
Marijuana and Hashish	183	156	38	40	76	78	164	126
Cocaine	81	87	6	9	28	24	76	84
Heroin	62	63	4	7	23	24	56	56
Hallucinogens	40	34	10	11	23	20	30	23
Inhalants	26	19	8	10	13	6	22	16
Nonmedical Use of Psychotherapeutics ^{2,3}	144	122	17	23	45	45	134	112
Pain Relievers	133	113	15	19	40	41	126	105
Tranquilizers	56	52	9	12	21	17	50	49
Stimulants ²	74	51	9	13	24	19	69	48
Sedatives	27	35	4	10	10	8	25	32
ALCOHOL	415	362	36	42	123	120	392	324
BOTH ILLICIT DRUGS AND ALCOHOL¹	140	138	21	29	58	58	126	117
ILLICIT DRUGS OR ALCOHOL¹	460	389	51	55	138	134	429	348

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.2D Substance Dependence or Abuse for Specific Substances in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Past Year Dependence or Abuse	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	0.09	0.08	0.17	0.18	0.25	0.26	0.11	0.09
Marijuana and Hashish	0.07	0.06	0.15	0.16	0.22	0.22	0.08	0.06
Cocaine	0.03	0.03	0.02	0.03	0.08	0.07	0.04	0.04
Heroin	0.02	0.02	0.02	0.03	0.07	0.07	0.03	0.03
Hallucinogens	0.02	0.01	0.04	0.04	0.07	0.06	0.01	0.01
Inhalants	0.01	0.01	0.03	0.04	0.04	0.02	0.01	0.01
Nonmedical Use of Psychotherapeutics ^{2,3}	0.05	0.05	0.07	0.09	0.13	0.13	0.07	0.05
Pain Relievers	0.05	0.04	0.06	0.08	0.11	0.12	0.06	0.05
Tranquilizers	0.02	0.02	0.04	0.05	0.06	0.05	0.02	0.02
Stimulants ²	0.03	0.02	0.03	0.05	0.07	0.05	0.03	0.02
Sedatives	0.01	0.01	0.01	0.04	0.03	0.02	0.01	0.02
ALCOHOL	0.16	0.14	0.14	0.17	0.35	0.34	0.19	0.16
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.05	0.05	0.09	0.12	0.17	0.16	0.06	0.06
ILLICIT DRUGS OR ALCOHOL¹	0.18	0.15	0.20	0.22	0.40	0.38	0.21	0.17

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.2P Substance Dependence or Abuse for Specific Substances in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Past Year Dependence or Abuse	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLCIT DRUGS¹	0.6367		0.9051		0.0515		0.1406	
Marijuana and Hashish	0.7610		0.3803		0.0989		0.4144	
Cocaine	0.6800		0.6439		0.1936		0.4133	
Heroin	0.4789		0.3710		0.6425		0.3743	
Hallucinogens	0.5210		0.5718		0.3735		0.9287	
Inhalants	0.2391		0.8930		0.0297		0.9094	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.5527		0.0264		0.0843		0.3609	
Pain Relievers	0.9157		0.1711		0.3197		0.7931	
Tranquilizers	0.5586		0.4089		0.1039		0.2717	
Stimulants ²	0.9782		0.2344		0.2259		0.7911	
Sedatives	0.3350		0.0920		0.4471		0.3958	
ALCOHOL	0.3824		0.7517		0.1249		0.7025	
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.8997		0.4807		0.0397		0.4687	
ILLCIT DRUGS OR ALCOHOL¹	0.6153		0.5021		0.0749		0.8487	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.3C Substance Dependence or Abuse in the Past Year, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol ¹ (2013)	Illicit Drugs or Alcohol ¹ (2014)
TOTAL	239	220	415	362	460	389
12	5	10	4	4	6	10
13	9	10	5	8	10	13
14	16	12	11	10	17	14
15	19	22	15	19	23	24
16	21	26	18	24	25	32
17	21	24	22	25	29	31
18	35	34	38	40	48	52
19	33	31	37	36	46	44
20	31	33	42	43	48	49
21	35	30	42	53	51	57
22	31	35	45	48	51	55
23	30	31	52	47	58	54
24	31	26	45	40	51	45
25	25	29	41	48	44	53
26-29	87	75	171	107	187	120
30-34	103	68	165	139	176	146
35-39	67	60	132	98	142	109
40-44	56	52	132	99	141	108
45-49	47	45	117	100	122	102
50-54	126	85	146	146	188	162
55-59	45	62	116	119	120	130
60-64	33	51	116	92	118	101
65 or Older	50	60	157	129	163	141

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.3D Substance Dependence or Abuse in the Past Year, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol ¹ (2013)	Illicit Drugs or Alcohol ¹ (2014)
TOTAL	0.09	0.08	0.16	0.14	0.18	0.15
12	0.14	0.26	0.11	0.10	0.16	0.26
13	0.22	0.25	0.11	0.20	0.24	0.31
14	0.36	0.29	0.26	0.24	0.38	0.34
15	0.46	0.51	0.37	0.43	0.53	0.55
16	0.47	0.56	0.41	0.53	0.56	0.69
17	0.50	0.56	0.52	0.58	0.65	0.70
18	0.70	0.71	0.71	0.81	0.86	1.02
19	0.77	0.72	0.81	0.82	0.98	0.98
20	0.70	0.73	0.90	0.92	1.00	1.04
21	0.76	0.64	0.92	1.07	1.06	1.10
22	0.67	0.74	0.91	0.95	1.02	1.06
23	0.67	0.66	1.07	0.99	1.17	1.09
24	0.68	0.60	0.93	0.86	1.01	0.95
25	0.60	0.67	0.90	1.05	0.95	1.13
26-29	0.49	0.43	0.90	0.59	0.96	0.64
30-34	0.49	0.32	0.76	0.62	0.80	0.65
35-39	0.35	0.30	0.67	0.47	0.72	0.52
40-44	0.26	0.26	0.59	0.47	0.62	0.51
45-49	0.23	0.22	0.56	0.47	0.58	0.48
50-54	0.54	0.35	0.62	0.60	0.79	0.66
55-59	0.22	0.30	0.56	0.57	0.58	0.62
60-64	0.19	0.29	0.63	0.52	0.64	0.57
65 or Older	0.12	0.13	0.36	0.29	0.37	0.31

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.3P Substance Dependence or Abuse in the Past Year, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol ¹ (2013)	Illicit Drugs or Alcohol ¹ (2014)
TOTAL	0.6367		0.3824		0.6153	
12	0.2263		0.7309		0.3237	
13	0.9312		0.3043		0.7998	
14	0.1593		0.1656		0.2265	
15	0.6342		0.6956		0.1782	
16	0.7263		0.6164		0.5838	
17	0.9454		0.5895		0.4506	
18	0.2513		0.2263		0.4665	
19	0.3454		0.0358		0.1137	
20	0.2641		0.8850		0.4307	
21	0.0878		0.3244		0.8207	
22	0.3662		0.3951		0.7642	
23	0.8159		0.2713		0.5836	
24	0.0273		0.0466		0.0121	
25	0.8708		0.2484		0.3315	
26-29	0.0809		0.0760		0.2375	
30-34	0.0336		0.3598		0.1888	
35-39	0.6836		0.9220		0.9584	
40-44	0.3960		0.6467		0.4929	
45-49	0.1571		0.7374		0.9145	
50-54	0.5251		0.2107		0.3054	
55-59	0.0875		0.2093		0.0703	
60-64	0.1699		0.2609		0.5792	
65 or Older	0.9237		0.7929		0.8216	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.4C Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	239	220	415	362	460	389
AGE						
12-17	41	46	36	42	51	55
18-25	88	92	123	120	138	134
26 or Older	219	187	392	324	429	348
GENDER						
Male	172	172	342	287	364	313
Female	170	125	243	191	277	216
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	224	200	362	322	409	350
White	204	159	339	288	387	310
Black or African American	91	106	120	118	138	134
American Indian or Alaska Native	17	10	45	46	46	47
Native Hawaiian or Other Pacific Islander	9	21	*	20	35	28
Asian	43	34	74	58	83	68
Two or More Races	35	32	58	48	62	55
Hispanic or Latino	83	97	191	157	197	173

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.4D Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.09	0.08	0.16	0.14	0.18	0.15
AGE						
12-17	0.17	0.18	0.14	0.17	0.20	0.22
18-25	0.25	0.26	0.35	0.34	0.40	0.38
26 or Older	0.11	0.09	0.19	0.16	0.21	0.17
GENDER						
Male	0.14	0.13	0.27	0.22	0.29	0.24
Female	0.13	0.09	0.18	0.14	0.20	0.16
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.10	0.09	0.16	0.14	0.18	0.16
White	0.12	0.09	0.20	0.17	0.23	0.18
Black or African American	0.29	0.34	0.39	0.37	0.44	0.42
American Indian or Alaska Native	1.05	0.69	2.89	2.55	2.89	2.54
Native Hawaiian or Other Pacific Islander	1.11	1.98	*	2.01	3.80	2.70
Asian	0.32	0.25	0.55	0.43	0.62	0.50
Two or More Races	0.83	0.73	1.32	1.03	1.43	1.20
Hispanic or Latino	0.20	0.23	0.46	0.37	0.48	0.41

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.4P Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.6367		0.3824		0.6153	
AGE						
12-17	0.9051		0.7517		0.5021	
18-25	0.0515		0.1249		0.0749	
26 or Older	0.1406		0.7025		0.8487	
GENDER						
Male	0.6823		0.5799		0.7683	
Female	0.7975		0.4513		0.6481	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.8489		0.5045		0.6259	
White	0.5442		0.2645		0.2338	
Black or African American	0.0725		0.2303		0.0665	
American Indian or Alaska Native	0.0915		0.6506		0.7629	
Native Hawaiian or Other Pacific Islander	0.4418		*		0.7729	
Asian	0.7222		0.9890		0.8465	
Two or More Races	0.8821		0.7162		0.9918	
Hispanic or Latino	0.4511		0.5105		0.8663	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.5C Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	41	46	36	42	51	55
GENDER						
Male	32	32	24	27	38	38
Female	26	32	26	31	33	38
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	35	36	32	34	43	45
White	31	32	30	33	39	42
Black or African American	16	16	10	11	18	18
American Indian or Alaska Native	3	2	2	2	3	2
Native Hawaiian or Other Pacific Islander	1	*	1	*	2	*
Asian	6	4	3	6	7	7
Two or More Races	4	5	4	2	5	5
Hispanic or Latino	20	27	16	23	24	31
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	23	22	19	22	27	29
Female, White, Not Hispanic	21	23	22	26	26	30
Male, Black, Not Hispanic	14	13	6	7	14	13
Female, Black, Not Hispanic	9	10	8	8	11	12
Male, Hispanic	18	20	13	16	21	23
Female, Hispanic	10	19	11	16	14	22

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.5D Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.17	0.18	0.14	0.17	0.20	0.22
GENDER						
Male	0.25	0.25	0.19	0.21	0.30	0.30
Female	0.21	0.26	0.21	0.25	0.27	0.31
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.18	0.19	0.16	0.18	0.22	0.23
White	0.22	0.24	0.22	0.24	0.27	0.30
Black or African American	0.45	0.45	0.29	0.30	0.49	0.50
American Indian or Alaska Native	1.63	1.15	1.43	1.10	1.87	1.34
Native Hawaiian or Other Pacific Islander	1.44	*	1.41	*	1.69	*
Asian	0.46	0.33	0.27	0.49	0.53	0.59
Two or More Races	0.59	0.66	0.55	0.30	0.72	0.71
Hispanic or Latino	0.36	0.47	0.30	0.41	0.44	0.55
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.32	0.32	0.27	0.31	0.38	0.41
Female, White, Not Hispanic	0.31	0.33	0.33	0.37	0.38	0.44
Male, Black, Not Hispanic	0.75	0.69	0.34	0.40	0.77	0.71
Female, Black, Not Hispanic	0.52	0.58	0.46	0.47	0.64	0.70
Male, Hispanic	0.61	0.66	0.44	0.55	0.71	0.78
Female, Hispanic	0.37	0.67	0.40	0.59	0.48	0.77

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.5P Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.9051		0.7517		0.5021	
GENDER						
Male	0.6044		0.9846		0.5509	
Female	0.6911		0.6507		0.7147	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4433		0.3951		0.2498	
White	0.7986		0.3485		0.3911	
Black or African American	0.3839		0.4868		0.5365	
American Indian or Alaska Native	0.5072		0.4934		0.3081	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.6107		0.3685		0.7902	
Two or More Races	0.6399		0.0235		0.3503	
Hispanic or Latino	0.3625		0.4302		0.5920	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.9214		0.8340		0.9144	
Female, White, Not Hispanic	0.7880		0.2899		0.2634	
Male, Black, Not Hispanic	0.6019		0.3589		0.6632	
Female, Black, Not Hispanic	0.4611		0.8880		0.6667	
Male, Hispanic	0.7497		0.9628		0.5521	
Female, Hispanic	0.0647		0.2765		0.1183	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.6C Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	88	92	123	120	138	134
GENDER						
Male	70	72	90	93	100	106
Female	46	54	76	69	81	77
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	81	80	103	107	118	118
White	73	71	111	107	130	120
Black or African American	34	39	34	36	45	50
American Indian or Alaska Native	6	4	7	8	9	8
Native Hawaiian or Other Pacific Islander	6	*	6	*	8	*
Asian	19	19	21	31	27	36
Two or More Races	18	12	18	17	21	20
Hispanic or Latino	35	41	57	57	63	65
EDUCATION						
< High School	42	36	46	45	59	54
High School Graduate	51	56	60	68	74	84
Some College	53	59	83	82	92	93
College Graduate	29	29	71	63	75	69
CURRENT EMPLOYMENT						
Full-Time	55	57	81	84	91	95
Part-Time	44	48	74	71	83	78
Unemployed	41	36	39	40	51	50
Other ²	39	36	52	56	63	67

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.6D Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.25	0.26	0.35	0.34	0.40	0.38
GENDER						
Male	0.40	0.41	0.52	0.53	0.57	0.61
Female	0.26	0.31	0.44	0.40	0.47	0.44
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.29	0.29	0.37	0.39	0.43	0.43
White	0.34	0.35	0.49	0.47	0.54	0.52
Black or African American	0.64	0.72	0.63	0.66	0.82	0.89
American Indian or Alaska Native	2.33	1.76	2.71	3.74	3.13	3.62
Native Hawaiian or Other Pacific Islander	3.13	*	2.85	*	3.95	*
Asian	0.99	0.88	1.12	1.40	1.37	1.61
Two or More Races	2.24	1.46	2.31	2.03	2.64	2.24
Hispanic or Latino	0.48	0.55	0.79	0.77	0.87	0.88
EDUCATION						
< High School	0.71	0.71	0.74	0.84	0.90	0.98
High School Graduate	0.42	0.45	0.46	0.53	0.54	0.64
Some College	0.41	0.45	0.58	0.56	0.63	0.63
College Graduate	0.56	0.50	1.14	0.99	1.18	1.04
CURRENT EMPLOYMENT						
Full-Time	0.39	0.38	0.54	0.54	0.58	0.59
Part-Time	0.43	0.48	0.65	0.67	0.70	0.72
Unemployed	0.93	0.89	0.89	0.96	1.11	1.18
Other ²	0.50	0.47	0.62	0.70	0.72	0.81

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.6P Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.0515		0.1249		0.0749	
GENDER						
Male	0.0060		0.1503		0.0576	
Female	0.6853		0.4538		0.5583	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0695		0.1438		0.0846	
White	0.0947		0.0278		0.0181	
Black or African American	0.9398		0.7585		0.7051	
American Indian or Alaska Native	0.0395		0.5377		0.7863	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7188		0.0860		0.2867	
Two or More Races	0.4652		0.7638		0.9604	
Hispanic or Latino	0.4536		0.5667		0.5547	
EDUCATION						
< High School	0.1239		0.6953		0.2428	
High School Graduate	0.0982		0.6372		0.3961	
Some College	0.9269		0.0466		0.1713	
College Graduate	0.5435		0.6723		0.6566	
CURRENT EMPLOYMENT						
Full-Time	0.3232		0.2597		0.3720	
Part-Time	0.4814		0.1114		0.0845	
Unemployed	0.1748		0.6800		0.7691	
Other ²	0.3470		0.5552		0.4216	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.7C Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	219	187	392	324	429	348
GENDER						
Male	149	149	332	265	349	288
Female	163	109	227	173	262	194
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	207	172	345	292	384	316
White	191	138	337	264	384	288
Black or African American	82	99	121	121	134	138
American Indian or Alaska Native	14	9	45	43	45	43
Native Hawaiian or Other Pacific Islander	6	19	*	19	*	26
Asian	38	26	74	48	81	55
Two or More Races	31	29	54	45	58	52
Hispanic or Latino	72	82	179	141	182	152
EDUCATION						
< High School	80	82	141	120	157	140
High School Graduate	114	100	219	170	233	188
Some College	157	110	208	173	251	190
College Graduate	81	87	279	213	288	226
CURRENT EMPLOYMENT						
Full-Time	168	132	347	257	383	281
Part-Time	83	71	128	132	140	141
Unemployed	65	50	92	73	104	82
Other ²	108	104	200	162	216	180

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.7D Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.11	0.09	0.19	0.16	0.21	0.17
GENDER						
Male	0.15	0.15	0.34	0.27	0.36	0.29
Female	0.15	0.10	0.21	0.16	0.25	0.18
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.12	0.10	0.20	0.17	0.22	0.18
White	0.14	0.10	0.23	0.19	0.26	0.20
Black or African American	0.36	0.41	0.51	0.49	0.56	0.55
American Indian or Alaska Native	1.30	0.80	3.91	3.05	3.86	3.07
Native Hawaiian or Other Pacific Islander	1.12	2.49	*	2.54	*	3.43
Asian	0.37	0.25	0.71	0.46	0.79	0.53
Two or More Races	1.13	1.02	1.89	1.50	2.06	1.76
Hispanic or Latino	0.25	0.28	0.63	0.48	0.64	0.52
EDUCATION						
< High School	0.29	0.29	0.52	0.42	0.57	0.48
High School Graduate	0.19	0.17	0.37	0.29	0.39	0.31
Some College	0.30	0.20	0.39	0.32	0.47	0.34
College Graduate	0.12	0.13	0.39	0.30	0.40	0.32
CURRENT EMPLOYMENT						
Full-Time	0.16	0.12	0.31	0.22	0.34	0.24
Part-Time	0.34	0.30	0.52	0.54	0.57	0.58
Unemployed	0.84	0.70	1.18	1.00	1.33	1.10
Other ²	0.16	0.15	0.30	0.24	0.33	0.27

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.7P Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.1406		0.7025		0.8487	
GENDER						
Male	0.0528		0.8819		0.7043	
Female	0.9361		0.6449		0.8370	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2435		0.8895		0.8275	
White	0.9164		0.7039		0.6891	
Black or African American	0.0264		0.2704		0.0538	
American Indian or Alaska Native	0.4037		0.6727		0.6245	
Native Hawaiian or Other Pacific Islander	0.2135		*		*	
Asian	0.8462		0.4281		0.4541	
Two or More Races	0.8370		0.8709		0.9832	
Hispanic or Latino	0.3095		0.5523		0.9823	
EDUCATION						
< High School	0.4244		0.9007		0.6488	
High School Graduate	0.9926		0.6415		0.9795	
Some College	0.7187		0.3855		0.5406	
College Graduate	0.0262		0.3141		0.5619	
CURRENT EMPLOYMENT						
Full-Time	0.1412		0.3074		0.5918	
Part-Time	0.4335		0.0736		0.0492	
Unemployed	0.9701		0.7100		0.9469	
Other ²	0.6934		0.5871		0.9338	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.8C Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	235	215	413	358	456	384
GENDER						
Male	168	170	341	286	363	311
Female	169	122	241	191	275	214
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	221	195	360	317	406	342
White	202	158	362	292	413	316
Black or African American	91	111	126	127	144	148
American Indian or Alaska Native	16	10	45	46	46	47
Native Hawaiian or Other Pacific Islander	9	19	*	20	*	27
Asian	42	34	74	58	83	67
Two or More Races	35	31	58	48	62	55
Hispanic or Latino	80	93	189	156	194	170
EDUCATION						
< High School	93	89	147	133	171	153
High School Graduate	123	118	229	182	245	202
Some College	161	126	223	194	268	217
College Graduate	85	95	291	224	300	236
CURRENT EMPLOYMENT						
Full-Time	175	149	352	273	388	299
Part-Time	93	86	154	152	167	164
Unemployed	79	63	99	85	117	96
Other ²	115	111	209	169	228	187

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.8D Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.10	0.09	0.17	0.15	0.19	0.16
GENDER						
Male	0.15	0.15	0.30	0.25	0.32	0.27
Female	0.14	0.10	0.20	0.15	0.22	0.17
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.11	0.10	0.18	0.16	0.20	0.17
White	0.13	0.10	0.22	0.18	0.25	0.19
Black or African American	0.32	0.38	0.43	0.41	0.49	0.47
American Indian or Alaska Native	1.13	0.77	3.23	2.78	3.19	2.78
Native Hawaiian or Other Pacific Islander	1.24	2.07	*	2.23	*	2.91
Asian	0.35	0.27	0.61	0.46	0.68	0.54
Two or More Races	1.01	0.88	1.62	1.24	1.76	1.45
Hispanic or Latino	0.22	0.25	0.53	0.42	0.54	0.46
EDUCATION						
< High School	0.28	0.27	0.44	0.40	0.51	0.45
High School Graduate	0.18	0.17	0.32	0.25	0.34	0.28
Some College	0.25	0.19	0.34	0.29	0.40	0.32
College Graduate	0.12	0.13	0.38	0.29	0.39	0.30
CURRENT EMPLOYMENT						
Full-Time	0.14	0.12	0.28	0.21	0.31	0.22
Part-Time	0.27	0.26	0.44	0.44	0.48	0.48
Unemployed	0.65	0.56	0.83	0.73	0.96	0.81
Other ²	0.16	0.15	0.28	0.23	0.31	0.25

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.8P Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.6075		0.3847		0.6604	
GENDER						
Male	0.6048		0.5615		0.8026	
Female	0.8486		0.4820		0.6823	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7410		0.5417		0.7087	
White	0.5699		0.2951		0.2629	
Black or African American	0.0512		0.2535		0.0547	
American Indian or Alaska Native	0.1040		0.6142		0.7003	
Native Hawaiian or Other Pacific Islander	0.4397		*		*	
Asian	0.7659		0.9410		0.8120	
Two or More Races	0.9304		0.8229		0.9466	
Hispanic or Latino	0.5849		0.4408		0.7920	
EDUCATION						
< High School	0.7826		0.8542		0.8201	
High School Graduate	0.5111		0.5847		0.8376	
Some College	0.7369		0.9508		0.9535	
College Graduate	0.0474		0.3420		0.5948	
CURRENT EMPLOYMENT						
Full-Time	0.2687		0.2194		0.4951	
Part-Time	0.7354		0.3890		0.3355	
Unemployed	0.5223		0.5949		0.9722	
Other ²	0.9846		0.4770		0.7248	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.9C Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	239	220	415	362	460	389
GEOGRAPHIC DIVISION						
Northeast	91	116	171	171	187	182
New England	55	43	70	73	93	84
Middle Atlantic	72	108	157	157	163	165
Midwest	90	79	163	158	183	167
East North Central	77	68	135	129	153	135
West North Central	46	36	89	74	99	79
South	169	132	262	225	299	242
South Atlantic	149	96	210	178	244	188
East South Central	60	37	75	78	94	82
West South Central	56	78	145	132	148	139
West	117	116	235	176	249	201
Mountain	57	51	106	97	116	104
Pacific	101	106	209	146	220	171
COUNTY TYPE						
Large Metro	171	194	372	304	395	339
Small Metro	167	119	235	202	289	222
250K – 1 Mil. Pop.	109	100	207	171	233	191
< 250K Pop.	127	69	116	118	177	127
Nonmetro	65	55	137	122	150	137
Urbanized	51	38	95	84	106	96
Less Urbanized	37	39	94	82	102	92
Completely Rural	20	14	33	28	43	31

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.9D Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.09	0.08	0.16	0.14	0.18	0.15
GEOGRAPHIC DIVISION						
Northeast	0.19	0.24	0.36	0.36	0.39	0.38
New England	0.44	0.35	0.56	0.58	0.74	0.67
Middle Atlantic	0.21	0.31	0.45	0.45	0.47	0.47
Midwest	0.16	0.14	0.29	0.28	0.33	0.30
East North Central	0.20	0.17	0.35	0.33	0.39	0.34
West North Central	0.26	0.21	0.52	0.42	0.58	0.45
South	0.17	0.13	0.27	0.23	0.31	0.24
South Atlantic	0.29	0.18	0.41	0.34	0.47	0.36
East South Central	0.39	0.24	0.48	0.50	0.61	0.52
West South Central	0.18	0.25	0.47	0.42	0.49	0.45
West	0.19	0.19	0.38	0.28	0.41	0.32
Mountain	0.30	0.27	0.57	0.51	0.62	0.55
Pacific	0.24	0.24	0.49	0.34	0.52	0.40
COUNTY TYPE						
Large Metro	0.12	0.13	0.24	0.20	0.25	0.22
Small Metro	0.20	0.15	0.29	0.24	0.34	0.25
250K – 1 Mil. Pop.	0.20	0.18	0.37	0.31	0.41	0.32
< 250K Pop.	0.47	0.27	0.44	0.42	0.62	0.44
Nonmetro	0.15	0.13	0.31	0.26	0.34	0.28
Urbanized	0.26	0.21	0.47	0.41	0.51	0.46
Less Urbanized	0.19	0.18	0.47	0.35	0.51	0.36
Completely Rural	0.42	0.30	0.66	0.62	0.87	0.66

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.9P Substance Dependence or Abuse in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.6367		0.3824		0.6153	
GEOGRAPHIC DIVISION						
Northeast	0.0780		0.9360		0.4623	
New England	0.7819		0.9454		0.8839	
Middle Atlantic	0.0319		0.9552		0.3647	
Midwest	0.8324		0.9452		0.8922	
East North Central	0.8862		0.9746		0.9090	
West North Central	0.4940		0.9334		0.6544	
South	0.8056		0.8210		0.7547	
South Atlantic	0.8586		0.8030		0.8631	
East South Central	0.1436		0.7332		0.3435	
West South Central	0.3910		0.8949		0.8062	
West	0.8634		0.1671		0.2886	
Mountain	0.4323		0.3906		0.6763	
Pacific	0.8287		0.0385		0.1468	
COUNTY TYPE						
Large Metro	0.4566		0.2958		0.5651	
Small Metro	0.8830		0.6844		0.6219	
250K – 1 Mil. Pop.	0.8412		0.2874		0.4335	
< 250K Pop.	0.6748		0.3268		0.7882	
Nonmetro	0.8302		0.6498		0.5743	
Urbanized	0.8061		0.5850		0.6778	
Less Urbanized	0.7864		0.7133		0.4578	
Completely Rural	0.6687		0.6648		0.5860	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.10C Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	41	46	36	42	51	55
GEOGRAPHIC DIVISION						
Northeast	15	14	15	15	19	18
New England	7	7	7	7	8	9
Middle Atlantic	13	12	13	13	17	16
Midwest	14	18	14	18	19	23
East North Central	11	16	11	17	16	22
West North Central	8	8	7	6	9	9
South	25	28	22	25	31	33
South Atlantic	18	21	16	15	23	23
East South Central	9	8	8	9	11	11
West South Central	15	17	11	17	18	21
West	23	29	19	22	27	32
Mountain	13	13	10	13	14	15
Pacific	19	26	16	18	23	27
COUNTY TYPE						
Large Metro	32	38	28	32	40	43
Small Metro	24	22	21	18	30	26
250K – 1 Mil. Pop.	19	18	17	15	24	22
< 250K Pop.	14	12	11	10	17	15
Nonmetro	11	16	13	20	15	23
Urbanized	8	10	7	14	10	16
Less Urbanized	7	12	9	12	10	14
Completely Rural	4	5	6	7	7	8

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.10D Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.17	0.18	0.14	0.17	0.20	0.22
GEOGRAPHIC DIVISION						
Northeast	0.35	0.33	0.36	0.36	0.44	0.44
New England	0.64	0.65	0.62	0.67	0.75	0.82
Middle Atlantic	0.42	0.38	0.44	0.43	0.53	0.51
Midwest	0.26	0.33	0.26	0.34	0.35	0.43
East North Central	0.30	0.43	0.30	0.45	0.41	0.58
West North Central	0.46	0.49	0.44	0.39	0.56	0.58
South	0.27	0.30	0.23	0.26	0.33	0.35
South Atlantic	0.39	0.45	0.35	0.32	0.49	0.49
East South Central	0.59	0.54	0.56	0.58	0.74	0.76
West South Central	0.47	0.51	0.34	0.53	0.55	0.64
West	0.39	0.49	0.32	0.38	0.45	0.54
Mountain	0.70	0.71	0.53	0.67	0.75	0.81
Pacific	0.46	0.64	0.40	0.45	0.56	0.68
COUNTY TYPE						
Large Metro	0.23	0.27	0.20	0.22	0.29	0.30
Small Metro	0.30	0.29	0.26	0.25	0.37	0.35
250K – 1 Mil. Pop.	0.36	0.36	0.31	0.30	0.44	0.43
< 250K Pop.	0.55	0.52	0.45	0.43	0.68	0.60
Nonmetro	0.30	0.42	0.31	0.51	0.39	0.58
Urbanized	0.50	0.66	0.43	0.91	0.58	1.03
Less Urbanized	0.38	0.59	0.47	0.59	0.57	0.69
Completely Rural	0.97	1.27	1.31	1.73	1.48	2.02

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.10P Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.9051		0.7517		0.5021	
GEOGRAPHIC DIVISION						
Northeast	0.2817		0.4766		0.1788	
New England	0.9188		0.4654		0.7347	
Middle Atlantic	0.2359		0.6726		0.1777	
Midwest	0.4336		0.7138		0.8383	
East North Central	0.8082		0.8353		0.8357	
West North Central	0.2411		0.1926		0.9620	
South	0.4646		0.6294		0.1685	
South Atlantic	0.9707		0.3548		0.3109	
East South Central	0.2275		0.1301		0.0754	
West South Central	0.5228		0.2659		0.9344	
West	0.5170		0.4525		0.3235	
Mountain	0.3767		0.9361		0.6365	
Pacific	0.2046		0.3424		0.1550	
COUNTY TYPE						
Large Metro	0.9110		0.9128		0.7944	
Small Metro	0.1468		0.0693		0.0427	
250K – 1 Mil. Pop.	0.3325		0.1566		0.1448	
< 250K Pop.	0.2485		0.2297		0.1323	
Nonmetro	0.1099		0.2387		0.1282	
Urbanized	0.6159		0.1362		0.2748	
Less Urbanized	0.1628		0.7437		0.3950	
Completely Rural	0.2595		0.9955		0.4441	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.11C Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	88	92	123	120	138	134
GEOGRAPHIC DIVISION						
Northeast	34	35	42	46	48	54
New England	20	20	21	23	25	30
Middle Atlantic	28	29	37	41	41	46
Midwest	34	39	45	55	52	59
East North Central	28	33	38	47	43	50
West North Central	18	19	25	28	28	29
South	53	52	81	74	88	80
South Atlantic	44	34	67	48	72	54
East South Central	16	18	24	29	29	31
West South Central	25	34	35	49	38	53
West	50	52	63	64	71	73
Mountain	26	21	30	29	37	33
Pacific	42	48	56	59	61	67
COUNTY TYPE						
Large Metro	70	78	120	108	134	127
Small Metro	50	48	68	73	85	81
250K – 1 Mil. Pop.	42	39	53	60	66	66
< 250K Pop.	29	28	47	44	57	50
Nonmetro	31	25	45	46	52	49
Urbanized	21	17	35	34	41	36
Less Urbanized	23	18	29	28	34	30
Completely Rural	8	6	12	12	12	13

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.11D Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.25	0.26	0.35	0.34	0.40	0.38
GEOGRAPHIC DIVISION						
Northeast	0.56	0.57	0.69	0.75	0.77	0.88
New England	1.20	1.21	1.27	1.41	1.51	1.82
Middle Atlantic	0.62	0.64	0.82	0.91	0.90	1.03
Midwest	0.45	0.53	0.61	0.75	0.70	0.79
East North Central	0.55	0.65	0.74	0.92	0.85	0.97
West North Central	0.78	0.81	1.08	1.19	1.23	1.27
South	0.41	0.40	0.63	0.57	0.68	0.62
South Atlantic	0.68	0.52	1.02	0.73	1.10	0.81
East South Central	0.78	0.87	1.17	1.39	1.40	1.50
West South Central	0.58	0.79	0.82	1.14	0.89	1.23
West	0.59	0.62	0.75	0.76	0.84	0.87
Mountain	1.01	0.83	1.18	1.12	1.46	1.29
Pacific	0.73	0.82	0.96	1.01	1.04	1.15
COUNTY TYPE						
Large Metro	0.36	0.39	0.53	0.49	0.57	0.57
Small Metro	0.44	0.41	0.53	0.60	0.63	0.63
250K – 1 Mil. Pop.	0.57	0.52	0.67	0.75	0.81	0.77
< 250K Pop.	0.65	0.72	0.86	1.06	1.00	1.15
Nonmetro	0.54	0.47	0.72	0.76	0.79	0.77
Urbanized	0.74	0.66	1.11	1.05	1.20	1.12
Less Urbanized	0.87	0.71	1.12	1.05	1.23	1.05
Completely Rural	1.52	1.52	1.89	2.56	1.96	2.70

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.11P Substance Dependence or Abuse in the Past Year among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.0515		0.1249		0.0749	
GEOGRAPHIC DIVISION						
Northeast	0.3426		0.2129		0.2787	
New England	0.5141		0.4693		0.9156	
Middle Atlantic	0.1087		0.3123		0.1820	
Midwest	0.6727		0.9828		0.6270	
East North Central	0.8563		0.3453		0.3852	
West North Central	0.2534		0.1338		0.6370	
South	0.2354		0.4188		0.2461	
South Atlantic	0.3503		0.4119		0.3590	
East South Central	0.7519		0.4621		0.4642	
West South Central	0.3086		0.9082		0.7077	
West	0.2042		0.2875		0.3850	
Mountain	0.2830		0.9568		0.4576	
Pacific	0.3865		0.2296		0.5709	
COUNTY TYPE						
Large Metro	0.1114		0.2223		0.2324	
Small Metro	0.8534		0.3235		0.3317	
250K – 1 Mil. Pop.	0.9679		0.1364		0.2191	
< 250K Pop.	0.7027		0.7215		0.9641	
Nonmetro	0.0320		0.8267		0.2185	
Urbanized	0.1222		0.6564		0.1226	
Less Urbanized	0.0598		0.9867		0.6218	
Completely Rural	0.6462		0.5775		0.4803	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.12C Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	219	187	392	324	429	348
GEOGRAPHIC DIVISION						
Northeast	83	108	163	170	176	182
New England	51	37	64	70	85	77
Middle Atlantic	65	101	150	156	155	166
Midwest	78	65	155	141	167	150
East North Central	68	56	128	116	139	122
West North Central	37	31	85	68	93	72
South	162	116	243	201	275	216
South Atlantic	145	86	195	161	223	170
East South Central	56	32	69	67	86	73
West South Central	46	67	139	116	142	124
West	101	91	228	154	236	173
Mountain	51	44	104	90	111	95
Pacific	87	81	203	122	209	143
COUNTY TYPE						
Large Metro	151	159	343	275	358	299
Small Metro	156	102	221	186	267	201
250K – 1 Mil. Pop.	95	85	194	161	215	175
< 250K Pop.	124	60	105	101	160	109
Nonmetro	54	46	125	106	135	117
Urbanized	45	33	84	71	93	81
Less Urbanized	25	32	88	71	92	79
Completely Rural	18	10	26	23	37	25

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.12D Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.11	0.09	0.19	0.16	0.21	0.17
GEOGRAPHIC DIVISION						
Northeast	0.22	0.29	0.44	0.46	0.48	0.49
New England	0.53	0.37	0.65	0.71	0.87	0.79
Middle Atlantic	0.24	0.37	0.55	0.57	0.57	0.61
Midwest	0.18	0.15	0.36	0.32	0.39	0.34
East North Central	0.23	0.18	0.43	0.38	0.46	0.40
West North Central	0.28	0.23	0.64	0.51	0.70	0.54
South	0.21	0.15	0.32	0.26	0.36	0.28
South Atlantic	0.36	0.21	0.48	0.39	0.55	0.42
East South Central	0.47	0.27	0.57	0.56	0.72	0.61
West South Central	0.20	0.28	0.60	0.49	0.62	0.53
West	0.22	0.19	0.49	0.32	0.50	0.36
Mountain	0.36	0.30	0.72	0.62	0.78	0.65
Pacific	0.27	0.24	0.62	0.37	0.64	0.43
COUNTY TYPE						
Large Metro	0.14	0.14	0.29	0.23	0.31	0.25
Small Metro	0.25	0.17	0.36	0.29	0.42	0.31
250K – 1 Mil. Pop.	0.23	0.20	0.45	0.38	0.50	0.39
< 250K Pop.	0.62	0.31	0.54	0.48	0.77	0.51
Nonmetro	0.16	0.14	0.37	0.30	0.40	0.32
Urbanized	0.31	0.24	0.55	0.48	0.61	0.54
Less Urbanized	0.17	0.20	0.57	0.41	0.60	0.43
Completely Rural	0.49	0.28	0.69	0.66	1.00	0.69

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.12P Substance Dependence or Abuse in the Past Year among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.1406		0.7025		0.8487	
GEOGRAPHIC DIVISION						
Northeast	0.0151		0.6322		0.2265	
New England	0.5627		0.7027		0.8696	
Middle Atlantic	0.0023		0.7234		0.1521	
Midwest	0.8453		0.9752		0.9947	
East North Central	0.9695		0.7878		0.6834	
West North Central	0.6313		0.6274		0.5138	
South	0.7335		0.9257		0.8045	
South Atlantic	0.8882		0.9298		0.8117	
East South Central	0.1363		0.9283		0.5999	
West South Central	0.0825		0.9862		0.6795	
West	0.7368		0.2316		0.3304	
Mountain	0.9219		0.3688		0.4400	
Pacific	0.6583		0.0571		0.1160	
COUNTY TYPE						
Large Metro	0.0947		0.4910		0.9102	
Small Metro	0.8564		0.9798		0.9961	
250K – 1 Mil. Pop.	0.6726		0.5536		0.7521	
< 250K Pop.	0.8806		0.2798		0.6348	
Nonmetro	0.7276		0.7370		0.4749	
Urbanized	0.8669		0.6926		0.5031	
Less Urbanized	0.2993		0.7384		0.4225	
Completely Rural	0.3643		0.5779		0.3931	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.13C Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	235	215	413	358	456	384
GEOGRAPHIC DIVISION						
Northeast	89	115	171	170	186	182
New England	54	43	71	73	93	84
Middle Atlantic	71	107	156	156	161	164
Midwest	87	77	164	156	182	163
East North Central	76	65	136	128	152	132
West North Central	44	36	89	73	99	78
South	169	130	261	222	297	239
South Atlantic	149	92	210	176	243	184
East South Central	59	37	74	78	93	82
West South Central	53	78	143	130	148	139
West	112	112	234	174	247	200
Mountain	55	52	106	96	114	103
Pacific	97	102	209	145	218	170
COUNTY TYPE						
Large Metro	166	188	370	302	390	335
Small Metro	164	116	233	201	286	218
250K – 1 Mil. Pop.	107	97	206	170	230	187
< 250K Pop.	126	67	115	117	175	125
Nonmetro	64	54	135	120	148	133
Urbanized	51	38	94	84	105	94
Less Urbanized	35	37	93	79	100	89
Completely Rural	20	13	31	27	41	29

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.13D Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.10	0.09	0.17	0.15	0.19	0.16
GEOGRAPHIC DIVISION						
Northeast	0.21	0.26	0.40	0.39	0.43	0.42
New England	0.48	0.37	0.62	0.64	0.82	0.73
Middle Atlantic	0.22	0.34	0.49	0.49	0.51	0.51
Midwest	0.17	0.15	0.32	0.30	0.36	0.32
East North Central	0.21	0.18	0.39	0.36	0.43	0.37
West North Central	0.28	0.23	0.57	0.46	0.64	0.50
South	0.19	0.14	0.30	0.25	0.34	0.27
South Atlantic	0.32	0.19	0.45	0.37	0.52	0.39
East South Central	0.42	0.26	0.53	0.55	0.66	0.58
West South Central	0.19	0.28	0.52	0.47	0.54	0.50
West	0.20	0.20	0.42	0.31	0.45	0.35
Mountain	0.33	0.31	0.63	0.56	0.68	0.60
Pacific	0.25	0.26	0.54	0.37	0.57	0.44
COUNTY TYPE						
Large Metro	0.13	0.14	0.26	0.22	0.28	0.24
Small Metro	0.22	0.16	0.32	0.27	0.38	0.28
250K – 1 Mil. Pop.	0.22	0.19	0.41	0.34	0.45	0.35
< 250K Pop.	0.51	0.29	0.48	0.46	0.68	0.48
Nonmetro	0.17	0.14	0.34	0.28	0.37	0.30
Urbanized	0.29	0.23	0.51	0.46	0.56	0.50
Less Urbanized	0.20	0.20	0.52	0.37	0.55	0.40
Completely Rural	0.46	0.31	0.68	0.66	0.93	0.71

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.13P Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.6075		0.3847		0.6604	
GEOGRAPHIC DIVISION						
Northeast	0.0532		0.8928		0.3873	
New England	0.7913		0.8934		0.9065	
Middle Atlantic	0.0195		0.9319		0.2989	
Midwest	0.7143		0.9663		0.8651	
East North Central	0.9177		0.9488		0.9355	
West North Central	0.3575		0.9831		0.6471	
South	0.9079		0.8497		0.8753	
South Atlantic	0.8544		0.8558		0.9426	
East South Central	0.2026		0.8593		0.4647	
West South Central	0.3044		0.9903		0.7999	
West	0.7499		0.1395		0.2270	
Mountain	0.5689		0.3836		0.6293	
Pacific	0.9571		0.0292		0.0985	
COUNTY TYPE						
Large Metro	0.4556		0.2839		0.5780	
Small Metro	0.9273		0.7802		0.7698	
250K – 1 Mil. Pop.	0.6959		0.3315		0.5181	
< 250K Pop.	0.7843		0.2824		0.6743	
Nonmetro	0.5481		0.7656		0.7311	
Urbanized	0.7187		0.7329		0.7974	
Less Urbanized	0.9255		0.7370		0.5240	
Completely Rural	0.4609		0.6488		0.4690	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.14C Substance Dependence for Specific Substances in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Past Year Dependence	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	211	186	29	34	73	80	197	157
Marijuana and Hashish	164	124	27	26	61	67	152	100
Cocaine	73	66	4	5	24	19	68	63
Heroin	57	59	2	6	20	23	52	53
Hallucinogens	20	19	8	6	12	12	14	13
Inhalants	10	13	5	6	7	3	*	11
Nonmedical Use of Psychotherapeutics ^{2,3}	129	103	12	19	40	39	121	94
Pain Relievers	121	96	11	15	35	36	115	89
Tranquilizers	47	44	7	7	17	13	42	42
Stimulants ²	71	44	5	8	21	17	68	41
Sedatives	23	33	2	9	9	8	21	31
ALCOHOL	286	246	22	25	78	78	278	228
BOTH ILLICIT DRUGS AND ALCOHOL¹	87	93	13	15	38	38	78	80
ILLICIT DRUGS OR ALCOHOL¹	336	291	35	39	102	104	319	264

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.14D Substance Dependence for Specific Substances in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Past Year Dependence	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	0.08	0.07	0.12	0.14	0.21	0.23	0.10	0.08
Marijuana and Hashish	0.06	0.05	0.11	0.10	0.18	0.19	0.07	0.05
Cocaine	0.03	0.02	0.01	0.02	0.07	0.06	0.03	0.03
Heroin	0.02	0.02	0.01	0.02	0.06	0.07	0.03	0.03
Hallucinogens	0.01	0.01	0.03	0.02	0.04	0.03	0.01	0.01
Inhalants	0.00	0.00	0.02	0.02	0.02	0.01	*	0.01
Nonmedical Use of Psychotherapeutics ^{2,3}	0.05	0.04	0.05	0.08	0.11	0.11	0.06	0.05
Pain Relievers	0.05	0.04	0.04	0.06	0.10	0.10	0.06	0.04
Tranquilizers	0.02	0.02	0.03	0.03	0.05	0.04	0.02	0.02
Stimulants ²	0.03	0.02	0.02	0.03	0.06	0.05	0.03	0.02
Sedatives	0.01	0.01	0.01	0.03	0.03	0.02	0.01	0.02
ALCOHOL	0.11	0.09	0.09	0.10	0.22	0.22	0.14	0.11
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.03	0.04	0.05	0.06	0.11	0.11	0.04	0.04
ILLICIT DRUGS OR ALCOHOL¹	0.13	0.11	0.14	0.16	0.29	0.30	0.16	0.13

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.14P Substance Dependence for Specific Substances in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Past Year Dependence	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLCIT DRUGS¹	0.9361		0.5862		0.1528		0.4040	
Marijuana and Hashish	0.3900		0.0374		0.2337		0.9231	
Cocaine	0.5481		0.9737		0.3760		0.7307	
Heroin	0.2273		0.1218		0.9092		0.2055	
Hallucinogens	0.7463		0.5738		0.9935		0.8717	
Inhalants	0.8302		0.9377		0.1718		*	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.6752		0.0310		0.1616		0.5108	
Pain Relievers	0.9808		0.3493		0.4159		0.8485	
Tranquilizers	0.4936		0.7024		0.3018		0.2955	
Stimulants ²	0.6721		0.4292		0.3363		0.8338	
Sedatives	0.1876		0.0513		0.6233		0.2636	
ALCOHOL	0.9627		0.5138		0.5922		0.8803	
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.8575		0.3866		0.1941		0.5571	
ILLCIT DRUGS OR ALCOHOL¹	0.9503		0.5765		0.8585		0.8267	

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.15C Received Substance Use Treatment in the Past Year, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	157	141	174	158	123	113	211	186
12	*	3	3	3	*	3	3	3
13	4	3	5	3	4	3	5	4
14	6	7	6	6	6	6	8	9
15	8	8	7	5	6	4	10	8
16	11	9	10	6	10	4	12	12
17	9	13	8	11	7	11	10	14
18	13	10	12	8	11	6	16	12
19	12	11	11	12	9	9	15	14
20	12	13	11	12	11	10	14	17
21	14	13	13	11	12	10	15	14
22	13	16	12	12	11	9	15	18
23	14	17	14	15	11	13	18	21
24	10	14	11	12	8	7	14	17
25	13	18	14	16	10	12	17	23
26-29	51	46	56	45	44	32	67	58
30-34	53	38	54	41	40	27	68	49
35-39	50	36	60	40	47	26	64	49
40-44	39	44	55	41	27	28	64	58
45-49	29	27	45	37	22	23	59	42
50-54	68	54	76	63	59	47	84	71
55-59	54	57	40	57	35	48	60	66
60-64	47	34	81	53	43	30	83	58
65 or Older	12	39	39	52	12	39	40	55

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.15D Received Substance Use Treatment in the Past Year, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.06	0.05	0.07	0.06	0.05	0.04	0.08	0.07
12	*	0.07	0.07	0.07	*	0.07	0.07	0.08
13	0.10	0.07	0.11	0.08	0.10	0.07	0.11	0.10
14	0.14	0.18	0.13	0.14	0.13	0.14	0.18	0.20
15	0.19	0.18	0.16	0.11	0.15	0.10	0.24	0.19
16	0.26	0.21	0.24	0.13	0.22	0.10	0.28	0.26
17	0.21	0.32	0.19	0.27	0.18	0.27	0.24	0.34
18	0.28	0.22	0.26	0.17	0.23	0.12	0.34	0.25
19	0.28	0.27	0.25	0.28	0.22	0.20	0.36	0.34
20	0.29	0.30	0.26	0.27	0.24	0.24	0.32	0.39
21	0.34	0.29	0.30	0.26	0.29	0.23	0.36	0.32
22	0.30	0.35	0.28	0.27	0.24	0.20	0.34	0.40
23	0.31	0.38	0.32	0.34	0.26	0.30	0.41	0.47
24	0.24	0.32	0.25	0.28	0.17	0.16	0.32	0.40
25	0.32	0.42	0.34	0.37	0.25	0.28	0.42	0.52
26-29	0.30	0.27	0.32	0.26	0.26	0.19	0.38	0.33
30-34	0.26	0.18	0.27	0.19	0.20	0.13	0.33	0.23
35-39	0.27	0.18	0.32	0.20	0.25	0.13	0.33	0.25
40-44	0.18	0.21	0.25	0.20	0.12	0.14	0.30	0.29
45-49	0.14	0.13	0.23	0.18	0.11	0.11	0.30	0.20
50-54	0.29	0.23	0.33	0.26	0.25	0.20	0.36	0.30
55-59	0.26	0.27	0.19	0.27	0.17	0.23	0.29	0.31
60-64	0.27	0.20	0.45	0.30	0.24	0.17	0.47	0.33
65 or Older	0.03	0.09	0.09	0.12	0.03	0.09	0.09	0.12

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.15P Received Substance Use Treatment in the Past Year, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.3401		0.7453		0.5827		0.8436	
12	*		0.9064		*		0.5919	
13	0.1817		0.2630		0.1966		0.5814	
14	0.7165		0.9314		0.9466		0.7818	
15	0.6087		0.3104		0.3485		0.1463	
16	0.1348		0.0237		0.0100		0.5762	
17	0.2844		0.5267		0.2833		0.2729	
18	0.1375		0.0687		0.0877		0.0349	
19	0.3891		0.9501		0.5927		0.4719	
20	0.8461		0.5197		0.7653		1.0000	
21	0.7606		0.5974		0.4458		0.8341	
22	0.8948		0.6262		0.2243		0.6403	
23	0.3593		0.7507		0.8719		0.5785	
24	0.9429		0.7608		0.7337		0.6675	
25	0.5460		0.5487		1.0000		0.4494	
26-29	0.7264		0.3134		0.5283		0.9537	
30-34	0.3644		0.8491		0.6433		0.4054	
35-39	0.7672		0.4791		0.5882		0.9542	
40-44	0.2302		0.8957		0.1742		0.9066	
45-49	0.6031		0.7132		0.3905		0.2465	
50-54	0.4279		0.2736		0.3083		0.1906	
55-59	0.4491		0.3029		0.2739		0.5560	
60-64	0.3523		0.1754		0.4192		0.3576	
65 or Older	0.0293		0.0574		0.0358		0.0340	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.16C Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	157	141	174	158	123	113	211	186
AGE								
12-17	17	20	16	16	14	15	20	23
18-25	37	42	36	36	31	28	46	52
26 or Older	149	131	168	151	118	108	204	172
GENDER								
Male	107	110	133	133	80	91	162	155
Female	109	81	113	80	93	65	131	97
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	140	130	156	139	114	105	186	167
White	133	107	145	117	107	83	174	142
Black or African American	46	64	54	67	36	60	66	79
American Indian or Alaska Native	13	8	15	12	13	8	15	12
Native Hawaiian or Other Pacific Islander	5	8	5	5	5	4	9	8
Asian	18	8	29	13	18	7	30	13
Two or More Races	13	7	21	23	13	6	22	24
Hispanic or Latino	66	59	75	71	42	42	99	89

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.16D Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.06	0.05	0.07	0.06	0.05	0.04	0.08	0.07
AGE								
12-17	0.07	0.08	0.06	0.06	0.06	0.06	0.08	0.09
18-25	0.11	0.12	0.10	0.10	0.09	0.08	0.13	0.15
26 or Older	0.07	0.06	0.08	0.07	0.06	0.05	0.10	0.08
GENDER								
Male	0.08	0.09	0.10	0.10	0.06	0.07	0.13	0.12
Female	0.08	0.06	0.08	0.06	0.07	0.05	0.10	0.07
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.06	0.06	0.07	0.06	0.05	0.05	0.08	0.08
White	0.08	0.06	0.09	0.07	0.06	0.05	0.10	0.08
Black or African American	0.15	0.20	0.17	0.21	0.11	0.19	0.21	0.25
American Indian or Alaska Native	0.88	0.53	1.03	0.76	0.88	0.50	1.05	0.77
Native Hawaiian or Other Pacific Islander	0.60	0.74	0.60	0.46	0.60	0.36	1.04	0.82
Asian	0.14	0.05	0.22	0.09	0.14	0.05	0.23	0.10
Two or More Races	0.30	0.16	0.49	0.52	0.29	0.13	0.51	0.53
Hispanic or Latino	0.16	0.14	0.18	0.17	0.10	0.10	0.24	0.21

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.16P Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.3401		0.7453		0.5827		0.8436	
AGE								
12-17	0.7487		0.1741		0.2953		0.8573	
18-25	0.7867		0.4007		0.1450		0.9703	
26 or Older	0.2541		0.9853		0.2736		0.8124	
GENDER								
Male	0.2773		0.9893		0.4854		0.6333	
Female	0.7957		0.5692		0.9457		0.7530	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.2906		0.8603		0.4640		0.6640	
White	0.6780		0.6976		0.9509		0.9475	
Black or African American	0.0451		0.2335		0.0336		0.1087	
American Indian or Alaska Native	0.6668		0.5977		0.3790		0.8422	
Native Hawaiian or Other Pacific Islander	0.9833		0.6516		0.3703		0.7511	
Asian	0.6481		0.2798		0.5434		0.1860	
Two or More Races	0.5341		0.5568		0.4073		0.6132	
Hispanic or Latino	1.0000		0.7006		0.7257		0.6825	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.17C Received Substance Use Treatment in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	17	20	16	16	14	15	20	23
GENDER								
Male	15	13	13	9	12	8	17	15
Female	11	16	11	13	10	12	13	17
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	16	17	15	14	14	14	18	19
White	14	16	13	13	12	13	17	18
Black or African American	7	5	6	4	6	4	7	6
American Indian or Alaska Native	3	1	3	1	3	*	3	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	4	*	0	*	0	1	4
Two or More Races	2	3	2	3	2	3	2	3
Hispanic or Latino	9	11	8	7	8	6	10	13
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	11	11	10	7	9	7	13	13
Female, White, Not Hispanic	9	11	9	11	8	10	10	12
Male, Black, Not Hispanic	7	4	5	4	5	4	7	5
Female, Black, Not Hispanic	2	3	2	2	2	2	2	4
Male, Hispanic	7	5	6	2	6	2	7	6
Female, Hispanic	6	9	6	6	5	5	7	10

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.17D Received Substance Use Treatment in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.07	0.08	0.06	0.06	0.06	0.06	0.08	0.09
GENDER								
Male	0.12	0.10	0.10	0.07	0.09	0.07	0.13	0.12
Female	0.09	0.13	0.09	0.11	0.08	0.10	0.10	0.14
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.08	0.09	0.08	0.07	0.07	0.07	0.10	0.10
White	0.10	0.12	0.10	0.10	0.09	0.09	0.12	0.13
Black or African American	0.20	0.15	0.17	0.12	0.17	0.12	0.20	0.18
American Indian or Alaska Native	1.71	0.71	1.71	0.57	1.71	*	1.72	0.74
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	0.32	*	0.01	*	0.01	0.07	0.32
Two or More Races	0.32	0.43	0.30	0.43	0.28	0.43	0.33	0.44
Hispanic or Latino	0.16	0.19	0.14	0.12	0.14	0.10	0.18	0.22
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	0.15	0.15	0.14	0.11	0.12	0.10	0.19	0.18
Female, White, Not Hispanic	0.13	0.17	0.13	0.16	0.12	0.16	0.15	0.19
Male, Black, Not Hispanic	0.38	0.23	0.31	0.22	0.31	0.22	0.39	0.25
Female, Black, Not Hispanic	0.10	0.18	0.10	0.10	0.10	0.10	0.10	0.25
Male, Hispanic	0.24	0.18	0.20	0.08	0.20	0.08	0.26	0.23
Female, Hispanic	0.23	0.33	0.21	0.22	0.19	0.20	0.25	0.37

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.17P Received Substance Use Treatment in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.7487		0.1741		0.2953		0.8573	
GENDER								
Male	0.0904		0.0243		0.0449		0.1675	
Female	0.2028		0.7774		0.6314		0.2312	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.9457		0.6341		0.9102		0.9001	
White	0.6626		0.7836		0.5154		0.8098	
Black or African American	0.3267		0.3357		0.3730		0.6795	
American Indian or Alaska Native	0.2248		0.1438		*		0.2233	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	*		*		*		0.2732	
Two or More Races	0.6803		0.6986		0.9646		0.5221	
Hispanic or Latino	0.6139		0.0468		0.0418		0.8852	
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	0.8772		0.4635		0.7493		0.8847	
Female, White, Not Hispanic	0.4325		0.3083		0.2612		0.5978	
Male, Black, Not Hispanic	0.1734		0.3942		0.4381		0.2116	
Female, Black, Not Hispanic	0.5080		0.6353		0.6353		0.1633	
Male, Hispanic	0.1203		0.0343		0.0313		0.2561	
Female, Hispanic	0.5638		0.3513		0.3934		0.4744	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.18C Received Substance Use Treatment in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	37	42	36	36	31	28	46	52
GENDER								
Male	31	34	30	30	26	23	38	42
Female	20	24	20	21	16	16	24	29
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	33	39	29	33	27	27	39	47
White	32	36	28	29	25	24	35	42
Black or African American	8	11	8	11	6	8	13	14
American Indian or Alaska Native	3	3	3	3	3	3	3	4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	6	5	5	8	5	5	10	8
Two or More Races	4	6	4	5	3	5	4	8
Hispanic or Latino	14	17	18	13	13	9	22	21
EDUCATION								
< High School	16	20	15	16	11	14	21	23
High School Graduate	27	23	25	20	24	15	30	28
Some College	20	25	20	22	16	18	26	30
College Graduate	4	10	8	10	2	4	8	16
CURRENT EMPLOYMENT								
Full-Time	19	26	19	22	15	17	25	34
Part-Time	21	17	19	17	17	13	24	22
Unemployed	17	21	15	15	14	14	18	22
Other ³	16	15	16	14	13	10	22	19

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.18D Received Substance Use Treatment in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.11	0.12	0.10	0.10	0.09	0.08	0.13	0.15
GENDER								
Male	0.18	0.19	0.17	0.17	0.15	0.13	0.22	0.24
Female	0.12	0.14	0.11	0.12	0.09	0.09	0.14	0.17
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.12	0.14	0.11	0.12	0.10	0.10	0.14	0.17
White	0.16	0.19	0.14	0.15	0.13	0.12	0.18	0.21
Black or African American	0.16	0.22	0.17	0.21	0.11	0.16	0.26	0.28
American Indian or Alaska Native	1.15	1.52	1.15	1.64	1.15	1.51	1.15	1.69
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	0.30	0.25	0.27	0.40	0.27	0.25	0.53	0.40
Two or More Races	0.50	0.75	0.49	0.64	0.42	0.60	0.56	0.98
Hispanic or Latino	0.19	0.23	0.25	0.17	0.17	0.13	0.30	0.28
EDUCATION								
< High School	0.29	0.39	0.26	0.32	0.20	0.28	0.36	0.44
High School Graduate	0.23	0.19	0.21	0.17	0.20	0.13	0.25	0.23
Some College	0.16	0.20	0.16	0.17	0.13	0.14	0.21	0.24
College Graduate	0.08	0.19	0.16	0.18	0.04	0.07	0.17	0.29
CURRENT EMPLOYMENT								
Full-Time	0.14	0.18	0.14	0.15	0.11	0.12	0.19	0.23
Part-Time	0.21	0.18	0.20	0.17	0.17	0.13	0.24	0.23
Unemployed	0.40	0.52	0.35	0.38	0.33	0.35	0.44	0.56
Other ³	0.21	0.21	0.21	0.19	0.17	0.14	0.29	0.25

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.18P Received Substance Use Treatment in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.7867		0.4007		0.1450		0.9703	
GENDER								
Male	0.5314		0.4132		0.0865		0.7976	
Female	0.6730		0.7724		0.9281		0.7511	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.6107		0.8759		0.2559		0.6037	
White	0.4498		0.6252		0.1398		0.7756	
Black or African American	0.4991		0.4833		0.5076		0.7092	
American Indian or Alaska Native	0.8222		0.5826		0.8835		0.3432	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.6168		0.2367		0.8798		0.6681	
Two or More Races	0.4837		0.8272		0.7097		0.3071	
Hispanic or Latino	0.6129		0.0270		0.2577		0.2652	
EDUCATION								
< High School	0.9365		0.4809		0.9405		0.6665	
High School Graduate	0.1873		0.1737		0.0201		0.3624	
Some College	0.3109		0.6272		0.6317		0.3562	
College Graduate	0.1892		0.4482		0.5300		0.1312	
CURRENT EMPLOYMENT								
Full-Time	0.5121		0.9792		0.4745		0.2346	
Part-Time	0.1757		0.2564		0.1743		0.1597	
Unemployed	0.7597		0.4797		0.6723		0.6756	
Other ³	0.8291		0.9070		0.7130		0.5631	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.19C Received Substance Use Treatment in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	149	131	168	151	118	108	204	172
GENDER								
Male	100	102	129	128	74	86	156	145
Female	106	76	110	75	91	61	127	89
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	134	120	153	132	111	99	180	155
White	125	99	141	110	103	78	167	131
Black or African American	44	64	54	67	35	60	65	79
American Indian or Alaska Native	12	7	14	11	12	6	14	12
Native Hawaiian or Other Pacific Islander	5	7	5	*	5	*	9	8
Asian	17	4	29	9	17	4	29	10
Two or More Races	12	2	21	22	12	1	21	22
Hispanic or Latino	63	56	72	69	39	39	95	82
EDUCATION								
< High School	72	67	74	78	51	50	97	92
High School Graduate	69	80	86	81	52	66	100	96
Some College	75	64	90	72	66	50	108	88
College Graduate	70	47	96	52	63	36	103	62
CURRENT EMPLOYMENT								
Full-Time	50	66	88	90	34	44	103	109
Part-Time	54	41	43	42	40	34	60	48
Unemployed	70	40	76	43	67	32	82	52
Other ³	102	99	120	103	81	85	141	120

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.19D Received Substance Use Treatment in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.07	0.06	0.08	0.07	0.06	0.05	0.10	0.08
GENDER								
Male	0.10	0.10	0.13	0.13	0.08	0.09	0.16	0.15
Female	0.10	0.07	0.10	0.07	0.09	0.06	0.12	0.08
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.08	0.07	0.09	0.08	0.06	0.06	0.10	0.09
White	0.09	0.07	0.10	0.08	0.08	0.06	0.12	0.10
Black or African American	0.20	0.27	0.23	0.28	0.15	0.25	0.28	0.33
American Indian or Alaska Native	1.16	0.61	1.38	0.96	1.16	0.57	1.41	0.98
Native Hawaiian or Other Pacific Islander	0.84	1.01	0.84	*	0.84	*	1.54	1.05
Asian	0.17	0.04	0.28	0.09	0.17	0.04	0.28	0.09
Two or More Races	0.44	0.08	0.74	0.77	0.43	0.04	0.76	0.78
Hispanic or Latino	0.22	0.19	0.25	0.23	0.14	0.13	0.33	0.28
EDUCATION								
< High School	0.27	0.24	0.28	0.28	0.19	0.19	0.36	0.33
High School Graduate	0.12	0.14	0.15	0.14	0.09	0.12	0.17	0.17
Some College	0.15	0.12	0.17	0.13	0.13	0.09	0.21	0.16
College Graduate	0.10	0.07	0.14	0.08	0.09	0.05	0.15	0.09
CURRENT EMPLOYMENT								
Full-Time	0.05	0.06	0.08	0.08	0.03	0.04	0.10	0.10
Part-Time	0.23	0.17	0.18	0.18	0.17	0.14	0.25	0.20
Unemployed	0.91	0.56	0.98	0.61	0.86	0.46	1.05	0.73
Other ³	0.16	0.15	0.18	0.16	0.12	0.13	0.22	0.18

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.19P Received Substance Use Treatment in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.2541		0.9853		0.2736		0.8124	
GENDER								
Male	0.1088		0.6956		0.1266		0.4761	
Female	0.9865		0.5733		0.9830		0.5684	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.2003		0.8641		0.2857		0.7359	
White	0.5340		0.7476		0.7060		0.9801	
Black or African American	0.0401		0.2354		0.0296		0.1068	
American Indian or Alaska Native	0.8217		0.6750		0.5459		0.8306	
Native Hawaiian or Other Pacific Islander	0.8781		*		*		0.6690	
Asian	0.5374		0.1317		0.5374		0.1538	
Two or More Races	0.3067		0.5497		0.2575		0.4648	
Hispanic or Latino	0.9618		0.7671		0.7644		0.8822	
EDUCATION								
< High School	0.6369		0.2617		0.5313		0.3937	
High School Graduate	0.1293		0.3633		0.0703		0.3707	
Some College	0.6571		0.9064		0.9435		0.6820	
College Graduate	0.7779		0.0246		0.6914		0.0524	
CURRENT EMPLOYMENT								
Full-Time	0.0069		0.2192		0.0267		0.0655	
Part-Time	0.4619		0.6687		0.6672		0.4296	
Unemployed	0.1609		0.0358		0.0363		0.1469	
Other ³	0.1745		0.6377		0.0369		0.8726	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.20C Received Substance Use Treatment in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	155	140	172	157	122	112	210	186
GENDER								
Male	106	109	133	133	79	90	161	154
Female	108	80	112	79	92	64	130	95
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	138	129	155	138	113	104	184	166
White	130	108	144	117	106	83	172	143
Black or African American	45	65	54	69	35	60	67	81
American Indian or Alaska Native	12	8	14	12	12	8	15	12
Native Hawaiian or Other Pacific Islander	5	8	5	5	5	4	9	8
Asian	18	7	29	13	18	7	30	13
Two or More Races	13	6	21	23	12	5	22	24
Hispanic or Latino	65	58	75	71	41	41	98	87
EDUCATION								
< High School	74	70	75	78	52	53	100	96
High School Graduate	73	83	89	85	58	68	103	102
Some College	78	70	94	76	68	54	113	96
College Graduate	70	48	96	53	63	36	103	64
CURRENT EMPLOYMENT								
Full-Time	52	72	90	94	37	48	106	116
Part-Time	59	45	48	45	45	36	65	54
Unemployed	73	45	78	45	68	35	84	57
Other ³	103	101	121	104	82	86	143	123

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.20D Received Substance Use Treatment in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.07	0.06	0.07	0.07	0.05	0.05	0.09	0.08
GENDER								
Male	0.09	0.09	0.12	0.11	0.07	0.08	0.14	0.13
Female	0.09	0.06	0.09	0.06	0.07	0.05	0.11	0.08
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.07	0.06	0.08	0.07	0.06	0.05	0.09	0.08
White	0.08	0.07	0.09	0.07	0.07	0.05	0.11	0.09
Black or African American	0.16	0.23	0.19	0.24	0.13	0.21	0.24	0.28
American Indian or Alaska Native	0.97	0.59	1.15	0.85	0.97	0.55	1.17	0.86
Native Hawaiian or Other Pacific Islander	0.62	0.82	0.62	0.51	0.62	0.40	1.13	0.91
Asian	0.15	0.05	0.24	0.10	0.15	0.05	0.25	0.10
Two or More Races	0.36	0.17	0.60	0.62	0.35	0.13	0.61	0.64
Hispanic or Latino	0.18	0.16	0.21	0.19	0.12	0.11	0.27	0.24
EDUCATION								
< High School	0.23	0.22	0.23	0.24	0.16	0.17	0.31	0.29
High School Graduate	0.11	0.12	0.13	0.12	0.08	0.10	0.15	0.15
Some College	0.12	0.11	0.15	0.12	0.11	0.08	0.17	0.14
College Graduate	0.10	0.06	0.13	0.07	0.09	0.05	0.14	0.09
CURRENT EMPLOYMENT								
Full-Time	0.04	0.06	0.08	0.08	0.03	0.04	0.09	0.09
Part-Time	0.17	0.14	0.14	0.14	0.13	0.11	0.19	0.16
Unemployed	0.61	0.41	0.65	0.41	0.57	0.32	0.70	0.51
Other ³	0.14	0.14	0.17	0.14	0.11	0.12	0.20	0.17

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.20P Received Substance Use Treatment in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.3156		0.8385		0.4938		0.8329	
GENDER								
Male	0.1901		0.8472		0.3446		0.5417	
Female	0.9373		0.5411		1.0000		0.6315	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.2827		0.8917		0.4522		0.6569	
White	0.7163		0.6729		0.9829		0.9714	
Black or African American	0.0331		0.2005		0.0252		0.0985	
American Indian or Alaska Native	0.8540		0.7683		0.5670		1.0000	
Native Hawaiian or Other Pacific Islander	0.7530		0.9237		0.5924		0.9552	
Asian	0.4537		0.2750		0.5354		0.1428	
Two or More Races	0.5951		0.5850		0.3947		0.6688	
Hispanic or Latino	0.9366		0.8480		0.9818		0.6889	
EDUCATION								
< High School	0.7043		0.3371		0.5807		0.4857	
High School Graduate	0.2996		0.6059		0.3318		0.5445	
Some College	0.4720		0.8170		0.8421		0.5286	
College Graduate	0.9097		0.0322		0.7180		0.0911	
CURRENT EMPLOYMENT								
Full-Time	0.0050		0.2334		0.0708		0.0366	
Part-Time	0.2444		0.4114		0.3735		0.2108	
Unemployed	0.2212		0.0265		0.0329		0.2036	
Other ³	0.1929		0.6288		0.0442		0.9442	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.21C Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	157	141	174	158	123	113	211	186
GEOGRAPHIC DIVISION								
Northeast	65	73	75	67	45	58	92	84
New England	38	29	31	27	20	22	47	34
Middle Atlantic	53	66	69	61	40	53	79	77
Midwest	55	59	67	58	42	44	82	78
East North Central	40	47	56	50	27	34	69	67
West North Central	38	34	40	31	33	26	47	42
South	86	87	106	100	73	70	121	118
South Atlantic	61	74	72	83	54	59	81	100
East South Central	38	24	37	21	24	17	48	31
West South Central	45	37	71	53	41	33	77	59
West	95	69	102	78	80	51	123	91
Mountain	46	29	38	42	12	24	58	44
Pacific	84	63	95	65	80	45	109	79
COUNTY TYPE								
Large Metro	120	119	133	125	98	96	160	152
Small Metro	67	70	95	91	52	55	110	107
250K – 1 Mil. Pop.	51	57	86	72	42	42	97	86
< 250K Pop.	43	38	39	53	30	34	51	62
Nonmetro	70	41	69	37	55	27	84	52
Urbanized	35	29	41	28	31	21	47	37
Less Urbanized	61	27	52	21	45	15	66	32
Completely Rural	8	10	22	12	2	8	23	17

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.21D Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.06	0.05	0.07	0.06	0.05	0.04	0.08	0.07
GEOGRAPHIC DIVISION								
Northeast	0.14	0.15	0.16	0.14	0.09	0.12	0.19	0.18
New England	0.31	0.23	0.25	0.22	0.16	0.18	0.38	0.27
Middle Atlantic	0.15	0.19	0.20	0.17	0.12	0.15	0.23	0.22
Midwest	0.10	0.10	0.12	0.10	0.07	0.08	0.15	0.14
East North Central	0.10	0.12	0.14	0.13	0.07	0.09	0.18	0.17
West North Central	0.22	0.19	0.23	0.18	0.19	0.15	0.27	0.24
South	0.09	0.09	0.11	0.10	0.07	0.07	0.12	0.12
South Atlantic	0.12	0.14	0.14	0.16	0.11	0.11	0.16	0.19
East South Central	0.25	0.16	0.24	0.13	0.16	0.11	0.31	0.20
West South Central	0.15	0.12	0.23	0.17	0.14	0.11	0.25	0.19
West	0.16	0.11	0.17	0.12	0.13	0.08	0.20	0.15
Mountain	0.24	0.15	0.20	0.22	0.06	0.13	0.31	0.23
Pacific	0.20	0.15	0.22	0.15	0.19	0.10	0.26	0.18
COUNTY TYPE								
Large Metro	0.09	0.08	0.09	0.08	0.07	0.06	0.11	0.10
Small Metro	0.09	0.09	0.12	0.12	0.07	0.07	0.14	0.14
250K – 1 Mil. Pop.	0.10	0.11	0.16	0.14	0.08	0.08	0.19	0.16
< 250K Pop.	0.17	0.15	0.15	0.21	0.12	0.14	0.20	0.24
Nonmetro	0.17	0.10	0.16	0.09	0.13	0.07	0.19	0.12
Urbanized	0.19	0.17	0.21	0.16	0.17	0.12	0.24	0.21
Less Urbanized	0.31	0.13	0.27	0.10	0.23	0.07	0.34	0.15
Completely Rural	0.18	0.23	0.47	0.28	0.04	0.20	0.50	0.39

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.21P Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.3401		0.7453		0.5827		0.8436	
GEOGRAPHIC DIVISION								
Northeast	0.2676		0.9620		0.1177		0.9187	
New England	0.8386		0.8089		0.5363		0.9146	
Middle Atlantic	0.1649		0.9556		0.1511		0.8633	
Midwest	0.8759		0.9652		0.8155		0.6398	
East North Central	0.4969		0.6833		0.4766		0.4488	
West North Central	0.5606		0.4994		0.6927		0.7564	
South	0.3169		0.9043		0.6886		0.3768	
South Atlantic	0.0723		0.2917		0.3385		0.0616	
East South Central	0.5110		0.3775		0.7410		0.3940	
West South Central	0.7241		0.4878		0.6201		0.6622	
West	0.7315		0.4580		0.4220		0.2995	
Mountain	0.7413		0.7085		0.0239		0.3730	
Pacific	0.5790		0.3134		0.1350		0.4839	
COUNTY TYPE								
Large Metro	0.3718		0.6823		0.5522		0.8122	
Small Metro	0.2916		0.5723		0.3853		0.3210	
250K – 1 Mil. Pop.	0.3044		0.8134		0.6655		0.8344	
< 250K Pop.	0.6653		0.1222		0.3739		0.1155	
Nonmetro	0.5026		0.3197		0.3394		0.6408	
Urbanized	0.7233		0.9325		0.8485		0.5907	
Less Urbanized	0.1741		0.1756		0.1519		0.2031	
Completely Rural	0.1776		0.8916		0.0696		0.5657	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.22C Substances for Which Last or Current Treatment Was Received among Persons Who Received Substance Use Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance for Which Last or Current Treatment Was Received ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	210	188	20	23	47	52	203	173
Marijuana	72	87	14	16	26	29	63	79
Cocaine	77	90	7	3	21	19	73	87
Heroin	69	75	*	5	19	22	66	70
Hallucinogens	48	53	7	4	18	17	42	51
Inhalants	39	39	4	4	14	13	35	38
Pain Relievers	85	72	7	6	22	26	82	64
Tranquilizers	57	65	5	7	17	21	53	61
Stimulants ²	73	60	6	7	17	20	70	56
Sedatives	41	38	*	6	10	9	39	36
Alcohol	168	150	14	13	33	34	164	144

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.22D Substances for Which Last or Current Treatment Was Received among Persons Who Received Substance Use Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Substance for Which Last or Current Treatment Was Received ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Marijuana	1.68	1.79	4.25	5.06	2.73	3.07	1.96	2.17
Cocaine	1.78	1.89	2.76	1.47	2.46	2.35	2.24	2.40
Heroin	1.57	1.64	*	2.31	2.17	2.49	2.01	2.02
Hallucinogens	1.15	1.19	2.82	1.72	2.16	2.06	1.36	1.50
Inhalants	0.94	0.91	1.79	1.73	1.79	1.60	1.11	1.16
Pain Relievers	1.90	1.58	2.77	2.59	2.41	2.81	2.43	1.90
Tranquilizers	1.35	1.48	2.11	3.03	2.00	2.52	1.68	1.84
Stimulants ²	1.66	1.37	2.34	2.82	1.99	2.47	2.12	1.69
Sedatives	1.00	0.89	*	2.47	1.21	1.14	1.27	1.12
Alcohol	2.29	2.19	4.34	4.58	2.87	3.15	2.85	2.71

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.22P Substances for Which Last or Current Treatment Was Received among Persons Who Received Substance Use Treatment in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance for Which Last or Current Treatment Was Received ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	N/A		N/A		N/A		N/A	
Marijuana	0.0974		0.6385		0.3426		0.0208	
Cocaine	0.0896		0.1110		0.2719		0.0321	
Heroin	0.4008		*		0.8388		0.4505	
Hallucinogens	0.6813		0.1521		0.4216		0.3583	
Inhalants	0.9624		0.8268		0.5423		0.8392	
Pain Relievers	0.9362		0.4958		0.7925		0.8187	
Tranquilizers	0.2551		0.5890		0.7960		0.2161	
Stimulants ²	0.5968		0.8995		0.8880		0.5613	
Sedatives	0.9218		*		0.7198		0.9120	
Alcohol	0.2035		0.1118		0.5146		0.3031	

*Low precision; no estimate reported.

N/A: Not Applicable.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.23C Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Location of Treatment¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	156	143	17	20	38	43	148	131
Hospital - Inpatient	65	64	6	7	19	19	61	60
Rehabilitation Facility - Inpatient	83	67	6	7	23	26	77	60
Rehabilitation Facility - Outpatient	104	88	10	9	29	29	97	83
Mental Health Center - Outpatient	70	73	9	9	20	21	66	69
Emergency Room	44	44	5	6	10	13	42	42
Private Doctor's Office	66	63	5	7	17	16	64	60
Self-Help Group	116	99	9	10	28	32	111	93
Prison/Jail	33	45	3	4	10	12	31	43

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment refers to treatment received in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.23D Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Hospital - Inpatient	2.60	2.30	3.46	4.03	3.11	3.28	3.63	2.97
Rehabilitation Facility - Inpatient	3.15	2.45	3.30	3.91	3.48	4.06	4.31	3.11
Rehabilitation Facility - Outpatient	3.31	2.65	4.59	4.82	3.49	4.04	4.61	3.47
Mental Health Center - Outpatient	2.68	2.49	4.38	4.77	3.20	3.44	3.71	3.30
Emergency Room	1.89	1.68	2.86	3.49	1.91	2.46	2.65	2.22
Private Doctor's Office	2.68	2.22	2.84	3.95	2.90	2.88	3.68	2.93
Self-Help Group	3.33	2.83	4.33	5.32	3.49	4.04	4.61	3.67
Prison/Jail	1.47	1.74	1.82	2.16	1.83	2.41	2.02	2.28

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment refers to treatment received in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.23P Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	N/A		N/A		N/A		N/A	
Hospital - Inpatient	0.6648		0.9354		0.5892		0.8055	
Rehabilitation Facility - Inpatient	0.6370		0.9856		0.9595		0.6072	
Rehabilitation Facility - Outpatient	0.2483		0.1942		0.1323		0.5147	
Mental Health Center - Outpatient	0.4838		0.6556		0.6143		0.5511	
Emergency Room	0.7669		0.8302		0.4880		0.6103	
Private Doctor's Office	0.7579		0.5109		0.4364		0.8081	
Self-Help Group	0.2704		0.8768		0.9379		0.2115	
Prison/Jail	0.1255		0.9894		0.8780		0.1059	

*Low precision; no estimate reported.

N/A: Not Applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment refers to treatment received in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.24C Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	173	159	16	*	36	36	168	151
Hospital - Inpatient	80	67	8	*	18	16	78	62
Rehabilitation Facility - Inpatient	86	62	8	*	17	17	83	57
Rehabilitation Facility - Outpatient	110	95	10	*	23	24	107	89
Mental Health Center - Outpatient	84	77	*	*	16	18	82	73
Emergency Room	58	50	*	*	10	9	57	48
Private Doctor's Office	65	60	6	*	13	10	63	58
Self-Help Group	151	117	10	*	25	26	148	112
Prison/Jail	33	33	5	*	9	9	31	32

*Low precision; no estimate reported.

NOTE: Received Alcohol Treatment refers to treatment received in order to reduce or stop alcohol use, or for medical problems associated with alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.24D Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	0.00	0.00	0.00	*	0.00	0.00	0.00	0.00
Hospital - Inpatient	2.62	2.20	4.96	*	3.23	3.35	3.31	2.56
Rehabilitation Facility - Inpatient	2.76	2.17	4.81	*	3.18	3.54	3.45	2.52
Rehabilitation Facility - Outpatient	3.16	2.68	5.76	*	3.83	4.15	3.98	3.16
Mental Health Center - Outpatient	2.80	2.37	*	*	3.02	3.68	3.51	2.83
Emergency Room	2.02	1.73	*	*	2.08	1.95	2.57	2.08
Private Doctor's Office	2.23	1.99	4.00	*	2.63	2.26	2.80	2.40
Self-Help Group	3.08	2.74	5.52	*	3.75	3.98	3.74	3.25
Prison/Jail	1.18	1.21	3.64	*	1.80	2.06	1.42	1.43

*Low precision; no estimate reported.

NOTE: Received Alcohol Treatment refers to treatment received in order to reduce or stop alcohol use, or for medical problems associated with alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.24P Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	N/A		N/A		N/A		N/A	
Hospital - Inpatient	0.8585		*		0.7752		0.9798	
Rehabilitation Facility - Inpatient	0.9396		*		0.9581		0.9265	
Rehabilitation Facility - Outpatient	0.8715		*		0.6157		0.9333	
Mental Health Center - Outpatient	0.4224		*		0.0948		0.6122	
Emergency Room	0.8085		*		0.5759		0.8419	
Private Doctor's Office	0.5291		*		0.5975		0.6040	
Self-Help Group	0.3534		*		0.2785		0.1929	
Prison/Jail	0.4501		*		0.7338		0.4115	

*Low precision; no estimate reported.

N/A: Not Applicable.

NOTE: Received Alcohol Treatment refers to treatment received in order to reduce or stop alcohol use, or for medical problems associated with alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.25C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Location of Treatment¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	210	188	20	23	47	52	203	173
Hospital - Inpatient	93	85	9	12	22	22	91	79
Rehabilitation Facility - Inpatient	107	85	9	10	24	29	101	76
Rehabilitation Facility - Outpatient	134	118	13	11	31	34	128	110
Mental Health Center - Outpatient	109	94	11	11	22	24	105	89
Emergency Room	74	60	9	12	16	16	71	57
Private Doctor's Office	86	78	7	11	19	18	83	75
Self-Help Group	172	136	12	13	31	38	169	130
Prison/Jail	42	51	6	8	11	13	40	49

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.25D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Location of Treatment¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Hospital - Inpatient	2.04	1.80	3.54	4.39	2.42	2.52	2.65	2.20
Rehabilitation Facility - Inpatient	2.29	1.87	3.56	4.09	2.60	3.11	2.89	2.26
Rehabilitation Facility - Outpatient	2.47	2.01	4.29	4.35	2.91	3.20	3.17	2.52
Mental Health Center - Outpatient	2.29	1.88	4.04	4.34	2.41	2.68	2.95	2.35
Emergency Room	1.69	1.37	3.42	4.41	1.94	2.00	2.19	1.69
Private Doctor's Office	1.91	1.65	2.86	4.14	2.13	2.10	2.48	2.08
Self-Help Group	2.44	2.16	4.31	4.63	2.79	3.21	3.07	2.71
Prison/Jail	1.03	1.17	2.52	3.22	1.37	1.66	1.30	1.47

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.25P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	N/A		N/A		N/A		N/A	
Hospital - Inpatient	0.8500		0.4716		0.8192		0.9635	
Rehabilitation Facility - Inpatient	0.9439		0.9176		0.7319		0.9919	
Rehabilitation Facility - Outpatient	0.6308		0.0433		0.4843		0.9221	
Mental Health Center - Outpatient	0.7013		0.4745		0.3579		0.5880	
Emergency Room	0.2882		0.6352		0.7949		0.2680	
Private Doctor's Office	0.9365		0.1782		0.4546		0.9374	
Self-Help Group	0.3092		0.5815		0.3213		0.1966	
Prison/Jail	0.1357		0.9936		0.5959		0.0878	

*Low precision; no estimate reported.

N/A: Not Applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 5.26C Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Dependence or Abuse Status	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol (2013)	Both Illicit Drugs and Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	157	141	174	158	123	113	211	186
DEPENDENCE								
None	127	106	145	118	104	83	172	144
Both Illicit Drugs and Alcohol	44	46	42	47	41	46	47	48
Illicit Drugs or Alcohol	82	90	99	98	63	72	120	115
Illicit Drugs	77	75	61	64	58	57	79	84
Alcohol	53	67	89	89	47	63	96	93
DEPENDENCE OR ABUSE								
None	123	97	141	102	101	74	166	129
Both Illicit Drugs and Alcohol	51	51	50	63	46	50	58	64
Illicit Drugs or Alcohol	87	97	107	113	68	79	132	130
Illicit Drugs	80	79	64	79	60	63	84	93
Alcohol	60	75	98	104	53	69	110	112

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 5.26D Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status:
Standard Errors of Percentages, 2013 and 2014**

Dependence or Abuse Status	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol (2013)	Both Illicit Drugs and Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.06	0.05	0.07	0.06	0.05	0.04	0.08	0.07
DEPENDENCE								
None	0.05	0.04	0.06	0.05	0.04	0.03	0.07	0.06
Both Illicit Drugs and Alcohol	3.56	3.42	3.43	3.48	3.36	3.45	3.71	3.51
Illicit Drugs or Alcohol	0.68	0.72	0.81	0.78	0.53	0.58	0.97	0.88
Illicit Drugs	1.49	1.38	1.19	1.19	1.15	1.08	1.53	1.47
Alcohol	0.65	0.80	1.07	1.03	0.59	0.75	1.14	1.07
DEPENDENCE OR ABUSE								
None	0.05	0.04	0.06	0.04	0.04	0.03	0.07	0.05
Both Illicit Drugs and Alcohol	1.84	1.74	1.82	2.11	1.69	1.70	2.05	2.04
Illicit Drugs or Alcohol	0.40	0.43	0.48	0.51	0.31	0.36	0.59	0.56
Illicit Drugs	1.11	1.05	0.91	1.05	0.85	0.85	1.16	1.17
Alcohol	0.34	0.43	0.55	0.58	0.31	0.40	0.61	0.63

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 5.26P Received Substance Use Treatment in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Dependence or Abuse Status	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol (2013)	Both Illicit Drugs and Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.3401		0.7453		0.5827		0.8436	
DEPENDENCE								
None	0.7954		0.3908		0.9974		0.6985	
Both Illicit Drugs and Alcohol	0.1098		0.0246		0.0279		0.0891	
Illicit Drugs or Alcohol	0.1821		0.5439		0.3347		0.3753	
Illicit Drugs	0.3208		0.5125		0.7083		0.1600	
Alcohol	0.0967		0.2284		0.0244		0.5164	
DEPENDENCE OR ABUSE								
None	0.8602		0.1216		0.6196		0.4083	
Both Illicit Drugs and Alcohol	0.4416		0.0391		0.1616		0.1214	
Illicit Drugs or Alcohol	0.0534		0.1597		0.1122		0.1438	
Illicit Drugs	0.4403		0.1896		0.6051		0.0977	
Alcohol	0.0452		0.0951		0.0221		0.2844	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.27C Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	123	108	125	121	75	73	160	148
12	*	1	2	3	*	1	2	3
13	3	*	4	1	3	*	4	1
14	2	5	5	1	2	1	5	5
15	5	5	4	4	3	3	6	6
16	8	7	7	5	7	3	9	8
17	7	7	6	9	5	4	8	12
18	11	6	8	5	6	3	12	8
19	10	8	7	9	6	5	11	12
20	8	11	6	8	6	8	8	11
21	11	11	11	8	10	7	13	12
22	10	10	6	10	6	7	11	13
23	13	15	12	13	8	11	15	17
24	9	12	8	10	5	6	11	14
25	12	16	10	11	9	7	12	18
26-29	47	40	41	35	34	26	53	47
30-34	45	34	43	31	32	20	53	41
35-39	45	33	46	29	37	19	53	41
40-44	32	38	30	35	14	22	44	49
45-49	25	23	32	32	10	18	40	35
50-54	31	36	45	35	18	23	51	50
55-59	40	48	23	43	6	32	50	57
60-64	43	13	58	47	26	13	70	47
65 or Older	*	21	39	38	*	21	39	38

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.27D Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.05	0.04	0.05	0.05	0.03	0.03	0.06	0.06
12	*	0.04	0.06	0.07	*	0.04	0.06	0.07
13	0.08	*	0.08	0.03	0.07	*	0.09	0.03
14	0.05	0.11	0.10	0.03	0.05	0.03	0.11	0.11
15	0.12	0.12	0.09	0.10	0.06	0.08	0.14	0.13
16	0.18	0.16	0.17	0.10	0.16	0.07	0.20	0.18
17	0.17	0.18	0.16	0.22	0.13	0.10	0.20	0.27
18	0.22	0.13	0.18	0.11	0.13	0.06	0.26	0.17
19	0.23	0.18	0.18	0.21	0.13	0.12	0.27	0.29
20	0.19	0.24	0.14	0.19	0.14	0.17	0.19	0.26
21	0.27	0.26	0.25	0.18	0.23	0.15	0.31	0.28
22	0.22	0.23	0.14	0.21	0.13	0.15	0.24	0.28
23	0.29	0.33	0.26	0.29	0.19	0.26	0.34	0.38
24	0.19	0.26	0.17	0.23	0.12	0.13	0.24	0.33
25	0.29	0.37	0.23	0.26	0.22	0.16	0.30	0.43
26-29	0.27	0.23	0.24	0.21	0.20	0.15	0.30	0.27
30-34	0.22	0.16	0.21	0.15	0.16	0.10	0.26	0.20
35-39	0.24	0.17	0.24	0.15	0.19	0.09	0.28	0.21
40-44	0.15	0.19	0.14	0.17	0.07	0.11	0.20	0.24
45-49	0.13	0.11	0.16	0.16	0.05	0.09	0.20	0.17
50-54	0.14	0.15	0.19	0.15	0.08	0.10	0.22	0.21
55-59	0.19	0.23	0.11	0.20	0.03	0.15	0.24	0.27
60-64	0.25	0.08	0.33	0.26	0.15	0.08	0.40	0.26
65 or Older	*	0.05	0.09	0.08	*	0.05	0.09	0.08

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.27P Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.5221		0.4371		0.2112		0.6008	
12	*		0.6995		*		0.6815	
13	*		0.1234		*		0.0733	
14	0.2618		0.3346		0.9338		0.9922	
15	0.9759		0.6863		0.4884		0.7343	
16	0.2640		0.1205		0.0394		0.2895	
17	0.5098		0.9075		0.2159		0.6868	
18	0.0316		0.0899		0.1146		0.0393	
19	0.1743		0.9322		0.8865		0.5614	
20	0.8567		0.9957		0.9336		0.9313	
21	0.5916		0.5236		0.3538		0.8264	
22	0.8597		0.1999		0.7679		0.8134	
23	0.5456		0.8877		0.3750		0.8752	
24	0.9645		0.4724		0.8208		0.5642	
25	0.7350		0.6087		0.3934		0.2374	
26-29	0.8650		0.8090		0.9153		0.6906	
30-34	0.7912		0.9378		0.6854		0.9809	
35-39	0.6033		0.6230		0.7407		0.6811	
40-44	0.2311		0.1341		0.0401		0.2704	
45-49	0.7553		0.5328		0.1293		0.9780	
50-54	0.2928		0.6185		0.1854		0.3196	
55-59	0.3331		0.1429		0.1266		0.3300	
60-64	0.0766		0.6421		0.6799		0.1504	
65 or Older	*		0.5646		*		0.5231	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.28C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	123	108	125	121	75	73	160	148
AGE								
12-17	12	12	13	12	10	6	15	16
18-25	31	34	26	27	22	20	35	41
26 or Older	115	100	121	115	70	69	153	139
GENDER								
Male	87	88	100	108	43	65	127	126
Female	83	56	77	51	61	32	99	73
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	107	97	113	106	71	65	143	134
White	101	83	99	90	63	51	129	114
Black or African American	37	45	44	50	22	37	56	62
American Indian or Alaska Native	12	8	13	9	12	8	13	9
Native Hawaiian or Other Pacific Islander	4	*	4	*	4	*	4	8
Asian	18	4	23	7	18	4	23	7
Two or More Races	3	6	14	14	2	5	18	15
Hispanic or Latino	57	51	56	52	23	28	77	66

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.28D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.05	0.04	0.05	0.05	0.03	0.03	0.06	0.06
AGE								
12-17	0.05	0.05	0.05	0.05	0.04	0.03	0.06	0.06
18-25	0.09	0.10	0.07	0.08	0.06	0.06	0.10	0.12
26 or Older	0.06	0.05	0.06	0.06	0.03	0.03	0.08	0.07
GENDER								
Male	0.07	0.07	0.08	0.08	0.03	0.05	0.10	0.10
Female	0.06	0.04	0.06	0.04	0.05	0.02	0.07	0.05
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.05	0.04	0.05	0.05	0.03	0.03	0.06	0.06
White	0.06	0.05	0.06	0.05	0.04	0.03	0.08	0.07
Black or African American	0.12	0.14	0.14	0.16	0.07	0.12	0.18	0.19
American Indian or Alaska Native	0.85	0.53	0.91	0.56	0.84	0.49	0.92	0.59
Native Hawaiian or Other Pacific Islander	0.48	*	0.48	*	0.48	*	0.48	0.74
Asian	0.14	0.03	0.17	0.05	0.14	0.03	0.17	0.05
Two or More Races	0.08	0.14	0.32	0.31	0.05	0.11	0.42	0.32
Hispanic or Latino	0.14	0.12	0.14	0.12	0.06	0.06	0.19	0.15

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.28P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.5221		0.4371		0.2112		0.6008	
AGE								
12-17	0.3555		0.2913		0.0433		0.5349	
18-25	0.7569		0.7881		0.5916		0.8445	
26 or Older	0.3745		0.3941		0.0862		0.5729	
GENDER								
Male	0.1618		0.3254		0.0217		0.4572	
Female	0.4966		0.9157		0.4804		0.8790	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.5671		0.4370		0.2170		0.5746	
White	0.9299		0.3791		0.5033		0.6829	
Black or African American	0.1647		0.4632		0.0525		0.3045	
American Indian or Alaska Native	0.9617		0.6644		0.7406		0.9042	
Native Hawaiian or Other Pacific Islander	*		*		*		0.6394	
Asian	0.3046		0.2480		0.3708		0.2088	
Two or More Races	0.1027		0.6855		0.1698		0.5219	
Hispanic or Latino	0.7723		0.8459		0.7291		0.9660	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.29C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	12	12	13	12	10	6	15	16
GENDER								
Male	9	7	11	6	8	4	12	9
Female	8	10	7	10	6	5	9	13
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	11	10	11	11	8	6	13	14
White	10	10	10	11	7	6	12	14
Black or African American	3	3	4	4	2	1	5	5
American Indian or Alaska Native	2	1	3	1	*	*	3	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	*	*	*	*	*	*	*
Two or More Races	2	1	1	1	1	1	2	1
Hispanic or Latino	6	7	6	2	5	1	7	7
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	8	7	9	5	7	3	10	8
Female, White, Not Hispanic	6	7	4	9	3	5	6	11
Male, Black, Not Hispanic	2	2	4	4	1	1	5	4
Female, Black, Not Hispanic	1	2	1	*	1	*	1	2
Male, Hispanic	4	2	3	1	3	1	4	2
Female, Hispanic	5	7	5	1	4	*	6	7

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.29D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.05	0.05	0.05	0.05	0.04	0.03	0.06	0.06
GENDER								
Male	0.07	0.06	0.08	0.05	0.06	0.03	0.09	0.07
Female	0.07	0.08	0.06	0.08	0.05	0.04	0.08	0.11
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.05	0.05	0.06	0.06	0.04	0.03	0.07	0.07
White	0.07	0.07	0.07	0.08	0.05	0.04	0.09	0.10
Black or African American	0.08	0.08	0.12	0.11	0.05	0.04	0.14	0.13
American Indian or Alaska Native	1.40	0.71	1.69	0.57	*	*	1.71	0.73
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	*	*	*	*	*	*	*	*
Two or More Races	0.21	0.18	0.20	0.17	0.18	0.17	0.22	0.18
Hispanic or Latino	0.11	0.12	0.10	0.03	0.09	0.03	0.13	0.13
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	0.11	0.10	0.13	0.07	0.10	0.05	0.14	0.12
Female, White, Not Hispanic	0.09	0.11	0.05	0.14	0.04	0.07	0.10	0.16
Male, Black, Not Hispanic	0.12	0.10	0.23	0.22	0.04	0.08	0.26	0.23
Female, Black, Not Hispanic	0.09	0.12	0.09	*	0.09	*	0.09	0.12
Male, Hispanic	0.13	0.07	0.10	0.05	0.10	0.05	0.14	0.07
Female, Hispanic	0.19	0.24	0.17	0.05	0.15	*	0.22	0.25

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.29P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.3555		0.2913		0.0433		0.5349	
GENDER								
Male	0.1413		0.0359		0.0348		0.0893	
Female	0.9418		0.5028		0.5013		0.4584	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.4694		0.9358		0.3187		0.9300	
White	0.6673		0.6417		0.5755		0.7366	
Black or African American	0.7562		0.7106		0.5178		0.7671	
American Indian or Alaska Native	0.5635		0.2472		*		0.3440	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	*		*		*		*	
Two or More Races	0.9467		0.7983		0.8903		0.6773	
Hispanic or Latino	0.5593		0.0048		0.0123		0.2376	
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	0.4786		0.0963		0.1080		0.4253	
Female, White, Not Hispanic	0.8852		0.0204		0.2112		0.2035	
Male, Black, Not Hispanic	0.6779		0.9766		0.6978		0.7510	
Female, Black, Not Hispanic	0.9964		*		*		0.9964	
Male, Hispanic	0.1861		0.1597		0.2162		0.0550	
Female, Hispanic	0.9852		0.0137		*		0.7219	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.30C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	31	34	26	27	22	20	35	41
GENDER								
Male	26	27	23	23	19	17	30	33
Female	15	20	10	15	9	10	16	23
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	28	32	22	26	19	20	31	37
White	27	29	20	22	18	18	30	33
Black or African American	6	9	5	9	2	7	8	11
American Indian or Alaska Native	1	3	2	3	1	*	2	3
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	5	*	5	6	5	*	5	6
Two or More Races	2	6	3	5	2	5	3	6
Hispanic or Latino	10	14	12	7	8	1	14	16
EDUCATION								
< High School	12	14	11	11	5	8	17	17
High School Graduate	23	18	19	16	18	12	24	21
Some College	17	21	13	17	13	13	17	23
College Graduate	3	8	4	8	0	4	5	12
CURRENT EMPLOYMENT								
Full-Time	14	20	13	16	10	12	17	24
Part-Time	16	15	12	14	11	10	18	18
Unemployed	16	18	13	13	11	11	17	20
Other ³	13	12	11	10	9	5	14	16

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.30D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.09	0.10	0.07	0.08	0.06	0.06	0.10	0.12
GENDER								
Male	0.15	0.15	0.13	0.13	0.11	0.10	0.17	0.19
Female	0.09	0.11	0.06	0.09	0.05	0.06	0.09	0.13
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.10	0.11	0.08	0.09	0.07	0.07	0.11	0.13
White	0.14	0.15	0.10	0.12	0.09	0.09	0.15	0.17
Black or African American	0.12	0.18	0.11	0.18	0.04	0.14	0.15	0.22
American Indian or Alaska Native	0.41	1.47	1.02	1.49	0.41	*	1.02	1.51
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	0.28	*	0.25	0.28	0.25	*	0.28	0.28
Two or More Races	0.27	0.70	0.43	0.60	0.24	0.59	0.45	0.71
Hispanic or Latino	0.14	0.19	0.16	0.10	0.11	0.02	0.19	0.21
EDUCATION								
< High School	0.21	0.27	0.20	0.22	0.09	0.17	0.29	0.34
High School Graduate	0.19	0.15	0.16	0.13	0.15	0.10	0.20	0.18
Some College	0.14	0.17	0.11	0.13	0.10	0.10	0.14	0.19
College Graduate	0.06	0.14	0.08	0.14	0.01	0.06	0.11	0.21
CURRENT EMPLOYMENT								
Full-Time	0.10	0.14	0.10	0.11	0.07	0.09	0.13	0.16
Part-Time	0.16	0.15	0.13	0.15	0.11	0.10	0.19	0.19
Unemployed	0.39	0.46	0.30	0.32	0.28	0.28	0.41	0.52
Other ³	0.17	0.16	0.14	0.14	0.12	0.06	0.19	0.21

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.30P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.7569		0.7881		0.5916		0.8445	
GENDER								
Male	0.4764		0.8106		0.5477		0.5847	
Female	0.6110		0.3220		0.9796		0.2142	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.6505		0.2850		0.9187		0.7256	
White	0.4345		0.5407		0.6035		0.9713	
Black or African American	0.4384		0.2053		0.0768		0.4239	
American Indian or Alaska Native	0.2856		0.8556		*		0.7592	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	*		0.6464		*		0.9369	
Two or More Races	0.1277		0.8482		0.2039		0.5280	
Hispanic or Latino	0.7645		0.0579		0.0224		0.7744	
EDUCATION								
< High School	0.8033		0.5628		0.4460		0.6221	
High School Graduate	0.1693		0.4856		0.1731		0.4053	
Some College	0.5064		0.2581		0.8480		0.2022	
College Graduate	0.1933		0.0720		0.2219		0.0714	
CURRENT EMPLOYMENT								
Full-Time	0.4086		0.9635		0.9634		0.6013	
Part-Time	0.3945		0.5144		0.9216		0.5640	
Unemployed	0.6442		0.4688		0.5176		0.9137	
Other ³	0.9436		0.3462		0.7431		0.4109	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.31C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	115	100	121	115	70	69	153	139
GENDER								
Male	80	83	96	105	38	63	121	120
Female	80	52	76	47	60	29	96	67
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	100	89	109	99	67	61	136	124
White	93	76	96	84	60	47	122	105
Black or African American	37	45	44	49	22	37	56	61
American Indian or Alaska Native	*	7	12	8	*	6	13	8
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	7
Asian	17	4	23	4	17	4	23	4
Two or More Races	2	2	13	13	*	1	17	13
Hispanic or Latino	56	48	55	51	21	28	76	65
EDUCATION								
< High School	62	48	46	59	24	29	75	74
High School Graduate	58	65	61	71	40	49	74	83
Some College	64	50	75	48	49	29	92	66
College Graduate	37	36	59	35	18	19	67	45
CURRENT EMPLOYMENT								
Full-Time	40	55	58	59	21	25	64	79
Part-Time	40	35	28	31	27	26	42	38
Unemployed	65	25	58	34	51	14	71	44
Other ³	72	74	85	81	34	55	111	98

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.31D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.06	0.05	0.06	0.06	0.03	0.03	0.08	0.07
GENDER								
Male	0.08	0.08	0.10	0.11	0.04	0.06	0.12	0.12
Female	0.08	0.05	0.07	0.04	0.06	0.03	0.09	0.06
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.06	0.05	0.06	0.06	0.04	0.03	0.08	0.07
White	0.07	0.05	0.07	0.06	0.04	0.03	0.09	0.08
Black or African American	0.16	0.19	0.19	0.21	0.10	0.16	0.24	0.26
American Indian or Alaska Native	*	0.61	1.22	0.66	*	0.57	1.24	0.70
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	1.01
Asian	0.17	0.04	0.22	0.04	0.17	0.04	0.22	0.04
Two or More Races	0.08	0.08	0.47	0.45	*	0.02	0.63	0.46
Hispanic or Latino	0.20	0.16	0.19	0.17	0.08	0.09	0.27	0.22
EDUCATION								
< High School	0.23	0.18	0.17	0.21	0.09	0.11	0.28	0.27
High School Graduate	0.10	0.11	0.11	0.12	0.07	0.09	0.13	0.14
Some College	0.12	0.09	0.15	0.09	0.09	0.05	0.18	0.12
College Graduate	0.06	0.05	0.09	0.05	0.03	0.03	0.10	0.07
CURRENT EMPLOYMENT								
Full-Time	0.04	0.05	0.05	0.05	0.02	0.02	0.06	0.07
Part-Time	0.17	0.15	0.12	0.13	0.11	0.11	0.18	0.16
Unemployed	0.84	0.36	0.75	0.49	0.66	0.20	0.92	0.63
Other ³	0.11	0.11	0.13	0.12	0.05	0.08	0.17	0.15

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.31P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.3745		0.3941		0.0862		0.5729	
GENDER								
Male	0.0605		0.2105		0.0035		0.2854	
Female	0.3931		0.6892		0.5160		0.5783	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.3999		0.5658		0.1638		0.6092	
White	0.6828		0.4685		0.3359		0.6937	
Black or African American	0.1983		0.5664		0.0865		0.3490	
American Indian or Alaska Native	*		0.7973		*		0.9967	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.5374		0.1731		0.5374		0.1731	
Two or More Races	0.4302		0.6423		*		0.4115	
Hispanic or Latino	0.7629		0.4488		0.2057		0.8287	
EDUCATION								
< High School	0.9185		0.0602		0.1753		0.3311	
High School Graduate	0.2023		0.0903		0.0714		0.1507	
Some College	0.9667		0.4484		0.7919		0.6287	
College Graduate	0.8031		0.1980		0.4527		0.2496	
CURRENT EMPLOYMENT								
Full-Time	0.0137		0.1371		0.0976		0.0090	
Part-Time	0.8417		0.8795		0.8478		0.5954	
Unemployed	0.0391		0.2663		0.0566		0.2342	
Other ³	0.1763		0.3167		0.0009		0.9329	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.32C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	122	107	124	120	74	72	159	148
GENDER								
Male	86	88	99	108	42	65	126	125
Female	82	55	77	51	61	32	98	72
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	107	96	112	105	70	64	142	133
White	99	83	98	89	63	51	127	114
Black or African American	37	45	44	50	22	37	56	63
American Indian or Alaska Native	12	8	13	9	*	7	13	9
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	8
Asian	18	4	23	7	18	4	23	7
Two or More Races	3	6	13	14	2	5	18	14
Hispanic or Latino	57	50	56	52	23	28	77	65
EDUCATION								
< High School	64	50	47	59	25	31	77	75
High School Graduate	62	67	64	73	43	51	78	86
Some College	67	55	76	53	50	32	94	72
College Graduate	37	37	59	36	18	20	67	47
CURRENT EMPLOYMENT								
Full-Time	42	59	59	62	22	28	67	84
Part-Time	44	39	31	35	29	28	46	44
Unemployed	68	31	59	36	52	18	74	49
Other ³	73	75	86	82	35	55	112	100

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.32D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.05	0.04	0.05	0.05	0.03	0.03	0.07	0.06
GENDER								
Male	0.08	0.08	0.09	0.09	0.04	0.06	0.11	0.11
Female	0.07	0.04	0.06	0.04	0.05	0.03	0.08	0.06
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.05	0.05	0.06	0.05	0.03	0.03	0.07	0.07
White	0.06	0.05	0.06	0.06	0.04	0.03	0.08	0.07
Black or African American	0.13	0.16	0.16	0.17	0.08	0.13	0.20	0.22
American Indian or Alaska Native	0.95	0.59	1.01	0.63	*	0.55	1.02	0.66
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	0.82
Asian	0.15	0.03	0.19	0.05	0.15	0.03	0.19	0.05
Two or More Races	0.09	0.16	0.38	0.38	0.05	0.13	0.51	0.39
Hispanic or Latino	0.16	0.14	0.16	0.14	0.06	0.07	0.22	0.18
EDUCATION								
< High School	0.20	0.16	0.15	0.18	0.08	0.10	0.24	0.23
High School Graduate	0.09	0.10	0.09	0.11	0.06	0.07	0.11	0.12
Some College	0.11	0.08	0.12	0.08	0.08	0.05	0.15	0.11
College Graduate	0.05	0.05	0.08	0.05	0.03	0.03	0.09	0.06
CURRENT EMPLOYMENT								
Full-Time	0.04	0.05	0.05	0.05	0.02	0.02	0.06	0.07
Part-Time	0.13	0.12	0.09	0.11	0.09	0.08	0.14	0.13
Unemployed	0.57	0.29	0.50	0.33	0.44	0.16	0.62	0.44
Other ³	0.10	0.10	0.12	0.11	0.05	0.07	0.15	0.14

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.32P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Illicit Drugs ¹ (2013)	Illicit Drugs ¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ¹ (2013)	Both Illicit Drugs and Alcohol ¹ (2014)	Illicit Drugs or Alcohol ^{1,2} (2013)	Illicit Drugs or Alcohol ^{1,2} (2014)
TOTAL	0.4600		0.3781		0.1380		0.5609	
GENDER								
Male	0.1238		0.2455		0.0108		0.3766	
Female	0.4859		0.8467		0.5248		0.7997	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.5182		0.4315		0.1774		0.5701	
White	0.8958		0.4062		0.4607		0.7105	
Black or African American	0.1566		0.4407		0.0469		0.2914	
American Indian or Alaska Native	0.9552		0.8290		*		0.9485	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.3037		0.2466		0.3698		0.2076	
Two or More Races	0.0806		0.7008		0.1553		0.5433	
Hispanic or Latino	0.7180		0.6822		0.4715		0.8756	
EDUCATION								
< High School	0.8935		0.0851		0.1352		0.4154	
High School Graduate	0.4332		0.1435		0.1986		0.2477	
Some College	0.8102		0.6387		0.8451		0.8811	
College Graduate	0.6525		0.3006		0.3684		0.3924	
CURRENT EMPLOYMENT								
Full-Time	0.0087		0.1543		0.1241		0.0083	
Part-Time	0.6208		0.9012		0.8857		0.4844	
Unemployed	0.0388		0.2037		0.0453		0.2429	
Other ³	0.1876		0.2666		0.0012		0.8414	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.33C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	123	108	125	121	75	73	160	148
GEOGRAPHIC DIVISION								
Northeast	56	52	58	60	22	33	76	76
New England	27	24	16	27	5	17	31	32
Middle Atlantic	49	46	55	54	22	29	70	69
Midwest	43	48	34	45	17	34	58	59
East North Central	33	37	28	35	14	23	45	48
West North Central	27	31	19	29	10	25	38	36
South	70	69	84	69	50	44	98	91
South Atlantic	53	58	60	62	41	39	71	79
East South Central	37	21	33	15	17	13	47	23
West South Central	27	28	49	27	23	17	50	37
West	69	51	64	58	48	28	83	70
Mountain	39	23	25	37	6	17	46	38
Pacific	58	45	59	44	48	21	69	58
COUNTY TYPE								
Large Metro	93	91	94	100	58	65	121	122
Small Metro	48	52	67	63	26	31	81	77
250K – 1 Mil. Pop.	33	45	62	55	23	25	68	69
< 250K Pop.	36	24	23	28	13	15	44	32
Nonmetro	61	31	50	27	39	13	68	40
Urbanized	26	21	23	21	13	8	32	29
Less Urbanized	55	22	39	15	37	8	56	27
Completely Rural	8	7	21	10	1	5	22	11

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.33D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.05	0.04	0.05	0.05	0.03	0.03	0.06	0.06
GEOGRAPHIC DIVISION								
Northeast	0.12	0.11	0.12	0.13	0.05	0.07	0.16	0.16
New England	0.22	0.19	0.13	0.21	0.04	0.13	0.25	0.25
Middle Atlantic	0.14	0.13	0.16	0.15	0.06	0.08	0.20	0.20
Midwest	0.08	0.09	0.06	0.08	0.03	0.06	0.10	0.10
East North Central	0.09	0.10	0.07	0.09	0.03	0.06	0.12	0.12
West North Central	0.16	0.18	0.11	0.17	0.06	0.14	0.22	0.21
South	0.07	0.07	0.09	0.07	0.05	0.04	0.10	0.09
South Atlantic	0.10	0.11	0.12	0.12	0.08	0.07	0.14	0.15
East South Central	0.24	0.14	0.21	0.10	0.11	0.08	0.30	0.15
West South Central	0.09	0.09	0.16	0.09	0.08	0.05	0.16	0.12
West	0.11	0.08	0.10	0.09	0.08	0.04	0.14	0.11
Mountain	0.21	0.12	0.13	0.19	0.03	0.09	0.25	0.20
Pacific	0.14	0.10	0.14	0.10	0.11	0.05	0.16	0.14
COUNTY TYPE								
Large Metro	0.07	0.06	0.07	0.07	0.04	0.04	0.09	0.08
Small Metro	0.06	0.07	0.09	0.08	0.03	0.04	0.10	0.10
250K – 1 Mil. Pop.	0.06	0.08	0.12	0.10	0.04	0.05	0.13	0.13
< 250K Pop.	0.14	0.10	0.09	0.11	0.05	0.06	0.17	0.13
Nonmetro	0.14	0.07	0.12	0.07	0.09	0.03	0.16	0.09
Urbanized	0.14	0.12	0.12	0.12	0.07	0.05	0.17	0.16
Less Urbanized	0.29	0.10	0.20	0.07	0.19	0.04	0.29	0.13
Completely Rural	0.17	0.16	0.46	0.23	0.03	0.12	0.49	0.25

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.33P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Illicit Drugs¹ (2013)	Illicit Drugs¹ (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol¹ (2013)	Both Illicit Drugs and Alcohol¹ (2014)	Illicit Drugs or Alcohol^{1,2} (2013)	Illicit Drugs or Alcohol^{1,2} (2014)
TOTAL	0.5221		0.4371		0.2112		0.6008	
GEOGRAPHIC DIVISION								
Northeast	0.4909		0.2768		0.0158		0.5329	
New England	0.9933		0.2271		0.0868		0.8483	
Middle Atlantic	0.4356		0.4932		0.0652		0.5495	
Midwest	0.4769		0.1209		0.0366		0.4891	
East North Central	0.3000		0.2084		0.0691		0.2382	
West North Central	0.8843		0.3649		0.2545		0.7017	
South	0.6088		0.7467		0.7101		0.9107	
South Atlantic	0.3584		0.5165		0.5199		0.4009	
East South Central	0.5378		0.3095		0.7546		0.1579	
West South Central	0.9138		0.3340		0.5307		0.6295	
West	0.6172		0.9279		0.2099		0.9607	
Mountain	0.9256		0.6376		0.2774		0.9750	
Pacific	0.5178		0.6957		0.0898		0.9700	
COUNTY TYPE								
Large Metro	0.4938		0.4131		0.1142		0.6776	
Small Metro	0.2716		0.6055		0.3906		0.4652	
250K – 1 Mil. Pop.	0.0804		0.9118		0.3747		0.3706	
< 250K Pop.	0.6718		0.2859		0.8304		0.9344	
Nonmetro	0.3384		0.6594		0.2862		0.6194	
Urbanized	0.8398		0.6313		0.8332		0.5287	
Less Urbanized	0.1766		0.3526		0.1912		0.2998	
Completely Rural	0.5149		0.9668		0.1035		0.9065	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.34C Substances for Which Last or Current Treatment Was Received among Persons Who Received Their Last or Current Substance Use Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance for Which Last or Current Treatment Was Received ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	124	121	*	*	29	33	118	113
Marijuana	55	67	*	*	20	19	49	63
Cocaine	55	73	*	*	17	16	51	71
Heroin	56	64	*	*	14	19	53	61
Hallucinogens	41	42	*	*	15	11	36	40
Inhalants	31	25	*	*	*	8	27	24
Pain Relievers	56	60	*	*	17	19	52	55
Tranquilizers	42	54	*	*	14	16	39	51
Stimulants ²	51	49	*	*	14	16	48	46
Sedatives	21	28	*	*	9	3	17	28
Alcohol	94	94	*	*	22	23	89	90

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.34D Substances for Which Last or Current Treatment Was Received among Persons Who Received Their Last or Current Substance Use Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Substance for Which Last or Current Treatment Was Received ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	0.00	0.00	*	*	0.00	0.00	0.00	0.00
Marijuana	2.89	2.97	*	*	4.39	4.46	3.47	3.63
Cocaine	2.89	3.30	*	*	4.31	4.36	3.67	4.06
Heroin	2.84	2.94	*	*	3.55	4.51	3.65	3.65
Hallucinogens	2.27	2.09	*	*	3.95	3.04	2.71	2.57
Inhalants	1.77	1.33	*	*	*	2.33	2.03	1.61
Pain Relievers	2.99	2.90	*	*	3.93	4.61	3.73	3.45
Tranquilizers	2.36	2.73	*	*	3.50	4.29	2.93	3.32
Stimulants ²	2.73	2.41	*	*	3.60	4.17	3.49	2.92
Sedatives	1.20	1.50	*	*	2.65	0.89	1.30	1.92
Alcohol	3.45	3.29	*	*	4.11	4.61	4.39	3.99

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.34P Substances for Which Last or Current Treatment Was Received among Persons Who Received Their Last or Current Substance Use Treatment at a Specialty Facility in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance for Which Last or Current Treatment Was Received ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	N/A		*		N/A		N/A	
Marijuana	0.6022		*		0.0789		0.2374	
Cocaine	0.1842		*		0.3344		0.1053	
Heroin	0.6553		*		0.3482		0.8700	
Hallucinogens	0.5551		*		0.1811		0.9556	
Inhalants	0.3028		*		*		0.5070	
Pain Relievers	0.8821		*		0.1688		0.5157	
Tranquilizers	0.4042		*		0.3659		0.2823	
Stimulants ²	0.6691		*		0.5920		0.8432	
Sedatives	0.6551		*		0.0481		0.2673	
Alcohol	0.9246		*		0.6135		0.8819	

*Low precision; no estimate reported.

N/A: Not Applicable.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.35C Source of Payment for Last or Current Illicit Drug Treatment among Persons Who Received Their Last or Current Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	99	91	*	*	26	29	93	85
Private Health Insurance	54	57	*	*	19	21	50	52
Medicare	48	52	*	*	10	11	46	51
Medicaid	47	58	*	*	9	11	46	56
Public Assistance Other Than Medicaid	59	47	*	*	14	11	56	45
Own Savings or Earnings	56	58	*	*	13	16	54	55
Family Members	46	43	*	*	16	*	41	39
Courts	27	36	*	*	10	9	25	35
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	18	15	*	*	5	4	17	14
Employer	7	13	*	*	3	5	7	12

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.35D Source of Payment for Last or Current Illicit Drug Treatment among Persons Who Received Their Last or Current Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	0.00	0.00	*	*	0.00	0.00	0.00	0.00
Private Health Insurance	4.18	3.85	*	*	5.27	6.00	5.50	4.68
Medicare	3.84	3.86	*	*	3.77	4.36	5.11	4.88
Medicaid	3.74	4.06	*	*	3.50	4.57	5.08	5.22
Public Assistance Other Than Medicaid	4.29	3.34	*	*	4.57	4.53	5.69	4.12
Own Savings or Earnings	4.21	3.69	*	*	4.62	5.25	5.61	4.55
Family Members	3.59	3.27	*	*	5.13	*	4.58	3.88
Courts	2.35	2.80	*	*	3.83	3.54	3.08	3.61
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	1.58	1.22	*	*	2.22	1.82	2.09	1.54
Employer	0.67	1.09	*	*	1.23	2.01	0.86	1.33

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.35P Source of Payment for Last or Current Illicit Drug Treatment among Persons Who Received Their Last or Current Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	N/A		*		N/A		N/A	
Private Health Insurance	0.3217		*		0.6695		0.3000	
Medicare	0.9124		*		0.9298		0.9243	
Medicaid	0.3647		*		0.5591		0.4799	
Public Assistance Other Than Medicaid	0.2692		*		0.3721		0.3462	
Own Savings or Earnings	0.7108		*		0.5500		0.7194	
Family Members	0.4497		*		*		0.2348	
Courts	0.4864		*		0.1887		0.3607	
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	0.8048		*		0.8902		0.8050	
Employer	0.2899		*		0.5881		0.3509	

*Low precision; no estimate reported.

N/A: Not Applicable.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.36C Source of Payment for Last or Current Alcohol Treatment among Persons Who Received Their Last or Current Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	87	94	*	*	20	*	83	90
Private Health Insurance	56	57	*	*	*	*	*	54
Medicare	44	45	*	*	5	*	*	43
Medicaid	53	48	*	*	4	*	*	46
Public Assistance Other Than Medicaid	31	46	*	*	*	*	29	46
Own Savings or Earnings	56	58	*	*	10	*	*	57
Family Members	35	34	*	*	*	*	31	30
Courts	20	33	*	*	*	*	18	32
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	*	28	*	*	*	*	*	28
Employer	9	20	*	*	3	*	8	20

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.36D Source of Payment for Last or Current Alcohol Treatment among Persons Who Received Their Last or Current Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	0.00	0.00	*	*	0.00	*	0.00	0.00
Private Health Insurance	4.95	4.15	*	*	*	*	*	5.05
Medicare	4.42	3.80	*	*	3.28	*	*	4.59
Medicaid	4.93	3.99	*	*	2.66	*	*	4.84
Public Assistance Other Than Medicaid	3.34	3.73	*	*	*	*	4.21	4.54
Own Savings or Earnings	4.81	4.22	*	*	5.79	*	*	5.20
Family Members	3.66	3.06	*	*	*	*	4.36	3.44
Courts	2.29	2.89	*	*	*	*	2.73	3.52
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	*	2.57	*	*	*	*	*	3.14
Employer	1.03	1.88	*	*	2.11	*	1.27	2.31

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.36P Source of Payment for Last or Current Alcohol Treatment among Persons Who Received Their Last or Current Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	N/A		*		N/A		N/A	
Private Health Insurance	0.8707		*		*		*	
Medicare	0.9811		*		*		*	
Medicaid	0.6999		*		*		*	
Public Assistance Other Than Medicaid	0.7234		*		*		0.5967	
Own Savings or Earnings	0.5920		*		*		*	
Family Members	0.8483		*		*		0.8648	
Courts	0.3804		*		*		0.3157	
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	*		*		*		*	
Employer	0.3882		*		*		0.3507	

*Low precision; no estimate reported.

N/A: Not Applicable.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.37C Source of Payment for Last or Current Illicit Drug or Alcohol Treatment among Persons Who Received Their Last or Current Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	124	121	*	*	29	33	118	113
Private Health Insurance	77	85	*	*	21	24	74	81
Medicare	66	76	*	*	12	13	64	74
Medicaid	70	79	*	*	10	14	68	78
Public Assistance Other Than Medicaid	68	67	*	*	15	14	65	64
Own Savings or Earnings	76	82	*	*	17	21	72	78
Family Members	49	55	*	*	18	24	43	49
Courts	33	42	*	*	10	10	31	41
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	42	41	*	*	6	5	42	40
Employer	11	26	*	*	4	5	11	25

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.37D Source of Payment for Last or Current Illicit Drug or Alcohol Treatment among Persons Who Received Their Last or Current Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	0.00	0.00	*	*	0.00	0.00	0.00	0.00
Private Health Insurance	3.52	3.20	*	*	4.42	4.83	4.58	3.93
Medicare	3.33	3.40	*	*	3.19	3.60	4.25	4.21
Medicaid	3.39	3.41	*	*	2.82	3.69	4.40	4.25
Public Assistance Other Than Medicaid	3.29	2.90	*	*	3.73	3.71	4.20	3.52
Own Savings or Earnings	3.64	3.12	*	*	4.19	4.43	4.64	3.87
Family Members	2.64	2.65	*	*	4.33	4.89	3.17	3.08
Courts	1.91	2.08	*	*	2.95	2.75	2.42	2.62
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	2.33	2.09	*	*	1.88	1.60	3.05	2.62
Employer	0.69	1.36	*	*	1.29	1.51	0.85	1.67

*Low precision; no estimate reported.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.37P Source of Payment for Last or Current Illicit Drug or Alcohol Treatment among Persons Who Received Their Last or Current Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Source of Payment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	N/A		*		N/A		N/A	
Private Health Insurance	0.1294		*		0.8319		0.1246	
Medicare	0.4565		*		0.8172		0.4905	
Medicaid	0.3129		*		0.2594		0.4656	
Public Assistance Other Than Medicaid	0.6257		*		0.4006		0.6855	
Own Savings or Earnings	0.5397		*		0.5495		0.5720	
Family Members	0.6561		*		0.1433		0.9411	
Courts	0.5086		*		0.2733		0.4457	
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	0.8291		*		0.8906		0.9333	
Employer	0.0197		*		0.9692		0.0318	

*Low precision; no estimate reported.

N/A: Not Applicable.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.38C Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	122	109	*	*	31	35	114	100
Hospital - Inpatient	65	64	*	*	19	19	61	60
Rehabilitation Facility - Inpatient	83	67	*	*	23	26	77	60
Rehabilitation Facility - Outpatient	104	88	*	*	29	29	97	83
Mental Health Center - Outpatient	70	73	*	*	20	21	66	69
Emergency Room ²	44	43	*	*	10	13	42	41
Private Doctor's Office ²	56	52	*	*	15	13	55	50
Self-Help Group ²	100	86	*	*	26	28	95	80
Prison/Jail ²	29	38	*	*	9	12	27	37

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.38D Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	0.00	0.00	*	*	0.00	0.00	0.00	0.00
Hospital - Inpatient	3.65	3.23	*	*	4.31	4.59	4.92	4.01
Rehabilitation Facility - Inpatient	4.12	3.39	*	*	4.54	5.03	5.48	4.21
Rehabilitation Facility - Outpatient	3.45	3.01	*	*	3.03	4.43	4.73	3.84
Mental Health Center - Outpatient	3.66	3.41	*	*	4.32	4.82	4.95	4.29
Emergency Room ²	2.76	2.46	*	*	2.57	3.65	3.78	3.08
Private Doctor's Office ²	3.27	2.75	*	*	3.80	3.45	4.45	3.56
Self-Help Group ²	3.75	3.37	*	*	4.32	4.51	4.97	4.23
Prison/Jail ²	1.93	2.24	*	*	2.49	3.39	2.58	2.82

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.38P Locations Received Illicit Drug Treatment in the Past Year among Persons Who Received Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	N/A		*		N/A		N/A	
Hospital - Inpatient	0.5531		*		0.5173		0.7224	
Rehabilitation Facility - Inpatient	0.7036		*		0.8789		0.6311	
Rehabilitation Facility - Outpatient	0.1892		*		0.0384		0.4785	
Mental Health Center - Outpatient	0.3488		*		0.5285		0.4377	
Emergency Room ²	0.7866		*		0.2436		0.5029	
Private Doctor's Office ²	0.9369		*		0.5241		0.9647	
Self-Help Group ²	0.4122		*		0.4377		0.1977	
Prison/Jail ²	0.1910		*		0.8566		0.1514	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.39C Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	125	121	*	*	26	28	121	115
Hospital - Inpatient	80	67	*	*	18	16	78	62
Rehabilitation Facility - Inpatient	86	62	*	*	17	17	83	57
Rehabilitation Facility - Outpatient	110	95	*	*	23	24	107	89
Mental Health Center - Outpatient	84	77	*	*	16	18	82	73
Emergency Room ²	58	49	*	*	8	9	57	47
Private Doctor's Office ²	53	48	*	*	*	8	52	47
Self-Help Group ²	109	89	*	*	19	22	107	84
Prison/Jail ²	21	29	*	*	8	*	18	27

*Low precision; no estimate reported.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.39D Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	0.00	0.00	*	*	0.00	0.00	0.00	0.00
Hospital - Inpatient	4.25	3.32	*	*	5.28	5.29	5.34	3.86
Rehabilitation Facility - Inpatient	4.42	3.47	*	*	5.23	5.52	5.47	4.06
Rehabilitation Facility - Outpatient	3.67	3.45	*	*	4.80	4.96	4.52	4.09
Mental Health Center - Outpatient	4.40	3.40	*	*	5.29	5.50	5.47	4.07
Emergency Room ²	3.67	2.72	*	*	3.32	3.33	4.58	3.27
Private Doctor's Office ²	3.39	2.67	*	*	*	3.19	4.21	3.20
Self-Help Group ²	4.09	3.39	*	*	5.34	5.28	4.92	3.95
Prison/Jail ²	1.50	1.75	*	*	3.26	*	1.61	2.02

*Low precision; no estimate reported.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.39P Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	N/A		*		N/A		N/A	
Hospital - Inpatient	0.5213		*		0.7261		0.4629	
Rehabilitation Facility - Inpatient	0.3438		*		0.4439		0.4030	
Rehabilitation Facility - Outpatient	0.1732		*		0.5056		0.2217	
Mental Health Center - Outpatient	0.9931		*		0.2419		0.8330	
Emergency Room ²	0.7530		*		0.8445		0.6365	
Private Doctor's Office ²	0.8834		*		*		0.9359	
Self-Help Group ²	0.4463		*		0.2419		0.2267	
Prison/Jail ²	0.3692		*		*		0.1919	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.40C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	158	149	15	*	36	41	151	139
Hospital - Inpatient	93	85	9	*	22	22	91	79
Rehabilitation Facility - Inpatient	107	85	9	*	24	29	101	76
Rehabilitation Facility - Outpatient	134	118	*	*	31	34	128	110
Mental Health Center - Outpatient	109	94	11	*	22	24	105	89
Emergency Room ²	73	60	*	*	14	16	71	56
Private Doctor's Office ²	70	65	6	*	16	14	68	62
Self-Help Group ²	133	112	10	*	28	33	128	106
Prison/Jail ²	34	44	*	*	10	12	31	42

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.40D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	0.00	0.00	0.00	*	0.00	0.00	0.00	0.00
Hospital - Inpatient	3.02	2.58	5.78	*	3.63	3.73	3.88	3.14
Rehabilitation Facility - Inpatient	3.26	2.74	5.75	*	3.74	4.26	4.10	3.30
Rehabilitation Facility - Outpatient	2.92	2.50	*	*	3.15	3.75	3.69	3.09
Mental Health Center - Outpatient	3.15	2.68	5.87	*	3.55	3.83	4.01	3.32
Emergency Room ²	2.62	2.07	*	*	2.75	3.06	3.36	2.51
Private Doctor's Office ²	2.51	2.14	4.46	*	2.98	2.70	3.24	2.67
Self-Help Group ²	3.08	2.63	5.83	*	3.62	3.86	3.81	3.22
Prison/Jail ²	1.36	1.60	*	*	2.08	2.41	1.67	1.96

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.40P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Location of Treatment ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL POPULATION	N/A		N/A		N/A		N/A	
Hospital - Inpatient	0.9428		*		0.9291		0.8165	
Rehabilitation Facility - Inpatient	0.8184		*		0.8435		0.7507	
Rehabilitation Facility - Outpatient	0.2270		*		0.1591		0.4935	
Mental Health Center - Outpatient	0.4262		*		0.4037		0.3282	
Emergency Room ²	0.2175		*		0.6362		0.1305	
Private Doctor's Office ²	0.9177		*		0.6187		0.9428	
Self-Help Group ²	0.7348		*		0.2972		0.4561	
Prison/Jail ²	0.1420		*		0.3925		0.0800	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

² Emergency Room, Private Doctor's Office, Self-Help Group, and Prison/Jail are not considered specialty facilities.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.41C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Numbers in Thousands, 2013 and 2014

Dependence or Abuse Status	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol (2013)	Both Illicit Drugs and Alcohol (2014)	Illicit Drugs or Alcohol ¹ (2013)	Illicit Drugs or Alcohol ¹ (2014)
TOTAL	123	108	125	121	75	73	160	148
DEPENDENCE								
None	95	76	93	82	55	48	120	102
Both Illicit Drugs and Alcohol	42	36	38	38	37	36	44	44
Illicit Drugs or Alcohol	72	75	82	81	49	50	100	102
Illicit Drugs	69	66	50	46	48	42	73	74
Alcohol	47	51	75	77	39	46	80	83
DEPENDENCE OR ABUSE								
None	94	71	90	74	55	44	118	92
Both Illicit Drugs and Alcohol	48	42	46	48	41	40	53	57
Illicit Drugs or Alcohol	74	78	86	89	50	54	105	111
Illicit Drugs	72	68	54	52	49	44	79	78
Alcohol	51	57	81	87	42	50	87	96

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.41D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Percentages, 2013 and 2014

Dependence or Abuse Status	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol (2013)	Both Illicit Drugs and Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.05	0.04	0.05	0.05	0.03	0.03	0.06	0.06
DEPENDENCE								
None	0.04	0.03	0.04	0.03	0.02	0.02	0.05	0.04
Both Illicit Drugs and Alcohol	3.46	3.16	3.21	3.25	3.11	3.14	3.55	3.38
Illicit Drugs or Alcohol	0.60	0.61	0.68	0.65	0.41	0.42	0.82	0.80
Illicit Drugs	1.34	1.27	1.00	0.89	0.95	0.82	1.42	1.34
Alcohol	0.58	0.62	0.91	0.91	0.48	0.56	0.96	0.98
DEPENDENCE OR ABUSE								
None	0.04	0.03	0.04	0.03	0.02	0.02	0.05	0.04
Both Illicit Drugs and Alcohol	1.74	1.57	1.67	1.72	1.51	1.47	1.88	1.89
Illicit Drugs or Alcohol	0.34	0.36	0.39	0.41	0.23	0.25	0.48	0.49
Illicit Drugs	1.00	0.94	0.77	0.71	0.70	0.61	1.08	1.02
Alcohol	0.29	0.33	0.46	0.50	0.24	0.29	0.49	0.54

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.41P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse Status: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Dependence or Abuse Status	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol (2013)	Both Illicit Drugs and Alcohol (2014)	Illicit Drugs or Alcohol¹ (2013)	Illicit Drugs or Alcohol¹ (2014)
TOTAL	0.5221		0.4371		0.2112		0.6008	
DEPENDENCE								
None	0.9581		0.8673		0.4610		0.8005	
Both Illicit Drugs and Alcohol	0.3305		0.1027		0.0747		0.1497	
Illicit Drugs or Alcohol	0.3418		0.3143		0.2776		0.2673	
Illicit Drugs	0.4875		0.6643		0.5610		0.3603	
Alcohol	0.2462		0.1147		0.0381		0.2165	
DEPENDENCE OR ABUSE								
None	0.7480		0.5669		0.7326		0.3483	
Both Illicit Drugs and Alcohol	0.7079		0.1898		0.2215		0.2436	
Illicit Drugs or Alcohol	0.1449		0.0783		0.1259		0.0630	
Illicit Drugs	0.6094		0.6915		0.6334		0.4571	
Alcohol	0.1556		0.0227		0.0300		0.0386	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.42C Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse of Specific Substances: Standard Errors of Numbers in Thousands, 2013 and 2014

Past Year Dependence or Abuse ¹	Illicit Drugs ² (2013)	Illicit Drugs ² (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ² (2013)	Both Illicit Drugs and Alcohol ² (2014)	Illicit Drugs or Alcohol ^{2,3} (2013)	Illicit Drugs or Alcohol ^{2,3} (2014)
ILLCIT DRUGS²	72	68	54	52	49	44	79	78
Marijuana and Hashish	41	41	33	38	29	27	47	55
Cocaine	35	46	28	32	25	32	37	51
Heroin	*	42	19	22	19	22	*	42
Hallucinogens	*	7	*	7	*	7	*	7
Inhalants	*	*	*	*	*	*	*	*
Nonmedical Use of Psychotherapeutics ^{4,5}	53	47	38	32	36	29	55	52
Pain Relievers	50	48	36	30	35	28	51	52
Tranquilizers	*	30	*	12	*	9	*	34
Stimulants ⁴	20	21	15	11	13	11	21	24
Sedatives	*	*	*	*	*	*	*	*
ALCOHOL	51	57	81	87	42	50	87	96
BOTH ILLICIT DRUGS AND ALCOHOL²	48	42	46	48	41	40	53	57
ILLCIT DRUGS OR ALCOHOL²	74	78	86	89	50	54	105	111

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

³ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

⁴ Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

⁵ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.42D Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse of Specific Substances: Standard Errors of Percentages, 2013 and 2014

Past Year Dependence or Abuse ¹	Illicit Drugs ² (2013)	Illicit Drugs ² (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ² (2013)	Both Illicit Drugs and Alcohol ² (2014)	Illicit Drugs or Alcohol ^{2,3} (2013)	Illicit Drugs or Alcohol ^{2,3} (2014)
ILLCIT DRUGS²	1.00	0.94	0.77	0.71	0.70	0.61	1.08	1.02
Marijuana and Hashish	0.94	0.96	0.77	0.87	0.68	0.65	1.07	1.23
Cocaine	3.96	4.32	3.02	3.31	2.75	3.29	4.13	4.62
Heroin	*	5.14	3.36	3.35	3.31	3.34	*	5.14
Hallucinogens	*	2.73	*	2.65	*	2.64	*	2.75
Inhalants	*	*	*	*	*	*	*	*
Nonmedical Use of Psychotherapeutics ^{4,5}	2.12	1.87	1.58	1.25	1.51	1.17	2.19	2.03
Pain Relievers	2.47	2.38	1.83	1.49	1.81	1.40	2.49	2.52
Tranquilizers	*	5.28	*	2.48	*	1.89	*	5.59
Stimulants ⁴	4.40	4.06	3.21	2.22	2.87	2.22	4.65	4.51
Sedatives	*	*	*	*	*	*	*	*
ALCOHOL	0.29	0.33	0.46	0.50	0.24	0.29	0.49	0.54
BOTH ILLICIT DRUGS AND ALCOHOL²	1.74	1.57	1.67	1.72	1.51	1.47	1.88	1.89
ILLCIT DRUGS OR ALCOHOL²	0.34	0.36	0.39	0.41	0.23	0.25	0.48	0.49

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

³ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

⁴ Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

⁵ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.42P Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Past Year Dependence or Abuse of Specific Substances: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Past Year Dependence or Abuse ¹	Illicit Drugs ² (2013)	Illicit Drugs ² (2014)	Alcohol (2013)	Alcohol (2014)	Both Illicit Drugs and Alcohol ² (2013)	Both Illicit Drugs and Alcohol ² (2014)	Illicit Drugs or Alcohol ^{2,3} (2013)	Illicit Drugs or Alcohol ^{2,3} (2014)
ILLCIT DRUGS²	0.6094		0.6915		0.6334		0.4571	
Marijuana and Hashish	0.3820		0.5423		0.9534		0.9878	
Cocaine	0.1242		0.3000		0.1067		0.1285	
Heroin	*		0.6149		0.5428		*	
Hallucinogens	*		*		*		*	
Inhalants	*		*		*		*	
Nonmedical Use of Psychotherapeutics ^{4,5}	0.9899		0.9489		0.9815		0.6673	
Pain Relievers	0.9475		0.8762		0.9508		0.7153	
Tranquilizers	*		*		*		*	
Stimulants ⁴	0.9648		0.5382		0.9050		0.7022	
Sedatives	*		*		*		*	
ALCOHOL	0.1556		0.0227		0.0300		0.0386	
BOTH ILLICIT DRUGS AND ALCOHOL²	0.7079		0.1898		0.2215		0.2436	
ILLCIT DRUGS OR ALCOHOL²	0.1449		0.0783		0.1259		0.0630	

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

³ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

⁴ Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

⁵ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 5.43C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year, by Detailed Age Category:
Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014**

Age Category	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	261	237	123	108	226	214	1.41	1.23
12	5	10	*	1	5	10	*	*
13	10	10	3	*	9	10	*	*
14	16	12	2	5	16	11	*	*
15	19	23	5	5	19	22	2.48	2.69
16	21	26	8	7	20	25	2.74	2.41
17	21	24	7	7	20	24	2.36	2.65
18	35	35	11	6	33	34	2.64	1.83
19	34	31	10	8	32	30	2.72	2.51
20	32	33	8	11	30	31	2.20	3.30
21	35	30	11	11	33	28	2.82	3.24
22	31	35	10	10	29	34	2.67	2.68
23	31	32	13	15	28	29	3.68	3.87
24	31	28	9	12	30	25	2.69	4.58
25	26	32	12	16	23	27	4.31	5.04
26-29	94	77	47	40	79	66	4.84	3.61
30-34	105	74	45	34	97	65	4.28	3.65
35-39	76	67	45	33	65	56	*	4.90
40-44	62	61	32	38	53	49	*	*
45-49	53	49	25	23	46	43	*	*
50-54	130	88	31	36	126	80	*	*
55-59	58	71	40	48	41	52	*	*
60-64	52	51	43	13	29	49	*	*
65 or Older	50	63	*	21	50	60	*	*

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 5.43D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year, by Detailed Age Category:
Standard Errors of Percentages, 2013 and 2014**

Age Category	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.10	0.09	0.05	0.04	0.09	0.08	1.41	1.23
12	0.14	0.26	*	0.04	0.14	0.25	*	*
13	0.23	0.25	0.08	*	0.21	0.25	*	*
14	0.36	0.29	0.05	0.11	0.36	0.26	*	*
15	0.46	0.51	0.12	0.12	0.44	0.50	2.48	2.69
16	0.48	0.57	0.18	0.16	0.45	0.55	2.74	2.41
17	0.50	0.56	0.17	0.18	0.47	0.55	2.36	2.65
18	0.70	0.72	0.22	0.13	0.67	0.71	2.64	1.83
19	0.78	0.73	0.23	0.18	0.75	0.71	2.72	2.51
20	0.71	0.73	0.19	0.24	0.67	0.70	2.20	3.30
21	0.77	0.65	0.27	0.26	0.73	0.61	2.82	3.24
22	0.68	0.75	0.22	0.23	0.63	0.73	2.67	2.68
23	0.67	0.69	0.29	0.33	0.62	0.62	3.68	3.87
24	0.69	0.63	0.19	0.26	0.67	0.58	2.69	4.58
25	0.62	0.71	0.29	0.37	0.56	0.62	4.31	5.04
26-29	0.53	0.44	0.27	0.23	0.45	0.38	4.84	3.61
30-34	0.50	0.35	0.22	0.16	0.46	0.30	4.28	3.65
35-39	0.40	0.34	0.24	0.17	0.34	0.28	*	4.90
40-44	0.28	0.30	0.15	0.19	0.24	0.24	*	*
45-49	0.26	0.23	0.13	0.11	0.23	0.21	*	*
50-54	0.56	0.36	0.14	0.15	0.54	0.33	*	*
55-59	0.28	0.34	0.19	0.23	0.20	0.25	*	*
60-64	0.30	0.29	0.25	0.08	0.17	0.28	*	*
65 or Older	0.12	0.14	*	0.05	0.12	0.13	*	*

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.43P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.5846		0.5221		0.7775		0.6462	
12	0.2233		*		0.2741		*	
13	0.7380		*		0.9331		*	
14	0.1525		0.2618		0.0702		*	
15	0.7806		0.9759		0.7803		0.9480	
16	0.8098		0.2640		0.5242		0.2088	
17	0.9593		0.5098		0.8651		0.5028	
18	0.2763		0.0316		0.5797		0.0530	
19	0.2575		0.1743		0.4333		0.3222	
20	0.2142		0.8567		0.1723		0.5580	
21	0.1264		0.5916		0.0660		0.2594	
22	0.3238		0.8597		0.2721		0.5823	
23	0.5777		0.5456		0.7564		0.6703	
24	0.0462		0.9645		0.0331		0.3719	
25	0.7427		0.7350		0.8567		0.8329	
26-29	0.2447		0.8650		0.1432		0.3765	
30-34	0.1451		0.7912		0.1414		0.5363	
35-39	0.6667		0.6033		0.8708		*	
40-44	0.1926		0.2311		0.4632		*	
45-49	0.2950		0.7553		0.3100		*	
50-54	0.5607		0.2928		0.7843		*	
55-59	0.1192		0.3331		0.2183		*	
60-64	0.8925		0.0766		0.1157		*	
65 or Older	0.7059		*		0.9084		*	

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.44C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	261	237	123	108	226	214	1.41	1.23
AGE								
12-17	41	46	12	12	39	45	1.27	1.34
18-25	90	93	31	34	83	88	1.08	1.31
26 or Older	238	207	115	100	209	180	2.52	1.93
GENDER								
Male	185	184	87	88	163	164	1.62	1.61
Female	181	132	83	56	163	121	2.58	1.74
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	240	215	107	97	212	193	1.48	1.33
White	217	173	101	83	190	151	1.85	1.56
Black or African American	96	109	37	45	86	103	2.91	3.01
American Indian or Alaska Native	21	12	12	8	17	9	*	*
Native Hawaiian or Other Pacific Islander	10	22	4	*	9	21	*	*
Asian	43	35	18	4	39	34	*	*
Two or More Races	35	32	3	6	35	31	1.81	3.05
Hispanic or Latino	97	102	57	51	80	91	4.13	3.46

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.44D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.10	0.09	0.05	0.04	0.09	0.08	1.41	1.23
AGE								
12-17	0.17	0.19	0.05	0.05	0.16	0.18	1.27	1.34
18-25	0.26	0.27	0.09	0.10	0.24	0.25	1.08	1.31
26 or Older	0.12	0.10	0.06	0.05	0.10	0.09	2.52	1.93
GENDER								
Male	0.15	0.14	0.07	0.07	0.13	0.13	1.62	1.61
Female	0.13	0.10	0.06	0.04	0.12	0.09	2.58	1.74
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.11	0.10	0.05	0.04	0.10	0.09	1.48	1.33
White	0.13	0.10	0.06	0.05	0.11	0.09	1.85	1.56
Black or African American	0.31	0.35	0.12	0.14	0.28	0.32	2.91	3.01
American Indian or Alaska Native	1.30	0.76	0.85	0.53	1.05	0.60	*	*
Native Hawaiian or Other Pacific Islander	1.14	2.08	0.48	*	1.01	1.98	*	*
Asian	0.32	0.25	0.14	0.03	0.29	0.25	*	*
Two or More Races	0.83	0.74	0.08	0.14	0.83	0.72	1.81	3.05
Hispanic or Latino	0.24	0.24	0.14	0.12	0.19	0.21	4.13	3.46

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.44P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.5846		0.5221		0.7775		0.6462	
AGE								
12-17	0.8400		0.3555		0.9524		0.3671	
18-25	0.0861		0.7569		0.0857		0.7361	
26 or Older	0.1608		0.3745		0.2663		0.8878	
GENDER								
Male	0.5965		0.1618		0.8797		0.1816	
Female	0.8059		0.4966		0.5443		0.3821	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.7408		0.5671		0.9344		0.6341	
White	0.7296		0.9299		0.6582		0.7779	
Black or African American	0.0992		0.1647		0.2362		0.4671	
American Indian or Alaska Native	0.0681		0.9617		0.0253		*	
Native Hawaiian or Other Pacific Islander	0.3469		*		0.3303		*	
Asian	0.7050		0.3046		0.9764		*	
Two or More Races	0.9902		0.1027		0.8017		0.1219	
Hispanic or Latino	0.5431		0.7723		0.5985		0.9310	

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.45C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	257	233	122	107	223	207	1.59	1.36
GENDER								
Male	181	181	86	88	159	161	1.80	1.77
Female	180	129	82	55	162	117	3.00	1.98
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	236	210	107	96	209	188	1.65	1.46
White	215	173	99	83	191	149	2.05	1.71
Black or African American	96	115	37	45	87	106	3.33	3.27
American Indian or Alaska Native	20	12	12	8	16	9	*	*
Native Hawaiian or Other Pacific Islander	9	20	*	*	9	19	*	*
Asian	43	34	18	4	39	34	*	*
Two or More Races	35	32	3	6	35	31	*	3.31
Hispanic or Latino	94	99	57	50	77	87	*	4.14
EDUCATION								
< High School	109	96	64	50	87	84	3.62	3.11
High School Graduate	129	130	62	67	114	111	2.47	2.66
Some College	172	134	67	55	157	118	2.84	2.12
College Graduate	88	96	37	37	81	92	3.91	3.08
CURRENT EMPLOYMENT								
Full-Time	178	157	42	59	172	144	1.55	1.72
Part-Time	98	89	44	39	88	80	3.13	2.80
Unemployed	97	67	68	31	65	57	4.56	3.29
Other ¹	132	124	73	75	108	104	3.52	3.50

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.45D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.11	0.10	0.05	0.04	0.09	0.09	1.59	1.36
GENDER								
Male	0.16	0.16	0.08	0.08	0.14	0.14	1.80	1.77
Female	0.15	0.10	0.07	0.04	0.13	0.09	3.00	1.98
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.12	0.10	0.05	0.05	0.10	0.09	1.65	1.46
White	0.14	0.11	0.06	0.05	0.12	0.09	2.05	1.71
Black or African American	0.34	0.39	0.13	0.16	0.31	0.36	3.33	3.27
American Indian or Alaska Native	1.42	0.85	0.95	0.59	1.13	0.67	*	*
Native Hawaiian or Other Pacific Islander	1.24	2.20	*	*	1.13	2.07	*	*
Asian	0.35	0.27	0.15	0.03	0.32	0.27	*	*
Two or More Races	1.01	0.88	0.09	0.16	1.00	0.87	*	3.31
Hispanic or Latino	0.26	0.27	0.16	0.14	0.21	0.23	*	4.14
EDUCATION								
< High School	0.33	0.29	0.20	0.16	0.27	0.25	3.62	3.11
High School Graduate	0.18	0.18	0.09	0.10	0.16	0.16	2.47	2.66
Some College	0.26	0.20	0.11	0.08	0.24	0.18	2.84	2.12
College Graduate	0.12	0.13	0.05	0.05	0.11	0.12	3.91	3.08
CURRENT EMPLOYMENT								
Full-Time	0.15	0.12	0.04	0.05	0.14	0.11	1.55	1.72
Part-Time	0.29	0.27	0.13	0.12	0.26	0.24	3.13	2.80
Unemployed	0.80	0.60	0.57	0.29	0.54	0.52	4.56	3.29
Other ¹	0.18	0.17	0.10	0.10	0.15	0.14	3.52	3.50

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.45P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.5497		0.4600		0.7743		0.5953	
GENDER								
Male	0.5008		0.1238		0.9402		0.1559	
Female	0.8750		0.4859		0.6030		0.3929	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.6214		0.5182		0.8220		0.6420	
White	0.7728		0.8958		0.6879		0.7570	
Black or African American	0.0707		0.1566		0.1788		0.5464	
American Indian or Alaska Native	0.1012		0.9552		0.0344		*	
Native Hawaiian or Other Pacific Islander	0.2907		*		0.3219		*	
Asian	0.7476		0.3037		0.9719		*	
Two or More Races	0.9825		0.0806		0.8256		*	
Hispanic or Latino	0.7055		0.7180		0.8335		*	
EDUCATION								
< High School	0.8245		0.8935		0.7204		0.7780	
High School Graduate	0.8675		0.4332		0.5142		0.3232	
Some College	0.8275		0.8102		0.8936		0.8804	
College Graduate	0.0669		0.6525		0.0791		0.8696	
CURRENT EMPLOYMENT								
Full-Time	0.1058		0.0087		0.3909		0.0440	
Part-Time	0.9633		0.6208		0.7662		0.5663	
Unemployed	0.1333		0.0388		0.8119		0.0543	
Other ¹	0.6766		0.1876		0.6738		0.1844	

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.46C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	261	237	123	108	226	214	1.41	1.23
GEOGRAPHIC DIVISION								
Northeast	101	119	56	52	83	109	3.21	2.87
New England	58	49	27	24	52	40	*	4.46
Middle Atlantic	83	109	49	46	64	102	3.95	3.59
Midwest	95	89	43	48	82	75	2.51	2.74
East North Central	79	75	33	37	70	67	2.79	3.02
West North Central	51	42	27	31	43	33	*	*
South	181	141	70	69	166	126	2.40	2.28
South Atlantic	157	105	53	58	145	93	3.33	3.33
East South Central	69	40	37	21	59	35	*	4.91
West South Central	60	81	27	28	54	74	3.59	3.41
West	129	128	69	51	108	112	3.04	2.22
Mountain	67	58	39	23	55	50	*	3.29
Pacific	108	115	58	45	92	102	3.81	2.86
COUNTY TYPE								
Large Metro	185	207	93	91	160	184	1.94	1.77
Small Metro	172	126	48	52	164	115	1.97	2.05
250K – 1 Mil. Pop.	111	104	33	45	105	94	2.02	2.62
< 250K Pop.	132	75	36	24	126	68	4.26	2.73
Nonmetro	89	66	61	31	61	54	4.45	2.71
Urbanized	56	45	26	21	48	39	4.35	4.12
Less Urbanized	68	46	55	22	34	37	*	4.11
Completely Rural	22	16	8	7	20	14	*	*

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.46D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.10	0.09	0.05	0.04	0.09	0.08	1.41	1.23
GEOGRAPHIC DIVISION								
Northeast	0.21	0.25	0.12	0.11	0.17	0.23	3.21	2.87
New England	0.46	0.39	0.22	0.19	0.42	0.32	*	4.46
Middle Atlantic	0.24	0.31	0.14	0.13	0.18	0.29	3.95	3.59
Midwest	0.17	0.16	0.08	0.09	0.15	0.13	2.51	2.74
East North Central	0.20	0.19	0.09	0.10	0.18	0.17	2.79	3.02
West North Central	0.29	0.24	0.16	0.18	0.25	0.19	*	*
South	0.19	0.14	0.07	0.07	0.17	0.13	2.40	2.28
South Atlantic	0.30	0.20	0.10	0.11	0.28	0.18	3.33	3.33
East South Central	0.44	0.25	0.24	0.14	0.38	0.22	*	4.91
West South Central	0.20	0.26	0.09	0.09	0.18	0.24	3.59	3.41
West	0.21	0.21	0.11	0.08	0.18	0.18	3.04	2.22
Mountain	0.36	0.30	0.21	0.12	0.29	0.26	*	3.29
Pacific	0.25	0.27	0.14	0.10	0.22	0.24	3.81	2.86
COUNTY TYPE								
Large Metro	0.13	0.14	0.07	0.06	0.11	0.12	1.94	1.77
Small Metro	0.21	0.15	0.06	0.07	0.20	0.14	1.97	2.05
250K – 1 Mil. Pop.	0.21	0.18	0.06	0.08	0.19	0.17	2.02	2.62
< 250K Pop.	0.49	0.29	0.14	0.10	0.47	0.26	4.26	2.73
Nonmetro	0.21	0.15	0.14	0.07	0.14	0.13	4.45	2.71
Urbanized	0.29	0.24	0.14	0.12	0.25	0.21	4.35	4.12
Less Urbanized	0.35	0.21	0.29	0.10	0.17	0.17	*	4.11
Completely Rural	0.46	0.35	0.17	0.16	0.42	0.30	*	*

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.46P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug Problem in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Geographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.5846		0.5221		0.7775		0.6462	
GEOGRAPHIC DIVISION								
Northeast	0.1557		0.4909		0.2152		1.0000	
New England	0.8995		0.9933		0.8805		*	
Middle Atlantic	0.0909		0.4356		0.1365		0.9370	
Midwest	0.9568		0.4769		0.7273		0.4232	
East North Central	0.6224		0.3000		0.9833		0.3556	
West North Central	0.4784		0.8843		0.4496		*	
South	0.8758		0.6088		0.6800		0.5203	
South Atlantic	0.9270		0.3584		0.7499		0.3378	
East South Central	0.1043		0.5378		0.1318		*	
West South Central	0.4446		0.9138		0.4280		0.8257	
West	1.0000		0.6172		0.7835		0.5672	
Mountain	0.5002		0.9256		0.3900		*	
Pacific	0.7066		0.5178		0.4374		0.3768	
COUNTY TYPE								
Large Metro	0.3328		0.4938		0.4660		0.7522	
Small Metro	0.9601		0.2716		0.6440		0.2347	
250K – 1 Mil. Pop.	0.8167		0.0804		0.6460		0.0783	
< 250K Pop.	0.7579		0.6718		0.8481		0.7887	
Nonmetro	0.5099		0.3384		0.9455		0.3630	
Urbanized	0.8768		0.8398		0.7719		0.7592	
Less Urbanized	0.5075		0.1766		0.5592		*	
Completely Rural	0.9057		0.5149		0.6663		*	

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.47C Perceived Need for Illicit Drug Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug Problem, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment¹ (2013)	Felt Need for Treatment¹ (2014)	Felt Need and Made Effort to Get Treatment¹ (2013)	Felt Need and Made Effort to Get Treatment¹ (2014)	Felt Need and Made No Effort to Get Treatment¹ (2013)	Felt Need and Made No Effort to Get Treatment¹ (2014)	Did Not Feel Need for Treatment¹ (2013)	Did Not Feel Need for Treatment¹ (2014)
TOTAL	231	221	52	48	35	31	38	36	227	215
AGE										
12-17	40	46	7	9	3	8	5	4	40	44
18-25	87	91	15	16	8	10	12	13	86	90
26 or Older	213	184	49	44	33	29	35	33	209	182
GENDER										
Male	164	169	45	32	31	22	32	23	160	163
Female	166	122	25	34	15	22	20	27	166	117

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but have not received treatment for an illicit drug problem at a specialty facility.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.47D Perceived Need for Illicit Drug Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug Problem, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment ¹ (2013)	Felt Need for Treatment ¹ (2014)	Felt Need and Made Effort to Get Treatment ¹ (2013)	Felt Need and Made Effort to Get Treatment ¹ (2014)	Felt Need and Made No Effort to Get Treatment ¹ (2013)	Felt Need and Made No Effort to Get Treatment ¹ (2014)	Did Not Feel Need for Treatment ¹ (2013)	Did Not Feel Need for Treatment ¹ (2014)
TOTAL	0.00	0.00	0.84	0.75	0.56	0.48	0.62	0.57	0.84	0.75
AGE										
12-17	0.00	0.00	0.81	1.03	0.37	0.95	0.58	0.44	0.81	1.03
18-25	0.00	0.00	0.63	0.74	0.36	0.45	0.52	0.59	0.63	0.74
26 or Older	0.00	0.00	1.65	1.32	1.13	0.84	1.19	1.01	1.65	1.32
GENDER										
Male	0.00	0.00	1.12	0.80	0.79	0.56	0.81	0.60	1.12	0.80
Female	0.00	0.00	1.19	1.37	0.70	0.90	0.92	1.08	1.19	1.37

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but have not received treatment for an illicit drug problem at a specialty facility.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.47P Perceived Need for Illicit Drug Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug Problem, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment ¹ (2013)	Felt Need for Treatment ¹ (2014)	Felt Need and Made Effort to Get Treatment ¹ (2013)	Felt Need and Made Effort to Get Treatment ¹ (2014)	Felt Need and Made No Effort to Get Treatment ¹ (2013)	Felt Need and Made No Effort to Get Treatment ¹ (2014)	Did Not Feel Need for Treatment ¹ (2013)	Did Not Feel Need for Treatment ¹ (2014)
TOTAL	N/A		0.9318		0.8767		0.9819		0.9318	
AGE										
12-17	N/A		0.6313		0.1285		0.2078		0.6313	
18-25	N/A		0.8079		0.8327		0.6492		0.8079	
26 or Older	N/A		0.6487		0.6170		0.8684		0.6487	
GENDER										
Male	N/A		0.2517		0.3869		0.4609		0.2517	
Female	N/A		0.1971		0.3383		0.3767		0.1971	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but have not received treatment for an illicit drug problem at a specialty facility.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.48C Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	429	375	125	121	406	351	0.65	0.64
AGE								
12-17	37	42	13	12	35	40	1.61	1.57
18-25	123	120	26	27	120	117	0.55	0.61
26 or Older	405	337	121	115	385	318	0.91	0.86
GENDER								
Male	351	298	100	108	336	280	0.82	0.90
Female	251	195	77	51	238	190	1.14	0.80
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	375	331	113	106	357	311	0.72	0.68
White	351	293	99	90	336	279	0.79	0.75
Black or African American	125	120	44	50	117	112	2.29	2.34
American Indian or Alaska Native	47	47	13	9	45	43	*	3.44
Native Hawaiian or Other Pacific Islander	*	20	4	*	*	20	*	*
Asian	75	58	23	7	71	58	*	1.34
Two or More Races	59	48	14	14	57	46	3.49	*
Hispanic or Latino	194	164	56	52	182	154	1.69	1.60

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.48D Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.16	0.14	0.05	0.05	0.15	0.13	0.65	0.64
AGE								
12-17	0.15	0.17	0.05	0.05	0.14	0.16	1.61	1.57
18-25	0.35	0.34	0.07	0.08	0.35	0.33	0.55	0.61
26 or Older	0.20	0.16	0.06	0.06	0.19	0.15	0.91	0.86
GENDER								
Male	0.28	0.23	0.08	0.08	0.26	0.22	0.82	0.90
Female	0.19	0.14	0.06	0.04	0.18	0.14	1.14	0.80
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.17	0.15	0.05	0.05	0.16	0.14	0.72	0.68
White	0.21	0.17	0.06	0.05	0.20	0.16	0.79	0.75
Black or African American	0.40	0.38	0.14	0.16	0.38	0.35	2.29	2.34
American Indian or Alaska Native	3.01	2.54	0.91	0.56	2.91	2.36	*	3.44
Native Hawaiian or Other Pacific Islander	*	2.01	0.48	*	*	1.98	*	*
Asian	0.56	0.43	0.17	0.05	0.53	0.42	*	1.34
Two or More Races	1.34	1.03	0.32	0.31	1.29	1.00	3.49	*
Hispanic or Latino	0.47	0.39	0.14	0.12	0.44	0.36	1.69	1.60

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.48P Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.3021		0.4371		0.1783		0.2603	
AGE								
12-17	0.5313		0.2913		0.7468		0.3562	
18-25	0.0919		0.7881		0.0732		0.4870	
26 or Older	0.6248		0.3941		0.4236		0.2918	
GENDER								
Male	0.4952		0.3254		0.2946		0.2172	
Female	0.3885		0.9157		0.3859		0.9055	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.4001		0.4370		0.2533		0.2856	
White	0.2647		0.3791		0.1506		0.2140	
Black or African American	0.3309		0.4632		0.4571		0.6484	
American Indian or Alaska Native	0.9723		0.6644		0.8724		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.9446		0.2480		0.8142		*	
Two or More Races	0.5887		0.6855		0.6537		*	
Hispanic or Latino	0.4971		0.8459		0.4312		0.6990	

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.49C Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	428	371	124	120	404	348	0.68	0.66
GENDER								
Male	350	296	99	108	336	278	0.84	0.92
Female	249	194	77	51	237	189	1.21	0.84
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	373	326	112	105	355	307	0.74	0.70
White	374	297	98	89	359	283	0.82	0.77
Black or African American	132	130	44	50	122	118	2.35	2.39
American Indian or Alaska Native	47	46	13	9	45	42	*	3.50
Native Hawaiian or Other Pacific Islander	*	20	*	*	*	20	*	*
Asian	75	58	23	7	71	57	*	1.37
Two or More Races	59	48	13	14	57	46	3.66	*
Hispanic or Latino	193	163	56	52	180	152	1.77	1.69
EDUCATION								
< High School	157	144	47	59	145	128	1.88	2.28
High School Graduate	234	188	64	73	224	172	1.32	1.52
Some College	238	197	76	53	219	186	1.39	0.96
College Graduate	293	226	59	36	287	221	1.09	0.69
CURRENT EMPLOYMENT								
Full-Time	356	278	59	62	351	270	0.58	0.61
Part-Time	156	153	31	35	153	149	1.20	1.30
Unemployed	109	86	59	36	89	77	3.71	2.70
Other ¹	219	180	86	82	203	156	2.36	2.20

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.49D Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.18	0.15	0.05	0.05	0.17	0.14	0.68	0.66
GENDER								
Male	0.31	0.26	0.09	0.09	0.29	0.24	0.84	0.92
Female	0.20	0.16	0.06	0.04	0.19	0.15	1.21	0.84
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.18	0.16	0.06	0.05	0.18	0.15	0.74	0.70
White	0.22	0.18	0.06	0.06	0.21	0.18	0.82	0.77
Black or African American	0.45	0.42	0.16	0.17	0.42	0.39	2.35	2.39
American Indian or Alaska Native	3.34	2.77	1.01	0.63	3.23	2.58	*	3.50
Native Hawaiian or Other Pacific Islander	*	2.23	*	*	*	2.20	*	*
Asian	0.62	0.47	0.19	0.05	0.58	0.46	*	1.37
Two or More Races	1.65	1.24	0.38	0.38	1.58	1.21	3.66	*
Hispanic or Latino	0.54	0.44	0.16	0.14	0.51	0.41	1.77	1.69
EDUCATION								
< High School	0.47	0.43	0.15	0.18	0.44	0.38	1.88	2.28
High School Graduate	0.33	0.26	0.09	0.11	0.31	0.24	1.32	1.52
Some College	0.36	0.29	0.12	0.08	0.33	0.27	1.39	0.96
College Graduate	0.38	0.29	0.08	0.05	0.37	0.29	1.09	0.69
CURRENT EMPLOYMENT								
Full-Time	0.28	0.21	0.05	0.05	0.28	0.20	0.58	0.61
Part-Time	0.45	0.45	0.09	0.11	0.44	0.44	1.20	1.30
Unemployed	0.90	0.74	0.50	0.33	0.74	0.67	3.71	2.70
Other ¹	0.30	0.24	0.12	0.11	0.27	0.21	2.36	2.20

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.49P Need for and Receipt of Treatment at a Specialty Facility for an Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.3182		0.3781		0.1789		0.2203	
GENDER								
Male	0.4982		0.2455		0.2709		0.1556	
Female	0.4206		0.8467		0.4349		0.9915	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.4368		0.4315		0.2809		0.2904	
White	0.2907		0.4062		0.1732		0.2411	
Black or African American	0.3481		0.4407		0.4889		0.6124	
American Indian or Alaska Native	0.8977		0.8290		0.8437		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.8750		0.2466		0.8837		*	
Two or More Races	0.6910		0.7008		0.7583		*	
Hispanic or Latino	0.4592		0.6822		0.3561		0.5270	
EDUCATION								
< High School	0.9079		0.0851		0.4120		0.0524	
High School Graduate	0.6737		0.1435		0.3339		0.0878	
Some College	0.7470		0.6387		0.8499		0.6799	
College Graduate	0.3205		0.3006		0.4217		0.3907	
CURRENT EMPLOYMENT								
Full-Time	0.2706		0.1543		0.1587		0.0897	
Part-Time	0.6069		0.9012		0.6188		0.9653	
Unemployed	0.9478		0.2037		0.4955		0.1507	
Other ¹	0.5312		0.2666		0.2245		0.1259	

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.50C Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment¹ (2013)	Felt Need for Treatment¹ (2014)	Felt Need and Made Effort to Get Treatment¹ (2013)	Felt Need and Made Effort to Get Treatment¹ (2014)	Felt Need and Made No Effort to Get Treatment¹ (2013)	Felt Need and Made No Effort to Get Treatment¹ (2014)	Did Not Feel Need for Treatment¹ (2013)	Did Not Feel Need for Treatment¹ (2014)
TOTAL	436	364	81	62	46	28	67	54	429	362
AGE										
12-17	35	40	4	5	3	*	2	1	35	40
18-25	142	133	15	15	12	11	9	11	141	133
26 or Older	408	328	80	60	44	25	66	53	400	325
GENDER										
Male	354	295	69	40	42	20	55	34	350	294
Female	248	192	42	48	18	19	38	42	243	189

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.50D Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment¹ (2013)	Felt Need for Treatment¹ (2014)	Felt Need and Made Effort to Get Treatment¹ (2013)	Felt Need and Made Effort to Get Treatment¹ (2014)	Felt Need and Made No Effort to Get Treatment¹ (2013)	Felt Need and Made No Effort to Get Treatment¹ (2014)	Did Not Feel Need for Treatment¹ (2013)	Did Not Feel Need for Treatment¹ (2014)
TOTAL	0.00	0.00	0.48	0.39	0.27	0.17	0.40	0.34	0.48	0.39
AGE										
12-17	0.00	0.00	0.56	0.84	0.46	*	0.33	0.13	0.56	0.84
18-25	0.00	0.00	0.34	0.37	0.27	0.25	0.21	0.27	0.34	0.37
26 or Older	0.00	0.00	0.67	0.52	0.38	0.22	0.56	0.46	0.67	0.52
GENDER										
Male	0.00	0.00	0.64	0.39	0.39	0.19	0.51	0.33	0.64	0.39
Female	0.00	0.00	0.70	0.81	0.31	0.34	0.63	0.72	0.70	0.81

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.50P Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment ¹ (2013)	Felt Need for Treatment ¹ (2014)	Felt Need and Made Effort to Get Treatment ¹ (2013)	Felt Need and Made Effort to Get Treatment ¹ (2014)	Felt Need and Made No Effort to Get Treatment ¹ (2013)	Felt Need and Made No Effort to Get Treatment ¹ (2014)	Did Not Feel Need for Treatment ¹ (2013)	Did Not Feel Need for Treatment ¹ (2014)
TOTAL	N/A		0.4431		0.1856		0.9329		0.4431	
AGE										
12-17	N/A		0.4739		*		0.0485		0.4739	
18-25	N/A		0.9913		0.7327		0.7260		0.9913	
26 or Older	N/A		0.4438		0.1970		0.9051		0.4438	
GENDER										
Male	N/A		0.0377		0.1048		0.1636		0.0377	
Female	N/A		0.1711		0.8769		0.1463		0.1711	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.51C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	477	404	160	148	444	373	0.66	0.61
AGE								
12-17	51	55	15	16	49	54	1.06	1.21
18-25	138	135	35	41	133	130	0.55	0.67
26 or Older	443	363	153	139	418	335	0.95	0.84
GENDER								
Male	376	323	127	126	354	302	0.83	0.82
Female	284	222	99	73	268	210	1.13	0.85
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	424	364	143	134	397	334	0.71	0.66
White	398	321	129	114	375	297	0.81	0.74
Black or African American	146	137	56	62	132	124	2.04	2.00
American Indian or Alaska Native	48	47	13	9	46	43	*	3.32
Native Hawaiian or Other Pacific Islander	36	29	4	8	*	28	*	*
Asian	84	68	23	7	80	68	*	1.12
Two or More Races	64	55	18	15	61	54	3.42	2.87
Hispanic or Latino	206	176	77	66	188	168	1.88	1.65

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.51D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.18	0.15	0.06	0.06	0.17	0.14	0.66	0.61
AGE								
12-17	0.21	0.22	0.06	0.06	0.19	0.22	1.06	1.21
18-25	0.40	0.39	0.10	0.12	0.38	0.37	0.55	0.67
26 or Older	0.22	0.18	0.08	0.07	0.21	0.16	0.95	0.84
GENDER								
Male	0.30	0.25	0.10	0.10	0.28	0.24	0.83	0.82
Female	0.21	0.16	0.07	0.05	0.20	0.15	1.13	0.85
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.19	0.16	0.06	0.06	0.18	0.15	0.71	0.66
White	0.23	0.19	0.08	0.07	0.22	0.17	0.81	0.74
Black or African American	0.47	0.43	0.18	0.19	0.42	0.39	2.04	2.00
American Indian or Alaska Native	2.99	2.53	0.92	0.59	2.88	2.34	*	3.32
Native Hawaiian or Other Pacific Islander	3.81	2.78	0.48	0.74	*	2.68	*	*
Asian	0.63	0.50	0.17	0.05	0.60	0.50	*	1.12
Two or More Races	1.48	1.20	0.42	0.32	1.40	1.18	3.42	2.87
Hispanic or Latino	0.50	0.42	0.19	0.15	0.46	0.40	1.88	1.65

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.51P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.4797		0.6008		0.3377		0.4181	
AGE								
12-17	0.4602		0.5349		0.5612		0.6796	
18-25	0.0913		0.8445		0.0701		0.4921	
26 or Older	0.9362		0.5729		0.7619		0.5261	
GENDER								
Male	0.6004		0.4572		0.3992		0.3343	
Female	0.6152		0.8790		0.6334		0.9887	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.5229		0.5746		0.3690		0.4087	
White	0.2559		0.6829		0.1735		0.3987	
Black or African American	0.1360		0.3045		0.2401		0.5486	
American Indian or Alaska Native	0.9592		0.9042		0.9852		*	
Native Hawaiian or Other Pacific Islander	0.8558		0.6394		*		*	
Asian	0.7742		0.2088		1.0000		*	
Two or More Races	0.7804		0.5219		0.9162		0.5782	
Hispanic or Latino	0.7406		0.9660		0.7092		0.8827	

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.52C Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	472	399	159	148	439	367	0.69	0.64
GENDER								
Male	374	319	126	125	352	299	0.86	0.85
Female	282	221	98	72	266	209	1.22	0.91
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	421	356	142	133	394	326	0.74	0.69
White	426	328	127	114	403	302	0.85	0.77
Black or African American	153	151	56	63	138	134	2.20	2.09
American Indian or Alaska Native	48	47	13	9	46	43	*	3.40
Native Hawaiian or Other Pacific Islander	*	28	*	8	*	27	*	*
Asian	83	67	23	7	80	67	*	1.16
Two or More Races	64	55	18	14	61	53	3.64	3.02
Hispanic or Latino	203	174	77	65	185	166	2.04	1.78
EDUCATION								
< High School	186	163	77	75	164	145	2.11	2.09
High School Graduate	250	210	78	86	237	190	1.27	1.37
Some College	284	223	94	72	262	201	1.34	0.99
College Graduate	304	239	67	47	295	232	1.12	0.79
CURRENT EMPLOYMENT								
Full-Time	393	308	67	84	384	293	0.56	0.68
Part-Time	169	166	46	44	164	159	1.37	1.27
Unemployed	131	100	74	49	106	85	3.04	2.40
Other ¹	243	199	112	100	217	172	2.23	2.00

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.52D Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.20	0.17	0.07	0.06	0.18	0.15	0.69	0.64
GENDER								
Male	0.33	0.28	0.11	0.11	0.31	0.26	0.86	0.85
Female	0.23	0.18	0.08	0.06	0.22	0.17	1.22	0.91
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.21	0.17	0.07	0.07	0.20	0.16	0.74	0.69
White	0.25	0.20	0.08	0.07	0.24	0.19	0.85	0.77
Black or African American	0.52	0.48	0.20	0.22	0.48	0.44	2.20	2.09
American Indian or Alaska Native	3.29	2.77	1.02	0.66	3.19	2.57	*	3.40
Native Hawaiian or Other Pacific Islander	*	3.01	*	0.82	*	2.89	*	*
Asian	0.69	0.54	0.19	0.05	0.66	0.54	*	1.16
Two or More Races	1.82	1.46	0.51	0.39	1.73	1.43	3.64	3.02
Hispanic or Latino	0.57	0.47	0.22	0.18	0.52	0.45	2.04	1.78
EDUCATION								
< High School	0.55	0.48	0.24	0.23	0.49	0.43	2.11	2.09
High School Graduate	0.35	0.29	0.11	0.12	0.33	0.26	1.27	1.37
Some College	0.42	0.32	0.15	0.11	0.39	0.29	1.34	0.99
College Graduate	0.39	0.31	0.09	0.06	0.38	0.30	1.12	0.79
CURRENT EMPLOYMENT								
Full-Time	0.31	0.23	0.06	0.07	0.30	0.22	0.56	0.68
Part-Time	0.48	0.48	0.14	0.13	0.47	0.46	1.37	1.27
Unemployed	1.06	0.84	0.62	0.44	0.87	0.73	3.04	2.40
Other ¹	0.33	0.27	0.15	0.14	0.29	0.23	2.23	2.00

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.52P Need for and Receipt of Treatment at a Specialty Facility for an Illicit Drug or Alcohol Problem in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Needed Treatment (2013)	Needed Treatment (2014)	Needed and Received Treatment (2013)	Needed and Received Treatment (2014)	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Percentage Who Received Treatment among Persons Who Needed Treatment (2013)	Percentage Who Received Treatment among Persons Who Needed Treatment (2014)
TOTAL	0.5217		0.5609		0.3615		0.3940	
GENDER								
Male	0.6344		0.3766		0.3976		0.2707	
Female	0.6498		0.7997		0.6972		0.9141	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.5961		0.5701		0.4321		0.4252	
White	0.2819		0.7105		0.1982		0.4322	
Black or African American	0.1123		0.2914		0.2067		0.5640	
American Indian or Alaska Native	0.9484		0.9485		0.9612		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.7402		0.2076		0.9631		*	
Two or More Races	0.8324		0.5433		0.9622		0.5838	
Hispanic or Latino	0.6814		0.8756		0.6130		0.7673	
EDUCATION								
< High School	0.8270		0.4154		0.5093		0.3119	
High School Graduate	0.9110		0.2477		0.5609		0.1964	
Some College	0.7850		0.8811		0.8097		0.9463	
College Graduate	0.5168		0.3924		0.6404		0.4686	
CURRENT EMPLOYMENT								
Full-Time	0.7004		0.0083		0.3113		0.0046	
Part-Time	0.5764		0.4844		0.4328		0.3566	
Unemployed	0.5964		0.2429		0.8800		0.2525	
Other ¹	0.5362		0.8414		0.4160		0.5868	

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.53C Perceived Need for Illicit Drug or Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug or Alcohol Problem, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment¹ (2013)	Felt Need for Treatment¹ (2014)	Felt Need and Made Effort to Get Treatment¹ (2013)	Felt Need and Made Effort to Get Treatment¹ (2014)	Felt Need and Made No Effort to Get Treatment¹ (2013)	Felt Need and Made No Effort to Get Treatment¹ (2014)	Did Not Feel Need for Treatment¹ (2013)	Did Not Feel Need for Treatment¹ (2014)
TOTAL	481	393	96	73	55	36	78	61	470	388
AGE										
12-17	50	55	7	9	4	8	5	4	50	54
18-25	162	150	21	25	14	14	16	19	160	149
26 or Older	448	351	93	68	53	32	76	58	436	347
GENDER										
Male	376	324	82	50	49	26	65	41	368	320
Female	283	214	49	53	24	24	43	46	278	209

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs or alcohol, but have not received treatment for an illicit drug or alcohol problem at a specialty facility.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug or alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.53D Perceived Need for Illicit Drug or Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug or Alcohol Problem, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment¹ (2013)	Felt Need for Treatment¹ (2014)	Felt Need and Made Effort to Get Treatment¹ (2013)	Felt Need and Made Effort to Get Treatment¹ (2014)	Felt Need and Made No Effort to Get Treatment¹ (2013)	Felt Need and Made No Effort to Get Treatment¹ (2014)	Did Not Feel Need for Treatment¹ (2013)	Did Not Feel Need for Treatment¹ (2014)
TOTAL	0.00	0.00	0.46	0.36	0.27	0.18	0.38	0.30	0.46	0.36
AGE										
12-17	0.00	0.00	0.59	0.74	0.31	0.68	0.42	0.31	0.59	0.74
18-25	0.00	0.00	0.37	0.45	0.24	0.26	0.28	0.35	0.37	0.45
26 or Older	0.00	0.00	0.68	0.50	0.40	0.24	0.56	0.43	0.68	0.50
GENDER										
Male	0.00	0.00	0.62	0.39	0.38	0.21	0.50	0.32	0.62	0.39
Female	0.00	0.00	0.66	0.71	0.33	0.34	0.58	0.62	0.66	0.71

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs or alcohol, but have not received treatment for an illicit drug or alcohol problem at a specialty facility.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug or alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.53P Perceived Need for Illicit Drug or Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Illicit Drug or Alcohol Problem, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Needed but Did Not Receive Treatment (2013)	Needed but Did Not Receive Treatment (2014)	Felt Need for Treatment ¹ (2013)	Felt Need for Treatment ¹ (2014)	Felt Need and Made Effort to Get Treatment ¹ (2013)	Felt Need and Made Effort to Get Treatment ¹ (2014)	Felt Need and Made No Effort to Get Treatment ¹ (2013)	Felt Need and Made No Effort to Get Treatment ¹ (2014)	Did Not Feel Need for Treatment ¹ (2013)	Did Not Feel Need for Treatment ¹ (2014)
TOTAL	N/A		0.4201		0.2204		0.8762		0.4201	
AGE										
12-17	N/A		0.8715		0.2119		0.1324		0.8715	
18-25	N/A		0.5693		0.9365		0.4200		0.5693	
26 or Older	N/A		0.2868		0.1436		0.7513		0.2868	
GENDER										
Male	N/A		0.0189		0.0663		0.1187		0.0189	
Female	N/A		0.0775		0.5227		0.0946		0.0775	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs or alcohol, but have not received treatment for an illicit drug or alcohol problem at a specialty facility.

¹ Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug or alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 5.54C Detailed Reasons for Not Receiving Illicit Drug Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2014

Reason Did Not Receive Illicit Drug Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	30	17	24
No Health Coverage and Could Not Afford Cost	22	13	17
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	11	7	8
No Transportation/Inconvenient	8	6	5
No Program Having Type of Treatment	8	7	5
Not Ready to Stop Using	16	7	14
No Openings in a Program	9	*	3
Did Not Know Where to Go for Treatment	11	6	9
Might Cause Neighbors/Community to Have Negative Opinion	14	6	13
Might Have Negative Effect on Job	13	6	12
Did Not Feel Need for Treatment at the Time	3	3	2
Could Handle the Problem Without Treatment	9	5	7
Treatment Would Not Help	6	0	6
Did Not Have Time	9	4	8
Did Not Want Others to Find Out	8	3	7
Some Other Reason	4	2	3

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but have not received treatment for an illicit drug problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 5.54D Detailed Reasons for Not Receiving Illicit Drug Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2011-2014

Reason Did Not Receive Illicit Drug Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	0.00	0.00	0.00
No Health Coverage and Could Not Afford Cost	3.36	5.31	4.27
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	2.17	3.70	2.67
No Transportation/Inconvenient	1.63	3.41	1.62
No Program Having Type of Treatment	1.73	3.68	1.63
Not Ready to Stop Using	2.90	3.76	3.94
No Openings in a Program	1.91	*	1.08
Did Not Know Where to Go for Treatment	2.25	3.45	2.91
Might Cause Neighbors/Community to Have Negative Opinion	2.75	3.11	3.83
Might Have Negative Effect on Job	2.53	3.21	3.44
Did Not Feel Need for Treatment at the Time	0.69	1.53	0.65
Could Handle the Problem Without Treatment	1.88	3.11	2.36
Treatment Would Not Help	1.31	0.11	2.04
Did Not Have Time	1.82	2.54	2.48
Did Not Want Others to Find Out	1.59	1.63	2.31
Some Other Reason	0.77	1.34	0.94

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but have not received treatment for an illicit drug problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 5.55C Detailed Reasons for Not Receiving Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Alcohol Treatment at a Specialty Facility and Who Felt a Need for Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2014

Reason Did Not Receive Alcohol Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	37	19	33
No Health Coverage and Could Not Afford Cost	18	*	14
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	9	7	5
No Transportation/Inconvenient	10	*	7
No Program Having Type of Treatment	14	*	11
Not Ready to Stop Using	30	10	28
No Openings in a Program	5	*	1
Did Not Know Where to Go for Treatment	9	*	7
Might Cause Neighbors/Community to Have Negative Opinion	8	*	7
Might Have Negative Effect on Job	10	5	8
Did Not Feel Need for Treatment at the Time	6	4	4
Could Handle the Problem Without Treatment	9	*	7
Treatment Would Not Help	4	0	4
Did Not Have Time	7	*	6
Did Not Want Others to Find Out	4	0	4
Some Other Reason	7	4	6

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 5.55D Detailed Reasons for Not Receiving Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Alcohol Treatment at a Specialty Facility and Who Felt a Need for Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2011-2014

Reason Did Not Receive Alcohol Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	0.00	0.00	0.00
No Health Coverage and Could Not Afford Cost	2.92	*	3.33
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	1.59	4.13	1.36
No Transportation/Inconvenient	1.75	*	1.86
No Program Having Type of Treatment	2.36	*	2.72
Not Ready to Stop Using	3.46	5.29	3.92
No Openings in a Program	0.87	*	0.30
Did Not Know Where to Go for Treatment	1.68	*	1.82
Might Cause Neighbors/Community to Have Negative Opinion	1.50	*	1.80
Might Have Negative Effect on Job	1.74	3.04	2.10
Did Not Feel Need for Treatment at the Time	1.10	2.61	1.11
Could Handle the Problem Without Treatment	1.59	*	1.77
Treatment Would Not Help	0.71	0.25	0.99
Did Not Have Time	1.20	*	1.52
Did Not Want Others to Find Out	0.82	0.26	1.15
Some Other Reason	1.22	2.28	1.45

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 5.56C Detailed Reasons for Not Receiving Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug or Alcohol Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug or Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2014

Reason Did Not Receive Illicit Drug or Alcohol Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	45	24	38
No Health Coverage and Could Not Afford Cost	26	16	21
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	13	8	10
No Transportation/Inconvenient	12	8	9
No Program Having Type of Treatment	15	9	12
Not Ready to Stop Using	31	11	30
No Openings in a Program	9	9	3
Did Not Know Where to Go for Treatment	14	8	11
Might Cause Neighbors/Community to Have Negative Opinion	16	6	15
Might Have Negative Effect on Job	15	6	14
Did Not Feel Need for Treatment at the Time	7	5	5
Could Handle the Problem Without Treatment	12	6	10
Treatment Would Not Help	7	1	7
Did Not Have Time	11	5	10
Did Not Want Others to Find Out	9	3	8
Some Other Reason	6	4	4

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs or alcohol, but have not received treatment for an illicit drug or alcohol problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug or alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 5.56D Detailed Reasons for Not Receiving Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Illicit Drug or Alcohol Treatment at a Specialty Facility and Who Felt a Need for Illicit Drug or Alcohol Treatment in the Past Year, by Whether Made an Effort to Get Treatment in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2011-2014

Reason Did Not Receive Illicit Drug or Alcohol Treatment¹	Total	Made Effort	Made No Effort
TOTAL POPULATION	0.00	0.00	0.00
No Health Coverage and Could Not Afford Cost	2.34	4.18	2.78
Had Health Coverage But Did Not Cover Treatment or Did Not Cover Cost	1.31	2.68	1.45
No Transportation/Inconvenient	1.24	2.54	1.37
No Program Having Type of Treatment	1.53	2.87	1.80
Not Ready to Stop Using	2.48	3.32	3.09
No Openings in a Program	1.00	2.89	0.53
Did Not Know Where to Go for Treatment	1.40	2.47	1.70
Might Cause Neighbors/Community to Have Negative Opinion	1.59	1.88	2.14
Might Have Negative Effect on Job	1.55	1.96	2.05
Did Not Feel Need for Treatment at the Time	0.74	1.58	0.80
Could Handle the Problem Without Treatment	1.24	2.02	1.56
Treatment Would Not Help	0.78	0.22	1.13
Did Not Have Time	1.13	1.53	1.49
Did Not Want Others to Find Out	0.94	0.96	1.29
Some Other Reason	0.62	1.46	0.60

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs or alcohol, but have not received treatment for an illicit drug or alcohol problem at a specialty facility.

NOTE: Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an illicit drug or alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

¹ Respondents could indicate multiple reasons for not receiving treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.1A Number of Days Used Marijuana in the Past Year among Past Year Users and the Number of Days Used Marijuana in the Past Month among Past Month Users, by Age Group: Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	32,952 ^b	35,124	3,346	3,262	11,000	11,154	18,606 ^b	20,707
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	9,967	10,191	1,232	1,117	3,255	3,159	5,480	5,914
12-49	6,133	6,283	687	742	1,970	1,918	3,475	3,624
50-99	3,230	3,688	349	396	1,064	1,060	1,816	2,231
100-299	7,894	8,454	779	750	2,544	2,792	4,571	4,911
300 or More	5,729 ^a	6,508	299	257	2,166	2,224	3,264 ^a	4,027
TOTAL PAST MONTH USERS	19,810 ^b	22,188	1,762	1,830	6,636	6,836	11,412 ^b	13,521
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	4,621	5,027	529	592	1,405	1,502	2,686	2,933
3-5	3,256	3,442	378	386	995	930	1,882	2,126
6-19	3,790 ^b	4,479	404	452	1,418	1,384	1,968 ^b	2,643
20 or More	8,144 ^b	9,240	451	400	2,817	3,020	4,875 ^a	5,820

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.1B Number of Days Used Marijuana in the Past Year among Past Year Users and the Number of Days Used Marijuana in the Past Month among Past Month Users, by Age Group: Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	12.6 ^a	13.2	13.4	13.1	31.6	31.9	9.2 ^b	10.1
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	30.2	29.0	36.8	34.2	29.6	28.3	29.5	28.6
12-49	18.6	17.9	20.5	22.7	17.9	17.2	18.7	17.5
50-99	9.8	10.5	10.4	12.2	9.7	9.5	9.8	10.8
100-299	24.0	24.1	23.3	23.0	23.1	25.0	24.6	23.7
300 or More	17.4	18.5	8.9	7.9	19.7	19.9	17.5	19.4
TOTAL PAST MONTH USERS	7.5 ^b	8.4	7.1	7.4	19.1	19.6	5.6 ^b	6.6
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	23.3	22.7	30.0	32.4	21.2	22.0	23.5	21.7
3-5	16.4	15.5	21.4	21.1	15.0	13.6	16.5	15.7
6-19	19.1	20.2	22.9	24.7	21.4	20.2	17.2	19.5
20 or More	41.1	41.6	25.6	21.8	42.5	44.2	42.7	43.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.2A Number of Days Used Cocaine in the Past Year among Past Year Users and the Number of Days Used Cocaine in the Past Month among Past Month Users, by Age Group: Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	4,182	4,553	131	168	1,543	1,604	2,508	2,782
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	2,418	2,598	89	107	1,032	1,041	1,298	1,449
12-49	855	939	20	36	285	332	550	570
50-99	337 ^a	502	*	9	118	129	208 ^a	364
100-299	421	449	11	14	103	93	307	342
300 or More	151	66	*	*	6	8	145	57
TOTAL PAST MONTH USERS	1,549	1,530	43	39	398	473	1,108	1,018
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	821	752	*	*	243	294	546	441
3-5	304	378	*	*	84	116	214	254
6-19	240	284	*	*	62	54	173	222
20 or More	185	116	*	*	8	8	*	100

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.2B Number of Days Used Cocaine in the Past Year among Past Year Users and the Number of Days Used Cocaine in the Past Month among Past Month Users, by Age Group: Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	1.6	1.7	0.5	0.7	4.4	4.6	1.2	1.4
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	57.8	57.1	67.7	64.0	66.9	64.9	51.7	52.1
12-49	20.4	20.6	15.1	21.6	18.4	20.7	22.0	20.5
50-99	8.0	11.0	*	5.2	7.6	8.1	8.3	13.1
100-299	10.1	9.9	8.5	8.5	6.7	5.8	12.2	12.3
300 or More	3.6	1.5	*	*	0.4	0.5	5.8	2.1
TOTAL PAST MONTH USERS	0.6	0.6	0.2	0.2	1.1	1.4	0.5	0.5
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	53.0	49.1	*	*	61.1	62.3	49.3	43.3
3-5	19.6	24.7	*	*	21.1	24.6	19.3	25.0
6-19	15.5	18.5	*	*	15.7	11.4	15.6	21.9
20 or More	11.9	7.6	*	*	2.1	1.7	*	9.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.3A Number of Days Used Hallucinogens in the Past Year among Past Year Users and the Number Days Used Hallucinogens in the Past Month among Past Month Users, by Age Group: Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	4,430	4,250	482	422	2,338	2,205	1,610	1,624
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	3,383	3,257	344	307	1,869	1,688	1,170	1,262
12-49	609	540	88	58	268	311	*	171
50-99	208	271	34	31	95	128	79	112
100-299	165	147	13	21	94	68	59	58
300 or More	64	35	3	5	12	9	*	21
TOTAL PAST MONTH USERS	1,333	1,173	154	136	617	502	562	535
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	997	864	124	*	489 ^a	370	*	401
3-5	192	210	13	*	91	99	*	82
6-19	82	54	13	*	34	23	*	21
20 or More	63	45	4	*	3	9	*	31

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.3B Number of Days Used Hallucinogens in the Past Year among Past Year Users and the Number Days Used Hallucinogens in the Past Month among Past Month Users, by Age Group: Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	1.7	1.6	1.9	1.7	6.7	6.3	0.8	0.8
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	76.4	76.6	71.4	72.8	79.9	76.6	72.7	77.7
12-49	13.8	12.7	18.3	13.8	11.5	14.1	*	10.5
50-99	4.7	6.4	7.0	7.3	4.1	5.8	4.9	6.9
100-299	3.7	3.5	2.7	5.0	4.0	3.1	3.6	3.6
300 or More	1.4	0.8	0.6	1.1	0.5	0.4	*	1.3
TOTAL PAST MONTH USERS	0.5	0.4	0.6	0.5	1.8	1.4	0.3	0.3
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	74.8	73.6	80.2	*	79.3	73.8	*	74.9
3-5	14.4	17.9	8.5	*	14.8	19.7	*	15.4
6-19	6.1	4.6	8.5	*	5.4	4.6	*	3.9
20 or More	4.7	3.9	2.8	*	0.6	1.9	*	5.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.4A Number of Days Used Inhalants in the Past Year among Past Year Users and the Number of Days Used Inhalants in the Past Month among Past Month Users, by Age Group: Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	1,533	1,617	480	512	478	379	575	725
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	955	910	299	268	351	273	*	*
12-49	363	402	87 ^a	140	88	80	*	*
50-99	132	154	53	45	17	16	*	*
100-299	74	138	40	51	16	6	*	81
300 or More	8	12	1	9	7	3	*	*
TOTAL PAST MONTH USERS	496	546	121	149	106	80	269	316
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	356	296	83	*	*	*	*	*
3-5	*	109	18	35	*	*	*	*
6-19	46	106	12	*	*	*	*	*
20 or More	16	*	*	*	*	*	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.4B Number of Days Used Inhalants in the Past Year among Past Year Users and the Number of Days Used Inhalants in the Past Month among Past Month Users, by Age Group: Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	0.6	0.6	1.9	2.1	1.4	1.1	0.3	0.4
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	62.3	56.3	62.3 ^a	52.2	73.4	72.1	*	*
12-49	23.7	24.9	18.0 ^a	27.3	18.3	21.2	*	*
50-99	8.6	9.5	11.0	8.7	3.5	4.3	*	*
100-299	4.8	8.6	8.4	10.0	3.3	1.7	*	11.1
300 or More	0.5	0.7	0.2	1.7	1.5	0.7	*	*
TOTAL PAST MONTH USERS	0.2	0.2	0.5	0.6	0.3	0.2	0.1	0.2
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	71.8 ^a	54.3	68.5	*	*	*	*	*
3-5	*	20.0	14.6	23.8	*	*	*	*
6-19	9.2	19.4	9.9	*	*	*	*	*
20 or More	3.2	*	*	*	*	*	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.5A Number of Days Used Cigarettes in the Past Month among Past Month Users and the Number of Cigarettes Smoked Per Day among Daily Smokers, by Age Group: Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST MONTH USERS	55,778	55,241	1,397 ^a	1,213	10,657 ^b	9,905	43,724	44,123
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	5,244	5,558	405	357	1,688	1,498	3,151 ^a	3,703
3-5	4,229	4,264	258	202	1,220	1,078	2,751	2,983
6-19	6,544	6,959	270	231	1,745	1,700	4,530	5,028
20-29	6,513	5,988	193 ^a	130	1,412	1,366	4,908	4,493
30	33,247	32,471	271	292	4,591	4,263	28,385	27,916
TOTAL DAILY SMOKERS¹	33,247	32,471	271	292	4,591 ^a	4,263	28,385	27,916
NUMBER OF CIGARETTES SMOKED PER DAY AMONG DAILY SMOKERS¹								
Fewer Than 6	6,862	6,675	127	145	1,463	1,438	5,272	5,091
6-15 (1/2 Pack)	12,635	12,662	111	112	2,106 ^a	1,860	10,419	10,689
16-25 (1 Pack)	10,407	9,936	22	27	887	870	9,497	9,039
26 or More (More Than 1 Pack)	3,314	3,123	10	8	134 ^a	89	3,170	3,026
Not Reported	29	77	*	*	1	6	28	71

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Daily Cigarette Use is defined as smoking on each of the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.5B Number of Days Used Cigarettes in the Past Month among Past Month Users and the Number of Cigarettes Smoked Per Day among Daily Smokers, by Age Group: Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST MONTH USERS	21.3	20.8	5.6 ^a	4.9	30.6 ^b	28.4	21.6	21.5
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	9.4	10.1	29.0	29.4	15.8	15.1	7.2 ^a	8.4
3-5	7.6	7.7	18.5	16.7	11.5	10.9	6.3	6.8
6-19	11.7	12.6	19.3	19.1	16.4	17.2	10.4	11.4
20-29	11.7	10.8	13.8	10.7	13.2	13.8	11.2	10.2
30	59.6	58.8	19.4 ^a	24.1	43.1	43.0	64.9	63.3
TOTAL DAILY SMOKERS¹	12.7	12.2	1.1	1.2	13.2 ^a	12.2	14.0	13.6
NUMBER OF CIGARETTES SMOKED PER DAY AMONG DAILY SMOKERS¹								
Fewer Than 6	20.6	20.6	46.8	49.7	31.9	33.7	18.6	18.2
6-15 (1/2 Pack)	38.0	39.0	40.9	38.3	45.9	43.6	36.7	38.3
16-25 (1 Pack)	31.3	30.6	8.1	9.2	19.3	20.4	33.5	32.4
26 or More (More Than 1 Pack)	10.0	9.6	3.8	2.7	2.9	2.1	11.2	10.8
Not Reported	0.1	0.2	*	*	0.0	0.1	0.1	0.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Daily Cigarette Use is defined as smoking on each of the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.6A Number of Days Used Alcohol in the Past Year among Past Year Users and the Number of Days Used Alcohol in the Past Month among Past Month Users, by Age Group: Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	173,966 ^a	176,556	6,133	5,979	26,707	26,713	141,126 ^a	143,864
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	37,751	36,571	3,372	3,219	5,590	5,611	28,789	27,741
12-49	45,110	45,441	1,556	1,548	7,774	7,563	35,781	36,330
50-99	26,900 ^a	28,753	687	674	4,844	5,029	21,369 ^a	23,050
100-299	53,618	54,552	475	477	7,901	7,866	45,242	46,209
300 or More	10,587	11,239	43	61	599	645	9,945	10,533
TOTAL PAST MONTH USERS	136,868 ^a	139,677	2,877	2,872	20,738	20,815	113,253 ^a	115,990
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	38,673	38,837	1,588	1,606	6,114	6,031	30,971	31,199
3-5	36,148	36,876	806	776	6,238	6,337	29,104	29,763
6-19	40,785	41,372	427	436	6,727	6,797	33,631	34,139
20 or More	21,262	22,593	55	55	1,660	1,649	19,547	20,889

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.6B Number of Days Used Alcohol in the Past Year among Past Year Users and the Number of Days Used Alcohol in the Past Month among Past Month Users, by Age Group: Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	66.3	66.6	24.6	24.0	76.8	76.5	69.6	70.1
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	21.7 ^a	20.7	55.0	53.8	20.9	21.0	20.4 ^a	19.3
12-49	25.9	25.7	25.4	25.9	29.1	28.3	25.4	25.3
50-99	15.5 ^a	16.3	11.2	11.3	18.1	18.8	15.1	16.0
100-299	30.8	30.9	7.7	8.0	29.6	29.4	32.1	32.1
300 or More	6.1	6.4	0.7	1.0	2.2	2.4	7.0	7.3
TOTAL PAST MONTH USERS	52.2	52.7	11.6	11.5	59.6	59.6	55.9	56.5
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	28.3	27.8	55.2	55.9	29.5	29.0	27.3	26.9
3-5	26.4	26.4	28.0	27.0	30.1	30.4	25.7	25.7
6-19	29.8	29.6	14.8	15.2	32.4	32.7	29.7	29.4
20 or More	15.5	16.2	1.9	1.9	8.0	7.9	17.3	18.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.7A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	13,470	13,901	11,104 ^b	13,082
Marijuana and Hashish	11,424	12,062	8,387 ^b	10,126
Cocaine	1,158	1,137	391	393
Crack	287	290	90	64
Heroin	248 ^a	384	42	52
Hallucinogens	822	784	511	389
LSD	146	197	82	90
PCP	19	10	14	*
Ecstasy	420	396	240	214
Inhalants	224	195	272	351
Nonmedical Use of Psychotherapeutics ^{2,3}	3,549	3,187	2,935	3,350
Pain Relievers	2,510	2,137	2,012	2,188
OxyContin [®]	344 ^a	203	148	153
Tranquilizers	1,057	1,106	648	770
Stimulants ³	934	871	431 ^b	724
Methamphetamine ³	471	346	124	223
Sedatives	83 ^a	190	168	140
ILLCIT DRUGS OTHER THAN MARIJUANA¹	4,917	4,541	3,748	4,178

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.7B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	24.1	25.2	5.4 ^b	6.2
Marijuana and Hashish	20.5	21.8	4.1 ^b	4.8
Cocaine	2.1	2.1	0.2	0.2
Crack	0.5	0.5	0.0	0.0
Heroin	0.4 ^a	0.7	0.0	0.0
Hallucinogens	1.5	1.4	0.2	0.2
LSD	0.3	0.4	0.0	0.0
PCP	0.0	0.0	0.0	*
Ecstasy	0.8	0.7	0.1	0.1
Inhalants	0.4	0.4	0.1	0.2
Nonmedical Use of Psychotherapeutics ^{2,3}	6.4	5.8	1.4	1.6
Pain Relievers	4.5	3.9	1.0	1.0
OxyContin [®]	0.6 ^a	0.4	0.1	0.1
Tranquilizers	1.9	2.0	0.3	0.4
Stimulants ³	1.7	1.6	0.2 ^b	0.3
Methamphetamine ³	0.8	0.6	0.1	0.1
Sedatives	0.1 ^a	0.3	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	8.8	8.2	1.8	2.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.8A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	752	683	1,445 ^a	1,655
Marijuana and Hashish	692	628	1,070	1,202
Cocaine	26	32	18	8
Crack	1	8	2	*
Heroin	9	13	3	3
Hallucinogens	101	72	53	64
LSD	41	38	10	27
PCP	5	3	2	*
Ecstasy	30	26	19	13
Inhalants	27	34	94	114
Nonmedical Use of Psychotherapeutics ^{2,3}	171	161	378 ^a	494
Pain Relievers	120	104	305	363
OxyContin [®]	14	14	15	21
Tranquilizers	51	54	40	49
Stimulants ³	40	51	44 ^b	118
Methamphetamine ³	11	19	11	26
Sedatives	7	7	18	34
ILLCIT DRUGS OTHER THAN MARIJUANA¹	262	249	494 ^b	636

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.8B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	53.9	56.3	6.1 ^a	7.0
Marijuana and Hashish	49.5	51.8	4.6	5.1
Cocaine	1.8	2.6	0.1	0.0
Crack	0.1	0.6	0.0	*
Heroin	0.7	1.0	0.0	0.0
Hallucinogens	7.2	6.0	0.2	0.3
LSD	3.0	3.2	0.0	0.1
PCP	0.4	0.3	0.0	*
Ecstasy	2.1	2.2	0.1	0.1
Inhalants	1.9	2.8	0.4	0.5
Nonmedical Use of Psychotherapeutics ^{2,3}	12.2	13.3	1.6 ^a	2.1
Pain Relievers	8.6	8.6	1.3	1.5
OxyContin [®]	1.0	1.2	0.1	0.1
Tranquilizers	3.6	4.5	0.2	0.2
Stimulants ³	2.9	4.2	0.2 ^b	0.5
Methamphetamine ³	0.8	1.6	0.0	0.1
Sedatives	0.5	0.6	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	18.8	20.5	2.1 ^b	2.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.9A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	4,443	4,376	3,047	3,300
Marijuana and Hashish	4,028	3,943	2,608	2,894
Cocaine	331	358	67	115
Crack	36	14	*	15
Heroin	89	81	3	2
Hallucinogens	428	381	189 ^a	120
LSD	76	97	29	22
PCP	2	3	2	2
Ecstasy	237	196	83	73
Inhalants	69	47	37	33
Nonmedical Use of Psychotherapeutics ^{2,3}	1,064	942	597	609
Pain Relievers	740	617	392	361
OxyContin [®]	92	86	15	16
Tranquilizers	317	288	117	128
Stimulants ³	313	246	149	160
Methamphetamine ³	104	62	16	23
Sedatives	14	25	21	31
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1,550	1,436	788	812

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.9B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	41.7 ^a	44.2	12.6	13.2
Marijuana and Hashish	37.8	39.8	10.8	11.6
Cocaine	3.1	3.6	0.3	0.5
Crack	0.3	0.1	*	0.1
Heroin	0.8	0.8	0.0	0.0
Hallucinogens	4.0	3.9	0.8 ^a	0.5
LSD	0.7	1.0	0.1	0.1
PCP	0.0	0.0	0.0	0.0
Ecstasy	2.2	2.0	0.3	0.3
Inhalants	0.6	0.5	0.2	0.1
Nonmedical Use of Psychotherapeutics ^{2,3}	10.0	9.5	2.5	2.4
Pain Relievers	6.9	6.2	1.6	1.4
OxyContin [®]	0.9	0.9	0.1	0.1
Tranquilizers	3.0	2.9	0.5	0.5
Stimulants ³	2.9	2.5	0.6	0.6
Methamphetamine ³	1.0	0.6	0.1	0.1
Sedatives	0.1	0.3	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	14.5	14.5	3.3	3.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.10A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	8,274	8,842	6,611 ^b	8,127
Marijuana and Hashish	6,704	7,491	4,708 ^b	6,030
Cocaine	801	748	306	270
Crack	250	268	88	49
Heroin	150 ^a	290	35	47
Hallucinogens	293	330	269	205
LSD	29	62	43	42
PCP	12	3	10	*
Ecstasy	153	173	138	127
Inhalants	128	114	141	203
Nonmedical Use of Psychotherapeutics ^{2,3}	2,314	2,083	1,960	2,247
Pain Relievers	1,649	1,416	1,315	1,464
OxyContin [®]	238 ^a	103	118	116
Tranquilizers	689	763	492	592
Stimulants ³	581	574	238 ^a	447
Methamphetamine ³	356	265	97	173
Sedatives	62	158	130	74
ILLCIT DRUGS OTHER THAN MARIJUANA¹	3,104	2,856	2,466	2,730

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.10B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	18.9	20.0	4.2 ^b	5.0
Marijuana and Hashish	15.3 ^a	17.0	3.0 ^b	3.7
Cocaine	1.8	1.7	0.2	0.2
Crack	0.6	0.6	0.1	0.0
Heroin	0.3 ^a	0.7	0.0	0.0
Hallucinogens	0.7	0.7	0.2	0.1
LSD	0.1	0.1	0.0	0.0
PCP	0.0	0.0	0.0	*
Ecstasy	0.4	0.4	0.1	0.1
Inhalants	0.3	0.3	0.1	0.1
Nonmedical Use of Psychotherapeutics ^{2,3}	5.3	4.7	1.2	1.4
Pain Relievers	3.8	3.2	0.8	0.9
OxyContin [®]	0.5 ^a	0.2	0.1	0.1
Tranquilizers	1.6	1.7	0.3	0.4
Stimulants ³	1.3	1.3	0.1 ^a	0.3
Methamphetamine ³	0.8	0.6	0.1	0.1
Sedatives	0.1	0.4	0.1	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA¹	7.1	6.5	1.6	1.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.11A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	5,566	5,684	8,043	8,591	5,603 ^b	6,693	5,360 ^a	6,015
Marijuana and Hashish	4,723	4,858	6,905	7,328	4,572 ^b	5,621	3,611 ^b	4,381
Cocaine	738	736	414	527	253	186	145	80
Crack	111	153	86	116	105	67	75 ^a	18
Heroin	33	87	68	120	79	105	110	123
Hallucinogens	450	450	466	419	248	180	170	124
LSD	90	114	101	80	13 ^a	57	24	36
PCP	3	8	15	5	*	*	14	*
Ecstasy	290	256	238	213	89	90	43	51
Inhalants	106	98	111	147	99	148	180	153
Nonmedical Use of Psychotherapeutics ^{3,4}	1,427	1,481	1,921	1,734	1,225	1,499	1,911	1,822
Pain Relievers	968	982	1,351	1,155	824	847	1,378	1,341
OxyContin [®]	113	129	139	111	59	71	181 ^a	46
Tranquilizers	442	465	574	581	279 ^a	472	410	357
Stimulants ⁴	421	432	400	387	256	401	288	375
Methamphetamine ⁴	170	119	138	89	106	161	181	200
Sedatives	40	72	36	85	73	67	102	105
ILLCIT DRUGS OTHER THAN MARIJUANA²	2,140	2,245	2,586	2,494	1,686	1,826	2,253	2,155

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.11B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	33.7	34.8	18.5	19.3	7.3 ^b	8.5	4.3 ^a	4.8
Marijuana and Hashish	28.6	29.7	15.9	16.4	6.0 ^b	7.1	2.9 ^b	3.5
Cocaine	4.5	4.5	1.0	1.2	0.3	0.2	0.1	0.1
Crack	0.7	0.9	0.2	0.3	0.1	0.1	0.1 ^a	0.0
Heroin	0.2	0.5	0.2	0.3	0.1	0.1	0.1	0.1
Hallucinogens	2.7	2.8	1.1	0.9	0.3	0.2	0.1	0.1
LSD	0.5	0.7	0.2	0.2	0.0 ^a	0.1	0.0	0.0
PCP	0.0	0.1	0.0	0.0	*	*	0.0	*
Ecstasy	1.8	1.6	0.5	0.5	0.1	0.1	0.0	0.0
Inhalants	0.6	0.6	0.3	0.3	0.1	0.2	0.1	0.1
Nonmedical Use of Psychotherapeutics ^{3,4}	8.6	9.1	4.4	3.9	1.6	1.9	1.5	1.5
Pain Relievers	5.8	6.0	3.1	2.6	1.1	1.1	1.1	1.1
OxyContin [®]	0.7	0.8	0.3	0.2	0.1	0.1	0.1 ^a	0.0
Tranquilizers	2.7	2.8	1.3	1.3	0.4 ^a	0.6	0.3	0.3
Stimulants ⁴	2.5	2.6	0.9	0.9	0.3	0.5	0.2	0.3
Methamphetamine ⁴	1.0	0.7	0.3	0.2	0.1	0.2	0.1	0.2
Sedatives	0.2	0.4	0.1	0.2	0.1	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	12.9	13.7	5.9	5.6	2.2	2.3	1.8	1.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.12A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	182	191	589	633	345	415	1,081	1,098
Marijuana and Hashish	169	177	545	579	311	362	736	712
Cocaine	18	21	7	17	2	2	16	*
Crack	0	*	0	5	1	*	*	*
Heroin	*	*	5	4	2	7	6	4
Hallucinogens	29	34	60	36	13	28	52	38
LSD	15	18	24	16	1	18	12	13
PCP	1	*	4	2	1	*	2	*
Ecstasy	11	11	21	13	1	6	14	8
Inhalants	10	4	20	29	6 ^b	27	85	89
Nonmedical Use of Psychotherapeutics ^{3,4}	46	58	110	135	63	93	329	369
Pain Relievers	37	38	83	92	44	55	262	282
OxyContin [®]	5	8	14	11	3	7	7	10
Tranquilizers	9 ^a	27	35	23	13	11	33	42
Stimulants ⁴	11	25	19	32	12 ^a	39	42	73
Methamphetamine ⁴	5	*	5	6	1 ^a	16	11	16
Sedatives	3	*	4	8	2	3	17	28
ILLCIT DRUGS OTHER THAN MARIJUANA²	75	84	158	188	77 ^a	133	448	479

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.12B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	62.3 ^a	74.1	46.6	50.5	26.1	30.5	4.9	5.0
Marijuana and Hashish	57.9	68.7	43.2	46.2	23.6	26.6	3.3	3.2
Cocaine	6.3	8.0	0.5	1.4	0.2	0.1	0.1	*
Crack	0.0	*	0.0	0.4	0.1	*	*	*
Heroin	*	*	0.4	0.3	0.1	0.5	0.0	0.0
Hallucinogens	10.0	13.2	4.8	2.9	1.0	2.1	0.2	0.2
LSD	5.0	7.0	1.9	1.3	0.1	1.3	0.1	0.1
PCP	0.2	*	0.3	0.1	0.0	*	0.0	*
Ecstasy	3.9	4.3	1.7	1.1	0.1	0.5	0.1	0.0
Inhalants	3.3	1.6	1.6	2.3	0.5 ^a	2.0	0.4	0.4
Nonmedical Use of Psychotherapeutics ^{3,4}	15.9	22.5	8.7	10.8	4.8	6.8	1.5	1.7
Pain Relievers	12.5	14.7	6.5	7.3	3.3	4.1	1.2	1.3
OxyContin [®]	1.8	3.1	1.1	0.9	0.2	0.5	0.0	0.0
Tranquilizers	3.1 ^a	10.6	2.8	1.8	1.0	0.8	0.1	0.2
Stimulants ⁴	3.7	9.8	1.5	2.5	0.9 ^a	2.9	0.2	0.3
Methamphetamine ⁴	1.7	*	0.4	0.5	0.1 ^a	1.2	0.1	0.1
Sedatives	0.9	*	0.3	0.6	0.1	0.2	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	25.5	32.8	12.5	14.9	5.8 ^a	9.8	2.0	2.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.13A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	1,917	1,926	3,093	2,998	1,283 ^a	1,497	1,196	1,256
Marijuana and Hashish	1,712	1,720	2,835	2,718	1,140 ^a	1,345	949	1,054
Cocaine	228	255	130	148	20	46	20	24
Crack	14	12	11	12	2	3	11	2
Heroin	14	14	16	8	22	14	40	47
Hallucinogens	249	196	250	223	47	55	71 ^a	28
LSD	40	50	47	45	9	17	9	7
PCP	2	3	*	1	*	*	2	*
Ecstasy	151	115	137	115	13	23	19	17
Inhalants	32	25	35	31	9	13	30	12
Nonmedical Use of Psychotherapeutics ^{3,4}	511	512	581	514	257	279	313	246
Pain Relievers	336	292	371	325	192	185	234	176
OxyContin [®]	49	34	38	39	14	17	6	12
Tranquilizers	146	172	166	134	51	57	71	53
Stimulants ⁴	212	167	139	140	59	62	52	37
Methamphetamine ⁴	57	39	33	19	18	18	13	10
Sedatives	11	10	13	18	3	15	8	13
ILLCIT DRUGS OTHER THAN MARIJUANA²	769	782	846	790	322	345	401	332

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.13B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	48.7	50.9	33.4	31.9	17.0 ^a	19.6	8.5	8.9
Marijuana and Hashish	43.5	45.5	30.7	28.9	15.1 ^a	17.6	6.8	7.5
Cocaine	5.8	6.7	1.4	1.6	0.3	0.6	0.1	0.2
Crack	0.4	0.3	0.1	0.1	0.0	0.0	0.1	0.0
Heroin	0.3	0.4	0.2	0.1	0.3	0.2	0.3	0.3
Hallucinogens	6.3	5.2	2.7	2.4	0.6	0.7	0.5 ^a	0.2
LSD	1.0	1.3	0.5	0.5	0.1	0.2	0.1	0.1
PCP	0.1	0.1	*	0.0	*	*	0.0	*
Ecstasy	3.8	3.0	1.5	1.2	0.2	0.3	0.1	0.1
Inhalants	0.8	0.7	0.4	0.3	0.1	0.2	0.2	0.1
Nonmedical Use of Psychotherapeutics ^{3,4}	13.0	13.5	6.3	5.5	3.4	3.7	2.2	1.7
Pain Relievers	8.5	7.7	4.0	3.5	2.5	2.4	1.7	1.2
OxyContin [®]	1.2	0.9	0.4	0.4	0.2	0.2	0.0	0.1
Tranquilizers	3.7	4.5	1.8	1.4	0.7	0.7	0.5	0.4
Stimulants ⁴	5.4	4.4	1.5	1.5	0.8	0.8	0.4	0.3
Methamphetamine ⁴	1.4	1.0	0.4	0.2	0.2	0.2	0.1	0.1
Sedatives	0.3	0.3	0.1	0.2	0.0	0.2	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	19.5	20.7	9.1	8.4	4.3	4.5	2.9	2.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.14A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	3,467	3,568	4,361	4,960	3,975 ^a	4,781	3,083	3,661
Marijuana and Hashish	2,841	2,961	3,525	4,032	3,120 ^b	3,914	1,926 ^b	2,615
Cocaine	491	460	277	362	231	138	109	56
Crack	98	138	75	99	102	64	64	16
Heroin	19	72	48	108	55	85	63	72
Hallucinogens	172	220	155	160	188	96	47	59
LSD	35	47	30	19	*	23	*	15
PCP	*	4	11	*	*	*	11	*
Ecstasy	128	130	80	84	75	61	10	25
Inhalants	65	69	56	87	83	108	65	52
Nonmedical Use of Psychotherapeutics ^{3,4}	870	911	1,230	1,086	906	1,127	1,268	1,207
Pain Relievers	595	652	898	737	588	607	883	884
OxyContin [®]	59	87	87	61	43	47	167 ^a	24
Tranquilizers	287	266	373	424	214 ^a	404	306	262
Stimulants ⁴	199	240	242	216	185	300	193	265
Methamphetamine ⁴	108	73	101	64	87	127	157	174
Sedatives	25	60	19	59	69	50	78	63
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,297	1,379	1,582	1,516	1,287	1,348	1,404	1,343

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.14B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	28.2	29.0	13.2	14.6	5.9 ^a	6.9	3.4 ^a	4.1
Marijuana and Hashish	23.1	24.1	10.7	11.9	4.6 ^a	5.6	2.2 ^b	2.9
Cocaine	4.0	3.7	0.8	1.1	0.3	0.2	0.1	0.1
Crack	0.8	1.1	0.2	0.3	0.2	0.1	0.1	0.0
Heroin	0.2 ^a	0.6	0.1	0.3	0.1	0.1	0.1	0.1
Hallucinogens	1.4	1.8	0.5	0.5	0.3	0.1	0.1	0.1
LSD	0.3	0.4	0.1	0.1	*	0.0	*	0.0
PCP	*	0.0	0.0	*	*	*	0.0	*
Ecstasy	1.0	1.1	0.2	0.2	0.1	0.1	0.0	0.0
Inhalants	0.5	0.6	0.2	0.3	0.1	0.2	0.1	0.1
Nonmedical Use of Psychotherapeutics ^{3,4}	7.1	7.4	3.7	3.2	1.3	1.6	1.4	1.4
Pain Relievers	4.8	5.3	2.7	2.2	0.9	0.9	1.0	1.0
OxyContin [®]	0.5	0.7	0.3	0.2	0.1	0.1	0.2 ^a	0.0
Tranquilizers	2.3	2.2	1.1	1.2	0.3 ^a	0.6	0.3	0.3
Stimulants ⁴	1.6	2.0	0.7	0.6	0.3	0.4	0.2	0.3
Methamphetamine ⁴	0.9	0.6	0.3	0.2	0.1	0.2	0.2	0.2
Sedatives	0.2	0.5	0.1	0.2	0.1	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	10.5	11.2	4.8	4.5	1.9	1.9	1.6	1.5

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.15A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Numbers in Thousands, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
ILLICIT DRUGS¹	19,213 ^b	20,968	3,417	3,512	15,796 ^b	17,457
Marijuana and Hashish	16,200 ^b	17,807	3,168	3,228	13,032 ^b	14,579
Cocaine (Including Crack)	1,404	1,450	150	174	1,254	1,275
Heroin	180 ^a	312	14	20	165 ^a	293
Hallucinogens	1,163	1,049	332	253	831	796
Inhalants	316	393	62	88	254	304
Pain Relievers ²	3,143	2,984	433	419	2,710	2,565
Tranquilizers ²	1,295	1,518	175	165	1,120	1,353
Stimulants ^{2,3}	1,078	1,220	174	197	904	1,023
Methamphetamine ³	414	369	42	56	372	313
Sedatives ²	149	225	21	19	127	206
ILLICIT DRUGS OTHER THAN MARIJUANA¹	6,411	6,565	966	986	5,446	5,579

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.15B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Percentages, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
ILLICIT DRUGS¹	14.0 ^a	15.0	39.5	40.5	12.3 ^a	13.3
Marijuana and Hashish	11.8 ^a	12.7	36.6	37.3	10.2 ^a	11.1
Cocaine (Including Crack)	1.0	1.0	1.7	2.0	1.0	1.0
Heroin	0.1 ^a	0.2	0.2	0.2	0.1 ^a	0.2
Hallucinogens	0.8	0.8	3.8	2.9	0.6	0.6
Inhalants	0.2	0.3	0.7	1.0	0.2	0.2
Pain Relievers ²	2.3	2.1	5.0	4.8	2.1	2.0
Tranquilizers ²	0.9	1.1	2.0	1.9	0.9	1.0
Stimulants ^{2,3}	0.8	0.9	2.0	2.3	0.7	0.8
Methamphetamine ³	0.3	0.3	0.5	0.6	0.3	0.2
Sedatives ²	0.1	0.2	0.2	0.2	0.1	0.2
ILLICIT DRUGS OTHER THAN MARIJUANA¹	4.7	4.7	11.2	11.4	4.2	4.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.16A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Numbers in Thousands, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
ILLCIT DRUGS¹	8,939 ^a	9,771	1,685	1,560	7,253 ^a	8,210
Marijuana and Hashish	8,082	8,780	1,643	1,490	6,440 ^a	7,290
Cocaine (Including Crack)	477	462	22	37	455	425
Heroin	85	92	1	7	84	86
Hallucinogens	149	157	14	22	135	135
Inhalants	14	20	3	8	11	12
Pain Relievers ²	547	600	27	37	519	563
Tranquilizers ²	104	193	14	14	90	179
Stimulants ²	60	77	21	10	40	67
Methamphetamine ³	131	136	11	18	121	119
Sedatives ²	26	37	4	3	22	34
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1,346	1,515	97	131	1,249	1,384

*Low precision; no estimate reported.

NOTE: Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.16B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Percentages, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
ILLCIT DRUGS¹	6.6	7.0	19.9	18.4	5.7 ^a	6.3
Marijuana and Hashish	5.9	6.3	19.5	17.7	5.0 ^a	5.6
Cocaine (Including Crack)	0.3	0.3	0.3	0.4	0.4	0.3
Heroin	0.1	0.1	0.0	0.1	0.1	0.1
Hallucinogens	0.1	0.1	0.2	0.3	0.1	0.1
Inhalants	0.0	0.0	0.0	0.1	0.0	0.0
Pain Relievers ²	0.4	0.4	0.3	0.4	0.4	0.4
Tranquilizers ²	0.1	0.1	0.2	0.2	0.1	0.1
Stimulants ²	0.0	0.1	0.2	0.1	0.0	0.1
Methamphetamine ³	0.1	0.1	0.1	0.2	0.1	0.1
Sedatives ²	0.0	0.0	0.0	0.0	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1.0	1.1	1.1	1.5	1.0	1.1

*Low precision; no estimate reported.

NOTE: Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.17A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	19,213 ^b	20,968	1,116	1,239	6,294	6,421	11,803 ^b	13,309
Marijuana and Hashish	16,200 ^b	17,807	1,026	1,118	5,687	5,782	9,486 ^b	10,907
Cocaine (Including Crack)	1,404	1,450	28	39	378	449	998	961
Heroin	180 ^a	312	6	12	52	35	122 ^a	265
Hallucinogens	1,163	1,049	102	98	546	474	515	477
Inhalants	316	393	36	60	76	68	204	264
Pain Relievers ²	3,143	2,984	163	185	899	802	2,081	1,996
Tranquilizers ²	1,295	1,518	57	61	363	363	874	1,093
Stimulants ^{2,3}	1,078	1,220	42 ^b	96	410	369	626	756
Methamphetamine ³	414	369	11	29	108	76	296	264
Sedatives ²	149	225	8	12	27	43	113	169
ILLICIT DRUGS OTHER THAN MARIJUANA¹	6,411	6,565	309 ^a	405	1,937	1,916	4,166	4,243

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.17B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	14.0 ^a	15.0	38.8 ^a	43.1	30.4	30.8	10.4 ^a	11.5
Marijuana and Hashish	11.8 ^a	12.7	35.7	38.9	27.4	27.8	8.4 ^a	9.4
Cocaine (Including Crack)	1.0	1.0	1.0	1.4	1.8	2.2	0.9	0.8
Heroin	0.1 ^a	0.2	0.2	0.4	0.2	0.2	0.1 ^a	0.2
Hallucinogens	0.8	0.8	3.6	3.4	2.6	2.3	0.5	0.4
Inhalants	0.2	0.3	1.2	2.1	0.4	0.3	0.2	0.2
Pain Relievers ²	2.3	2.1	5.7	6.5	4.3	3.9	1.8	1.7
Tranquilizers ²	0.9	1.1	2.0	2.1	1.8	1.7	0.8	0.9
Stimulants ^{2,3}	0.8	0.9	1.5 ^b	3.3	2.0	1.8	0.6	0.7
Methamphetamine ³	0.3	0.3	0.4	1.0	0.5	0.4	0.3	0.2
Sedatives ²	0.1	0.2	0.3	0.4	0.1	0.2	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA¹	4.7	4.7	10.7 ^a	14.1	9.3	9.2	3.7	3.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.18A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLCIT DRUGS¹	8,939 ^a	9,771	499	506	3,209	3,149	5,231 ^a	6,115
Marijuana and Hashish	8,082	8,780	486	470	3,026	3,003	4,570 ^a	5,308
Cocaine (Including Crack)	477	462	3	12	109	114	365	336
Heroin	85	92	*	7	25 ^a	3	60	82
Hallucinogens	149	157	4	15	40	51	105	91
Inhalants	14	20	*	5	5	5	8	10
Pain Relievers ²	547	600	14	21	110	97	423	482
Tranquilizers ²	104	193	7	9	29	22	67 ^a	163
Stimulants ²	60	77	4	9	26	14	29	54
Methamphetamine ³	131	136	2	7	24	30	105	99
Sedatives ²	26	37	*	3	8	10	18	24
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1,346	1,515	32 ^a	68	327	306	987	1,141

*Low precision; no estimate reported.

NOTE: Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.18B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLCIT DRUGS¹	6.6	7.0	18.0	18.3	15.6	15.2	4.6 ^a	5.3
Marijuana and Hashish	5.9	6.3	17.6	17.2	14.8	14.6	4.0	4.6
Cocaine (Including Crack)	0.3	0.3	0.1	0.4	0.5	0.6	0.3	0.3
Heroin	0.1	0.1	*	0.2	0.1 ^a	0.0	0.1	0.1
Hallucinogens	0.1	0.1	0.1	0.5	0.2	0.2	0.1	0.1
Inhalants	0.0	0.0	*	0.2	0.0	0.0	0.0	0.0
Pain Relievers ²	0.4	0.4	0.5	0.7	0.5	0.5	0.4	0.4
Tranquilizers ²	0.1	0.1	0.3	0.3	0.1	0.1	0.1 ^a	0.1
Stimulants ²	0.0	0.1	0.1	0.3	0.1	0.1	0.0	0.0
Methamphetamine ³	0.1	0.1	0.1	0.3	0.1	0.1	0.1	0.1
Sedatives ²	0.0	0.0	*	0.1	0.0	0.0	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1.0	1.1	1.1 ^a	2.4	1.6	1.5	0.9	1.0

*Low precision; no estimate reported.

NOTE: Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.19A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	10,898	11,064	2,571	2,837	8,315 ^b	9,904	2,789	3,178
Marijuana and Hashish	9,432	9,720	1,992	2,343	6,768 ^b	8,087	1,619 ^a	2,038
Cocaine	1,074	1,076	84	61	330	373	61	20
Crack	255	272	32	18	47	64	43	*
Heroin	177	265	71	118	*	47	38	*
Hallucinogens	750	699	73	86	414	350	97	39
LSD	129	165	17	32	75	86	7	*
PCP	17	10	2	*	*	*	12	*
Ecstasy	398	353	21	42	218	205	22	8
Inhalants	182	168	42	27	134	224	137	126
Nonmedical Use of Psychotherapeutics ^{2,3}	2,769	2,511	780	676	1,804 ^a	2,204	1,131	1,147
Pain Relievers	1,957	1,651	553	486	1,186	1,332	826	856
OxyContin [®]	245	183	99 ^a	20	66	128	82	25
Tranquilizers	835	916	222	190	460	602	188	168
Stimulants ³	762	693	172	178	316 ^b	527	116	197
Methamphetamine ³	352	233	118	113	62	135	62	88
Sedatives	67	156	15	34	81	68	87	71
ILLCIT DRUGS OTHER THAN MARIJUANA¹	3,990	3,683	927	859	2,421 ^a	2,882	1,326	1,296

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.19B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Percentages, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS¹	30.0	30.4	13.2	15.0	8.3 ^b	9.6	2.6	3.0
Marijuana and Hashish	26.0	26.7	10.3 ^a	12.4	6.7 ^b	7.8	1.5 ^a	1.9
Cocaine	3.0	3.0	0.4	0.3	0.3	0.4	0.1	0.0
Crack	0.7	0.7	0.2	0.1	0.0	0.1	0.0	*
Heroin	0.5	0.7	0.4	0.6	*	0.0	0.0	*
Hallucinogens	2.1	1.9	0.4	0.5	0.4	0.3	0.1	0.0
LSD	0.4	0.5	0.1	0.2	0.1	0.1	0.0	*
PCP	0.0	0.0	0.0	*	*	*	0.0	*
Ecstasy	1.1	1.0	0.1	0.2	0.2	0.2	0.0	0.0
Inhalants	0.5	0.5	0.2	0.1	0.1	0.2	0.1	0.1
Nonmedical Use of Psychotherapeutics ^{2,3}	7.6	6.9	4.0	3.6	1.8	2.1	1.1	1.1
Pain Relievers	5.4	4.5	2.8	2.6	1.2	1.3	0.8	0.8
OxyContin [®]	0.7	0.5	0.5 ^a	0.1	0.1	0.1	0.1	0.0
Tranquilizers	2.3	2.5	1.1	1.0	0.5	0.6	0.2	0.2
Stimulants ³	2.1	1.9	0.9	0.9	0.3 ^b	0.5	0.1	0.2
Methamphetamine ³	1.0	0.6	0.6	0.6	0.1	0.1	0.1	0.1
Sedatives	0.2	0.4	0.1	0.2	0.1	0.1	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA¹	11.0	10.1	4.8	4.5	2.4	2.8	1.3	1.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.20A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	504	483	248	200	612 ^a	756	833	898
Marijuana and Hashish	467	440	225	188	559 ^a	678	511	524
Cocaine	20	32	5	*	7	8	11	*
Crack	1	8	*	*	1	*	*	*
Heroin	5	12	4	1	1	*	2	3
Hallucinogens	76	54	25	18	27	44	27	20
LSD	34	29	7	9	6	23	4	4
PCP	4	3	1	*	1	*	*	*
Ecstasy	21	19	8	7	13	12	6	1
Inhalants	18	29	8	5	17	31	77	84
Nonmedical Use of Psychotherapeutics ^{2,3}	135	134	36	27	84 ^b	152	294	342
Pain Relievers	97	89	24	15	67	97	238	267
OxyContin [®]	14	14	0	0	8	12	7	9
Tranquilizers	49	41	2 ^a	13	9	20	31	29
Stimulants ³	27	40	14	10	15 ^b	55	29 ^a	63
Methamphetamine ³	6	15	5	*	5	14	6	12
Sedatives	6	7	1	*	3	5	15	28
ILLCIT DRUGS OTHER THAN MARIJUANA¹	197	203	66	46	112 ^b	202	382	433

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.20B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Percentages, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS¹	64.5	68.5	40.4	39.4	29.2 ^a	34.9	3.9	4.2
Marijuana and Hashish	59.7	62.4	36.6	37.1	26.7 ^a	31.3	2.4	2.4
Cocaine	2.6	4.5	0.9	*	0.3	0.4	0.0	*
Crack	0.1	1.1	*	*	0.1	*	*	*
Heroin	0.6	1.6	0.7	0.2	0.1	*	0.0	0.0
Hallucinogens	9.7	7.6	4.1	3.6	1.3	2.0	0.1	0.1
LSD	4.4	4.2	1.2	1.8	0.3	1.0	0.0	0.0
PCP	0.5	0.5	0.2	*	0.1	*	*	*
Ecstasy	2.7	2.7	1.4	1.4	0.6	0.6	0.0	0.0
Inhalants	2.4	4.1	1.3	1.0	0.8	1.4	0.4	0.4
Nonmedical Use of Psychotherapeutics ^{2,3}	17.3	19.0	5.8	5.4	4.0 ^b	7.0	1.4	1.6
Pain Relievers	12.4	12.6	3.8	3.0	3.2	4.5	1.1	1.2
OxyContin [®]	1.8	2.0	0.1	0.0	0.4	0.5	0.0	0.0
Tranquilizers	6.3	5.8	0.3 ^a	2.6	0.4	0.9	0.1	0.1
Stimulants ³	3.4	5.7	2.2	2.1	0.7 ^b	2.5	0.1 ^a	0.3
Methamphetamine ³	0.8	2.1	0.8	*	0.2	0.6	0.0	0.1
Sedatives	0.7	1.0	0.2	*	0.1	0.2	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA¹	25.1	28.8	10.7	9.0	5.4 ^b	9.3	1.8	2.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.21A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	3,871	3,785	572	592	2,423	2,636	625	664
Marijuana and Hashish	3,529	3,435	499	508	2,159	2,348	449	546
Cocaine	314	335	17	23	64 ^a	114	3	1
Crack	26	12	11	2	*	15	*	*
Heroin	50	34	39	47	2	2	1	*
Hallucinogens	389	365	39	17	157	109	32	11
LSD	70	89	6	7	26	22	3	*
PCP	2	3	*	*	*	2	2	*
Ecstasy	224	186	13	10	77	66	6	7
Inhalants	53	47	16	0	23	21	14	12
Nonmedical Use of Psychotherapeutics ^{2,3}	934	820	131	122	415	485	182 ^a	124
Pain Relievers	649	530	91	88	249	273	142 ^a	88
OxyContin [®]	85	75	6	10	15	14	*	2
Tranquilizers	267	256	50	31	96	107	21	22
Stimulants ³	286	225	27	21	124	144	25	16
Methamphetamine ³	94	55	10	7	14	21	2	3
Sedatives	14	20	*	5	13	23	8	9
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1,367	1,254	183	182	570	662	218	150

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.21B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Percentages, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS¹	46.8	49.4	24.0	26.3	19.4	20.0	5.4	5.6
Marijuana and Hashish	42.7	44.9	20.9	22.6	17.3	17.8	3.9	4.6
Cocaine	3.8	4.4	0.7	1.0	0.5	0.9	0.0	0.0
Crack	0.3	0.2	0.4	0.1	*	0.1	*	*
Heroin	0.6	0.4	1.6	2.1	0.0	0.0	0.0	*
Hallucinogens	4.7	4.8	1.6	0.7	1.3	0.8	0.3	0.1
LSD	0.8	1.2	0.2	0.3	0.2	0.2	0.0	*
PCP	0.0	0.0	*	*	*	0.0	0.0	*
Ecstasy	2.7	2.4	0.5	0.4	0.6	0.5	0.1	0.1
Inhalants	0.6	0.6	0.7	0.0	0.2	0.2	0.1	0.1
Nonmedical Use of								
Psychotherapeutics ^{2,3}	11.3	10.7	5.5	5.4	3.3	3.7	1.6 ^a	1.0
Pain Relievers	7.8	6.9	3.8	3.9	2.0	2.1	1.2 ^a	0.7
OxyContin [®]	1.0	1.0	0.3	0.5	0.1	0.1	*	0.0
Tranquilizers	3.2	3.3	2.1	1.4	0.8	0.8	0.2	0.2
Stimulants ³	3.5	2.9	1.1	0.9	1.0	1.1	0.2	0.1
Methamphetamine ³	1.1	0.7	0.4	0.3	0.1	0.2	0.0	0.0
Sedatives	0.2	0.3	*	0.2	0.1	0.2	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA¹	16.5	16.4	7.7	8.1	4.6	5.0	1.9 ^a	1.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.22A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS¹	6,523	6,797	1,751	2,046	5,280 ^b	6,512	1,331	1,615
Marijuana and Hashish	5,436	5,845	1,268	1,646	4,050 ^b	5,062	658 ^a	968
Cocaine	739	710	62	38	260	251	47	19
Crack	229	252	21	16	46	49	42	*
Heroin	122	220	28	70	*	45	35	*
Hallucinogens	284	280	9 ^a	50	230	197	38	8
LSD	25	46	4	15	43	42	*	*
PCP	11	3	1	*	*	*	10	*
Ecstasy	153	148	*	25	129	127	10	*
Inhalants	110	92	18	22	94	172	47	31
Nonmedical Use of								
Psychotherapeutics ^{2,3}	1,700	1,557	614	526	1,306	1,566	655	682
Pain Relievers	1,211	1,033	438	383	870	963	445	501
OxyContin [®]	145	93	92 ^a	10	43	102	75	14
Tranquilizers	519	618	170	145	355	475	136	117
Stimulants ³	450	428	131	146	176 ^a	328	62	118
Methamphetamine ³	253	163	103	102	43	101	54	73
Sedatives	48	129	14	29	66	40	65	34
ILLICIT DRUGS OTHER THAN MARIJUANA¹	2,427	2,226	678	631	1,739	2,018	727	713

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.22B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Percentages, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS¹	23.9	24.3	10.7	12.7	6.1 ^b	7.4	1.8	2.2
Marijuana and Hashish	19.9	20.9	7.7 ^a	10.2	4.7 ^b	5.8	0.9 ^a	1.3
Cocaine	2.7	2.5	0.4	0.2	0.3	0.3	0.1	0.0
Crack	0.8	0.9	0.1	0.1	0.1	0.1	0.1	*
Heroin	0.4	0.8	0.2	0.4	*	0.1	0.0	*
Hallucinogens	1.0	1.0	0.1 ^a	0.3	0.3	0.2	0.1	0.0
LSD	0.1	0.2	0.0	0.1	0.1	0.0	*	*
PCP	0.0	0.0	0.0	*	*	*	0.0	*
Ecstasy	0.6	0.5	*	0.2	0.1	0.1	0.0	*
Inhalants	0.4	0.3	0.1	0.1	0.1	0.2	0.1	0.0
Nonmedical Use of								
Psychotherapeutics ^{2,3}	6.2	5.6	3.7	3.3	1.5	1.8	0.9	0.9
Pain Relievers	4.4	3.7	2.7	2.4	1.0	1.1	0.6	0.7
OxyContin [®]	0.5	0.3	0.6 ^a	0.1	0.0	0.1	0.1	0.0
Tranquilizers	1.9	2.2	1.0	0.9	0.4	0.5	0.2	0.2
Stimulants ³	1.6	1.5	0.8	0.9	0.2 ^a	0.4	0.1	0.2
Methamphetamine ³	0.9	0.6	0.6	0.6	0.1	0.1	0.1	0.1
Sedatives	0.2	0.5	0.1	0.2	0.1	0.0	0.1	0.0
ILLICIT DRUGS OTHER THAN MARIJUANA¹	8.9	8.0	4.1	3.9	2.0	2.3	1.0	1.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.23A Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Numbers in Thousands, 2013 and 2014

Cigarette Use/Level of Alcohol Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	24,573 ^b	26,983	2,197	2,338	7,490	7,676	14,886 ^b	16,969
Heavy Alcohol Use ¹	5,566	5,684	182	191	1,917	1,926	3,467	3,568
Binge Alcohol Use But Not Heavy Use ¹	8,043	8,591	589	633	3,093	2,998	4,361	4,960
Alcohol Use But Not Binge Use ¹	5,603 ^b	6,693	345	415	1,283 ^a	1,497	3,975 ^a	4,781
No Alcohol Use	5,360 ^a	6,015	1,081	1,098	1,196	1,256	3,083	3,661
PAST MONTH CIGARETTE USE	13,470	13,901	752	683	4,443	4,376	8,274	8,842
Heavy Alcohol Use ¹	3,759	3,660	118	*	1,399	1,303	2,242	2,256
Binge Alcohol Use But Not Heavy Use ¹	4,672	4,639	278	271	1,836	1,737	2,557	2,630
Alcohol Use But Not Binge Use ¹	2,468	2,765	108	110	636	745	1,724	1,910
No Alcohol Use	2,571	2,837	248	200	572	592	1,751	2,046
NO PAST MONTH CIGARETTE USE	11,104 ^b	13,082	1,445 ^a	1,655	3,047	3,300	6,611 ^b	8,127
Heavy Alcohol Use ¹	1,808	2,024	*	*	518	623	1,225	1,311
Binge Alcohol Use But Not Heavy Use ¹	3,372 ^a	3,952	311	362	1,257	1,260	1,804 ^b	2,330
Alcohol Use But Not Binge Use ¹	3,136 ^b	3,928	237	305	647	752	2,251 ^a	2,871
No Alcohol Use	2,789	3,178	833	898	625	664	1,331	1,615

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.23B Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Percentages, 2013 and 2014

Cigarette Use/Level of Alcohol Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	9.4 ^b	10.2	8.8	9.4	21.5	22.0	7.3 ^b	8.3
Heavy Alcohol Use ¹	33.7	34.8	62.3 ^a	74.1	48.7	50.9	28.2	29.0
Binge Alcohol Use But Not Heavy Use ¹	18.5	19.3	46.6	50.5	33.4	31.9	13.2	14.6
Alcohol Use But Not Binge Use ¹	7.3 ^b	8.5	26.1	30.5	17.0 ^a	19.6	5.9 ^a	6.9
No Alcohol Use	4.3 ^a	4.8	4.9	5.0	8.5	8.9	3.4 ^a	4.1
PAST MONTH CIGARETTE USE	24.1	25.2	53.9	56.3	41.7 ^a	44.2	18.9	20.0
Heavy Alcohol Use ¹	42.8	43.1	70.6	*	58.4	60.4	36.1	36.4
Binge Alcohol Use But Not Heavy Use ¹	30.9	30.1	66.1	70.3	45.9	46.0	23.9	23.4
Alcohol Use But Not Binge Use ¹	19.8	22.2	55.6	58.7	34.0 ^b	43.3	16.6	18.1
No Alcohol Use	13.2	15.0	40.4	39.4	24.0	26.3	10.7	12.7
NO PAST MONTH CIGARETTE USE	5.4 ^b	6.2	6.1 ^a	7.0	12.6	13.2	4.2 ^b	5.0
Heavy Alcohol Use ¹	23.3	25.8	*	*	33.6	38.3	20.1	21.5
Binge Alcohol Use But Not Heavy Use ¹	11.9 ^a	13.5	36.9	41.6	24.0	22.4	8.1 ^a	10.3
Alcohol Use But Not Binge Use ¹	4.9 ^a	5.9	21.1	26.0	11.4	12.7	3.9 ^a	4.8
No Alcohol Use	2.6	3.0	3.9	4.2	5.4	5.6	1.8	2.2

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.24A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	11,101	11,659
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	3,383	3,136	5,430	5,526
Cigars	6,846 ^a	5,997	5,592	5,967
Pipe Tobacco	1,360	1,197	910	1,049
ALCOHOL	36,345	36,340	100,523	103,336
Binge Alcohol Use ²	23,912	23,901	36,152	37,040
Heavy Alcohol Use ²	8,782	8,485	7,759	7,857

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.24B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	5.4	5.6
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	6.1	5.7	2.6	2.6
Cigars	12.3 ^a	10.9	2.7	2.8
Pipe Tobacco	2.4	2.2	0.4	0.5
ALCOHOL	65.2	65.8	48.7	49.2
Binge Alcohol Use ²	42.9	43.3	17.5	17.6
Heavy Alcohol Use ²	15.7	15.4	3.8	3.7

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.25A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	554	529
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	211	222	276	268
Cigars	311	279	265	242
Pipe Tobacco	90	107	70	72
ALCOHOL	782	705	2,095	2,167
Binge Alcohol Use ²	588	517	968	995
Heavy Alcohol Use ²	167	131	125	126

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.25B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	2.4	2.2
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	15.1	18.3	1.2	1.1
Cigars	22.3	23.0	1.1	1.0
Pipe Tobacco	6.4	8.8	0.3	0.3
ALCOHOL	55.9	58.1	8.9	9.2
Binge Alcohol Use ²	42.1	42.6	4.1	4.2
Heavy Alcohol Use ²	12.0	10.8	0.5	0.5

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.26A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	2,226	2,339
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,209	1,091	810	873
Cigars	2,138	1,925	1,335	1,467
Pipe Tobacco	419	406	339	259
ALCOHOL	8,270 ^a	7,656	12,469	13,159
Binge Alcohol Use ²	6,399 ^a	5,936	6,788	7,249
Heavy Alcohol Use ²	2,397	2,157	1,542	1,625

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.26B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	9.2	9.3
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	11.3	11.0	3.4	3.5
Cigars	20.1	19.4	5.5	5.9
Pipe Tobacco	3.9	4.1	1.4	1.0
ALCOHOL	77.6	77.3	51.7	52.6
Binge Alcohol Use ²	60.0	59.9	28.1	29.0
Heavy Alcohol Use ²	22.5	21.8	6.4	6.5

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.27A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	8,322	8,790
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,963	1,823	4,344	4,386
Cigars	4,397	3,793	3,992	4,258
Pipe Tobacco	851	684	502	718
ALCOHOL	27,293	27,979	85,959	88,011
Binge Alcohol Use ²	16,925	17,448	28,396	28,797
Heavy Alcohol Use ²	6,218	6,197	6,092	6,106

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.27B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	5.2	5.5
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	4.5	4.1	2.7	2.7
Cigars	10.1 ^a	8.6	2.5	2.6
Pipe Tobacco	1.9	1.6	0.3	0.4
ALCOHOL	62.4	63.4	54.1	54.6
Binge Alcohol Use ²	38.7	39.5	17.9	17.9
Heavy Alcohol Use ²	14.2	14.0	3.8	3.8

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.28A Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	10,385	10,208	18,669	18,958	15,601	15,623	22,224	22,111
Cigarettes	8,782	8,485	15,130	15,416	12,432	12,439	19,433	18,900
Smokeless Tobacco	2,005	1,971	2,729	2,521	1,540	1,496	2,540	2,675
Cigars	2,552	2,219	4,669	4,270	2,973	3,157	2,244	2,318
Pipe Tobacco	389	292	728	647	528	492	626	815

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.28B Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	62.8	62.5	42.9	42.5	20.3	19.8	17.7	17.6
Cigarettes	53.1	51.9	34.8	34.6	16.2	15.8	15.5	15.1
Smokeless Tobacco	12.1	12.1	6.3	5.7	2.0	1.9	2.0	2.1
Cigars	15.4	13.6	10.7	9.6	3.9	4.0	1.8	1.8
Pipe Tobacco	2.4	1.8	1.7	1.5	0.7	0.6	0.5	0.6

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.29A Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	209	174	580	532	284	267	879	770
Cigarettes	167	131	421	386	194	188	615 ^a	508
Smokeless Tobacco	78	72	142	122	61	67	206	229
Cigars	95	76	200	194	74	67	207	185
Pipe Tobacco	19	15	57	58	18	29	66	76

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.29B Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	71.3	67.6	45.9	42.4	21.5	19.6	4.0	3.5
Cigarettes	57.1	51.0	33.3	30.7	14.7	13.8	2.8 ^a	2.3
Smokeless Tobacco	26.8	28.0	11.2	9.7	4.6	4.9	0.9	1.0
Cigars	32.6	29.3	15.8	15.5	5.6	4.9	0.9	0.8
Pipe Tobacco	6.4	6.0	4.5	4.6	1.4	2.2	0.3	0.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.30A Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	2,805	2,584	4,857	4,644	2,356	2,201	2,864	2,815
Cigarettes	2,397	2,157	4,002	3,778	1,871	1,720	2,387	2,248
Smokeless Tobacco	677	568	733	707	249	235	359	454
Cigars	929	803	1,370	1,363	602	640	572	586
Pipe Tobacco	151	124	306	288	160	132	141	122

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.30B Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	71.2	68.3	52.5 ^a	49.4	31.2	28.8	20.4	19.9
Cigarettes	60.9	57.0	43.3 ^a	40.2	24.8	22.5	17.0	15.9
Smokeless Tobacco	17.2	15.0	7.9	7.5	3.3	3.1	2.6	3.2
Cigars	23.6	21.2	14.8	14.5	8.0	8.4	4.1	4.1
Pipe Tobacco	3.8	3.3	3.3	3.1	2.1	1.7	1.0	0.9

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.31A Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	7,372	7,450	13,232	13,782	12,961	13,155	18,481	18,526
Cigarettes	6,218	6,197	10,708	11,252	10,368	10,531	16,431	16,144
Smokeless Tobacco	1,250	1,331	1,854	1,692	1,230	1,194	1,974	1,992
Cigars	1,528	1,341	3,099	2,713	2,297	2,450	1,465	1,547
Pipe Tobacco	220	153	365	301	349	331	419	616

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.31B Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	59.9	60.6	40.1	40.6	19.1	18.9	20.7	20.7
Cigarettes	50.5	50.4	32.4	33.1	15.3	15.1	18.4	18.1
Smokeless Tobacco	10.2	10.8	5.6	5.0	1.8	1.7	2.2	2.2
Cigars	12.4	10.9	9.4	8.0	3.4	3.5	1.6	1.7
Pipe Tobacco	1.8	1.2	1.1	0.9	0.5	0.5	0.5	0.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.32A Marijuana Use in the Prior Year and Past Year Marijuana Use Status among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Marijuana Use in Prior Year		Marijuana Use in Prior Year and Past Year		Marijuana Use in Prior Year But No Use in Past Year		Percentage Who Did Not Use Marijuana in the Past Year among Prior Year Marijuana Users	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	31,972 ^b	34,344	24,767 ^a	26,505	7,205	7,839	22.5	22.8
AGE								
12-17	2,083	1,993	1,721	1,642	361	351	17.3	17.6
12-13	77	68	56	61	20 ^a	6	*	*
14-15	553 ^a	460	444	381	109	79	19.8	17.1
16-17	1,452	1,466	1,221	1,200	231	266	15.9	18.2
18-25	10,310	10,657	8,186	8,394	2,123	2,264	20.6	21.2
18-20	3,778	3,618	3,101	3,022	677	596	17.9	16.5
21-25	6,531 ^a	7,039	5,085	5,372	1,446 ^a	1,668	22.1	23.7
26 or Older	19,579 ^b	21,694	14,859 ^a	16,469	4,720	5,225	24.1	24.1
26-34	7,822	7,847	6,196	6,029	1,626	1,818	20.8	23.2
35-49	6,035 ^b	7,047	4,469 ^b	5,325	1,566	1,722	25.9	24.4
50 or Older	5,723 ^a	6,800	4,194 ^a	5,115	1,528	1,685	26.7	24.8
GENDER								
Male	19,273 ^b	20,969	15,321 ^b	16,789	3,952	4,180	20.5	19.9
Female	12,699	13,375	9,446	9,716	3,253	3,660	25.6	27.4
GENDER/AGE								
Male 12-17	1,139	1,098	965	916	174	182	15.3	16.5
Male 18-25	6,073	5,988	4,985	4,901	1,088	1,087	17.9	18.2
Male 26 or Older	12,060 ^b	13,883	9,370 ^b	10,972	2,690	2,911	22.3	21.0
Female 12-17	943	895	756	726	187	170	19.8	18.9
Female 18-25	4,237 ^b	4,669	3,201 ^a	3,493	1,036	1,176	24.4	25.2
Female 26 or Older	7,519	7,811	5,489	5,497	2,030	2,314	27.0	29.6

*Low precision; no estimate reported.

NOTE: Prior Year Marijuana Use is defined as using marijuana/hashish 12 to 23 months prior to the interview date.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.32B Marijuana Use in the Prior Year and Past Year Marijuana Use Status among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Marijuana Use in Prior Year		Marijuana Use in Prior Year and Past Year		Marijuana Use in Prior Year But No Use in Past Year		Percentage Who Did Not Use Marijuana in the Past Year among Prior Year Marijuana Users	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	12.2 ^a	13.0	9.4 ^a	10.0	2.7	3.0	22.5	22.8
AGE								
12-17	8.4	8.0	6.9	6.6	1.5	1.4	17.3	17.6
12-13	1.0	0.9	0.7	0.8	0.3 ^a	0.1	*	*
14-15	6.5 ^a	5.4	5.2	4.5	1.3	0.9	19.8	17.1
16-17	17.3	17.2	14.5	14.1	2.8	3.1	15.9	18.2
18-25	29.6	30.5	23.5	24.0	6.1	6.5	20.6	21.2
18-20	28.6	27.6	23.5	23.1	5.1	4.5	17.9	16.5
21-25	30.2 ^a	32.2	23.6	24.6	6.7 ^a	7.6	22.1	23.7
26 or Older	9.7 ^a	10.6	7.3 ^a	8.0	2.3	2.5	24.1	24.1
26-34	20.9	20.7	16.6	15.9	4.4	4.8	20.8	23.2
35-49	10.0 ^b	11.7	7.4 ^b	8.8	2.6	2.9	25.9	24.4
50 or Older	5.5	6.3	4.0	4.8	1.5	1.6	26.7	24.8
GENDER								
Male	15.2 ^a	16.3	12.1 ^a	13.1	3.1	3.3	20.5	19.9
Female	9.4	9.8	7.0	7.1	2.4	2.7	25.6	27.4
GENDER/AGE								
Male 12-17	9.0	8.7	7.6	7.2	1.4	1.4	15.3	16.5
Male 18-25	34.8	34.2	28.6	28.0	6.2	6.2	17.9	18.2
Male 26 or Older	12.4 ^b	14.1	9.7 ^b	11.2	2.8	3.0	22.3	21.0
Female 12-17	7.7	7.3	6.2	6.0	1.5	1.4	19.8	18.9
Female 18-25	24.5 ^b	26.8	18.5	20.1	6.0	6.8	24.4	25.2
Female 26 or Older	7.1	7.3	5.2	5.1	1.9	2.2	27.0	29.6

*Low precision; no estimate reported.

NOTE: Prior Year Marijuana Use is defined as using marijuana/hashish 12 to 23 months prior to the interview date.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.33A Cigarette Use and Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Detailed Age Category:
Numbers in Thousands and Percentages, 2013 and 2014**

Age Category	Cigarette Use in Past Month		Nicotine Dependence in Past Month		Percentage with Nicotine Dependence among Past Month Cigarette Smokers	
	2013	2014	2013	2014	2013	2014
TOTAL	55,778	55,241	31,033	31,011	55.6	56.1
12	16	12	6	3	*	*
13	55	45	9	11	*	*
14	115	86	40 ^a	18	*	*
15	252	202	74	60	29.5	29.4
16	372	374	124	129	33.3	34.4
17	587 ^a	493	223	198	38.0	40.3
18	992	892	363	316	36.6	35.4
19	1,188 ^a	993	474 ^a	365	39.9	36.8
20	1,391	1,261	557	492	40.0	39.0
21	1,336	1,285	520	523	38.9	40.7
22	1,468	1,442	654	620	44.5	43.0
23	1,489	1,358	636	650	42.7	47.9
24	1,463	1,305	649	565	44.4	43.3
25	1,329	1,369	654	651	49.2	47.5
26-29	5,623	5,089	2,609	2,473	46.4	48.6
30-34	6,689 ^a	6,039	3,483	3,144	52.1	52.1
35-39	4,876	5,244	2,631	2,713	54.0	51.7
40-44	5,018	4,797	3,199	2,869	63.8	59.8
45-49	4,882	4,813	3,005	3,138	61.6	65.2
50-54	5,323	5,764	3,766	3,873	70.7	67.2
55-59	4,036 ^a	4,784	2,782	3,236	68.9	67.6
60-64	2,915	3,272	1,836	2,081	63.0	63.6
65 or Older	4,363	4,321	2,737	2,885	62.7	66.8

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.33B Cigarette Use and Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Cigarette Use in Past Month		Nicotine Dependence in Past Month		Percentage with Nicotine Dependence among Past Month Cigarette Smokers	
	2013	2014	2013	2014	2013	2014
TOTAL	21.3	20.8	11.8	11.7	55.6	56.1
12	0.4	0.3	0.2	0.1	*	*
13	1.3	1.1	0.2	0.3	*	*
14	2.6	2.1	0.9 ^a	0.4	*	*
15	6.1 ^a	4.7	1.8	1.4	29.5	29.4
16	8.7	8.6	2.9	3.0	33.3	34.4
17	14.2 ^a	11.8	5.4	4.8	38.0	40.3
18	21.0	19.4	7.7	6.9	36.6	35.4
19	28.5 ^b	24.0	11.4 ^a	8.8	39.9	36.8
20	32.4	28.9	13.0	11.3	40.0	39.0
21	31.5	29.6	12.3	12.0	38.9	40.7
22	32.8	32.0	14.6	13.7	44.5	43.0
23	34.0	30.9	14.5	14.8	42.7	47.9
24	33.3	30.2	14.8	13.1	44.4	43.3
25	32.3	32.2	15.9	15.3	49.2	47.5
26-29	32.7	29.9	15.2	14.5	46.4	48.6
30-34	33.2 ^b	29.0	17.3 ^a	15.1	52.1	52.1
35-39	25.6	26.8	13.8	13.9	54.0	51.7
40-44	23.4	23.8	14.9	14.2	63.8	59.8
45-49	24.4	23.4	15.0	15.3	61.6	65.2
50-54	23.1	24.3	16.3	16.3	70.7	67.2
55-59	19.6 ^a	22.9	13.5	15.5	68.9	67.6
60-64	16.4	18.6	10.3	11.9	63.0	63.6
65 or Older	10.0	9.6	6.3	6.4	62.7	66.8

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.34A Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	31,033	31,011	477	418	4,507 ^a	4,182	26,049	26,411
GENDER								
Male	16,597	16,411	270	220	2,650	2,441	13,678	13,751
Female	14,436	14,599	207	199	1,858	1,741	12,371	12,660
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	28,503	28,542	420	364	4,085 ^a	3,760	23,998	24,418
White	23,253	22,945	357	312	3,233 ^a	2,957	19,663	19,675
Black or African American	3,879	4,130	37	34	569	553	3,273	3,544
American Indian or Alaska Native	272	214	5	3	43	28	224	184
Native Hawaiian or Other Pacific Islander	*	136	3	*	18	*	*	111
Asian	302 ^a	523	3	2	107	111	192 ^a	409
Two or More Races	697	595	15	13	115	87	566	495
Hispanic or Latino	2,530	2,468	57	54	423	422	2,050	1,993
FAMILY INCOME								
Less Than \$20,000	9,561	9,863	131	124	1,765 ^b	1,491	7,665	8,249
\$20,000 - \$49,999	11,405	11,437	185	153	1,508	1,561	9,712	9,722
\$50,000 - \$74,999	4,884	4,267	71	73	537	512	4,275	3,681
\$75,000 or More	5,183	5,444	89	67	697	617	4,397	4,759

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.34B Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	11.8	11.7	1.9	1.7	13.0 ^a	12.0	12.9	12.9
GENDER								
Male	13.1	12.8	2.1	1.7	15.2	13.9	14.1	14.0
Female	10.7	10.7	1.7	1.6	10.7	10.0	11.7	11.8
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	12.9	12.8	2.2	1.9	14.8 ^a	13.6	13.8	13.9
White	13.7	13.5	2.6	2.3	16.6 ^a	15.3	14.4	14.3
Black or African American	12.4	13.0	1.1	1.0	11.4	11.0	14.4	15.3
American Indian or Alaska Native	19.4	14.9	2.9	1.8	18.8	13.6	22.1	17.1
Native Hawaiian or Other Pacific Islander	*	13.3	2.9	*	8.9	*	*	15.0
Asian	2.3 ^a	3.8	0.2	0.1	5.7	5.5	1.9 ^a	3.9
Two or More Races	16.2	13.5	2.0	1.7	15.7	11.1	20.2	17.3
Hispanic or Latino	6.1	5.8	1.0	1.0	5.8	5.7	7.2	6.8
FAMILY INCOME								
Less Than \$20,000	19.9	20.5	2.9	2.9	15.6 ^a	13.7	23.7	25.1
\$20,000 - \$49,999	13.9	14.0	2.5	2.1	13.1	13.7	15.3	15.4
\$50,000 - \$74,999	10.9	9.7	1.7	1.9	11.4	10.7	11.9	10.4
\$75,000 or More	5.9	5.9	1.0	0.7	9.6 ^a	7.8	6.2	6.4

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.35A Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	31,033	31,011	477	418	4,507 ^a	4,182	26,049	26,411
GEOGRAPHIC DIVISION								
Northeast	4,974	5,275	67	62	737	686	4,170	4,527
New England	1,284	1,275	18	15	194	201	1,072	1,058
Middle Atlantic	3,690	4,001	50	47	543	485	3,098	3,469
Midwest	8,100	7,969	115	130	1,158	1,101	6,828	6,738
East North Central	5,579	5,644	81	93	757	738	4,741	4,813
West North Central	2,521	2,325	34	38	401	363	2,087	1,925
South	12,413	12,669	186	160	1,735	1,674	10,492	10,836
South Atlantic	6,320	6,547	92	69	847	767	5,381	5,712
East South Central	2,490	2,580	42	51	390	352	2,059	2,177
West South Central	3,603	3,541	52	40	498	554	3,053	2,946
West	5,546	5,098	109 ^a	66	878 ^a	721	4,559	4,311
Mountain	1,892	2,058	38	30	334	294	1,519	1,734
Pacific	3,654 ^a	3,040	71	36	544	427	3,040	2,576
COUNTY TYPE								
Large Metro	13,707	14,331	199	164	1,956	1,948	11,552	12,220
Small Metro	10,140	9,950	153	148	1,483	1,371	8,504	8,431
250K – 1 Mil. Pop.	6,311	6,386	103	98	902	843	5,306	5,446
< 250K Pop.	3,829	3,564	50	50	581	528	3,198	2,985
Nonmetro	7,187	6,729	126	106	1,068 ^a	863	5,993	5,760
Urbanized	3,083	2,693	53	40	464	371	2,566	2,282
Less Urbanized	3,446	3,396	62	50	527 ^a	405	2,856	2,942
Completely Rural	658	640	10	16	77	87	571	537

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.35B Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	11.8	11.7	1.9	1.7	13.0 ^a	12.0	12.9	12.9
GEOGRAPHIC DIVISION								
Northeast	10.5	11.1	1.6	1.5	12.0	11.2	11.3	12.1
New England	10.3	10.1	1.6	1.4	11.9	12.3	11.0	10.8
Middle Atlantic	10.6	11.4	1.6	1.5	12.0	10.8	11.4	12.6
Midwest	14.4	14.1	2.1	2.4	15.6	14.8	15.7	15.4
East North Central	14.3	14.4	2.2	2.5	14.8	14.4	15.7	15.9
West North Central	14.6	13.4	2.0	2.3	17.4	15.6	15.7	14.4
South	12.7	12.8	2.0	1.7	13.5	12.9	13.9	14.2
South Atlantic	12.3	12.6	2.0	1.5	12.9	11.6	13.4	14.0
East South Central	16.1	16.5	2.8	3.4	19.1	17.2	17.2	18.0
West South Central	11.8	11.4	1.6	1.2	11.7	12.9	13.2	12.5
West	9.1	8.2	1.8 ^a	1.1	10.5 ^a	8.6	9.7	9.0
Mountain	10.1	10.8	2.0	1.6	13.2	11.5	10.6	11.9
Pacific	8.6 ^a	7.0	1.7	0.9	9.3	7.3	9.3	7.7
COUNTY TYPE								
Large Metro	9.7	9.8	1.5	1.2	10.4	10.2	10.6	10.8
Small Metro	12.9	12.9	2.1	2.1	13.6	12.9	14.0	14.1
250K – 1 Mil. Pop.	11.8	12.2	2.0	2.0	12.9	12.3	12.9	13.4
< 250K Pop.	15.0	14.2	2.2	2.3	15.0	14.0	16.6	15.7
Nonmetro	16.9	16.4	3.2	2.8	20.7 ^b	16.8	17.9	17.9
Urbanized	16.2	15.6	3.2	2.6	18.8	15.5	17.3	17.1
Less Urbanized	18.1	17.2	3.5	2.7	23.7 ^b	17.3	19.1	19.0
Completely Rural	14.7	15.2	2.2	4.4	16.2	22.3	16.1	15.5

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.36A Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	First Used at Age 14 or Younger (2013)	First Used at Age 14 or Younger (2014)	First Used at Age 15-17 (2013)	First Used at Age 15-17 (2014)	First Used at Age 18 or Older (2013)	First Used at Age 18 or Older (2014)
TOTAL	30,556	30,593	13,237	13,059	10,717	10,978	6,602	6,555
AGE								
18-25	4,507	4,182	2,062 ^a	1,794	1,672	1,640	774	747
26-34	6,092	5,616	2,861	2,764	2,108	1,775	1,124	1,077
35-49	8,836	8,719	4,123	3,936	2,852	2,975	1,861	1,809
50 or Older	11,121	12,075	4,192	4,565	4,085	4,588	2,844	2,922
GENDER								
Male	16,327	16,192	7,437	7,245	5,599	5,766	3,291	3,180
Female	14,229	14,401	5,800	5,814	5,118	5,212	3,311	3,375
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	28,083	28,178	12,213	11,980	9,898	10,162	5,972	6,037
White	22,895	22,633	10,597	10,287	7,988	8,178	4,310	4,168
Black or African American	3,842	4,096	1,073	1,166	1,395	1,514	1,374	1,417
Other or Two or More Races	1,346	1,449	543	527	*	470	288 ^a	452
Hispanic or Latino	2,473	2,414	1,024	1,079	819	816	630	519
EDUCATION								
< High School	7,032	6,591	3,643	3,535	2,211	2,146	1,179	909
High School Graduate	12,024	12,191	4,950	4,851	4,549	4,724	2,525	2,616
Some College	8,478	8,645	3,490	3,625	2,830	2,926	2,158	2,094
College Graduate	3,022	3,166	1,154	1,047	1,127	1,182	741	937

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.36B Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	First Used at Age 14 or Younger (2013)	First Used at Age 14 or Younger (2014)	First Used at Age 15-17 (2013)	First Used at Age 15-17 (2014)	First Used at Age 18 or Older (2013)	First Used at Age 18 or Older (2014)
TOTAL	56.2	56.6	67.2	65.3	53.7	56.3	44.8	45.1
AGE								
18-25	42.3	42.2	57.6	55.5	38.6	40.5	28.2	28.5
26-34	49.5	50.5	57.7	62.5	47.0	46.7	39.2	37.1
35-49	59.8	58.7	70.7	66.2	58.7	60.0	45.5	45.8
50 or Older	66.8	66.6	78.5 ^a	71.4	65.2	68.6	56.6	57.8
GENDER								
Male	55.7	55.5	68.5 ^a	64.5	52.3	54.7	42.4	43.0
Female	56.8	57.9	65.5	66.3	55.4	58.2	47.5	47.4
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	58.9	59.7	69.1	67.4	56.6	58.7	47.8	50.0
White	60.8	61.0	69.8	67.9	57.6	59.1	50.1	51.3
Black or African American	54.2	58.2	69.2	67.7	53.9	61.1	46.7	49.8
Other or Two or More Races	46.5	48.1	58.7	58.5	*	47.0	30.6	40.6
Hispanic or Latino	36.7	35.2	50.1	48.5	33.4	37.4	28.2	21.2
EDUCATION								
< High School	65.1	65.6	74.8	74.8	59.7	64.8	52.7	45.2
High School Graduate	62.1	61.9	70.5	69.3	61.1	61.4	51.6	52.3
Some College	52.5	53.1	64.8	62.9	47.8	52.0	44.5	42.8
College Graduate	37.5	39.6	47.4	41.7	39.2	41.3	26.9 ^a	35.7

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.37A Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes Daily and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total		AGE FIRST USED CIGARETTES DAILY ¹						Never Used Cigarettes Daily ¹	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	30,556	30,593	5,219	5,421	11,036	11,184	13,395	12,977	907	1,010
AGE										
18-25	4,507	4,182	703	663	1,839 ^a	1,600	1,609	1,641	357	278
26-34	6,092	5,616	1,069	978	2,339	2,098	2,553	2,381	131	159
35-49	8,836	8,719	1,710	1,637	3,083	3,269	3,773	3,622	270	191
50 or Older	11,121	12,075	1,737	2,143	3,775	4,218	5,459	5,332	*	382
GENDER										
Male	16,327	16,192	2,895	2,890	5,899	6,187	6,995	6,529	538	586
Female	14,229	14,401	2,325	2,531	5,136	4,997	6,400	6,448	368	425
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	28,083	28,178	4,834	4,977	10,386	10,514	12,181	11,869	682	818
White	22,895	22,633	4,148	4,353	8,837	8,907	9,539	9,042	371	330
Black or African American	3,842	4,096	*	*	*	1,163	2,126	2,163	273	374
Other or Two or More Races	1,346	1,449	*	*	*	444	517	663	38 ^a	114
Hispanic or Latino	2,473	2,414	*	444	650	670	1,213	1,108	224	192
EDUCATION										
< High School	7,032	6,591	1,933	1,787	2,730	2,677	2,145	1,841	223	285
High School Graduate	12,024	12,191	1,836	2,083	4,559	4,605	5,176	5,125	453	378
Some College	8,478	8,645	1,159	1,273	2,836	2,971	4,327	4,146	155	256
College Graduate	3,022	3,166	*	*	910	931	1,747	1,865	75	92

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.37B Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes Daily and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		AGE FIRST USED CIGARETTES DAILY ²						Never Used Cigarettes Daily ²	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	56.2	56.6	83.2 ^a	78.6	70.0	72.9	55.4	56.1	11.1	11.7
AGE										
18-25	42.3	42.2	76.5	77.2	64.4	61.6	47.5	49.6	10.2	8.9
26-34	49.5	50.5	76.9	79.3	64.3	68.9	46.6	51.0	7.3	7.3
35-49	59.8	58.7	85.8	80.1	75.2	76.3	54.6	54.2	15.2	10.4
50 or Older	66.8	66.6	88.2 ^a	77.7	72.8	78.0	65.0	62.9	*	25.6
GENDER										
Male	55.7	55.5	83.6	78.5	72.0	73.9	54.9	54.2	10.9	11.6
Female	56.8	57.9	82.7	78.8	67.8	71.7	56.0	58.1	11.4	11.9
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	58.9	59.7	83.6	79.3	71.5	74.5	57.0	57.8	11.4	13.1
White	60.8	61.0	83.9	79.7	72.5	74.4	57.5	57.6	9.4	8.3
Black or African American	54.2	58.2	*	*	*	79.5	59.9	62.7	17.8	23.0
Other or Two or More Races	46.5	48.1	*	*	*	65.4	41.6	48.1	7.8 ^a	17.6
Hispanic or Latino	36.7	35.2	*	71.8	51.9	54.9	43.1	42.1	10.3	8.1
EDUCATION										
< High School	65.1	65.6	87.0 ^a	80.1	73.6	79.1	62.2	63.2	15.7	18.8
High School Graduate	62.1	61.9	81.4	81.6	73.9	74.7	61.8	60.3	17.7	15.2
Some College	52.5	53.1	84.0 ^a	75.1	67.7	69.6	53.8	54.3	6.1	9.5
College Graduate	37.5	39.6	*	*	53.7	61.2	40.4	45.3	4.6	4.7

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Percentages in the Total column are among all persons in the specified subgroup who used cigarettes in the past month, regardless of whether they ever used cigarettes daily.

² Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.38A Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total		AGE FIRST USED MARIJUANA						Never Used Marijuana	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	5,976	6,210	2,315	2,277	2,024 ^b	2,484	1,308	1,156	331	293
AGE										
18-25	2,557	2,320	1,000	858	1,045	1,018	443	396	69	48
26-34	1,606	1,566	724	661	533	581	285	283	64	41
35-49	1,028	1,217	457	445	246 ^b	454	189	269	136	50
50 or Older	785	1,107	134 ^a	314	200 ^a	430	391	209	61	154
GENDER										
Male	3,798	3,966	1,595	1,493	1,307 ^a	1,626	754	742	141	105
Female	2,179	2,244	720	784	716	858	554	414	189	188
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	5,130	5,269	1,978	1,931	1,781 ^a	2,099	1,130	995	242	244
White	3,799	3,665	1,541	1,466	1,304	1,475	824	632	129	92
Black or African American	906 ^a	1,192	324	403	326	450	223	223	34	116
Other or Two or More Races	425	412	112	62	151	174	83	140	79	36
Hispanic or Latino	846	941	337	346	243 ^a	385	178	161	89	49
EDUCATION										
< High School	1,215	1,183	575	550	388	376	126	153	126	104
High School Graduate	2,000	1,860	899	757	702	807	257	243	141	53
Some College	1,933	2,068	620	691	660 ^a	853	612	404	41	120
College Graduate	829 ^a	1,099	220	279	274 ^b	448	312	357	23	16

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.38B Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		AGE FIRST USED MARIJUANA						Never Used Marijuana	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	2.5	2.6	11.5	11.5	5.0 ^a	6.0	2.6	2.2	0.3	0.2
AGE										
18-25	7.4	6.6	23.2	20.3	13.3	12.7	7.5	6.4	0.4	0.3
26-34	4.3	4.1	14.4	13.8	6.3	7.0	3.6	3.5	0.4	0.2
35-49	1.7	2.0	7.5	7.5	2.1 ^b	3.9	1.6	2.1	0.4	0.2
50 or Older	0.7	1.0	2.8 ^a	6.4	1.6 ^a	3.2	1.6	0.8	0.1	0.2
GENDER										
Male	3.3	3.4	13.6	12.6	5.9 ^a	7.1	2.9	2.8	0.3	0.2
Female	1.8	1.8	8.6	9.8	3.9	4.6	2.3	1.7	0.3	0.3
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	2.5	2.6	11.4	11.3	4.9	5.7	2.5	2.1	0.2	0.2
White	2.4	2.3	10.6	10.5	4.4	4.9	2.3	1.7	0.2	0.1
Black or African American	3.3	4.2	17.3	18.4	7.0	9.8	3.7	3.6	0.2	0.8
Other or Two or More Races	2.4	2.2	11.4	6.7	9.0	9.2	3.2	4.6	0.6	0.3
Hispanic or Latino	2.4	2.6	12.2	12.7	5.4	8.2	3.4	3.1	0.4	0.2
EDUCATION										
< High School	3.8	3.7	16.5	15.4	9.1	8.7	3.6	4.1	0.6	0.5
High School Graduate	2.9	2.7	14.4	11.9	5.5	6.4	2.2	2.0	0.4	0.1
Some College	3.0	3.2	10.4	11.4	5.4	6.4	4.1	2.7	0.1	0.4
College Graduate	1.2 ^a	1.5	4.9	7.3	2.4 ^b	4.0	1.6	1.7	0.1	0.0

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Percentages in the Total column are among all persons in the specified subgroup, regardless of whether they ever used marijuana.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.39A Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total		AGE FIRST USED ALCOHOL					
			14 or Younger		15-17		18 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	16,602	16,316	6,006	5,771	7,167	6,789	3,429	3,755
AGE								
18-25	4,529	4,284	1,660 ^a	1,419	2,081	1,984	789	880
26-34	4,212	3,748	1,564	1,368	1,888 ^b	1,369	759 ^a	1,011
35-49	4,233	4,227	1,785	1,637	1,618	1,761	830	829
50 or Older	3,628	4,056	997	1,347	1,580	1,676	1,051	1,034
GENDER								
Male	10,769	10,633	4,052	3,807	4,552	4,474	2,165	2,352
Female	5,833	5,682	1,954	1,964	2,616	2,315	1,264	1,403
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	13,752	13,568	4,931	4,879	5,977	5,670	2,845	3,020
White	11,063	10,629	4,113	4,116	4,862	4,461	2,088	2,052
Black or African American	1,617	1,834	495	495	632	736	490	602
Other or Two or More Races	1,072	1,106	323	267	483	473	266	365
Hispanic or Latino	2,850	2,747	1,075	893	1,190	1,119	584	735
EDUCATION								
< High School	2,158	2,126	886	770	771	928	500	428
High School Graduate	4,408	4,189	1,682	1,500	1,790	1,727	937	962
Some College	4,819	5,003	1,641	1,781	2,253	2,003	926 ^a	1,219
College Graduate	5,216	4,997	1,797	1,720	2,353	2,132	1,066	1,146

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.39B Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		AGE FIRST USED ALCOHOL					
			14 or Younger		15-17		18 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	7.0	6.8	15.4	14.6	9.3	8.9	3.8	4.0
AGE								
18-25	13.0	12.3	27.5 ^a	24.0	16.5	16.1	7.5	8.1
26-34	11.3	9.9	22.3	18.6	14.4 ^a	11.4	5.6	6.7
35-49	7.0	7.0	14.2	13.5	8.0	8.7	3.9	3.8
50 or Older	3.5	3.8	7.5	9.6	5.1	5.3	2.4	2.2
GENDER								
Male	9.4	9.2	17.1	15.9	10.9	10.7	5.7	6.0
Female	4.7	4.6	12.8	12.7	7.4	6.7	2.4	2.5
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	6.8	6.7	14.6	14.1	8.8	8.5	3.7	3.8
White	7.1	6.8	14.1	14.0	8.5	8.0	3.7	3.5
Black or African American	5.8	6.5	16.8	15.2	9.0	10.7	4.0	4.7
Other or Two or More Races	6.1	6.0	18.3	13.2	13.8	12.9	3.6	4.5
Hispanic or Latino	8.0	7.5	20.8	18.5	12.7	11.4	4.2	4.9
EDUCATION								
< High School	6.7	6.6	17.6	16.1	10.1	11.2	4.7	4.0
High School Graduate	6.3	6.1	16.6	14.2	8.2	8.0	3.5	3.6
Some College	7.6	7.6	15.0	15.3	10.2	9.2	3.8	4.6
College Graduate	7.2	6.8	13.8	13.8	9.2	8.6	3.8	3.7

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Percentages in the Total column are among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.40A Alcohol Dependence or Abuse in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Numbers in Thousands, 2013 and 2014

Age Group	Total		AGE FIRST USED ALCOHOL							
			14 or Younger		15-17		18-20		21 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
21 OR OLDER	15,153	15,042	5,439	5,270	6,438	6,157	2,412	2,720	864	895
21 to 25	3,080	3,011	1,093 ^a	919	1,351	1,352	512	586	124	154
26 or Older	12,072	12,032	4,346	4,352	5,087	4,805	1,900	2,134	740	741
26 to 49	8,444	7,975	3,349	3,005	3,506	3,129	1,197	1,292	392	549
26 to 34	4,212	3,748	1,564	1,368	1,888 ^b	1,369	594	704	166 ^a	307
35 to 49	4,233	4,227	1,785	1,637	1,618	1,761	603	588	227	242
50 or Older	3,628	4,056	997	1,347	1,580	1,676	704	842	347	192

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.40B Alcohol Dependence or Abuse in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Percentages, 2013 and 2014

Age Group	Total ¹		AGE FIRST USED ALCOHOL							
			14 or Younger		15-17		18-20		21 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
21 OR OLDER	6.8	6.6	14.8	14.1	8.9	8.6	4.8	5.1	2.3	2.3
21 to 25	14.3	13.8	28.6 ^b	24.0	18.0	18.3	9.9	11.4	4.0	4.5
26 or Older	6.0	5.9	13.2	13.0	7.9	7.5	4.2	4.4	2.2	2.1
26 to 49	8.6	8.1	17.1	15.5	10.5	9.7	6.3	6.3	2.5	3.3
26 to 34	11.3	9.9	22.3	18.6	14.4 ^a	11.4	8.0	8.4	2.7	4.5
35 to 49	7.0	7.0	14.2	13.5	8.0	8.7	5.3	4.9	2.3	2.5
50 or Older	3.5	3.8	7.5	9.6	5.1	5.3	2.7	3.0	1.9	1.0

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Percentages in the Total column are among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.41A Illicit Drug Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total		AGE FIRST USED MARIJUANA						Never Used Marijuana	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	4,437	4,536	1,823	1,746	1,477 ^b	1,867	999	760	138	163
AGE										
18-25	1,808	1,661	762	658	694	713	317	249	35	42
26-34	1,322	1,268	619	545	440	486	235	197	27	40
35-49	693 ^a	927	347	354	192 ^b	389	103	170	51	14
50 or Older	614	680	94	189	150	279	344	145	25	67
GENDER										
Male	2,697	2,818	1,205	1,095	884 ^b	1,199	556	465	52	59
Female	1,741	1,717	619	651	593	668	443	294	86	104
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	3,890	3,912	1,569	1,493	1,319 ^a	1,603	876	661	125	155
White	2,937	2,705	1,216	1,135	988	1,086	657	413	75	71
Black or African American	652 ^a	897	261	304	220 ^a	379	150	155	21	57
Other or Two or More Races	301	310	92	53	111	137	70	93	29	26
Hispanic or Latino	547	624	254	253	158 ^a	264	123	99	12	8
EDUCATION										
< High School	878	841	466	404	282	271	105	113	25	53
High School Graduate	1,462	1,402	687	598	506	572	194	196	75	36
Some College	1,477	1,490	522	527	474 ^a	653	461	242	20	68
College Graduate	620	803	148	217	215 ^b	371	239	209	18	7

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.41B Illicit Drug Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		AGE FIRST USED MARIJUANA						Never Used Marijuana	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	1.9	1.9	9.1	8.8	3.6 ^a	4.5	2.0	1.5	0.1	0.1
AGE										
18-25	5.2	4.8	17.7	15.6	8.8	8.9	5.4	4.0	0.2	0.3
26-34	3.5	3.3	12.3	11.4	5.2	5.8	3.0	2.5	0.2	0.2
35-49	1.1 ^a	1.5	5.7	6.0	1.6 ^b	3.4	0.9	1.3	0.2	0.0
50 or Older	0.6	0.6	2.0	3.9	1.2	2.0	1.4	0.6	0.0	0.1
GENDER										
Male	2.4	2.4	10.3	9.2	4.0 ^a	5.2	2.2	1.7	0.1	0.1
Female	1.4	1.4	7.4	8.2	3.2	3.6	1.8	1.2	0.1	0.1
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	1.9	1.9	9.0	8.7	3.7	4.4	2.0	1.4	0.1	0.2
White	1.9	1.7	8.4	8.1	3.3	3.6	1.8	1.1	0.1	0.1
Black or African American	2.3	3.2	13.9	13.9	4.7 ^a	8.3	2.5	2.5	0.1	0.4
Other or Two or More Races	1.7	1.7	9.3	5.8	6.6	7.3	2.7	3.1	0.2	0.2
Hispanic or Latino	1.5	1.7	9.2	9.3	3.5	5.6	2.3	1.9	0.1	0.0
EDUCATION										
< High School	2.7	2.6	13.4	11.3	6.6	6.3	3.0	3.0	0.1	0.3
High School Graduate	2.1	2.0	11.0	9.4	4.0	4.5	1.6	1.6	0.2	0.1
Some College	2.3	2.3	8.8	8.7	3.8	4.9	3.1	1.6	0.1	0.2
College Graduate	0.9	1.1	3.3	5.7	1.9 ^b	3.3	1.2	1.0	0.0	0.0

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Percentages in the Total column are among all persons in the specified subgroup, regardless of whether they ever used marijuana.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.42A Alcohol Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total		AGE FIRST USED ALCOHOL					
			14 or Younger		15-17		18 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	7,736	7,824	3,004	2,898	3,120	3,139	1,613	1,787
AGE								
18-25	1,922	1,989	738	694	899	898	285 ^a	397
26-34	2,014	1,748	795	705	868	622	351	421
35-49	2,025	2,071	1,018	795	574 ^b	849	433	427
50 or Older	1,776	2,016	454	704	778	770	544	542
GENDER								
Male	4,813	4,736	1,934	1,762	1,859	1,926	1,020	1,048
Female	2,923	3,088	1,070	1,137	1,261	1,213	593	739
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	6,537	6,628	2,555	2,450	2,552	2,673	1,430	1,505
White	5,165	4,945	2,044	2,000	2,128	1,980	992	965
Black or African American	825	1,028	298	272	253 ^a	408	275	348
Other or Two or More Races	547	655	213	178	171 ^a	284	164	193
Hispanic or Latino	1,199	1,196	448	448	568	466	183	282
EDUCATION								
< High School	1,178	1,261	551	475	376 ^a	556	251	230
High School Graduate	2,044	2,025	817	819	774	773	453	433
Some College	2,129	2,503	757	917	913	924	459 ^a	662
College Graduate	2,385	2,036	878	687	1,056	886	450	462

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.42B Alcohol Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		AGE FIRST USED ALCOHOL					
			14 or Younger		15-17		18 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	3.3	3.3	7.7	7.3	4.0	4.1	1.8	1.9
AGE								
18-25	5.5	5.7	12.2	11.7	7.2	7.3	2.7 ^a	3.6
26-34	5.4	4.6	11.3	9.6	6.6	5.2	2.6	2.8
35-49	3.3	3.4	8.1	6.6	2.8 ^b	4.2	2.0	2.0
50 or Older	1.7	1.9	3.4	5.0	2.5	2.4	1.2	1.2
GENDER								
Male	4.2	4.1	8.2	7.3	4.4	4.6	2.7	2.7
Female	2.4	2.5	7.0	7.4	3.6	3.5	1.1	1.3
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	3.2	3.3	7.5	7.1	3.8	4.0	1.9	1.9
White	3.3	3.2	7.0	6.8	3.7	3.5	1.8	1.6
Black or African American	3.0	3.6	10.1	8.3	3.6 ^a	5.9	2.2	2.7
Other or Two or More Races	3.1	3.5	12.1	8.8	4.9	7.7	2.2	2.4
Hispanic or Latino	3.4	3.2	8.7	9.3	6.1	4.8	1.3	1.9
EDUCATION								
< High School	3.7	3.9	11.0	9.9	4.9	6.7	2.4	2.1
High School Graduate	2.9	2.9	8.1	7.8	3.5	3.6	1.7	1.6
Some College	3.4	3.8	6.9	7.9	4.2	4.3	1.9	2.5
College Graduate	3.3	2.8	6.8	5.5	4.1	3.6	1.6	1.5

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Percentages in the Total column are among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.43A Alcohol Dependence in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Numbers in Thousands, 2013 and 2014

Age Group	Total		AGE FIRST USED ALCOHOL							
			14 or Younger		15-17		18-20		21 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
21 OR OLDER	7,146	7,257	2,751	2,660	2,839	2,867	1,126	1,304	430	425
21 to 25	1,331	1,422	485	456	619	626	181 ^a	258	47	82
26 or Older	5,815	5,835	2,266	2,204	2,220	2,241	945	1,047	383	343
26 to 49	4,038	3,819	1,812	1,500	1,442	1,471	568	612	216	236
26 to 34	2,014	1,748	795	705	868	622	265	321	86	100
35 to 49	2,025	2,071	1,018	795	574 ^b	849	303	292	130	135
50 or Older	1,776	2,016	454	704	778	770	377	435	167	108

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.43B Alcohol Dependence in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Percentages, 2013 and 2014

Age Group	Total ¹		AGE FIRST USED ALCOHOL							
			14 or Younger		15-17		18-20		21 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
21 OR OLDER	3.2	3.2	7.5	7.1	3.9	4.0	2.2	2.4	1.2	1.1
21 to 25	6.2	6.5	12.7	11.9	8.3	8.5	3.5 ^a	5.0	1.5	2.4
26 or Older	2.9	2.8	6.9	6.6	3.4	3.5	2.1	2.2	1.1	1.0
26 to 49	4.1	3.9	9.2	7.7	4.3	4.6	3.0	3.0	1.4	1.4
26 to 34	5.4	4.6	11.3	9.6	6.6	5.2	3.6	3.8	1.4	1.5
35 to 49	3.3	3.4	8.1	6.6	2.8 ^b	4.2	2.6	2.4	1.3	1.4
50 or Older	1.7	1.9	3.4	5.0	2.5	2.4	1.4	1.6	0.9	0.6

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Percentages in the Total column are among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.44A Method of Obtaining Most Recently Used Marijuana among Past Year Marijuana Users, by Age Group: Numbers in Thousands, 2013 and 2014

Method of Obtaining Most Recently Used Marijuana	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS	32,952 ^a	35,124	3,346	3,262	11,000	11,154	18,606 ^b	20,707
Bought It	14,706 ^b	16,641	1,266	1,236	4,958	5,243	8,483 ^b	10,161
Traded for It	538	384	37	37	113	86	389	261
Got It Free or Shared Someone Else's	16,382	16,957	1,849	1,747	5,696	5,578	8,837	9,632
Grew It Yourself	764 ^a	403	17	31	66	53	680 ^a	319
Method Unspecified	561 ^a	739	177	210	167	194	217	335

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.44B Method of Obtaining Most Recently Used Marijuana among Past Year Marijuana Users, by Age Group: Percentages, 2013 and 2014

Method of Obtaining Most Recently Used Marijuana	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Bought It	44.6 ^a	47.4	37.8	37.9	45.1	47.0	45.6 ^a	49.1
Traded for It	1.6 ^a	1.1	1.1	1.1	1.0	0.8	2.1	1.3
Got It Free or Shared Someone Else's	49.7	48.3	55.3	53.6	51.8	50.0	47.5	46.5
Grew It Yourself	2.3 ^b	1.1	0.5	1.0	0.6	0.5	3.7 ^b	1.5
Method Unspecified	1.7	2.1	5.3	6.4	1.5	1.7	1.2	1.6

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.45A Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Bought Their Most Recently Used Marijuana, by Age Group: Numbers in Thousands, 2013 and 2014

Source and Location of Bought Marijuana	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS WHO BOUGHT MOST RECENTLY USED MARIJUANA	14,706 ^b	16,641	1,266	1,236	4,958	5,243	8,483 ^b	10,161
SOURCE								
Friend	11,017 ^a	12,082	966	1,005	3,700	3,836	6,350 ^a	7,240
Relative or Family Member	793	755	40	26	138	187	615	541
Someone Just Met or Did Not Know Well	2,291 ^a	2,780	203	153	879	982	1,210 ^a	1,645
Source Unspecified	606 ^b	1,024	57	52	241	238	308 ^b	734
LOCATION								
Inside Public Building ¹	1,030 ^b	1,523	45	71	320 ^a	425	664 ^b	1,027
Outside in Public Area ²	3,182	3,330	476	451	1,122	1,165	1,584	1,715
Inside School Building	126	133	67	57	48	50	12	26
Outside on School Property	121	144	32	26	37	68	52	50
Inside Home, Apartment, or Dorm	8,186	8,859	448	479	2,703	2,838	5,035	5,543
Medical Marijuana Dispensary/Club ³	338 ^a	584	1	1	110	105	227 ^a	478
Some Other Place ⁴	144	177	23	14	35	45	86	118
Location Unspecified ⁵	1,580	1,890	174	138	583	548	823 ^a	1,204

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ For example, a store, restaurant, sports arena, bar, or club.

² For example, a parking lot, street, or park.

³ Medical Marijuana Dispensary/Club includes respondents who reported that they bought marijuana in some other place and specified that this other location was a medical marijuana dispensary or a marijuana club.

⁴ Some Other Place includes respondents who reported that they bought marijuana in some other place and specified a location other than a medical marijuana dispensary/club that did not correspond to an existing category. Examples of these other locations include private backyard, prison, and hotel room. Because of revisions to this category to remove Medical Marijuana Dispensary/Club in 2013, these estimates are not comparable with the estimates presented before the 2013 NSDUH.

⁵ Location Unspecified includes respondents with unknown, invalid, or nonspecific responses to the other-specify question (e.g., in a car, at a concert, or at a party where the specific locations were not provided).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.45B Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Bought Their Most Recently Used Marijuana, by Age Group: Percentages, 2013 and 2014

Source and Location of Bought Marijuana	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS WHO BOUGHT MOST RECENTLY USED MARIJUANA	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SOURCE								
Friend	74.9	72.6	76.3 ^a	81.3	74.6	73.2	74.9	71.3
Relative or Family Member	5.4	4.5	3.2	2.1	2.8	3.6	7.2	5.3
Someone Just Met or Did Not Know Well	15.6	16.7	16.0	12.4	17.7	18.7	14.3	16.2
Source Unspecified	4.1 ^b	6.2	4.5	4.2	4.9	4.5	3.6 ^b	7.2
LOCATION								
Inside Public Building ¹	7.0 ^a	9.2	3.6	5.7	6.5	8.1	7.8	10.1
Outside in Public Area ²	21.6	20.0	37.6	36.4	22.6	22.2	18.7	16.9
Inside School Building	0.9	0.8	5.3	4.6	1.0	1.0	0.1	0.3
Outside on School Property	0.8	0.9	2.5	2.1	0.7	1.3	0.6	0.5
Inside Home, Apartment, or Dorm	55.7	53.2	35.4	38.7	54.5	54.1	59.4	54.6
Medical Marijuana Dispensary/Club ³	2.3	3.5	0.0	0.1	2.2	2.0	2.7	4.7
Some Other Place ⁴	1.0	1.1	1.9	1.1	0.7	0.9	1.0	1.2
Location Unspecified ⁵	10.7	11.4	13.7	11.1	11.8	10.4	9.7	11.9

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ For example, a store, restaurant, sports arena, bar, or club.

² For example, a parking lot, street, or park.

³ Medical Marijuana Dispensary/Club includes respondents who reported that they bought marijuana in some other place and specified that this other location was a medical marijuana dispensary or a marijuana club.

⁴ Some Other Place includes respondents who reported that they bought marijuana in some other place and specified a location other than a medical marijuana dispensary/club that did not correspond to an existing category. Examples of these other locations include private backyard, prison, and hotel room. Because of revisions to this category to remove Medical Marijuana Dispensary/Club in 2013, these estimates are not comparable with the estimates presented before the 2013 NSDUH.

⁵ Location Unspecified includes respondents with unknown, invalid, or nonspecific responses to the other-specify question (e.g., in a car, at a concert, or at a party where the specific locations were not provided).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.46A Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Obtained for Free Their Most Recently Used Marijuana and Who Had Not Bought or Traded for Marijuana in the Past Year, by Age Group: Numbers in Thousands, 2013 and 2014

Source and Location of Free Marijuana ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS WHO OBTAINED FOR FREE MOST RECENTLY USED MARIJUANA AND WHO HAD NOT BOUGHT OR TRADED FOR MARIJUANA IN PAST YEAR	12,591	12,781	1,316	1,240	4,166	4,050	7,109	7,491
SOURCE								
Friend	10,110	9,959	1,087	991	3,492	3,306	5,531	5,663
Relative or Family Member	1,612	1,902	132	131	362	422	1,117	1,350
Someone Just Met or Did Not Know Well	644	708	62	86	258	262	324	360
Source Unspecified	225	211	35	31	54	61	137	119
LOCATION								
Inside Public Building ²	466	447	29	18	126	88	311	341
Outside in Public Area ³	1,567	1,578	351	281	574	539	643	758
Inside School Building	115	68	38	24	25	29	52	14
Outside on School Property	130	139	62	57	55	73	13	9
Inside Home, Apartment, or Dorm	8,956	9,169	670	692	3,076	2,960	5,210	5,517
Some Other Place ⁴	220	228	22	12	41	44	157	173
Location Unspecified ⁵	1,137	1,151	143	155	271	317	723	679

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents who reported that they had bought or traded for marijuana in the past year or who had missing data for whether they bought or traded for marijuana in the past year were excluded from this analysis.

² For example, a store, restaurant, sports arena, bar, or club.

³ For example, a parking lot, street, or park.

⁴ Some Other Place includes respondents who reported that they obtained marijuana for free in some other place and specified a location (e.g., private backyard, prison, hotel room) that did not correspond to an existing category.

⁵ Location Unspecified includes respondents with unknown, invalid, or nonspecific responses to the other-specify question (e.g., in a car, at a concert, or at a party where the specific locations were not provided).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.46B Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Obtained for Free Their Most Recently Used Marijuana and Who Had Not Bought or Traded for Marijuana in the Past Year, by Age Group: Percentages, 2013 and 2014

Source and Location of Free Marijuana ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS WHO OBTAINED FOR FREE MOST RECENTLY USED MARIJUANA AND WHO HAD NOT BOUGHT OR TRADED FOR MARIJUANA IN PAST YEAR	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SOURCE								
Friend	80.3	77.9	82.6	79.9	83.8	81.6	77.8	75.6
Relative or Family Member	12.8	14.9	10.1	10.6	8.7	10.4	15.7	18.0
Someone Just Met or Did Not Know Well	5.1	5.5	4.7	7.0	6.2	6.5	4.6	4.8
Source Unspecified	1.8	1.7	2.6	2.5	1.3	1.5	1.9	1.6
LOCATION								
Inside Public Building ²	3.7	3.5	2.2	1.4	3.0	2.2	4.4	4.6
Outside in Public Area ³	12.4	12.3	26.7	22.7	13.8	13.3	9.0	10.1
Inside School Building	0.9	0.5	2.9	2.0	0.6	0.7	0.7	0.2
Outside on School Property	1.0	1.1	4.7	4.6	1.3	1.8	0.2	0.1
Inside Home, Apartment, or Dorm	71.1	71.7	51.0	55.8	73.8	73.1	73.3	73.6
Some Other Place ⁴	1.7	1.8	1.7	0.9	1.0	1.1	2.2	2.3
Location Unspecified ⁵	9.0	9.0	10.9	12.5	6.5	7.8	10.2	9.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents who reported that they had bought or traded for marijuana in the past year or who had missing data for whether they bought or traded for marijuana in the past year were excluded from this analysis.

² For example, a store, restaurant, sports arena, bar, or club.

³ For example, a parking lot, street, or park.

⁴ Some Other Place includes respondents who reported that they obtained marijuana for free in some other place and specified a location (e.g., private backyard, prison, hotel room) that did not correspond to an existing category.

⁵ Location Unspecified includes respondents with unknown, invalid, or nonspecific responses to the other-specify question (e.g., in a car, at a concert, or at a party where the specific locations were not provided).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.47A Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	2,133	2,142	241	211	481	450	1,412	1,481
From More Than One Doctor	198	303	25	30	46	50	127	222
Wrote Fake Prescription	23 ^a	9	8	2	10	5	5	1
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	91	38	12	4	17	6	62	28
From Friend or Relative for Free	5,838 ^b	4,895	515 ^b	397	1,688 ^b	1,336	3,636 ^a	3,162
Bought from Friend or Relative	1,179	1,069	103	87	499 ^b	363	577	619
Took from Friend or Relative without Asking	431	428	107	83	145	140	180	205
Bought from Drug Dealer or Other Stranger	468	464	37	38	223	204	208	223
Bought on the Internet	21	7	1	1	5	5	14	1
Some Other Way	433	347	70	68	114	113	249	166

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.47B Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	19.7	22.1	21.5	22.9	14.9	16.8	21.8	24.2
From More Than One Doctor	1.8 ^a	3.1	2.2	3.3	1.4	1.9	2.0	3.6
Wrote Fake Prescription	0.2 ^a	0.1	0.7	0.2	0.3	0.2	0.1	0.0
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8	0.4	1.0	0.4	0.5	0.2	1.0	0.5
From Friend or Relative for Free	54.0 ^a	50.5	46.1	43.1	52.3	50.0	56.2 ^a	51.8
Bought from Friend or Relative	10.9	11.0	9.2	9.4	15.5	13.6	8.9	10.1
Took from Friend or Relative without Asking	4.0	4.4	9.6	9.0	4.5	5.2	2.8	3.3
Bought from Drug Dealer or Other Stranger	4.3	4.8	3.3	4.1	6.9	7.6	3.2	3.6
Bought on the Internet	0.2	0.1	0.1	0.1	0.2	0.2	0.2	0.0
Some Other Way	4.0	3.6	6.3	7.4	3.5	4.2	3.9	2.7

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.48A Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Pain Relievers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	4,331 ^b	3,803	262 ^a	214	1,124 ^b	956	2,945	2,633
From More Than One Doctor	188	150	31	21	71 ^b	37	86	93
Wrote Fake Prescription	17	2	*	0	3	2	14	0
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	19 ^b	3	3	1	14 ^a	3	2	0
From Friend or Relative for Free	286	215	37	26	107 ^a	74	142	114
Bought from Friend or Relative	216	169	31	20	109 ^a	65	77	84
Took from Friend or Relative without Asking	68	42	28	21	32	14	8	7
Bought from Drug Dealer or Other Stranger	74	58	13	14	39	23	21	21
Bought on the Internet	11	9	0	2	1	5	10	2
Some Other Way	60	52	17	13	14	14	28	26

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.48B Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Pain Relievers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	82.2	84.4	62.0	64.7	74.2 ^b	80.1	88.3	88.3
From More Than One Doctor	3.6	3.3	7.3	6.2	4.7	3.1	2.6	3.1
Wrote Fake Prescription	0.3	0.1	*	0.1	0.2	0.1	0.4	0.0
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4 ^b	0.1	0.7	0.2	0.9 ^a	0.2	0.1	0.0
From Friend or Relative for Free	5.4	4.8	8.7	7.9	7.0	6.2	4.3	3.8
Bought from Friend or Relative	4.1	3.8	7.3	6.2	7.2	5.5	2.3	2.8
Took from Friend or Relative without Asking	1.3	0.9	6.6	6.3	2.1	1.2	0.2	0.2
Bought from Drug Dealer or Other Stranger	1.4	1.3	3.2	4.1	2.6	1.9	0.6	0.7
Bought on the Internet	0.2	0.2	0.1	0.6	0.1	0.4	0.3	0.1
Some Other Way	1.1	1.2	4.1	3.8	0.9	1.2	0.9	0.9

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.49A Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	1,152	1,109	114	103	256	202	782	803
From More Than One Doctor	163 ^a	276	16	23	44	42	103 ^a	211
Wrote Fake Prescription	30	14	7	2	2	2	21	10
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	66	36	9 ^a	2	12 ^a	3	44	31
From Friend or Relative for Free	2,537 ^b	2,074	216 ^b	150	734 ^b	520	1,587	1,404
Bought from Friend or Relative	1,093 ^a	884	72 ^a	46	370 ^b	270	651	568
Took from Friend or Relative without Asking	427	340	63 ^a	42	134	102	230	196
Bought from Drug Dealer or Other Stranger	515	496	40	33	246 ^b	168	229	295
Bought on the Internet	46 ^a	9	3	*	15 ^a	1	29	8
Some Other Way	196	180	37	35	77	59	82	86

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.49B Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	27.3	27.9	25.9	30.4	21.9	21.1	29.9	30.0
From More Than One Doctor	3.9 ^a	6.9	3.7	6.7	3.7	4.4	3.9 ^a	7.9
Wrote Fake Prescription	0.7	0.4	1.6	0.6	0.2	0.2	0.8	0.4
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.6	0.9	2.0 ^a	0.7	1.1	0.3	1.7	1.1
From Friend or Relative for Free	60.0 ^b	52.2	49.0	43.5	62.7 ^b	54.3	60.6 ^a	52.5
Bought from Friend or Relative	25.9	22.2	16.4	13.5	31.7	28.2	24.9	21.2
Took from Friend or Relative without Asking	10.1	8.6	14.4	12.3	11.5	10.7	8.8	7.3
Bought from Drug Dealer or Other Stranger	12.2	12.5	9.2	9.7	21.0	17.5	8.7	11.0
Bought on the Internet	1.1 ^a	0.2	0.6	*	1.3 ^a	0.1	1.1	0.3
Some Other Way	4.7	4.6	8.7	10.5	6.7	6.3	3.2	3.3

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.50A Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Pain Relievers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	1,874 ^a	1,600	113	89	504 ^b	381	1,257	1,130
From More Than One Doctor	175 ^a	111	23 ^a	12	75 ^b	33	76	67
Wrote Fake Prescription	44	13	2	*	3	4	39	6
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	27	18	3	*	8	8	16	7
From Friend or Relative for Free	318 ^b	196	26	21	105 ^a	69	187 ^a	106
Bought from Friend or Relative	288 ^b	180	25	16	115 ^a	69	148	96
Took from Friend or Relative without Asking	99	58	18	9	45 ^a	18	37	31
Bought from Drug Dealer or Other Stranger	176	129	15	10	81	52	80	67
Bought on the Internet	22	11	*	*	5	*	16	2
Some Other Way	64	37	10	8	22	8	33	20

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.50B Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Pain Relievers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	79.2 ^b	85.3	62.3	69.2	74.5 ^b	82.4	83.3	88.0
From More Than One Doctor	7.4	6.0	12.9	9.3	11.2	7.1	5.1	5.3
Wrote Fake Prescription	1.9	0.7	0.9	*	0.5	1.0	2.6	0.5
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.2	1.0	1.8	*	1.2	1.7	1.1	0.6
From Friend or Relative for Free	13.5	10.6	14.5	16.5	15.7	14.9	12.5	8.4
Bought from Friend or Relative	12.3	9.7	13.9	12.5	17.2	14.9	9.9	7.5
Took from Friend or Relative without Asking	4.2	3.1	9.8	6.8	6.8	3.9	2.4	2.4
Bought from Drug Dealer or Other Stranger	7.5	7.0	8.6	8.0	12.0	11.3	5.3	5.3
Bought on the Internet	0.9	0.6	*	*	0.8	*	1.1	0.1
Some Other Way	2.8	2.0	5.7	7.0	3.3	1.9	2.2	1.6

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.51A Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	659	628	29	27	111	112	519	489
From More Than One Doctor	38	56	3	6	18	15	17	35
Wrote Fake Prescription	3	6	2	*	1	1	*	3
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	11	22	1	1	6	6	4	15
From Friend or Relative for Free	3,157	2,930	189 ^a	143	779	729	2,190	2,058
Bought from Friend or Relative	708 ^a	574	67	58	309	264	332	253
Took from Friend or Relative without Asking	159	205	38	30	65	74	56	102
Bought from Drug Dealer or Other Stranger	303	261	35	25	166	133	102	102
Bought on the Internet	1 ^a	10	*	2	1	3	*	6
Some Other Way	147	152	13	18	52	52	82	81

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.51B Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	12.7	13.0	7.6	8.7	7.3	8.1	15.7	15.6
From More Than One Doctor	0.7	1.1	0.8	1.8	1.2	1.1	0.5	1.1
Wrote Fake Prescription	0.1	0.1	0.6	*	0.0	0.1	*	0.1
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2	0.5	0.2	0.3	0.4	0.4	0.1	0.5
From Friend or Relative for Free	60.9	60.5	50.0	46.0	51.7	52.5	66.3	65.5
Bought from Friend or Relative	13.7	11.9	17.7	18.5	20.5	19.0	10.1	8.1
Took from Friend or Relative without Asking	3.1	4.2	10.0	9.6	4.3	5.3	1.7	3.2
Bought from Drug Dealer or Other Stranger	5.8	5.4	9.4	8.2	11.0	9.6	3.1	3.3
Bought on the Internet	0.0 ^a	0.2	*	0.6	0.0	0.2	*	0.2
Some Other Way	2.8	3.1	3.6	5.9	3.4	3.8	2.5	2.6

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.52A Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Tranquilizers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	2,340	2,213	74	65	500	469	1,765	1,679
From More Than One Doctor	95	67	7	6	29	17	59	44
Wrote Fake Prescription	1	2	*	1	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	10	3	1	*	3	2	6	1
From Friend or Relative for Free	209	189	24	19	68	60	117	110
Bought from Friend or Relative	122	88	21 ^a	8	41	45	60	36
Took from Friend or Relative without Asking	33 ^a	13	8	5	13	4	11	4
Bought from Drug Dealer or Other Stranger	43 ^a	80	11	8	21 ^a	41	11	31
Bought on the Internet	1	14	*	*	*	5	*	8
Some Other Way	25	23	3	2	15	6	8	16

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.52B Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Tranquilizers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	81.3	82.2	49.6	57.6	72.2	72.2	86.7	87.0
From More Than One Doctor	3.3	2.5	4.5	4.9	4.2	2.7	2.9	2.3
Wrote Fake Prescription	0.0	0.1	*	0.6	0.2	0.2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3	0.1	0.5	*	0.5	0.2	0.3	0.1
From Friend or Relative for Free	7.2	7.0	15.8	16.8	9.8	9.2	5.7	5.7
Bought from Friend or Relative	4.2	3.3	13.9	6.9	5.9	6.9	2.9	1.9
Took from Friend or Relative without Asking	1.1 ^a	0.5	5.6	4.2	1.9	0.7	0.5	0.2
Bought from Drug Dealer or Other Stranger	1.5 ^b	3.0	7.5	7.5	3.1 ^a	6.3	0.5	1.6
Bought on the Internet	0.0	0.5	*	*	*	0.8	*	0.4
Some Other Way	0.9	0.9	2.0	1.5	2.1	0.9	0.4	0.8

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.53A Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	353	307	12	*	57	40	285	252
From More Than One Doctor	15	21	2	3	10	8	3	9
Wrote Fake Prescription	3	2	2	*	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	21	7	*	*	4	4	*	4
From Friend or Relative for Free	1,092	947	60	*	250	208	783	706
Bought from Friend or Relative	349	279	28	*	124	111	198	149
Took from Friend or Relative without Asking	73	110	12	10	31	23	30 ^a	76
Bought from Drug Dealer or Other Stranger	156	133	18	*	74 ^a	43	64	79
Bought on the Internet	*	12	*	*	0	*	*	11
Some Other Way	57	43	7	3	16	7	34	33

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.53B Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	18.8	18.5	9.4	*	11.5	10.1	22.6	21.3
From More Than One Doctor	0.8	1.2	1.7	4.2	2.0	2.1	0.2	0.8
Wrote Fake Prescription	0.2	0.1	1.3	*	0.3	0.2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.1	0.4	*	*	0.9	0.9	*	0.3
From Friend or Relative for Free	58.3	57.2	48.1	*	50.6	52.9	62.3	59.7
Bought from Friend or Relative	18.6	16.8	22.2	*	25.1	28.2	15.7	12.6
Took from Friend or Relative without Asking	3.9	6.6	9.3	13.1	6.4	5.9	2.4 ^a	6.5
Bought from Drug Dealer or Other Stranger	8.3	8.1	14.8	*	15.0	11.0	5.1	6.6
Bought on the Internet	*	0.7	*	*	0.0	*	*	0.9
Some Other Way	3.1	2.6	5.8	4.1	3.3	1.8	2.7	2.8

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.54A Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Tranquilizers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	773	708	*	*	162	134	586	561
From More Than One Doctor	66	37	*	*	10	9	54	25
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	7	8	*	*	3	*	*	8
From Friend or Relative for Free	92	104	*	*	29	27	52	68
Bought from Friend or Relative	56	86	*	*	27	34	17 ^a	51
Took from Friend or Relative without Asking	22	9	*	*	8	1	11	7
Bought from Drug Dealer or Other Stranger	23 ^a	58	*	*	14	11	5 ^b	45
Bought on the Internet	1	17	*	*	*	*	*	13
Some Other Way	16	13	*	*	*	3	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.54B Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Tranquilizers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	76.7	79.3	*	*	70.1	69.3	80.5	83.5
From More Than One Doctor	6.6	4.2	*	*	4.2	4.6	7.4	3.8
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7	0.9	*	*	1.3	*	*	1.1
From Friend or Relative for Free	9.2	11.6	*	*	12.5	14.1	7.2	10.2
Bought from Friend or Relative	5.6 ^a	9.7	*	*	11.7	17.4	2.4 ^a	7.6
Took from Friend or Relative without Asking	2.2	1.0	*	*	3.4	0.7	1.5	1.1
Bought from Drug Dealer or Other Stranger	2.3 ^b	6.5	*	*	6.2	5.9	0.7 ^b	6.7
Bought on the Internet	0.1	1.9	*	*	*	*	*	1.9
Some Other Way	1.6	1.4	*	*	*	1.7	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.55A Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Stimulant Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	241	290	23	35	64	68	153	187
From More Than One Doctor	27	20	3	5	19	11	6	4
Wrote Fake Prescription	3	7	*	1	1	3	*	3
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	10	3	2	*	5	1	3	1
From Friend or Relative for Free	1,151 ^a	1,387	108	96	497	548	546 ^a	743
Bought from Friend or Relative	477	573	42	45	284	318	152	210
Took from Friend or Relative without Asking	71 ^a	130	19	23	19 ^a	38	34	69
Bought from Drug Dealer or Other Stranger	136	121	16	15	77	55	43	51
Bought on the Internet	11	30	*	1	3 ^a	15	8	15
Some Other Way	59	58	4	9	30	32	25	17

*Low precision; no estimate reported.

NOTE: Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.55B Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Stimulant Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	11.0	11.1	10.7	15.2	6.5	6.2	15.8	14.4
From More Than One Doctor	1.2	0.8	1.2	2.1	1.9	1.0	0.6	0.3
Wrote Fake Prescription	0.1	0.3	*	0.5	0.1	0.3	*	0.2
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4	0.1	0.9	*	0.5	0.1	0.3	0.1
From Friend or Relative for Free	52.7	53.0	49.6	41.8	49.8	50.3	56.3	57.2
Bought from Friend or Relative	21.8	21.9	19.1	19.7	28.4	29.2	15.7	16.2
Took from Friend or Relative without Asking	3.3	5.0	8.6	10.0	1.9	3.5	3.5	5.3
Bought from Drug Dealer or Other Stranger	6.2	4.6	7.4	6.3	7.7 ^a	5.1	4.4	3.9
Bought on the Internet	0.5	1.1	*	0.3	0.3 ^a	1.4	0.8	1.1
Some Other Way	2.7	2.2	1.8	4.1	3.0	2.9	2.6	1.3

*Low precision; no estimate reported.

NOTE: Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.56A Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Stimulants	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	846	948	54	59	346	391	446	498
From More Than One Doctor	33	42	7	3	22	14	4	24
Wrote Fake Prescription	*	0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	0	*	0	2	0	*	*
From Friend or Relative for Free	76 ^a	126	12	*	35	35	29 ^a	76
Bought from Friend or Relative	55	66	6	2	35	31	14	33
Took from Friend or Relative without Asking	12	8	5	2	7	2	*	4
Bought from Drug Dealer or Other Stranger	23	47	5	2	15	16	3 ^a	30
Bought on the Internet	3	3	1	*	0	1	*	*
Some Other Way	3	5	1	1	2	2	0	2

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.56B Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Stimulants	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	80.3	76.1	59.9	69.2	74.6	79.3	89.4 ^b	74.6
From More Than One Doctor	3.1	3.4	7.7	3.5	4.8	2.9	0.8	3.6
Wrote Fake Prescription	*	0.0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2	0.0	*	0.1	0.4	0.1	*	*
From Friend or Relative for Free	7.2	10.1	12.8	*	7.6	7.0	5.9	11.4
Bought from Friend or Relative	5.2	5.3	6.5	1.8	7.5	6.3	2.8	5.0
Took from Friend or Relative without Asking	1.1	0.6	5.4	1.9	1.4	0.4	*	0.6
Bought from Drug Dealer or Other Stranger	2.1	3.8	5.2	1.8	3.3	3.3	0.6 ^a	4.5
Bought on the Internet	0.3	0.2	1.1	*	0.1	0.3	*	*
Some Other Way	0.3	0.4	1.4	1.6	0.4	0.4	0.0	0.3

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.57A Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Stimulant Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	140	158	13	20	25	34	*	103
From More Than One Doctor	19	11	3	3	4	8	*	*
Wrote Fake Prescription	*	1	*	0	0	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	12	3	1	0	3	2	*	1
From Friend or Relative for Free	356 ^b	548	39	39	163	187	*	322
Bought from Friend or Relative	236	297	23	19	115	130	*	148
Took from Friend or Relative without Asking	45	54	7	7	18	19	*	28
Bought from Drug Dealer or Other Stranger	103	70	11	12	44	29	47	29
Bought on the Internet	18	18	*	*	2	7	*	11
Some Other Way	39	31	3	2	12	15	25	14

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.57B Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Stimulant Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	18.9	15.9	17.7	24.0	8.3	9.8	*	18.4
From More Than One Doctor	2.5	1.1	4.2	3.0	1.4	2.3	*	*
Wrote Fake Prescription	*	0.1	*	0.3	0.0	0.3	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.6	0.3	1.3	0.3	0.9	0.5	*	0.2
From Friend or Relative for Free	47.9	55.2	52.3	46.0	53.5	53.9	*	57.3
Bought from Friend or Relative	31.9	30.1	31.4	23.3	38.1	37.3	*	26.6
Took from Friend or Relative without Asking	6.1	5.5	9.8	8.3	6.1	5.5	*	5.0
Bought from Drug Dealer or Other Stranger	13.9 ^b	7.1	15.3	14.1	14.5 ^a	8.4	13.0	5.2
Bought on the Internet	2.5	1.8	*	*	0.7	2.1	*	1.9
Some Other Way	5.4	3.3	3.7	2.5	4.0	4.4	6.8	2.6

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.58A Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Stimulants	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	271	359	*	*	131	151	*	*
From More Than One Doctor	19	24	*	*	4	7	*	*
Wrote Fake Prescription	*	*	*	*	*	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0	*	*	*	*	*	*
From Friend or Relative for Free	33 ^a	73	*	*	13	14	*	53
Bought from Friend or Relative	31	51	*	*	11	12	*	*
Took from Friend or Relative without Asking	17	10	*	*	5	*	*	7
Bought from Drug Dealer or Other Stranger	36	43	*	*	8	6	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	3	6	*	*	1	3	*	*

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.58B Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Stimulants	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	83.4 ^a	71.1	*	*	86.2	87.5	*	*
From More Than One Doctor	5.9	4.8	*	*	2.8	4.1	*	*
Wrote Fake Prescription	*	*	*	*	*	0.1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0.0	*	*	*	*	*	*
From Friend or Relative for Free	10.2	14.6	*	*	8.5	8.1	*	17.8
Bought from Friend or Relative	9.6	10.1	*	*	7.1	6.7	*	*
Took from Friend or Relative without Asking	5.4	2.1	*	*	3.5	*	*	2.4
Bought from Drug Dealer or Other Stranger	11.1	8.5	*	*	5.4	3.7	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	1.0	1.3	*	*	0.6	1.9	*	*

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.59A Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	407	425	39	*	129	130	239	265
Bought from Friend or Relative	228	222	*	*	36	53	183	161
Took from Friend or Relative without Asking	23	43	2	*	11	11	10	*
Bought from Drug Dealer or Other Stranger	223 ^a	338	*	*	66	73	137 ^a	253
Bought on the Internet	4	7	*	*	3	2	*	4
Some Other Way	52	45	*	*	9	13	*	28

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.59B Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	43.4	39.4	52.2	*	50.7	46.0	39.3	36.1
Bought from Friend or Relative	24.3	20.6	*	*	14.1	18.9	30.1	21.9
Took from Friend or Relative without Asking	2.4	4.0	2.3	*	4.5	3.8	1.6	*
Bought from Drug Dealer or Other Stranger	23.8	31.3	*	*	26.0	26.0	22.5	34.4
Bought on the Internet	0.5	0.6	*	*	1.2	0.8	*	0.6
Some Other Way	5.5	4.2	*	*	3.5	4.4	*	3.8

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.60A Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Methamphetamine	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	118	92	*	*	37	38	*	*
Bought from Friend or Relative	114	128	*	*	*	34	*	*
Took from Friend or Relative without Asking	3	5	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	52	89	*	*	26	33	*	*
Bought on the Internet	2	*	*	*	2	*	*	*
Some Other Way	38	23	*	*	9	*	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.60B Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Methamphetamine	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	36.2	27.3	*	*	31.3	33.1	*	*
Bought from Friend or Relative	35.0	38.0	*	*	*	29.3	*	*
Took from Friend or Relative without Asking	0.8	1.6	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	15.8 ^a	26.3	*	*	22.3	28.3	*	*
Bought on the Internet	0.5	*	*	*	1.5	*	*	*
Some Other Way	11.6	6.8	*	*	8.1	*	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.61A Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	*	213	*	*	*	45	*	153
Bought from Friend or Relative	*	171	*	*	*	*	*	*
Took from Friend or Relative without Asking	10	*	*	*	*	6	*	*
Bought from Drug Dealer or Other Stranger	87 ^a	161	*	*	*	27	*	131
Bought on the Internet	4	*	*	*	*	1	*	*
Some Other Way	*	*	*	*	*	5	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.61B Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	*	43.6	*	*	*	52.9	*	40.1
Bought from Friend or Relative	*	35.0	*	*	*	*	*	*
Took from Friend or Relative without Asking	2.8	*	*	*	*	7.6	*	*
Bought from Drug Dealer or Other Stranger	23.4	32.9	*	*	*	31.6	*	34.2
Bought on the Internet	1.0	*	*	*	*	0.7	*	*
Some Other Way	*	*	*	*	*	5.6	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.62A Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Methamphetamine	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	4	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.62B Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Methamphetamine	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	2.1	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.63A Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Sedative Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	122	130	15	*	14	25	*	92
From More Than One Doctor	9	11	*	*	3	*	*	4
Wrote Fake Prescription	1	*	*	*	0	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	*	3	*	2	*	*	*
From Friend or Relative for Free	271	330	26	*	64	81	*	224
Bought from Friend or Relative	41	59	7	*	13	13	*	44
Took from Friend or Relative without Asking	17	25	5	*	*	*	*	9
Bought from Drug Dealer or Other Stranger	12	19	2	*	8	3	*	12
Bought on the Internet	0	1	0	*	0	*	*	*
Some Other Way	18	19	0	*	4	0	*	17

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.63B Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Sedative Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	24.6	20.7	24.2	*	12.3	18.5	*	21.2
From More Than One Doctor	1.8	1.7	*	*	2.2	*	*	0.9
Wrote Fake Prescription	0.3	*	*	*	0.2	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.0	*	5.3	*	1.4	*	*	*
From Friend or Relative for Free	54.5	52.4	40.8	*	55.7	61.3	*	51.6
Bought from Friend or Relative	8.2	9.4	10.7	*	11.4	9.5	*	10.0
Took from Friend or Relative without Asking	3.4	3.9	8.4	*	*	*	*	2.2
Bought from Drug Dealer or Other Stranger	2.5	3.0	3.5	*	6.7	2.5	*	2.8
Bought on the Internet	0.0	0.1	0.1	*	0.1	*	*	*
Some Other Way	3.7	3.1	0.7	*	3.9	0.1	*	4.0

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.64A Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Sedatives	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	227	245	*	*	*	*	*	*
From More Than One Doctor	7	7	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	4	*	*	*	*	*	*
From Friend or Relative for Free	12	30	*	*	*	*	*	*
Bought from Friend or Relative	3	1	*	*	*	*	*	*
Took from Friend or Relative without Asking	6	8	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	5	6	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	1	2	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.64B Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Sedatives	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	86.8	79.4	*	*	*	*	*	*
From More Than One Doctor	2.8	2.3	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	1.4	*	*	*	*	*	*
From Friend or Relative for Free	4.6	9.8	*	*	*	*	*	*
Bought from Friend or Relative	1.0	0.4	*	*	*	*	*	*
Took from Friend or Relative without Asking	2.2	2.6	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	1.9	2.1	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	0.4	0.8	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.65A Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Sedative Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	7	*	*	*	*	*	*
Wrote Fake Prescription	1	0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	10	9	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	8	*	*	*	*	*	*
Bought on the Internet	0	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.65B Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Sedative Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	2.7	*	*	*	*	*	*
Wrote Fake Prescription	0.5	0.2	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.1	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	5.0	3.3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	3.2	*	*	*	*	*	*
Bought on the Internet	0.0	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.66A Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Sedatives	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.66B Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Sedatives	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.67A Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Numbers in Thousands, 2013 and 2014

Social Context and Location of Most Recent Alcohol Use	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	8,651	8,661	4,482	4,424	4,170	4,236
SOCIAL CONTEXT OF MOST RECENT ALCOHOL USE²						
Drank Alone	512	529	336	290	176 ^a	239
Drank with One Other Person	1,369	1,414	702	760	667	654
Drank with Two or More Other People	6,506	6,439	3,292	3,213	3,215	3,225
LOCATION OF MOST RECENT ALCOHOL USE^{2,3}						
Car or Other Vehicle	286	291	150	131	136	160
Home	2,837	2,757	1,479	1,461	1,358	1,296
Someone Else's Home	4,328	4,415	2,187	2,183	2,141	2,231
Park, Beach, or Parking Lot	354	308	170	130	184	178
Restaurant, Bar, or Club	546	599	190	251	356	349
Concert or Sports Game	169	192	91	113	77	79
School	158	206	80	84	78	122
Some Other Place ⁴	448	490	244 ^a	339	204	152
Party, Wedding, or Celebration	130	178	75	115	55	64
Outside (Location not Specified)	53	65	25	50	28	14
Hotel, Motel, or Resort	60	47	23	33	37	15
Camping, Hunting, or Fishing	34	42	25	29	9	13
Cabin, Cottage, Vacation Home, etc.	13	8	5	4	8	4
Dorm Room	13	14	3	11	10	3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents could indicate multiple locations for the most recent time they used alcohol; thus, these response categories are not mutually exclusive.

⁴ Some Other Place includes only valid responses from the other-specify questions, including these six types of commonly reported locations.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.67B Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Percentages, 2013 and 2014

Social Context and Location of Most Recent Alcohol Use	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	100.0	100.0	100.0	100.0	100.0	100.0
SOCIAL CONTEXT OF MOST RECENT ALCOHOL USE²						
Drank Alone	6.1	6.3	7.8	6.8	4.3	5.8
Drank with One Other Person	16.3	16.9	16.2	17.8	16.4	15.9
Drank with Two or More Other People	77.6	76.8	76.0	75.4	79.2	78.3
LOCATION OF MOST RECENT ALCOHOL USE^{2,3}						
Car or Other Vehicle	3.4	3.5	3.5	3.1	3.4	3.9
Home	34.2	33.3	34.6	34.7	33.7	31.9
Someone Else's Home	52.2	53.4	51.2	51.9	53.2	55.0
Park, Beach, or Parking Lot	4.3	3.7	4.0	3.1	4.6	4.4
Restaurant, Bar, or Club	6.6	7.2	4.5	6.0	8.8	8.6
Concert or Sports Game	2.0	2.3	2.1	2.7	1.9	1.9
School	1.9	2.5	1.9	2.0	1.9	3.0
Some Other Place ⁴	5.5	6.0	5.8 ^a	8.2	5.1	3.8
Party, Wedding, or Celebration	1.6	2.2	1.8	2.8	1.4	1.6
Outside (Location not Specified)	0.6	0.8	0.6	1.2	0.7	0.4
Hotel, Motel, or Resort	0.7	0.6	0.6	0.8	0.9	0.4
Camping, Hunting, or Fishing	0.4	0.5	0.6	0.7	0.2	0.3
Cabin, Cottage, Vacation Home, etc.	0.2	0.1	0.1	0.1	0.2	0.1
Dorm Room	0.2	0.2	0.1	0.3	0.2	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents could indicate multiple locations for the most recent time they used alcohol; thus, these response categories are not mutually exclusive.

⁴ Some Other Place includes only valid responses from the other-specify questions, including these six types of commonly reported locations.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.68A Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Numbers in Thousands, 2013 and 2014

Social Context and Location of Most Recent Alcohol Use	Total (2013)	Total (2014)	Aged 12-14 (2013)	Aged 12-14 (2014)	Aged 15-17 (2013)	Aged 15-17 (2014)	Aged 18-20 (2013)	Aged 18-20 (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	8,651	8,661	473 ^a	388	2,404	2,484	5,774	5,788
SOCIAL CONTEXT OF MOST RECENT ALCOHOL USE²								
Drank Alone	512	529	62	38	180	173	270	318
Drank with One Other Person	1,369	1,414	80	82	349	354	939	978
Drank with Two or More Other People	6,506	6,439	287	225	1,774	1,852	4,445	4,361
LOCATION OF MOST RECENT ALCOHOL USE^{2,3}								
Car or Other Vehicle	286	291	24 ^a	8	103	93	159	190
Home	2,837	2,757	177	154	624	665	2,036	1,938
Someone Else's Home	4,328	4,415	174	161	1,353	1,377	2,802	2,876
Park, Beach, or Parking Lot	354	308	25	12	143	103	186	193
Restaurant, Bar, or Club	546	599	6	5	49 ^a	82	491	513
Concert or Sports Game	169	192	12	5	37	46	120	142
School	158	206	9	10	20	32	130	165
Some Other Place ⁴	448	490	24 ^a	11	165	186	259	293
Party, Wedding, or Celebration	130	178	8	4	60	84	61	90
Outside (Location not Specified)	53	65	3	2	37	28	12	35
Hotel, Motel, or Resort	60	47	*	*	13	12	47	35
Camping, Hunting, or Fishing	34	42	3	1	13	9	18	33
Cabin, Cottage, Vacation Home, etc.	13	8	0	1	4	7	9	0
Dorm Room	13	14	*	*	*	5	13	9

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents could indicate multiple locations for the most recent time they used alcohol; thus, these response categories are not mutually exclusive.

⁴ Some Other Place includes only valid responses from the other-specify questions, including these six types of commonly reported locations.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.68B Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Percentages, 2013 and 2014

Social Context and Location of Most Recent Alcohol Use	Total (2013)	Total (2014)	Aged 12-14 (2013)	Aged 12-14 (2014)	Aged 15-17 (2013)	Aged 15-17 (2014)	Aged 18-20 (2013)	Aged 18-20 (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
SOCIAL CONTEXT OF MOST RECENT ALCOHOL USE²								
Drank Alone	6.1	6.3	14.5	10.9	7.8	7.3	4.8	5.6
Drank with One Other Person	16.3	16.9	18.7	23.7	15.2	14.9	16.6	17.3
Drank with Two or More Other People	77.6	76.8	66.8	65.3	77.0	77.9	78.6	77.1
LOCATION OF MOST RECENT ALCOHOL USE^{2,3}								
Car or Other Vehicle	3.4	3.5	5.8	2.3	4.5	4.0	2.8	3.4
Home	34.2	33.3	42.4	46.5	27.5	28.5	36.3	34.6
Someone Else's Home	52.2	53.4	41.7	48.7	59.5	59.0	50.0	51.3
Park, Beach, or Parking Lot	4.3	3.7	5.9	3.7	6.3 ^a	4.4	3.3	3.4
Restaurant, Bar, or Club	6.6	7.2	1.4	1.5	2.2 ^a	3.5	8.8	9.1
Concert or Sports Game	2.0	2.3	2.9	1.4	1.6	2.0	2.1	2.5
School	1.9	2.5	2.0	2.9	0.9	1.4	2.3	2.9
Some Other Place ⁴	5.5	6.0	6.0	3.5	7.4	8.1	4.7	5.3
Party, Wedding, or Celebration	1.6	2.2	2.1	1.3	2.7	3.7	1.1	1.6
Outside (Location not Specified)	0.6	0.8	0.8	0.7	1.6	1.2	0.2	0.6
Hotel, Motel, or Resort	0.7	0.6	*	*	0.6	0.5	0.8	0.6
Camping, Hunting, or Fishing	0.4	0.5	0.7	0.3	0.6	0.4	0.3	0.6
Cabin, Cottage, Vacation Home, etc.	0.2	0.1	0.1	0.3	0.2	0.3	0.2	0.0
Dorm Room	0.2	0.2	*	*	*	0.2	0.2	0.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents could indicate multiple locations for the most recent time they used alcohol; thus, these response categories are not mutually exclusive.

⁴ Some Other Place includes only valid responses from the other-specify questions, including these six types of commonly reported locations.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.69A Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Numbers in Thousands, 2013 and 2014

Source of Most Recent Alcohol Use in the Past Month	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	8,651	8,661	4,482	4,424	4,170	4,236
UNDERAGE DRINKER PAID²	2,403	2,331	1,536	1,381	867	951
Purchased It Himself or Herself	653	702	457	429	197 ^a	273
From Store, Restaurant, Bar, Club, or Event	562	582	390	351	172	231
Liquor, Convenience, or Grocery Store	394	383	301	257	92	126
Restaurant, Bar, or Club	157	186	79	87	78	99
Concert, Sports, or Other Event	11	12	10	6	1	5
From Another Person	86	118	61	76	25	42
From Person under Age 21	30	28	22	12	8	16
From Person Aged 21 or Older	54	89	39	64	14	25
Purchased by Someone Else	1,712	1,609	1,050	939	663	670
Parent or Guardian	64	88	38	46	27	42
Another Family Member Aged 21 or Older	171	168	93	101	78	68
Someone Not Related Aged 21 or Older	1,243	1,130	767	653	475	478
Someone under Age 21	164	174	102	106	61	68
UNDERAGE DRINKER DID NOT PAY²	5,982	6,055	2,796	2,890	3,186	3,165
Got It from Parent or Guardian	635	682	276	314	359	368
Got It from Another Family Member Aged 21 or Older	829	890	343	428	487	461
Got It from Someone Not Related Aged 21 or Older	2,189	2,091	976	894	1,213	1,197
Got It from Someone under Age 21	984	1,017	459	487	524	530
Took It from Own Home	469 ^a	351	256	190	214	160
Took It from Someone Else's Home	175	215	98	103	77	112
Got It Some Other Way	360	409	211	235	149	173
From Friend or Acquaintance, Unspecified Age and Method ³	160	171	91	89	69	82

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents were permitted to specify other sources for most recent alcohol use in the past month. This source is one of the most commonly reported other sources for most recent alcohol use in the past month.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.69B Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Percentages, 2013 and 2014

Source of Most Recent Alcohol Use in the Past Month	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	100.0	100.0	100.0	100.0	100.0	100.0
UNDERAGE DRINKER PAID²	28.7	27.8	35.5	32.3	21.4	23.1
Purchased It Himself or Herself	7.8	8.4	10.6	10.1	4.9 ^a	6.6
From Store, Restaurant, Bar, Club, or Event	6.7	7.0	9.1	8.3	4.2	5.6
Liquor, Convenience, or Grocery Store	4.7	4.6	7.0	6.0	2.3	3.1
Restaurant, Bar, or Club	1.9	2.2	1.8	2.0	1.9	2.4
Concert, Sports, or Other Event	0.1	0.1	0.2	0.1	0.0	0.1
From Another Person	1.0	1.4	1.4	1.8	0.6	1.0
From Person under Age 21	0.4	0.3	0.5	0.3	0.2	0.4
From Person Aged 21 or Older	0.6	1.1	0.9	1.5	0.4	0.6
Purchased by Someone Else	20.5	19.2	24.4	22.0	16.4	16.3
Parent or Guardian	0.8	1.1	0.9	1.1	0.7	1.0
Another Family Member Aged 21 or Older	2.1	2.0	2.2	2.4	1.9	1.7
Someone Not Related Aged 21 or Older	15.0	13.6	18.0	15.4	11.8	11.7
Someone under Age 21	2.0	2.1	2.4	2.5	1.5	1.7
UNDERAGE DRINKER DID NOT PAY²	71.3	72.2	64.5	67.7	78.6	76.9
Got It from Parent or Guardian	7.9	8.5	6.6	7.8	9.2	9.3
Got It from Another Family Member Aged 21 or Older	10.3	11.1	8.2 ^a	10.6	12.5	11.7
Got It from Someone Not Related Aged 21 or Older	27.2	26.2	23.5	22.2	31.2	30.3
Got It from Someone under Age 21	12.2	12.7	11.1	12.1	13.5	13.4
Took It from Own Home	5.8 ^a	4.4	6.2	4.7	5.5 ^a	4.1
Took It from Someone Else's Home	2.2	2.7	2.4	2.6	2.0	2.8
Got It Some Other Way	4.5	5.1	5.1	5.8	3.8	4.4
From Friend or Acquaintance, Unspecified Age and Method ³	2.0	2.1	2.2	2.2	1.8	2.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents were permitted to specify other sources for most recent alcohol use in the past month. This source is one of the most commonly reported other sources for most recent alcohol use in the past month.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.70A Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Numbers in Thousands, 2013 and 2014

Source of Most Recent Alcohol Use in the Past Month	Total (2013)	Total (2014)	Aged 12-14 (2013)	Aged 12-14 (2014)	Aged 15-17 (2013)	Aged 15-17 (2014)	Aged 18-20 (2013)	Aged 18-20 (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	8,651	8,661	473 ^a	388	2,404	2,484	5,774	5,788
UNDERAGE DRINKER PAID²	2,403	2,331	27	23	478	423	1,899	1,886
Purchased It Himself or Herself	653	702	9	*	93	93	551	603
From Store, Restaurant, Bar, Club, or Event	562	582	4	*	64	66	494	511
Liquor, Convenience, or Grocery Store	394	383	3	*	55	54	336	323
Restaurant, Bar, or Club	157	186	1	*	7	11	149	175
Concert, Sports, or Other Event	11	12	*	*	2	1	9	11
From Another Person	86	118	5	*	29	25	52	92
From Person under Age 21	30	28	3	*	14	7	14	21
From Person Aged 21 or Older	54	89	2	*	13	18	38	72
Purchased by Someone Else	1,712	1,609	17	16	375	322	1,321	1,271
Parent or Guardian	64	88	*	*	9	10	55	77
Another Family Member Aged 21 or Older	171	168	2	1	35	21	134	146
Someone Not Related Aged 21 or Older	1,243	1,130	13	12	253	203	977	915
Someone under Age 21	164	174	2	*	62	72	100	101
UNDERAGE DRINKER DID NOT PAY²	5,982	6,055	398	324	1,823	1,950	3,761	3,780
Got It from Parent or Guardian	635	682	65	55	185	243	385	384
Got It from Another Family Member Aged 21 or Older	829	890	65	68	217	249	547	572
Got It from Someone Not Related Aged 21 or Older	2,189	2,091	52	38	435	442	1,702	1,610
Got It from Someone under Age 21	984	1,017	63	36	456	471	465	511
Took It from Own Home	469 ^a	351	69 ^b	33	210	165	190	153
Took It from Someone Else's Home	175	215	19	19	80	91	76	106
Got It Some Other Way	360	409	31	36	138	180	191	193
From Friend or Acquaintance, Unspecified Age and Method ³	160	171	12	19	69	99	79	52

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents were permitted to specify other sources for most recent alcohol use in the past month. This source is one of the most commonly reported other sources for most recent alcohol use in the past month.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.70B Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Percentages, 2013 and 2014

Source of Most Recent Alcohol Use in the Past Month	Total (2013)	Total (2014)	Aged 12-14 (2013)	Aged 12-14 (2014)	Aged 15-17 (2013)	Aged 15-17 (2014)	Aged 18-20 (2013)	Aged 18-20 (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
UNDERAGE DRINKER PAID²	28.7	27.8	6.3	6.5	20.8	17.8	33.6	33.3
Purchased It Himself or Herself	7.8	8.4	2.1	*	4.1	3.9	9.8	10.7
From Store, Restaurant, Bar, Club, or Event	6.7	7.0	0.9	*	2.8	2.8	8.8	9.0
Liquor, Convenience, or Grocery Store	4.7	4.6	0.7	*	2.4	2.3	6.0	5.7
Restaurant, Bar, or Club	1.9	2.2	0.2	*	0.3	0.5	2.7	3.1
Concert, Sports, or Other Event	0.1	0.1	*	*	0.1	0.0	0.2	0.2
From Another Person	1.0	1.4	1.2	*	1.3	1.1	0.9	1.6
From Person under Age 21	0.4	0.3	0.7	*	0.6	0.3	0.2	0.4
From Person Aged 21 or Older	0.6	1.1	0.6	*	0.6	0.7	0.7	1.3
Purchased by Someone Else	20.5	19.2	4.0	4.6	16.4	13.6	23.5	22.5
Parent or Guardian	0.8	1.1	*	*	0.4	0.4	1.0	1.4
Another Family Member Aged 21 or Older	2.1	2.0	0.4	0.3	1.6	0.9	2.4	2.6
Someone Not Related Aged 21 or Older	15.0	13.6	3.0	3.5	11.1	8.7	17.5	16.3
Someone under Age 21	2.0	2.1	0.4	*	2.7	3.1	1.8	1.8
UNDERAGE DRINKER DID NOT PAY²	71.3	72.2	93.7	93.5	79.2	82.2	66.4	66.7
Got It from Parent or Guardian	7.9	8.5	16.7	17.9	8.4	10.7	7.1	7.1
Got It from Another Family Member Aged 21 or Older	10.3	11.1	16.6	22.1	9.9	11.0	10.0	10.6
Got It from Someone Not Related Aged 21 or Older	27.2	26.2	13.2	12.3	19.8	19.5	31.2	29.7
Got It from Someone under Age 21	12.2	12.7	16.2	11.8	20.7	20.8	8.5	9.4
Took It from Own Home	5.8 ^a	4.4	17.7 ^a	10.6	9.6	7.3	3.5	2.8
Took It from Someone Else's Home	2.2	2.7	4.8	6.2	3.6	4.0	1.4	1.9
Got It Some Other Way	4.5	5.1	8.0	11.7	6.3	7.9	3.5	3.6
From Friend or Acquaintance, Unspecified Age and Method ³	2.0	2.1	3.1	6.3	3.1	4.4	1.4	1.0

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents were permitted to specify other sources for most recent alcohol use in the past month. This source is one of the most commonly reported other sources for most recent alcohol use in the past month.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.71A Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
ILLCIT DRUGS²	6,574 ^b	7,527	130	123	6,415 ^b	7,367
Marijuana and Hashish	5,122 ^b	5,967	115	96	4,985 ^b	5,844
Cocaine	344	389	4	7	338	381
Crack	47	73	*	2	45	71
Heroin	83	109	4	5	79	100
Hallucinogens	335	374	2	4	330	364
LSD	37 ^a	79	*	*	37 ^a	79
PCP	*	13	0	*	*	13
Ecstasy	226	228	1	3	225	224
Inhalants	102	102	1	8	101	94
Nonmedical Use of Psychotherapeutics ^{3,4}	2,162	2,268	33	30	2,116	2,222
Pain Relievers	1,473	1,432	19	25	1,442	1,396
OxyContin [®]	93	98	3	2	88	96
Tranquilizers	737	665	7	11	728	643
Stimulants ⁴	383 ^b	549	9	3	374 ^b	539
Methamphetamine ⁴	145	138	6	3	138	128
Sedatives	72	98	0	1	72	96
ILLCIT DRUGS OTHER THAN MARIJUANA²	2,599	2,803	38	47	2,547	2,736

*Low precision; no estimate reported.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.71B Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
ILLCIT DRUGS²	10.5 ^b	11.9	5.9	5.3	10.7 ^b	12.1
Marijuana and Hashish	8.2 ^b	9.4	5.2	4.1	8.3 ^b	9.6
Cocaine	0.6	0.6	0.2	0.3	0.6	0.6
Crack	0.1	0.1	*	0.1	0.1	0.1
Heroin	0.1	0.2	0.2	0.2	0.1	0.2
Hallucinogens	0.5	0.6	0.1	0.2	0.5	0.6
LSD	0.1 ^a	0.1	*	*	0.1 ^a	0.1
PCP	*	0.0	0.0	*	*	0.0
Ecstasy	0.4	0.4	0.0	0.1	0.4	0.4
Inhalants	0.2	0.2	0.0	0.3	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	3.5	3.6	1.5	1.3	3.5	3.7
Pain Relievers	2.4	2.3	0.9	1.1	2.4	2.3
OxyContin [®]	0.1	0.2	0.1	0.1	0.1	0.2
Tranquilizers	1.2	1.0	0.3	0.5	1.2	1.1
Stimulants ⁴	0.6 ^b	0.9	0.4	0.1	0.6 ^b	0.9
Methamphetamine ⁴	0.2	0.2	0.3	0.1	0.2	0.2
Sedatives	0.1	0.2	0.0	0.1	0.1	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA²	4.2	4.4	1.7	2.0	4.2	4.5

*Low precision; no estimate reported.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.72A Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	6,574 ^b	7,527	130	123	6,415 ^b	7,367
AGE						
15-17	869	793	11	*	852	781
18-25	2,961	3,098	74	74	2,866	3,009
26-44	2,743 ^b	3,636	45	40	2,697 ^b	3,576
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5,503 ^b	6,401	105	98	5,379 ^b	6,267
White	4,045 ^b	4,764	68	67	3,962 ^b	4,669
Black or African American	1,010	1,156	33	22	972	1,128
American Indian or Alaska Native	56	42	*	*	55	40
Native Hawaiian or Other Pacific Islander	33	38	*	*	32	38
Asian	182	157	*	*	182	150
Two or More Races	178 ^a	244	*	*	175 ^a	242
Hispanic or Latino	1,071	1,126	25	25	1,037	1,100
TRIMESTER²						
First	N/A	N/A	82	63	N/A	N/A
Second	N/A	N/A	31	44	N/A	N/A
Third	N/A	N/A	16	14	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.72B Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	10.5 ^b	11.9	5.9	5.3	10.7 ^b	12.1
AGE						
15-17	13.9 ^a	12.6	18.3	*	13.8 ^a	12.5
18-25	17.2	17.8	9.0	9.1	17.6	18.3
26-44	7.0 ^b	9.2	3.4	2.7	7.2 ^b	9.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	10.9 ^b	12.6	6.0	5.2	11.1 ^b	12.9
White	11.2 ^b	13.3	5.5	5.3	11.4 ^b	13.5
Black or African American	11.8	13.1	9.6	5.7	11.8	13.4
American Indian or Alaska Native	13.7	11.6	*	*	14.1	11.8
Native Hawaiian or Other Pacific Islander	12.6	14.2	*	*	13.1	15.0
Asian	4.8	3.8	*	*	4.9	3.8
Two or More Races	15.6	19.9	*	*	16.0	20.6
Hispanic or Latino	8.8	8.9	5.6	5.9	8.9	9.1
TRIMESTER²						
First	N/A	N/A	11.6	8.9	N/A	N/A
Second	N/A	N/A	4.0	5.3	N/A	N/A
Third	N/A	N/A	2.3	1.9	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.73A Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
	2011-2012	2013-2014	Pregnant		Not Pregnant	
			2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	5,122 ^b	5,967	115	96	4,985 ^b	5,844
AGE						
15-17	709	659	10	*	694	649
18-25	2,485	2,648	62	57	2,406	2,578
26-44	1,928 ^b	2,660	43	30	1,884 ^b	2,616
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4,372 ^b	5,136	97	75	4,261 ^b	5,035
White	3,217 ^b	3,815	64	52	3,144 ^b	3,742
Black or African American	804 ^a	960	30	20	770 ^a	935
American Indian or Alaska Native	39	27	*	*	38	25
Native Hawaiian or Other Pacific Islander	25	31	*	*	25	31
Asian	132	101	*	*	132	101
Two or More Races	154	201	*	*	152	200
Hispanic or Latino	750	831	18	21	724	809
TRIMESTER²						
First	N/A	N/A	76	45	N/A	N/A
Second	N/A	N/A	23	42	N/A	N/A
Third	N/A	N/A	16	8	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.73B Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	8.2 ^b	9.4	5.2	4.1	8.3 ^b	9.6
AGE						
15-17	11.3	10.5	16.5	*	11.3	10.4
18-25	14.5	15.3	7.5	7.1	14.8	15.7
26-44	4.9 ^b	6.7	3.3	2.1	5.0 ^b	6.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8.7 ^b	10.1	5.5	4.0	8.8 ^b	10.4
White	8.9 ^b	10.6	5.2	4.1	9.1 ^b	10.9
Black or African American	9.4	10.9	8.7	5.3	9.4 ^a	11.1
American Indian or Alaska Native	9.6	7.5	*	*	9.9	7.4
Native Hawaiian or Other Pacific Islander	9.7	11.8	*	*	10.0	12.5
Asian	3.4	2.4	*	*	3.6	2.6
Two or More Races	13.5	16.4	*	*	13.8	17.0
Hispanic or Latino	6.2	6.6	3.9	4.9	6.2	6.7
TRIMESTER²						
First	N/A	N/A	10.7	6.5	N/A	N/A
Second	N/A	N/A	2.8	4.9	N/A	N/A
Third	N/A	N/A	2.3	1.1	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.74A Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
ILLICIT DRUGS²	6,574 ^b	7,527	130	123	6,415 ^b	7,367
TOBACCO PRODUCTS³	16,173	15,645	363	309	15,747	15,250
Cigarettes	15,148 ^a	14,457	352	301	14,739 ^a	14,076
Smokeless Tobacco	279	299	11	11	267	288
Cigars	2,104	2,014	33	23	2,057	1,971
Pipe Tobacco	339	374	6	5	330	359
ALCOHOL	33,625	34,183	187	214	33,310	33,819
Binge Alcohol Use ⁴	14,913	14,996	59	64	14,798	14,863
Heavy Alcohol Use ⁴	3,171	3,071	6	15	3,150	3,048
ILLICIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL^{2,3}	39,074	39,684	547	508	38,377	38,990
ILLICIT DRUGS OR ALCOHOL²	34,976	35,655	285	290	34,556	35,209
TOBACCO PRODUCTS OR ALCOHOL³	38,413	38,992	499	481	37,766	38,327

*Low precision; no estimate reported.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.74B Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
ILLCIT DRUGS²	10.5 ^b	11.9	5.9	5.3	10.7 ^b	12.1
TOBACCO PRODUCTS³	25.9 ^a	24.7	16.4	13.4	26.2 ^a	25.1
Cigarettes	24.2 ^b	22.8	15.9	13.1	24.6 ^b	23.2
Smokeless Tobacco	0.4	0.5	0.5	0.5	0.4	0.5
Cigars	3.4	3.2	1.5	1.0	3.4	3.2
Pipe Tobacco	0.5	0.6	0.2	0.2	0.5	0.6
ALCOHOL	53.8	54.0	8.5	9.3	55.5	55.7
Binge Alcohol Use ⁴	23.9	23.7	2.7	2.8	24.7	24.5
Heavy Alcohol Use ⁴	5.1	4.8	0.3	0.6	5.2	5.0
ILLCIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL^{2,3}	62.5	62.6	24.8	22.1	63.9	64.2
ILLCIT DRUGS OR ALCOHOL²	56.0	56.3	12.9	12.6	57.6	57.9
TOBACCO PRODUCTS OR ALCOHOL³	61.5	61.5	22.6	20.9	62.9	63.1

*Low precision; no estimate reported.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.75A Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
	2011-2012	2013-2014	Pregnant		Not Pregnant	
			2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	15,148 ^a	14,457	352	301	14,739 ^a	14,076
AGE						
15-17	691 ^b	528	*	*	670 ^b	518
18-25	4,786 ^b	4,166	172	155	4,591 ^b	3,979
26-44	9,671	9,763	166	138	9,479	9,579
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	13,234	12,688	335	277	12,847	12,334
White	10,637 ^a	10,072	251	214	10,344 ^a	9,801
Black or African American	1,618	1,807	54	49	1,558	1,745
American Indian or Alaska Native	188	162	*	*	177	151
Native Hawaiian or Other Pacific Islander	*	47	*	*	*	45
Asian	335	273	*	*	335	271
Two or More Races	379	328	*	*	359	319
Hispanic or Latino	1,913	1,769	17	24	1,893	1,742
TRIMESTER²						
First	N/A	N/A	164	116	N/A	N/A
Second	N/A	N/A	108	91	N/A	N/A
Third	N/A	N/A	78	90	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.75B Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	24.2 ^b	22.8	15.9	13.1	24.6 ^b	23.2
AGE						
15-17	11.0 ^b	8.4	*	*	10.9 ^b	8.3
18-25	27.8 ^b	24.0	20.9	19.2	28.2 ^b	24.2
26-44	24.8	24.6	12.5	9.5	25.2	25.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	26.3 ^a	25.0	19.1 ^a	14.7	26.5 ^a	25.4
White	29.4 ^a	28.1	20.3	16.9	29.8 ^a	28.4
Black or African American	18.9	20.4	15.8	12.6	19.0	20.8
American Indian or Alaska Native	46.3	44.7	*	*	45.5	44.1
Native Hawaiian or Other Pacific Islander	*	17.5	*	*	*	17.9
Asian	8.7	6.6	*	*	9.0	6.9
Two or More Races	33.2	26.8	*	*	32.7	27.2
Hispanic or Latino	15.8 ^a	14.0	3.8	5.7	16.3 ^a	14.4
TRIMESTER²						
First	N/A	N/A	23.2	16.5	N/A	N/A
Second	N/A	N/A	13.6	10.9	N/A	N/A
Third	N/A	N/A	11.1	12.3	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.76A Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	33,625	34,183	187	214	33,310	33,819
AGE						
15-17	1,348 ^a	1,236	8	*	1,331 ^a	1,227
18-25	9,934	9,942	60	49	9,827	9,841
26-44	22,344	23,005	119	160	22,152	22,751
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	28,413	28,703	162	195	28,136	28,381
White	22,000	21,794	121	143	21,788	21,569
Black or African American	4,065	4,395	31	40	4,018	4,332
American Indian or Alaska Native	197	168	*	*	196	163
Native Hawaiian or Other Pacific Islander	*	105	*	*	*	105
Asian	1,374	1,507	*	*	1,371	1,487
Two or More Races	672	734	*	*	658	725
Hispanic or Latino	5,212	5,480	25	19	5,174	5,438
TRIMESTER²						
First	N/A	N/A	127	147	N/A	N/A
Second	N/A	N/A	33	41	N/A	N/A
Third	N/A	N/A	26	24	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.76B Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	53.8	54.0	8.5	9.3	55.5	55.7
AGE						
15-17	21.5 ^a	19.7	13.4	*	21.6 ^a	19.7
18-25	57.8	57.3	7.3	6.1	60.4	59.8
26-44	57.2	57.9	9.0	11.0	58.9	59.8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	56.4	56.6	9.2	10.4	58.1	58.4
White	60.9	60.7	9.7	11.3	62.7	62.5
Black or African American	47.4	49.7	9.2	10.4	49.0	51.5
American Indian or Alaska Native	48.7	46.4	*	*	50.4	47.6
Native Hawaiian or Other Pacific Islander	*	39.4	*	*	*	41.6
Asian	35.8	36.5	*	*	36.9	37.5
Two or More Races	58.9	60.0	*	*	60.0	61.8
Hispanic or Latino	43.0	43.4	5.5	4.5	44.5	44.8
TRIMESTER²						
First	N/A	N/A	17.9	20.9	N/A	N/A
Second	N/A	N/A	4.2	4.9	N/A	N/A
Third	N/A	N/A	3.7	3.3	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.77A Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	14,913	14,996	59	64	14,798	14,863
AGE						
15-17	718 ^b	607	*	*	710 ^b	603
18-25	5,707	5,485	25	25	5,658	5,430
26-44	8,488	8,904	30	37	8,431	8,830
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	12,397	12,363	52	56	12,295	12,251
White	9,817	9,627	35	28	9,737	9,568
Black or African American	1,665	1,823	10	17	1,650	1,787
American Indian or Alaska Native	123	105	*	*	122	100
Native Hawaiian or Other Pacific Islander	59	36	*	*	*	36
Asian	459	490	*	*	458	486
Two or More Races	274	282	*	*	268	274
Hispanic or Latino	2,516	2,633	7	8	2,503	2,612
TRIMESTER²						
First	N/A	N/A	47	47	N/A	N/A
Second	N/A	N/A	9	15	N/A	N/A
Third	N/A	N/A	3	2	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.77B Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	23.9	23.7	2.7	2.8	24.7	24.5
AGE						
15-17	11.5 ^b	9.7	*	*	11.5 ^b	9.7
18-25	33.2 ^a	31.6	3.0	3.1	34.8 ^b	33.0
26-44	21.7	22.4	2.3	2.5	22.4	23.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	24.6	24.4	3.0	3.0	25.4	25.2
White	27.2	26.8	2.8	2.2	28.0	27.7
Black or African American	19.4	20.6	3.1	4.3	20.1	21.3
American Indian or Alaska Native	30.3	28.9	*	*	31.3	29.2
Native Hawaiian or Other Pacific Islander	22.6	13.7	*	*	*	14.5
Asian	12.0	11.9	*	*	12.3	12.3
Two or More Races	24.0	23.0	*	*	24.5	23.3
Hispanic or Latino	20.7	20.8	1.5	1.9	21.5	21.5
TRIMESTER²						
First	N/A	N/A	6.6	6.7	N/A	N/A
Second	N/A	N/A	1.1	1.8	N/A	N/A
Third	N/A	N/A	0.4	0.3	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.78A Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	3,171	3,071	6	15	3,150	3,048
AGE						
15-17	130 ^a	101	*	*	130 ^a	99
18-25	1,360	1,283	4	5	1,349	1,270
26-44	1,680	1,687	2	8	1,671	1,679
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,786	2,645	5	13	2,768	2,624
White	2,304	2,205	4	8	2,287	2,193
Black or African American	278	292	0	4	277	285
American Indian or Alaska Native	44	23	*	*	44	22
Native Hawaiian or Other Pacific Islander	*	7	*	*	*	7
Asian	75	61	*	*	75	60
Two or More Races	59	58	*	*	59	58
Hispanic or Latino	385	426	1	*	382	424
TRIMESTER²						
First	N/A	N/A	5	10	N/A	N/A
Second	N/A	N/A	0	4	N/A	N/A
Third	N/A	N/A	*	0	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.78B Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	5.1	4.8	0.3	0.6	5.2	5.0
AGE						
15-17	2.1 ^a	1.6	*	*	2.1 ^a	1.6
18-25	7.9	7.4	0.4	0.6	8.3	7.7
26-44	4.3	4.2	0.2	0.6	4.4	4.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5.5	5.2	0.3	0.7	5.7	5.4
White	6.4	6.1	0.3	0.6	6.6	6.4
Black or African American	3.2	3.3	0.1	1.1	3.4	3.4
American Indian or Alaska Native	10.9	6.2	*	*	11.3	6.3
Native Hawaiian or Other Pacific Islander	*	2.5	*	*	*	2.7
Asian	1.9	1.5	*	*	2.0	1.5
Two or More Races	5.2	4.7	*	*	5.3	4.9
Hispanic or Latino	3.2	3.4	0.3	*	3.3	3.5
TRIMESTER²						
First	N/A	N/A	0.8	1.5	N/A	N/A
Second	N/A	N/A	0.1	0.5	N/A	N/A
Third	N/A	N/A	*	0.0	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2013-2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013-2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.79A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	4,979	5,057	1,934	1,961	3,045	3,096
Marijuana and Hashish	4,463	4,543	1,717	1,774	2,746	2,769
Cocaine	251	299	87 ^a	154	164	145
Crack	15	19	1	8	14	11
Heroin	41	38	14	4	26	35
Hallucinogens	435	339	182	131	253	208
LSD	96	74	44	21	52	53
PCP	2	2	*	2	2	*
Ecstasy	218	186	86	75	132	111
Inhalants	83	45	39	19	45	26
Nonmedical Use of Psychotherapeutics ^{4,5}	1,045	1,001	369	379	676	622
Pain Relievers	688	632	193	200	496	431
OxyContin [®]	73	62	35 ^a	6	38	56
Tranquilizers	265	273	84	83	181	190
Stimulants ⁵	294	261	145	132	149	129
Methamphetamine ⁵	73	60	12	14	61	45
Sedatives	21	29	6	6	15	23
ILLCIT DRUGS OTHER THAN MARIJUANA³	1,527	1,448	576	575	951	873

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 to 22, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.79B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	22.7	23.0	22.3	22.4	23.0	23.4
Marijuana and Hashish	20.4	20.7	19.8	20.3	20.7	21.0
Cocaine	1.1	1.4	1.0 ^a	1.8	1.2	1.1
Crack	0.1	0.1	0.0	0.1	0.1	0.1
Heroin	0.2	0.2	0.2	0.0	0.2	0.3
Hallucinogens	2.0	1.5	2.1	1.5	1.9	1.6
LSD	0.4	0.3	0.5	0.2	0.4	0.4
PCP	0.0	0.0	*	0.0	0.0	*
Ecstasy	1.0	0.8	1.0	0.9	1.0	0.8
Inhalants	0.4	0.2	0.4	0.2	0.3	0.2
Nonmedical Use of Psychotherapeutics ^{4,5}	4.8	4.6	4.3	4.3	5.1	4.7
Pain Relievers	3.1	2.9	2.2	2.3	3.7	3.3
OxyContin [®]	0.3	0.3	0.4 ^a	0.1	0.3	0.4
Tranquilizers	1.2	1.2	1.0	1.0	1.4	1.4
Stimulants ⁵	1.3	1.2	1.7	1.5	1.1	1.0
Methamphetamine ⁵	0.3	0.3	0.1	0.2	0.5	0.3
Sedatives	0.1	0.1	0.1	0.1	0.1	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA³	7.0	6.6	6.7	6.6	7.2	6.6

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 to 22, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.80A Types of Illicit Drug Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Male College Students		Other Males Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	2,929	2,896	1,043	1,015	1,886	1,881
Marijuana and Hashish	2,677	2,661	948	930	1,729	1,730
Cocaine	157	176	44	81	113	94
Crack	7	14	*	8	7	6
Heroin	25	23	12	*	14	23
Hallucinogens	315 ^b	187	129	77	186 ^a	111
LSD	70 ^a	33	30	8	40	25
PCP	1	2	*	2	1	*
Ecstasy	145 ^a	88	57	38	89	51
Inhalants	64 ^a	27	26	13	38 ^a	14
Nonmedical Use of Psychotherapeutics ^{4,5}	560	516	201	188	359	329
Pain Relievers	368	336	109	98	259	238
OxyContin [®]	41	45	23	6	18	40
Tranquilizers	133	149	38	46	95	104
Stimulants ⁵	161	131	83	64	78	67
Methamphetamine ⁵	38	31	7	4	32	27
Sedatives	12	8	3	2	9	6
ILLCIT DRUGS OTHER THAN MARIJUANA³	893	753	335	284	558	469

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all males aged 18 to 22, including those with unknown enrollment status.

² Other Males include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.80B Types of Illicit Drug Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Male College Students		Other Males Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	26.4	25.6	26.0	24.6	26.6	26.2
Marijuana and Hashish	24.1	23.5	23.6	22.5	24.4	24.1
Cocaine	1.4	1.6	1.1	2.0	1.6	1.3
Crack	0.1	0.1	*	0.2	0.1	0.1
Heroin	0.2	0.2	0.3	*	0.2	0.3
Hallucinogens	2.8 ^b	1.7	3.2	1.9	2.6 ^a	1.5
LSD	0.6 ^a	0.3	0.8 ^a	0.2	0.6	0.3
PCP	0.0	0.0	*	0.0	0.0	*
Ecstasy	1.3 ^a	0.8	1.4	0.9	1.2	0.7
Inhalants	0.6 ^a	0.2	0.6	0.3	0.5 ^a	0.2
Nonmedical Use of Psychotherapeutics ^{4,5}	5.0	4.6	5.0	4.5	5.1	4.6
Pain Relievers	3.3	3.0	2.7	2.4	3.7	3.3
OxyContin [®]	0.4	0.4	0.6	0.1	0.3	0.6
Tranquilizers	1.2	1.3	0.9	1.1	1.3	1.4
Stimulants ⁵	1.4	1.2	2.1	1.6	1.1	0.9
Methamphetamine ⁵	0.3	0.3	0.2	0.1	0.4	0.4
Sedatives	0.1	0.1	0.1	0.0	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA³	8.0 ^a	6.7	8.3	6.9	7.9	6.5

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all males aged 18 to 22, including those with unknown enrollment status.

² Other Males include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.81A Types of Illicit Drug Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Female College Students		Other Females Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	2,050	2,161	891	946	1,159	1,215
Marijuana and Hashish	1,786	1,883	768	844	1,017	1,039
Cocaine	93	123	43	72	50	51
Crack	9	5	1	*	8	5
Heroin	15	15	3	4	13	11
Hallucinogens	120	151	53	54	67	97
LSD	26	42	13	14	13	28
PCP	*	*	*	*	0	*
Ecstasy	73	97	30	37	43	60
Inhalants	20	18	13	6	7	12
Nonmedical Use of Psychotherapeutics ^{4,5}	485	485	168	191	317	294
Pain Relievers	320	296	83	102	236	193
OxyContin [®]	32	16	12	0	20	16
Tranquilizers	132	124	46	38	86	86
Stimulants ⁵	134	130	62	68	72	62
Methamphetamine ⁵	34	29	5	11	29	19
Sedatives	9	21	3	4	6	17
ILLCIT DRUGS OTHER THAN MARIJUANA³	635	695	241	291	394	404

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 18 to 22, including those with unknown enrollment status.

² Other Females include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.81B Types of Illicit Drug Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Female College Students		Other Females Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	19.0	20.3	19.2	20.4	18.9	20.2
Marijuana and Hashish	16.5	17.7	16.6	18.2	16.5	17.3
Cocaine	0.9	1.2	0.9	1.6	0.8	0.9
Crack	0.1	0.0	0.0	*	0.1	0.1
Heroin	0.1	0.1	0.1	0.1	0.2	0.2
Hallucinogens	1.1	1.4	1.1	1.2	1.1	1.6
LSD	0.2	0.4	0.3	0.3	0.2	0.5
PCP	*	*	*	*	0.0	*
Ecstasy	0.7	0.9	0.6	0.8	0.7	1.0
Inhalants	0.2	0.2	0.3	0.1	0.1	0.2
Nonmedical Use of Psychotherapeutics ^{4,5}	4.5	4.6	3.6	4.1	5.2	4.9
Pain Relievers	3.0	2.8	1.8	2.2	3.8	3.2
OxyContin [®]	0.3	0.2	0.3	0.0	0.3	0.3
Tranquilizers	1.2	1.2	1.0	0.8	1.4	1.4
Stimulants ⁵	1.2	1.2	1.3	1.5	1.2	1.0
Methamphetamine ⁵	0.3	0.3	0.1	0.2	0.5	0.3
Sedatives	0.1	0.2	0.1	0.1	0.1	0.3
ILLCIT DRUGS OTHER THAN MARIJUANA³	5.9	6.5	5.2	6.3	6.4	6.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 18 to 22, including those with unknown enrollment status.

² Other Females include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.82A Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	4,979	5,057	1,934	1,961	3,045	3,096
AGE						
18	993	963	290	323	703	640
19	948	971	456	418	492	553
20	1,036	1,039	431	485	605	554
21	1,022	1,016	485	394	537	622
22	980	1,068	272	341	708	727
GENDER						
Male	2,929	2,896	1,043	1,015	1,886	1,881
Female	2,050	2,161	891	946	1,159	1,215
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4,016	4,149	1,626	1,679	2,390	2,469
White	3,018	2,974	1,303	1,233	1,715	1,741
Black or African American	665 ^a	807	201	244	463	563
American Indian or Alaska Native	23	*	*	*	20	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	144	152	67	118	78 ^a	33
Two or More Races	137	153	*	71	94	83
Hispanic or Latino	963	908	309	282	655	626
CURRENT EMPLOYMENT						
Full-Time	1,418 ^b	1,753	313	388	1,105 ^b	1,365
Part-Time	1,744	1,650	882	891	862	759
Unemployed	843 ^a	705	193	138	650	567
Other ³	974	949	546	544	428	405

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.82B Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	22.7	23.0	22.3	22.4	23.0	23.4
AGE						
18	21.0	20.9	18.3	19.4	22.3	21.8
19	22.8	23.5	22.1	21.5	23.4	25.2
20	24.1	23.8	21.8	24.8	26.1	23.1
21	24.1	23.4	28.0 ^b	21.3	21.4	24.9
22	21.9	23.7	21.0	25.5	22.3	22.9
GENDER						
Male	26.4	25.6	26.0	24.6	26.6	26.2
Female	19.0	20.3	19.2	20.4	18.9	20.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	23.4	24.2	22.5	23.4	24.0	24.7
White	24.8	24.8	25.1	24.6	24.7	24.9
Black or African American	21.4 ^a	25.3	19.7	23.0	22.2	26.5
American Indian or Alaska Native	17.7	*	*	*	18.8	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	12.6	12.5	9.4	14.6	18.0 ^a	8.3
Two or More Races	27.1	29.4	*	32.1	31.5	27.4
Hispanic or Latino	20.4	19.0	21.5	18.0	19.9	19.5
CURRENT EMPLOYMENT						
Full-Time	22.5 ^a	25.3	22.7	25.9	22.4	25.1
Part-Time	24.2	23.6	23.6	23.1	25.0	24.2
Unemployed	28.1	25.8	26.4	20.3	28.7	27.6
Other ³	18.0	17.9	19.5	20.0	16.4	15.7

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.83A Marijuana Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	4,463	4,543	1,717	1,774	2,746	2,769
AGE						
18	897	872	260	294	637	578
19	873	895	409	382	464	513
20	915	969	374	465	541	503
21	919	891	430	342	488	549
22	860	917	244	291	616	626
GENDER						
Male	2,677	2,661	948	930	1,729	1,730
Female	1,786	1,883	768	844	1,017	1,039
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	3,601	3,765	1,438	1,522	2,163	2,243
White	2,729	2,681	1,160	1,133	1,569	1,548
Black or African American	596 ^b	780	172	233	424 ^a	546
American Indian or Alaska Native	17	*	*	*	15	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	104	113	54	85	50	28
Two or More Races	127	141	*	62	84	79
Hispanic or Latino	862	778	279	252	583	526
CURRENT EMPLOYMENT						
Full-Time	1,267 ^b	1,571	281	354	987 ^b	1,217
Part-Time	1,568	1,483	787	806	781	677
Unemployed	770	648	163	130	607	518
Other ³	857	842	486	485	372	358

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.83B Marijuana Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	20.4	20.7	19.8	20.3	20.7	21.0
AGE						
18	18.9	18.9	16.4	17.6	20.2	19.7
19	20.9	21.6	19.8	19.6	22.1	23.4
20	21.3	22.2	18.9 ^a	23.8	23.4	20.9
21	21.7	20.5	24.9 ^b	18.5	19.5	22.0
22	19.2	20.3	18.8	21.8	19.4	19.7
GENDER						
Male	24.1	23.5	23.6	22.5	24.4	24.1
Female	16.5	17.7	16.6	18.2	16.5	17.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	21.0	21.9	19.9	21.2	21.7	22.5
White	22.5	22.3	22.3	22.6	22.6	22.1
Black or African American	19.1 ^b	24.4	16.8	21.9	20.3 ^a	25.7
American Indian or Alaska Native	12.8	*	*	*	14.1	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	9.1	9.3	7.6	10.4	11.6	7.0
Two or More Races	25.0	27.0	*	28.0	28.3	26.3
Hispanic or Latino	18.2	16.2	19.4	16.1	17.7	16.4
CURRENT EMPLOYMENT						
Full-Time	20.1 ^a	22.7	20.4	23.6	20.0	22.4
Part-Time	21.8	21.2	21.0	20.9	22.7	21.6
Unemployed	25.7	23.7	22.3	19.1	26.8	25.2
Other ³	15.9	15.9	17.3	17.8	14.3	13.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.84A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	7,815	7,415	2,490 ^a	2,174	5,325	5,240
Cigarettes	6,375 ^a	5,873	1,816 ^a	1,568	4,560	4,305
Smokeless Tobacco	1,351	1,284	430	352	921	933
Cigars	2,325	2,295	864	770	1,461	1,525
Pipe Tobacco	575 ^a	446	258	186	317	260
ALCOHOL	11,847	12,033	5,144	5,240	6,701	6,794
Binge Alcohol Use ⁴	7,800	7,750	3,374	3,323	4,424	4,427
Heavy Alcohol Use ⁴	2,330	2,317	1,097	1,066	1,233	1,251

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 to 22, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.84B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	35.7 ^a	33.8	28.8 ^b	24.8	40.2	39.7
Cigarettes	29.1 ^b	26.7	21.0 ^b	17.9	34.4	32.6
Smokeless Tobacco	6.2	5.8	5.0	4.0	7.0	7.1
Cigars	10.6	10.4	10.0	8.8	11.0	11.6
Pipe Tobacco	2.6 ^a	2.0	3.0	2.1	2.4	2.0
ALCOHOL	54.1	54.8	59.4	59.8	50.6	51.5
Binge Alcohol Use ⁴	35.6	35.3	39.0	37.9	33.4	33.5
Heavy Alcohol Use ⁴	10.6	10.5	12.7	12.2	9.3	9.5

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 to 22, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.85A Tobacco Product and Alcohol Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Male College Students		Other Males Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	4,922	4,777	1,503	1,333	3,419	3,443
Cigarettes	3,817	3,611	1,016	909	2,800	2,702
Smokeless Tobacco	1,261	1,233	393	341	868	891
Cigars	1,736	1,652	631	523	1,105	1,129
Pipe Tobacco	409 ^a	290	180 ^a	108	229	182
ALCOHOL	6,200	6,350	2,440	2,490	3,760	3,860
Binge Alcohol Use ⁴	4,505	4,509	1,801	1,795	2,705	2,714
Heavy Alcohol Use ⁴	1,520	1,485	665	638	855	847

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all males aged 18 to 22, including those with unknown enrollment status.

² Other Males include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.85B Tobacco Product and Alcohol Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Male College Students		Other Males Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	44.3	42.2	37.4 ^a	32.3	48.2	47.9
Cigarettes	34.3 ^a	31.9	25.3	22.0	39.5	37.6
Smokeless Tobacco	11.3	10.9	9.8	8.3	12.2	12.4
Cigars	15.6	14.6	15.7 ^a	12.7	15.6	15.7
Pipe Tobacco	3.7 ^b	2.6	4.5 ^a	2.6	3.2	2.5
ALCOHOL	55.8	56.1	60.8	60.3	53.0	53.7
Binge Alcohol Use ⁴	40.5	39.8	44.8	43.5	38.1	37.8
Heavy Alcohol Use ⁴	13.7	13.1	16.5	15.4	12.1	11.8

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all males aged 18 to 22, including those with unknown enrollment status.

² Other Males include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.86A Tobacco Product and Alcohol Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Numbers in Thousands, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Female College Students		Other Females Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	2,893	2,638	987	841	1,906	1,797
Cigarettes	2,559 ^a	2,262	799 ^a	658	1,760	1,604
Smokeless Tobacco	91 ^a	52	37 ^a	10	54	41
Cigars	589	643	233	247	356	397
Pipe Tobacco	166	156	78	78	88	78
ALCOHOL	5,647	5,683	2,704	2,750	2,941	2,933
Binge Alcohol Use ⁴	3,294	3,241	1,573	1,528	1,719	1,713
Heavy Alcohol Use ⁴	810	832	432	429	378	404

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 18 to 22, including those with unknown enrollment status.

² Other Females include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.86B Tobacco Product and Alcohol Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Percentages, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Female College Students		Other Females Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	26.8	24.8	21.3 ^a	18.2	31.0	29.9
Cigarettes	23.7 ^a	21.3	17.2 ^a	14.2	28.6	26.7
Smokeless Tobacco	0.8	0.5	0.8 ^a	0.2	0.9	0.7
Cigars	5.5	6.0	5.0	5.3	5.8	6.6
Pipe Tobacco	1.5	1.5	1.7	1.7	1.4	1.3
ALCOHOL	52.3	53.4	58.2	59.4	47.8	48.8
Binge Alcohol Use ⁴	30.5	30.4	33.9	33.0	28.0	28.5
Heavy Alcohol Use ⁴	7.5	7.8	9.3	9.3	6.1	6.7

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 18 to 22, including those with unknown enrollment status.

² Other Females include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.87A Cigarette Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	6,375 ^a	5,873	1,816 ^a	1,568	4,560	4,305
AGE						
18	992	892	224	218	768	674
19	1,188 ^a	993	422 ^b	276	767	716
20	1,391	1,261	456	403	934	858
21	1,336	1,285	424	370	912	915
22	1,468	1,442	290	301	1,178	1,142
GENDER						
Male	3,817	3,611	1,016	909	2,800	2,702
Female	2,559 ^a	2,262	799 ^a	658	1,760	1,604
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5,235 ^a	4,815	1,553	1,352	3,681	3,462
White	4,195 ^a	3,824	1,314 ^a	1,089	2,880	2,735
Black or African American	640	648	121	106	519	542
American Indian or Alaska Native	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	148	163	72	113	76	50
Two or More Races	165	117	38	*	127 ^a	83
Hispanic or Latino	1,141	1,058	263	215	878	843
CURRENT EMPLOYMENT						
Full-Time	2,287	2,337	310	302	1,978	2,035
Part-Time	1,849 ^b	1,543	779	671	1,069 ^a	872
Unemployed	1,054 ^a	878	184 ^b	85	871	793
Other ³	1,185	1,115	543	510	642	605

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.87B Cigarette Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	29.1 ^b	26.7	21.0 ^b	17.9	34.4	32.6
AGE						
18	21.0	19.4	14.1	13.1	24.4	23.0
19	28.5 ^b	24.0	20.4 ^b	14.2	36.5	32.7
20	32.4	28.9	23.0	20.6	40.4	35.7
21	31.5	29.6	24.5	20.0	36.4	36.6
22	32.8	32.0	22.3	22.5	37.1	36.0
GENDER						
Male	34.3 ^a	31.9	25.3	22.0	39.5	37.6
Female	23.7 ^a	21.3	17.2 ^a	14.2	28.6	26.7
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	30.5 ^b	28.0	21.5 ^a	18.8	37.0	34.7
White	34.5 ^a	31.8	25.3 ^a	21.7	41.5	39.1
Black or African American	20.6	20.3	11.9	9.9	24.8	25.5
American Indian or Alaska Native	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	13.0	13.4	10.1	13.9	17.6	12.4
Two or More Races	32.5 ^a	22.5	18.1	*	42.6 ^a	27.6
Hispanic or Latino	24.1	22.1	18.3	13.7	26.6	26.2
CURRENT EMPLOYMENT						
Full-Time	36.3	33.7	22.5	20.2	40.2	37.5
Part-Time	25.7 ^b	22.1	20.8 ^a	17.4	31.0	27.8
Unemployed	35.1	32.1	25.0 ^b	12.4	38.4	38.6
Other ³	21.9	21.0	19.4	18.8	24.6	23.4

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.88A Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	11,847	12,033	5,144	5,240	6,701	6,794
AGE						
18	1,661	1,678	610	711	1,051	967
19	1,894	1,783	1,028	917	867	866
20	2,219	2,327	1,195	1,166	1,025	1,161
21	2,941	3,095	1,326	1,436	1,614	1,659
22	3,132	3,150	986	1,009	2,145	2,140
GENDER						
Male	6,200	6,350	2,440	2,490	3,760	3,860
Female	5,647	5,683	2,704	2,750	2,941	2,933
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	9,512	9,717	4,363	4,379	5,148	5,338
White	7,359	7,357	3,471	3,328	3,886	4,030
Black or African American	1,351	1,442	503	522	849	920
American Indian or Alaska Native	56	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	401	501	250	350	151	151
Two or More Races	269	304	109	135	161	169
Hispanic or Latino	2,335	2,316	781	860	1,553	1,456
CURRENT EMPLOYMENT						
Full-Time	3,879 ^b	4,446	841	983	3,038 ^b	3,463
Part-Time	4,152	3,956	2,377	2,406	1,773	1,550
Unemployed	1,497 ^a	1,303	423	352	1,074	951
Other ³	2,320	2,329	1,503	1,498	817	830

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.88B Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	54.1	54.8	59.4	59.8	50.6	51.5
AGE						
18	35.1	36.4	38.4	42.7	33.4	32.9
19	45.5	43.1	49.8	47.1	41.2	39.5
20	51.7	53.4	60.3	59.6	44.3	48.3
21	69.4	71.2	76.7	77.7	64.4	66.4
22	70.0	69.8	76.0	75.5	67.5	67.4
GENDER						
Male	55.8	56.1	60.8	60.3	53.0	53.7
Female	52.3	53.4	58.2	59.4	47.8	48.8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	55.4	56.6	60.4	60.9	51.8	53.4
White	60.6	61.3	66.7	66.3	56.0	57.6
Black or African American	43.4	45.2	49.2	49.0	40.6	43.3
American Indian or Alaska Native	42.9	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	35.1	41.4	35.1	43.3	35.1	37.5
Two or More Races	53.2	58.3	52.2	61.1	53.8	56.3
Hispanic or Latino	49.3	48.4	54.4	54.9	47.1	45.3
CURRENT EMPLOYMENT						
Full-Time	61.6	64.2	61.0	65.6	61.7	63.8
Part-Time	57.7	56.5	63.5	62.3	51.4	49.4
Unemployed	49.9	47.7	57.6	51.7	47.4	46.3
Other ³	42.9	43.9	53.6	55.2	31.3	32.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.89A Binge Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	7,800	7,750	3,374	3,323	4,424	4,427
AGE						
18	1,053	1,098	384	457	670	640
19	1,312	1,131	710	578	601	553
20	1,478	1,510	764	713	714	797
21	1,914	2,046	866	938	1,048	1,108
22	2,042	1,965	650	637	1,390	1,328
GENDER						
Male	4,505	4,509	1,801	1,795	2,705	2,714
Female	3,294	3,241	1,573	1,528	1,719	1,713
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	6,255	6,229	2,862	2,792	3,391	3,437
White	5,012	4,924	2,384	2,229	2,626	2,695
Black or African American	745	792	261	283	484	510
American Indian or Alaska Native	35	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	247	275	129	186	118	90
Two or More Races	168	176	70	80	98	96
Hispanic or Latino	1,545	1,521	512	531	1,032	990
CURRENT EMPLOYMENT						
Full-Time	2,622 ^a	2,944	563	621	2,060 ^a	2,323
Part-Time	2,673	2,454	1,549	1,495	1,122	960
Unemployed	1,005	892	248	227	757	665
Other ³	1,499	1,460	1,014	980	485	480

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.89B Binge Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	35.6	35.3	39.0	37.9	33.4	33.5
AGE						
18	22.3	23.8	24.2	27.4	21.3	21.8
19	31.5 ^a	27.3	34.4	29.7	28.6	25.2
20	34.4	34.6	38.6	36.4	30.9	33.1
21	45.2	47.1	50.1	50.7	41.8	44.4
22	45.6	43.6	50.1	47.6	43.8	41.8
GENDER						
Male	40.5	39.8	44.8	43.5	38.1	37.8
Female	30.5	30.4	33.9	33.0	28.0	28.5
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	36.4	36.3	39.6	38.8	34.1	34.4
White	41.3	41.0	45.8	44.4	37.8	38.5
Black or African American	23.9	24.8	25.5	26.6	23.2	24.0
American Indian or Alaska Native	27.2	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	21.6	22.7	18.2	22.9	27.3	22.3
Two or More Races	33.1	33.7	33.5	36.2	32.9	31.8
Hispanic or Latino	32.6	31.8	35.7	33.9	31.3	30.8
CURRENT EMPLOYMENT						
Full-Time	41.6	42.5	40.8	41.4	41.8	42.8
Part-Time	37.1	35.1	41.4	38.7	32.6	30.6
Unemployed	33.5	32.6	33.8	33.4	33.4	32.4
Other ³	27.7	27.5	36.2	36.1	18.6	18.6

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.90A Heavy Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	2,330	2,317	1,097	1,066	1,233	1,251
AGE						
18	284	270	123	101	160	169
19	356	325	237	185	119	140
20	480	449	268	261	212	188
21	585	675	275	314	310	362
22	625	599	193	206	432	393
GENDER						
Male	1,520	1,485	665	638	855	847
Female	810	832	432	429	378	404
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,944	1,895	962	909	981	986
White	1,706	1,596	865	771	841	825
Black or African American	135	160	48	51	87	109
American Indian or Alaska Native	5	8	*	*	5	5
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	48	59	37	53	12	*
Two or More Races	38	65	9	*	28	36
Hispanic or Latino	386	422	135	157	252	265
CURRENT EMPLOYMENT						
Full-Time	833	899	192	231	640	668
Part-Time	801	716	502	438	299	278
Unemployed	268	255	80	50	187	205
Other ³	428	447	322	347	107	100

*Low precision; no estimate reported.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.90B Heavy Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	10.6	10.5	12.7	12.2	9.3	9.5
AGE						
18	6.0	5.9	7.8	6.1	5.1	5.7
19	8.5	7.8	11.5	9.5	5.6	6.4
20	11.2	10.3	13.6	13.3	9.2	7.8
21	13.8	15.5	15.9	17.0	12.4	14.5
22	14.0	13.3	14.9	15.4	13.6	12.4
GENDER						
Male	13.7	13.1	16.5	15.4	12.1	11.8
Female	7.5	7.8	9.3	9.3	6.1	6.7
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	11.3	11.0	13.3	12.6	9.9	9.9
White	14.0	13.3	16.6	15.4	12.1	11.8
Black or African American	4.3	5.0	4.7	4.8	4.2	5.1
American Indian or Alaska Native	4.1	6.1	*	*	4.7	4.9
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	4.2	4.9	5.1	6.5	2.7	*
Two or More Races	7.5	12.5	4.5	*	9.5	11.9
Hispanic or Latino	8.2	8.8	9.4	10.0	7.6	8.2
CURRENT EMPLOYMENT						
Full-Time	13.2	13.0	14.0	15.4	13.0	12.3
Part-Time	11.1	10.2	13.4	11.3	8.7	8.9
Unemployed	8.9	9.3	11.0	7.4	8.3	10.0
Other ³	7.9	8.4	11.5	12.8	4.1	3.9

*Low precision; no estimate reported.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.91A Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	1,733	1,562	554	491	1,179	1,071
AGE						
18	375	312	78	78	297	233
19	328	284	128	112	200	172
20	339	295	127	121	212	174
21	370	306	153 ^a	88	217	218
22	322	365	68	92	253	274
GENDER						
Male	1,123	989	343 ^a	255	779	734
Female	611	573	211	236	400	337
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,445	1,286	478	428	967	858
White	1,056	912	402 ^a	294	655	618
Black or African American	230	240	48	60	183	180
American Indian or Alaska Native	15 ^a	4	*	*	*	3
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	72	71	16 ^a	53	56 ^a	17
Two or More Races	59	54	12	18	47	36
Hispanic or Latino	288	276	76	64	212	213
CURRENT EMPLOYMENT						
Full-Time	522	530	105	67	417	463
Part-Time	532	458	229	232	303 ^a	227
Unemployed	365 ^a	265	72	39	293	226
Other ³	315	308	149	153	166	155

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.91B Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	7.9	7.1	6.4	5.6	8.9	8.1
AGE						
18	7.9	6.8	4.9	4.7	9.4	7.9
19	7.9	6.9	6.2	5.8	9.5	7.9
20	7.9	6.8	6.4	6.2	9.2	7.2
21	8.7	7.0	8.8 ^b	4.7	8.7	8.7
22	7.2	8.1	5.3	6.9	8.0	8.6
GENDER						
Male	10.1 ^a	8.7	8.6 ^a	6.2	11.0	10.2
Female	5.7	5.4	4.5	5.1	6.5	5.6
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8.4	7.5	6.6	5.9	9.7	8.6
White	8.7	7.6	7.7 ^a	5.9	9.4	8.8
Black or African American	7.4	7.5	4.7	5.6	8.7	8.5
American Indian or Alaska Native	11.3 ^a	2.8	*	*	*	3.5
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	6.3	5.8	2.2 ^a	6.6	13.0 ^a	4.3
Two or More Races	11.7	10.3	5.7	8.0	15.8	12.0
Hispanic or Latino	6.1	5.8	5.3	4.1	6.4	6.6
CURRENT EMPLOYMENT						
Full-Time	8.3	7.6	7.6 ^a	4.5	8.5	8.5
Part-Time	7.4	6.6	6.1	6.0	8.8	7.2
Unemployed	12.2	9.7	9.8	5.7	12.9	11.0
Other ³	5.8	5.8	5.3	5.6	6.4	6.0

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.92A Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	2,731	2,587	1,170	1,156	1,561	1,431
AGE						
18	440	368	146	173	294 ^a	196
19	460 ^a	357	229	185	231	172
20	549	549	304	281	245	268
21	609	686	288	313	321	373
22	673	628	202	205	470	423
GENDER						
Male	1,558	1,487	609	593	949	894
Female	1,173	1,100	561	564	612	536
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,141	2,011	958	950	1,183	1,061
White	1,746	1,578	806	730	940	848
Black or African American	215	223	57	87	158	136
American Indian or Alaska Native	15	*	*	*	14	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	88	119	56	95	33	24
Two or More Races	67	53	32	*	34	28
Hispanic or Latino	590	576	212	206	379	369
CURRENT EMPLOYMENT						
Full-Time	919	956	207	236	711	720
Part-Time	951	835	551	522	401	313
Unemployed	336	325	69	76	267	249
Other ³	525	471	342	322	182	149

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.92B Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	12.5	11.8	13.5	13.2	11.8	10.8
AGE						
18	9.3	8.0	9.2	10.4	9.4 ^a	6.7
19	11.0 ^a	8.6	11.1	9.5	11.0	7.8
20	12.8	12.6	15.3	14.4	10.6	11.1
21	14.4	15.8	16.7	16.9	12.8	14.9
22	15.0	13.9	15.6	15.3	14.8	13.3
GENDER						
Male	14.0	13.1	15.2	14.4	13.4	12.4
Female	10.9	10.3	12.1	12.2	10.0	8.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	12.5	11.7	13.3	13.2	11.9	10.6
White	14.4	13.1	15.5	14.5	13.5	12.1
Black or African American	6.9	7.0	5.6	8.1	7.6	6.4
American Indian or Alaska Native	11.5	*	*	*	13.3	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	7.7	9.8	7.8	11.7	7.5	6.1
Two or More Races	13.2	10.2	15.5	*	11.5	9.2
Hispanic or Latino	12.5	12.0	14.7	13.2	11.5	11.5
CURRENT EMPLOYMENT						
Full-Time	14.6	13.8	15.1	15.8	14.5	13.3
Part-Time	13.2	11.9	14.7	13.5	11.6	10.0
Unemployed	11.2	11.9	9.5	11.2	11.8	12.1
Other ³	9.7	8.9	12.2	11.8	7.0	5.8

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.93A Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	3,735	3,533	1,486	1,427	2,249	2,107
AGE						
18	660	597	192	227	467	370
19	642	547	299	248	343	299
20	746	708	368	345	378	363
21	832	839	392	347	440	492
22	855	842	234	260	620	582
GENDER						
Male	2,212	2,107	800	741	1,412	1,366
Female	1,523	1,426	686	686	838	740
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,978	2,793	1,226	1,194	1,752	1,599
White	2,325	2,111	1,025	893	1,300	1,217
Black or African American	368	384	91	128	277	255
American Indian or Alaska Native	23	25	*	*	21	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	141	159	68	120	73	40
Two or More Races	100	96	35	39	66	57
Hispanic or Latino	757	740	260	232	497	508
CURRENT EMPLOYMENT						
Full-Time	1,201	1,273	260	271	941	1,002
Part-Time	1,247	1,094	669	643	579 ^a	451
Unemployed	573	492	127	101	446	390
Other ³	713	674	430	411	284	263

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.93B Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	17.1	16.1	17.2	16.3	17.0	16.0
AGE						
18	13.9	13.0	12.1	13.6	14.9	12.6
19	15.4	13.2	14.5	12.7	16.3	13.6
20	17.4	16.2	18.6	17.6	16.3	15.1
21	19.6	19.3	22.7	18.8	17.6	19.7
22	19.1	18.7	18.1	19.4	19.5	18.3
GENDER						
Male	19.9	18.6	19.9	17.9	19.9	19.0
Female	14.1	13.4	14.8	14.8	13.6	12.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	17.3	16.3	17.0	16.6	17.6	16.0
White	19.1	17.6	19.7	17.8	18.7	17.4
Black or African American	11.8	12.0	8.9	12.0	13.3	12.0
American Indian or Alaska Native	17.5	18.4	*	*	20.0	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	12.3	13.2	9.5	14.8	17.0	9.9
Two or More Races	19.8	18.4	16.7	17.7	22.0	18.9
Hispanic or Latino	16.0	15.5	18.1	14.8	15.1	15.8
CURRENT EMPLOYMENT						
Full-Time	19.1	18.4	18.9	18.1	19.1	18.5
Part-Time	17.3	15.6	17.9	16.7	16.8	14.4
Unemployed	19.1	18.0	17.3	14.9	19.7	19.0
Other ³	13.2	12.7	15.3	15.1	10.9	10.2

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.94A Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year, by Detailed Age Category: Numbers in Thousands, 2013 and 2014

Age Category	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	9,929	10,217	28,671	27,730	31,432	30,909
12-20	1,915	1,865	1,787	1,660	2,564	2,454
12	2	7	0	3	2	7
13	8	5	1	2	8	5
14	17	17	16	8	23	17
15	37	53	23	40	43	61
16	122	139	95	102	156	171
17	289	274	221	202	360	337
18	465	402	385	345	595	527
19	490	416	486 ^a	381	638	545
20	485	552	560	578	739	783
21-25	2,261	2,340	4,245	4,136	4,825	4,783
21	548	546	808	834	967	973
22	507	558	887	876	1,009	1,027
23	436	482	858	832	979	957
24	429	362	874	840	983	952
25	340	392	817	755	888	875
26-29	1,390	1,353	3,556 ^a	3,004	3,837	3,348
30-34	1,284	1,152	3,535	3,165	3,769	3,464
35-39	742	852	2,720	2,620	2,904	2,836
40-44	597	561	2,719	2,410	2,918	2,582
45-49	349 ^b	637	2,507	2,449	2,581	2,689
50-54	679	609	2,842	2,680	3,064	2,843
55-59	332	504	1,675 ^a	2,284	1,802 ^a	2,470
60-64	176	269	1,495	1,489	1,507	1,588
65 or Older	203	75	1,590	1,831	1,661	1,852

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.94B Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year, by Detailed Age Category: Percentages, 2013 and 2014

Age Category	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	3.8	3.9	10.9	10.5	12.0	11.7
12-20	5.0	4.9	4.7	4.4	6.7	6.5
12	0.0	0.2	0.0	0.1	0.0	0.2
13	0.2	0.1	0.0	0.0	0.2	0.1
14	0.4	0.4	0.4	0.2	0.5	0.4
15	0.9	1.2	0.6	0.9	1.0	1.4
16	2.9	3.2	2.2	2.3	3.6	3.9
17	7.0	6.6	5.4	4.8	8.7	8.1
18	9.8	8.7	8.1	7.5	12.6	11.5
19	11.8	10.0	11.7 ^a	9.2	15.3	13.2
20	11.3	12.7	13.0	13.2	17.2	18.0
21-25	10.5	10.7	19.7	18.9	22.3	21.9
21	12.9	12.6	19.1	19.2	22.8	22.4
22	11.3	12.4	19.8	19.4	22.5	22.8
23	10.0	11.0	19.6	18.9	22.4	21.8
24	9.8	8.4	19.9	19.4	22.4	22.0
25	8.3	9.2	19.9	17.8	21.6	20.6
26-29	8.1	7.9	20.7 ^a	17.6	22.3 ^a	19.7
30-34	6.4	5.5	17.6 ^a	15.2	18.7	16.6
35-39	3.9	4.4	14.3	13.4	15.2	14.5
40-44	2.8	2.8	12.7	11.9	13.6	12.8
45-49	1.7 ^b	3.1	12.5	11.9	12.9	13.1
50-54	2.9	2.6	12.3	11.3	13.3	12.0
55-59	1.6	2.4	8.2 ^a	10.9	8.8 ^a	11.8
60-64	1.0	1.5	8.4	8.5	8.5	9.0
65 or Older	0.5	0.2	3.7	4.1	3.8	4.1

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.95A Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	9,929	10,217	28,671	27,730	31,432	30,909
AGE						
12-17	475	495	357	356	593	598
18-25	3,701	3,709	5,675	5,441	6,797	6,639
26 or Older	5,753	6,012	22,639	21,933	24,043	23,672
GENDER						
Male	6,753	7,015	17,917	17,611	19,685	19,733
Female	3,177	3,201	10,754	10,118	11,747	11,177
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8,632	8,906	25,254	24,529	27,673	27,400
White	7,123	7,002	21,599	20,521	23,632	22,814
Black or African American	964 ^b	1,300	2,231	2,570	2,457 ^a	2,931
American Indian or Alaska Native	46	60	94	119	108	138
Native Hawaiian or Other Pacific Islander	19 ^b	66	90	127	95	152
Asian	143	223	726	715	783	805
Two or More Races	336	255	513	476	599	559
Hispanic or Latino	1,298	1,311	3,417	3,201	3,759	3,510

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.95B Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	3.8	3.9	10.9	10.5	12.0	11.7
AGE						
12-17	1.9	2.0	1.4	1.4	2.4	2.4
18-25	10.6	10.6	16.3	15.6	19.5	19.0
26 or Older	2.8	2.9	11.2	10.7	11.9	11.5
GENDER						
Male	5.3	5.5	14.1	13.7	15.5	15.4
Female	2.3	2.3	7.9	7.4	8.7	8.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	3.9	4.0	11.4	11.0	12.5	12.3
White	4.2	4.1	12.7	12.1	13.9	13.4
Black or African American	3.1 ^a	4.1	7.1	8.1	7.9 ^a	9.3
American Indian or Alaska Native	3.3	4.2	6.7	8.3	7.7	9.6
Native Hawaiian or Other Pacific Islander	2.2 ^a	6.5	10.4	12.5	10.9	14.9
Asian	1.1	1.6	5.4	5.2	5.9	5.8
Two or More Races	7.8	5.8	12.0	10.8	14.0	12.7
Hispanic or Latino	3.1	3.1	8.3	7.5	9.1	8.3

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.96A Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	9,454	9,722	28,314	27,374	30,840	30,311
GENDER						
Male	6,457	6,733	17,731	17,426	19,337	19,401
Female	2,997	2,989	10,583	9,947	11,503	10,910
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8,237	8,482	24,954	24,229	27,175	26,892
White	6,795	6,662	21,341	20,268	23,209	22,400
Black or African American	924 ^a	1,242	2,207	2,545	2,412 ^a	2,871
American Indian or Alaska Native	41	58	92	117	102	136
Native Hawaiian or Other Pacific Islander	16 ^b	65	88	*	92	150
Asian	141	215	724	705	777	793
Two or More Races	321	239	502	469	582	543
Hispanic or Latino	1,217	1,239	3,361	3,144	3,664	3,419
EDUCATION						
< High School	1,066	1,019	1,487	1,570	1,876	1,958
High School Graduate	2,841	2,874	6,382	6,158	7,342	7,227
Some College	3,189	3,466	8,310	8,462	9,102	9,466
College Graduate	2,358	2,363	12,135	11,184	12,519	11,661
CURRENT EMPLOYMENT						
Full-Time	5,229	5,704	19,318	18,864	20,464	20,365
Part-Time	2,000	1,901	4,320	3,957	4,928	4,563
Unemployed	876	699	1,211	1,096	1,464	1,294
Other ¹	1,350	1,418	3,465	3,457	3,984	4,089

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.96B Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	4.0	4.0	11.9	11.4	13.0	12.6
GENDER						
Male	5.6	5.8	15.5	15.1	16.9	16.8
Female	2.4	2.4	8.6	8.0	9.3	8.8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4.1	4.2	12.4	11.9	13.5	13.2
White	4.3	4.3	13.6	12.9	14.8	14.3
Black or African American	3.3 ^a	4.4	7.9	9.0	8.7 ^a	10.2
American Indian or Alaska Native	3.3	4.6	7.4	9.2	8.2	10.7
Native Hawaiian or Other Pacific Islander	2.1 ^a	7.1	11.3	*	11.9	16.3
Asian	1.2	1.7	6.0	5.6	6.4	6.3
Two or More Races	9.1	6.6	14.2	12.8	16.5	14.9
Hispanic or Latino	3.4	3.4	9.4	8.5	10.3	9.3
EDUCATION						
< High School	3.3	3.2	4.6	4.9	5.8	6.1
High School Graduate	4.1	4.2	9.1	8.9	10.5	10.5
Some College	5.0	5.3	13.1	12.9	14.4	14.4
College Graduate	3.3	3.2	16.9 ^a	15.2	17.4 ^a	15.9
CURRENT EMPLOYMENT						
Full-Time	4.4	4.6	16.2	15.3	17.2	16.6
Part-Time	5.9	5.8	12.8	12.0	14.6	13.8
Unemployed	7.4	6.5	10.3	10.2	12.4	12.0
Other ¹	1.8	1.9	4.7	4.7	5.5	5.6

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.97A Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	9,454	9,722	28,314	27,374	30,840	30,311
GEOGRAPHIC DIVISION						
Northeast	1,535	1,762	4,865	4,829	5,287	5,341
New England	556	583	1,462	1,563	1,592	1,752
Middle Atlantic	978	1,179	3,403	3,266	3,695	3,589
Midwest	2,241	2,262	7,035	7,054	7,675	7,732
East North Central	1,468	1,519	4,722	4,855	5,105	5,322
West North Central	773	744	2,313	2,198	2,569	2,410
South	3,276	3,491	9,613	9,419	10,508	10,536
South Atlantic	1,823	1,867	5,062	4,961	5,569	5,582
East South Central	498	538	1,157	1,360	1,293	1,536
West South Central	955	1,087	3,394	3,098	3,646	3,418
West	2,402	2,206	6,800	6,072	7,369	6,703
Mountain	762	688	2,160 ^a	1,754	2,368	1,962
Pacific	1,640	1,518	4,641	4,318	5,001	4,741
COUNTY TYPE						
Large Metro	5,093	5,503	15,829	15,861	17,128	17,428
Small Metro	3,059	2,927	8,869 ^a	7,819	9,698	8,810
250K – 1 Mil. Pop.	2,038	1,857	6,215 ^a	5,251	6,741 ^a	5,874
< 250K Pop.	1,021	1,070	2,654	2,568	2,957	2,936
Nonmetro	1,302	1,292	3,616	3,693	4,013	4,073
Urbanized	702	656	1,912	1,706	2,079	1,905
Less Urbanized	483	543	1,360	1,684	1,554	1,832
Completely Rural	117	92	345	302	380	336

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.97B Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	4.0	4.0	11.9	11.4	13.0	12.6
GEOGRAPHIC DIVISION						
Northeast	3.6	4.1	11.3	11.1	12.2	12.3
New England	4.9	5.1	12.8	13.6	14.0	15.3
Middle Atlantic	3.1	3.7	10.7	10.2	11.6	11.2
Midwest	4.4	4.4	13.8	13.8	15.1	15.1
East North Central	4.2	4.3	13.4	13.7	14.5	15.0
West North Central	5.0	4.7	14.8	14.0	16.5	15.3
South	3.7	3.9	10.9	10.5	11.9	11.8
South Atlantic	3.9	3.9	10.8	10.4	11.9	11.8
East South Central	3.6	3.8	8.3	9.6	9.2	10.9
West South Central	3.5	3.9	12.4	11.1	13.3	12.3
West	4.3	3.9	12.3 ^a	10.8	13.3	11.9
Mountain	4.5	4.0	12.8 ^a	10.2	14.1 ^a	11.5
Pacific	4.3	3.9	12.1	11.0	13.0	12.1
COUNTY TYPE						
Large Metro	4.0	4.2	12.4	12.0	13.4	13.2
Small Metro	4.3	4.2	12.4 ^a	11.1	13.6	12.5
250K – 1 Mil. Pop.	4.2	3.9	12.9 ^a	11.1	14.0 ^a	12.4
< 250K Pop.	4.4	4.7	11.4	11.2	12.8	12.9
Nonmetro	3.4	3.5	9.4	9.9	10.4	10.9
Urbanized	4.0	4.2	11.0	10.9	12.0	12.1
Less Urbanized	2.8	3.0	7.9	9.4	9.0	10.3
Completely Rural	2.9	2.4	8.6	7.9	9.4	8.7

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.98A Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status: Numbers in Thousands, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
ILLCIT DRUGS²	22,376 ^b	24,646	1,413	1,283	20,885 ^b	23,289
Marijuana and Hashish	18,048 ^b	20,357	1,057	978	16,961 ^b	19,320
Cocaine	1,505	1,491	174	151	1,325	1,331
Crack	375	347	54	50	317	294
Heroin	277 ^a	419	59	83	212 ^a	337
Hallucinogens	1,179	1,037	112	82	1,064	945
LSD	177	222	9	39	169	183
PCP	26	*	11	3	14	*
Ecstasy	611	570	60	46	549	524
Inhalants	375	397	21	4	355	386
Nonmedical Use of Psychotherapeutics ^{3,4}	5,935	5,882	555	473	5,314	5,396
Pain Relievers	4,096	3,858	381	295	3,692	3,550
OxyContin [®]	462	321	74 ^a	20	388	301
Tranquilizers	1,615	1,772	215	177	1,393	1,589
Stimulants ⁴	1,281	1,427	177	183	1,065	1,242
Methamphetamine ⁴	573	524	137	111	396	412
Sedatives	226	289	68	28	158	260
ILLCIT DRUGS OTHER THAN MARIJUANA²	7,908	7,835	654	621	7,182	7,177

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 or older, including those with unknown probation status in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.98B Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status: Percentages, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
ILLCIT DRUGS²	9.4 ^b	10.3	31.4	29.1	9.0 ^b	9.9
Marijuana and Hashish	7.6 ^b	8.5	23.5	22.2	7.3 ^b	8.2
Cocaine	0.6	0.6	3.9	3.4	0.6	0.6
Crack	0.2	0.1	1.2	1.1	0.1	0.1
Heroin	0.1 ^a	0.2	1.3	1.9	0.1	0.1
Hallucinogens	0.5	0.4	2.5	1.9	0.5	0.4
LSD	0.1	0.1	0.2	0.9	0.1	0.1
PCP	0.0	*	0.2	0.1	0.0	*
Ecstasy	0.3	0.2	1.3	1.0	0.2	0.2
Inhalants	0.2	0.2	0.5	0.1	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	2.5	2.4	12.3	10.7	2.3	2.3
Pain Relievers	1.7	1.6	8.5	6.7	1.6	1.5
OxyContin [®]	0.2	0.1	1.6 ^a	0.5	0.2	0.1
Tranquilizers	0.7	0.7	4.8	4.0	0.6	0.7
Stimulants ⁴	0.5	0.6	3.9	4.2	0.5	0.5
Methamphetamine ⁴	0.2	0.2	3.1	2.5	0.2	0.2
Sedatives	0.1	0.1	1.5	0.6	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	3.3	3.3	14.5	14.1	3.1	3.1

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 or older, including those with unknown probation status in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.99A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Probation Status: Numbers in Thousands, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
ILLCIT DRUGS²	18,073 ^a	19,122	1,238	1,148	16,793 ^a	17,907
Marijuana and Hashish	14,941 ^b	16,189	956	895	13,954 ^b	15,242
Cocaine	1,181	1,237	124	147	1,051	1,082
Crack	198	227	27	46	167	179
Heroin	235	332	59	81	170	250
Hallucinogens	1,005	970	99	81	902	880
LSD	177	220	9	39	169	182
PCP	26	8	11	3	14	*
Ecstasy	576	553	60	46	513	508
Inhalants	315	334	21	4	294	324
Nonmedical Use of Psychotherapeutics ^{3,4}	4,822	4,545	490	383	4,300	4,149
Pain Relievers	3,311	2,929	358 ^a	235	2,931	2,681
OxyContin [®]	322	234	61	20	261	214
Tranquilizers	1,373	1,416	195	144	1,172	1,266
Stimulants ⁴	1,157	1,080	177	124	977	955
Methamphetamine ⁴	453	337	137	75	312	260
Sedatives	160	165	27	3	134	161
ILLCIT DRUGS OTHER THAN MARIJUANA²	6,286	6,164	560	529	5,689	5,598

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 to 49, including those with unknown probation status in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.99B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Probation Status: Percentages, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
ILLCIT DRUGS²	13.6 ^a	14.4	32.7	31.1	13.1 ^a	13.9
Marijuana and Hashish	11.3 ^b	12.2	25.3	24.2	10.9 ^b	11.8
Cocaine	0.9	0.9	3.3	4.0	0.8	0.8
Crack	0.1	0.2	0.7	1.2	0.1	0.1
Heroin	0.2	0.2	1.6	2.2	0.1	0.2
Hallucinogens	0.8	0.7	2.6	2.2	0.7	0.7
LSD	0.1	0.2	0.2	1.0	0.1	0.1
PCP	0.0	0.0	0.3	0.1	0.0	*
Ecstasy	0.4	0.4	1.6	1.2	0.4	0.4
Inhalants	0.2	0.3	0.5	0.1	0.2	0.3
Nonmedical Use of Psychotherapeutics ^{3,4}	3.6	3.4	13.0	10.4	3.3	3.2
Pain Relievers	2.5	2.2	9.5 ^a	6.4	2.3	2.1
OxyContin [®]	0.2	0.2	1.6	0.5	0.2	0.2
Tranquilizers	1.0	1.1	5.2	3.9	0.9	1.0
Stimulants ⁴	0.9	0.8	4.7	3.3	0.8	0.7
Methamphetamine ⁴	0.3	0.3	3.6	2.0	0.2	0.2
Sedatives	0.1	0.1	0.7	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	4.7	4.6	14.8	14.3	4.4	4.3

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 to 49, including those with unknown probation status in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.100A Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	22,376 ^b	24,646	1,413	1,283	20,885 ^b	23,289
AGE						
18-49	18,073 ^a	19,122	1,238	1,148	16,793 ^a	17,907
18-25	7,490	7,676	602	526	6,869	7,132
26-34	5,699	5,728	355	390	5,341	5,320
35-49	4,883 ^b	5,718	281	233	4,583 ^b	5,455
GENDER/AGE						
Male 18 or Older	13,451 ^b	15,201	1,036	940	12,343 ^b	14,226
Male 18-49	10,908 ^a	11,670	885	823	9,987 ^a	10,813
Female 18 or Older	8,926	9,445	376	343	8,542	9,063
Female 18-49	7,166	7,452	353	325	6,806	7,095

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.100B Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	9.4 ^b	10.3	31.4	29.1	9.0 ^b	9.9
AGE						
18-49	13.6 ^a	14.4	32.7	31.1	13.1 ^a	13.9
18-25	21.5	22.0	40.2	40.1	20.7	21.3
26-34	15.3	15.1	32.6	30.9	14.7	14.6
35-49	8.1 ^b	9.5	23.5	20.7	7.8 ^b	9.2
GENDER/AGE						
Male 18 or Older	11.8 ^b	13.1	32.1	31.3	11.1 ^b	12.7
Male 18-49	16.6 ^a	17.8	32.9	33.4	15.9 ^a	17.2
Female 18 or Older	7.3	7.6	29.6	24.5	7.0	7.4
Female 18-49	10.7	11.1	32.3	26.4	10.3	10.8

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.101A Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	5,976	6,210	654	671	5,307	5,531
AGE						
18-49	5,191	5,103	625	583	4,552	4,512
18-25	2,557	2,320	366 ^b	239	2,181	2,079
26-34	1,606	1,566	199	220	1,403	1,342
35-49	1,028	1,217	59	124	968	1,090
GENDER/AGE						
Male 18 or Older	3,798	3,966	458	490	3,329	3,471
Male 18-49	3,401	3,299	429	407	2,962	2,887
Female 18 or Older	2,179	2,244	196	181	1,978	2,060
Female 18-49	1,790	1,804	196	176	1,590	1,625

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.101B Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	2.5	2.6	14.5	15.2	2.3	2.4
AGE						
18-49	3.9	3.8	16.5	15.8	3.5	3.5
18-25	7.4	6.6	24.5 ^a	18.3	6.6	6.2
26-34	4.3	4.1	18.3	17.4	3.9	3.7
35-49	1.7	2.0	5.0	11.0	1.6	1.8
GENDER/AGE						
Male 18 or Older	3.3	3.4	14.2	16.3	3.0	3.1
Male 18-49	5.2	5.0	15.9	16.5	4.7	4.6
Female 18 or Older	1.8	1.8	15.4	12.9	1.6	1.7
Female 18-49	2.7	2.7	18.0	14.3	2.4	2.5

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.102A Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	20,259	20,230	1,574	1,388	18,629	18,827
AGE						
18-49	16,082	15,434	1,431	1,217	14,596	14,203
18-25	6,014	5,696	636 ^b	481	5,364	5,213
26-34	5,185	4,732	427	403	4,753	4,319
35-49	4,884	5,007	368	334	4,479	4,671
GENDER/AGE						
Male 18 or Older	13,059	13,091	1,179	1,007	11,830	12,079
Male 18-49	10,407	9,944	1,074	852	9,283	9,088
Female 18 or Older	7,200	7,139	395	382	6,799	6,748
Female 18-49	5,675	5,490	357	366	5,313	5,115

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.102B Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	8.5	8.4	35.0	31.5	8.0	8.0
AGE						
18-49	12.1	11.6	37.8	32.9	11.4	11.0
18-25	17.3	16.3	42.5	36.7	16.2	15.6
26-34	13.9	12.5	39.3	31.9	13.1	11.8
35-49	8.1	8.3	30.8	29.6	7.6	7.9
GENDER/AGE						
Male 18 or Older	11.4	11.3	36.5	33.5	10.7	10.7
Male 18-49	15.9	15.1	39.9	34.5	14.8	14.4
Female 18 or Older	5.8	5.7	31.1	27.2	5.6	5.5
Female 18-49	8.5	8.2	32.7	29.7	8.1	7.8

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.103A Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status: Numbers in Thousands, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
ILLCIT DRUGS³	22,376 ^b	24,646	460	334	21,851 ^b	24,226
Marijuana and Hashish	18,048 ^b	20,357	342	266	17,695 ^b	20,015
Cocaine	1,505	1,491	86	44	1,413	1,439
Crack	375	347	17	22	354	322
Heroin	277 ^a	419	24	30	249 ^a	389
Hallucinogens	1,179	1,037	51	17	1,125	1,010
LSD	177	222	4	5	173	217
PCP	26	*	*	1	26	*
Ecstasy	611	570	*	10	569	560
Inhalants	375	397	15	0	361	390
Nonmedical Use of Psychotherapeutics ^{4,5}	5,935	5,882	203	131	5,675	5,730
Pain Relievers	4,096	3,858	134	74	3,949	3,762
OxyContin [®]	462	321	24	3	438	318
Tranquilizers	1,615	1,772	71	61	1,535	1,705
Stimulants ⁵	1,281	1,427	79	45	1,166	1,382
Methamphetamine ⁵	573	524	75	20	461	504
Sedatives	226	289	2	4	224	284
ILLCIT DRUGS OTHER THAN MARIJUANA³	7,908	7,835	265 ^a	150	7,580	7,640

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 or older, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.103B Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status: Percentages, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
ILLCIT DRUGS³	9.4 ^b	10.3	27.4	23.0	9.3 ^b	10.2
Marijuana and Hashish	7.6 ^b	8.5	20.4	18.4	7.5 ^b	8.4
Cocaine	0.6	0.6	5.1	3.0	0.6	0.6
Crack	0.2	0.1	1.0	1.5	0.2	0.1
Heroin	0.1 ^a	0.2	1.4	2.1	0.1 ^a	0.2
Hallucinogens	0.5	0.4	3.0	1.2	0.5	0.4
LSD	0.1	0.1	0.2	0.3	0.1	0.1
PCP	0.0	*	*	0.1	0.0	*
Ecstasy	0.3	0.2	*	0.7	0.2	0.2
Inhalants	0.2	0.2	0.9	0.0	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{4,5}	2.5	2.4	12.1	9.0	2.4	2.4
Pain Relievers	1.7	1.6	8.0	5.1	1.7	1.6
OxyContin [®]	0.2	0.1	1.5	0.2	0.2	0.1
Tranquilizers	0.7	0.7	4.2	4.2	0.7	0.7
Stimulants ⁵	0.5	0.6	4.7	3.1	0.5	0.6
Methamphetamine ⁵	0.2	0.2	4.4	1.4	0.2	0.2
Sedatives	0.1	0.1	0.1	0.3	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA³	3.3	3.3	15.8	10.3	3.2	3.2

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 or older, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.104A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Parole/Supervised Release Status: Numbers in Thousands, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
ILLCIT DRUGS³	18,073 ^a	19,122	389	306	17,654 ^a	18,751
Marijuana and Hashish	14,941 ^b	16,189	292	240	14,639 ^a	15,894
Cocaine	1,181	1,237	56	34	1,119	1,194
Crack	198	227	9	13	185	212
Heroin	235	332	24	20	207	312
Hallucinogens	1,005	970	51	16	950	945
LSD	177	220	4	5	173	215
PCP	26	8	*	1	26	7
Ecstasy	576	553	*	10	533	543
Inhalants	315	334	15	0	300	327
Nonmedical Use of Psychotherapeutics ^{4,5}	4,822	4,545	193	119	4,607	4,418
Pain Relievers	3,311	2,929	123	63	3,175	2,858
OxyContin [®]	322	234	24	3	298	231
Tranquilizers	1,373	1,416	63	61	1,300	1,349
Stimulants ⁵	1,157	1,080	79	44	1,077	1,037
Methamphetamine ⁵	453	337	75	19	377	318
Sedatives	160	165	2	3	158	161
ILLCIT DRUGS OTHER THAN MARIJUANA³	6,286	6,164	232	137	6,026	5,995

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 to 49, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.104B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Parole/Supervised Release Status: Percentages, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
ILLCIT DRUGS³	13.6 ^a	14.4	29.0	26.3	13.5 ^a	14.3
Marijuana and Hashish	11.3 ^b	12.2	21.8	20.6	11.2 ^b	12.1
Cocaine	0.9	0.9	4.2	3.0	0.9	0.9
Crack	0.1	0.2	0.7	1.1	0.1	0.2
Heroin	0.2	0.2	1.8	1.7	0.2	0.2
Hallucinogens	0.8	0.7	3.8	1.4	0.7	0.7
LSD	0.1	0.2	0.3	0.4	0.1	0.2
PCP	0.0	0.0	*	0.1	0.0	0.0
Ecstasy	0.4	0.4	*	0.9	0.4	0.4
Inhalants	0.2	0.3	1.1	0.0	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{4,5}	3.6	3.4	14.4	10.2	3.5	3.4
Pain Relievers	2.5	2.2	9.2	5.4	2.4	2.2
OxyContin [®]	0.2	0.2	1.8	0.2	0.2	0.2
Tranquilizers	1.0	1.1	4.7	5.3	1.0	1.0
Stimulants ⁵	0.9	0.8	5.9	3.7	0.8	0.8
Methamphetamine ⁵	0.3	0.3	5.6	1.7	0.3	0.2
Sedatives	0.1	0.1	0.2	0.2	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA³	4.7	4.6	17.3	11.7	4.6	4.6

*Low precision; no estimate reported.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons aged 18 to 49, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.105A Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	22,376 ^b	24,646	460	334	21,851 ^b	24,226
AGE						
18-49	18,073 ^a	19,122	389	306	17,654 ^a	18,751
18-25	7,490	7,676	152	113	7,325	7,543
26-34	5,699	5,728	*	138	5,581	5,571
35-49	4,883 ^b	5,718	*	*	4,748 ^b	5,636
GENDER/AGE						
Male 18 or Older	13,451 ^b	15,201	362 ^a	220	13,028 ^b	14,941
Male 18-49	10,908 ^a	11,670	309	192	10,573 ^a	11,451
Female 18 or Older	8,926	9,445	*	*	8,823	9,285
Female 18-49	7,166	7,452	*	*	7,081	7,300

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.105B Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	9.4 ^b	10.3	27.4	23.0	9.3 ^b	10.2
AGE						
18-49	13.6 ^a	14.4	29.0	26.3	13.5 ^a	14.3
18-25	21.5	22.0	37.9	36.6	21.4	21.8
26-34	15.3	15.1	*	33.2	15.1	14.9
35-49	8.1 ^b	9.5	*	*	7.9 ^b	9.4
GENDER/AGE						
Male 18 or Older	11.8 ^b	13.1	26.9	20.0	11.5 ^b	13.1
Male 18-49	16.6 ^a	17.8	29.4	22.0	16.4 ^a	17.7
Female 18 or Older	7.3	7.6	*	*	7.2	7.5
Female 18-49	10.7	11.1	*	*	10.6	10.9

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.106A Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	5,976	6,210	251	238	5,720	5,966
AGE						
18-49	5,191	5,103	248	210	4,939	4,886
18-25	2,557	2,320	92	57	2,460	2,263
26-34	1,606	1,566	*	104	1,500	1,458
35-49	1,028	1,217	*	*	979	1,164
GENDER/AGE						
Male 18 or Older	3,798	3,966	178	166	3,616	3,797
Male 18-49	3,401	3,299	175	152	3,223	3,143
Female 18 or Older	2,179	2,244	*	*	2,104	2,169
Female 18-49	1,790	1,804	*	*	1,716	1,743

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.106B Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	2.5	2.6	15.0	16.4	2.4	2.5
AGE						
18-49	3.9	3.8	18.5	18.0	3.8	3.7
18-25	7.4	6.6	23.1	18.2	7.2	6.6
26-34	4.3	4.1	*	25.1	4.1	3.9
35-49	1.7	2.0	*	*	1.6	2.0
GENDER/AGE						
Male 18 or Older	3.3	3.4	13.2	15.1	3.2	3.3
Male 18-49	5.2	5.0	16.6	17.4	5.0	4.9
Female 18 or Older	1.8	1.8	*	*	1.7	1.8
Female 18-49	2.7	2.7	*	*	2.6	2.6

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.107A Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	20,259	20,230	575	460	19,654	19,743
AGE						
18-49	16,082	15,434	478	394	15,575	15,028
18-25	6,014	5,696	160	123	5,845	5,573
26-34	5,185	4,732	*	176	5,007	4,545
35-49	4,884	5,007	*	*	4,723	4,910
GENDER/AGE						
Male 18 or Older	13,059	13,091	433	331	12,599	12,743
Male 18-49	10,407	9,944	368	278	10,012	9,662
Female 18 or Older	7,200	7,139	*	*	7,055	7,001
Female 18-49	5,675	5,490	*	*	5,562	5,365

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.107B Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	8.5	8.4	34.3	31.7	8.4	8.3
AGE						
18-49	12.1	11.6	35.7	33.7	11.9	11.4
18-25	17.3	16.3	40.0	39.6	17.0	16.1
26-34	13.9	12.5	*	42.4	13.6	12.2
35-49	8.1	8.3	*	*	7.9	8.2
GENDER/AGE						
Male 18 or Older	11.4	11.3	32.1	30.1	11.2	11.1
Male 18-49	15.9	15.1	35.0	31.9	15.6	14.9
Female 18 or Older	5.8	5.7	*	*	5.8	5.6
Female 18-49	8.5	8.2	*	*	8.3	8.0

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.1C Number of Days Used Marijuana in the Past Year among Past Year Users and the Number of Days Used Marijuana in the Past Month among Past Month Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	577	538	78	82	163	186	519	451
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	309	269	50	52	115	116	280	228
12-49	257	197	37	41	80	88	233	178
50-99	186	175	26	32	56	61	175	157
100-299	290	239	39	41	89	98	278	209
300 or More	261	231	24	24	88	94	241	198
TOTAL PAST MONTH USERS	442	424	56	67	135	157	407	361
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	215	190	32	41	69	82	198	161
3-5	164	143	28	29	53	54	154	131
6-19	163	182	26	33	67	71	151	161
20 or More	318	273	28	31	99	114	292	235

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.1D Number of Days Used Marijuana in the Past Year among Past Year Users and the Number of Days Used Marijuana in the Past Month among Past Month Users, by Age Group: Standard Errors of Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	0.22	0.20	0.31	0.33	0.47	0.53	0.26	0.22
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	0.76	0.63	1.16	1.29	0.76	0.84	1.26	0.94
12-49	0.69	0.50	0.99	1.08	0.62	0.68	1.14	0.76
50-99	0.53	0.46	0.73	0.92	0.47	0.50	0.89	0.71
100-299	0.76	0.56	1.02	1.10	0.67	0.73	1.29	0.85
300 or More	0.71	0.56	0.67	0.71	0.72	0.74	1.17	0.82
TOTAL PAST MONTH USERS	0.17	0.16	0.23	0.27	0.39	0.45	0.20	0.18
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	0.91	0.74	1.52	1.82	0.88	1.00	1.47	1.04
3-5	0.79	0.60	1.39	1.43	0.74	0.70	1.29	0.89
6-19	0.76	0.68	1.26	1.51	0.85	0.90	1.23	1.01
20 or More	1.13	0.86	1.35	1.41	1.09	1.19	1.80	1.25

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.1P Number of Days Used Marijuana in the Past Year among Past Year Users and the Number of Days Used Marijuana in the Past Month among Past Month Users, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	0.0214		0.4690		0.6654		0.0073	
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	0.2115		0.1416		0.2608		0.5714	
12-49	0.3984		0.1352		0.4325		0.3902	
50-99	0.3214		0.1424		0.8057		0.3729	
100-299	0.9041		0.8460		0.0551		0.5846	
300 or More	0.2061		0.2712		0.8085		0.1817	
TOTAL PAST MONTH USERS	0.0004		0.4269		0.4081		0.0004	
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	0.5683		0.3296		0.5495		0.3043	
3-5	0.3548		0.8666		0.1701		0.6224	
6-19	0.3029		0.3697		0.3632		0.1480	
20 or More	0.7065		0.0562		0.2860		0.8832	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.2C Number of Days Used Cocaine in the Past Year among Past Year Users and the Number of Days Used Cocaine in the Past Month among Past Month Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	189	186	15	23	72	81	172	165
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	146	133	13	18	60	65	132	112
12-49	84	94	5	9	29	39	78	85
50-99	54	62	*	4	19	20	49	59
100-299	61	56	4	5	21	19	57	53
300 or More	50	22	*	*	3	4	50	21
TOTAL PAST MONTH USERS	133	105	9	10	34	40	129	98
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	97	67	*	*	26	31	93	61
3-5	55	55	*	*	15	22	52	50
6-19	49	52	*	*	17	12	46	50
20 or More	55	26	*	*	4	4	*	25

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.2D Number of Days Used Cocaine in the Past Year among Past Year Users and the Number of Days Used Cocaine in the Past Month among Past Month Users, by Age Group: Standard Errors of Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	0.07	0.07	0.06	0.09	0.21	0.23	0.08	0.08
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	2.33	2.08	5.21	5.56	2.09	2.44	3.60	2.90
12-49	1.77	1.77	3.77	4.68	1.72	2.08	2.73	2.65
50-99	1.26	1.24	*	2.53	1.18	1.16	1.95	1.93
100-299	1.39	1.15	2.93	3.01	1.28	1.12	2.18	1.78
300 or More	1.16	0.47	*	*	0.17	0.24	1.93	0.76
TOTAL PAST MONTH USERS	0.05	0.04	0.04	0.04	0.10	0.11	0.06	0.05
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	4.12	3.37	*	*	4.57	4.20	5.51	4.58
3-5	3.24	2.98	*	*	3.44	3.88	4.32	4.02
6-19	3.02	2.97	*	*	3.93	2.48	3.98	4.25
20 or More	3.28	1.70	*	*	1.06	0.84	*	2.50

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.2P Number of Days Used Cocaine in the Past Year among Past Year Users and the Number of Days Used Cocaine in the Past Month among Past Month Users, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	0.2191		0.1860		0.6188		0.3133	
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	0.8055		0.6290		0.5489		0.9405	
12-49	0.9455		0.2822		0.3968		0.7011	
50-99	0.0940		*		0.7934		0.0821	
100-299	0.9075		1.0000		0.5948		0.9832	
300 or More	0.0851		*		0.7008		0.0730	
TOTAL PAST MONTH USERS	0.8370		0.7663		0.1629		0.5229	
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	0.4706		*		0.8460		0.4077	
3-5	0.2480		*		0.5013		0.3345	
6-19	0.4669		*		0.3508		0.2841	
20 or More	0.2418		*		0.7970		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.3C Number of Days Used Hallucinogens in the Past Year among Past Year Users and the Number Days Used Hallucinogens in the Past Month among Past Month Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	204	159	29	31	89	90	177	120
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	174	134	24	27	86	80	145	100
12-49	94	52	13	12	27	34	*	36
50-99	37	39	8	9	17	20	31	32
100-299	39	26	4	7	22	16	32	19
300 or More	31	14	2	2	6	5	*	13
TOTAL PAST MONTH USERS	132	88	17	19	48	47	123	74
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	119	70	16	*	43	36	*	61
3-5	43	35	4	*	14	21	*	27
6-19	24	14	5	*	15	9	*	10
20 or More	31	17	3	*	2	5	*	16

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.3D Number of Days Used Hallucinogens in the Past Year among Past Year Users and the Number Days Used Hallucinogens in the Past Month among Past Month Users, by Age Group: Standard Errors of Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	0.08	0.06	0.12	0.13	0.25	0.26	0.09	0.06
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	2.16	1.39	2.78	3.31	1.52	1.61	5.34	2.74
12-49	1.92	1.09	2.41	2.68	1.08	1.39	*	2.05
50-99	0.82	0.88	1.53	2.03	0.72	0.89	1.93	1.89
100-299	0.86	0.60	0.88	1.69	0.92	0.70	1.95	1.16
300 or More	0.69	0.33	0.39	0.56	0.24	0.22	*	0.78
TOTAL PAST MONTH USERS	0.05	0.03	0.07	0.08	0.14	0.13	0.06	0.04
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	3.98	2.93	4.03	*	3.03	3.57	*	4.94
3-5	3.07	2.53	2.49	*	2.20	3.34	*	4.39
6-19	1.81	1.16	3.00	*	2.39	1.69	*	1.82
20 or More	2.27	1.37	1.66	*	0.33	0.94	*	2.80

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.3P Number of Days Used Hallucinogens in the Past Year among Past Year Users and the Number Days Used Hallucinogens in the Past Month among Past Month Users, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	0.3865		0.1625		0.2608		0.9728	
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	0.9162		0.7428		0.1282		0.4000	
12-49	0.6337		0.2121		0.1333		*	
50-99	0.1638		0.9088		0.1321		0.4623	
100-299	0.8011		0.2241		0.4329		0.9716	
300 or More	0.4174		0.4826		0.7743		*	
TOTAL PAST MONTH USERS	0.2769		0.4918		0.0780		0.8146	
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	0.8155		*		0.2478		*	
3-5	0.3750		*		0.2177		*	
6-19	0.4750		*		0.7829		*	
20 or More	0.7530		*		0.1927		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.4C Number of Days Used Inhalants in the Past Year among Past Year Users and the Number of Days Used Inhalants in the Past Month among Past Month Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	105	95	29	36	41	37	90	85
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	74	70	25	27	35	32	*	*
12-49	66	52	12	19	17	19	*	*
50-99	25	37	9	9	7	6	*	*
100-299	16	29	9	10	7	4	*	27
300 or More	5	6	1	5	5	2	*	*
TOTAL PAST MONTH USERS	64	65	15	19	18	16	59	59
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	56	48	12	*	*	*	*	*
3-5	*	25	5	8	*	*	*	*
6-19	17	28	4	*	*	*	*	*
20 or More	6	*	*	*	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.4D Number of Days Used Inhalants in the Past Year among Past Year Users and the Number of Days Used Inhalants in the Past Month among Past Month Users, by Age Group: Standard Errors of Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	0.04	0.04	0.12	0.14	0.12	0.11	0.04	0.04
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	3.41	3.23	3.01	3.49	3.51	4.76	*	*
12-49	3.45	2.84	2.35	3.17	3.14	4.31	*	*
50-99	1.62	2.13	1.91	1.75	1.44	1.62	*	*
100-299	1.07	1.72	1.81	1.89	1.45	0.93	*	3.57
300 or More	0.31	0.35	0.17	1.02	0.97	0.52	*	*
TOTAL PAST MONTH USERS	0.02	0.02	0.06	0.07	0.05	0.05	0.03	0.03
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	5.38	5.79	5.66	*	*	*	*	*
3-5	*	4.26	4.14	4.80	*	*	*	*
6-19	3.28	4.61	3.46	*	*	*	*	*
20 or More	1.28	*	*	*	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.4P Number of Days Used Inhalants in the Past Year among Past Year Users and the Number of Days Used Inhalants in the Past Month among Past Month Users, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	0.6320		0.4803		0.0698		0.2517	
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	0.2014		0.0281		0.8229		*	
12-49	0.7901		0.0188		0.5913		*	
50-99	0.7390		0.3736		0.6935		*	
100-299	0.0646		0.5295		0.3472		*	
300 or More	0.7000		0.1475		0.4812		*	
TOTAL PAST MONTH USERS	0.6223		0.2319		0.2937		0.6026	
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	0.0266		*		*		*	
3-5	*		0.1484		*		*	
6-19	0.0730		*		*		*	
20 or More	*		*		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.5C Number of Days Used Cigarettes in the Past Month among Past Month Users and the Number of Cigarettes Smoked Per Day among Daily Smokers, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST MONTH USERS	798	700	51	52	162	185	770	651
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	221	208	28	27	83	75	199	193
3-5	233	184	21	21	63	59	222	174
6-19	271	255	21	22	74	86	254	234
20-29	290	212	20	16	66	69	277	199
30	707	578	21	27	115	127	691	545
TOTAL DAILY SMOKERS¹	656	528	21	27	109	119	648	502
NUMBER OF CIGARETTES SMOKED PER DAY AMONG DAILY SMOKERS¹								
Fewer Than 6	305	223	15	18	58	72	298	206
6-15 (1/2 Pack)	395	346	13	15	78	76	385	333
16-25 (1 Pack)	393	304	5	7	45	51	388	295
26 or More (More Than 1 Pack)	216	195	5	4	16	16	215	193
Not Reported	14	35	*	*	1	4	14	34

*Low precision; no estimate reported.

¹ Daily Cigarette Use is defined as smoking on each of the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.5D Number of Days Used Cigarettes in the Past Month among Past Month Users and the Number of Cigarettes Smoked Per Day among Daily Smokers, by Age Group: Standard Errors of Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST MONTH USERS	0.30	0.26	0.20	0.21	0.46	0.53	0.38	0.32
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	0.37	0.35	1.66	1.92	0.69	0.70	0.43	0.41
3-5	0.40	0.32	1.36	1.52	0.53	0.55	0.49	0.38
6-19	0.45	0.42	1.37	1.69	0.61	0.73	0.55	0.49
20-29	0.49	0.36	1.27	1.24	0.55	0.62	0.60	0.43
30	0.73	0.59	1.35	1.89	0.83	0.91	0.88	0.72
TOTAL DAILY SMOKERS¹	0.25	0.20	0.08	0.11	0.31	0.34	0.32	0.24
NUMBER OF CIGARETTES SMOKED PER DAY AMONG DAILY SMOKERS¹								
Fewer Than 6	0.81	0.62	3.91	4.38	1.09	1.29	0.93	0.68
6-15 (1/2 Pack)	0.95	0.81	3.66	3.92	1.12	1.28	1.09	0.91
16-25 (1 Pack)	0.91	0.77	1.94	2.21	0.85	1.06	1.06	0.87
26 or More (More Than 1 Pack)	0.61	0.56	1.64	1.36	0.35	0.37	0.71	0.64
Not Reported	0.04	0.11	*	*	0.01	0.09	0.05	0.12

*Low precision; no estimate reported.

¹ Daily Cigarette Use is defined as smoking on each of the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.5P Number of Days Used Cigarettes in the Past Month among Past Month Users and the Number of Cigarettes Smoked Per Day among Daily Smokers, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST MONTH USERS	0.2954		0.0115		0.0013		0.8733	
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	0.1959		0.8599		0.4612		0.0455	
3-5	0.7913		0.3795		0.4563		0.4485	
6-19	0.1601		0.9118		0.4048		0.1572	
20-29	0.1713		0.0786		0.5128		0.1585	
30	0.3802		0.0432		0.9722		0.1467	
TOTAL DAILY SMOKERS¹	0.1860		0.5228		0.0318		0.3138	
NUMBER OF CIGARETTES SMOKED PER DAY AMONG DAILY SMOKERS¹								
Fewer Than 6	0.9351		0.6204		0.2684		0.7711	
6-15 (1/2 Pack)	0.4285		0.6337		0.1903		0.2633	
16-25 (1 Pack)	0.5559		0.7182		0.4254		0.4301	
26 or More (More Than 1 Pack)	0.6705		0.6264		0.1015		0.7322	
Not Reported	0.1962		*		0.1862		0.2368	

*Low precision; no estimate reported.

¹ Daily Cigarette Use is defined as smoking on each of the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.6C Number of Days Used Alcohol in the Past Year among Past Year Users and the Number of Days Used Alcohol in the Past Month among Past Month Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	1,007	833	100	103	158	165	954	764
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	764	596	88	90	145	143	726	558
12-49	789	656	59	61	173	186	750	616
50-99	643	527	39	37	131	129	618	504
100-299	1,010	773	30	34	204	202	967	733
300 or More	417	377	10	14	40	48	413	374
TOTAL PAST MONTH USERS	1,076	869	73	81	183	197	1,021	802
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	738	601	56	63	146	152	711	588
3-5	698	602	41	44	149	168	679	570
6-19	833	665	29	31	179	178	794	644
20 or More	643	547	9	11	74	79	638	541

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.6D Number of Days Used Alcohol in the Past Year among Past Year Users and the Number of Days Used Alcohol in the Past Month among Past Month Users, by Age Group: Standard Errors of Percentages, 2013 and 2014

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	0.38	0.31	0.40	0.42	0.45	0.47	0.47	0.37
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	0.38	0.29	0.92	1.01	0.47	0.48	0.45	0.34
12-49	0.38	0.30	0.79	0.85	0.48	0.53	0.45	0.35
50-99	0.32	0.26	0.60	0.59	0.39	0.41	0.39	0.30
100-299	0.43	0.34	0.47	0.54	0.53	0.56	0.51	0.40
300 or More	0.23	0.20	0.16	0.23	0.15	0.18	0.28	0.24
TOTAL PAST MONTH USERS	0.41	0.33	0.29	0.33	0.53	0.56	0.50	0.39
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	0.45	0.35	1.25	1.43	0.55	0.60	0.53	0.41
3-5	0.41	0.36	1.15	1.28	0.51	0.62	0.49	0.42
6-19	0.45	0.37	0.91	0.99	0.55	0.62	0.53	0.44
20 or More	0.41	0.33	0.31	0.37	0.32	0.35	0.49	0.39

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.6P Number of Days Used Alcohol in the Past Year among Past Year Users and the Number of Days Used Alcohol in the Past Month among Past Month Users, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Frequency of Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR USERS	0.5535		0.2985		0.6344		0.4515	
NUMBER OF DAYS USED IN PAST YEAR AMONG PAST YEAR USERS								
1-11	0.0405		0.4001		0.9111		0.0479	
12-49	0.6900		0.6569		0.2654		0.8607	
50-99	0.0464		0.9217		0.2240		0.0730	
100-299	0.8877		0.7418		0.8588		0.9234	
300 or More	0.3556		0.2529		0.4503		0.4545	
TOTAL PAST MONTH USERS	0.3203		0.9818		0.9625		0.3269	
NUMBER OF DAYS USED IN PAST MONTH AMONG PAST MONTH USERS								
1-2	0.4292		0.7070		0.5372		0.5036	
3-5	0.9848		0.5478		0.6503		0.9532	
6-19	0.7594		0.8074		0.7925		0.7034	
20 or More	0.2215		0.9867		0.8664		0.2278	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.7C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	372	339	355	326
Marijuana and Hashish	344	305	296	282
Cocaine	112	89	73	54
Crack	54	49	35	25
Heroin	38	51	18	21
Hallucinogens	79	68	98	50
LSD	21	30	31	28
PCP	11	4	10	*
Ecstasy	57	48	53	40
Inhalants	46	37	44	52
Nonmedical Use of Psychotherapeutics ^{2,3}	190	154	171	160
Pain Relievers	156	120	135	128
OxyContin [®]	59	33	56	37
Tranquilizers	99	88	69	76
Stimulants ³	98	82	60	75
Methamphetamine ³	76	55	37	46
Sedatives	27	43	62	31
ILLCIT DRUGS OTHER THAN MARIJUANA¹	226	190	208	176

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.7D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	0.58	0.52	0.17	0.15
Marijuana and Hashish	0.54	0.48	0.14	0.13
Cocaine	0.20	0.16	0.04	0.03
Crack	0.10	0.09	0.02	0.01
Heroin	0.07	0.09	0.01	0.01
Hallucinogens	0.14	0.12	0.05	0.02
LSD	0.04	0.05	0.01	0.01
PCP	0.02	0.01	0.00	*
Ecstasy	0.10	0.09	0.03	0.02
Inhalants	0.08	0.07	0.02	0.02
Nonmedical Use of Psychotherapeutics ^{2,3}	0.33	0.26	0.08	0.08
Pain Relievers	0.27	0.21	0.07	0.06
OxyContin [®]	0.11	0.06	0.03	0.02
Tranquilizers	0.18	0.16	0.03	0.04
Stimulants ³	0.17	0.15	0.03	0.04
Methamphetamine ³	0.14	0.10	0.02	0.02
Sedatives	0.05	0.08	0.03	0.01
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.39	0.32	0.10	0.08

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.7P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	0.1923		0.0002	
Marijuana and Hashish	0.0611		0.0001	
Cocaine	0.9459		0.9616	
Crack	0.9368		0.5270	
Heroin	0.0300		0.7319	
Hallucinogens	0.7683		0.2456	
LSD	0.1499		0.8755	
PCP	0.4397		*	
Ecstasy	0.7873		0.6537	
Inhalants	0.6484		0.2797	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.1616		0.1162	
Pain Relievers	0.0682		0.4382	
OxyContin [®]	0.0418		0.9729	
Tranquilizers	0.6482		0.2845	
Stimulants ³	0.6663		0.0031	
Methamphetamine ³	0.1962		0.1036	
Sedatives	0.0341		0.6535	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.2387		0.1793	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.8C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	36	41	55	65
Marijuana and Hashish	34	40	47	57
Cocaine	7	9	6	5
Crack	1	4	1	*
Heroin	4	6	2	2
Hallucinogens	14	13	10	14
LSD	9	10	4	10
PCP	3	2	1	*
Ecstasy	7	8	6	7
Inhalants	6	8	14	17
Nonmedical Use of Psychotherapeutics ^{2,3}	17	20	28	35
Pain Relievers	15	16	26	29
OxyContin [®]	4	5	5	8
Tranquilizers	9	11	8	12
Stimulants ³	8	11	9	20
Methamphetamine ³	4	8	4	9
Sedatives	3	3	6	10
ILLCIT DRUGS OTHER THAN MARIJUANA¹	22	25	33	39

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.8D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	1.70	2.15	0.22	0.26
Marijuana and Hashish	1.74	2.15	0.20	0.24
Cocaine	0.48	0.71	0.03	0.02
Crack	0.04	0.35	0.01	*
Heroin	0.32	0.47	0.01	0.01
Hallucinogens	0.93	1.06	0.04	0.06
LSD	0.63	0.83	0.02	0.04
PCP	0.22	0.17	0.01	*
Ecstasy	0.49	0.63	0.02	0.03
Inhalants	0.45	0.63	0.06	0.07
Nonmedical Use of Psychotherapeutics ^{2,3}	1.15	1.50	0.12	0.15
Pain Relievers	1.00	1.25	0.11	0.12
OxyContin [®]	0.29	0.39	0.02	0.03
Tranquilizers	0.60	0.89	0.04	0.05
Stimulants ³	0.55	0.89	0.04	0.08
Methamphetamine ³	0.31	0.64	0.02	0.04
Sedatives	0.22	0.28	0.02	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1.38	1.78	0.13	0.16

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.8P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	0.3713		0.0148	
Marijuana and Hashish	0.4144		0.0877	
Cocaine	0.3702		0.2136	
Crack	0.0980		*	
Heroin	0.5090		0.9917	
Hallucinogens	0.3764		0.5563	
LSD	0.8405		0.1288	
PCP	0.7313		*	
Ecstasy	0.9707		0.5181	
Inhalants	0.2218		0.3593	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.5723		0.0108	
Pain Relievers	1.0000		0.1476	
OxyContin [®]	0.7727		0.5331	
Tranquilizers	0.4248		0.5221	
Stimulants ³	0.2112		0.0007	
Methamphetamine ³	0.2941		0.1243	
Sedatives	0.8838		0.1563	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.4454		0.0058	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.9C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	126	142	103	121
Marijuana and Hashish	120	134	94	115
Cocaine	29	34	16	19
Crack	12	5	*	8
Heroin	16	16	2	2
Hallucinogens	39	44	28	19
LSD	15	17	9	8
PCP	1	2	2	2
Ecstasy	28	31	16	16
Inhalants	15	13	10	10
Nonmedical Use of Psychotherapeutics ^{2,3}	59	59	41	45
Pain Relievers	51	49	35	35
OxyContin [®]	17	18	7	8
Tranquilizers	30	30	18	19
Stimulants ³	29	27	22	23
Methamphetamine ³	18	13	6	8
Sedatives	6	9	7	11
ILLCIT DRUGS OTHER THAN MARIJUANA¹	72	80	50	52

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.9D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	0.84	0.92	0.37	0.43
Marijuana and Hashish	0.82	0.92	0.34	0.42
Cocaine	0.27	0.33	0.07	0.08
Crack	0.11	0.05	*	0.03
Heroin	0.15	0.17	0.01	0.01
Hallucinogens	0.36	0.43	0.12	0.07
LSD	0.14	0.17	0.04	0.03
PCP	0.01	0.02	0.01	0.01
Ecstasy	0.26	0.31	0.07	0.06
Inhalants	0.14	0.13	0.04	0.04
Nonmedical Use of Psychotherapeutics ^{2,3}	0.50	0.54	0.17	0.18
Pain Relievers	0.44	0.46	0.14	0.14
OxyContin [®]	0.16	0.18	0.03	0.03
Tranquilizers	0.27	0.29	0.07	0.08
Stimulants ³	0.27	0.27	0.09	0.09
Methamphetamine ³	0.16	0.13	0.02	0.03
Sedatives	0.06	0.09	0.03	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.59	0.71	0.20	0.20

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.9P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	0.0468		0.3286	
Marijuana and Hashish	0.1038		0.1636	
Cocaine	0.2385		0.0721	
Crack	0.1045		*	
Heroin	0.9252		0.5611	
Hallucinogens	0.7626		0.0274	
LSD	0.2415		0.4617	
PCP	0.6223		0.8495	
Ecstasy	0.5511		0.5814	
Inhalants	0.3609		0.7551	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.5194		0.8732	
Pain Relievers	0.2647		0.3657	
OxyContin [®]	0.9850		0.9709	
Tranquilizers	0.8571		0.8088	
Stimulants ³	0.2343		0.8782	
Methamphetamine ³	0.1019		0.5151	
Sedatives	0.2414		0.4500	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.9605		0.9458	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.10C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	348	295	320	293
Marijuana and Hashish	317	260	268	246
Cocaine	108	81	71	50
Crack	52	49	35	24
Heroin	35	48	18	21
Hallucinogens	68	53	94	44
LSD	12	22	29	25
PCP	11	2	10	*
Ecstasy	49	35	52	38
Inhalants	42	34	40	49
Nonmedical Use of Psychotherapeutics ^{2,3}	181	143	163	149
Pain Relievers	148	110	131	119
OxyContin [®]	55	27	55	36
Tranquilizers	93	81	65	73
Stimulants ³	92	79	53	69
Methamphetamine ³	73	53	36	45
Sedatives	27	42	62	27
ILLCIT DRUGS OTHER THAN MARIJUANA¹	213	174	199	166

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.10D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	0.70	0.59	0.20	0.18
Marijuana and Hashish	0.64	0.53	0.17	0.15
Cocaine	0.25	0.18	0.04	0.03
Crack	0.12	0.11	0.02	0.01
Heroin	0.08	0.11	0.01	0.01
Hallucinogens	0.15	0.12	0.06	0.03
LSD	0.03	0.05	0.02	0.02
PCP	0.03	0.01	0.01	*
Ecstasy	0.11	0.08	0.03	0.02
Inhalants	0.10	0.08	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{2,3}	0.40	0.31	0.10	0.09
Pain Relievers	0.33	0.24	0.08	0.07
OxyContin [®]	0.13	0.06	0.03	0.02
Tranquilizers	0.21	0.18	0.04	0.05
Stimulants ³	0.21	0.18	0.03	0.04
Methamphetamine ³	0.17	0.12	0.02	0.03
Sedatives	0.06	0.10	0.04	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.47	0.38	0.13	0.10

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.10P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
ILLCIT DRUGS¹	0.2221		0.0010	
Marijuana and Hashish	0.0480		0.0006	
Cocaine	0.6527		0.6426	
Crack	0.8230		0.3408	
Heroin	0.0188		0.6905	
Hallucinogens	0.6860		0.5208	
LSD	0.1973		0.9601	
PCP	0.4375		*	
Ecstasy	0.7562		0.8369	
Inhalants	0.7725		0.3480	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.2624		0.2406	
Pain Relievers	0.1715		0.4601	
OxyContin [®]	0.0276		0.9558	
Tranquilizers	0.5747		0.3398	
Stimulants ³	0.9202		0.0191	
Methamphetamine ³	0.2974		0.1932	
Sedatives	0.0546		0.3995	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.2984		0.3773	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.11C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	232	212	267	243	270	236	245	217
Marijuana and Hashish	209	180	244	214	239	214	207	180
Cocaine	96	76	58	56	58	34	45	24
Crack	35	38	26	32	38	21	27	9
Heroin	13	25	21	30	22	28	26	23
Hallucinogens	57	54	57	47	101	34	35	23
LSD	21	23	29	18	5	19	8	12
PCP	1	4	11	3	*	*	10	*
Ecstasy	45	45	42	32	66	27	13	16
Inhalants	31	29	25	29	36	35	34	30
Nonmedical Use of Psychotherapeutics ^{3,4}	124	105	129	103	119	109	143	123
Pain Relievers	108	85	104	82	90	74	117	108
OxyContin [®]	28	35	38	25	18	20	64	14
Tranquilizers	59	50	69	66	45	65	62	47
Stimulants ⁴	60	49	65	51	56	57	49	61
Methamphetamine ⁴	42	23	40	26	43	37	44	51
Sedatives	20	23	15	31	44	24	44	28
ILLCIT DRUGS OTHER THAN MARIJUANA²	147	135	145	131	165	117	154	128

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.11D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	1.17	0.99	0.57	0.49	0.34	0.29	0.19	0.17
Marijuana and Hashish	1.09	0.90	0.53	0.44	0.30	0.26	0.16	0.14
Cocaine	0.56	0.45	0.13	0.12	0.08	0.04	0.04	0.02
Crack	0.21	0.23	0.06	0.07	0.05	0.03	0.02	0.01
Heroin	0.08	0.15	0.05	0.07	0.03	0.04	0.02	0.02
Hallucinogens	0.34	0.32	0.13	0.10	0.13	0.04	0.03	0.02
LSD	0.13	0.14	0.07	0.04	0.01	0.02	0.01	0.01
PCP	0.01	0.02	0.03	0.01	*	*	0.01	*
Ecstasy	0.27	0.27	0.10	0.07	0.09	0.03	0.01	0.01
Inhalants	0.19	0.18	0.06	0.07	0.05	0.04	0.03	0.02
Nonmedical Use of Psychotherapeutics ^{3,4}	0.71	0.59	0.29	0.22	0.16	0.14	0.11	0.10
Pain Relievers	0.63	0.50	0.24	0.18	0.12	0.09	0.09	0.09
OxyContin [®]	0.17	0.21	0.09	0.06	0.02	0.03	0.05	0.01
Tranquilizers	0.36	0.29	0.16	0.15	0.06	0.08	0.05	0.04
Stimulants ⁴	0.36	0.29	0.15	0.11	0.07	0.07	0.04	0.05
Methamphetamine ⁴	0.25	0.14	0.09	0.06	0.06	0.05	0.03	0.04
Sedatives	0.12	0.14	0.03	0.07	0.06	0.03	0.04	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.83	0.73	0.33	0.28	0.21	0.15	0.12	0.10

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.11P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	0.4600		0.2968		0.0069		0.0382	
Marijuana and Hashish	0.4070		0.4068		0.0031		0.0042	
Cocaine	0.9517		0.2051		0.2895		0.2043	
Crack	0.4013		0.5030		0.3574		0.0480	
Heroin	0.0516		0.1766		0.5093		0.6933	
Hallucinogens	0.9464		0.4374		0.4953		0.2799	
LSD	0.4204		0.5053		0.0289		0.4392	
PCP	0.1544		0.3649		*		*	
Ecstasy	0.6253		0.5658		0.9881		0.7101	
Inhalants	0.8608		0.3836		0.3609		0.5588	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.6347		0.1561		0.1366		0.6412	
Pain Relievers	0.8416		0.0847		0.9863		0.8209	
OxyContin [®]	0.7052		0.4960		0.7180		0.0383	
Tranquilizers	0.7056		0.9440		0.0206		0.4978	
Stimulants ⁴	0.8330		0.7891		0.0863		0.2646	
Methamphetamine ⁴	0.2970		0.2759		0.3638		0.7698	
Sedatives	0.2784		0.1618		0.8874		0.9575	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.4720		0.4195		0.6294		0.6240	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.12C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	17	22	35	43	24	32	47	49
Marijuana and Hashish	17	21	33	41	24	28	37	40
Cocaine	6	8	3	6	1	1	6	*
Crack	0	*	0	3	1	*	*	*
Heroin	*	*	3	2	1	5	3	2
Hallucinogens	7	10	10	9	5	10	10	9
LSD	6	7	7	7	1	9	5	6
PCP	1	*	3	2	1	*	1	*
Ecstasy	4	6	6	6	1	5	5	4
Inhalants	4	2	6	7	3	8	13	15
Nonmedical Use of Psychotherapeutics ^{3,4}	8	13	14	20	11	17	27	30
Pain Relievers	7	10	13	17	8	13	24	25
OxyContin [®]	3	5	4	5	1	3	3	5
Tranquilizers	3	9	8	6	6	4	8	11
Stimulants ⁴	3	9	6	8	5	12	8	15
Methamphetamine ⁴	2	*	3	4	1	8	5	7
Sedatives	2	*	2	4	1	3	6	9
ILLCIT DRUGS OTHER THAN MARIJUANA²	11	16	16	23	12	21	31	33

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.12D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	4.26	3.71	2.08	2.33	1.61	1.94	0.20	0.22
Marijuana and Hashish	4.27	4.41	2.04	2.35	1.58	1.81	0.17	0.18
Cocaine	2.02	2.84	0.21	0.51	0.11	0.09	0.03	*
Crack	0.03	*	0.03	0.26	0.09	*	*	*
Heroin	*	*	0.27	0.17	0.09	0.36	0.01	0.01
Hallucinogens	2.39	3.50	0.80	0.75	0.38	0.74	0.05	0.04
LSD	1.90	2.70	0.51	0.52	0.08	0.65	0.02	0.03
PCP	0.21	*	0.23	0.13	0.05	*	0.01	*
Ecstasy	1.43	2.07	0.48	0.47	0.10	0.37	0.02	0.02
Inhalants	1.24	0.78	0.44	0.59	0.20	0.55	0.06	0.07
Nonmedical Use of Psychotherapeutics ^{3,4}	2.62	4.34	1.10	1.46	0.80	1.19	0.12	0.13
Pain Relievers	2.36	3.53	0.99	1.29	0.61	0.95	0.11	0.11
OxyContin [®]	0.86	1.72	0.36	0.36	0.11	0.25	0.02	0.02
Tranquilizers	1.03	3.07	0.64	0.49	0.46	0.32	0.03	0.05
Stimulants ⁴	1.15	3.33	0.44	0.65	0.36	0.88	0.04	0.07
Methamphetamine ⁴	0.80	*	0.23	0.28	0.07	0.56	0.02	0.03
Sedatives	0.65	*	0.16	0.31	0.08	0.21	0.03	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA²	3.28	4.87	1.25	1.66	0.87	1.44	0.14	0.15

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.12P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	0.0361		0.2206		0.0808		0.7844	
Marijuana and Hashish	0.0796		0.3397		0.2000		0.6668	
Cocaine	0.6183		0.1403		0.6693		*	
Crack	*		0.1623		*		*	
Heroin	*		0.7827		0.3471		0.6401	
Hallucinogens	0.4489		0.0845		0.1994		0.3171	
LSD	0.5531		0.4086		0.0606		0.8349	
PCP	*		0.5095		*		*	
Ecstasy	0.8535		0.3579		0.3583		0.3296	
Inhalants	0.2368		0.3007		0.0102		0.8408	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.1878		0.2525		0.1571		0.3082	
Pain Relievers	0.6014		0.6228		0.5155		0.5560	
OxyContin [®]	0.5198		0.6209		0.2644		0.7434	
Tranquilizers	0.0212		0.2366		0.7510		0.4901	
Stimulants ⁴	0.0832		0.1942		0.0443		0.0679	
Methamphetamine ⁴	*		0.7814		0.0467		0.5707	
Sedatives	*		0.3608		0.6859		0.2596	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.2120		0.2363		0.0185		0.4668	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.13C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	82	94	103	109	64	74	61	68
Marijuana and Hashish	77	88	99	104	60	70	51	62
Cocaine	27	31	18	20	7	12	6	8
Crack	10	5	4	7	1	3	5	2
Heroin	6	6	5	4	8	6	12	14
Hallucinogens	30	29	27	28	11	15	19	8
LSD	10	12	13	11	5	9	5	4
PCP	1	2	*	1	*	*	2	*
Ecstasy	22	21	21	22	4	10	7	7
Inhalants	9	9	10	11	4	6	12	5
Nonmedical Use of Psychotherapeutics ^{3,4}	42	45	40	41	27	30	28	27
Pain Relievers	35	34	33	34	23	25	24	24
OxyContin [®]	13	14	12	10	6	9	4	5
Tranquilizers	21	24	22	20	11	13	13	11
Stimulants ⁴	27	23	19	21	13	13	11	11
Methamphetamine ⁴	15	11	7	6	6	7	4	4
Sedatives	6	5	5	7	2	8	4	7
ILLCIT DRUGS OTHER THAN MARIJUANA²	50	58	48	52	30	34	35	34

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.13D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	1.35	1.53	0.86	0.89	0.79	0.87	0.41	0.45
Marijuana and Hashish	1.37	1.53	0.83	0.88	0.74	0.83	0.36	0.42
Cocaine	0.68	0.77	0.19	0.21	0.09	0.15	0.04	0.06
Crack	0.26	0.14	0.04	0.07	0.01	0.04	0.04	0.02
Heroin	0.14	0.16	0.05	0.04	0.11	0.08	0.08	0.10
Hallucinogens	0.73	0.74	0.29	0.29	0.14	0.20	0.13	0.06
LSD	0.25	0.31	0.14	0.11	0.06	0.12	0.03	0.02
PCP	0.03	0.06	*	0.02	*	*	0.01	*
Ecstasy	0.56	0.52	0.22	0.23	0.05	0.13	0.05	0.05
Inhalants	0.23	0.25	0.11	0.11	0.06	0.08	0.08	0.04
Nonmedical Use of Psychotherapeutics ^{3,4}	0.94	1.03	0.42	0.42	0.35	0.39	0.19	0.19
Pain Relievers	0.84	0.84	0.34	0.35	0.30	0.32	0.17	0.17
OxyContin [®]	0.32	0.36	0.13	0.11	0.08	0.11	0.03	0.03
Tranquilizers	0.51	0.60	0.23	0.21	0.14	0.17	0.09	0.08
Stimulants ⁴	0.64	0.57	0.20	0.22	0.18	0.17	0.08	0.08
Methamphetamine ⁴	0.37	0.30	0.08	0.07	0.08	0.09	0.03	0.03
Sedatives	0.16	0.14	0.06	0.08	0.03	0.10	0.03	0.05
ILLCIT DRUGS OTHER THAN MARIJUANA²	1.09	1.26	0.49	0.52	0.39	0.42	0.24	0.23

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.13P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	0.2741		0.2087		0.0252		0.5385	
Marijuana and Hashish	0.3272		0.1478		0.0238		0.1980	
Cocaine	0.3590		0.5686		0.0525		0.6797	
Crack	0.9077		0.8697		0.6452		0.1297	
Heroin	0.9490		0.1689		0.4000		0.7215	
Hallucinogens	0.2729		0.4171		0.6745		0.0340	
LSD	0.4623		0.8754		0.4835		0.7858	
PCP	0.6057		*		*		*	
Ecstasy	0.2943		0.4199		0.3401		0.8323	
Inhalants	0.6724		0.7431		0.6736		0.1591	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.6802		0.1673		0.6171		0.0745	
Pain Relievers	0.5026		0.2626		0.7848		0.0801	
OxyContin [®]	0.4725		0.9768		0.7960		0.3168	
Tranquilizers	0.2866		0.2446		0.7800		0.2776	
Stimulants ⁴	0.2594		0.9503		0.8854		0.3118	
Methamphetamine ⁴	0.3988		0.1359		0.9803		0.6031	
Sedatives	0.9178		0.5551		0.1439		0.4920	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.4926		0.2952		0.6532		0.1353	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.14C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	217	179	237	207	261	223	231	192
Marijuana and Hashish	193	150	217	177	229	201	191	162
Cocaine	93	68	54	52	58	32	43	22
Crack	33	38	26	31	38	21	27	9
Heroin	12	24	20	30	21	27	23	18
Hallucinogens	48	46	49	39	100	29	28	21
LSD	17	19	25	13	*	14	*	10
PCP	*	3	11	*	*	*	10	*
Ecstasy	39	39	35	23	66	24	10	13
Inhalants	30	28	22	26	35	34	29	26
Nonmedical Use of Psychotherapeutics ^{3,4}	118	92	124	92	117	102	138	116
Pain Relievers	102	77	99	73	86	70	114	101
OxyContin [®]	22	32	36	22	18	18	63	12
Tranquilizers	55	43	64	63	44	64	60	44
Stimulants ⁴	52	42	62	46	53	55	46	58
Methamphetamine ⁴	38	20	38	24	43	36	43	50
Sedatives	19	22	14	30	45	22	44	26
ILLCIT DRUGS OTHER THAN MARIJUANA²	138	120	138	115	165	110	147	120

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.14D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	1.49	1.17	0.67	0.56	0.37	0.30	0.25	0.21
Marijuana and Hashish	1.37	1.04	0.62	0.49	0.33	0.28	0.21	0.18
Cocaine	0.73	0.54	0.16	0.15	0.08	0.05	0.05	0.02
Crack	0.27	0.30	0.08	0.09	0.06	0.03	0.03	0.01
Heroin	0.10	0.20	0.06	0.09	0.03	0.04	0.03	0.02
Hallucinogens	0.39	0.37	0.15	0.11	0.15	0.04	0.03	0.02
LSD	0.14	0.15	0.08	0.04	*	0.02	*	0.01
PCP	*	0.02	0.03	*	*	*	0.01	*
Ecstasy	0.32	0.32	0.11	0.07	0.10	0.04	0.01	0.01
Inhalants	0.24	0.22	0.07	0.08	0.05	0.05	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{3,4}	0.90	0.70	0.37	0.26	0.17	0.14	0.15	0.13
Pain Relievers	0.79	0.60	0.30	0.21	0.13	0.10	0.13	0.11
OxyContin [®]	0.18	0.26	0.11	0.06	0.03	0.03	0.07	0.01
Tranquilizers	0.44	0.34	0.19	0.18	0.06	0.09	0.07	0.05
Stimulants ⁴	0.42	0.33	0.19	0.13	0.08	0.08	0.05	0.07
Methamphetamine ⁴	0.30	0.16	0.11	0.07	0.06	0.05	0.05	0.06
Sedatives	0.15	0.18	0.04	0.09	0.07	0.03	0.05	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA²	1.05	0.89	0.41	0.33	0.24	0.16	0.16	0.13

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.14P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS²	0.6585		0.1071		0.0369		0.0468	
Marijuana and Hashish	0.5669		0.1256		0.0175		0.0052	
Cocaine	0.7865		0.3053		0.1435		0.2764	
Crack	0.4176		0.5926		0.3554		0.0907	
Heroin	0.0489		0.1026		0.4182		0.7594	
Hallucinogens	0.4681		0.9903		0.3673		0.7381	
LSD	0.6623		0.6772		*		*	
PCP	*		*		*		*	
Ecstasy	0.9633		0.9551		0.8222		0.3398	
Inhalants	0.9292		0.3893		0.6477		0.7440	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.7658		0.2471		0.2080		0.7402	
Pain Relievers	0.6407		0.1342		0.9793		0.9889	
OxyContin [®]	0.4670		0.5112		0.8939		0.0266	
Tranquilizers	0.7612		0.6531		0.0186		0.5593	
Stimulants ⁴	0.5284		0.6760		0.1543		0.3318	
Methamphetamine ⁴	0.4061		0.3928		0.5090		0.7871	
Sedatives	0.2283		0.2352		0.6906		0.7696	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.6253		0.5379		0.8934		0.7529	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.15C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
ILLCIT DRUGS¹	458	441	106	123	439	411
Marijuana and Hashish	407	393	102	114	387	360
Cocaine (Including Crack)	127	104	19	23	125	102
Heroin	33	49	5	7	33	48
Hallucinogens	129	85	30	29	126	78
Inhalants	54	57	11	15	52	54
Pain Relievers ²	174	141	35	37	173	137
Tranquilizers ²	100	107	20	22	99	102
Stimulants ^{2,3}	106	91	23	24	103	87
Methamphetamine ³	73	50	9	13	72	48
Sedatives ²	50	45	6	6	51	44
ILLCIT DRUGS OTHER THAN MARIJUANA¹	269	233	50	60	267	224

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.15D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
ILLCIT DRUGS¹	0.32	0.29	0.87	0.98	0.32	0.29
Marijuana and Hashish	0.28	0.26	0.87	0.94	0.29	0.26
Cocaine (Including Crack)	0.09	0.07	0.22	0.25	0.10	0.08
Heroin	0.02	0.03	0.05	0.09	0.03	0.04
Hallucinogens	0.09	0.06	0.34	0.33	0.10	0.06
Inhalants	0.04	0.04	0.13	0.17	0.04	0.04
Pain Relievers ²	0.13	0.10	0.39	0.41	0.13	0.10
Tranquilizers ²	0.07	0.08	0.23	0.25	0.08	0.08
Stimulants ^{2,3}	0.08	0.06	0.25	0.27	0.08	0.07
Methamphetamine ³	0.05	0.04	0.10	0.15	0.06	0.04
Sedatives ²	0.04	0.03	0.07	0.07	0.04	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.19	0.16	0.53	0.62	0.20	0.16

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.15P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
ILLCIT DRUGS¹	0.0229		0.4241		0.0211	
Marijuana and Hashish	0.0173		0.6057		0.0126	
Cocaine (Including Crack)	0.9198		0.4048		0.9700	
Heroin	0.0310		0.5531		0.0360	
Hallucinogens	0.3747		0.0549		0.7209	
Inhalants	0.3716		0.1626		0.5551	
Pain Relievers ²	0.3195		0.7597		0.3565	
Tranquilizers ²	0.1787		0.7227		0.1415	
Stimulants ^{2,3}	0.3914		0.4718		0.4642	
Methamphetamine ³	0.5480		0.3843		0.4452	
Sedatives ²	0.2863		0.8119		0.2744	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.9507		0.7870		0.9671	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.16C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
ILLCIT DRUGS¹	300	281	69	71	288	264
Marijuana and Hashish	284	263	69	68	271	246
Cocaine (Including Crack)	79	61	6	11	79	60
Heroin	25	26	1	5	25	25
Hallucinogens	75	29	6	8	75	28
Inhalants	9	9	2	4	9	8
Pain Relievers ²	90	66	7	11	89	65
Tranquilizers ²	31	35	7	6	30	35
Stimulants ²	17	18	9	4	15	18
Methamphetamine ³	40	31	5	7	40	31
Sedatives ²	19	14	4	2	19	14
ILLCIT DRUGS OTHER THAN MARIJUANA¹	140	108	15	20	139	106

*Low precision; no estimate reported.

NOTE: Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.16D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
ILLCIT DRUGS¹	0.21	0.19	0.72	0.76	0.22	0.20
Marijuana and Hashish	0.20	0.18	0.72	0.74	0.21	0.18
Cocaine (Including Crack)	0.06	0.04	0.07	0.12	0.06	0.05
Heroin	0.02	0.02	0.01	0.06	0.02	0.02
Hallucinogens	0.05	0.02	0.07	0.10	0.06	0.02
Inhalants	0.01	0.01	0.02	0.05	0.01	0.01
Pain Relievers ²	0.07	0.05	0.09	0.12	0.07	0.05
Tranquilizers ²	0.02	0.03	0.08	0.07	0.02	0.03
Stimulants ²	0.01	0.01	0.10	0.05	0.01	0.01
Methamphetamine ³	0.03	0.02	0.05	0.09	0.03	0.02
Sedatives ²	0.01	0.01	0.05	0.03	0.01	0.01
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.10	0.08	0.18	0.24	0.11	0.08

*Low precision; no estimate reported.

NOTE: Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.16P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
ILLCIT DRUGS¹	0.1024		0.1464		0.0367	
Marijuana and Hashish	0.1513		0.0864		0.0469	
Cocaine (Including Crack)	0.8091		0.2111		0.6887	
Heroin	0.8791		0.2755		1.0000	
Hallucinogens	0.9516		0.4526		0.9747	
Inhalants	0.6380		0.2713		0.9387	
Pain Relievers ²	0.7104		0.4840		0.7704	
Tranquilizers ²	0.0638		0.9321		0.0602	
Stimulants ²	0.5479		0.2791		0.2643	
Methamphetamine ³	0.9688		0.4238		0.9242	
Sedatives ²	0.6935		0.7649		0.6422	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.4257		0.1764		0.5394	

*Low precision; no estimate reported.

NOTE: Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.17C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLCIT DRUGS¹	458	441	46	60	156	182	421	379
Marijuana and Hashish	407	393	45	55	148	172	376	328
Cocaine (Including Crack)	127	104	7	10	33	40	122	96
Heroin	33	49	4	6	11	9	32	47
Hallucinogens	129	85	13	17	44	47	122	70
Inhalants	54	57	7	11	14	16	51	53
Pain Relievers ²	174	141	17	23	56	54	165	128
Tranquilizers ²	100	107	9	12	33	35	94	98
Stimulants ^{2,3}	106	91	8	17	37	34	97	85
Methamphetamine ³	73	50	4	9	18	15	69	47
Sedatives ²	50	45	3	5	8	12	51	43
ILLCIT DRUGS OTHER THAN MARIJUANA¹	269	233	23	34	79	92	258	211

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.17D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	0.32	0.29	1.28	1.51	0.58	0.64	0.35	0.30
Marijuana and Hashish	0.28	0.26	1.25	1.42	0.55	0.64	0.32	0.27
Cocaine (Including Crack)	0.09	0.07	0.24	0.35	0.16	0.18	0.11	0.08
Heroin	0.02	0.03	0.12	0.19	0.05	0.04	0.03	0.04
Hallucinogens	0.09	0.06	0.46	0.59	0.21	0.22	0.11	0.06
Inhalants	0.04	0.04	0.24	0.37	0.07	0.07	0.05	0.05
Pain Relievers ²	0.13	0.10	0.59	0.76	0.26	0.24	0.15	0.11
Tranquilizers ²	0.07	0.08	0.31	0.40	0.15	0.17	0.08	0.08
Stimulants ^{2,3}	0.08	0.06	0.28	0.59	0.17	0.16	0.09	0.07
Methamphetamine ³	0.05	0.04	0.13	0.32	0.09	0.07	0.06	0.04
Sedatives ²	0.04	0.03	0.10	0.17	0.04	0.06	0.05	0.04
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.19	0.16	0.75	1.07	0.35	0.39	0.23	0.18

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.17P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	0.0229		0.0284		0.5648		0.0245	
Marijuana and Hashish	0.0173		0.0844		0.6737		0.0133	
Cocaine (Including Crack)	0.9198		0.3335		0.1722		0.6968	
Heroin	0.0310		0.4347		0.2468		0.0144	
Hallucinogens	0.3747		0.8471		0.2404		0.7218	
Inhalants	0.3716		0.0549		0.7095		0.4617	
Pain Relievers ²	0.3195		0.4182		0.1813		0.5212	
Tranquilizers ²	0.1787		0.7829		0.9718		0.1442	
Stimulants ^{2,3}	0.3914		0.0043		0.3745		0.3767	
Methamphetamine ³	0.5480		0.0758		0.1698		0.6455	
Sedatives ²	0.2863		0.5135		0.2877		0.4321	
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.9507		0.0103		0.8019		0.9443	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.18C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	300	281	31	36	107	120	277	241
Marijuana and Hashish	284	263	31	34	103	117	261	219
Cocaine (Including Crack)	79	61	2	6	22	20	77	57
Heroin	25	26	*	5	8	2	23	25
Hallucinogens	75	29	3	7	12	13	74	25
Inhalants	9	9	*	3	3	3	8	7
Pain Relievers ²	90	66	5	8	16	16	88	65
Tranquilizers ²	31	35	5	5	11	7	28	35
Stimulants ²	17	18	3	4	9	5	14	17
Methamphetamine ³	40	31	2	4	7	10	40	29
Sedatives ²	19	14	*	2	6	7	18	12
ILLICIT DRUGS OTHER THAN MARIJUANA¹	140	108	9	15	34	34	135	103

*Low precision; no estimate reported.

NOTE: Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.18D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2013 and 2014

Drug	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLCIT DRUGS¹	0.21	0.19	1.01	1.15	0.45	0.51	0.24	0.20
Marijuana and Hashish	0.20	0.18	1.03	1.10	0.43	0.50	0.23	0.18
Cocaine (Including Crack)	0.06	0.04	0.07	0.23	0.11	0.10	0.07	0.05
Heroin	0.02	0.02	*	0.17	0.04	0.01	0.02	0.02
Hallucinogens	0.05	0.02	0.10	0.26	0.06	0.06	0.07	0.02
Inhalants	0.01	0.01	*	0.12	0.01	0.02	0.01	0.01
Pain Relievers ²	0.07	0.05	0.17	0.30	0.08	0.08	0.08	0.06
Tranquilizers ²	0.02	0.03	0.19	0.17	0.05	0.03	0.02	0.03
Stimulants ²	0.01	0.01	0.11	0.15	0.04	0.02	0.01	0.01
Methamphetamine ³	0.03	0.02	0.06	0.13	0.03	0.05	0.04	0.03
Sedatives ²	0.01	0.01	*	0.08	0.03	0.03	0.02	0.01
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.10	0.08	0.30	0.52	0.16	0.16	0.12	0.09

*Low precision; no estimate reported.

NOTE: Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.18P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
ILLICIT DRUGS¹	0.1024		0.8076		0.6020		0.0353	
Marijuana and Hashish	0.1513		0.7776		0.8030		0.0604	
Cocaine (Including Crack)	0.8091		0.1480		0.8708		0.6931	
Heroin	0.8791		*		0.0120		0.5444	
Hallucinogens	0.9516		0.1813		0.5244		0.8366	
Inhalants	0.6380		*		0.9903		0.9287	
Pain Relievers ²	0.7104		0.4525		0.5514		0.6582	
Tranquilizers ²	0.0638		0.8227		0.5303		0.0364	
Stimulants ²	0.5479		0.3582		0.2218		0.2977	
Methamphetamine ³	0.9688		0.1681		0.6773		0.8598	
Sedatives ²	0.6935		*		0.8466		0.8148	
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.4257		0.0300		0.6309		0.4472	

*Low precision; no estimate reported.

NOTE: Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.19C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	323	302	187	150	309	283	159	157
Marijuana and Hashish	296	272	171	132	268	248	112	128
Cocaine	106	88	36	20	70	52	25	14
Crack	52	48	12	9	25	25	24	*
Heroin	33	44	19	23	*	21	18	*
Hallucinogens	78	66	14	21	92	50	32	9
LSD	20	27	7	12	30	28	3	*
PCP	11	4	1	*	*	*	10	*
Ecstasy	56	45	7	15	52	42	11	6
Inhalants	41	35	21	12	35	45	27	28
Nonmedical Use of Psychotherapeutics ^{2,3}	165	135	92	77	133	128	109	96
Pain Relievers	139	102	78	68	104	95	89	83
OxyContin [®]	46	32	38	8	22	35	51	12
Tranquilizers	81	80	48	33	60	68	38	34
Stimulants ³	91	70	37	39	50	57	33	46
Methamphetamine ³	68	40	34	35	25	30	27	35
Sedatives	26	40	8	17	45	21	44	22
ILLCIT DRUGS OTHER THAN MARIJUANA¹	201	172	103	83	173	144	115	98

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.19D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	0.74	0.65	0.89	0.75	0.30	0.26	0.15	0.15
Marijuana and Hashish	0.69	0.61	0.82	0.67	0.26	0.24	0.10	0.12
Cocaine	0.29	0.24	0.19	0.10	0.07	0.05	0.02	0.01
Crack	0.14	0.13	0.06	0.05	0.02	0.02	0.02	*
Heroin	0.09	0.12	0.10	0.12	*	0.02	0.02	*
Hallucinogens	0.21	0.18	0.07	0.11	0.09	0.05	0.03	0.01
LSD	0.06	0.07	0.04	0.06	0.03	0.03	0.00	*
PCP	0.03	0.01	0.01	*	*	*	0.01	*
Ecstasy	0.15	0.12	0.04	0.08	0.05	0.04	0.01	0.01
Inhalants	0.11	0.09	0.11	0.06	0.03	0.04	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{2,3}	0.43	0.35	0.47	0.40	0.13	0.12	0.10	0.09
Pain Relievers	0.37	0.27	0.40	0.36	0.10	0.09	0.08	0.08
OxyContin [®]	0.13	0.09	0.19	0.04	0.02	0.03	0.05	0.01
Tranquilizers	0.22	0.21	0.25	0.17	0.06	0.07	0.04	0.03
Stimulants ³	0.25	0.19	0.19	0.21	0.05	0.05	0.03	0.04
Methamphetamine ³	0.19	0.11	0.18	0.19	0.02	0.03	0.03	0.03
Sedatives	0.07	0.11	0.04	0.09	0.04	0.02	0.04	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.51	0.43	0.52	0.43	0.17	0.14	0.11	0.09

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.19P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	0.6410		0.1268		0.0011		0.0897	
Marijuana and Hashish	0.3876		0.0441		0.0020		0.0145	
Cocaine	0.9834		0.5998		0.7058		0.1588	
Crack	0.8192		0.3981		0.6541		*	
Heroin	0.1083		0.0908		*		*	
Hallucinogens	0.6102		0.5593		0.4857		0.0787	
LSD	0.2899		0.2816		0.8293		*	
PCP	0.5363		*		*		*	
Ecstasy	0.5339		0.1829		0.7765		0.2897	
Inhalants	0.7998		0.5485		0.1308		0.7557	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.2009		0.4776		0.0628		0.9373	
Pain Relievers	0.0673		0.6080		0.4297		0.8261	
OxyContin [®]	0.2656		0.0433		0.1553		0.2814	
Tranquilizers	0.4649		0.6438		0.1592		0.6779	
Stimulants ³	0.5419		0.8433		0.0084		0.1508	
Methamphetamine ³	0.1295		0.9580		0.0698		0.5735	
Sedatives	0.0619		0.3090		0.7646		0.7379	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.2089		0.7376		0.0866		0.8127	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.20C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	29	37	21	19	35	47	42	45
Marijuana and Hashish	28	35	20	19	34	43	32	36
Cocaine	6	9	3	*	3	5	5	*
Crack	1	4	*	*	1	*	*	*
Heroin	3	6	3	1	1	*	1	2
Hallucinogens	11	12	7	6	7	13	8	6
LSD	8	9	4	5	3	10	3	3
PCP	3	2	1	*	1	*	*	*
Ecstasy	6	7	3	3	4	7	4	1
Inhalants	5	7	4	3	5	8	13	14
Nonmedical Use of Psychotherapeutics ^{2,3}	15	19	8	7	12	21	25	29
Pain Relievers	13	15	7	6	10	17	23	25
OxyContin [®]	4	5	0	0	4	5	3	5
Tranquilizers	8	10	1	5	3	6	8	10
Stimulants ³	6	10	5	5	6	14	6	14
Methamphetamine ³	2	7	4	*	3	6	3	6
Sedatives	3	3	1	*	1	4	6	9
ILLCIT DRUGS OTHER THAN MARIJUANA¹	18	23	11	10	13	24	29	32

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.20D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	2.20	2.75	2.63	3.03	1.40	1.72	0.19	0.20
Marijuana and Hashish	2.32	2.91	2.56	2.99	1.37	1.61	0.15	0.16
Cocaine	0.76	1.21	0.49	*	0.15	0.24	0.03	*
Crack	0.06	0.61	*	*	0.05	*	*	*
Heroin	0.44	0.78	0.47	0.22	0.05	*	0.01	0.01
Hallucinogens	1.39	1.59	1.07	1.25	0.34	0.58	0.04	0.03
LSD	1.01	1.24	0.62	0.98	0.16	0.44	0.01	0.01
PCP	0.38	0.30	0.17	*	0.04	*	*	*
Ecstasy	0.78	0.97	0.53	0.66	0.20	0.31	0.02	0.01
Inhalants	0.66	1.02	0.58	0.54	0.23	0.36	0.06	0.07
Nonmedical Use of Psychotherapeutics ^{2,3}	1.71	2.30	1.29	1.41	0.55	0.91	0.12	0.13
Pain Relievers	1.54	1.94	1.08	1.11	0.50	0.77	0.11	0.11
OxyContin [®]	0.51	0.67	0.05	0.02	0.17	0.25	0.02	0.02
Tranquilizers	1.03	1.34	0.17	0.96	0.15	0.29	0.04	0.05
Stimulants ³	0.74	1.35	0.81	1.01	0.26	0.62	0.03	0.06
Methamphetamine ³	0.30	0.96	0.59	*	0.14	0.30	0.02	0.03
Sedatives	0.34	0.48	0.24	*	0.06	0.17	0.03	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1.96	2.65	1.69	1.80	0.63	1.05	0.13	0.14

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.20P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS¹	0.2550		0.8072		0.0106		0.3007	
Marijuana and Hashish	0.4759		0.8984		0.0296		0.8202	
Cocaine	0.1914		*		0.9632		*	
Crack	0.0987		*		*		*	
Heroin	0.2683		0.3543		*		0.5888	
Hallucinogens	0.3297		0.7977		0.2488		0.4720	
LSD	0.9051		0.6047		0.1009		0.9478	
PCP	0.9007		*		*		*	
Ecstasy	0.9641		0.9286		0.8848		0.2058	
Inhalants	0.1425		0.7255		0.1599		0.7403	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.5576		0.8413		0.0047		0.2183	
Pain Relievers	0.9210		0.6057		0.1653		0.4149	
OxyContin [®]	0.8155		0.6058		0.6248		0.7223	
Tranquilizers	0.7979		0.0169		0.1039		0.8434	
Stimulants ³	0.1340		0.9103		0.0075		0.0270	
Methamphetamine ³	0.1913		*		0.2021		0.3841	
Sedatives	0.6758		*		0.5210		0.2033	
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.2721		0.4949		0.0013		0.2349	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.21C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	117	133	40	46	91	108	45	51
Marijuana and Hashish	111	123	37	42	84	104	36	46
Cocaine	29	34	6	8	16	19	2	1
Crack	11	4	5	2	*	8	*	*
Heroin	11	9	12	14	1	2	1	*
Hallucinogens	37	44	11	6	23	18	15	6
LSD	15	17	4	4	8	8	2	*
PCP	1	2	*	*	*	2	2	*
Ecstasy	27	31	6	5	15	15	4	6
Inhalants	12	13	10	0	8	9	6	5
Nonmedical Use of Psychotherapeutics ^{2,3}	57	55	18	20	36	41	22	20
Pain Relievers	49	45	15	17	28	30	20	17
OxyContin [®]	17	18	4	5	7	8	*	2
Tranquilizers	29	29	11	9	17	18	7	7
Stimulants ³	28	25	8	7	21	22	7	8
Methamphetamine ³	17	13	4	4	6	8	2	2
Sedatives	6	7	*	4	6	9	4	6
ILLCIT DRUGS OTHER THAN MARIJUANA¹	67	76	23	27	42	47	27	22

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.21D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	0.98	1.06	1.45	1.66	0.59	0.67	0.37	0.41
Marijuana and Hashish	0.96	1.05	1.37	1.57	0.55	0.67	0.31	0.38
Cocaine	0.34	0.42	0.24	0.35	0.13	0.15	0.02	0.01
Crack	0.13	0.06	0.21	0.10	*	0.06	*	*
Heroin	0.13	0.12	0.48	0.59	0.01	0.01	0.01	*
Hallucinogens	0.44	0.55	0.46	0.26	0.19	0.13	0.13	0.05
LSD	0.18	0.22	0.17	0.16	0.07	0.06	0.02	*
PCP	0.01	0.03	*	*	*	0.01	0.02	*
Ecstasy	0.32	0.39	0.25	0.22	0.12	0.11	0.04	0.05
Inhalants	0.14	0.16	0.41	0.01	0.07	0.07	0.05	0.04
Nonmedical Use of Psychotherapeutics ^{2,3}	0.61	0.65	0.73	0.85	0.27	0.30	0.18	0.17
Pain Relievers	0.55	0.56	0.60	0.73	0.22	0.22	0.17	0.14
OxyContin [®]	0.20	0.23	0.15	0.20	0.06	0.06	*	0.02
Tranquilizers	0.33	0.36	0.47	0.39	0.13	0.14	0.06	0.06
Stimulants ³	0.33	0.32	0.33	0.33	0.16	0.16	0.06	0.07
Methamphetamine ³	0.21	0.16	0.16	0.17	0.05	0.06	0.01	0.02
Sedatives	0.07	0.10	*	0.20	0.05	0.07	0.03	0.05
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.70	0.86	0.91	1.11	0.32	0.34	0.22	0.18

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.21P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS¹	0.0690		0.2865		0.5027		0.6666	
Marijuana and Hashish	0.1234		0.4229		0.5440		0.1271	
Cocaine	0.2856		0.4468		0.0662		0.3709	
Crack	0.2693		0.1409		*		*	
Heroin	0.3516		0.5569		0.9356		*	
Hallucinogens	0.9389		0.0946		0.0631		0.1872	
LSD	0.2572		0.7388		0.6040		*	
PCP	0.6199		*		*		*	
Ecstasy	0.5983		0.7329		0.4923		0.8829	
Inhalants	0.9051		0.1004		0.8153		0.8172	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.5138		0.9747		0.3681		0.0360	
Pain Relievers	0.2359		0.9452		0.8164		0.0302	
OxyContin [®]	0.8782		0.4538		0.8445		*	
Tranquilizers	0.8099		0.2466		0.8386		0.9969	
Stimulants ³	0.2631		0.6697		0.6726		0.3719	
Methamphetamine ³	0.1239		0.5767		0.5483		0.8578	
Sedatives	0.4274		*		0.4176		0.9100	
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.8946		0.7708		0.3198		0.0367	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.22C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	301	262	181	140	281	253	146	135
Marijuana and Hashish	274	229	164	124	244	216	99	106
Cocaine	102	81	36	16	68	48	25	14
Crack	51	48	11	9	25	24	24	*
Heroin	32	42	15	18	*	21	18	*
Hallucinogens	67	49	6	20	90	44	27	7
LSD	11	19	4	10	29	25	*	*
PCP	11	2	1	*	*	*	10	*
Ecstasy	49	32	*	13	51	38	10	*
Inhalants	39	31	18	11	34	43	23	23
Nonmedical Use of Psychotherapeutics ^{2,3}	155	124	90	73	128	120	104	88
Pain Relievers	129	91	77	65	101	89	84	75
OxyContin [®]	41	26	37	7	20	34	51	11
Tranquilizers	74	74	47	31	57	65	37	31
Stimulants ³	85	68	36	38	44	51	28	43
Methamphetamine ³	64	38	34	35	24	28	27	34
Sedatives	25	39	8	16	44	18	43	20
ILLCIT DRUGS OTHER THAN MARIJUANA¹	188	156	99	77	168	134	108	90

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.22D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	0.92	0.78	1.03	0.83	0.32	0.27	0.20	0.18
Marijuana and Hashish	0.86	0.70	0.95	0.75	0.28	0.24	0.13	0.14
Cocaine	0.37	0.28	0.22	0.10	0.08	0.05	0.03	0.02
Crack	0.19	0.17	0.07	0.05	0.03	0.03	0.03	*
Heroin	0.12	0.15	0.09	0.11	*	0.02	0.02	*
Hallucinogens	0.24	0.17	0.03	0.12	0.10	0.05	0.04	0.01
LSD	0.04	0.07	0.02	0.06	0.03	0.03	*	*
PCP	0.04	0.01	0.01	*	*	*	0.01	*
Ecstasy	0.18	0.11	*	0.08	0.06	0.04	0.01	*
Inhalants	0.14	0.11	0.11	0.07	0.04	0.05	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{2,3}	0.54	0.42	0.54	0.45	0.15	0.13	0.14	0.12
Pain Relievers	0.46	0.32	0.46	0.40	0.12	0.10	0.11	0.10
OxyContin [®]	0.15	0.09	0.23	0.04	0.02	0.04	0.07	0.01
Tranquilizers	0.27	0.26	0.28	0.19	0.07	0.07	0.05	0.04
Stimulants ³	0.31	0.24	0.22	0.23	0.05	0.06	0.04	0.06
Methamphetamine ³	0.23	0.13	0.20	0.21	0.03	0.03	0.04	0.05
Sedatives	0.09	0.14	0.05	0.10	0.05	0.02	0.06	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.65	0.52	0.59	0.47	0.20	0.15	0.15	0.12

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.22P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2013	2014	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS¹	0.7451		0.1291		0.0031		0.1544	
Marijuana and Hashish	0.3784		0.0398		0.0049		0.0325	
Cocaine	0.7138		0.5454		0.8649		0.3183	
Crack	0.8048		0.7393		0.9511		*	
Heroin	0.0736		0.0589		*		*	
Hallucinogens	0.8909		0.0401		0.7003		0.2806	
LSD	0.3589		0.3084		0.9513		*	
PCP	0.4869		*		*		*	
Ecstasy	0.8794		*		0.9389		*	
Inhalants	0.6729		0.8423		0.1673		0.6056	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.3347		0.4965		0.1954		0.8509	
Pain Relievers	0.1818		0.6316		0.5939		0.6254	
OxyContin [®]	0.2550		0.0313		0.1473		0.2418	
Tranquilizers	0.4026		0.6869		0.2028		0.6986	
Stimulants ³	0.7613		0.7397		0.0299		0.2712	
Methamphetamine ³	0.2029		0.9909		0.1327		0.6686	
Sedatives	0.0862		0.3906		0.5848		0.5192	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.2574		0.7734		0.2749		0.9117	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.23C Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Cigarette Use/Level of Alcohol Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	509	476	62	73	143	160	472	415
Heavy Alcohol Use ¹	232	212	17	22	82	94	217	179
Binge Alcohol Use But Not Heavy Use ¹	267	243	35	43	103	109	237	207
Alcohol Use But Not Binge Use ¹	270	236	24	32	64	74	261	223
No Alcohol Use	245	217	47	49	61	68	231	192
PAST MONTH CIGARETTE USE	372	339	36	41	126	142	348	295
Heavy Alcohol Use ¹	176	163	13	*	67	75	168	140
Binge Alcohol Use But Not Heavy Use ¹	194	174	23	27	77	77	175	149
Alcohol Use But Not Binge Use ¹	173	143	12	16	43	50	165	133
No Alcohol Use	187	150	21	19	40	46	181	140
NO PAST MONTH CIGARETTE USE	355	326	55	65	103	121	320	293
Heavy Alcohol Use ¹	140	126	*	*	44	48	130	115
Binge Alcohol Use But Not Heavy Use ¹	170	151	27	34	62	65	149	133
Alcohol Use But Not Binge Use ¹	208	192	20	28	44	55	200	181
No Alcohol Use	159	157	42	45	45	51	146	135

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.23D Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Percentages, 2013 and 2014

Cigarette Use/Level of Alcohol Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	0.19	0.18	0.25	0.30	0.41	0.46	0.23	0.20
Heavy Alcohol Use ¹	1.17	0.99	4.26	3.71	1.35	1.53	1.49	1.17
Binge Alcohol Use But Not Heavy Use ¹	0.57	0.49	2.08	2.33	0.86	0.89	0.67	0.56
Alcohol Use But Not Binge Use ¹	0.34	0.29	1.61	1.94	0.79	0.87	0.37	0.30
No Alcohol Use	0.19	0.17	0.20	0.22	0.41	0.45	0.25	0.21
PAST MONTH CIGARETTE USE	0.58	0.52	1.70	2.15	0.84	0.92	0.70	0.59
Heavy Alcohol Use ¹	1.65	1.44	4.43	*	1.65	1.93	2.19	1.75
Binge Alcohol Use But Not Heavy Use ¹	1.10	0.92	3.36	3.69	1.38	1.49	1.40	1.13
Alcohol Use But Not Binge Use ¹	1.24	1.04	4.73	5.25	1.91	2.08	1.43	1.15
No Alcohol Use	0.89	0.75	2.63	3.03	1.45	1.66	1.03	0.83
NO PAST MONTH CIGARETTE USE	0.17	0.15	0.22	0.26	0.37	0.43	0.20	0.18
Heavy Alcohol Use ¹	1.62	1.36	*	*	2.15	2.16	1.94	1.63
Binge Alcohol Use But Not Heavy Use ¹	0.59	0.50	2.41	2.93	0.96	0.98	0.65	0.56
Alcohol Use But Not Binge Use ¹	0.32	0.28	1.58	2.06	0.75	0.85	0.34	0.29
No Alcohol Use	0.15	0.15	0.19	0.20	0.37	0.41	0.20	0.18

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.23P Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Cigarette Use/Level of Alcohol Use	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	0.0022		0.1397		0.4747		0.0029	
Heavy Alcohol Use ¹	0.4600		0.0361		0.2741		0.6585	
Binge Alcohol Use But Not Heavy Use ¹	0.2968		0.2206		0.2087		0.1071	
Alcohol Use But Not Binge Use ¹	0.0069		0.0808		0.0252		0.0369	
No Alcohol Use	0.0382		0.7844		0.5385		0.0468	
PAST MONTH CIGARETTE USE	0.1923		0.3713		0.0468		0.2221	
Heavy Alcohol Use ¹	0.8785		*		0.4247		0.8990	
Binge Alcohol Use But Not Heavy Use ¹	0.5847		0.4007		0.9638		0.7794	
Alcohol Use But Not Binge Use ¹	0.1433		0.6575		0.0011		0.4118	
No Alcohol Use	0.1268		0.8072		0.2865		0.1291	
NO PAST MONTH CIGARETTE USE	0.0002		0.0148		0.3286		0.0010	
Heavy Alcohol Use ¹	0.2438		*		0.1218		0.5893	
Binge Alcohol Use But Not Heavy Use ¹	0.0305		0.2100		0.2613		0.0118	
Alcohol Use But Not Binge Use ¹	0.0125		0.0570		0.2378		0.0364	
No Alcohol Use	0.0897		0.3007		0.6666		0.1544	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.24C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	359	303
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	160	139	256	212
Cigars	272	209	244	215
Pipe Tobacco	120	93	93	101
ALCOHOL	675	592	1,519	1,240
Binge Alcohol Use ²	517	460	762	596
Heavy Alcohol Use ²	308	258	313	270

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.24D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	0.17	0.14
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.28	0.24	0.12	0.10
Cigars	0.46	0.35	0.12	0.10
Pipe Tobacco	0.21	0.16	0.04	0.05
ALCOHOL	0.74	0.63	0.48	0.37
Binge Alcohol Use ²	0.71	0.62	0.32	0.26
Heavy Alcohol Use ²	0.52	0.43	0.15	0.13

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.24P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A		0.4030	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.2970		0.9754	
Cigars	0.0140		0.3722	
Pipe Tobacco	0.3112		0.3681	
ALCOHOL	0.5225		0.3341	
Binge Alcohol Use ²	0.6740		0.7127	
Heavy Alcohol Use ²	0.5679		0.9522	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.25C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	34	34
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	19	22	24	24
Cigars	23	25	23	22
Pipe Tobacco	12	16	13	12
ALCOHOL	39	43	70	73
Binge Alcohol Use ²	33	36	48	49
Heavy Alcohol Use ²	16	18	18	16

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.25D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	0.14	0.14
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1.24	1.68	0.10	0.10
Cigars	1.50	1.82	0.10	0.09
Pipe Tobacco	0.81	1.22	0.05	0.05
ALCOHOL	1.81	2.12	0.28	0.29
Binge Alcohol Use ²	1.72	2.14	0.19	0.20
Heavy Alcohol Use ²	1.08	1.41	0.07	0.07

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.25P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A		0.5427	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.1266		0.7561	
Cigars	0.7531		0.4318	
Pipe Tobacco	0.1013		0.9193	
ALCOHOL	0.4309		0.5483	
Binge Alcohol Use ²	0.8441		0.7569	
Heavy Alcohol Use ²	0.5236		0.9921	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.26C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	86	93
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	58	65	45	55
Cigars	81	83	65	74
Pipe Tobacco	35	34	37	30
ALCOHOL	173	196	280	274
Binge Alcohol Use ²	155	170	195	191
Heavy Alcohol Use ²	90	96	77	85

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.26D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	0.33	0.34
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.50	0.60	0.19	0.22
Cigars	0.63	0.72	0.26	0.28
Pipe Tobacco	0.32	0.34	0.15	0.12
ALCOHOL	0.64	0.78	0.65	0.67
Binge Alcohol Use ²	0.81	0.91	0.59	0.59
Heavy Alcohol Use ²	0.67	0.79	0.29	0.32

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.26P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A		0.7986	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.6794		0.6461	
Cigars	0.5138		0.3856	
Pipe Tobacco	0.7182		0.0501	
ALCOHOL	0.7638		0.3403	
Binge Alcohol Use ²	0.9233		0.3198	
Heavy Alcohol Use ²	0.4923		0.8128	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.27C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	345	294
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	148	117	251	207
Cigars	258	187	233	199
Pipe Tobacco	114	86	86	95
ALCOHOL	640	541	1,461	1,204
Binge Alcohol Use ²	488	401	718	555
Heavy Alcohol Use ²	298	235	295	252

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.27D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2013 and 2014

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A	N/A	0.21	0.17
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.33	0.26	0.15	0.13
Cigars	0.55	0.40	0.14	0.12
Pipe Tobacco	0.25	0.19	0.05	0.06
ALCOHOL	0.93	0.76	0.58	0.45
Binge Alcohol Use ²	0.87	0.71	0.40	0.31
Heavy Alcohol Use ²	0.64	0.50	0.18	0.15

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.27P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2013	2014	2013	2014
TOBACCO PRODUCTS¹	N/A		0.4242	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.3966		0.9543	
Cigars	0.0323		0.4871	
Pipe Tobacco	0.2146		0.1064	
ALCOHOL	0.4079		0.4669	
Binge Alcohol Use ²	0.4573		0.9927	
Heavy Alcohol Use ²	0.8270		0.8556	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.28C Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	337	283	445	397	453	386	590	501
Cigarettes	308	258	406	363	398	348	560	466
Smokeless Tobacco	122	108	154	121	133	114	175	160
Cigars	154	123	219	165	182	164	152	133
Pipe Tobacco	56	39	83	62	74	60	85	102

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.28D Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	1.32	1.07	0.80	0.66	0.55	0.45	0.41	0.35
Cigarettes	1.27	1.12	0.77	0.64	0.49	0.42	0.40	0.34
Smokeless Tobacco	0.73	0.62	0.35	0.26	0.17	0.14	0.14	0.13
Cigars	0.87	0.68	0.48	0.35	0.24	0.20	0.12	0.10
Pipe Tobacco	0.33	0.23	0.19	0.14	0.10	0.08	0.07	0.08

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.28P Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	0.8484		0.7099		0.5064		0.8839	
Cigarettes	0.4907		0.8435		0.5463		0.4249	
Smokeless Tobacco	0.9462		0.1559		0.6376		0.5598	
Cigars	0.0931		0.0511		0.6556		0.7050	
Pipe Tobacco	0.1670		0.3438		0.6149		0.1542	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.29C Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	18	21	35	35	22	25	41	40
Cigarettes	16	18	28	31	18	21	34	31
Smokeless Tobacco	12	13	15	16	10	12	21	23
Cigars	12	14	22	20	10	11	19	20
Pipe Tobacco	5	6	11	11	5	8	11	13

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.29D Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	4.13	4.62	1.98	2.09	1.49	1.64	0.18	0.18
Cigarettes	4.14	4.75	1.86	1.98	1.29	1.46	0.15	0.14
Smokeless Tobacco	3.51	4.12	1.16	1.20	0.73	0.87	0.09	0.10
Cigars	3.61	4.63	1.53	1.50	0.78	0.82	0.09	0.09
Pipe Tobacco	1.68	2.13	0.88	0.84	0.38	0.57	0.05	0.06

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.29P Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	0.5533		0.2226		0.3945		0.0519	
Cigarettes	0.3324		0.3448		0.6734		0.0197	
Smokeless Tobacco	0.8183		0.3711		0.7992		0.4573	
Cigars	0.5773		0.8641		0.5442		0.4192	
Pipe Tobacco	0.8875		0.9187		0.2603		0.5407	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.30C Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	100	104	124	133	83	90	92	97
Cigarettes	90	96	110	124	70	78	83	87
Smokeless Tobacco	40	41	44	48	26	28	30	41
Cigars	57	51	60	71	41	48	38	44
Pipe Tobacco	22	19	31	31	23	17	20	20

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.30D Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	1.33	1.48	0.94	0.94	0.89	1.02	0.57	0.62
Cigarettes	1.39	1.53	0.93	0.96	0.81	0.93	0.52	0.57
Smokeless Tobacco	0.94	1.01	0.45	0.49	0.34	0.36	0.21	0.28
Cigars	1.19	1.19	0.59	0.69	0.53	0.61	0.27	0.30
Pipe Tobacco	0.53	0.50	0.32	0.32	0.29	0.22	0.14	0.14

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150410

Table 6.30P Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	0.1446		0.0184		0.0818		0.5901	
Cigarettes	0.0649		0.0215		0.0715		0.1679	
Smokeless Tobacco	0.1153		0.5421		0.6523		0.0617	
Cigars	0.1629		0.7226		0.6021		0.8422	
Pipe Tobacco	0.4424		0.5896		0.2820		0.4963	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.31C Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	322	256	419	355	438	376	571	479
Cigarettes	298	235	384	322	388	342	543	446
Smokeless Tobacco	114	99	146	106	130	109	173	154
Cigars	139	112	207	145	175	156	149	125
Pipe Tobacco	51	35	78	51	69	58	81	98

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.31D Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2013 and 2014

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	1.71	1.34	1.01	0.80	0.61	0.49	0.56	0.47
Cigarettes	1.65	1.41	0.97	0.77	0.54	0.46	0.54	0.45
Smokeless Tobacco	0.90	0.76	0.43	0.30	0.19	0.15	0.19	0.17
Cigars	1.06	0.83	0.59	0.40	0.26	0.22	0.16	0.14
Pipe Tobacco	0.41	0.28	0.24	0.15	0.10	0.08	0.09	0.11

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.31P Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2013	2014	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS²	0.7581		0.6846		0.7798		0.9113	
Cigarettes	0.9470		0.5628		0.8181		0.6765	
Smokeless Tobacco	0.5753		0.2334		0.6892		0.9279	
Cigars	0.2603		0.0531		0.6971		0.6612	
Pipe Tobacco	0.2774		0.4351		0.7681		0.1203	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.32C Marijuana Use in the Prior Year and Past Year Marijuana Use Status among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Marijuana Use in Prior Year		Marijuana Use in Prior Year and Past Year		Marijuana Use in Prior Year But No Use in Past Year		Percentage Who Did Not Use Marijuana in the Past Year among Prior Year Marijuana Users	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	597	546	517	479	266	247	0.72	0.63
AGE								
12-17	62	67	56	61	28	28	1.22	1.30
12-13	13	12	12	12	6	3	*	*
14-15	32	32	29	30	15	13	2.47	2.68
16-17	51	62	47	55	23	25	1.45	1.52
18-25	161	182	148	168	70	87	0.63	0.75
18-20	117	125	105	115	43	45	1.03	1.15
21-25	170	186	147	163	61	76	0.80	0.95
26 or Older	549	471	476	410	256	233	1.13	0.94
26-34	258	219	243	189	118	102	1.41	1.12
35-49	233	207	209	186	122	106	1.81	1.34
50 or Older	385	347	331	296	194	170	2.91	2.13
GENDER								
Male	437	411	381	372	203	179	0.92	0.77
Female	359	311	320	252	176	161	1.24	0.98
GENDER/AGE								
Male 12-17	45	49	43	45	19	20	1.60	1.71
Male 18-25	114	126	110	117	54	60	0.84	0.93
Male 26 or Older	408	371	357	331	193	169	1.40	1.09
Female 12-17	43	42	37	39	20	19	1.82	1.96
Female 18-25	97	120	89	112	48	57	1.04	1.14
Female 26 or Older	339	260	302	209	168	151	2.00	1.56

*Low precision; no estimate reported.

NOTE: Prior Year Marijuana Use is defined as using marijuana/hashish 12 to 23 months prior to the interview date.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.32D Marijuana Use in the Prior Year and Past Year Marijuana Use Status among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Marijuana Use in Prior Year		Marijuana Use in Prior Year and Past Year		Marijuana Use in Prior Year But No Use in Past Year		Percentage Who Did Not Use Marijuana in the Past Year among Prior Year Marijuana Users	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.23	0.21	0.20	0.18	0.10	0.09	0.72	0.63
AGE								
12-17	0.25	0.27	0.23	0.25	0.11	0.11	1.22	1.30
12-13	0.17	0.15	0.15	0.15	0.08	0.03	*	*
14-15	0.36	0.37	0.34	0.34	0.17	0.16	2.47	2.68
16-17	0.55	0.64	0.53	0.58	0.27	0.29	1.45	1.52
18-25	0.46	0.52	0.43	0.48	0.20	0.25	0.63	0.75
18-20	0.70	0.80	0.64	0.74	0.32	0.34	1.03	1.15
21-25	0.57	0.64	0.53	0.60	0.26	0.33	0.80	0.95
26 or Older	0.27	0.23	0.23	0.20	0.13	0.11	1.13	0.94
26-34	0.69	0.58	0.65	0.50	0.31	0.27	1.41	1.12
35-49	0.38	0.34	0.35	0.31	0.20	0.18	1.81	1.34
50 or Older	0.37	0.32	0.32	0.28	0.18	0.16	2.91	2.13
GENDER								
Male	0.34	0.32	0.30	0.29	0.16	0.14	0.92	0.77
Female	0.27	0.23	0.24	0.18	0.13	0.12	1.24	0.98
GENDER/AGE								
Male 12-17	0.36	0.39	0.34	0.36	0.15	0.16	1.60	1.71
Male 18-25	0.65	0.72	0.63	0.67	0.31	0.34	0.84	0.93
Male 26 or Older	0.42	0.38	0.37	0.34	0.20	0.17	1.40	1.09
Female 12-17	0.35	0.34	0.31	0.32	0.16	0.16	1.82	1.96
Female 18-25	0.56	0.69	0.51	0.64	0.28	0.33	1.04	1.14
Female 26 or Older	0.32	0.24	0.29	0.20	0.16	0.14	2.00	1.56

*Low precision; no estimate reported.

NOTE: Prior Year Marijuana Use is defined as using marijuana/hashish 12 to 23 months prior to the interview date.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.32P Marijuana Use in the Prior Year and Past Year Marijuana Use Status among Persons Aged 12 or Older, by Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Marijuana Use in Prior Year		Marijuana Use in Prior Year and Past Year		Marijuana Use in Prior Year But No Use in Past Year		Percentage Who Did Not Use Marijuana in the Past Year among Prior Year Marijuana Users	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.0124		0.0371		0.1263		0.7613	
AGE								
12-17	0.3360		0.3494		0.8054		0.8784	
12-13	0.6417		0.7365		0.0328		*	
14-15	0.0401		0.1418		0.1302		0.4690	
16-17	0.9380		0.5774		0.3420		0.2899	
18-25	0.2128		0.4438		0.2435		0.5131	
18-20	0.3304		0.6482		0.2105		0.3459	
21-25	0.0198		0.1851		0.0253		0.2127	
26 or Older	0.0108		0.0252		0.2031		0.9869	
26-34	0.8002		0.4107		0.2826		0.1873	
35-49	0.0009		0.0018		0.3144		0.5008	
50 or Older	0.0701		0.0656		0.6376		0.5930	
GENDER								
Male	0.0133		0.0146		0.4912		0.6343	
Female	0.2571		0.6789		0.1209		0.2679	
GENDER/AGE								
Male 12-17	0.5587		0.4519		0.7780		0.5897	
Male 18-25	0.5206		0.5163		0.9519		0.8449	
Male 26 or Older	0.0028		0.0026		0.4717		0.4506	
Female 12-17	0.4162		0.5658		0.5169		0.7356	
Female 18-25	0.0077		0.0530		0.0698		0.6286	
Female 26 or Older	0.6485		0.8684		0.2570		0.3009	

*Low precision; no estimate reported.

NOTE: Prior Year Marijuana Use is defined as using marijuana/hashish 12 to 23 months prior to the interview date.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.33C Cigarette Use and Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Detailed Age Category:
Standard Errors of Numbers in Thousands and Standard Errors of Percentages, 2013 and 2014**

Age Category	Cigarette Use in Past Month		Nicotine Dependence in Past Month		Percentage with Nicotine Dependence among Past Month Cigarette Smokers	
	2013	2014	2013	2014	2013	2014
TOTAL	798	700	647	532	0.74	0.60
12	7	4	3	2	*	*
13	10	11	3	6	*	*
14	14	14	9	5	*	*
15	21	20	12	11	4.04	4.81
16	26	28	14	17	3.08	3.67
17	33	32	21	21	2.74	3.36
18	54	60	31	32	2.61	3.04
19	62	57	38	34	2.39	2.79
20	69	68	41	43	2.36	2.75
21	60	66	33	40	2.08	2.49
22	67	74	44	50	2.21	2.60
23	66	65	41	43	2.16	2.46
24	67	64	41	40	2.04	2.31
25	58	73	38	49	2.14	2.50
26-29	260	175	163	116	2.21	1.63
30-34	263	192	180	129	1.87	1.60
35-39	222	190	167	134	2.41	1.82
40-44	230	183	191	138	2.21	1.80
45-49	232	190	176	155	2.38	1.84
50-54	338	300	288	242	2.89	2.26
55-59	279	248	241	213	3.22	2.53
60-64	224	218	173	172	3.82	3.19
65 or Older	303	236	246	196	3.54	2.74

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.33D Cigarette Use and Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Detailed Age Category:
Standard Errors of Percentages, 2013 and 2014**

Age Category	Cigarette Use in Past Month		Nicotine Dependence in Past Month		Percentage with Nicotine Dependence among Past Month Cigarette Smokers	
	2013	2014	2013	2014	2013	2014
TOTAL	0.30	0.26	0.25	0.20	0.74	0.60
12	0.17	0.12	0.09	0.06	*	*
13	0.23	0.26	0.08	0.14	*	*
14	0.32	0.33	0.20	0.11	*	*
15	0.51	0.47	0.30	0.27	4.04	4.81
16	0.60	0.61	0.33	0.38	3.08	3.67
17	0.74	0.71	0.49	0.48	2.74	3.36
18	1.05	1.14	0.67	0.68	2.61	3.04
19	1.25	1.21	0.85	0.78	2.39	2.79
20	1.24	1.31	0.90	0.93	2.36	2.75
21	1.15	1.28	0.76	0.87	2.08	2.49
22	1.25	1.33	0.90	1.02	2.21	2.60
23	1.19	1.22	0.89	0.90	2.16	2.46
24	1.27	1.23	0.87	0.86	2.04	2.31
25	1.22	1.31	0.92	1.01	2.14	2.50
26-29	1.17	0.85	0.88	0.64	2.21	1.63
30-34	1.07	0.77	0.82	0.59	1.87	1.60
35-39	1.04	0.85	0.84	0.65	2.41	1.82
40-44	0.98	0.78	0.83	0.64	2.21	1.80
45-49	1.00	0.82	0.82	0.69	2.38	1.84
50-54	1.30	1.08	1.14	0.92	2.89	2.26
55-59	1.24	1.07	1.11	0.95	3.22	2.53
60-64	1.22	1.15	0.96	0.92	3.82	3.19
65 or Older	0.70	0.52	0.57	0.44	3.54	2.74

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.33P Cigarette Use and Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Detailed Age Category:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Age Category	Cigarette Use in Past Month		Nicotine Dependence in Past Month		Percentage with Nicotine Dependence among Past Month Cigarette Smokers	
	2013	2014	2013	2014	2013	2014
TOTAL	0.2954		0.6818		0.5983	
12	0.6144		0.4693		*	
13	0.5854		0.7930		*	
14	0.2258		0.0331		*	
15	0.0486		0.3020		0.9841	
16	0.9254		0.8928		0.8177	
17	0.0189		0.3372		0.6104	
18	0.3032		0.3853		0.7570	
19	0.0093		0.0272		0.3944	
20	0.0550		0.1950		0.7794	
21	0.2517		0.8403		0.5771	
22	0.6462		0.5215		0.6496	
23	0.0663		0.8333		0.1131	
24	0.0776		0.1606		0.7238	
25	0.9440		0.6528		0.6042	
26-29	0.0538		0.5592		0.4224	
30-34	0.0013		0.0274		0.9934	
35-39	0.3668		0.9602		0.4580	
40-44	0.7451		0.5166		0.1651	
45-49	0.4476		0.8155		0.2268	
50-54	0.4800		0.9921		0.3332	
55-59	0.0452		0.1784		0.7559	
60-64	0.1789		0.2498		0.9041	
65 or Older	0.6160		0.8757		0.3689	

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.34C Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	647	532	31	31	111	121	632	510
GENDER								
Male	432	359	24	22	80	89	426	350
Female	420	350	19	20	66	75	412	337
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	612	510	28	28	103	112	597	491
White	548	455	28	27	96	102	566	465
Black or African American	219	192	8	7	44	56	233	211
American Indian or Alaska Native	44	38	2	1	9	6	42	38
Native Hawaiian or Other Pacific Islander	*	33	1	*	7	*	*	32
Asian	53	76	3	1	18	20	51	75
Two or More Races	95	60	4	4	17	13	93	58
Hispanic or Latino	201	155	13	11	38	40	197	148
FAMILY INCOME								
Less Than \$20,000	368	331	15	15	73	75	353	308
\$20,000 - \$49,999	410	329	19	18	68	71	401	324
\$50,000 - \$74,999	275	188	11	12	38	45	273	184
\$75,000 or More	267	226	14	12	42	44	260	219

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.34D Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	0.25	0.20	0.13	0.12	0.32	0.35	0.31	0.25
GENDER								
Male	0.34	0.28	0.19	0.18	0.46	0.51	0.44	0.36
Female	0.31	0.26	0.16	0.17	0.38	0.43	0.39	0.31
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.28	0.23	0.15	0.15	0.37	0.41	0.34	0.28
White	0.32	0.27	0.20	0.20	0.44	0.47	0.39	0.32
Black or African American	0.70	0.61	0.22	0.21	0.78	1.00	0.96	0.79
American Indian or Alaska Native	2.91	2.57	1.45	0.75	3.66	2.95	3.86	3.38
Native Hawaiian or Other Pacific Islander	*	3.10	1.46	*	3.17	*	*	4.13
Asian	0.39	0.54	0.22	0.10	0.95	0.96	0.49	0.70
Two or More Races	1.99	1.27	0.57	0.50	2.08	1.56	2.98	1.92
Hispanic or Latino	0.49	0.36	0.23	0.20	0.52	0.54	0.69	0.50
FAMILY INCOME								
Less Than \$20,000	0.67	0.59	0.34	0.34	0.63	0.66	0.96	0.79
\$20,000 - \$49,999	0.45	0.36	0.25	0.25	0.57	0.57	0.57	0.47
\$50,000 - \$74,999	0.58	0.40	0.25	0.32	0.74	0.89	0.72	0.49
\$75,000 or More	0.29	0.24	0.16	0.13	0.55	0.53	0.35	0.29

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.34P Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	0.6818		0.1812		0.0366		0.9728	
GENDER								
Male	0.5283		0.1259		0.0687		0.8507	
Female	0.9841		0.7576		0.2088		0.8055	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.8443		0.1744		0.0304		0.7957	
White	0.6321		0.2753		0.0438		0.9266	
Black or African American	0.5054		0.6935		0.7612		0.4515	
American Indian or Alaska Native	0.2454		0.5342		0.2692		0.3298	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.0252		0.6485		0.9001		0.0210	
Two or More Races	0.2396		0.6584		0.0759		0.4071	
Hispanic or Latino	0.5994		0.8416		0.8874		0.6043	
FAMILY INCOME								
Less Than \$20,000	0.4706		0.8885		0.0396		0.2728	
\$20,000 - \$49,999	0.8322		0.2948		0.5156		0.9161	
\$50,000 - \$74,999	0.1082		0.5799		0.5929		0.1021	
\$75,000 or More	1.0000		0.1363		0.0216		0.6079	

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.35C Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Geographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	647	532	31	31	111	121	632	510
GEOGRAPHIC DIVISION								
Northeast	227	236	9	9	43	47	221	223
New England	103	101	5	3	21	26	102	99
Middle Atlantic	202	212	8	9	38	39	196	198
Midwest	277	259	13	17	45	55	268	248
East North Central	215	220	11	15	36	43	206	211
West North Central	164	143	7	8	28	35	162	134
South	452	339	21	20	74	82	440	324
South Atlantic	357	257	16	13	58	49	347	251
East South Central	165	153	7	11	27	37	162	139
West South Central	202	167	10	9	36	51	197	161
West	297	212	18	12	54	54	291	202
Mountain	153	116	8	7	28	28	149	109
Pacific	255	180	16	9	46	45	250	173
COUNTY TYPE								
Large Metro	458	421	23	19	81	99	444	400
Small Metro	417	321	17	17	69	68	405	297
250K – 1 Mil. Pop.	287	256	14	14	51	53	278	238
< 250K Pop.	304	202	8	10	47	42	292	185
Nonmetro	319	306	14	15	59	63	299	276
Urbanized	210	191	10	10	40	47	197	166
Less Urbanized	228	229	9	9	45	38	210	210
Completely Rural	108	96	3	6	14	16	101	87

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.35D Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Geographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Geographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	0.25	0.20	0.13	0.12	0.32	0.35	0.31	0.25
GEOGRAPHIC DIVISION								
Northeast	0.48	0.50	0.22	0.22	0.70	0.77	0.60	0.60
New England	0.83	0.80	0.43	0.31	1.30	1.58	1.05	1.01
Middle Atlantic	0.58	0.60	0.26	0.28	0.83	0.86	0.72	0.72
Midwest	0.49	0.46	0.23	0.31	0.61	0.74	0.62	0.57
East North Central	0.55	0.56	0.28	0.39	0.71	0.85	0.68	0.70
West North Central	0.95	0.82	0.42	0.51	1.22	1.53	1.22	1.00
South	0.46	0.34	0.22	0.22	0.58	0.64	0.58	0.42
South Atlantic	0.69	0.49	0.35	0.27	0.89	0.75	0.86	0.61
East South Central	1.07	0.98	0.51	0.75	1.33	1.83	1.35	1.15
West South Central	0.66	0.54	0.31	0.28	0.84	1.19	0.85	0.68
West	0.49	0.34	0.30	0.20	0.65	0.64	0.62	0.42
Mountain	0.82	0.61	0.44	0.38	1.12	1.11	1.04	0.75
Pacific	0.60	0.42	0.38	0.23	0.78	0.76	0.77	0.52
COUNTY TYPE								
Large Metro	0.31	0.27	0.16	0.13	0.41	0.48	0.39	0.33
Small Metro	0.48	0.36	0.22	0.24	0.57	0.58	0.61	0.44
250K – 1 Mil. Pop.	0.52	0.43	0.27	0.28	0.67	0.69	0.65	0.52
< 250K Pop.	0.92	0.68	0.37	0.45	1.04	1.04	1.18	0.82
Nonmetro	0.62	0.56	0.35	0.38	0.91	1.05	0.77	0.66
Urbanized	0.92	0.82	0.60	0.66	1.20	1.62	1.14	0.94
Less Urbanized	0.91	0.83	0.51	0.48	1.52	1.35	1.11	1.02
Completely Rural	1.84	1.78	0.76	1.54	2.29	3.18	2.24	2.06

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.35P Nicotine (Cigarette) Dependence in the Past Month among Persons Aged 12 or Older, by Age Group and Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Geographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	0.6818		0.1812		0.0366		0.9728	
GEOGRAPHIC DIVISION								
Northeast	0.4005		0.7125		0.4297		0.3006	
New England	0.9027		0.6977		0.8636		0.8751	
Middle Atlantic	0.3167		0.8263		0.2958		0.2133	
Midwest	0.6603		0.4364		0.3955		0.7233	
East North Central	0.8815		0.4967		0.7299		0.8717	
West North Central	0.3296		0.7032		0.3649		0.4014	
South	0.8801		0.3497		0.5126		0.7482	
South Atlantic	0.7509		0.2496		0.2753		0.5738	
East South Central	0.7440		0.5013		0.3928		0.6310	
West South Central	0.6405		0.3755		0.4175		0.5201	
West	0.1508		0.0440		0.0363		0.3519	
Mountain	0.4873		0.4403		0.2769		0.3183	
Pacific	0.0351		0.0580		0.0658		0.0916	
COUNTY TYPE								
Large Metro	0.8819		0.1606		0.6622		0.7607	
Small Metro	1.0000		0.8993		0.3546		0.9109	
250K – 1 Mil. Pop.	0.5779		0.9967		0.4994		0.5097	
< 250K Pop.	0.4964		0.8157		0.5317		0.5234	
Nonmetro	0.4892		0.4167		0.0061		0.9332	
Urbanized	0.6214		0.5114		0.1068		0.9368	
Less Urbanized	0.4601		0.2269		0.0016		0.9640	
Completely Rural	0.8561		0.2072		0.1165		0.8539	

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.36C Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	First Used at Age 14 or Younger (2013)	First Used at Age 14 or Younger (2014)	First Used at Age 15-17 (2013)	First Used at Age 15-17 (2014)	First Used at Age 18 or Older (2013)	First Used at Age 18 or Older (2014)
TOTAL	681	567	423	352	385	311	294	248
AGE								
18-25	120	131	76	76	71	80	48	50
26-34	246	174	160	125	141	96	107	74
35-49	327	253	214	166	177	141	141	100
50 or Older	494	412	316	253	292	240	221	205
GENDER								
Male	455	387	312	248	271	229	205	158
Female	443	373	276	219	269	216	212	182
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	637	546	405	338	365	297	263	240
White	583	487	379	310	321	261	225	198
Black or African American	239	221	130	119	138	125	137	117
Other or Two or More Races	124	108	74	56	*	63	47	69
Hispanic or Latino	211	163	133	102	104	97	131	64
EDUCATION								
< High School	308	254	233	169	150	147	145	88
High School Graduate	412	343	230	202	260	209	177	156
Some College	353	273	226	176	207	162	157	136
College Graduate	206	174	126	104	133	102	88	97

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.36D Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	First Used at Age 14 or Younger (2013)	First Used at Age 14 or Younger (2014)	First Used at Age 15-17 (2013)	First Used at Age 15-17 (2014)	First Used at Age 18 or Older (2013)	First Used at Age 18 or Older (2014)
TOTAL	0.75	0.61	1.08	0.95	1.24	0.99	1.46	1.22
AGE								
18-25	0.84	0.96	1.38	1.55	1.26	1.52	1.46	1.69
26-34	1.45	1.17	2.27	1.81	2.39	1.91	2.91	2.15
35-49	1.38	1.10	1.85	1.63	2.27	1.73	2.68	1.97
50 or Older	1.70	1.29	2.45	2.16	2.95	2.07	2.98	2.78
GENDER								
Male	1.00	0.82	1.44	1.28	1.65	1.37	1.99	1.62
Female	1.09	0.89	1.63	1.34	1.85	1.43	2.11	1.78
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.80	0.64	1.12	0.97	1.33	1.03	1.54	1.33
White	0.86	0.73	1.21	1.05	1.40	1.16	1.78	1.59
Black or African American	2.33	1.75	4.26	3.32	4.03	2.96	3.43	2.80
Other or Two or More Races	3.04	2.39	4.60	3.87	*	4.39	4.44	4.53
Hispanic or Latino	2.25	1.77	3.95	3.20	3.23	3.20	4.57	2.26
EDUCATION								
< High School	1.65	1.40	2.08	1.80	2.91	2.17	4.11	3.59
High School Graduate	1.26	0.96	1.75	1.46	2.15	1.54	2.40	2.00
Some College	1.40	1.10	2.12	1.71	2.35	1.93	2.40	2.11
College Graduate	1.92	1.63	3.36	3.09	3.43	2.60	2.86	2.82

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.36P Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total (2013)	Total (2014)	First Used at Age 14 or Younger (2013)	First Used at Age 14 or Younger (2014)	First Used at Age 15-17 (2013)	First Used at Age 15-17 (2014)	First Used at Age 18 or Older (2013)	First Used at Age 18 or Older (2014)
TOTAL	0.6535		0.1923		0.1068		0.8705	
AGE								
18-25	0.9515		0.3014		0.3324		0.8841	
26-34	0.5934		0.0984		0.9120		0.5754	
35-49	0.5325		0.0636		0.6567		0.9195	
50 or Older	0.8946		0.0299		0.3371		0.7714	
GENDER								
Male	0.8855		0.0388		0.2733		0.8175	
Female	0.4066		0.7254		0.2251		0.9511	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.4358		0.2358		0.2155		0.2840	
White	0.8600		0.2325		0.4035		0.5970	
Black or African American	0.1791		0.7935		0.1493		0.4813	
Other or Two or More Races	0.6802		0.9832		*		0.1146	
Hispanic or Latino	0.5986		0.7535		0.3760		0.1719	
EDUCATION								
< High School	0.8120		0.9908		0.1568		0.1667	
High School Graduate	0.8821		0.5789		0.9138		0.8214	
Some College	0.7269		0.4826		0.1702		0.6046	
College Graduate	0.4049		0.2165		0.6325		0.0292	

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.37C Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes Daily and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total		AGE FIRST USED CIGARETTES DAILY ¹						Never Used Cigarettes Daily ¹	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	681	567	250	231	381	306	460	357	91	100
AGE										
18-25	120	131	42	44	71	79	65	77	36	31
26-34	246	174	91	71	143	107	165	111	31	30
35-49	327	253	136	101	177	149	209	150	68	34
50 or Older	494	412	184	181	290	229	348	287	*	86
GENDER										
Male	455	387	198	163	277	238	314	235	72	82
Female	443	373	155	151	251	202	326	248	55	58
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	637	546	238	218	375	294	427	341	70	89
White	583	487	217	195	349	280	387	288	52	54
Black or African American	239	221	*	*	*	103	172	148	47	64
Other or Two or More Races	124	108	*	*	*	60	71	82	9	28
Hispanic or Latino	211	163	*	63	78	84	168	106	57	47
EDUCATION										
< High School	308	254	151	112	180	162	192	125	40	63
High School Graduate	412	343	139	138	248	193	266	228	70	51
Some College	353	273	132	108	198	154	260	195	35	51
College Graduate	206	174	*	*	109	94	162	126	25	31

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

¹ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.37D Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes Daily and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		AGE FIRST USED CIGARETTES DAILY ²						Never Used Cigarettes Daily ²	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.75	0.61	1.44	1.52	1.34	0.98	1.16	0.97	1.05	1.05
AGE										
18-25	0.84	0.96	2.14	2.44	1.49	1.73	1.45	1.66	0.95	0.94
26-34	1.45	1.17	3.53	2.42	2.59	1.92	2.16	1.69	1.69	1.34
35-49	1.38	1.10	2.13	2.19	2.07	1.58	1.97	1.56	3.46	1.77
50 or Older	1.70	1.29	2.98	3.20	3.02	2.04	2.41	2.02	*	4.76
GENDER										
Male	1.00	0.82	1.89	2.19	1.71	1.32	1.62	1.33	1.38	1.47
Female	1.09	0.89	2.20	2.20	1.99	1.43	1.73	1.41	1.61	1.53
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	0.80	0.64	1.48	1.61	1.38	1.00	1.22	1.01	1.11	1.30
White	0.86	0.73	1.60	1.67	1.37	1.07	1.41	1.14	1.24	1.29
Black or African American	2.33	1.75	*	*	*	3.75	3.13	2.41	2.87	3.34
Other or Two or More Races	3.04	2.39	*	*	*	4.82	4.51	3.98	1.93	4.07
Hispanic or Latino	2.25	1.77	*	4.74	4.40	4.35	3.93	2.96	2.42	1.84
EDUCATION										
< High School	1.65	1.40	2.11	2.70	2.68	1.88	3.08	2.76	2.74	3.52
High School Graduate	1.26	0.96	2.47	2.16	2.10	1.49	1.91	1.54	2.43	1.85
Some College	1.40	1.10	3.07	3.03	2.40	1.89	2.05	1.67	1.33	1.80
College Graduate	1.92	1.63	*	*	4.64	3.56	2.69	2.21	1.49	1.56

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

¹ Standard errors in the Total column correspond to percentages among all persons in the specified subgroup who used cigarettes in the past month, regardless of whether they ever used cigarettes daily.

² Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.37P Nicotine (Cigarette) Dependence in the Past Month among Past Month Cigarette Smokers Aged 18 or Older, by Age First Used Cigarettes Daily and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total ¹		AGE FIRST USED CIGARETTES DAILY ²						Never Used Cigarettes Daily ²	
			14 or Younger		15-17		18 or Older		2013	2014
	2013	2014	2013	2014	2013	2014	2013	2014		
TOTAL	0.6535		0.0291		0.0774		0.6593		0.7001	
AGE										
18-25	0.9515		0.8311		0.2070		0.3539		0.3153	
26-34	0.5934		0.5694		0.1610		0.1058		0.9792	
35-49	0.5325		0.0650		0.6909		0.8665		0.2148	
50 or Older	0.8946		0.0165		0.1579		0.4996		*	
GENDER										
Male	0.8855		0.0781		0.3722		0.7319		0.7610	
Female	0.4066		0.2063		0.1120		0.3340		0.8212	
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	0.4358		0.0509		0.0864		0.5946		0.3317	
White	0.8600		0.0686		0.2775		0.9562		0.5421	
Black or African American	0.1791		*		*		0.4779		0.2389	
Other or Two or More Races	0.6802		*		*		0.2830		0.0302	
Hispanic or Latino	0.5986		*		0.6244		0.8346		0.4646	
EDUCATION										
< High School	0.8120		0.0441		0.0908		0.7984		0.4944	
High School Graduate	0.8821		0.9550		0.7633		0.5440		0.4120	
Some College	0.7269		0.0398		0.5179		0.8470		0.1276	
College Graduate	0.4049		*		0.1975		0.1580		0.9607	

*Low precision; no estimate reported.

NOTE: Nicotine (Cigarette) Dependence is based on criteria derived from the Nicotine Dependence Syndrome Scale (NDSS) or the Fagerstrom Test of Nicotine Dependence (FTND). See Section B.4.2 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: National Findings*.

¹ *P* Values in the Total column correspond to tests of differences of percentages among all persons in the specified subgroup who used cigarettes in the past month, regardless of whether they ever used cigarettes daily.

² Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.38C Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total		AGE FIRST USED MARIJUANA						Never Used Marijuana	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	235	215	141	128	111	133	148	87	60	55
AGE										
18-25	88	92	54	56	57	60	38	36	18	13
26-34	130	97	100	69	71	58	55	38	21	14
35-49	99	92	67	54	45	60	38	44	43	14
50 or Older	146	130	39	73	53	81	127	58	32	48
GENDER										
Male	168	170	113	110	88	106	82	73	36	23
Female	169	122	90	62	66	79	126	46	48	50
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	221	195	134	114	106	120	142	84	50	49
White	202	158	122	91	90	101	136	66	31	24
Black or African American	91	111	53	65	50	62	49	36	14	41
Other or Two or More Races	55	49	24	11	28	29	26	35	36	12
Hispanic or Latino	80	93	48	56	37	59	39	33	33	25
EDUCATION										
< High School	93	89	57	64	47	55	34	33	38	30
High School Graduate	123	118	83	67	70	80	45	38	43	15
Some College	161	126	83	72	61	72	128	45	15	41
College Graduate	85	95	44	52	42	51	55	58	10	8

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.38D Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		AGE FIRST USED MARIJUANA						Never Used Marijuana	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.10	0.09	0.67	0.59	0.28	0.31	0.29	0.17	0.05	0.04
AGE										
18-25	0.25	0.26	1.11	1.18	0.66	0.70	0.60	0.56	0.11	0.08
26-34	0.35	0.26	1.79	1.31	0.81	0.67	0.69	0.47	0.13	0.08
35-49	0.16	0.15	1.08	0.86	0.38	0.50	0.32	0.35	0.14	0.04
50 or Older	0.14	0.12	0.84	1.42	0.42	0.58	0.52	0.23	0.05	0.08
GENDER										
Male	0.15	0.15	0.94	0.86	0.41	0.45	0.32	0.27	0.07	0.04
Female	0.14	0.10	1.02	0.75	0.37	0.41	0.51	0.18	0.07	0.07
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	0.11	0.10	0.74	0.62	0.30	0.31	0.31	0.18	0.05	0.05
White	0.13	0.10	0.80	0.62	0.30	0.32	0.37	0.17	0.04	0.03
Black or African American	0.32	0.38	2.66	2.62	1.07	1.30	0.78	0.59	0.09	0.26
Other or Two or More Races	0.32	0.26	2.50	1.19	1.73	1.51	1.00	1.13	0.29	0.10
Hispanic or Latino	0.22	0.25	1.69	1.91	0.84	1.21	0.75	0.62	0.14	0.10
EDUCATION										
< High School	0.28	0.27	1.55	1.64	1.08	1.19	0.94	0.86	0.18	0.15
High School Graduate	0.18	0.17	1.26	0.98	0.56	0.62	0.38	0.31	0.11	0.04
Some College	0.25	0.19	1.32	1.10	0.50	0.53	0.84	0.30	0.05	0.13
College Graduate	0.12	0.13	0.99	1.28	0.38	0.44	0.28	0.28	0.03	0.02

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Standard Errors in the Total column correspond to percentages among all persons in the specified subgroup, regardless of whether they ever used marijuana.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.38P Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total ¹		AGE FIRST USED MARIJUANA						Never Used Marijuana	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.6075		0.9940		0.0189		0.2565		0.6377	
AGE										
18-25	0.0515		0.0767		0.5534		0.1694		0.3535	
26-34	0.6986		0.7824		0.5417		0.8998		0.3234	
35-49	0.1522		1.0000		0.0037		0.2502		0.0583	
50 or Older	0.1229		0.0295		0.0314		0.1750		0.1089	
GENDER										
Male	0.6048		0.4487		0.0498		0.6899		0.4121	
Female	0.8486		0.3276		0.2013		0.2534		0.9571	
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	0.7410		0.9213		0.0833		0.2842		0.9535	
White	0.5699		0.8834		0.2947		0.1485		0.3760	
Black or African American	0.0512		0.7556		0.0921		0.9805		0.0572	
Other or Two or More Races	0.6723		0.0910		0.9126		0.3355		0.2548	
Hispanic or Latino	0.5849		0.8283		0.0606		0.7555		0.3071	
EDUCATION										
< High School	0.7826		0.6258		0.8011		0.7062		0.6775	
High School Graduate	0.5111		0.1121		0.2936		0.7051		0.0579	
Some College	0.7369		0.5514		0.1522		0.1039		0.0708	
College Graduate	0.0474		0.1458		0.0079		0.7064		0.5378	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ P Values in the Total column correspond to tests of differences of percentages among all persons in the specified subgroup, regardless of whether they ever used marijuana.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.39C Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total		AGE FIRST USED ALCOHOL					
			14 or Younger		15-17		18 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	413	358	246	201	294	233	204	180
AGE								
18-25	123	120	71	72	89	91	56	60
26-34	220	169	142	99	165	95	93	87
35-49	203	159	143	107	133	111	90	75
50 or Older	270	239	132	132	180	161	151	128
GENDER								
Male	341	286	204	171	250	193	160	148
Female	241	191	136	106	159	125	127	97
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	360	317	211	190	258	205	185	157
White	362	292	191	171	234	179	171	132
Black or African American	126	127	72	72	81	83	62	67
Other or Two or More Races	98	92	63	42	73	63	51	47
Hispanic or Latino	189	156	127	85	125	100	85	90
EDUCATION								
< High School	147	133	84	69	89	94	81	55
High School Graduate	229	182	125	100	158	115	113	98
Some College	223	194	130	116	149	114	95	93
College Graduate	291	224	151	117	188	139	121	97

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.39D Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		AGE FIRST USED ALCOHOL					
			14 or Younger		15-17		18 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.17	0.15	0.59	0.48	0.36	0.29	0.22	0.19
AGE								
18-25	0.35	0.34	0.95	1.01	0.61	0.63	0.48	0.52
26-34	0.59	0.45	1.66	1.19	1.12	0.74	0.66	0.54
35-49	0.34	0.26	1.05	0.82	0.63	0.53	0.42	0.34
50 or Older	0.26	0.22	0.98	0.91	0.57	0.49	0.34	0.27
GENDER								
Male	0.30	0.25	0.82	0.68	0.57	0.44	0.42	0.36
Female	0.20	0.15	0.84	0.65	0.43	0.35	0.24	0.18
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.18	0.16	0.60	0.51	0.37	0.30	0.24	0.20
White	0.22	0.18	0.63	0.55	0.39	0.31	0.30	0.23
Black or African American	0.43	0.41	2.35	2.00	1.07	1.14	0.51	0.51
Other or Two or More Races	0.57	0.49	3.21	1.94	1.92	1.59	0.69	0.58
Hispanic or Latino	0.53	0.42	2.19	1.59	1.25	0.97	0.61	0.59
EDUCATION								
< High School	0.44	0.40	1.57	1.37	1.09	1.07	0.76	0.50
High School Graduate	0.32	0.25	1.17	0.92	0.70	0.52	0.41	0.36
Some College	0.34	0.29	1.15	0.92	0.65	0.50	0.38	0.35
College Graduate	0.38	0.29	1.10	0.89	0.70	0.54	0.42	0.31

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Standard Errors in the Total column correspond to percentages among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.39P Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total ¹		AGE FIRST USED ALCOHOL					
			14 or Younger		15-17		18 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.3847		0.3227		0.4005		0.5936	
AGE								
18-25	0.1249		0.0127		0.6527		0.4260	
26-34	0.0617		0.0699		0.0238		0.2042	
35-49	0.9663		0.6329		0.3836		0.8341	
50 or Older	0.3405		0.1151		0.7777		0.7434	
GENDER								
Male	0.5615		0.2564		0.8305		0.6522	
Female	0.4820		0.9559		0.1972		0.6981	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.5417		0.5568		0.5323		0.8602	
White	0.2951		0.9395		0.2946		0.5716	
Black or African American	0.2535		0.5887		0.2711		0.3251	
Other or Two or More Races	0.9060		0.1704		0.7146		0.2946	
Hispanic or Latino	0.4408		0.3860		0.4245		0.4211	
EDUCATION								
< High School	0.8542		0.4460		0.4578		0.4250	
High School Graduate	0.5847		0.1100		0.8396		0.8414	
Some College	0.9508		0.8811		0.2256		0.0950	
College Graduate	0.3420		0.9875		0.5048		0.8286	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ P Values in the Total column correspond to tests of differences of percentages among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150403

Table 6.40C Alcohol Dependence or Abuse in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Group	Total		AGE FIRST USED ALCOHOL							
			14 or Younger		15-17		18-20		21 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
21 OR OLDER	435	362	244	197	286	228	160	153	122	82
21 to 25	123	116	56	59	70	78	48	47	18	25
26 or Older	392	324	237	187	276	211	154	146	121	80
26 to 49	302	225	202	140	207	146	108	93	68	66
26 to 34	220	169	142	99	165	95	82	66	43	55
35 to 49	203	159	143	107	133	111	75	61	49	42
50 or Older	270	239	132	132	180	161	111	117	102	44

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.40D Alcohol Dependence or Abuse in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Standard Errors of Percentages, 2013 and 2014

Age Group	Total ¹		AGE FIRST USED ALCOHOL							
			14 or Younger		15-17		18-20		21 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
21 OR OLDER	0.18	0.15	0.63	0.50	0.38	0.30	0.31	0.28	0.33	0.21
21 to 25	0.48	0.47	1.21	1.29	0.80	0.89	0.85	0.87	0.56	0.71
26 or Older	0.19	0.16	0.68	0.53	0.41	0.31	0.33	0.30	0.35	0.22
26 to 49	0.31	0.23	0.91	0.66	0.58	0.43	0.55	0.43	0.43	0.39
26 to 34	0.59	0.45	1.66	1.19	1.12	0.74	1.05	0.75	0.70	0.79
35 to 49	0.34	0.26	1.05	0.82	0.63	0.53	0.64	0.48	0.51	0.42
50 or Older	0.26	0.22	0.98	0.91	0.57	0.49	0.41	0.42	0.56	0.23

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Standard Errors in the Total column correspond to percentages among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.40P Alcohol Dependence or Abuse in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Group	Total ¹		AGE FIRST USED ALCOHOL							
			14 or Younger		15-17		18-20		21 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
21 OR OLDER	0.5760		0.4034		0.5228		0.4692		0.9736	
21 to 25	0.4785		0.0096		0.8240		0.2177		0.5789	
26 or Older	0.7025		0.8203		0.4732		0.6263		0.8620	
26 to 49	0.1908		0.1459		0.2602		0.9702		0.1373	
26 to 34	0.0617		0.0699		0.0238		0.7480		0.0786	
35 to 49	0.9663		0.6329		0.3836		0.6128		0.8117	
50 or Older	0.3405		0.1151		0.7777		0.5459		0.1410	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ P Values in the Total column correspond to tests of differences of percentages among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.41C Illicit Drug Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total		AGE FIRST USED MARIJUANA						Never Used Marijuana	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	208	182	127	107	94	113	141	76	35	35
AGE										
18-25	73	80	46	49	45	51	32	29	10	13
26-34	119	87	93	62	64	53	52	33	12	14
35-49	80	82	58	49	39	56	24	35	27	7
50 or Older	137	100	33	50	46	61	125	47	16	30
GENDER										
Male	142	138	98	90	70	86	75	61	17	18
Female	160	103	85	57	63	67	123	40	30	32
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	200	168	121	95	92	103	137	71	34	35
White	188	132	111	81	79	81	132	52	21	23
Black or African American	77	97	48	48	41	60	41	33	12	24
Other or Two or More Races	45	45	22	10	25	27	25	32	25	12
Hispanic or Latino	59	71	40	47	28	44	32	28	5	5
EDUCATION										
< High School	79	70	53	45	43	44	33	28	12	20
High School Graduate	102	101	73	60	59	64	41	36	29	12
Some College	154	106	80	63	54	63	124	32	10	26
College Graduate	75	81	36	44	37	46	51	46	10	7

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.41D Illicit Drug Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		AGE FIRST USED MARIJUANA						Never Used Marijuana	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.09	0.08	0.60	0.50	0.24	0.26	0.28	0.15	0.03	0.03
AGE										
18-25	0.21	0.23	0.98	1.06	0.54	0.62	0.53	0.45	0.06	0.08
26-34	0.32	0.23	1.69	1.20	0.73	0.62	0.65	0.40	0.07	0.08
35-49	0.13	0.14	0.93	0.79	0.33	0.47	0.20	0.27	0.09	0.02
50 or Older	0.13	0.09	0.69	1.01	0.37	0.44	0.51	0.19	0.02	0.05
GENDER										
Male	0.12	0.12	0.82	0.73	0.32	0.37	0.29	0.23	0.03	0.03
Female	0.13	0.08	0.97	0.69	0.35	0.35	0.50	0.16	0.04	0.04
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	0.10	0.08	0.67	0.52	0.26	0.27	0.30	0.15	0.03	0.03
White	0.12	0.08	0.73	0.55	0.27	0.26	0.36	0.14	0.03	0.03
Black or African American	0.27	0.33	2.43	2.03	0.87	1.27	0.67	0.54	0.08	0.15
Other or Two or More Races	0.26	0.24	2.28	1.11	1.52	1.38	0.97	1.05	0.20	0.09
Hispanic or Latino	0.16	0.19	1.43	1.64	0.64	0.94	0.62	0.53	0.02	0.02
EDUCATION										
< High School	0.24	0.22	1.41	1.24	0.98	0.98	0.91	0.73	0.06	0.10
High School Graduate	0.14	0.14	1.12	0.90	0.47	0.50	0.34	0.29	0.07	0.03
Some College	0.24	0.16	1.27	0.98	0.44	0.47	0.82	0.21	0.03	0.08
College Graduate	0.10	0.11	0.81	1.10	0.33	0.40	0.26	0.22	0.03	0.02

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Standard Errors in the Total column correspond to percentages among all persons in the specified subgroup, regardless of whether they ever used marijuana.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.41P Illicit Drug Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Marijuana and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total ¹		AGE FIRST USED MARIJUANA						Never Used Marijuana	
			14 or Younger		15-17		18 or Older			
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.8659		0.7520		0.0171		0.0932		0.6131	
AGE										
18-25	0.1528		0.1482		0.9249		0.0504		0.6376	
26-34	0.6256		0.6474		0.5250		0.4787		0.5455	
35-49	0.0401		0.8204		0.0027		0.1638		0.1874	
50 or Older	0.7597		0.1274		0.1452		0.1236		0.2115	
GENDER										
Male	0.6494		0.3554		0.0116		0.2474		0.7618	
Female	0.8203		0.5101		0.4780		0.2191		0.6906	
HISPANIC ORIGIN AND RACE										
Not Hispanic or Latino	0.9698		0.7172		0.0662		0.1131		0.5352	
White	0.2923		0.7645		0.5044		0.0656		0.9153	
Black or African American	0.0538		1.0000		0.0206		0.9388		0.1748	
Other or Two or More Races	0.9333		0.1631		0.7424		0.7829		0.9153	
Hispanic or Latino	0.5204		0.9546		0.0658		0.5784		0.5150	
EDUCATION										
< High School	0.7057		0.2714		0.8045		1.0000		0.2310	
High School Graduate	0.7882		0.2562		0.4178		0.9352		0.2343	
Some College	0.8378		0.9781		0.1015		0.0753		0.0939	
College Graduate	0.1219		0.0864		0.0082		0.5601		0.3302	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ P Values in the Total column correspond to tests of differences of percentages among all persons in the specified subgroup, regardless of whether they ever used marijuana.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.42C Alcohol Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Total		AGE FIRST USED ALCOHOL					
			14 or Younger		15-17		18 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	286	244	161	138	191	153	141	115
AGE								
18-25	78	78	46	47	59	61	29	37
26-34	159	102	98	67	111	63	63	50
35-49	134	117	99	72	72	76	67	55
50 or Older	187	165	81	101	124	103	108	86
GENDER								
Male	231	190	132	108	156	121	118	93
Female	165	146	99	82	111	93	76	71
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	259	227	148	133	165	141	133	107
White	238	195	130	116	151	117	118	90
Black or African American	87	104	58	61	43	62	49	50
Other or Two or More Races	73	68	52	38	37	44	36	34
Hispanic or Latino	124	94	69	58	93	57	48	46
EDUCATION								
< High School	104	104	69	53	51	70	60	37
High School Graduate	155	123	85	75	97	78	82	63
Some College	138	144	77	92	87	77	75	68
College Graduate	186	132	101	70	131	85	73	59

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.42D Alcohol Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		AGE FIRST USED ALCOHOL					
			14 or Younger		15-17		18 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.12	0.10	0.41	0.34	0.24	0.20	0.16	0.12
AGE								
18-25	0.22	0.22	0.70	0.73	0.44	0.46	0.27	0.33
26-34	0.43	0.27	1.30	0.85	0.81	0.50	0.46	0.32
35-49	0.22	0.19	0.75	0.58	0.36	0.37	0.31	0.25
50 or Older	0.18	0.15	0.60	0.70	0.39	0.32	0.24	0.18
GENDER								
Male	0.20	0.16	0.54	0.45	0.36	0.28	0.31	0.23
Female	0.13	0.12	0.64	0.51	0.31	0.26	0.14	0.13
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.13	0.11	0.43	0.37	0.24	0.21	0.17	0.13
White	0.15	0.12	0.44	0.38	0.26	0.20	0.21	0.15
Black or African American	0.31	0.35	1.93	1.77	0.61	0.88	0.40	0.38
Other or Two or More Races	0.41	0.36	2.68	1.74	1.00	1.14	0.48	0.43
Hispanic or Latino	0.35	0.26	1.30	1.15	0.96	0.57	0.35	0.30
EDUCATION								
< High School	0.32	0.31	1.33	1.05	0.65	0.81	0.57	0.35
High School Graduate	0.22	0.18	0.82	0.69	0.44	0.36	0.30	0.23
Some College	0.22	0.22	0.70	0.76	0.40	0.35	0.30	0.26
College Graduate	0.25	0.18	0.76	0.55	0.50	0.34	0.26	0.19

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Standard Errors in the Total column correspond to percentages among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.42P Alcohol Dependence in the Past Year among Persons Aged 18 or Older, by Age First Used Alcohol and Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Total ¹		AGE FIRST USED ALCOHOL					
			14 or Younger		15-17		18 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014
TOTAL	0.9959		0.5095		0.8234		0.6322	
AGE								
18-25	0.5922		0.6427		0.8064		0.0300	
26-34	0.1222		0.2623		0.1267		0.7296	
35-49	0.7592		0.1125		0.0077		0.8299	
50 or Older	0.4266		0.0827		0.8925		0.8446	
GENDER								
Male	0.6582		0.2508		0.7048		0.9225	
Female	0.5541		0.6579		0.8718		0.3010	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.9121		0.4109		0.4278		0.9607	
White	0.4441		0.7437		0.5905		0.6494	
Black or African American	0.1499		0.4910		0.0282		0.3864	
Other or Two or More Races	0.4209		0.3046		0.0610		0.7692	
Hispanic or Latino	0.7971		0.7320		0.2440		0.2281	
EDUCATION								
< High School	0.5721		0.5300		0.0823		0.7343	
High School Graduate	0.9520		0.7798		0.9394		0.8655	
Some College	0.1361		0.3782		0.8301		0.1036	
College Graduate	0.0750		0.1855		0.3619		0.7125	

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ P Values in the Total column correspond to tests of differences of percentages among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

150403

Table 6.43C Alcohol Dependence in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Group	Total		AGE FIRST USED ALCOHOL							
			14 or Younger		15-17		18-20		21 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
21 OR OLDER	287	244	158	136	189	149	113	99	79	52
21 to 25	72	75	37	37	50	54	23	29	10	19
26 or Older	278	228	156	130	181	139	114	97	78	50
26 to 49	209	155	137	96	133	99	77	61	46	40
26 to 34	159	102	98	67	111	63	57	44	27	24
35 to 49	134	117	99	72	72	76	57	42	37	33
50 or Older	187	165	81	101	124	103	88	80	64	31

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.43D Alcohol Dependence in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: Standard Errors of Percentages, 2013 and 2014

Age Group	Total ¹		AGE FIRST USED ALCOHOL							
			14 or Younger		15-17		18-20		21 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
21 OR OLDER	0.13	0.11	0.42	0.35	0.26	0.20	0.22	0.18	0.21	0.13
21 to 25	0.31	0.32	0.90	0.89	0.62	0.66	0.43	0.56	0.33	0.55
26 or Older	0.14	0.11	0.46	0.37	0.28	0.21	0.25	0.20	0.23	0.14
26 to 49	0.21	0.16	0.66	0.48	0.39	0.30	0.40	0.29	0.29	0.24
26 to 34	0.43	0.27	1.30	0.85	0.81	0.50	0.75	0.51	0.43	0.34
35 to 49	0.22	0.19	0.75	0.58	0.36	0.37	0.49	0.34	0.38	0.33
50 or Older	0.18	0.15	0.60	0.70	0.39	0.32	0.33	0.28	0.35	0.17

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Standard Errors in the Total column correspond to percentages among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.43P Alcohol Dependence in the Past Year among Persons Aged 21 or Older, by Age First Used Alcohol and Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Group	Total ¹		AGE FIRST USED ALCOHOL							
			14 or Younger		15-17		18-20		21 or Older	
	2013	2014	2013	2014	2013	2014	2013	2014	2013	2014
21 OR OLDER	0.9564		0.5215		0.8188		0.4740		0.8205	
21 to 25	0.4310		0.5413		0.8118		0.0323		0.1611	
26 or Older	0.8803		0.6190		0.8588		0.8097		0.5772	
26 to 49	0.3737		0.0599		0.6282		0.9722		0.8564	
26 to 34	0.1222		0.2623		0.1267		0.7778		0.8763	
35 to 49	0.7592		0.1125		0.0077		0.6970		0.9146	
50 or Older	0.4266		0.0827		0.8925		0.7673		0.3738	

*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ P Values in the Total column correspond to tests of differences of percentages among all persons in the specified subgroup, regardless of whether they ever used alcohol.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.44C Method of Obtaining Most Recently Used Marijuana among Past Year Marijuana Users, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Method of Obtaining Most Recently Used Marijuana	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS	628	569	84	90	231	242	554	476
Bought It	395	382	53	55	130	152	359	327
Traded for It	78	49	8	8	17	18	77	45
Got It Free or Shared Someone Else's	411	355	60	67	159	161	363	301
Grew It Yourself	134	60	5	10	13	12	134	59
Method Unspecified	55	65	19	23	21	24	47	57

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.44D Method of Obtaining Most Recently Used Marijuana among Past Year Marijuana Users, by Age Group: Standard Errors of Percentages, 2013 and 2014

Method of Obtaining Most Recently Used Marijuana	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Bought It	0.83	0.71	1.18	1.35	0.81	0.90	1.37	1.04
Traded for It	0.24	0.14	0.24	0.25	0.15	0.16	0.41	0.22
Got It Free or Shared Someone Else's	0.84	0.70	1.24	1.35	0.82	0.93	1.36	1.04
Grew It Yourself	0.40	0.17	0.15	0.29	0.12	0.11	0.71	0.28
Method Unspecified	0.17	0.18	0.56	0.68	0.19	0.21	0.25	0.27

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.44P Method of Obtaining Most Recently Used Marijuana among Past Year Marijuana Users, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Method of Obtaining Most Recently Used Marijuana	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS	N/A		N/A		N/A		N/A	
Bought It	0.0123		0.9710		0.1115		0.0433	
Traded for It	0.0477		0.9309		0.2547		0.0727	
Got It Free or Shared Someone Else's	0.1890		0.3567		0.1520		0.5655	
Grew It Yourself	0.0074		0.1831		0.4413		0.0057	
Method Unspecified	0.1047		0.1887		0.4531		0.2232	

*Low precision; no estimate reported.

N/A: Not applicable.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.45C Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Bought Their Most Recently Used Marijuana, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Source and Location of Bought Marijuana	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS WHO BOUGHT MOST RECENTLY USED MARIJUANA	395	382	53	55	130	152	359	327
SOURCE								
Friend	339	306	45	50	111	123	312	263
Relative or Family Member	113	76	10	8	18	25	111	71
Someone Just Met or Did Not Know Well	140	145	20	19	52	61	124	128
Source Unspecified	65	86	11	10	30	26	58	79
LOCATION								
Inside Public Building ¹	95	112	9	13	32	41	88	104
Outside in Public Area ²	153	146	30	30	57	65	136	124
Inside School Building	17	21	11	12	11	13	8	12
Outside on School Property	24	25	9	8	8	15	20	19
Inside Home, Apartment, or Dorm	320	269	29	35	94	108	303	238
Medical Marijuana Dispensary/Club ³	71	80	1	1	24	20	64	77
Some Other Place ⁴	31	32	9	5	11	12	29	29
Location Unspecified ⁵	115	125	19	19	45	44	104	114

*Low precision; no estimate reported.

¹ For example, a store, restaurant, sports arena, bar, or club.

² For example, a parking lot, street, or park.

³ Medical Marijuana Dispensary/Club includes respondents who reported that they bought marijuana in some other place and specified that this other location was a medical marijuana dispensary or a marijuana club.

⁴ Some Other Place includes respondents who reported that they bought marijuana in some other place and specified a location other than a medical marijuana dispensary/club that did not correspond to an existing category. Examples of these other locations include private backyard, prison, and hotel room. Because of revisions to this category to remove Medical Marijuana Dispensary/Club in 2013, these estimates are not comparable with the estimates presented before the 2013 NSDUH.

⁵ Location Unspecified includes respondents with unknown, invalid, or nonspecific responses to the other-specify question (e.g., in a car, at a concert, or at a party where the specific locations were not provided).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.45D Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Bought Their Most Recently Used Marijuana, by Age Group: Standard Errors of Percentages, 2013 and 2014

Source and Location of Bought Marijuana	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS WHO BOUGHT MOST RECENTLY USED MARIJUANA	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
SOURCE								
Friend	1.11	0.92	1.67	1.67	1.08	1.12	1.80	1.32
Relative or Family Member	0.74	0.43	0.74	0.66	0.35	0.47	1.25	0.66
Someone Just Met or Did Not Know Well	0.88	0.77	1.44	1.44	0.95	0.98	1.37	1.14
Source Unspecified	0.43	0.50	0.84	0.81	0.59	0.48	0.67	0.75
LOCATION								
Inside Public Building ¹	0.63	0.64	0.69	1.02	0.62	0.75	1.02	0.96
Outside in Public Area ²	0.97	0.77	1.81	1.90	1.02	1.06	1.51	1.12
Inside School Building	0.12	0.13	0.84	0.94	0.22	0.24	0.09	0.12
Outside on School Property	0.17	0.15	0.67	0.61	0.17	0.29	0.23	0.19
Inside Home, Apartment, or Dorm	1.27	1.05	1.76	2.12	1.20	1.28	2.07	1.59
Medical Marijuana Dispensary/Club ³	0.47	0.47	0.04	0.07	0.48	0.38	0.75	0.74
Some Other Place ⁴	0.21	0.19	0.68	0.39	0.22	0.23	0.34	0.28
Location Unspecified ⁵	0.75	0.70	1.33	1.44	0.83	0.80	1.17	1.04

*Low precision; no estimate reported.

¹ For example, a store, restaurant, sports arena, bar, or club.

² For example, a parking lot, street, or park.

³ Medical Marijuana Dispensary/Club includes respondents who reported that they bought marijuana in some other place and specified that this other location was a medical marijuana dispensary or a marijuana club.

⁴ Some Other Place includes respondents who reported that they bought marijuana in some other place and specified a location other than a medical marijuana dispensary/club that did not correspond to an existing category. Examples of these other locations include private backyard, prison, and hotel room. Because of revisions to this category to remove Medical Marijuana Dispensary/Club in 2013, these estimates are not comparable with the estimates presented before the 2013 NSDUH.

⁵ Location Unspecified includes respondents with unknown, invalid, or nonspecific responses to the other-specify question (e.g., in a car, at a concert, or at a party where the specific locations were not provided).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.45P Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Bought Their Most Recently Used Marijuana, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Source and Location of Bought Marijuana	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS WHO BOUGHT MOST RECENTLY USED MARIJUANA	N/A		N/A		N/A		N/A	
SOURCE								
Friend	0.1106		0.0357		0.3456		0.1079	
Relative or Family Member	0.3179		0.3028		0.1835		0.1749	
Someone Just Met or Did Not Know Well	0.3352		0.0737		0.4631		0.2781	
Source Unspecified	0.0021		0.7994		0.6834		0.0004	
LOCATION								
Inside Public Building ¹	0.0167		0.0803		0.0921		0.1044	
Outside in Public Area ²	0.1876		0.6626		0.7799		0.3374	
Inside School Building	0.7428		0.6166		0.9727		0.4268	
Outside on School Property	0.8358		0.6554		0.0991		0.6966	
Inside Home, Apartment, or Dorm	0.1424		0.2297		0.8217		0.0663	
Medical Marijuana Dispensary/Club ³	0.0691		0.5102		0.7319		0.0538	
Some Other Place ⁴	0.7672		0.3357		0.6276		0.7238	
Location Unspecified ⁵	0.5508		0.1886		0.2511		0.1709	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ For example, a store, restaurant, sports arena, bar, or club.

² For example, a parking lot, street, or park.

³ Medical Marijuana Dispensary/Club includes respondents who reported that they bought marijuana in some other place and specified that this other location was a medical marijuana dispensary or a marijuana club.

⁴ Some Other Place includes respondents who reported that they bought marijuana in some other place and specified a location other than a medical marijuana dispensary/club that did not correspond to an existing category. Examples of these other locations include private backyard, prison, and hotel room. Because of revisions to this category to remove Medical Marijuana Dispensary/Club in 2013, these estimates are not comparable with the estimates presented before the 2013 NSDUH.

⁵ Location Unspecified includes respondents with unknown, invalid, or nonspecific responses to the other-specify question (e.g., in a car, at a concert, or at a party where the specific locations were not provided).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.46C Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Obtained for Free Their Most Recently Used Marijuana and Who Had Not Bought or Traded for Marijuana in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Source and Location of Free Marijuana ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS WHO OBTAINED FOR FREE MOST RECENTLY USED MARIJUANA AND WHO HAD NOT BOUGHT OR TRADED FOR MARIJUANA IN PAST YEAR	359	311	53	56	132	134	324	268
SOURCE								
Friend	317	278	48	50	121	119	284	238
Relative or Family Member	128	124	16	17	34	41	121	116
Someone Just Met or Did Not Know Well	79	65	11	15	25	31	75	54
Source Unspecified	43	37	9	8	11	13	42	32
LOCATION								
Inside Public Building ²	85	54	8	6	19	16	82	51
Outside in Public Area ³	102	97	26	25	43	46	90	80
Inside School Building	30	16	9	7	7	11	28	9
Outside on School Property	20	20	13	11	13	15	9	5
Inside Home, Apartment, or Dorm	304	263	35	40	110	114	272	232
Some Other Place ⁴	46	37	7	4	11	11	44	36
Location Unspecified ⁵	113	92	17	20	25	35	109	84

*Low precision; no estimate reported.

¹ Respondents who reported that they had bought or traded for marijuana in the past year or who had missing data for whether they bought or traded for marijuana in the past year were excluded from this analysis.

² For example, a store, restaurant, sports arena, bar, or club.

³ For example, a parking lot, street, or park.

⁴ Some Other Place includes respondents who reported that they obtained marijuana for free in some other place and specified a location (e.g., private backyard, prison, hotel room) that did not correspond to an existing category.

⁵ Location Unspecified includes respondents with unknown, invalid, or nonspecific responses to the other-specify question (e.g., in a car, at a concert, or at a party where the specific locations were not provided).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.46D Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Obtained for Free Their Most Recently Used Marijuana and Who Had Not Bought or Traded for Marijuana in the Past Year, by Age Group: Standard Errors of Percentages, 2013 and 2014

Source and Location of Free Marijuana ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS WHO OBTAINED FOR FREE MOST RECENTLY USED MARIJUANA AND WHO HAD NOT BOUGHT OR TRADED FOR MARIJUANA IN PAST YEAR	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
SOURCE								
Friend	1.15	1.02	1.52	1.73	0.95	1.18	1.91	1.57
Relative or Family Member	0.93	0.88	1.19	1.29	0.76	0.94	1.57	1.40
Someone Just Met or Did Not Know Well	0.61	0.51	0.81	1.14	0.57	0.73	1.02	0.72
Source Unspecified	0.34	0.29	0.64	0.66	0.25	0.32	0.59	0.42
LOCATION								
Inside Public Building ²	0.66	0.42	0.63	0.51	0.44	0.40	1.13	0.66
Outside in Public Area ³	0.79	0.70	1.69	1.79	0.93	1.03	1.23	1.03
Inside School Building	0.24	0.12	0.64	0.58	0.17	0.26	0.39	0.12
Outside on School Property	0.16	0.15	0.92	0.88	0.31	0.37	0.13	0.07
Inside Home, Apartment, or Dorm	1.26	1.00	1.87	2.08	1.15	1.35	2.07	1.51
Some Other Place ⁴	0.36	0.29	0.49	0.31	0.26	0.27	0.61	0.48
Location Unspecified ⁵	0.84	0.69	1.16	1.50	0.58	0.82	1.42	1.06

*Low precision; no estimate reported.

¹ Respondents who reported that they had bought or traded for marijuana in the past year or who had missing data for whether they bought or traded for marijuana in the past year were excluded from this analysis.

² For example, a store, restaurant, sports arena, bar, or club.

³ For example, a parking lot, street, or park.

⁴ Some Other Place includes respondents who reported that they obtained marijuana for free in some other place and specified a location (e.g., private backyard, prison, hotel room) that did not correspond to an existing category.

⁵ Location Unspecified includes respondents with unknown, invalid, or nonspecific responses to the other-specify question (e.g., in a car, at a concert, or at a party where the specific locations were not provided).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.46P Source and Location Where Marijuana Was Obtained among Past Year Marijuana Users Who Obtained for Free Their Most Recently Used Marijuana and Who Had Not Bought or Traded for Marijuana in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Source and Location of Free Marijuana ¹	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL PAST YEAR MARIJUANA USERS WHO OBTAINED FOR FREE MOST RECENTLY USED MARIJUANA AND WHO HAD NOT BOUGHT OR TRADED FOR MARIJUANA IN PAST YEAR	N/A		N/A		N/A		N/A	
SOURCE								
Friend	0.1225		0.2488		0.1477		0.3714	
Relative or Family Member	0.1048		0.7703		0.1561		0.2752	
Someone Just Met or Did Not Know Well	0.5926		0.1065		0.7758		0.8416	
Source Unspecified	0.7570		0.9021		0.6038		0.6436	
LOCATION								
Inside Public Building ²	0.7902		0.3501		0.1510		0.8930	
Outside in Public Area ³	0.9248		0.1030		0.7414		0.5001	
Inside School Building	0.1502		0.2635		0.6820		0.1853	
Outside on School Property	0.7856		0.9437		0.3125		0.6883	
Inside Home, Apartment, or Dorm	0.7023		0.0810		0.6785		0.8871	
Some Other Place ⁴	0.9310		0.2106		0.7617		0.9084	
Location Unspecified ⁵	0.9841		0.3806		0.1895		0.5316	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Respondents who reported that they had bought or traded for marijuana in the past year or who had missing data for whether they bought or traded for marijuana in the past year were excluded from this analysis.

² For example, a store, restaurant, sports arena, bar, or club.

³ For example, a parking lot, street, or park.

⁴ Some Other Place includes respondents who reported that they obtained marijuana for free in some other place and specified a location (e.g., private backyard, prison, hotel room) that did not correspond to an existing category.

⁵ Location Unspecified includes respondents with unknown, invalid, or nonspecific responses to the other-specify question (e.g., in a car, at a concert, or at a party where the specific locations were not provided).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.47C Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	108	106	17	16	27	27	103	101
From More Than One Doctor	44	43	5	5	9	10	43	42
Wrote Fake Prescription	6	3	4	1	4	2	3	1
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	26	14	4	2	6	3	26	14
From Friend or Relative for Free	164	139	22	21	52	47	154	130
Bought from Friend or Relative	61	67	9	9	26	25	54	60
Took from Friend or Relative without Asking	39	37	10	10	16	15	35	33
Bought from Drug Dealer or Other Stranger	33	36	6	7	17	19	28	30
Bought on the Internet	13	3	1	1	4	3	12	1
Some Other Way	48	32	8	8	11	14	46	29

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.47D Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.87	0.91	1.27	1.47	0.77	0.92	1.37	1.35
From More Than One Doctor	0.40	0.44	0.44	0.57	0.29	0.37	0.65	0.67
Wrote Fake Prescription	0.06	0.03	0.32	0.12	0.11	0.09	0.05	0.02
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.24	0.15	0.34	0.18	0.18	0.10	0.39	0.23
From Friend or Relative for Free	1.00	1.00	1.44	1.63	1.04	1.23	1.60	1.50
Bought from Friend or Relative	0.52	0.64	0.81	0.95	0.74	0.86	0.79	0.94
Took from Friend or Relative without Asking	0.36	0.38	0.81	0.99	0.47	0.55	0.53	0.53
Bought from Drug Dealer or Other Stranger	0.31	0.36	0.53	0.72	0.52	0.68	0.42	0.49
Bought on the Internet	0.12	0.03	0.09	0.09	0.12	0.09	0.19	0.02
Some Other Way	0.44	0.33	0.69	0.84	0.34	0.52	0.70	0.47

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

**Table 6.47P Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.0617		0.4660		0.1110		0.2063	
From More Than One Doctor	0.0302		0.1496		0.3203		0.0766	
Wrote Fake Prescription	0.0470		0.1995		0.4131		0.2371	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1146		0.1294		0.1119		0.2763	
From Friend or Relative for Free	0.0118		0.1747		0.1483		0.0394	
Bought from Friend or Relative	0.8909		0.8865		0.1036		0.3324	
Took from Friend or Relative without Asking	0.4152		0.6895		0.2970		0.4446	
Bought from Drug Dealer or Other Stranger	0.3298		0.3688		0.3979		0.4963	
Bought on the Internet	0.3556		0.8536		0.7990		0.2982	
Some Other Way	0.4324		0.3259		0.2532		0.1736	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.48C Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Pain Relievers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	148	130	16	15	41	41	142	122
From More Than One Doctor	28	25	5	4	11	7	23	23
Wrote Fake Prescription	11	1	*	0	2	1	10	0
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	6	1	1	0	5	1	2	0
From Friend or Relative for Free	29	23	6	6	12	9	25	20
Bought from Friend or Relative	23	21	5	5	13	11	19	18
Took from Friend or Relative without Asking	12	8	5	5	8	5	7	4
Bought from Drug Dealer or Other Stranger	14	11	4	4	8	5	11	9
Bought on the Internet	8	5	0	1	1	4	8	2
Some Other Way	14	12	4	4	4	4	13	10

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.48D Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Pain Relievers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	1.03	0.93	2.18	2.70	1.36	1.40	1.37	1.23
From More Than One Doctor	0.51	0.54	1.15	1.26	0.70	0.56	0.69	0.77
Wrote Fake Prescription	0.20	0.03	*	0.10	0.13	0.09	0.31	0.01
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.11	0.03	0.32	0.11	0.33	0.12	0.06	0.01
From Friend or Relative for Free	0.53	0.51	1.29	1.66	0.75	0.76	0.76	0.67
Bought from Friend or Relative	0.44	0.46	1.12	1.47	0.83	0.90	0.57	0.58
Took from Friend or Relative without Asking	0.23	0.18	1.11	1.48	0.56	0.42	0.21	0.12
Bought from Drug Dealer or Other Stranger	0.27	0.24	1.00	1.07	0.49	0.43	0.34	0.29
Bought on the Internet	0.15	0.11	0.05	0.45	0.08	0.34	0.23	0.08
Some Other Way	0.27	0.26	0.85	1.14	0.26	0.34	0.40	0.33

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.48P Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source Where Friend or Relative Obtained Pain Relievers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.0994		0.4443		0.0029		1.0000	
From More Than One Doctor	0.7497		0.4910		0.0758		0.6027	
Wrote Fake Prescription	0.1764		*		0.8969		0.1669	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.0079		0.0957		0.0446		0.2848	
From Friend or Relative for Free	0.3823		0.7143		0.4566		0.6665	
Bought from Friend or Relative	0.5876		0.5421		0.1545		0.5280	
Took from Friend or Relative without Asking	0.2308		0.8727		0.1880		0.9889	
Bought from Drug Dealer or Other Stranger	0.7720		0.5263		0.3147		0.8301	
Bought on the Internet	0.9186		0.2549		0.4110		0.3953	
Some Other Way	0.9425		0.8232		0.5353		0.9917	

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.49C Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	74	69	10	11	21	18	70	65
From More Than One Doctor	27	43	4	5	9	9	25	41
Wrote Fake Prescription	14	7	3	1	1	1	13	6
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	20	15	3	1	4	2	19	15
From Friend or Relative for Free	108	91	13	12	36	30	100	86
Bought from Friend or Relative	64	62	8	7	23	22	60	58
Took from Friend or Relative without Asking	43	35	7	6	14	13	41	32
Bought from Drug Dealer or Other Stranger	37	40	5	6	18	16	31	37
Bought on the Internet	17	5	1	*	6	1	16	5
Some Other Way	25	22	5	6	11	11	22	19

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.49D Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	1.46	1.47	1.94	2.54	1.54	1.71	2.22	2.06
From More Than One Doctor	0.63	1.02	0.96	1.34	0.74	0.94	0.95	1.45
Wrote Fake Prescription	0.32	0.17	0.76	0.33	0.11	0.10	0.49	0.24
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.46	0.37	0.60	0.32	0.33	0.22	0.72	0.55
From Friend or Relative for Free	1.64	1.65	2.17	2.67	1.72	2.04	2.50	2.33
Bought from Friend or Relative	1.38	1.40	1.62	1.91	1.60	1.90	2.12	1.95
Took from Friend or Relative without Asking	0.97	0.85	1.56	1.72	1.09	1.27	1.47	1.16
Bought from Drug Dealer or Other Stranger	0.88	0.97	1.17	1.73	1.37	1.47	1.18	1.31
Bought on the Internet	0.39	0.12	0.27	*	0.50	0.08	0.59	0.17
Some Other Way	0.60	0.55	1.15	1.59	0.96	1.10	0.84	0.73

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.49P Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.7557		0.1605		0.7505		0.9643	
From More Than One Doctor	0.0103		0.0703		0.5951		0.0233	
Wrote Fake Prescription	0.3183		0.2485		0.9153		0.4381	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2739		0.0477		0.0578		0.5407	
From Friend or Relative for Free	0.0008		0.1130		0.0016		0.0166	
Bought from Friend or Relative	0.0707		0.2509		0.1698		0.2128	
Took from Friend or Relative without Asking	0.2277		0.3742		0.6258		0.4333	
Bought from Drug Dealer or Other Stranger	0.8150		0.7939		0.0748		0.1926	
Bought on the Internet	0.0368		*		0.0258		0.1992	
Some Other Way	0.8862		0.3579		0.7788		0.9405	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.50C Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Pain Relievers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	96	80	9	10	28	25	91	76
From More Than One Doctor	25	17	4	4	13	7	21	15
Wrote Fake Prescription	17	6	1	*	2	3	17	6
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	11	6	1	*	5	3	10	5
From Friend or Relative for Free	33	23	4	5	14	11	29	19
Bought from Friend or Relative	29	23	4	4	14	12	25	19
Took from Friend or Relative without Asking	18	13	3	3	11	6	14	12
Bought from Drug Dealer or Other Stranger	23	18	4	3	12	9	19	15
Bought on the Internet	12	5	*	*	3	*	11	2
Some Other Way	15	9	3	3	6	3	13	9

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.50D Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Pain Relievers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	1.64	1.34	3.27	4.19	2.12	2.12	2.26	1.73
From More Than One Doctor	1.05	0.90	2.23	2.64	1.78	1.38	1.39	1.19
Wrote Fake Prescription	0.73	0.35	0.64	*	0.30	0.59	1.12	0.44
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.46	0.33	0.79	*	0.69	0.73	0.66	0.37
From Friend or Relative for Free	1.34	1.18	2.26	3.43	1.86	2.24	1.87	1.49
Bought from Friend or Relative	1.16	1.17	2.13	3.08	1.88	2.30	1.59	1.44
Took from Friend or Relative without Asking	0.77	0.71	1.77	2.55	1.62	1.18	0.92	0.91
Bought from Drug Dealer or Other Stranger	0.95	0.96	2.27	2.42	1.67	1.85	1.24	1.20
Bought on the Internet	0.50	0.28	*	*	0.51	*	0.74	0.13
Some Other Way	0.64	0.51	1.40	2.09	0.91	0.67	0.90	0.68

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.50P Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source Where Friend or Relative Obtained Pain Relievers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.0028		0.2004		0.0097		0.0908	
From More Than One Doctor	0.3047		0.2810		0.0644		0.9169	
Wrote Fake Prescription	0.1475		*		0.5030		0.0882	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7525		*		0.5967		0.5273	
From Friend or Relative for Free	0.1056		0.6244		0.7829		0.0927	
Bought from Friend or Relative	0.1324		0.7143		0.4404		0.2871	
Took from Friend or Relative without Asking	0.2859		0.3436		0.1561		0.9965	
Bought from Drug Dealer or Other Stranger	0.6930		0.8495		0.7537		0.9744	
Bought on the Internet	0.5262		*		*		0.2133	
Some Other Way	0.3510		0.6013		0.1994		0.5903	

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.51C Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	61	60	6	6	12	13	59	59
From More Than One Doctor	11	17	2	2	7	6	8	15
Wrote Fake Prescription	1	3	1	*	1	1	*	3
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	12	1	1	3	3	4	12
From Friend or Relative for Free	131	114	14	13	33	33	126	107
Bought from Friend or Relative	53	40	8	8	21	22	48	31
Took from Friend or Relative without Asking	22	25	6	6	12	11	17	22
Bought from Drug Dealer or Other Stranger	29	24	6	5	15	15	23	18
Bought on the Internet	1	4	*	1	1	2	*	3
Some Other Way	24	28	3	4	11	9	21	26

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.51D Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	1.08	1.13	1.41	1.79	0.77	0.88	1.63	1.69
From More Than One Doctor	0.20	0.34	0.44	0.66	0.43	0.40	0.25	0.49
Wrote Fake Prescription	0.03	0.06	0.31	*	0.04	0.07	*	0.08
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.09	0.25	0.13	0.20	0.20	0.22	0.11	0.36
From Friend or Relative for Free	1.43	1.38	2.40	3.01	1.48	1.61	2.04	2.00
Bought from Friend or Relative	0.95	0.80	1.97	2.33	1.21	1.38	1.36	0.97
Took from Friend or Relative without Asking	0.42	0.51	1.47	1.77	0.76	0.76	0.52	0.68
Bought from Drug Dealer or Other Stranger	0.57	0.50	1.53	1.63	0.96	1.02	0.69	0.58
Bought on the Internet	0.01	0.09	*	0.40	0.04	0.18	*	0.11
Some Other Way	0.45	0.56	0.79	1.35	0.72	0.63	0.62	0.81

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

**Table 6.51P Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.8661		0.6348		0.5443		0.9444	
From More Than One Doctor	0.2878		0.2125		0.9184		0.2855	
Wrote Fake Prescription	0.4284		*		0.4396		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3419		0.7274		0.9943		0.3197	
From Friend or Relative for Free	0.8419		0.2859		0.7255		0.7660	
Bought from Friend or Relative	0.1501		0.7898		0.4091		0.2326	
Took from Friend or Relative without Asking	0.0615		0.8350		0.3538		0.0557	
Bought from Drug Dealer or Other Stranger	0.5429		0.5913		0.3072		0.8472	
Bought on the Internet	0.0372		*		0.4304		*	
Some Other Way	0.6762		0.1306		0.7251		0.9190	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.52C Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Tranquilizers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	118	99	8	8	27	26	115	95
From More Than One Doctor	22	16	2	2	6	4	21	15
Wrote Fake Prescription	1	1	*	0	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	2	1	*	2	1	4	1
From Friend or Relative for Free	26	23	4	5	10	9	23	20
Bought from Friend or Relative	19	13	6	2	7	8	16	10
Took from Friend or Relative without Asking	7	4	2	2	4	2	6	2
Bought from Drug Dealer or Other Stranger	8	14	3	3	5	7	5	11
Bought on the Internet	1	8	*	*	*	4	*	6
Some Other Way	9	9	2	1	6	2	6	8

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.52D Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Tranquilizers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	1.37	1.37	4.22	4.63	1.98	2.12	1.73	1.68
From More Than One Doctor	0.76	0.59	1.38	1.84	0.91	0.65	1.03	0.78
Wrote Fake Prescription	0.05	0.05	*	0.44	0.20	0.17	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.16	0.06	0.38	*	0.23	0.19	0.20	0.06
From Friend or Relative for Free	0.89	0.84	2.81	3.84	1.38	1.26	1.13	1.02
Bought from Friend or Relative	0.64	0.49	3.80	1.91	0.99	1.18	0.78	0.54
Took from Friend or Relative without Asking	0.26	0.14	1.39	1.84	0.64	0.33	0.28	0.12
Bought from Drug Dealer or Other Stranger	0.27	0.50	2.07	2.24	0.69	1.11	0.25	0.59
Bought on the Internet	0.03	0.28	*	*	*	0.65	*	0.33
Some Other Way	0.31	0.32	1.03	1.06	0.81	0.35	0.30	0.42

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.52P Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source Where Friend or Relative Obtained Tranquilizers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.6389		0.1997		1.0000		0.8937	
From More Than One Doctor	0.3970		0.8683		0.1690		0.6325	
Wrote Fake Prescription	0.7697		*		0.9170		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1742		*		0.4679		0.3190	
From Friend or Relative for Free	0.8448		0.8301		0.7296		0.9817	
Bought from Friend or Relative	0.2546		0.0923		0.5481		0.2779	
Took from Friend or Relative without Asking	0.0247		0.5626		0.0753		0.2641	
Bought from Drug Dealer or Other Stranger	0.0099		1.0000		0.0142		0.0889	
Bought on the Internet	0.0961		*		*		*	
Some Other Way	0.9653		0.7528		0.1667		0.4035	

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.53C Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	52	37	4	*	9	7	51	36
From More Than One Doctor	5	7	1	1	4	4	2	5
Wrote Fake Prescription	1	2	1	*	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	13	5	*	*	3	3	*	4
From Friend or Relative for Free	81	61	8	*	19	17	77	57
Bought from Friend or Relative	36	30	5	*	13	12	34	26
Took from Friend or Relative without Asking	17	19	3	3	8	5	14	18
Bought from Drug Dealer or Other Stranger	21	19	5	*	10	8	18	17
Bought on the Internet	*	7	*	*	0	*	*	7
Some Other Way	15	11	2	2	7	3	13	11

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.53D Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	2.39	1.99	3.18	*	1.63	1.64	3.41	2.68
From More Than One Doctor	0.27	0.39	0.97	1.81	0.89	0.93	0.15	0.44
Wrote Fake Prescription	0.07	0.10	0.80	*	0.19	0.15	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.69	0.28	*	*	0.55	0.63	*	0.32
From Friend or Relative for Free	2.63	2.34	4.33	*	2.67	2.90	3.66	3.11
Bought from Friend or Relative	1.80	1.67	3.67	*	2.20	2.62	2.51	2.06
Took from Friend or Relative without Asking	0.88	1.11	2.15	3.91	1.57	1.25	1.11	1.49
Bought from Drug Dealer or Other Stranger	1.14	1.11	3.57	*	1.91	1.84	1.42	1.38
Bought on the Internet	*	0.40	*	*	0.02	*	*	0.56
Some Other Way	0.80	0.68	2.00	2.23	1.33	0.71	1.04	0.90

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.53P Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.9238		*		0.5606		0.7573	
From More Than One Doctor	0.3270		0.2213		0.9372		0.2312	
Wrote Fake Prescription	0.8120		*		0.6967		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3770		*		0.9830		*	
From Friend or Relative for Free	0.7724		*		0.5597		0.5922	
Bought from Friend or Relative	0.4908		*		0.3554		0.3614	
Took from Friend or Relative without Asking	0.0551		0.3910		0.8024		0.0293	
Bought from Drug Dealer or Other Stranger	0.8570		*		0.1298		0.4278	
Bought on the Internet	*		*		*		*	
Some Other Way	0.6915		0.5626		0.3380		0.9346	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.54C Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Tranquilizers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	69	53	*	*	14	14	67	50
From More Than One Doctor	20	12	*	*	3	3	19	11
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	6	*	*	2	*	*	6
From Friend or Relative for Free	17	18	*	*	6	7	16	17
Bought from Friend or Relative	11	16	*	*	5	8	9	13
Took from Friend or Relative without Asking	7	5	*	*	3	1	6	5
Bought from Drug Dealer or Other Stranger	5	14	*	*	3	4	3	14
Bought on the Internet	1	9	*	*	*	*	*	8
Some Other Way	8	8	*	*	*	2	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.54D Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Tranquilizers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	2.72	2.45	*	*	3.63	4.01	3.56	2.92
From More Than One Doctor	1.85	1.28	*	*	1.44	1.68	2.52	1.61
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.41	0.67	*	*	0.79	*	*	0.88
From Friend or Relative for Free	1.72	1.96	*	*	2.63	3.22	2.17	2.40
Bought from Friend or Relative	1.11	1.68	*	*	2.08	3.86	1.18	1.89
Took from Friend or Relative without Asking	0.68	0.55	*	*	1.40	0.50	0.80	0.75
Bought from Drug Dealer or Other Stranger	0.51	1.55	*	*	1.47	1.97	0.43	1.99
Bought on the Internet	0.09	1.02	*	*	*	*	*	1.21
Some Other Way	0.80	0.85	*	*	*	1.01	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.54P Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source Where Friend or Relative Obtained Tranquilizers	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.4824		*		0.8753		0.5143	
From More Than One Doctor	0.2773		*		0.8652		0.2234	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8190		*		*		*	
From Friend or Relative for Free	0.3645		*		0.7049		0.3590	
Bought from Friend or Relative	0.0453		*		0.1978		0.0201	
Took from Friend or Relative without Asking	0.1848		*		0.0795		0.6871	
Bought from Drug Dealer or Other Stranger	0.0092		*		0.9275		0.0036	
Bought on the Internet	0.0812		*		*		*	
Some Other Way	0.8866		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.55C Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Stimulant Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	35	33	4	6	10	9	33	31
From More Than One Doctor	8	6	1	2	7	6	4	3
Wrote Fake Prescription	2	4	*	1	1	2	*	3
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	1	1	*	2	1	3	1
From Friend or Relative for Free	65	75	10	11	29	31	59	68
Bought from Friend or Relative	39	38	6	7	26	23	28	30
Took from Friend or Relative without Asking	15	20	4	5	5	8	14	18
Bought from Drug Dealer or Other Stranger	17	17	4	4	11	8	13	14
Bought on the Internet	5	10	*	1	1	5	5	9
Some Other Way	12	13	1	3	7	8	10	9

*Low precision; no estimate reported.

NOTE: Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.55D Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Stimulant Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	1.47	1.18	1.95	2.52	0.88	0.79	3.10	2.23
From More Than One Doctor	0.38	0.25	0.67	0.84	0.67	0.50	0.45	0.22
Wrote Fake Prescription	0.08	0.14	*	0.39	0.08	0.21	*	0.20
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.18	0.05	0.55	*	0.21	0.09	0.32	0.07
From Friend or Relative for Free	1.99	1.83	3.07	3.61	2.09	1.88	3.89	3.21
Bought from Friend or Relative	1.55	1.33	2.51	2.88	2.00	1.73	2.66	2.14
Took from Friend or Relative without Asking	0.67	0.75	1.65	2.16	0.45	0.67	1.45	1.32
Bought from Drug Dealer or Other Stranger	0.78	0.64	1.66	1.72	1.05	0.73	1.32	1.09
Bought on the Internet	0.24	0.37	*	0.26	0.14	0.41	0.52	0.65
Some Other Way	0.55	0.47	0.58	1.15	0.68	0.73	1.01	0.70

*Low precision; no estimate reported.

NOTE: Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

**Table 6.55P Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.9789		0.1615		0.8466		0.7166	
From More Than One Doctor	0.3049		0.3940		0.3157		0.6007	
Wrote Fake Prescription	0.3488		*		0.3862		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.0733		*		0.1677		0.4863	
From Friend or Relative for Free	0.9101		0.1049		0.8569		0.8629	
Bought from Friend or Relative	0.9763		0.8700		0.7772		0.8884	
Took from Friend or Relative without Asking	0.0927		0.6096		0.0506		0.3536	
Bought from Drug Dealer or Other Stranger	0.1074		0.6409		0.0320		0.7890	
Bought on the Internet	0.1422		*		0.0150		0.7248	
Some Other Way	0.4994		0.0720		0.9427		0.2765	

*Low precision; no estimate reported.

NOTE: Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.56C Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Stimulants	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	60	59	7	9	24	25	54	53
From More Than One Doctor	7	13	3	2	6	5	3	12
Wrote Fake Prescription	*	0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	0	*	0	2	0	*	*
From Friend or Relative for Free	14	19	3	*	7	7	12	17
Bought from Friend or Relative	10	15	3	1	7	7	6	13
Took from Friend or Relative without Asking	4	3	3	1	2	1	*	3
Bought from Drug Dealer or Other Stranger	5	13	2	1	5	5	2	12
Bought on the Internet	2	2	1	*	0	1	*	*
Some Other Way	2	3	1	1	1	1	0	2

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.56D Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Stimulants	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	1.94	2.20	4.59	5.62	2.34	2.12	2.75	3.70
From More Than One Doctor	0.71	1.06	2.70	1.93	1.22	0.95	0.59	1.84
Wrote Fake Prescription	*	0.03	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.15	0.03	*	0.05	0.33	0.08	*	*
From Friend or Relative for Free	1.31	1.50	2.83	*	1.44	1.33	2.29	2.48
Bought from Friend or Relative	0.94	1.16	2.78	0.80	1.42	1.34	1.26	1.92
Took from Friend or Relative without Asking	0.34	0.28	2.72	1.15	0.53	0.28	*	0.45
Bought from Drug Dealer or Other Stranger	0.52	1.00	1.81	1.09	1.00	0.94	0.41	1.73
Bought on the Internet	0.21	0.13	0.68	*	0.07	0.23	*	*
Some Other Way	0.15	0.21	0.91	1.12	0.29	0.29	0.03	0.28

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.56P Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source Where Friend or Relative Obtained Stimulants	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.1534		0.2027		0.1274		0.0014	
From More Than One Doctor	0.8610		0.2071		0.2441		0.1414	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3099		*		0.3112		*	
From Friend or Relative for Free	0.1468		*		0.7771		0.1068	
Bought from Friend or Relative	0.9684		0.1024		0.5349		0.3573	
Took from Friend or Relative without Asking	0.2685		0.2403		0.0741		*	
Bought from Drug Dealer or Other Stranger	0.1415		0.1088		0.9959		0.0285	
Bought on the Internet	0.6206		*		0.3525		*	
Some Other Way	0.6334		0.9091		0.9198		0.3770	

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.57C Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Stimulant Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	29	25	3	5	5	6	*	24
From More Than One Doctor	9	5	2	1	2	5	*	*
Wrote Fake Prescription	*	1	*	0	0	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	9	1	1	0	2	1	*	1
From Friend or Relative for Free	32	47	6	7	17	18	*	42
Bought from Friend or Relative	27	34	4	5	14	13	*	31
Took from Friend or Relative without Asking	12	11	2	2	5	5	*	10
Bought from Drug Dealer or Other Stranger	17	12	3	3	8	5	16	10
Bought on the Internet	10	7	*	*	1	3	*	6
Some Other Way	11	9	1	1	5	5	10	8

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.57D Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Stimulant Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	3.33	2.27	3.86	5.06	1.43	1.66	*	3.86
From More Than One Doctor	1.23	0.49	2.11	1.39	0.63	1.35	*	*
Wrote Fake Prescription	*	0.09	*	0.32	0.03	0.26	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.14	0.14	0.97	0.32	0.54	0.29	*	0.15
From Friend or Relative for Free	3.34	3.03	5.26	5.76	3.13	3.25	*	4.71
Bought from Friend or Relative	3.08	2.82	4.68	5.13	3.19	3.04	*	4.62
Took from Friend or Relative without Asking	1.54	1.09	2.86	2.79	1.47	1.28	*	1.69
Bought from Drug Dealer or Other Stranger	2.19	1.18	3.96	3.92	2.33	1.48	4.03	1.77
Bought on the Internet	1.30	0.72	*	*	0.43	0.96	*	1.13
Some Other Way	1.46	0.95	1.45	1.17	1.63	1.47	2.59	1.40

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.57P Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source for Past Month Nonmedical Use among Past Month Stimulant Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.4536		0.3212		0.4922		*	
From More Than One Doctor	0.2742		0.6460		0.5272		*	
Wrote Fake Prescription	*		*		0.3059		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2605		0.3380		0.5315		*	
From Friend or Relative for Free	0.1122		0.4181		0.9279		*	
Bought from Friend or Relative	0.6596		0.2454		0.8663		*	
Took from Friend or Relative without Asking	0.7276		0.7227		0.7562		*	
Bought from Drug Dealer or Other Stranger	0.0066		0.8227		0.0262		0.0783	
Bought on the Internet	0.6644		*		0.1882		*	
Some Other Way	0.2273		0.5159		0.8700		0.1607	

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.58C Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Stimulants	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	29	34	*	*	16	16	*	*
From More Than One Doctor	9	11	*	*	2	4	*	*
Wrote Fake Prescription	*	*	*	*	*	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0	*	*	*	*	*	*
From Friend or Relative for Free	10	15	*	*	3	4	*	15
Bought from Friend or Relative	10	15	*	*	3	4	*	*
Took from Friend or Relative without Asking	9	5	*	*	2	*	*	4
Bought from Drug Dealer or Other Stranger	11	14	*	*	3	2	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	2	3	*	*	1	2	*	*

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.58D Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Stimulants	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	2.93	4.04	*	*	2.76	2.90	*	*
From More Than One Doctor	2.69	2.16	*	*	1.16	1.96	*	*
Wrote Fake Prescription	*	*	*	*	*	0.14	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0.01	*	*	*	*	*	*
From Friend or Relative for Free	2.82	2.84	*	*	2.26	2.33	*	4.55
Bought from Friend or Relative	2.82	2.76	*	*	1.84	2.04	*	*
Took from Friend or Relative without Asking	2.63	0.92	*	*	1.60	*	*	1.45
Bought from Drug Dealer or Other Stranger	3.26	2.53	*	*	1.81	1.27	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	0.49	0.67	*	*	0.46	1.22	*	*

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.58P Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source Where Friend or Relative Obtained Stimulants	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.0149		*		0.7494		*	
From More Than One Doctor	0.7405		*		0.5665		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	0.2678		*		0.8825		*	
Bought from Friend or Relative	0.8969		*		0.8861		*	
Took from Friend or Relative without Asking	0.2313		*		*		*	
Bought from Drug Dealer or Other Stranger	0.5258		*		0.4471		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.6844		*		0.3409		*	

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.59C Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	43	39	6	*	16	14	40	35
Bought from Friend or Relative	41	30	*	*	9	10	40	29
Took from Friend or Relative without Asking	8	19	1	*	6	4	6	*
Bought from Drug Dealer or Other Stranger	37	41	*	*	11	10	35	39
Bought on the Internet	2	4	*	*	2	1	*	4
Some Other Way	23	16	*	*	4	4	*	16

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.59D Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	3.86	2.87	5.79	*	4.84	3.73	5.50	3.83
Bought from Friend or Relative	3.61	2.52	*	*	2.97	3.03	5.41	3.56
Took from Friend or Relative without Asking	0.91	1.68	0.95	*	2.33	1.31	0.98	*
Bought from Drug Dealer or Other Stranger	3.42	2.98	*	*	3.31	3.03	4.99	4.10
Bought on the Internet	0.21	0.36	*	*	0.67	0.53	*	0.49
Some Other Way	2.34	1.48	*	*	1.43	1.33	*	2.10

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.59P Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	0.3948		*		0.4573		0.6268	
Bought from Friend or Relative	0.4006		*		0.2558		0.2169	
Took from Friend or Relative without Asking	0.4197		*		0.8076		*	
Bought from Drug Dealer or Other Stranger	0.1002		*		0.9849		0.0688	
Bought on the Internet	0.7413		*		0.6569		*	
Some Other Way	0.6167		*		0.6356		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.60C Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Methamphetamine	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	23	15	*	*	8	8	*	*
Bought from Friend or Relative	21	22	*	*	*	7	*	*
Took from Friend or Relative without Asking	2	4	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	10	16	*	*	6	7	*	*
Bought on the Internet	1	*	*	*	1	*	*	*
Some Other Way	13	8	*	*	4	*	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.60D Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Methamphetamine	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	5.37	3.89	*	*	5.16	5.64	*	*
Bought from Friend or Relative	5.27	4.81	*	*	*	5.42	*	*
Took from Friend or Relative without Asking	0.56	1.11	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	3.07	4.03	*	*	4.71	5.08	*	*
Bought on the Internet	0.40	*	*	*	1.11	*	*	*
Some Other Way	3.73	2.27	*	*	3.04	*	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.60P Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source Where Friend or Relative Obtained Methamphetamine	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	0.1829		*		0.8159		*	
Bought from Friend or Relative	0.6743		*		*		*	
Took from Friend or Relative without Asking	0.5362		*		*		*	
Bought from Drug Dealer or Other Stranger	0.0349		*		0.3805		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.2673		*		*		*	

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.61C Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	*	32	*	*	*	8	*	30
Bought from Friend or Relative	*	28	*	*	*	*	*	*
Took from Friend or Relative without Asking	5	*	*	*	*	3	*	*
Bought from Drug Dealer or Other Stranger	18	27	*	*	*	6	*	26
Bought on the Internet	2	*	*	*	*	1	*	*
Some Other Way	*	*	*	*	*	2	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.61D Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	*	4.87	*	*	*	6.05	*	5.87
Bought from Friend or Relative	*	4.85	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.26	*	*	*	*	2.94	*	*
Bought from Drug Dealer or Other Stranger	4.65	4.48	*	*	*	5.64	*	5.61
Bought on the Internet	0.47	*	*	*	*	0.60	*	*
Some Other Way	*	*	*	*	*	2.41	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.61P Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	0.1572		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.62C Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Methamphetamine	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	2	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.62D Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Methamphetamine	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.02	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.62P Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source Where Friend or Relative Obtained Methamphetamine	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

**Table 6.63C Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014**

Source for Most Recent Nonmedical Use among Past Year Sedative Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	27	23	3	*	4	6	*	22
From More Than One Doctor	4	5	*	*	2	*	*	3
Wrote Fake Prescription	1	*	*	*	0	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	*	2	*	1	*	*	*
From Friend or Relative for Free	31	38	5	*	9	11	*	36
Bought from Friend or Relative	10	16	2	*	4	4	*	15
Took from Friend or Relative without Asking	5	8	2	*	*	*	*	6
Bought from Drug Dealer or Other Stranger	4	7	1	*	2	1	*	7
Bought on the Internet	0	1	0	*	0	*	*	*
Some Other Way	10	8	0	*	2	0	*	8

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.63D Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Most Recent Nonmedical Use among Past Year Sedative Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	4.36	3.35	4.71	*	3.62	4.18	*	4.64
From More Than One Doctor	0.81	0.73	*	*	1.38	*	*	0.67
Wrote Fake Prescription	0.19	*	*	*	0.23	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.45	*	2.66	*	1.00	*	*	*
From Friend or Relative for Free	4.21	4.31	5.71	*	5.30	5.23	*	5.89
Bought from Friend or Relative	2.01	2.44	3.36	*	3.59	3.21	*	3.35
Took from Friend or Relative without Asking	0.92	1.23	2.84	*	*	*	*	1.43
Bought from Drug Dealer or Other Stranger	0.72	1.16	1.56	*	2.05	1.12	*	1.59
Bought on the Internet	0.03	0.14	0.06	*	0.15	*	*	*
Some Other Way	1.90	1.28	0.59	*	2.07	0.12	*	1.84

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

**Table 6.63P Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012**

Source for Most Recent Nonmedical Use among Past Year Sedative Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.4843		*		0.2634		*	
From More Than One Doctor	0.9664		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	0.7199		*		0.4521		*	
Bought from Friend or Relative	0.7044		*		0.6980		*	
Took from Friend or Relative without Asking	0.7087		*		*		*	
Bought from Drug Dealer or Other Stranger	0.6604		*		0.0708		*	
Bought on the Internet	0.5013		*		*		*	
Some Other Way	0.7915		*		0.0687		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.64C Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Sedatives	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	29	35	*	*	*	*	*	*
From More Than One Doctor	3	3	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	3	*	*	*	*	*	*
From Friend or Relative for Free	4	11	*	*	*	*	*	*
Bought from Friend or Relative	1	1	*	*	*	*	*	*
Took from Friend or Relative without Asking	3	4	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	3	3	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	1	2	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.64D Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Sedatives	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	2.72	4.07	*	*	*	*	*	*
From More Than One Doctor	1.27	0.84	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0.89	*	*	*	*	*	*
From Friend or Relative for Free	1.71	3.33	*	*	*	*	*	*
Bought from Friend or Relative	0.56	0.24	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.12	1.27	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	1.05	1.04	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	0.31	0.53	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.64P Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source Where Friend or Relative Obtained Sedatives	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	0.1299		*		*		*	
From More Than One Doctor	0.7153		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	0.1632		*		*		*	
Bought from Friend or Relative	0.3098		*		*		*	
Took from Friend or Relative without Asking	0.8111		*		*		*	
Bought from Drug Dealer or Other Stranger	0.8862		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.5575		*		*		*	

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.65C Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Sedative Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	4	*	*	*	*	*	*
Wrote Fake Prescription	1	0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	4	4	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	4	*	*	*	*	*	*
Bought on the Internet	0	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.65D Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source for Past Month Nonmedical Use among Past Month Sedative Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	1.48	*	*	*	*	*	*
Wrote Fake Prescription	0.43	0.17	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.61	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.96	1.57	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	1.63	*	*	*	*	*	*
Bought on the Internet	0.02	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.65P Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source for Past Month Nonmedical Use among Past Month Sedative Users	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	*		*		*		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	0.4492		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	0.5143		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.66C Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Sedatives	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.66D Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Source Where Friend or Relative Obtained Sedatives	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.66P Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Source Where Friend or Relative Obtained Sedatives	Total (2011-2012)	Total (2013-2014)	Aged 12-17 (2011-2012)	Aged 12-17 (2013-2014)	Aged 18-25 (2011-2012)	Aged 18-25 (2013-2014)	Aged 26+ (2011-2012)	Aged 26+ (2013-2014)
From One Doctor	*		*		*		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.67C Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Standard Errors of Numbers in Thousands, 2013 and 2014

Social Context and Location of Most Recent Alcohol Use	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	194	211	124	138	122	140
SOCIAL CONTEXT OF MOST RECENT ALCOHOL USE²						
Drank Alone	34	43	28	34	20	25
Drank with One Other Person	62	70	46	51	42	44
Drank with Two or More Other People	167	190	108	115	106	128
LOCATION OF MOST RECENT ALCOHOL USE^{2,3}						
Car or Other Vehicle	25	31	18	20	17	23
Home	98	104	68	75	64	65
Someone Else's Home	128	142	82	93	85	95
Park, Beach, or Parking Lot	28	30	19	21	21	21
Restaurant, Bar, or Club	42	47	21	31	34	35
Concert or Sports Game	19	25	14	20	13	15
School	22	39	14	17	13	30
Some Other Place ⁴	34	39	25	33	23	20
Party, Wedding, or Celebration	16	24	13	19	10	13
Outside (Location not Specified)	11	13	7	12	9	5
Hotel, Motel, or Resort	14	12	7	10	12	6
Camping, Hunting, or Fishing	8	9	8	9	3	5
Cabin, Cottage, Vacation Home, etc.	6	4	4	4	4	2
Dorm Room	5	6	2	6	5	2

*Low precision; no estimate reported.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents could indicate multiple locations for the most recent time they used alcohol; thus, these response categories are not mutually exclusive.

⁴ Some Other Place includes only valid responses from the other-specify questions, including these six types of commonly reported locations.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.67D Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender: Standard Errors of Percentages, 2013 and 2014

Social Context and Location of Most Recent Alcohol Use	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	0.00	0.00	0.00	0.00	0.00	0.00
SOCIAL CONTEXT OF MOST RECENT ALCOHOL USE²						
Drank Alone	0.39	0.51	0.64	0.76	0.49	0.60
Drank with One Other Person	0.66	0.78	0.98	1.06	0.93	0.99
Drank with Two or More Other People	0.72	0.91	1.07	1.22	1.00	1.12
LOCATION OF MOST RECENT ALCOHOL USE^{2,3}						
Car or Other Vehicle	0.29	0.37	0.40	0.47	0.41	0.57
Home	0.89	1.03	1.24	1.41	1.19	1.31
Someone Else's Home	0.90	1.08	1.24	1.48	1.27	1.38
Park, Beach, or Parking Lot	0.33	0.36	0.43	0.48	0.52	0.52
Restaurant, Bar, or Club	0.47	0.54	0.48	0.72	0.78	0.80
Concert or Sports Game	0.23	0.29	0.34	0.46	0.31	0.37
School	0.26	0.45	0.33	0.40	0.30	0.69
Some Other Place ⁴	0.40	0.47	0.58	0.76	0.55	0.48
Party, Wedding, or Celebration	0.19	0.29	0.30	0.46	0.26	0.33
Outside (Location not Specified)	0.13	0.16	0.16	0.29	0.21	0.13
Hotel, Motel, or Resort	0.17	0.14	0.17	0.24	0.29	0.15
Camping, Hunting, or Fishing	0.10	0.11	0.18	0.22	0.07	0.12
Cabin, Cottage, Vacation Home, etc.	0.07	0.05	0.09	0.09	0.11	0.06
Dorm Room	0.07	0.07	0.04	0.13	0.13	0.05

*Low precision; no estimate reported.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents could indicate multiple locations for the most recent time they used alcohol; thus, these response categories are not mutually exclusive.

⁴ Some Other Place includes only valid responses from the other-specify questions, including these six types of commonly reported locations.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.67P Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Social Context and Location of Most Recent Alcohol Use	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	N/A		N/A		N/A	
SOCIAL CONTEXT OF MOST RECENT ALCOHOL USE²						
Drank Alone	0.7495		0.3336		0.0587	
Drank with One Other Person	0.5923		0.2641		0.6814	
Drank with Two or More Other People	0.5168		0.6879		0.5468	
LOCATION OF MOST RECENT ALCOHOL USE^{2,3}						
Car or Other Vehicle	0.8771		0.5030		0.4112	
Home	0.5261		0.9759		0.3084	
Someone Else's Home	0.3909		0.7439		0.3441	
Park, Beach, or Parking Lot	0.2635		0.1612		0.8080	
Restaurant, Bar, or Club	0.3490		0.0821		0.8278	
Concert or Sports Game	0.4327		0.3306		0.9579	
School	0.2511		0.8272		0.1480	
Some Other Place ⁴	0.3782		0.0149		0.0733	
Party, Wedding, or Celebration	0.0847		0.0761		0.5955	
Outside (Location not Specified)	0.4752		0.0637		0.1707	
Hotel, Motel, or Resort	0.4779		0.4151		0.0839	
Camping, Hunting, or Fishing	0.4726		0.7108		0.4365	
Cabin, Cottage, Vacation Home, etc.	0.4754		0.8221		0.4251	
Dorm Room	0.8495		0.1553		0.2194	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents could indicate multiple locations for the most recent time they used alcohol; thus, these response categories are not mutually exclusive.

⁴ Some Other Place includes only valid responses from the other-specify questions, including these six types of commonly reported locations.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.68C Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Social Context and Location of Most Recent Alcohol Use	Total (2013)	Total (2014)	Aged 12-14 (2013)	Aged 12-14 (2014)	Aged 15-17 (2013)	Aged 15-17 (2014)	Aged 18-20 (2013)	Aged 18-20 (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	194	211	31	30	73	81	168	186
SOCIAL CONTEXT OF MOST RECENT ALCOHOL USE²								
Drank Alone	34	43	11	9	17	21	27	36
Drank with One Other Person	62	70	12	13	27	30	53	61
Drank with Two or More Other People	167	190	25	25	61	71	147	166
LOCATION OF MOST RECENT ALCOHOL USE^{2,3}								
Car or Other Vehicle	25	31	6	4	14	15	19	26
Home	98	104	18	19	37	43	86	91
Someone Else's Home	128	142	20	21	53	61	109	120
Park, Beach, or Parking Lot	28	30	7	4	16	15	23	27
Restaurant, Bar, or Club	42	47	3	3	9	14	41	45
Concert or Sports Game	19	25	6	3	8	11	17	22
School	22	39	4	4	5	11	21	37
Some Other Place ⁴	34	39	5	4	18	22	28	30
Party, Wedding, or Celebration	16	24	3	2	11	16	11	17
Outside (Location not Specified)	11	13	2	2	9	7	5	10
Hotel, Motel, or Resort	14	12	*	*	5	5	13	11
Camping, Hunting, or Fishing	8	9	1	1	4	4	7	8
Cabin, Cottage, Vacation Home, etc.	6	4	0	1	3	4	5	0
Dorm Room	5	6	*	*	*	4	5	5

*Low precision; no estimate reported.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents could indicate multiple locations for the most recent time they used alcohol; thus, these response categories are not mutually exclusive.

⁴ Some Other Place includes only valid responses from the other-specify questions, including these six types of commonly reported locations.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.68D Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Standard Errors of Percentages, 2013 and 2014

Social Context and Location of Most Recent Alcohol Use	Total (2013)	Total (2014)	Aged 12-14 (2013)	Aged 12-14 (2014)	Aged 15-17 (2013)	Aged 15-17 (2014)	Aged 18-20 (2013)	Aged 18-20 (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
SOCIAL CONTEXT OF MOST RECENT ALCOHOL USE²								
Drank Alone	0.39	0.51	2.30	2.38	0.72	0.84	0.46	0.62
Drank with One Other Person	0.66	0.78	2.58	3.46	1.06	1.17	0.86	1.02
Drank with Two or More Other People	0.72	0.91	3.04	3.80	1.22	1.36	0.92	1.17
LOCATION OF MOST RECENT ALCOHOL USE^{2,3}								
Car or Other Vehicle	0.29	0.37	1.48	1.06	0.61	0.63	0.33	0.46
Home	0.89	1.03	3.30	4.19	1.36	1.52	1.13	1.33
Someone Else's Home	0.90	1.08	3.42	4.18	1.45	1.65	1.13	1.35
Park, Beach, or Parking Lot	0.33	0.36	1.56	1.20	0.69	0.63	0.41	0.46
Restaurant, Bar, or Club	0.47	0.54	0.63	0.84	0.37	0.58	0.65	0.75
Concert or Sports Game	0.23	0.29	1.38	0.94	0.33	0.48	0.30	0.38
School	0.26	0.45	0.94	1.09	0.24	0.45	0.36	0.62
Some Other Place ⁴	0.40	0.47	1.20	1.09	0.77	0.93	0.50	0.54
Party, Wedding, or Celebration	0.19	0.29	0.73	0.66	0.49	0.69	0.20	0.31
Outside (Location not Specified)	0.13	0.16	0.40	0.56	0.42	0.33	0.10	0.19
Hotel, Motel, or Resort	0.17	0.14	*	*	0.24	0.20	0.23	0.19
Camping, Hunting, or Fishing	0.10	0.11	0.37	0.25	0.17	0.16	0.13	0.15
Cabin, Cottage, Vacation Home, etc.	0.07	0.05	0.06	0.31	0.11	0.18	0.09	0.01
Dorm Room	0.07	0.07	*	*	*	0.16	0.10	0.09

*Low precision; no estimate reported.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents could indicate multiple locations for the most recent time they used alcohol; thus, these response categories are not mutually exclusive.

⁴ Some Other Place includes only valid responses from the other-specify questions, including these six types of commonly reported locations.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.68P Social Context and Location of Most Recent Alcohol Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Social Context and Location of Most Recent Alcohol Use	Total (2013)	Total (2014)	Aged 12-14 (2013)	Aged 12-14 (2014)	Aged 15-17 (2013)	Aged 15-17 (2014)	Aged 18-20 (2013)	Aged 18-20 (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	N/A		N/A		N/A		N/A	
SOCIAL CONTEXT OF MOST RECENT ALCOHOL USE²								
Drank Alone	0.7495		0.2877		0.6276		0.2734	
Drank with One Other Person	0.5923		0.2458		0.8505		0.6080	
Drank with Two or More Other People	0.5168		0.7588		0.6474		0.3049	
LOCATION OF MOST RECENT ALCOHOL USE^{2,3}								
Car or Other Vehicle	0.8771		0.0605		0.5403		0.3260	
Home	0.5261		0.4436		0.6097		0.3142	
Someone Else's Home	0.3909		0.1942		0.8166		0.4495	
Park, Beach, or Parking Lot	0.2635		0.2487		0.0430		0.8495	
Restaurant, Bar, or Club	0.3490		0.9255		0.0499		0.6926	
Concert or Sports Game	0.4327		0.3715		0.5792		0.4080	
School	0.2511		0.5502		0.3193		0.3883	
Some Other Place ⁴	0.3782		0.1288		0.5193		0.4054	
Party, Wedding, or Celebration	0.0847		0.4504		0.2519		0.1537	
Outside (Location not Specified)	0.4752		0.8077		0.4112		0.0535	
Hotel, Motel, or Resort	0.4779		*		0.8054		0.4871	
Camping, Hunting, or Fishing	0.4726		0.4839		0.3961		0.1816	
Cabin, Cottage, Vacation Home, etc.	0.4754		0.4390		0.6540		0.1035	
Dorm Room	0.8495		*		*		0.6261	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents could indicate multiple locations for the most recent time they used alcohol; thus, these response categories are not mutually exclusive.

⁴ Some Other Place includes only valid responses from the other-specify questions, including these six types of commonly reported locations.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.69C Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Source of Most Recent Alcohol Use in the Past Month	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	194	211	124	138	122	140
UNDERAGE DRINKER PAID²	92	97	70	74	51	61
Purchased It Himself or Herself	45	48	38	37	23	30
From Store, Restaurant, Bar, Club, or Event	41	45	34	34	22	29
Liquor, Convenience, or Grocery Store	33	36	29	29	16	21
Restaurant, Bar, or Club	20	26	14	17	14	19
Concert, Sports, or Other Event	5	5	5	4	1	3
From Another Person	17	19	16	16	6	12
From Person under Age 21	9	10	8	6	3	8
From Person Aged 21 or Older	15	16	14	14	4	8
Purchased by Someone Else	73	79	54	61	44	51
Parent or Guardian	14	17	10	12	10	12
Another Family Member Aged 21 or Older	20	23	14	18	14	14
Someone Not Related Aged 21 or Older	62	66	45	49	37	43
Someone under Age 21	19	23	15	19	11	13
UNDERAGE DRINKER DID NOT PAY²	143	163	89	105	101	113
Got It from Parent or Guardian	41	48	26	34	29	32
Got It from Another Family Member Aged 21 or Older	49	57	31	40	37	38
Got It from Someone Not Related Aged 21 or Older	88	96	54	57	62	73
Got It from Someone under Age 21	51	60	33	38	37	43
Took It from Own Home	35	35	28	26	20	20
Took It from Someone Else's Home	18	26	14	18	11	18
Got It Some Other Way	28	36	22	28	18	22
From Friend or Acquaintance, Unspecified Age and Method ³	17	23	13	16	11	15

*Low precision; no estimate reported.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents were permitted to specify other sources for most recent alcohol use in the past month. This source is one of the most commonly reported other sources for most recent alcohol use in the past month.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.69D Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender:
Standard Errors of Percentages, 2013 and 2014**

Source of Most Recent Alcohol Use in the Past Month	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	0.00	0.00	0.00	0.00	0.00	0.00
UNDERAGE DRINKER PAID²	0.80	0.88	1.18	1.35	1.02	1.17
Purchased It Himself or Herself	0.49	0.53	0.80	0.82	0.53	0.70
From Store, Restaurant, Bar, Club, or Event	0.45	0.50	0.72	0.76	0.52	0.66
Liquor, Convenience, or Grocery Store	0.38	0.41	0.62	0.65	0.39	0.50
Restaurant, Bar, or Club	0.23	0.30	0.31	0.40	0.34	0.45
Concert, Sports, or Other Event	0.06	0.06	0.12	0.10	0.03	0.07
From Another Person	0.21	0.23	0.38	0.36	0.15	0.28
From Person under Age 21	0.11	0.11	0.20	0.14	0.07	0.18
From Person Aged 21 or Older	0.17	0.19	0.32	0.33	0.11	0.19
Purchased by Someone Else	0.70	0.78	1.04	1.20	0.92	1.07
Parent or Guardian	0.16	0.20	0.22	0.28	0.24	0.28
Another Family Member Aged 21 or Older	0.24	0.27	0.34	0.41	0.35	0.33
Someone Not Related Aged 21 or Older	0.63	0.68	0.93	1.04	0.82	0.92
Someone under Age 21	0.22	0.27	0.36	0.44	0.27	0.31
UNDERAGE DRINKER DID NOT PAY²	0.80	0.88	1.18	1.35	1.02	1.17
Got It from Parent or Guardian	0.48	0.57	0.59	0.81	0.74	0.79
Got It from Another Family Member Aged 21 or Older	0.58	0.66	0.70	0.93	0.90	0.93
Got It from Someone Not Related Aged 21 or Older	0.85	0.97	1.12	1.26	1.16	1.39
Got It from Someone under Age 21	0.59	0.66	0.78	0.87	0.87	0.96
Took It from Own Home	0.42	0.43	0.66	0.63	0.51	0.49
Took It from Someone Else's Home	0.22	0.32	0.33	0.45	0.29	0.44
Got It Some Other Way	0.34	0.43	0.51	0.66	0.46	0.56
From Friend or Acquaintance, Unspecified Age and Method ³	0.21	0.28	0.31	0.40	0.29	0.37

*Low precision; no estimate reported.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents were permitted to specify other sources for most recent alcohol use in the past month. This source is one of the most commonly reported other sources for most recent alcohol use in the past month.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.69P Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Gender:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Source of Most Recent Alcohol Use in the Past Month	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	N/A		N/A		N/A	
UNDERAGE DRINKER PAID²	0.4700		0.0808		0.2703	
Purchased It Himself or Herself	0.4333		0.6427		0.0428	
From Store, Restaurant, Bar, Club, or Event	0.7380		0.4321		0.1037	
Liquor, Convenience, or Grocery Store	0.8037		0.2817		0.2177	
Restaurant, Bar, or Club	0.3722		0.6992		0.3922	
Concert, Sports, or Other Event	0.9313		0.6229		0.1966	
From Another Person	0.2253		0.4933		0.2070	
From Person under Age 21	0.8758		0.3247		0.3129	
From Person Aged 21 or Older	0.0998		0.1961		0.2363	
Purchased by Someone Else	0.2210		0.1392		0.9639	
Parent or Guardian	0.2574		0.5515		0.3084	
Another Family Member Aged 21 or Older	0.8947		0.7328		0.5573	
Someone Not Related Aged 21 or Older	0.1271		0.0630		0.9112	
Someone under Age 21	0.7301		0.8473		0.7309	
UNDERAGE DRINKER DID NOT PAY²	0.4700		0.0808		0.2703	
Got It from Parent or Guardian	0.3841		0.2517		0.9421	
Got It from Another Family Member Aged 21 or Older	0.3479		0.0417		0.5175	
Got It from Someone Not Related Aged 21 or Older	0.4255		0.4286		0.6216	
Got It from Someone under Age 21	0.5621		0.3820		0.9587	
Took It from Own Home	0.0160		0.1162		0.0419	
Took It from Someone Else's Home	0.1777		0.7209		0.1043	
Got It Some Other Way	0.2435		0.3643		0.4435	
From Friend or Acquaintance, Unspecified Age and Method ³	0.6755		0.9586		0.5481	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents were permitted to specify other sources for most recent alcohol use in the past month. This source is one of the most commonly reported other sources for most recent alcohol use in the past month.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.70C Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Standard Errors of Numbers in Thousands, 2013 and 2014

Source of Most Recent Alcohol Use in the Past Month	Total (2013)	Total (2014)	Aged 12-14 (2013)	Aged 12-14 (2014)	Aged 15-17 (2013)	Aged 15-17 (2014)	Aged 18-20 (2013)	Aged 18-20 (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	194	211	31	30	73	81	168	186
UNDERAGE DRINKER PAID²	92	97	5	8	32	30	85	92
Purchased It Himself or Herself	45	48	3	*	14	14	42	47
From Store, Restaurant, Bar, Club, or Event	41	45	2	*	10	12	39	44
Liquor, Convenience, or Grocery Store	33	36	2	*	10	11	32	34
Restaurant, Bar, or Club	20	26	1	*	3	5	20	25
Concert, Sports, or Other Event	5	5	*	*	1	1	5	5
From Another Person	17	19	3	*	10	7	14	18
From Person under Age 21	9	10	2	*	8	4	4	8
From Person Aged 21 or Older	15	16	1	*	6	6	13	15
Purchased by Someone Else	73	79	4	5	27	26	68	73
Parent or Guardian	14	17	*	*	4	4	13	16
Another Family Member Aged 21 or Older	20	23	1	1	8	7	19	22
Someone Not Related Aged 21 or Older	62	66	4	5	23	20	57	61
Someone under Age 21	19	23	1	*	11	13	16	19
UNDERAGE DRINKER DID NOT PAY²	143	163	28	28	61	74	121	134
Got It from Parent or Guardian	41	48	11	11	20	25	32	38
Got It from Another Family Member Aged 21 or Older	49	57	12	14	21	26	39	49
Got It from Someone Not Related Aged 21 or Older	88	96	11	9	30	34	81	86
Got It from Someone under Age 21	51	60	11	8	30	36	36	47
Took It from Own Home	35	35	12	7	21	20	25	24
Took It from Someone Else's Home	18	26	5	5	13	15	12	20
Got It Some Other Way	28	36	7	9	15	22	22	26
From Friend or Acquaintance, Unspecified Age and Method ³	17	23	4	6	10	16	14	13

*Low precision; no estimate reported.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents were permitted to specify other sources for most recent alcohol use in the past month. This source is one of the most commonly reported other sources for most recent alcohol use in the past month.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.70D Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: Standard Errors of Percentages, 2013 and 2014

Source of Most Recent Alcohol Use in the Past Month	Total (2013)	Total (2014)	Aged 12-14 (2013)	Aged 12-14 (2014)	Aged 15-17 (2013)	Aged 15-17 (2014)	Aged 18-20 (2013)	Aged 18-20 (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
UNDERAGE DRINKER PAID²	0.80	0.88	1.26	2.27	1.20	1.15	1.05	1.15
Purchased It Himself or Herself	0.49	0.53	0.75	*	0.59	0.57	0.67	0.75
From Store, Restaurant, Bar, Club, or Event	0.45	0.50	0.44	*	0.43	0.48	0.64	0.72
Liquor, Convenience, or Grocery Store	0.38	0.41	0.40	*	0.42	0.44	0.53	0.58
Restaurant, Bar, or Club	0.23	0.30	0.19	*	0.13	0.19	0.33	0.44
Concert, Sports, or Other Event	0.06	0.06	*	*	0.07	0.04	0.09	0.09
From Another Person	0.21	0.23	0.60	*	0.43	0.29	0.25	0.31
From Person under Age 21	0.11	0.11	0.49	*	0.34	0.16	0.07	0.14
From Person Aged 21 or Older	0.17	0.19	0.35	*	0.25	0.25	0.23	0.26
Purchased by Someone Else	0.70	0.78	1.01	1.52	1.10	1.03	0.94	1.02
Parent or Guardian	0.16	0.20	*	*	0.18	0.18	0.23	0.28
Another Family Member Aged 21 or Older	0.24	0.27	0.28	0.31	0.35	0.30	0.33	0.38
Someone Not Related Aged 21 or Older	0.63	0.68	0.92	1.39	0.94	0.84	0.85	0.91
Someone under Age 21	0.22	0.27	0.28	*	0.47	0.53	0.28	0.32
UNDERAGE DRINKER DID NOT PAY²	0.80	0.88	1.26	2.27	1.20	1.15	1.05	1.15
Got It from Parent or Guardian	0.48	0.57	2.52	3.16	0.85	1.04	0.56	0.67
Got It from Another Family Member Aged 21 or Older	0.58	0.66	2.78	3.93	0.93	1.07	0.70	0.84
Got It from Someone Not Related Aged 21 or Older	0.85	0.97	2.57	2.88	1.24	1.33	1.10	1.27
Got It from Someone under Age 21	0.59	0.66	2.65	2.46	1.24	1.37	0.62	0.80
Took It from Own Home	0.42	0.43	2.70	2.35	0.91	0.86	0.44	0.45
Took It from Someone Else's Home	0.22	0.32	1.33	1.67	0.58	0.63	0.22	0.36
Got It Some Other Way	0.34	0.43	1.81	2.60	0.67	0.91	0.39	0.47
From Friend or Acquaintance, Unspecified Age and Method ³	0.21	0.28	1.08	2.01	0.46	0.69	0.25	0.24

*Low precision; no estimate reported.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents were permitted to specify other sources for most recent alcohol use in the past month. This source is one of the most commonly reported other sources for most recent alcohol use in the past month.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.70P Source Where Alcohol Was Obtained for Most Recent Use in the Past Month among Past Month Alcohol Users Aged 12 to 20, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Source of Most Recent Alcohol Use in the Past Month	Total (2013)	Total (2014)	Aged 12-14 (2013)	Aged 12-14 (2014)	Aged 15-17 (2013)	Aged 15-17 (2014)	Aged 18-20 (2013)	Aged 18-20 (2014)
TOTAL PAST MONTH ALCOHOL USERS¹	N/A		N/A		N/A		N/A	
UNDERAGE DRINKER PAID²	0.4700		0.9190		0.0765		0.8641	
Purchased It Himself or Herself	0.4333		*		0.8809		0.3839	
From Store, Restaurant, Bar, Club, or Event	0.7380		*		0.9938		0.7965	
Liquor, Convenience, or Grocery Store	0.8037		*		0.8346		0.7524	
Restaurant, Bar, or Club	0.3722		*		0.5263		0.4157	
Concert, Sports, or Other Event	0.9313		*		0.7575		0.8733	
From Another Person	0.2253		*		0.7016		0.0715	
From Person under Age 21	0.8758		*		0.4464		0.4305	
From Person Aged 21 or Older	0.0998		*		0.6090		0.0940	
Purchased by Someone Else	0.2210		0.7214		0.0691		0.4856	
Parent or Guardian	0.2574		*		0.9871		0.2718	
Another Family Member Aged 21 or Older	0.8947		0.8891		0.1358		0.6931	
Someone Not Related Aged 21 or Older	0.1271		0.7925		0.0532		0.3170	
Someone under Age 21	0.7301		*		0.6339		0.9887	
UNDERAGE DRINKER DID NOT PAY²	0.4700		0.9190		0.0765		0.8641	
Got It from Parent or Guardian	0.3841		0.7605		0.0870		0.9617	
Got It from Another Family Member Aged 21 or Older	0.3479		0.2524		0.4226		0.6270	
Got It from Someone Not Related Aged 21 or Older	0.4255		0.8112		0.8904		0.3869	
Got It from Someone under Age 21	0.5621		0.2212		0.9707		0.3707	
Took It from Own Home	0.0160		0.0485		0.0725		0.2881	
Took It from Someone Else's Home	0.1777		0.4927		0.6806		0.1957	
Got It Some Other Way	0.2435		0.2461		0.1381		0.9138	
From Friend or Acquaintance, Unspecified Age and Method ³	0.6755		0.1652		0.1419		0.1676	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ The total row represents all respondents in the specified subgroup, including those with missing underage drinking information.

² Respondents with unknown responses were excluded.

³ Respondents were permitted to specify other sources for most recent alcohol use in the past month. This source is one of the most commonly reported other sources for most recent alcohol use in the past month.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.71C Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
ILLCIT DRUGS²	151	162	19	15	148	161
Marijuana and Hashish	122	139	19	13	119	138
Cocaine	30	42	2	3	30	42
Crack	12	16	*	2	12	16
Heroin	16	18	3	2	16	17
Hallucinogens	26	32	1	2	26	31
LSD	6	17	*	*	6	17
PCP	*	7	0	*	*	7
Ecstasy	23	27	1	2	23	27
Inhalants	16	18	1	4	16	18
Nonmedical Use of Psychotherapeutics ^{3,4}	82	92	10	7	82	92
Pain Relievers	64	73	6	7	64	72
OxyContin [®]	13	14	1	2	14	14
Tranquilizers	50	47	3	4	50	46
Stimulants ⁴	30	43	7	2	29	43
Methamphetamine ⁴	22	17	6	2	21	17
Sedatives	13	25	0	1	13	25
ILLCIT DRUGS OTHER THAN MARIJUANA²	90	101	10	9	89	100

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.71D Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
ILLCIT DRUGS²	0.22	0.24	0.86	0.65	0.22	0.25
Marijuana and Hashish	0.18	0.21	0.82	0.57	0.18	0.21
Cocaine	0.05	0.07	0.11	0.14	0.05	0.07
Crack	0.02	0.03	*	0.09	0.02	0.03
Heroin	0.03	0.03	0.14	0.10	0.03	0.03
Hallucinogens	0.04	0.05	0.05	0.10	0.04	0.05
LSD	0.01	0.03	*	*	0.01	0.03
PCP	*	0.01	0.01	*	*	0.01
Ecstasy	0.04	0.04	0.03	0.10	0.04	0.04
Inhalants	0.02	0.03	0.03	0.19	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{3,4}	0.13	0.14	0.43	0.30	0.13	0.15
Pain Relievers	0.10	0.11	0.28	0.28	0.10	0.12
OxyContin [®]	0.02	0.02	0.07	0.07	0.02	0.02
Tranquilizers	0.08	0.07	0.15	0.19	0.08	0.07
Stimulants ⁴	0.05	0.07	0.30	0.08	0.05	0.07
Methamphetamine ⁴	0.04	0.03	0.29	0.08	0.04	0.03
Sedatives	0.02	0.04	0.01	0.05	0.02	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.14	0.15	0.44	0.39	0.14	0.16

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.71P Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
ILLCIT DRUGS²	0.0000		0.5887		0.0000	
Marijuana and Hashish	0.0000		0.2858		0.0000	
Cocaine	0.4163		0.4749		0.4349	
Crack	0.2016		*		0.2062	
Heroin	0.2951		0.8905		0.3821	
Hallucinogens	0.4197		0.3923		0.4631	
LSD	0.0202		*		0.0191	
PCP	*		*		*	
Ecstasy	0.9674		0.2998		0.9225	
Inhalants	0.9433		0.0942		0.7103	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.5277		0.7009		0.4993	
Pain Relievers	0.5199		0.6154		0.5084	
OxyContin [®]	0.8642		0.7252		0.7429	
Tranquilizers	0.2218		0.4525		0.1634	
Stimulants ⁴	0.0023		0.3939		0.0019	
Methamphetamine ⁴	0.7471		0.6429		0.6498	
Sedatives	0.3836		0.4537		0.4165	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.1990		0.6040		0.2192	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.72C Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	151	162	19	15	148	161
AGE						
15-17	31	29	3	*	30	29
18-25	65	67	12	10	81	76
26-44	115	136	15	11	112	136
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	137	150	17	14	135	149
White	119	131	14	12	117	130
Black or African American	57	63	10	5	54	62
American Indian or Alaska Native	10	7	*	*	10	7
Native Hawaiian or Other Pacific Islander	9	12	*	*	9	12
Asian	29	21	*	*	29	21
Two or More Races	17	24	*	*	17	24
Hispanic or Latino	61	57	9	7	60	56
TRIMESTER²						
First	N/A	N/A	16	12	N/A	N/A
Second	N/A	N/A	9	9	N/A	N/A
Third	N/A	N/A	4	4	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.72D Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.22	0.24	0.86	0.65	0.22	0.25
AGE						
15-17	0.44	0.44	4.44	*	0.44	0.44
18-25	0.38	0.39	1.42	1.17	0.40	0.40
26-44	0.28	0.32	1.10	0.77	0.28	0.33
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.24	0.27	0.97	0.71	0.25	0.28
White	0.29	0.33	1.13	0.91	0.30	0.34
Black or African American	0.57	0.63	2.74	1.41	0.57	0.67
American Indian or Alaska Native	2.21	1.69	*	*	2.29	1.74
Native Hawaiian or Other Pacific Islander	3.49	3.84	*	*	3.68	4.03
Asian	0.73	0.51	*	*	0.76	0.52
Two or More Races	1.46	1.88	*	*	1.51	1.96
Hispanic or Latino	0.48	0.43	1.89	1.72	0.49	0.44
TRIMESTER²						
First	N/A	N/A	2.19	1.59	N/A	N/A
Second	N/A	N/A	1.15	1.04	N/A	N/A
Third	N/A	N/A	0.63	0.55	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.72P Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.0000		0.5887		0.0000	
AGE						
15-17	0.0366		*		0.0349	
18-25	0.2604		0.9337		0.2321	
26-44	0.0000		0.6100		0.0000	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0000		0.5133		0.0000	
White	0.0000		0.8521		0.0000	
Black or African American	0.1273		0.2096		0.0786	
American Indian or Alaska Native	0.4475		*		0.4089	
Native Hawaiian or Other Pacific Islander	0.7549		*		0.7202	
Asian	0.2819		*		0.2253	
Two or More Races	0.0620		*		0.0540	
Hispanic or Latino	0.8936		0.9069		0.8143	
TRIMESTER²						
First	N/A		0.3100		N/A	
Second	N/A		0.4014		N/A	
Third	N/A		0.6540		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.73C Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	122	139	19	13	119	138
AGE						
15-17	27	26	3	*	27	26
18-25	60	65	11	9	73	71
26-44	92	115	15	10	90	114
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	113	130	17	12	110	129
White	97	113	14	11	96	113
Black or African American	51	54	10	5	48	54
American Indian or Alaska Native	6	5	*	*	6	4
Native Hawaiian or Other Pacific Islander	7	11	*	*	7	11
Asian	22	16	*	*	22	16
Two or More Races	16	21	*	*	16	21
Hispanic or Latino	46	44	8	7	45	43
TRIMESTER²						
First	N/A	N/A	16	10	N/A	N/A
Second	N/A	N/A	8	9	N/A	N/A
Third	N/A	N/A	4	2	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.73D Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.18	0.21	0.82	0.57	0.18	0.21
AGE						
15-17	0.39	0.40	4.28	*	0.39	0.40
18-25	0.35	0.37	1.31	1.03	0.36	0.39
26-44	0.23	0.27	1.09	0.69	0.23	0.28
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.20	0.24	0.96	0.62	0.21	0.25
White	0.25	0.29	1.12	0.83	0.25	0.30
Black or African American	0.52	0.56	2.71	1.37	0.51	0.59
American Indian or Alaska Native	1.38	1.28	*	*	1.44	1.32
Native Hawaiian or Other Pacific Islander	2.88	3.71	*	*	3.04	3.90
Asian	0.57	0.40	*	*	0.59	0.41
Two or More Races	1.39	1.62	*	*	1.45	1.69
Hispanic or Latino	0.36	0.34	1.64	1.59	0.37	0.35
TRIMESTER²						
First	N/A	N/A	2.17	1.38	N/A	N/A
Second	N/A	N/A	1.00	1.01	N/A	N/A
Third	N/A	N/A	0.63	0.33	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.73P Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.0000		0.2858		0.0000	
AGE						
15-17	0.1181		*		0.1235	
18-25	0.1174		0.8214		0.0939	
26-44	0.0000		0.3502		0.0000	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0000		0.1669		0.0000	
White	0.0000		0.4526		0.0000	
Black or African American	0.0523		0.2487		0.0267	
American Indian or Alaska Native	0.2426		*		0.1959	
Native Hawaiian or Other Pacific Islander	0.6442		*		0.6090	
Asian	0.1512		*		0.1654	
Two or More Races	0.1625		*		0.1449	
Hispanic or Latino	0.4177		0.6612		0.3686	
TRIMESTER²						
First	N/A		0.0955		N/A	
Second	N/A		0.1401		N/A	
Third	N/A		0.0924		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.74C Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
ILLCIT DRUGS²	151	162	19	15	148	161
TOBACCO PRODUCTS³	271	241	32	26	265	239
Cigarettes	259	230	31	26	254	228
Smokeless Tobacco	28	28	6	4	28	28
Cigars	86	74	8	6	86	73
Pipe Tobacco	28	24	2	3	27	24
ALCOHOL	459	390	21	25	459	387
Binge Alcohol Use ⁴	259	241	10	11	258	240
Heavy Alcohol Use ⁴	118	105	3	5	118	105
ILLCIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL^{2,3}	493	413	39	35	490	410
ILLCIT DRUGS OR ALCOHOL²	469	392	28	27	467	389
TOBACCO PRODUCTS OR ALCOHOL³	487	411	37	35	485	408

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.74D Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
ILLCIT DRUGS²	0.22	0.24	0.86	0.65	0.22	0.25
TOBACCO PRODUCTS³	0.37	0.31	1.36	1.09	0.37	0.33
Cigarettes	0.36	0.30	1.34	1.08	0.36	0.31
Smokeless Tobacco	0.05	0.04	0.28	0.19	0.05	0.05
Cigars	0.13	0.11	0.38	0.26	0.14	0.12
Pipe Tobacco	0.04	0.04	0.10	0.11	0.05	0.04
ALCOHOL	0.41	0.38	0.93	1.02	0.42	0.39
Binge Alcohol Use ⁴	0.32	0.31	0.46	0.48	0.34	0.32
Heavy Alcohol Use ⁴	0.18	0.16	0.13	0.22	0.19	0.16
ILLCIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL^{2,3}	0.39	0.36	1.60	1.39	0.40	0.37
ILLCIT DRUGS OR ALCOHOL²	0.41	0.37	1.22	1.11	0.42	0.38
TOBACCO PRODUCTS OR ALCOHOL³	0.40	0.37	1.53	1.37	0.40	0.37

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.74P Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
ILLCIT DRUGS²	0.0000		0.5887		0.0000	
TOBACCO PRODUCTS³	0.0148		0.0788		0.0237	
Cigarettes	0.0025		0.0943		0.0040	
Smokeless Tobacco	0.6911		0.8939		0.6594	
Cigars	0.2788		0.2927		0.3057	
Pipe Tobacco	0.4217		0.8703		0.4872	
ALCOHOL	0.8052		0.5588		0.7550	
Binge Alcohol Use ⁴	0.6577		0.8550		0.6893	
Heavy Alcohol Use ⁴	0.3394		0.1367		0.3508	
ILLCIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL^{2,3}	0.8577		0.1914		0.6486	
ILLCIT DRUGS OR ALCOHOL²	0.5840		0.8267		0.4852	
TOBACCO PRODUCTS OR ALCOHOL³	0.9074		0.3887		0.7501	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.75C Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	259	230	31	26	254	228
AGE						
15-17	25	24	*	*	24	23
18-25	72	73	16	14	92	82
26-44	227	208	25	22	223	207
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	236	212	31	25	230	211
White	208	185	26	20	204	183
Black or African American	80	86	11	14	78	84
American Indian or Alaska Native	21	21	*	*	20	20
Native Hawaiian or Other Pacific Islander	*	10	*	*	*	10
Asian	39	34	*	*	39	34
Two or More Races	36	27	*	*	35	27
Hispanic or Latino	94	78	5	8	94	77
TRIMESTER²						
First	N/A	N/A	23	15	N/A	N/A
Second	N/A	N/A	16	14	N/A	N/A
Third	N/A	N/A	12	16	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.75D Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.36	0.30	1.34	1.08	0.36	0.31
AGE						
15-17	0.37	0.36	*	*	0.36	0.36
18-25	0.42	0.42	1.69	1.59	0.43	0.43
26-44	0.51	0.44	1.82	1.44	0.52	0.45
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.39	0.35	1.63	1.25	0.40	0.36
White	0.47	0.42	1.96	1.51	0.48	0.43
Black or African American	0.77	0.81	3.14	3.34	0.78	0.82
American Indian or Alaska Native	3.58	3.83	*	*	3.59	4.00
Native Hawaiian or Other Pacific Islander	*	3.48	*	*	*	3.64
Asian	0.97	0.80	*	*	1.00	0.84
Two or More Races	2.45	2.12	*	*	2.50	2.18
Hispanic or Latino	0.72	0.58	1.10	1.90	0.75	0.59
TRIMESTER²						
First	N/A	N/A	2.85	2.06	N/A	N/A
Second	N/A	N/A	1.97	1.59	N/A	N/A
Third	N/A	N/A	1.72	2.06	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.75P Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.0025		0.0943		0.0040	
AGE						
15-17	0.0000		*		0.0000	
18-25	0.0000		0.4571		0.0000	
26-44	0.7862		0.1986		0.9417	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0161		0.0339		0.0284	
White	0.0289		0.1684		0.0355	
Black or African American	0.1416		0.4949		0.1013	
American Indian or Alaska Native	0.7489		*		0.7923	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0948		*		0.1034	
Two or More Races	0.0515		*		0.0976	
Hispanic or Latino	0.0488		0.3665		0.0404	
TRIMESTER²						
First	N/A		0.0533		N/A	
Second	N/A		0.2726		N/A	
Third	N/A		0.6536		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.76C Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	459	390	21	25	459	387
AGE						
15-17	36	39	3	*	36	39
18-25	88	87	7	8	185	153
26-44	393	341	20	23	394	338
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	410	355	19	24	410	353
White	358	302	17	22	358	302
Black or African American	165	145	8	9	164	143
American Indian or Alaska Native	21	21	*	*	21	21
Native Hawaiian or Other Pacific Islander	*	18	*	*	*	18
Asian	95	90	*	*	95	86
Two or More Races	47	51	*	*	46	51
Hispanic or Latino	182	145	7	6	181	145
TRIMESTER²						
First	N/A	N/A	17	21	N/A	N/A
Second	N/A	N/A	8	10	N/A	N/A
Third	N/A	N/A	9	9	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.76D Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.41	0.38	0.93	1.02	0.42	0.39
AGE						
15-17	0.50	0.55	4.11	*	0.51	0.56
18-25	0.51	0.50	0.89	0.97	0.53	0.51
26-44	0.59	0.54	1.46	1.54	0.61	0.55
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.44	0.42	1.08	1.22	0.45	0.42
White	0.48	0.45	1.34	1.59	0.48	0.46
Black or African American	1.10	0.99	2.18	2.34	1.15	1.00
American Indian or Alaska Native	3.45	3.67	*	*	3.43	3.77
Native Hawaiian or Other Pacific Islander	*	4.93	*	*	*	5.10
Asian	1.96	1.64	*	*	1.99	1.67
Two or More Races	2.52	2.40	*	*	2.58	2.43
Hispanic or Latino	1.06	0.86	1.48	1.31	1.10	0.89
TRIMESTER²						
First	N/A	N/A	2.30	2.64	N/A	N/A
Second	N/A	N/A	0.97	1.12	N/A	N/A
Third	N/A	N/A	1.19	1.17	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.76P Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.8052		0.5588		0.7550	
AGE						
15-17	0.0104		*		0.0108	
18-25	0.4292		0.3428		0.4071	
26-44	0.3631		0.3429		0.2937	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7879		0.4873		0.6934	
White	0.7858		0.4525		0.7832	
Black or African American	0.1182		0.7098		0.0945	
American Indian or Alaska Native	0.6537		*		0.5858	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7974		*		0.7882	
Two or More Races	0.7558		*		0.6051	
Hispanic or Latino	0.7688		0.5822		0.8080	
TRIMESTER²						
First	N/A		0.3922		N/A	
Second	N/A		0.6388		N/A	
Third	N/A		0.7942		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.77C Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	259	241	10	11	258	240
AGE						
15-17	24	26	*	*	24	26
18-25	83	84	4	6	129	110
26-44	214	199	9	9	214	199
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	229	223	10	11	228	222
White	204	195	8	7	203	195
Black or African American	88	86	3	6	87	85
American Indian or Alaska Native	17	18	*	*	17	17
Native Hawaiian or Other Pacific Islander	15	8	*	*	*	8
Asian	48	45	*	*	49	45
Two or More Races	27	26	*	*	27	26
Hispanic or Latino	109	93	3	4	109	92
TRIMESTER²						
First	N/A	N/A	9	10	N/A	N/A
Second	N/A	N/A	3	6	N/A	N/A
Third	N/A	N/A	2	1	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.77D Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.32	0.31	0.46	0.48	0.34	0.32
AGE						
15-17	0.36	0.40	*	*	0.37	0.40
18-25	0.48	0.48	0.55	0.74	0.50	0.50
26-44	0.46	0.42	0.67	0.62	0.47	0.44
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.36	0.36	0.56	0.56	0.37	0.37
White	0.42	0.43	0.67	0.53	0.43	0.45
Black or African American	0.84	0.80	0.86	1.56	0.88	0.83
American Indian or Alaska Native	3.22	3.68	*	*	3.28	3.78
Native Hawaiian or Other Pacific Islander	5.02	2.74	*	*	*	2.89
Asian	1.21	1.02	*	*	1.24	1.07
Two or More Races	2.16	1.91	*	*	2.23	1.96
Hispanic or Latino	0.76	0.65	0.71	0.83	0.79	0.67
TRIMESTER²						
First	N/A	N/A	1.32	1.32	N/A	N/A
Second	N/A	N/A	0.44	0.65	N/A	N/A
Third	N/A	N/A	0.22	0.19	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.77P Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.6577		0.8550		0.6893	
AGE						
15-17	0.0007		*		0.0007	
18-25	0.0143		0.9049		0.0095	
26-44	0.2735		0.7714		0.2342	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6202		0.9742		0.6923	
White	0.5546		0.4576		0.6394	
Black or African American	0.3118		0.5167		0.3548	
American Indian or Alaska Native	0.7820		*		0.6702	
Native Hawaiian or Other Pacific Islander	0.1235		*		*	
Asian	0.9391		*		0.9765	
Two or More Races	0.7402		*		0.7043	
Hispanic or Latino	0.9213		0.7238		1.0000	
TRIMESTER²						
First	N/A		0.9718		N/A	
Second	N/A		0.4222		N/A	
Third	N/A		0.8333		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.78C Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	118	105	3	5	118	105
AGE						
15-17	11	11	*	*	11	10
18-25	52	46	2	3	59	49
26-44	99	91	2	4	99	91
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	105	100	3	5	105	99
White	94	90	3	3	93	90
Black or African American	37	33	0	3	37	33
American Indian or Alaska Native	12	5	*	*	12	5
Native Hawaiian or Other Pacific Islander	*	2	*	*	*	2
Asian	17	18	*	*	17	18
Two or More Races	9	9	*	*	9	9
Hispanic or Latino	47	35	1	*	47	35
TRIMESTER²						
First	N/A	N/A	3	4	N/A	N/A
Second	N/A	N/A	0	3	N/A	N/A
Third	N/A	N/A	*	0	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.78D Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.18	0.16	0.13	0.22	0.19	0.16
AGE						
15-17	0.17	0.17	*	*	0.17	0.17
18-25	0.30	0.27	0.21	0.32	0.31	0.28
26-44	0.25	0.22	0.18	0.26	0.25	0.23
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.20	0.19	0.15	0.24	0.20	0.19
White	0.24	0.24	0.21	0.26	0.25	0.25
Black or African American	0.42	0.36	0.15	0.78	0.44	0.38
American Indian or Alaska Native	2.72	1.48	*	*	2.82	1.53
Native Hawaiian or Other Pacific Islander	*	0.94	*	*	*	1.00
Asian	0.43	0.42	*	*	0.44	0.44
Two or More Races	0.80	0.76	*	*	0.84	0.79
Hispanic or Latino	0.38	0.27	0.22	*	0.40	0.28
TRIMESTER²						
First	N/A	N/A	0.41	0.56	N/A	N/A
Second	N/A	N/A	0.04	0.36	N/A	N/A
Third	N/A	N/A	*	0.03	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.78P Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2013-2014 Versus 2011-2012

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2011-2012	2013-2014	2011-2012	2013-2014	2011-2012	2013-2014
TOTAL	0.3394		0.1367		0.3508	
AGE						
15-17	0.0407		*		0.0266	
18-25	0.1718		0.6456		0.1530	
26-44	0.8673		0.2353		0.9183	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2367		0.1481		0.2505	
White	0.4768		0.3705		0.5100	
Black or African American	0.9128		0.2289		0.9919	
American Indian or Alaska Native	0.1376		*		0.1227	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4312		*		0.4412	
Two or More Races	0.6919		*		0.7143	
Hispanic or Latino	0.6639		*		0.6699	
TRIMESTER²						
First	N/A		0.3084		N/A	
Second	N/A		0.2141		N/A	
Third	N/A		*		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 6.79C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	137	155	98	107	97	110
Marijuana and Hashish	130	146	93	99	90	105
Cocaine	26	32	18	24	20	21
Crack	5	8	1	6	5	5
Heroin	11	11	9	4	6	10
Hallucinogens	40	40	26	25	30	26
LSD	17	14	13	7	11	12
PCP	1	2	*	2	1	*
Ecstasy	25	30	16	17	20	21
Inhalants	17	12	12	8	11	9
Nonmedical Use of Psychotherapeutics ^{4,5}						
Pain Relievers	46	49	24	31	38	39
OxyContin [®]	17	16	13	5	11	15
Tranquilizers	26	30	15	17	21	25
Stimulants ⁵	30	30	23	21	18	21
Methamphetamine ⁵	13	13	6	7	12	11
Sedatives	7	11	3	3	6	11
ILLCIT DRUGS OTHER THAN MARIJUANA³	68	78	45	54	53	55

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 22, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.79D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	0.51	0.57	0.84	0.87	0.63	0.71
Marijuana and Hashish	0.49	0.55	0.82	0.86	0.60	0.70
Cocaine	0.12	0.14	0.20	0.26	0.15	0.16
Crack	0.02	0.04	0.01	0.07	0.04	0.04
Heroin	0.05	0.05	0.10	0.04	0.04	0.07
Hallucinogens	0.18	0.18	0.29	0.28	0.22	0.20
LSD	0.08	0.06	0.15	0.08	0.09	0.09
PCP	0.01	0.01	*	0.02	0.01	*
Ecstasy	0.11	0.13	0.18	0.19	0.15	0.16
Inhalants	0.08	0.05	0.14	0.09	0.08	0.07
Nonmedical Use of Psychotherapeutics ^{4,5}	0.25	0.26	0.39	0.43	0.32	0.34
Pain Relievers	0.20	0.21	0.28	0.33	0.28	0.29
OxyContin [®]	0.08	0.07	0.15	0.06	0.08	0.12
Tranquilizers	0.12	0.13	0.18	0.19	0.16	0.18
Stimulants ⁵	0.13	0.13	0.26	0.23	0.14	0.16
Methamphetamine ⁵	0.06	0.06	0.07	0.07	0.09	0.08
Sedatives	0.03	0.05	0.04	0.04	0.05	0.08
ILLCIT DRUGS OTHER THAN MARIJUANA³	0.29	0.33	0.47	0.55	0.38	0.40

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 22, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.79P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	0.7004		0.9638		0.6313	
Marijuana and Hashish	0.6743		0.7141		0.7962	
Cocaine	0.2480		0.0248		0.5315	
Crack	0.6798		0.2111		0.6024	
Heroin	0.8726		0.2716		0.4665	
Hallucinogens	0.0812		0.1318		0.2589	
LSD	0.3257		0.1192		0.9532	
PCP	0.9392		*		*	
Ecstasy	0.3905		0.5827		0.4720	
Inhalants	0.0571		0.1663		0.1882	
Nonmedical Use of Psychotherapeutics ^{4,5}	0.5558		0.9084		0.4001	
Pain Relievers	0.3665		0.8891		0.2398	
OxyContin [®]	0.6198		0.0317		0.3375	
Tranquilizers	0.8520		0.9372		0.7709	
Stimulants ⁵	0.4104		0.6316		0.4756	
Methamphetamine ⁵	0.4777		0.7966		0.3533	
Sedatives	0.5069		0.9624		0.4854	
ILLCIT DRUGS OTHER THAN MARIJUANA³	0.3887		0.9041		0.3046	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 22, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.80C Types of Illicit Drug Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Male College Students		Other Males Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	103	113	69	70	79	83
Marijuana and Hashish	98	108	66	66	74	81
Cocaine	19	25	9	17	17	17
Crack	3	7	*	6	3	4
Heroin	10	9	9	*	4	9
Hallucinogens	36	25	24	16	27	18
LSD	15	8	11	3	10	7
PCP	1	2	*	2	1	*
Ecstasy	21	17	13	11	16	13
Inhalants	15	9	11	7	10	5
Nonmedical Use of Psychotherapeutics ^{4,5}	42	44	26	29	34	34
Pain Relievers	35	39	18	24	30	30
OxyContin [®]	13	13	11	5	7	12
Tranquilizers	19	22	12	13	15	18
Stimulants ⁵	21	21	16	14	13	15
Methamphetamine ⁵	10	9	5	3	9	9
Sedatives	5	6	2	1	5	6
ILLCIT DRUGS OTHER THAN MARIJUANA³	54	53	34	35	44	41

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all males aged 18 to 22, including those with unknown enrollment status.

² Other Males include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.80D Types of Illicit Drug Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Male College Students		Other Males Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	0.74	0.82	1.27	1.33	0.93	0.98
Marijuana and Hashish	0.72	0.80	1.24	1.29	0.89	0.97
Cocaine	0.17	0.22	0.22	0.40	0.24	0.24
Crack	0.03	0.06	*	0.14	0.05	0.05
Heroin	0.09	0.08	0.22	*	0.06	0.12
Hallucinogens	0.32	0.22	0.58	0.39	0.37	0.25
LSD	0.13	0.07	0.28	0.07	0.14	0.10
PCP	0.01	0.01	*	0.04	0.02	*
Ecstasy	0.19	0.15	0.33	0.27	0.23	0.19
Inhalants	0.13	0.08	0.26	0.16	0.14	0.07
Nonmedical Use of Psychotherapeutics ^{4,5}	0.36	0.37	0.60	0.66	0.46	0.46
Pain Relievers	0.31	0.33	0.45	0.56	0.41	0.40
OxyContin [®]	0.11	0.12	0.26	0.13	0.09	0.17
Tranquilizers	0.17	0.19	0.29	0.31	0.21	0.25
Stimulants ⁵	0.19	0.18	0.39	0.33	0.19	0.21
Methamphetamine ⁵	0.09	0.08	0.13	0.07	0.12	0.13
Sedatives	0.05	0.05	0.06	0.03	0.07	0.08
ILLCIT DRUGS OTHER THAN MARIJUANA³	0.45	0.44	0.77	0.77	0.58	0.55

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all males aged 18 to 22, including those with unknown enrollment status.

² Other Males include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.80P Types of Illicit Drug Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Male College Students		Other Males Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLICIT DRUGS³	0.4824		0.4463		0.7695	
Marijuana and Hashish	0.5834		0.5443		0.8281	
Cocaine	0.6244		0.0556		0.3945	
Crack	0.3445		*		0.8363	
Heroin	0.8451		*		0.3335	
Hallucinogens	0.0025		0.0539		0.0158	
LSD	0.0252		0.0473		0.2285	
PCP	0.8917		*		*	
Ecstasy	0.0307		0.2449		0.0658	
Inhalants	0.0317		0.2867		0.0425	
Nonmedical Use of Psychotherapeutics ^{4,5}	0.3614		0.6144		0.4560	
Pain Relievers	0.4403		0.6243		0.5552	
OxyContin [®]	0.8449		0.1438		0.1286	
Tranquilizers	0.6345		0.7008		0.7558	
Stimulants ⁵	0.2781		0.3206		0.5710	
Methamphetamine ⁵	0.5351		0.5572		0.6906	
Sedatives	0.6286		0.6525		0.7328	
ILLICIT DRUGS OTHER THAN MARIJUANA³	0.0297		0.1767		0.0957	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all males aged 18 to 22, including those with unknown enrollment status.

² Other Males include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.81C Types of Illicit Drug Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Female College Students		Other Females Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	83	95	58	63	55	71
Marijuana and Hashish	78	90	54	61	52	67
Cocaine	17	18	15	15	10	11
Crack	3	4	1	*	3	4
Heroin	5	6	2	4	4	4
Hallucinogens	17	25	11	13	13	18
LSD	8	11	6	5	5	10
PCP	*	*	*	*	0	*
Ecstasy	14	22	8	12	11	15
Inhalants	7	8	6	3	4	7
Nonmedical Use of Psychotherapeutics ^{4,5}						
Pain Relievers	36	38	23	25	28	28
OxyContin [®]	29	28	15	18	24	23
Tranquilizers	11	6	7	0	8	6
Stimulants ⁵	18	19	10	10	15	16
Methamphetamine ⁵	19	20	14	15	13	13
Sedatives	9	9	3	6	8	6
Sedatives	4	10	2	3	4	9
ILLCIT DRUGS OTHER THAN MARIJUANA³	41	50	27	33	30	34

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 18 to 22, including those with unknown enrollment status.

² Other Females include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.81D Types of Illicit Drug Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Female College Students		Other Females Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	0.67	0.77	1.04	1.11	0.83	1.03
Marijuana and Hashish	0.65	0.75	1.00	1.13	0.79	1.00
Cocaine	0.16	0.17	0.32	0.32	0.16	0.19
Crack	0.03	0.04	0.02	*	0.06	0.07
Heroin	0.04	0.05	0.04	0.08	0.07	0.07
Hallucinogens	0.16	0.23	0.24	0.29	0.21	0.30
LSD	0.08	0.11	0.14	0.12	0.08	0.17
PCP	*	*	*	*	0.01	*
Ecstasy	0.13	0.21	0.17	0.25	0.18	0.25
Inhalants	0.06	0.07	0.12	0.08	0.06	0.12
Nonmedical Use of Psychotherapeutics ^{4,5}	0.33	0.34	0.48	0.52	0.45	0.45
Pain Relievers	0.27	0.26	0.33	0.38	0.39	0.36
OxyContin [®]	0.10	0.05	0.15	0.01	0.14	0.10
Tranquilizers	0.17	0.18	0.22	0.21	0.25	0.27
Stimulants ⁵	0.17	0.19	0.30	0.33	0.21	0.21
Methamphetamine ⁵	0.08	0.08	0.05	0.13	0.14	0.10
Sedatives	0.04	0.09	0.04	0.07	0.06	0.15
ILLCIT DRUGS OTHER THAN MARIJUANA³	0.37	0.45	0.56	0.68	0.48	0.55

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 18 to 22, including those with unknown enrollment status.

² Other Females include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.81P Types of Illicit Drug Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Female College Students		Other Females Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
ILLCIT DRUGS³	0.1998		0.4104		0.3126	
Marijuana and Hashish	0.2479		0.2631		0.5720	
Cocaine	0.2123		0.1613		0.8948	
Crack	0.5247		*		0.6306	
Heroin	1.0000		0.7865		0.8648	
Hallucinogens	0.2601		0.9246		0.1473	
LSD	0.2544		0.9751		0.1625	
PCP	*		*		*	
Ecstasy	0.3330		0.5880		0.3472	
Inhalants	0.8597		0.3313		0.5630	
Nonmedical Use of Psychotherapeutics ^{4,5}	0.9014		0.4732		0.6648	
Pain Relievers	0.6186		0.4087		0.2353	
OxyContin [®]	0.2092		0.0729		0.7544	
Tranquilizers	0.8055		0.5430		0.9215	
Stimulants ⁵	0.9382		0.7752		0.6515	
Methamphetamine ⁵	0.7142		0.3792		0.3376	
Sedatives	0.2317		0.6863		0.2585	
ILLCIT DRUGS OTHER THAN MARIJUANA³	0.2637		0.2101		0.6764	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 18 to 22, including those with unknown enrollment status.

² Other Females include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.82C Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	137	155	98	107	97	110
AGE						
18	56	70	33	49	45	47
19	55	59	40	39	37	46
20	61	65	41	45	43	47
21	57	63	42	41	37	48
22	55	65	32	38	45	52
GENDER						
Male	103	113	69	70	79	83
Female	83	95	58	63	55	71
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	125	138	89	96	84	96
White	108	119	81	79	70	82
Black or African American	44	55	27	31	35	46
American Indian or Alaska Native	6	*	*	*	5	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	22	27	14	24	18	12
Two or More Races	18	21	*	16	14	14
Hispanic or Latino	58	61	36	35	49	51
CURRENT EMPLOYMENT						
Full-Time	64	82	31	37	56	74
Part-Time	79	84	61	62	48	54
Unemployed	52	47	28	20	44	42
Other ³	63	65	49	55	37	35

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.82D Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.51	0.57	0.84	0.87	0.63	0.71
AGE						
18	1.00	1.23	1.63	2.25	1.28	1.35
19	1.14	1.25	1.65	1.63	1.54	1.81
20	1.17	1.23	1.72	1.84	1.57	1.71
21	1.12	1.17	1.86	1.65	1.34	1.59
22	1.07	1.21	2.03	2.27	1.26	1.41
GENDER						
Male	0.74	0.82	1.27	1.33	0.93	0.98
Female	0.67	0.77	1.04	1.11	0.83	1.03
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.58	0.64	0.90	0.95	0.74	0.83
White	0.70	0.77	1.06	1.16	0.89	1.00
Black or African American	1.18	1.46	2.05	2.30	1.45	1.82
American Indian or Alaska Native	3.75	*	*	*	4.66	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.83	1.95	1.85	2.51	3.78	2.81
Two or More Races	3.11	3.27	*	5.50	3.97	4.00
Hispanic or Latino	1.07	1.14	2.14	1.92	1.28	1.42
CURRENT EMPLOYMENT						
Full-Time	0.87	0.99	1.93	2.06	0.98	1.15
Part-Time	0.90	0.96	1.29	1.20	1.22	1.50
Unemployed	1.42	1.53	3.20	2.71	1.60	1.82
Other ³	0.99	1.00	1.39	1.56	1.30	1.26

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.82P Illicit Drug Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.7004		0.9638		0.6313	
AGE						
18	0.9621		0.6919		0.7729	
19	0.6853		0.7892		0.4530	
20	0.8668		0.2312		0.1860	
21	0.6460		0.0069		0.0941	
22	0.2739		0.1358		0.7491	
GENDER						
Male	0.4824		0.4463		0.7695	
Female	0.1998		0.4104		0.3126	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3751		0.5210		0.5308	
White	0.9416		0.7631		0.8786	
Black or African American	0.0366		0.2872		0.0661	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.9731		0.0927		0.0407	
Two or More Races	0.6054		*		0.4709	
Hispanic or Latino	0.3785		0.2255		0.8431	
CURRENT EMPLOYMENT						
Full-Time	0.0342		0.2635		0.0740	
Part-Time	0.6229		0.7875		0.6851	
Unemployed	0.2693		0.1509		0.6601	
Other ³	0.9249		0.7963		0.6713	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.83C Marijuana Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	130	146	93	99	90	105
AGE						
18	52	66	32	45	41	45
19	53	56	38	37	36	44
20	56	63	39	44	40	45
21	54	57	39	36	35	45
22	52	61	31	35	41	48
GENDER						
Male	98	108	66	66	74	81
Female	78	90	54	61	52	67
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	118	132	84	91	78	91
White	103	112	76	77	67	76
Black or African American	42	54	25	29	34	46
American Indian or Alaska Native	4	*	*	*	4	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	15	22	12	19	11	11
Two or More Races	17	20	*	14	13	14
Hispanic or Latino	55	57	35	32	47	47
CURRENT EMPLOYMENT						
Full-Time	60	78	30	35	53	71
Part-Time	76	79	59	59	46	51
Unemployed	50	45	25	20	43	41
Other ³	59	60	46	50	33	33

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.83D Marijuana Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.49	0.55	0.82	0.86	0.60	0.70
AGE						
18	0.95	1.19	1.56	2.14	1.21	1.32
19	1.11	1.21	1.60	1.62	1.51	1.74
20	1.11	1.21	1.66	1.85	1.49	1.66
21	1.08	1.09	1.78	1.53	1.27	1.51
22	1.02	1.15	2.01	2.19	1.17	1.33
GENDER						
Male	0.72	0.80	1.24	1.29	0.89	0.97
Female	0.65	0.75	1.00	1.13	0.79	1.00
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.56	0.63	0.89	0.96	0.70	0.81
White	0.69	0.75	1.06	1.17	0.87	0.95
Black or African American	1.15	1.43	1.96	2.22	1.41	1.81
American Indian or Alaska Native	2.91	*	*	*	3.57	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.34	1.81	1.68	2.35	2.53	2.66
Two or More Races	3.01	3.14	*	5.18	3.76	3.94
Hispanic or Latino	1.02	1.07	2.09	1.79	1.22	1.35
CURRENT EMPLOYMENT						
Full-Time	0.85	0.97	1.89	2.02	0.95	1.13
Part-Time	0.88	0.93	1.26	1.18	1.18	1.43
Unemployed	1.41	1.51	3.00	2.64	1.58	1.79
Other ³	0.93	0.97	1.32	1.51	1.20	1.20

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.83P Marijuana Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.6743		0.7141		0.7962	
AGE						
18	0.9928		0.6295		0.7573	
19	0.6852		0.9304		0.5649	
20	0.5824		0.0499		0.2735	
21	0.4446		0.0069		0.2084	
22	0.4731		0.3241		0.8537	
GENDER						
Male	0.5834		0.5443		0.8281	
Female	0.2479		0.2631		0.5720	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2617		0.3369		0.5023	
White	0.8937		0.8551		0.7224	
Black or African American	0.0040		0.0826		0.0192	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.9356		0.3286		0.2127	
Two or More Races	0.6393		*		0.7205	
Hispanic or Latino	0.1870		0.2307		0.4615	
CURRENT EMPLOYMENT						
Full-Time	0.0481		0.2374		0.1097	
Part-Time	0.6421		0.9305		0.5696	
Unemployed	0.3382		0.4294		0.5154	
Other ³	0.9874		0.8029		0.8062	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.83P Marijuana Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.6743		0.7141		0.7962	
AGE						
18	0.9928		0.6295		0.7573	
19	0.6852		0.9304		0.5649	
20	0.5824		0.0499		0.2735	
21	0.4446		0.0069		0.2084	
22	0.4731		0.3241		0.8537	
GENDER						
Male	0.5834		0.5443		0.8281	
Female	0.2479		0.2631		0.5720	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2617		0.3369		0.5023	
White	0.8937		0.8551		0.7224	
Black or African American	0.0040		0.0826		0.0192	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.9356		0.3286		0.2127	
Two or More Races	0.6393		*		0.7205	
Hispanic or Latino	0.1870		0.2307		0.4615	
CURRENT EMPLOYMENT						
Full-Time	0.0481		0.2374		0.1097	
Part-Time	0.6421		0.9305		0.5696	
Unemployed	0.3382		0.4294		0.5154	
Other ³	0.9874		0.8029		0.8062	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.84C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	174	182	111	107	136	145
Cigarettes	154	161	90	87	127	133
Smokeless Tobacco	59	68	37	40	47	57
Cigars	88	90	52	51	68	73
Pipe Tobacco	43	37	31	28	30	29
ALCOHOL	256	270	199	222	156	167
Binge Alcohol Use ⁴	200	214	147	174	125	127
Heavy Alcohol Use ⁴	95	107	70	78	63	70

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 22, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.84D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	0.58	0.65	0.91	0.89	0.73	0.83
Cigarettes	0.57	0.62	0.83	0.81	0.73	0.82
Smokeless Tobacco	0.26	0.30	0.40	0.42	0.34	0.41
Cigars	0.36	0.38	0.51	0.53	0.46	0.51
Pipe Tobacco	0.19	0.17	0.33	0.31	0.22	0.22
ALCOHOL	0.64	0.71	1.06	1.22	0.76	0.84
Binge Alcohol Use ⁴	0.64	0.71	1.02	1.23	0.74	0.80
Heavy Alcohol Use ⁴	0.37	0.44	0.65	0.74	0.43	0.50

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 22, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.84P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	0.0276		0.0021		0.6425	
Cigarettes	0.0051		0.0084		0.0965	
Smokeless Tobacco	0.4135		0.1021		0.8402	
Cigars	0.7499		0.1035		0.4507	
Pipe Tobacco	0.0184		0.0580		0.1746	
ALCOHOL	0.4643		0.7943		0.4489	
Binge Alcohol Use ⁴	0.7357		0.5213		0.9044	
Heavy Alcohol Use ⁴	0.8806		0.6210		0.8057	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 22, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.85C Tobacco Product and Alcohol Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Male College Students		Other Males Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	137	143	83	81	110	116
Cigarettes	119	124	66	66	99	104
Smokeless Tobacco	58	66	35	40	47	56
Cigars	79	75	44	40	62	63
Pipe Tobacco	36	30	25	20	25	23
ALCOHOL	161	180	113	131	113	124
Binge Alcohol Use ⁴	132	148	93	110	92	102
Heavy Alcohol Use ⁴	74	80	53	56	51	56

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all males aged 18 to 22, including those with unknown enrollment status.

² Other Males include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.85D Tobacco Product and Alcohol Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Male College Students		Other Males Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	0.84	0.95	1.38	1.45	1.05	1.17
Cigarettes	0.83	0.90	1.29	1.31	1.01	1.15
Smokeless Tobacco	0.48	0.56	0.78	0.86	0.62	0.74
Cigars	0.62	0.60	0.94	0.89	0.77	0.78
Pipe Tobacco	0.31	0.27	0.58	0.46	0.35	0.32
ALCOHOL	0.86	0.95	1.37	1.69	1.06	1.11
Binge Alcohol Use ⁴	0.84	0.94	1.41	1.70	1.00	1.11
Heavy Alcohol Use ⁴	0.58	0.63	1.08	1.13	0.65	0.73

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all males aged 18 to 22, including those with unknown enrollment status.

² Other Males include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.85P Tobacco Product and Alcohol Use in the Past Month among Males Aged 18 to 22, by College Enrollment Status: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Male College Students		Other Males Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	0.0984		0.0105		0.8676	
Cigarettes	0.0447		0.0739		0.2255	
Smokeless Tobacco	0.5324		0.1888		0.8526	
Cigars	0.2302		0.0192		0.9015	
Pipe Tobacco	0.0065		0.0118		0.1411	
ALCOHOL	0.8166		0.8278		0.6324	
Binge Alcohol Use ⁴	0.5711		0.5330		0.8181	
Heavy Alcohol Use ⁴	0.5148		0.4794		0.7893	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all males aged 18 to 22, including those with unknown enrollment status.

² Other Males include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.86C Tobacco Product and Alcohol Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Standard Errors of Numbers in Thousands, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Female College Students		Other Females Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	94	101	60	56	72	81
Cigarettes	85	90	52	47	68	72
Smokeless Tobacco	16	11	10	4	13	10
Cigars	38	48	23	29	29	39
Pipe Tobacco	21	21	15	19	15	17
ALCOHOL	155	170	122	136	93	102
Binge Alcohol Use ⁴	116	126	84	97	74	75
Heavy Alcohol Use ⁴	53	57	40	43	34	36

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 18 to 22, including those with unknown enrollment status.

² Other Females include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.86D Tobacco Product and Alcohol Use in the Past Month among Females Aged 18 to 22, by College Enrollment Status: Standard Errors of Percentages, 2013 and 2014

Substance	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time Female College Students		Other Females Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOBACCO PRODUCTS³	0.71	0.79	1.06	1.04	0.95	1.09
Cigarettes	0.69	0.75	1.02	0.92	0.92	1.02
Smokeless Tobacco	0.15	0.10	0.21	0.09	0.21	0.17
Cigars	0.33	0.43	0.45	0.61	0.45	0.61
Pipe Tobacco	0.19	0.19	0.32	0.41	0.24	0.28
ALCOHOL	0.85	0.95	1.39	1.53	1.00	1.14
Binge Alcohol Use ⁴	0.82	0.91	1.30	1.44	1.00	1.04
Heavy Alcohol Use ⁴	0.45	0.51	0.77	0.85	0.53	0.58

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 18 to 22, including those with unknown enrollment status.

² Other Females include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.87C Cigarette Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	154	161	90	87	127	133
AGE						
18	54	60	26	31	46	50
19	62	57	38	31	47	48
20	69	68	41	41	56	54
21	60	66	37	38	47	54
22	67	74	30	31	59	66
GENDER						
Male	119	124	66	66	99	104
Female	85	90	52	47	68	72
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	135	143	83	79	106	116
White	118	128	77	70	90	103
Black or African American	49	49	21	18	44	46
American Indian or Alaska Native	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	20	26	14	21	14	14
Two or More Races	20	17	10	*	17	13
Hispanic or Latino	66	70	32	29	62	62
CURRENT EMPLOYMENT						
Full-Time	85	97	28	31	81	92
Part-Time	78	80	53	52	58	58
Unemployed	55	55	26	16	49	53
Other ³	61	69	46	54	39	46

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.87D Cigarette Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.57	0.62	0.83	0.81	0.73	0.82
AGE						
18	1.05	1.14	1.58	1.61	1.29	1.51
19	1.25	1.21	1.63	1.44	1.80	1.77
20	1.24	1.31	1.73	1.77	1.75	1.86
21	1.15	1.28	1.83	1.73	1.52	1.75
22	1.25	1.33	1.98	2.01	1.51	1.59
GENDER						
Male	0.83	0.90	1.29	1.31	1.01	1.15
Female	0.69	0.75	1.02	0.92	0.92	1.02
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.61	0.70	0.90	0.89	0.80	0.93
White	0.73	0.85	1.08	1.12	0.96	1.11
Black or African American	1.27	1.37	1.74	1.54	1.71	1.82
American Indian or Alaska Native	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.77	1.95	2.04	2.37	3.03	3.17
Two or More Races	3.30	2.88	4.21	*	4.55	3.80
Hispanic or Latino	1.15	1.32	2.00	1.65	1.49	1.74
CURRENT EMPLOYMENT						
Full-Time	1.01	1.16	1.74	1.88	1.18	1.35
Part-Time	0.89	0.96	1.17	1.17	1.34	1.51
Unemployed	1.49	1.74	2.92	2.30	1.75	2.12
Other ³	1.02	1.06	1.46	1.60	1.38	1.54

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.87P Cigarette Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.0051		0.0084		0.0965	
AGE						
18	0.3032		0.6412		0.4697	
19	0.0093		0.0043		0.1324	
20	0.0550		0.3177		0.0681	
21	0.2517		0.0757		0.9260	
22	0.6462		0.9594		0.6049	
GENDER						
Male	0.0447		0.0739		0.2255	
Female	0.0158		0.0297		0.1511	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0088		0.0332		0.0562	
White	0.0165		0.0222		0.1094	
Black or African American	0.8836		0.4094		0.7966	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.8546		0.2220		0.2373	
Two or More Races	0.0221		*		0.0116	
Hispanic or Latino	0.2542		0.0798		0.8556	
CURRENT EMPLOYMENT						
Full-Time	0.0961		0.3684		0.1343	
Part-Time	0.0055		0.0380		0.1128	
Unemployed	0.1886		0.0007		0.9298	
Other ³	0.5382		0.7745		0.5506	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.88C Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	256	270	199	222	156	167
AGE						
18	87	97	64	78	57	61
19	85	86	66	67	51	56
20	92	102	73	76	56	66
21	105	121	78	92	69	77
22	106	117	56	68	84	96
GENDER						
Male	161	180	113	131	113	124
Female	155	170	122	136	93	102
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	230	236	182	196	130	143
White	200	207	160	161	109	126
Black or African American	72	84	47	53	50	63
American Indian or Alaska Native	10	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	42	55	34	48	23	26
Two or More Races	26	29	16	21	20	20
Hispanic or Latino	98	106	57	69	83	79
CURRENT EMPLOYMENT						
Full-Time	113	132	48	58	103	119
Part-Time	140	149	110	121	78	83
Unemployed	70	65	39	35	59	57
Other ³	118	125	105	113	47	54

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.88D Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.64	0.71	1.06	1.22	0.76	0.84
AGE						
18	1.22	1.50	2.07	2.87	1.48	1.64
19	1.38	1.47	1.98	2.19	1.82	1.95
20	1.41	1.52	2.06	2.19	1.75	1.95
21	1.22	1.34	1.85	1.94	1.60	1.69
22	1.20	1.34	2.18	2.26	1.45	1.69
GENDER						
Male	0.86	0.95	1.37	1.69	1.06	1.11
Female	0.85	0.95	1.39	1.53	1.00	1.14
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.70	0.75	1.18	1.27	0.83	0.92
White	0.80	0.85	1.20	1.35	0.97	1.05
Black or African American	1.48	1.64	2.84	2.79	1.71	2.09
American Indian or Alaska Native	5.90	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.70	3.19	3.46	3.77	4.27	4.86
Two or More Races	3.73	3.58	6.07	5.88	4.49	4.38
Hispanic or Latino	1.37	1.55	2.46	2.62	1.69	1.85
CURRENT EMPLOYMENT						
Full-Time	1.04	1.11	2.19	2.27	1.16	1.28
Part-Time	1.09	1.21	1.49	1.55	1.46	1.75
Unemployed	1.57	1.88	3.18	3.65	1.82	2.19
Other ³	1.33	1.44	1.91	2.11	1.53	1.69

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.88P Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.4643		0.7943		0.4489	
AGE						
18	0.4899		0.2328		0.8358	
19	0.2352		0.3645		0.5140	
20	0.4137		0.8038		0.1230	
21	0.3175		0.7077		0.3879	
22	0.9133		0.8761		0.9583	
GENDER						
Male	0.8166		0.8278		0.6324	
Female	0.4012		0.5682		0.5487	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2545		0.7733		0.1724	
White	0.5511		0.8251		0.2428	
Black or African American	0.4290		0.9649		0.3274	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1331		0.1102		0.7029	
Two or More Races	0.3185		0.2917		0.6973	
Hispanic or Latino	0.6477		0.8881		0.4654	
CURRENT EMPLOYMENT						
Full-Time	0.0861		0.1466		0.2288	
Part-Time	0.4713		0.5870		0.3782	
Unemployed	0.3684		0.2234		0.7151	
Other ³	0.6123		0.5957		0.7306	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.89C Binge Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2013 and 2014**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	200	214	147	174	125	127
AGE						
18	63	78	41	60	47	50
19	71	68	53	51	44	44
20	71	79	54	57	45	54
21	81	95	59	73	53	62
22	83	91	46	55	64	70
GENDER						
Male	132	148	93	110	92	102
Female	116	126	84	97	74	75
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	177	187	138	155	102	110
White	156	165	125	132	87	99
Black or African American	50	57	30	33	37	46
American Indian or Alaska Native	7	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	29	37	20	31	21	18
Two or More Races	20	22	13	17	15	14
Hispanic or Latino	80	82	45	52	66	62
CURRENT EMPLOYMENT						
Full-Time	91	105	40	47	82	94
Part-Time	110	116	87	97	61	61
Unemployed	58	55	28	28	51	49
Other ³	87	97	76	87	37	42

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.89D Binge Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
Standard Errors of Percentages, 2013 and 2014**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.64	0.71	1.02	1.23	0.74	0.80
AGE						
18	1.05	1.35	1.85	2.55	1.32	1.48
19	1.28	1.33	1.86	1.94	1.70	1.71
20	1.28	1.39	1.90	2.10	1.62	1.84
21	1.31	1.43	2.14	2.29	1.58	1.80
22	1.29	1.45	2.24	2.78	1.50	1.68
GENDER						
Male	0.84	0.94	1.41	1.70	1.00	1.11
Female	0.82	0.91	1.30	1.44	1.00	1.04
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.69	0.78	1.13	1.32	0.81	0.90
White	0.81	0.89	1.20	1.49	0.99	1.08
Black or African American	1.21	1.47	2.23	2.37	1.48	1.84
American Indian or Alaska Native	4.92	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.32	2.41	2.75	2.96	4.02	3.70
Two or More Races	3.39	3.40	5.51	5.52	4.20	3.94
Hispanic or Latino	1.38	1.42	2.39	2.46	1.64	1.68
CURRENT EMPLOYMENT						
Full-Time	1.09	1.13	2.17	2.37	1.21	1.26
Part-Time	1.07	1.20	1.47	1.65	1.41	1.57
Unemployed	1.57	1.75	3.09	3.51	1.78	2.05
Other ³	1.18	1.32	1.87	2.06	1.28	1.44

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.89P Binge Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.7357		0.5213		0.9044	
AGE						
18	0.3563		0.3037		0.7836	
19	0.0248		0.0800		0.1605	
20	0.9181		0.4454		0.3513	
21	0.3286		0.8359		0.2879	
22	0.2840		0.4921		0.3901	
GENDER						
Male	0.5711		0.5330		0.8181	
Female	0.9523		0.6522		0.7235	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.8773		0.6461		0.7897	
White	0.8374		0.4606		0.6126	
Black or African American	0.6435		0.7548		0.7359	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7340		0.2361		0.3660	
Two or More Races	0.9077		0.7290		0.8583	
Hispanic or Latino	0.6606		0.6054		0.8218	
CURRENT EMPLOYMENT						
Full-Time	0.5787		0.8490		0.5900	
Part-Time	0.1991		0.2327		0.3555	
Unemployed	0.7119		0.9230		0.7102	
Other ³	0.9064		0.9598		0.9897	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.90C Heavy Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	95	107	70	78	63	70
AGE						
18	30	35	20	23	22	26
19	33	34	27	25	18	21
20	40	39	32	30	24	24
21	42	51	30	37	27	35
22	44	51	24	30	36	38
GENDER						
Male	74	80	53	56	51	56
Female	53	57	40	43	34	36
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	85	92	66	69	51	60
White	79	83	62	61	48	54
Black or African American	20	24	12	12	15	20
American Indian or Alaska Native	2	4	*	*	2	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	12	16	10	15	5	*
Two or More Races	7	14	3	*	7	9
Hispanic or Latino	40	43	24	28	32	31
CURRENT EMPLOYMENT						
Full-Time	52	60	23	30	45	50
Part-Time	54	53	43	42	32	31
Unemployed	29	29	17	14	24	26
Other ³	38	48	35	44	16	18

*Low precision; no estimate reported.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.90D Heavy Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.37	0.44	0.65	0.74	0.43	0.50
AGE						
18	0.57	0.72	1.01	1.23	0.69	0.85
19	0.71	0.78	1.18	1.20	0.83	0.94
20	0.85	0.83	1.40	1.42	0.98	0.96
21	0.86	1.03	1.47	1.63	1.01	1.30
22	0.88	1.02	1.69	2.02	1.03	1.10
GENDER						
Male	0.58	0.63	1.08	1.13	0.65	0.73
Female	0.45	0.51	0.77	0.85	0.53	0.58
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.42	0.48	0.72	0.79	0.47	0.56
White	0.54	0.59	0.89	0.96	0.63	0.71
Black or African American	0.59	0.72	1.12	1.08	0.71	0.93
American Indian or Alaska Native	1.57	2.66	*	*	1.91	1.70
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.04	1.27	1.53	1.81	1.09	*
Two or More Races	1.38	2.44	1.38	*	2.17	2.76
Hispanic or Latino	0.78	0.86	1.58	1.67	0.91	0.94
CURRENT EMPLOYMENT						
Full-Time	0.76	0.80	1.56	1.85	0.82	0.86
Part-Time	0.67	0.69	0.98	0.96	0.85	0.92
Unemployed	0.94	0.98	2.25	1.92	1.00	1.21
Other ³	0.64	0.81	1.06	1.37	0.59	0.70

*Low precision; no estimate reported.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.90P Heavy Alcohol Use in the Past Month among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.8806		0.6210		0.8057	
AGE						
18	0.8814		0.2824		0.5548	
19	0.5099		0.2338		0.5556	
20	0.4550		0.9154		0.3265	
21	0.1977		0.6190		0.2047	
22	0.6027		0.8473		0.4148	
GENDER						
Male	0.5148		0.4794		0.7893	
Female	0.6453		0.9665		0.4776	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6547		0.5275		0.9944	
White	0.3474		0.3363		0.7376	
Black or African American	0.4661		0.9308		0.4176	
American Indian or Alaska Native	0.5142		*		0.9452	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.6971		0.5630		*	
Two or More Races	0.0723		*		0.4954	
Hispanic or Latino	0.5734		0.7733		0.6458	
CURRENT EMPLOYMENT						
Full-Time	0.8288		0.5487		0.5549	
Part-Time	0.3452		0.1326		0.8834	
Unemployed	0.7611		0.2287		0.2748	
Other ³	0.6183		0.4537		0.8161	

*Low precision; no estimate reported.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.91C Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	73	75	45	46	58	58
AGE						
18	35	34	15	17	31	28
19	33	31	22	18	23	25
20	31	33	22	23	22	23
21	35	30	24	17	25	26
22	31	35	13	16	27	30
GENDER						
Male	58	59	34	29	48	50
Female	41	43	27	31	31	31
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	68	66	41	40	53	51
White	58	56	38	33	42	44
Black or African American	24	27	11	13	21	24
American Indian or Alaska Native	5	1	*	*	*	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	19	18	6	16	17	8
Two or More Races	14	10	5	6	13	8
Hispanic or Latino	26	34	17	16	25	29
CURRENT EMPLOYMENT						
Full-Time	37	41	15	14	34	39
Part-Time	39	41	29	30	26	27
Unemployed	36	29	17	13	31	27
Other ³	33	32	23	23	24	22

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.91D Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.32	0.33	0.48	0.49	0.43	0.43
AGE						
18	0.70	0.71	0.88	0.93	0.95	0.91
19	0.77	0.72	1.05	0.88	1.08	1.10
20	0.70	0.73	1.03	1.11	0.91	0.93
21	0.76	0.64	1.31	0.90	0.95	0.96
22	0.67	0.74	1.01	1.20	0.84	0.90
GENDER						
Male	0.49	0.50	0.78	0.66	0.65	0.68
Female	0.37	0.40	0.56	0.65	0.51	0.50
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.37	0.37	0.52	0.52	0.52	0.49
White	0.44	0.44	0.66	0.62	0.60	0.60
Black or African American	0.73	0.82	1.00	1.17	0.97	1.07
American Indian or Alaska Native	3.55	1.12	*	*	*	1.49
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.60	1.40	0.87	1.82	3.67	2.01
Two or More Races	2.56	1.87	2.37	2.78	3.83	2.52
Hispanic or Latino	0.55	0.70	1.16	1.01	0.74	0.90
CURRENT EMPLOYMENT						
Full-Time	0.55	0.58	1.05	0.90	0.65	0.68
Part-Time	0.51	0.56	0.72	0.73	0.75	0.83
Unemployed	1.12	1.04	2.24	1.83	1.24	1.26
Other ³	0.59	0.56	0.79	0.80	0.87	0.81

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.91P Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.0818		0.2488		0.1922	
AGE						
18	0.2513		0.8744		0.2602	
19	0.3454		0.7469		0.2833	
20	0.2641		0.8845		0.1407	
21	0.0878		0.0098		0.9619	
22	0.3662		0.3112		0.6052	
GENDER						
Male	0.0497		0.0201		0.4149	
Female	0.6207		0.5104		0.2056	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0765		0.3620		0.1141	
White	0.0793		0.0408		0.4851	
Black or African American	0.9229		0.5507		0.8515	
American Indian or Alaska Native	0.0225		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.8364		0.0297		0.0380	
Two or More Races	0.6677		0.5268		0.4015	
Hispanic or Latino	0.7235		0.4123		0.8667	
CURRENT EMPLOYMENT						
Full-Time	0.4299		0.0247		0.9563	
Part-Time	0.2716		0.9162		0.1662	
Unemployed	0.1095		0.1625		0.2853	
Other ³	0.9795		0.7753		0.7476	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.92C Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	99	104	70	77	68	68
AGE						
18	38	40	26	31	29	28
19	37	36	26	25	26	25
20	42	43	33	32	27	29
21	42	53	32	35	29	36
22	45	48	26	29	34	38
GENDER						
Male	69	78	47	55	53	54
Female	65	61	45	47	44	42
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	85	89	61	69	57	59
White	77	81	57	60	52	55
Black or African American	23	26	11	16	20	19
American Indian or Alaska Native	4	*	*	*	4	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	18	24	14	22	11	10
Two or More Races	12	12	9	*	8	7
Hispanic or Latino	48	52	31	31	35	38
CURRENT EMPLOYMENT						
Full-Time	49	55	23	30	44	47
Part-Time	53	60	44	45	32	37
Unemployed	32	31	15	17	29	27
Other ³	46	50	39	44	23	23

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.92D Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.40	0.42	0.65	0.72	0.48	0.48
AGE						
18	0.71	0.81	1.21	1.71	0.88	0.92
19	0.81	0.82	1.11	1.22	1.17	1.10
20	0.90	0.92	1.52	1.44	1.08	1.12
21	0.92	1.07	1.60	1.64	1.10	1.30
22	0.91	0.95	1.78	1.84	1.02	1.09
GENDER						
Male	0.57	0.62	1.01	1.13	0.69	0.72
Female	0.55	0.51	0.83	0.87	0.68	0.65
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.44	0.46	0.69	0.78	0.54	0.55
White	0.55	0.58	0.86	0.97	0.70	0.72
Black or African American	0.70	0.74	0.99	1.40	0.92	0.88
American Indian or Alaska Native	3.17	*	*	*	3.84	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.49	1.79	1.92	2.47	2.34	2.26
Two or More Races	2.34	2.23	4.10	*	2.53	2.30
Hispanic or Latino	0.96	1.02	1.91	1.80	1.01	1.14
CURRENT EMPLOYMENT						
Full-Time	0.73	0.74	1.53	1.82	0.83	0.79
Part-Time	0.66	0.77	1.00	0.99	0.87	1.10
Unemployed	1.01	1.06	1.98	2.35	1.18	1.22
Other ³	0.72	0.86	1.04	1.41	0.87	0.87

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.92P Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.2336		0.7538		0.1626	
AGE						
18	0.2263		0.5804		0.0345	
19	0.0358		0.3271		0.0503	
20	0.8850		0.6453		0.7234	
21	0.3244		0.9157		0.2136	
22	0.3951		0.9160		0.3170	
GENDER						
Male	0.2955		0.5912		0.3554	
Female	0.4767		0.9302		0.2645	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2322		0.9609		0.1028	
White	0.1233		0.4599		0.1630	
Black or African American	0.9585		0.1413		0.3588	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.3664		0.2157		0.6499	
Two or More Races	0.3523		*		0.4901	
Hispanic or Latino	0.7513		0.5505		0.9961	
CURRENT EMPLOYMENT						
Full-Time	0.4523		0.7668		0.3005	
Part-Time	0.2047		0.4030		0.2387	
Unemployed	0.6438		0.5777		0.8369	
Other ³	0.4628		0.8358		0.3158	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.93C Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	117	120	82	85	81	83
AGE						
18	48	52	30	35	38	38
19	46	44	32	29	32	33
20	48	49	36	37	32	33
21	51	57	39	37	34	42
22	51	55	27	30	39	44
GENDER						
Male	85	92	57	59	65	69
Female	71	68	50	51	49	48
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	103	102	71	78	71	69
White	93	91	67	67	61	62
Black or African American	30	37	14	20	27	28
American Indian or Alaska Native	5	7	*	*	5	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	24	30	15	24	19	15
Two or More Races	16	15	9	11	13	10
Hispanic or Latino	54	58	34	32	40	45
CURRENT EMPLOYMENT						
Full-Time	55	65	25	31	50	57
Part-Time	63	66	50	50	38	41
Unemployed	44	39	21	20	38	35
Other ³	55	58	44	49	31	30

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.93D Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.46	0.47	0.71	0.77	0.56	0.58
AGE						
18	0.86	1.02	1.34	1.79	1.13	1.22
19	0.98	0.98	1.35	1.35	1.38	1.42
20	1.00	1.04	1.60	1.65	1.25	1.30
21	1.06	1.10	1.82	1.68	1.24	1.44
22	1.02	1.06	1.88	1.88	1.16	1.25
GENDER						
Male	0.67	0.72	1.14	1.17	0.83	0.90
Female	0.60	0.57	0.91	0.95	0.75	0.75
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.51	0.51	0.75	0.85	0.66	0.64
White	0.63	0.62	0.93	1.05	0.81	0.80
Black or African American	0.90	1.05	1.25	1.72	1.18	1.22
American Indian or Alaska Native	3.95	4.48	*	*	4.72	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.02	2.24	2.05	2.65	3.87	3.48
Two or More Races	3.01	2.67	4.21	4.52	3.94	3.15
Hispanic or Latino	1.05	1.14	2.08	1.83	1.13	1.35
CURRENT EMPLOYMENT						
Full-Time	0.80	0.86	1.66	1.90	0.91	0.94
Part-Time	0.75	0.83	1.09	1.07	1.02	1.16
Unemployed	1.33	1.32	2.66	2.75	1.48	1.55
Other ³	0.85	0.96	1.16	1.54	1.11	1.10

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.93P Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total ¹		COLLEGE ENROLLMENT STATUS			
			Full-Time College Students		Other Persons Aged 18-22 ²	
	2013	2014	2013	2014	2013	2014
TOTAL	0.1435		0.4079		0.2052	
AGE						
18	0.4665		0.5078		0.1802	
19	0.1137		0.3662		0.1729	
20	0.4307		0.6792		0.4909	
21	0.8207		0.1149		0.2640	
22	0.7642		0.6095		0.4840	
GENDER						
Male	0.1889		0.2246		0.4754	
Female	0.3867		0.9663		0.2094	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1364		0.7496		0.0827	
White	0.0796		0.1740		0.2554	
Black or African American	0.8927		0.1382		0.4557	
American Indian or Alaska Native	0.8847		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7811		0.1153		0.1748	
Two or More Races	0.7233		0.8696		0.5387	
Hispanic or Latino	0.7255		0.2395		0.6912	
CURRENT EMPLOYMENT						
Full-Time	0.5583		0.7602		0.6144	
Part-Time	0.1288		0.4331		0.1200	
Unemployed	0.5528		0.5221		0.7602	
Other ³	0.7019		0.9086		0.6521	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.94C Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2013 and 2014

Age Category	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	304	270	587	471	605	485
12-20	74	80	73	74	88	92
12	1	4	0	3	1	4
13	5	3	1	1	5	3
14	4	6	5	4	6	6
15	7	12	6	10	8	12
16	13	18	12	16	15	20
17	24	25	22	22	27	29
18	35	37	33	36	41	43
19	39	38	38	37	43	44
20	37	48	40	44	46	54
21-25	89	97	134	129	143	138
21	41	45	51	57	56	59
22	38	46	54	59	57	64
23	37	39	48	53	52	55
24	36	35	49	53	52	57
25	33	40	55	49	56	56
26-29	126	92	208	133	216	141
30-34	118	82	207	140	209	145
35-39	87	78	181	132	186	136
40-44	77	63	184	129	193	131
45-49	54	66	160	127	161	134
50-54	128	92	262	202	277	209
55-59	80	88	183	182	189	192
60-64	46	71	178	155	179	159
65 or Older	92	36	185	164	190	165

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.94D Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year, by Detailed Age Category: Standard Errors of Percentages, 2013 and 2014

Age Category	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	0.12	0.10	0.22	0.18	0.23	0.18
12-20	0.18	0.20	0.18	0.19	0.21	0.23
12	0.03	0.11	0.01	0.08	0.03	0.11
13	0.12	0.06	0.02	0.03	0.12	0.06
14	0.10	0.14	0.10	0.09	0.13	0.14
15	0.18	0.27	0.14	0.22	0.19	0.28
16	0.30	0.40	0.28	0.35	0.35	0.45
17	0.56	0.59	0.51	0.52	0.62	0.67
18	0.67	0.76	0.62	0.76	0.76	0.89
19	0.85	0.88	0.83	0.83	0.93	0.99
20	0.80	0.99	0.84	0.92	0.93	1.08
21-25	0.38	0.40	0.48	0.50	0.51	0.52
21	0.88	0.94	1.01	1.13	1.11	1.14
22	0.82	0.92	1.02	1.13	1.07	1.19
23	0.80	0.83	0.99	1.11	1.07	1.12
24	0.76	0.78	0.99	1.05	1.04	1.12
25	0.77	0.87	1.10	1.03	1.11	1.14
26-29	0.68	0.51	1.04	0.69	1.05	0.72
30-34	0.56	0.39	0.91	0.60	0.92	0.62
35-39	0.45	0.39	0.87	0.61	0.89	0.63
40-44	0.35	0.31	0.79	0.60	0.82	0.60
45-49	0.27	0.32	0.74	0.56	0.75	0.59
50-54	0.54	0.38	1.05	0.81	1.10	0.83
55-59	0.39	0.42	0.85	0.82	0.88	0.87
60-64	0.26	0.40	0.92	0.83	0.93	0.85
65 or Older	0.21	0.08	0.43	0.36	0.44	0.37

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.94P Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Age Category	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	0.6529		0.1020		0.2764	
12-20	0.6641		0.2251		0.3896	
12	0.2212		0.3323		0.2085	
13	0.5765		0.6309		0.5392	
14	0.9333		0.2009		0.5282	
15	0.2899		0.1667		0.2620	
16	0.4966		0.7737		0.6085	
17	0.6052		0.4733		0.4810	
18	0.2718		0.5226		0.3354	
19	0.1626		0.0365		0.1125	
20	0.2813		0.8642		0.5927	
21-25	0.6552		0.3071		0.5511	
21	0.7694		0.9332		0.7806	
22	0.4058		0.7809		0.8885	
23	0.3815		0.6469		0.7024	
24	0.1981		0.7370		0.8099	
25	0.4186		0.1604		0.5234	
26-29	0.8750		0.0154		0.0393	
30-34	0.2082		0.0290		0.0581	
35-39	0.4408		0.4052		0.4954	
40-44	0.9958		0.4681		0.4370	
45-49	0.0012		0.5112		0.8461	
50-54	0.5660		0.4366		0.3431	
55-59	0.1611		0.0189		0.0132	
60-64	0.2567		0.9529		0.6491	
65 or Older	0.1848		0.4637		0.6075	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.95C Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	304	270	587	471	605	485
AGE						
12-17	30	33	26	28	33	37
18-25	104	114	126	131	134	137
26 or Older	282	229	563	446	580	458
GENDER						
Male	246	224	454	371	469	384
Female	172	127	345	261	356	270
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	280	249	526	450	539	465
White	253	214	489	402	504	414
Black or African American	85	94	144	147	148	147
American Indian or Alaska Native	9	12	20	22	20	23
Native Hawaiian or Other Pacific Islander	6	17	31	44	31	46
Asian	29	43	116	85	118	90
Two or More Races	84	47	95	62	98	67
Hispanic or Latino	117	98	219	163	229	167

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.95D Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	0.12	0.10	0.22	0.18	0.23	0.18
AGE						
12-17	0.12	0.13	0.11	0.11	0.13	0.15
18-25	0.30	0.33	0.36	0.38	0.39	0.39
26 or Older	0.14	0.11	0.28	0.22	0.29	0.22
GENDER						
Male	0.19	0.17	0.36	0.29	0.37	0.30
Female	0.13	0.09	0.26	0.19	0.26	0.20
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.13	0.11	0.24	0.20	0.24	0.21
White	0.15	0.13	0.29	0.24	0.30	0.24
Black or African American	0.27	0.30	0.46	0.47	0.47	0.46
American Indian or Alaska Native	0.64	0.86	1.39	1.31	1.42	1.38
Native Hawaiian or Other Pacific Islander	0.69	1.67	3.34	4.01	3.33	4.06
Asian	0.22	0.32	0.84	0.62	0.86	0.66
Two or More Races	1.86	1.04	2.04	1.30	2.10	1.40
Hispanic or Latino	0.28	0.23	0.53	0.38	0.55	0.39

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.95P Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 12 or Older, by Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	0.6529		0.1020		0.2764	
AGE						
12-17	0.6505		0.9954		0.9050	
18-25	0.9612		0.1549		0.3308	
26 or Older	0.6129		0.1692		0.3626	
GENDER						
Male	0.5627		0.4073		0.8003	
Female	0.9666		0.0858		0.1232	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.5681		0.1962		0.5169	
White	0.6913		0.0806		0.1919	
Black or African American	0.0119		0.1407		0.0366	
American Indian or Alaska Native	0.4036		0.4179		0.3359	
Native Hawaiian or Other Pacific Islander	0.0160		0.6855		0.4460	
Asian	0.1643		0.7904		0.9613	
Two or More Races	0.3311		0.6226		0.6107	
Hispanic or Latino	0.8725		0.2559		0.2130	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.96C Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	304	266	582	466	600	479
GENDER						
Male	246	221	453	369	468	382
Female	172	125	343	260	355	268
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	279	246	523	445	536	460
White	259	213	566	433	588	451
Black or African American	87	98	155	158	159	160
American Indian or Alaska Native	9	12	20	22	20	23
Native Hawaiian or Other Pacific Islander	5	17	31	*	31	46
Asian	29	42	116	85	118	89
Two or More Races	84	47	95	62	98	66
Hispanic or Latino	116	97	218	162	228	165
EDUCATION						
< High School	82	82	113	107	128	119
High School Graduate	161	147	265	229	287	244
Some College	154	148	328	246	337	254
College Graduate	189	133	472	322	482	329
CURRENT EMPLOYMENT						
Full-Time	231	205	558	410	575	421
Part-Time	144	117	232	192	245	199
Unemployed	82	56	98	79	107	84
Other ¹	113	98	207	192	221	203

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.96D Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	0.13	0.11	0.25	0.19	0.25	0.20
GENDER						
Male	0.22	0.19	0.40	0.32	0.41	0.33
Female	0.14	0.10	0.28	0.21	0.29	0.22
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.14	0.12	0.26	0.22	0.27	0.23
White	0.16	0.13	0.31	0.25	0.32	0.26
Black or African American	0.31	0.33	0.52	0.52	0.53	0.52
American Indian or Alaska Native	0.70	0.96	1.58	1.45	1.62	1.53
Native Hawaiian or Other Pacific Islander	0.73	1.86	3.76	*	3.76	4.48
Asian	0.25	0.34	0.93	0.68	0.94	0.72
Two or More Races	2.25	1.26	2.47	1.56	2.55	1.68
Hispanic or Latino	0.33	0.26	0.61	0.44	0.64	0.45
EDUCATION						
< High School	0.26	0.26	0.35	0.32	0.39	0.36
High School Graduate	0.23	0.21	0.37	0.32	0.40	0.34
Some College	0.24	0.22	0.48	0.35	0.49	0.36
College Graduate	0.26	0.18	0.56	0.39	0.57	0.39
CURRENT EMPLOYMENT						
Full-Time	0.19	0.16	0.41	0.29	0.42	0.30
Part-Time	0.41	0.34	0.64	0.54	0.67	0.55
Unemployed	0.68	0.52	0.81	0.69	0.88	0.73
Other ¹	0.15	0.13	0.28	0.25	0.30	0.27

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.96P Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	0.6973		0.0915		0.2523	
GENDER						
Male	0.5479		0.3856		0.7919	
Female	0.8438		0.0812		0.1057	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6324		0.1806		0.4801	
White	0.6481		0.0747		0.1817	
Black or African American	0.0172		0.1405		0.0435	
American Indian or Alaska Native	0.2847		0.4140		0.2713	
Native Hawaiian or Other Pacific Islander	0.0130		*		0.4542	
Asian	0.1991		0.7297		0.9020	
Two or More Races	0.3302		0.6460		0.6077	
Hispanic or Latino	0.9186		0.2445		0.2091	
EDUCATION						
< High School	0.6741		0.6049		0.6497	
High School Graduate	0.7231		0.7042		0.9901	
Some College	0.4366		0.7281		0.8992	
College Graduate	0.8396		0.0154		0.0264	
CURRENT EMPLOYMENT						
Full-Time	0.3271		0.0728		0.2130	
Part-Time	0.7563		0.3303		0.3664	
Unemployed	0.2654		0.9254		0.7178	
Other ¹	0.6956		0.9055		0.7932	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.97C Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Geographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	304	266	582	466	600	479
GEOGRAPHIC DIVISION						
Northeast	103	117	215	204	224	216
New England	63	69	111	121	115	135
Middle Atlantic	83	96	184	167	194	172
Midwest	141	111	225	224	238	231
East North Central	91	92	181	193	185	199
West North Central	109	58	131	106	153	109
South	174	165	392	308	401	311
South Atlantic	140	114	301	226	312	228
East South Central	61	60	121	112	125	116
West South Central	84	99	202	166	205	169
West	175	127	322	234	333	238
Mountain	78	62	168	118	174	121
Pacific	156	114	275	202	282	207
COUNTY TYPE						
Large Metro	231	217	537	429	558	441
Small Metro	174	146	348	277	369	296
250K – 1 Mil. Pop.	148	108	324	236	341	247
< 250K Pop.	96	104	186	166	199	184
Nonmetro	121	83	202	179	214	187
Urbanized	96	59	154	123	160	129
Less Urbanized	63	56	127	129	135	134
Completely Rural	37	20	59	47	67	51

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.97D Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Geographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	0.13	0.11	0.25	0.19	0.25	0.20
GEOGRAPHIC DIVISION						
Northeast	0.24	0.27	0.50	0.47	0.52	0.50
New England	0.55	0.60	0.98	1.06	1.01	1.18
Middle Atlantic	0.26	0.30	0.58	0.52	0.61	0.54
Midwest	0.28	0.22	0.44	0.44	0.47	0.45
East North Central	0.26	0.26	0.52	0.55	0.53	0.56
West North Central	0.70	0.37	0.84	0.68	0.98	0.69
South	0.20	0.18	0.44	0.34	0.45	0.35
South Atlantic	0.30	0.24	0.64	0.48	0.67	0.48
East South Central	0.43	0.42	0.87	0.79	0.89	0.82
West South Central	0.31	0.36	0.74	0.60	0.75	0.61
West	0.32	0.23	0.58	0.42	0.60	0.42
Mountain	0.46	0.36	1.00	0.69	1.03	0.71
Pacific	0.40	0.29	0.71	0.52	0.73	0.53
COUNTY TYPE						
Large Metro	0.17	0.16	0.36	0.29	0.37	0.29
Small Metro	0.24	0.20	0.44	0.36	0.47	0.38
250K – 1 Mil. Pop.	0.30	0.22	0.59	0.45	0.62	0.46
< 250K Pop.	0.41	0.43	0.72	0.65	0.76	0.72
Nonmetro	0.30	0.21	0.47	0.40	0.49	0.41
Urbanized	0.52	0.34	0.75	0.65	0.76	0.67
Less Urbanized	0.36	0.29	0.66	0.58	0.71	0.59
Completely Rural	0.87	0.47	1.19	1.13	1.36	1.22

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.97P Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Persons Aged 18 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Geographic Characteristic	Illicit Drugs (2013)	Illicit Drugs (2014)	Alcohol (2013)	Alcohol (2014)	Illicit Drugs or Alcohol (2013)	Illicit Drugs or Alcohol (2014)
TOTAL	0.6973		0.0915		0.2523	
GEOGRAPHIC DIVISION						
Northeast	0.1647		0.8214		0.9494	
New England	0.8081		0.5887		0.4068	
Middle Atlantic	0.1264		0.5302		0.6222	
Midwest	0.9601		0.9510		0.9612	
East North Central	0.7336		0.6737		0.4772	
West North Central	0.7750		0.4360		0.3446	
South	0.4877		0.5073		0.8076	
South Atlantic	0.9243		0.6466		0.8609	
East South Central	0.6742		0.2410		0.1721	
West South Central	0.3830		0.1759		0.2748	
West	0.2786		0.0368		0.0565	
Mountain	0.3863		0.0334		0.0374	
Pacific	0.4450		0.2482		0.3333	
COUNTY TYPE						
Large Metro	0.5022		0.3344		0.5483	
Small Metro	0.6986		0.0237		0.0832	
250K – 1 Mil. Pop.	0.3963		0.0140		0.0387	
< 250K Pop.	0.6369		0.8348		0.9236	
Nonmetro	0.8283		0.4128		0.4443	
Urbanized	0.8362		0.8722		0.8932	
Less Urbanized	0.6076		0.0802		0.1785	
Completely Rural	0.6114		0.6641		0.7034	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.98C Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
ILLCIT DRUGS²	503	462	114	88	516	474
Marijuana and Hashish	437	414	90	76	453	422
Cocaine	132	105	33	26	130	102
Crack	64	55	23	18	60	54
Heroin	42	55	20	21	37	50
Hallucinogens	131	84	34	20	126	81
LSD	36	38	4	16	36	34
PCP	15	*	11	2	10	*
Ecstasy	91	63	28	17	86	61
Inhalants	61	61	12	4	60	61
Nonmedical Use of Psychotherapeutics ^{3,4}	255	217	70	58	243	213
Pain Relievers	205	170	53	44	198	167
OxyContin [®]	80	49	24	8	76	49
Tranquilizers	120	113	41	37	113	107
Stimulants ⁴	116	114	39	45	101	103
Methamphetamine ⁴	85	77	36	35	67	63
Sedatives	66	53	44	24	50	47
ILLCIT DRUGS OTHER THAN MARIJUANA²	305	255	75	63	294	248

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 or older, including those with unknown probation status in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.98D Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status: Standard Errors of Percentages, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
ILLCIT DRUGS²	0.21	0.19	2.09	1.65	0.21	0.19
Marijuana and Hashish	0.18	0.17	1.80	1.48	0.19	0.17
Cocaine	0.06	0.04	0.73	0.59	0.06	0.04
Crack	0.03	0.02	0.50	0.40	0.03	0.02
Heroin	0.02	0.02	0.45	0.48	0.02	0.02
Hallucinogens	0.06	0.04	0.76	0.46	0.05	0.03
LSD	0.02	0.02	0.09	0.37	0.02	0.01
PCP	0.01	*	0.24	0.04	0.00	*
Ecstasy	0.04	0.03	0.63	0.39	0.04	0.03
Inhalants	0.03	0.03	0.26	0.09	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{3,4}	0.11	0.09	1.44	1.24	0.10	0.09
Pain Relievers	0.09	0.07	1.13	0.97	0.08	0.07
OxyContin [®]	0.03	0.02	0.53	0.18	0.03	0.02
Tranquilizers	0.05	0.05	0.89	0.81	0.05	0.05
Stimulants ⁴	0.05	0.05	0.86	0.99	0.04	0.04
Methamphetamine ⁴	0.04	0.03	0.81	0.77	0.03	0.03
Sedatives	0.03	0.02	0.97	0.54	0.02	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.13	0.11	1.52	1.33	0.12	0.10

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 or older, including those with unknown probation status in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.98P Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Total ¹ (2013)	Total ¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
ILLCIT DRUGS²	0.0036		0.3956		0.0014	
Marijuana and Hashish	0.0006		0.5805		0.0003	
Cocaine	0.8494		0.6368		0.9517	
Crack	0.7016		0.9084		0.7418	
Heroin	0.0454		0.3919		0.0509	
Hallucinogens	0.3199		0.4773		0.3838	
LSD	0.4220		0.0726		0.8030	
PCP	*		0.4518		*	
Ecstasy	0.6653		0.6828		0.7682	
Inhalants	0.8427		0.1826		0.7511	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.7175		0.3986		0.9571	
Pain Relievers	0.2866		0.2362		0.4674	
OxyContin [®]	0.1227		0.0337		0.3115	
Tranquilizers	0.4051		0.5169		0.2547	
Stimulants ⁴	0.4230		0.8676		0.2559	
Methamphetamine ⁴	0.6325		0.6296		0.9067	
Sedatives	0.4819		0.4263		0.1494	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.6814		0.8267		0.8082	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 or older, including those with unknown probation status in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.99C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Probation Status: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
ILLCIT DRUGS²	378	322	95	77	405	338
Marijuana and Hashish	332	301	81	68	359	311
Cocaine	109	87	24	26	109	82
Crack	39	40	16	17	35	36
Heroin	37	45	20	21	31	39
Hallucinogens	85	78	32	20	80	74
LSD	36	38	4	16	36	34
PCP	15	4	11	2	10	*
Ecstasy	69	59	28	17	63	57
Inhalants	54	55	12	4	53	54
Nonmedical Use of Psychotherapeutics ^{3,4}	207	159	55	46	203	157
Pain Relievers	173	118	51	33	166	116
OxyContin [®]	50	33	20	8	45	32
Tranquilizers	104	89	38	27	98	85
Stimulants ⁴	102	78	39	31	94	72
Methamphetamine ⁴	69	48	36	26	56	39
Sedatives	49	31	16	2	46	30
ILLCIT DRUGS OTHER THAN MARIJUANA²	236	191	58	52	235	189

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 49, including those with unknown probation status in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.99D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Probation Status: Standard Errors of Percentages, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
ILLCIT DRUGS²	0.29	0.24	2.04	1.70	0.28	0.24
Marijuana and Hashish	0.25	0.23	1.85	1.55	0.25	0.22
Cocaine	0.08	0.06	0.64	0.69	0.08	0.06
Crack	0.03	0.03	0.42	0.46	0.03	0.03
Heroin	0.03	0.03	0.54	0.57	0.02	0.03
Hallucinogens	0.06	0.06	0.83	0.54	0.06	0.06
LSD	0.03	0.03	0.10	0.44	0.03	0.03
PCP	0.01	0.00	0.29	0.05	0.01	*
Ecstasy	0.05	0.04	0.74	0.47	0.05	0.04
Inhalants	0.04	0.04	0.30	0.10	0.04	0.04
Nonmedical Use of Psychotherapeutics ^{3,4}	0.16	0.12	1.35	1.17	0.15	0.12
Pain Relievers	0.13	0.09	1.27	0.88	0.13	0.09
OxyContin [®]	0.04	0.02	0.53	0.22	0.03	0.02
Tranquilizers	0.08	0.07	0.98	0.72	0.08	0.07
Stimulants ⁴	0.08	0.06	1.01	0.82	0.07	0.05
Methamphetamine ⁴	0.05	0.04	0.96	0.70	0.04	0.03
Sedatives	0.04	0.02	0.43	0.06	0.04	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.18	0.14	1.43	1.30	0.18	0.14

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 49, including those with unknown probation status in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.99P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Probation Status: *P* Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
ILLCIT DRUGS²	0.0475		0.5248		0.0272	
Marijuana and Hashish	0.0078		0.6508		0.0045	
Cocaine	0.7096		0.4670		0.8381	
Crack	0.5993		0.4063		0.8202	
Heroin	0.1004		0.4178		0.1089	
Hallucinogens	0.7416		0.6608		0.8132	
LSD	0.4221		0.0720		0.8047	
PCP	0.2466		0.4515		*	
Ecstasy	0.7896		0.6833		0.9289	
Inhalants	0.8121		0.1822		0.7086	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.2648		0.1432		0.5071	
Pain Relievers	0.0611		0.0448		0.1912	
OxyContin [®]	0.1344		0.0612		0.3781	
Tranquilizers	0.7780		0.2984		0.4849	
Stimulants ⁴	0.5302		0.3059		0.8336	
Methamphetamine ⁴	0.1633		0.1793		0.4445	
Sedatives	0.9383		0.1450		0.6288	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.6410		0.7996		0.7024	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 49, including those with unknown probation status in the past year.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.100C Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	503	462	114	88	516	474
AGE						
18-49	378	322	95	77	405	338
18-25	143	160	43	43	164	187
26-34	254	184	63	48	277	193
35-49	220	192	55	45	218	195
GENDER/AGE						
Male 18 or Older	356	351	105	80	351	361
Male 18-49	254	246	87	68	273	259
Female 18 or Older	342	263	48	40	355	267
Female 18-49	252	189	46	38	267	197

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.100D Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	0.21	0.19	2.09	1.65	0.21	0.19
AGE						
18-49	0.29	0.24	2.04	1.70	0.28	0.24
18-25	0.41	0.46	2.15	2.46	0.41	0.45
26-34	0.68	0.49	4.53	3.07	0.68	0.48
35-49	0.36	0.32	4.04	3.53	0.36	0.31
GENDER/AGE						
Male 18 or Older	0.31	0.30	2.67	2.16	0.31	0.31
Male 18-49	0.39	0.37	2.60	2.19	0.39	0.38
Female 18 or Older	0.28	0.21	3.38	2.56	0.28	0.21
Female 18-49	0.38	0.28	3.52	2.68	0.38	0.28

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

**Table 6.100P Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics:
P Values from Tests of Differences of Percentages, 2014 Versus 2013**

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	0.0036		0.3956		0.0014	
AGE						
18-49	0.0475		0.5248		0.0272	
18-25	0.4747		0.9739		0.3365	
26-34	0.8673		0.7523		0.8441	
35-49	0.0035		0.6034		0.0017	
GENDER/AGE						
Male 18 or Older	0.0016		0.8220		0.0005	
Male 18-49	0.0403		0.8979		0.0237	
Female 18 or Older	0.3382		0.2224		0.3164	
Female 18-49	0.4089		0.1855		0.3668	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.101C Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	235	215	58	67	229	206
AGE						
18-49	183	168	55	56	179	160
18-25	88	92	35	27	83	90
26-34	130	97	37	36	129	92
35-49	99	92	19	33	97	89
GENDER/AGE						
Male 18 or Older	168	170	55	61	161	163
Male 18-49	147	135	51	49	144	132
Female 18 or Older	169	122	33	28	169	119
Female 18-49	113	87	33	27	110	83

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.101D Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	0.10	0.09	1.29	1.40	0.10	0.09
AGE						
18-49	0.14	0.13	1.39	1.41	0.14	0.12
18-25	0.25	0.26	2.03	1.84	0.24	0.26
26-34	0.35	0.26	3.03	2.59	0.35	0.24
35-49	0.16	0.15	1.63	2.74	0.16	0.15
GENDER/AGE						
Male 18 or Older	0.15	0.15	1.65	1.84	0.14	0.14
Male 18-49	0.22	0.21	1.79	1.82	0.22	0.20
Female 18 or Older	0.14	0.10	2.57	1.92	0.14	0.10
Female 18-49	0.17	0.13	2.85	2.12	0.16	0.12

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.101P Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	0.6075		0.7156		0.6007	
AGE						
18-49	0.6704		0.7046		0.8130	
18-25	0.0515		0.0239		0.3007	
26-34	0.6986		0.8283		0.6474	
35-49	0.1522		0.0588		0.3408	
GENDER/AGE						
Male 18 or Older	0.6048		0.3857		0.6719	
Male 18-49	0.5718		0.8250		0.6311	
Female 18 or Older	0.8486		0.4255		0.7674	
Female 18-49	0.9572		0.3039		0.8081	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.102C Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	456	384	107	93	477	391
AGE						
18-49	345	281	99	82	367	297
18-25	138	134	44	39	156	149
26-34	235	177	59	49	259	188
35-49	213	171	64	51	216	177
GENDER/AGE						
Male 18 or Older	363	311	98	83	369	322
Male 18-49	270	231	91	70	291	245
Female 18 or Older	275	214	51	41	285	215
Female 18-49	204	154	48	40	215	160

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.102D Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	0.19	0.16	2.11	1.76	0.19	0.16
AGE						
18-49	0.26	0.21	2.14	1.84	0.26	0.21
18-25	0.40	0.38	2.22	2.45	0.38	0.38
26-34	0.63	0.47	4.20	3.25	0.64	0.47
35-49	0.35	0.28	4.67	3.62	0.34	0.28
GENDER/AGE						
Male 18 or Older	0.32	0.27	2.60	2.23	0.32	0.27
Male 18-49	0.41	0.35	2.63	2.29	0.41	0.35
Female 18 or Older	0.22	0.17	3.58	2.67	0.22	0.17
Female 18-49	0.30	0.23	3.67	2.82	0.30	0.23

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.102P Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	0.6604		0.2073		0.9603	
AGE						
18-49	0.1163		0.0818		0.3006	
18-25	0.0749		0.0837		0.2662	
26-34	0.0758		0.1661		0.1054	
35-49	0.6004		0.8480		0.4411	
GENDER/AGE						
Male 18 or Older	0.8026		0.3834		0.8540	
Male 18-49	0.1648		0.1210		0.4725	
Female 18 or Older	0.6823		0.3845		0.7197	
Female 18-49	0.4243		0.5198		0.4096	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.103C Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
ILLCIT DRUGS³	503	462	64	46	524	487
Marijuana and Hashish	437	414	57	42	456	432
Cocaine	132	105	34	16	133	103
Crack	64	55	9	12	63	54
Heroin	42	55	10	15	41	53
Hallucinogens	131	84	28	8	127	83
LSD	36	38	3	4	36	38
PCP	15	*	*	1	15	*
Ecstasy	91	63	*	7	86	63
Inhalants	61	61	11	0	60	61
Nonmedical Use of Psychotherapeutics ^{4,5}	255	217	44	27	253	220
Pain Relievers	205	170	32	21	204	171
OxyContin [®]	80	49	14	2	78	49
Tranquilizers	120	113	23	20	119	111
Stimulants ⁵	116	114	32	16	105	113
Methamphetamine ⁵	85	77	31	8	71	77
Sedatives	66	53	2	3	67	53
ILLCIT DRUGS OTHER THAN MARIJUANA³	305	255	50	29	304	261

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 or older, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.103D Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status: Standard Errors of Percentages, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
ILLCIT DRUGS³	0.21	0.19	3.40	2.74	0.21	0.19
Marijuana and Hashish	0.18	0.17	3.06	2.55	0.18	0.17
Cocaine	0.06	0.04	1.97	1.12	0.06	0.04
Crack	0.03	0.02	0.57	0.80	0.03	0.02
Heroin	0.02	0.02	0.58	1.04	0.02	0.02
Hallucinogens	0.06	0.04	1.64	0.54	0.05	0.03
LSD	0.02	0.02	0.19	0.26	0.02	0.02
PCP	0.01	*	*	0.09	0.01	*
Ecstasy	0.04	0.03	*	0.46	0.04	0.03
Inhalants	0.03	0.03	0.65	0.03	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{4,5}	0.11	0.09	2.50	1.84	0.11	0.09
Pain Relievers	0.09	0.07	1.85	1.40	0.09	0.07
OxyContin [®]	0.03	0.02	0.84	0.12	0.03	0.02
Tranquilizers	0.05	0.05	1.38	1.34	0.05	0.05
Stimulants ⁵	0.05	0.05	1.85	1.08	0.04	0.05
Methamphetamine ⁵	0.04	0.03	1.83	0.59	0.03	0.03
Sedatives	0.03	0.02	0.14	0.18	0.03	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA³	0.13	0.11	2.78	1.92	0.13	0.11

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 or older, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.103P Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
ILLCIT DRUGS³	0.0036		0.3159		0.0022	
Marijuana and Hashish	0.0006		0.6127		0.0005	
Cocaine	0.8494		0.3440		0.9635	
Crack	0.7016		0.6083		0.6592	
Heroin	0.0454		0.5926		0.0413	
Hallucinogens	0.3199		0.2835		0.4010	
LSD	0.4220		0.7621		0.4315	
PCP	*		*		*	
Ecstasy	0.6653		*		0.8799	
Inhalants	0.8427		0.1963		0.7729	
Nonmedical Use of Psychotherapeutics ^{4,5}	0.7175		0.3155		0.9591	
Pain Relievers	0.2866		0.2153		0.3703	
OxyContin [®]	0.1227		0.1315		0.1807	
Tranquilizers	0.4051		1.0000		0.3564	
Stimulants ⁵	0.4230		0.4553		0.1939	
Methamphetamine ⁵	0.6325		0.1152		0.7279	
Sedatives	0.4819		0.6278		0.5059	
ILLCIT DRUGS OTHER THAN MARIJUANA³	0.6814		0.1051		0.9271	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 or older, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.104C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Parole/Supervised Release Status: Standard Errors of Numbers in Thousands, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
ILLCIT DRUGS³	378	322	56	41	416	351
Marijuana and Hashish	332	301	49	37	363	323
Cocaine	109	87	29	14	112	86
Crack	39	40	5	7	38	39
Heroin	37	45	10	12	35	43
Hallucinogens	85	78	28	8	82	77
LSD	36	38	3	4	36	38
PCP	15	4	*	1	15	3
Ecstasy	69	59	*	7	63	59
Inhalants	54	55	11	0	53	54
Nonmedical Use of Psychotherapeutics ^{4,5}	207	159	43	26	210	164
Pain Relievers	173	118	31	19	172	120
OxyContin [®]	50	33	14	2	47	33
Tranquilizers	104	89	22	20	104	87
Stimulants ⁵	102	78	32	16	98	77
Methamphetamine ⁵	69	48	31	8	60	47
Sedatives	49	31	2	2	49	30
ILLCIT DRUGS OTHER THAN MARIJUANA³	236	191	46	27	243	198

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 49, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.104D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Parole/Supervised Release Status: Standard Errors of Percentages, 2013 and 2014

Drug	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
ILLCIT DRUGS³	0.29	0.24	3.56	3.04	0.29	0.24
Marijuana and Hashish	0.25	0.23	3.21	2.80	0.25	0.23
Cocaine	0.08	0.06	2.09	1.16	0.08	0.06
Crack	0.03	0.03	0.42	0.62	0.03	0.03
Heroin	0.03	0.03	0.73	1.04	0.03	0.03
Hallucinogens	0.06	0.06	2.03	0.66	0.06	0.06
LSD	0.03	0.03	0.24	0.33	0.03	0.03
PCP	0.01	0.00	*	0.12	0.01	0.00
Ecstasy	0.05	0.04	*	0.57	0.05	0.04
Inhalants	0.04	0.04	0.81	0.04	0.04	0.04
Nonmedical Use of Psychotherapeutics ^{4,5}	0.16	0.12	2.96	2.10	0.16	0.12
Pain Relievers	0.13	0.09	2.19	1.55	0.13	0.09
OxyContin [®]	0.04	0.02	1.05	0.15	0.04	0.02
Tranquilizers	0.08	0.07	1.61	1.64	0.08	0.07
Stimulants ⁵	0.08	0.06	2.28	1.33	0.07	0.06
Methamphetamine ⁵	0.05	0.04	2.27	0.72	0.05	0.04
Sedatives	0.04	0.02	0.18	0.20	0.04	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA³	0.18	0.14	3.12	2.20	0.18	0.14

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 49, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.104P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 49, by Past Year Parole/Supervised Release Status: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Drug	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
ILLCIT DRUGS³	0.0475		0.5561		0.0381	
Marijuana and Hashish	0.0078		0.7799		0.0074	
Cocaine	0.7096		0.6118		0.6129	
Crack	0.5993		0.5866		0.6299	
Heroin	0.1004		0.9215		0.0632	
Hallucinogens	0.7416		0.2572		0.9294	
LSD	0.4221		0.7661		0.4315	
PCP	0.2466		*		0.2105	
Ecstasy	0.7896		*		0.9300	
Inhalants	0.8121		0.1949		0.7331	
Nonmedical Use of Psychotherapeutics ^{4,5}	0.2648		0.2516		0.4208	
Pain Relievers	0.0611		0.1524		0.1101	
OxyContin [®]	0.1344		0.1291		0.2353	
Tranquilizers	0.7780		0.8132		0.7464	
Stimulants ⁵	0.5302		0.4230		0.7170	
Methamphetamine ⁵	0.1633		0.1012		0.4329	
Sedatives	0.9383		0.8646		0.9682	
ILLCIT DRUGS OTHER THAN MARIJUANA³	0.6410		0.1446		0.8533	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all persons aged 18 to 49, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁵ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.105C Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	503	462	64	46	524	487
AGE						
18-49	378	322	56	41	416	351
18-25	143	160	20	22	170	191
26-34	254	184	*	29	288	197
35-49	220	192	*	*	224	201
GENDER/AGE						
Male 18 or Older	356	351	61	38	358	369
Male 18-49	254	246	53	31	281	269
Female 18 or Older	342	263	*	*	357	269
Female 18-49	252	189	*	*	269	200

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.105D Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	0.21	0.19	3.40	2.74	0.21	0.19
AGE						
18-49	0.29	0.24	3.56	3.04	0.29	0.24
18-25	0.41	0.46	3.93	5.24	0.41	0.45
26-34	0.68	0.49	*	5.56	0.69	0.48
35-49	0.36	0.32	*	*	0.36	0.32
GENDER/AGE						
Male 18 or Older	0.31	0.30	3.97	3.06	0.31	0.31
Male 18-49	0.39	0.37	4.22	3.25	0.39	0.38
Female 18 or Older	0.28	0.21	*	*	0.28	0.21
Female 18-49	0.38	0.28	*	*	0.38	0.28

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.105P Illicit Drug Use in the Past Month among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	0.0036		0.3159		0.0022	
AGE						
18-49	0.0475		0.5561		0.0381	
18-25	0.4747		0.8465		0.4240	
26-34	0.8673		*		0.7921	
35-49	0.0035		*		0.0018	
GENDER/AGE						
Male 18 or Older	0.0016		0.1719		0.0005	
Male 18-49	0.0403		0.1659		0.0188	
Female 18 or Older	0.3382		*		0.4039	
Female 18-49	0.4089		*		0.5353	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.106C Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	235	215	43	39	234	219
AGE						
18-49	183	168	43	34	184	168
18-25	88	92	16	12	90	94
26-34	130	97	*	25	132	96
35-49	99	92	*	*	98	91
GENDER/AGE						
Male 18 or Older	168	170	38	32	165	172
Male 18-49	147	135	38	30	147	137
Female 18 or Older	169	122	*	*	170	121
Female 18-49	113	87	*	*	113	87

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.106D Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	0.10	0.09	2.47	2.43	0.10	0.09
AGE						
18-49	0.14	0.13	2.94	2.68	0.14	0.12
18-25	0.25	0.26	3.53	3.72	0.25	0.26
26-34	0.35	0.26	*	5.07	0.35	0.25
35-49	0.16	0.15	*	*	0.16	0.15
GENDER/AGE						
Male 18 or Older	0.15	0.15	2.73	2.65	0.15	0.15
Male 18-49	0.22	0.21	3.31	3.06	0.22	0.20
Female 18 or Older	0.14	0.10	*	*	0.14	0.10
Female 18-49	0.17	0.13	*	*	0.17	0.13

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.106P Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	0.6075		0.6830		0.5844	
AGE						
18-49	0.6704		0.9092		0.7639	
18-25	0.0515		0.3423		0.0892	
26-34	0.6986		*		0.7007	
35-49	0.1522		*		0.1553	
GENDER/AGE						
Male 18 or Older	0.6048		0.6234		0.5819	
Male 18-49	0.5718		0.8584		0.6233	
Female 18 or Older	0.8486		*		0.8439	
Female 18-49	0.9572		*		0.8773	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.107C Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	456	384	70	54	487	401
AGE						
18-49	345	281	60	47	376	306
18-25	138	134	20	19	164	153
26-34	235	177	*	33	262	192
35-49	213	171	*	*	221	180
GENDER/AGE						
Male 18 or Older	363	311	62	46	378	331
Male 18-49	270	231	55	40	299	253
Female 18 or Older	275	214	*	*	288	217
Female 18-49	204	154	*	*	219	162

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.107D Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: Standard Errors of Percentages, 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	0.19	0.16	3.63	3.02	0.19	0.16
AGE						
18-49	0.26	0.21	3.64	3.28	0.26	0.21
18-25	0.40	0.38	3.94	5.04	0.39	0.38
26-34	0.63	0.47	*	5.79	0.63	0.47
35-49	0.35	0.28	*	*	0.35	0.28
GENDER/AGE						
Male 18 or Older	0.32	0.27	4.06	3.42	0.32	0.27
Male 18-49	0.41	0.35	4.13	3.72	0.41	0.35
Female 18 or Older	0.22	0.17	*	*	0.22	0.17
Female 18-49	0.30	0.23	*	*	0.30	0.23

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 6.107P Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2014 Versus 2013

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	0.6604		0.5865		0.7886	
AGE						
18-49	0.1163		0.6977		0.1654	
18-25	0.0749		0.9523		0.0998	
26-34	0.0758		*		0.0740	
35-49	0.6004		*		0.4569	
GENDER/AGE						
Male 18 or Older	0.8026		0.7091		0.9472	
Male 18-49	0.1648		0.5778		0.2578	
Female 18 or Older	0.6823		*		0.6975	
Female 18-49	0.4243		*		0.3941	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 7.1A Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	108,255 ^b	110,205 ^b	110,057 ^b	112,085 ^b	111,747 ^b	114,509 ^b	117,815 ^b	119,259 ^b	119,933 ^b	121,078 ^b	124,808 ^b	127,458 ^a	130,332
Marijuana and Hashish	94,946 ^b	96,611 ^b	96,772 ^b	97,545 ^b	97,792 ^b	100,737 ^b	102,857 ^b	104,950 ^b	106,613 ^b	107,842 ^b	111,239 ^b	114,712	117,213
Cocaine	33,910 ^b	34,891 ^b	34,153 ^b	33,673 ^b	35,303 ^b	35,947 ^b	36,916 ^a	36,742 ^b	37,361 ^a	36,921 ^a	37,688	37,634	39,200
Crack	8,402 ^a	7,949 ^b	7,840 ^b	7,928 ^b	8,559	8,615	8,454 ^a	8,390 ^a	9,208	8,214 ^b	9,015	8,870	9,424
Heroin	3,668 ^b	3,744 ^b	3,145 ^b	3,534 ^b	3,788 ^b	3,806 ^b	3,795 ^b	3,680 ^b	4,144 ^a	4,162 ^a	4,565	4,812	4,813
Hallucinogens	34,314 ^b	34,363 ^b	34,333 ^b	33,728 ^b	35,290 ^b	34,278 ^b	36,064 ^b	37,404 ^a	37,544 ^a	36,362 ^b	37,908	39,736	39,647
LSD	24,516	24,424	23,398 ^a	22,433 ^b	23,365 ^a	22,720 ^b	23,604	23,714	23,375 ^a	23,000 ^b	23,735	24,770	25,035
PCP	7,418 ^b	7,107	6,762	6,603	6,633	6,179	6,657	6,271	6,255	6,103	6,552	6,462	6,388
Ecstasy	10,150 ^b	10,904 ^b	11,130 ^b	11,495 ^b	12,282 ^b	12,418 ^b	12,964 ^b	14,280 ^b	15,929 ^b	14,570 ^b	16,162 ^a	17,815	17,548
Inhalants	22,870 ^a	22,995 ^b	22,798 ^a	22,745 ^a	22,856 ^a	22,558	22,430	22,552 ^a	21,778	20,523	21,095	21,068	21,293
Nonmedical Use of Psychotherapeutics ^{2,3}	47,958 ^b	49,001 ^b	49,157 ^b	49,571 ^b	50,983 ^b	50,541 ^b	52,154 ^a	51,991 ^a	51,832 ^a	51,243 ^b	54,389	53,172	54,395
Pain Relievers	29,611 ^b	31,207 ^b	31,768 ^b	32,692 ^b	33,478 ^b	33,132 ^b	34,992	35,197	34,908	34,247 ^a	37,045	35,473	36,064
OxyContin [®]	1,924 ^b	2,832 ^b	3,072 ^b	3,481 ^b	4,116 ^b	4,379 ^b	4,857 ^b	5,863 ^b	6,160 ^b	5,917 ^b	6,579	6,973	7,031
Tranquilizers	19,267 ^b	20,220 ^b	19,852 ^b	21,041 ^b	21,318 ^b	20,253 ^b	21,551 ^b	21,848 ^b	22,187 ^b	21,655 ^b	23,639	23,493	24,851
Stimulants ³	23,496	23,004	22,297	20,983 ^a	22,495	21,669	21,285	22,033	21,739	20,379 ^b	21,498	21,656	22,530
Methamphetamine ³	15,365 ^b	15,139 ^b	14,512 ^b	12,663	14,226 ^a	13,081	12,634	12,908	13,060	11,928	12,259	12,257	12,943
Sedatives	9,960 ^b	9,510 ^b	9,891 ^b	8,982 ^a	8,842 ^a	8,401	8,920 ^a	8,633	7,653	7,515	7,939	7,480	7,826
ILLCIT DRUGS OTHER THAN MARIJUANA¹	70,300 ^b	71,128 ^b	70,657 ^b	71,822 ^b	72,934 ^b	73,654 ^b	75,907 ^b	76,080 ^b	76,472 ^b	75,447 ^b	78,034	78,076	80,119

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.1B Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	46.0 ^b	46.4 ^b	45.8 ^b	46.1 ^b	45.4 ^b	46.2 ^b	47.2 ^b	47.4 ^b	47.3 ^b	47.0 ^b	48.0 ^a	48.6	49.2
Marijuana and Hashish	40.4 ^b	40.6 ^b	40.2 ^b	40.1 ^b	39.7 ^b	40.6 ^b	41.2 ^b	41.7 ^b	42.0 ^b	41.9 ^b	42.8 ^b	43.7	44.2
Cocaine	14.4	14.7	14.2	13.8 ^b	14.3	14.5	14.8	14.6	14.7	14.3	14.5	14.3	14.8
Crack	3.6	3.3	3.3	3.3	3.5	3.5	3.4	3.3	3.6	3.2 ^a	3.5	3.4	3.6
Heroin	1.6 ^a	1.6 ^a	1.3 ^b	1.5 ^b	1.5 ^a	1.5 ^a	1.5 ^a	1.5 ^b	1.6	1.6	1.8	1.8	1.8
Hallucinogens	14.6	14.5	14.3	13.9 ^b	14.3	13.8 ^b	14.4	14.9	14.8	14.1 ^a	14.6	15.1	15.0
LSD	10.4 ^b	10.3 ^b	9.7	9.2	9.5	9.2	9.4	9.4	9.2	8.9	9.1	9.4	9.4
PCP	3.2 ^b	3.0 ^b	2.8 ^a	2.7	2.7	2.5	2.7	2.5	2.5	2.4	2.5	2.5	2.4
Ecstasy	4.3 ^b	4.6 ^b	4.6 ^b	4.7 ^b	5.0 ^b	5.0 ^b	5.2 ^b	5.7 ^b	6.3	5.7 ^b	6.2	6.8	6.6
Inhalants	9.7 ^b	9.7 ^b	9.5 ^b	9.4 ^b	9.3 ^b	9.1 ^b	9.0 ^b	9.0 ^b	8.6 ^a	8.0	8.1	8.0	8.0
Nonmedical Use of Psychotherapeutics ^{2,3}	20.4	20.6	20.4	20.4	20.7	20.4	20.9	20.6	20.4	19.9	20.9	20.3	20.5
Pain Relievers	12.6 ^b	13.1	13.2	13.4	13.6	13.4	14.0	14.0	13.8	13.3	14.2 ^a	13.5	13.6
OxyContin [®]	0.8 ^b	1.2 ^b	1.3 ^b	1.4 ^b	1.7 ^b	1.8 ^b	1.9 ^b	2.3 ^b	2.4	2.3 ^b	2.5	2.7	2.7
Tranquilizers	8.2 ^b	8.5 ^b	8.3 ^b	8.7 ^b	8.7 ^b	8.2 ^b	8.6 ^b	8.7 ^b	8.7 ^a	8.4 ^b	9.1	9.0	9.4
Stimulants ³	10.0 ^b	9.7 ^b	9.3 ^b	8.6	9.1 ^a	8.7	8.5	8.7	8.6	7.9 ^a	8.3	8.3	8.5
Methamphetamine ³	6.5 ^b	6.4 ^b	6.0 ^b	5.2	5.8 ^b	5.3 ^a	5.1	5.1	5.1	4.6	4.7	4.7	4.9
Sedatives	4.2 ^b	4.0 ^b	4.1 ^b	3.7 ^b	3.6 ^b	3.4 ^a	3.6 ^b	3.4 ^a	3.0	2.9	3.1	2.9	3.0
ILLCIT DRUGS OTHER THAN MARIJUANA¹	29.9	29.9	29.4	29.5	29.6	29.7	30.4	30.2	30.2	29.3 ^a	30.0	29.8	30.2

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.2A Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	35,132 ^b	34,993 ^b	34,807 ^b	35,041 ^b	35,810 ^b	35,789 ^b	35,643 ^b	38,123 ^b	38,908 ^b	38,287 ^b	41,479 ^b	41,591 ^b	44,157
Marijuana and Hashish	25,755 ^b	25,231 ^b	25,451 ^b	25,375 ^b	25,432 ^b	25,136 ^b	25,891 ^b	28,688 ^b	29,301 ^b	29,739 ^b	31,513 ^b	32,952 ^b	35,124
Cocaine	5,902 ^b	5,908 ^b	5,658 ^b	5,523 ^b	6,075 ^b	5,748 ^b	5,283 ^b	4,806	4,533	3,857 ^b	4,671	4,182	4,553
Crack	1,554 ^b	1,406 ^b	1,304 ^b	1,381 ^b	1,478 ^b	1,452 ^b	1,113 ^b	1,008	885	625	921	632	773
Heroin	404 ^b	314 ^b	398 ^b	379 ^b	560 ^b	373 ^b	455 ^b	582 ^b	621 ^b	620 ^b	669 ^a	681 ^a	914
Hallucinogens	4,749 ^a	3,936	3,878	3,809 ^a	3,970	3,759 ^a	3,670 ^b	4,536	4,543	4,069	4,306	4,430	4,250
LSD	999 ^b	558 ^b	592 ^b	563 ^b	667 ^b	621 ^b	808 ^b	790 ^b	881 ^b	880 ^b	1,057 ^a	1,111	1,290
PCP	235 ^b	219 ^b	210 ^b	164 ^b	187 ^a	139	99	125	96	119	172 ^a	90	90
Ecstasy	3,167 ^b	2,119	1,915 ^b	1,960 ^a	2,138	2,122	2,116	2,804 ^b	2,653	2,422	2,610	2,588	2,342
Inhalants	2,084 ^b	2,075 ^b	2,255 ^b	2,187 ^b	2,219 ^b	2,085 ^b	2,065 ^b	2,109 ^b	2,048 ^b	1,861	1,693	1,533	1,617
Nonmedical Use of Psychotherapeutics ^{2,3}	14,795	15,163	14,849	15,346	16,482 ^b	16,334 ^b	15,190	16,064 ^a	16,051 ^a	14,657	16,666 ^b	15,348	14,966
Pain Relievers	10,992	11,671 ^b	11,256 ^a	11,815 ^b	12,650 ^b	12,517 ^b	11,916 ^b	12,450 ^b	12,242 ^b	11,143 ^a	12,489 ^b	11,082	10,337
OxyContin [®]	--	--	1,213	1,226	1,324	1,427	1,462	1,685 ^b	1,879 ^b	1,623 ^b	1,477	1,442	1,237
Tranquilizers	4,849	5,051	5,068	5,249	5,066	5,293	5,099	5,493	5,583	5,109	6,073 ^b	5,269	5,202
Stimulants ³	3,380	3,031 ^b	3,254 ^a	3,088 ^b	3,789	2,994 ^b	2,648 ^b	3,093 ^b	2,902 ^b	2,700 ^b	3,317	3,492	3,715
Methamphetamine ³	1,755 ^b	1,602 ^a	1,808 ^b	1,603 ^a	1,885 ^b	1,345	854 ^b	1,167	959 ^a	1,033	1,155	1,186	1,301
Sedatives	981	831	737	750	927	864	626	816	906	526 ^a	590	639	775
ILLCIT DRUGS OTHER THAN MARIJUANA¹	20,423	20,305	19,658	20,109	21,262 ^b	21,205 ^a	20,018	21,077 ^a	20,625	18,959	21,267 ^b	19,868	19,719

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.2B Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	14.9 ^b	14.7 ^b	14.5 ^b	14.4 ^b	14.6 ^b	14.4 ^b	14.3 ^b	15.1 ^b	15.3 ^b	14.9 ^b	16.0 ^a	15.9 ^a	16.7
Marijuana and Hashish	11.0 ^b	10.6 ^b	10.6 ^b	10.4 ^b	10.3 ^b	10.1 ^b	10.4 ^b	11.4 ^b	11.6 ^b	11.5 ^b	12.1 ^b	12.6 ^a	13.2
Cocaine	2.5 ^b	2.5 ^b	2.4 ^b	2.3 ^b	2.5 ^b	2.3 ^b	2.1 ^b	1.9	1.8	1.5 ^a	1.8	1.6	1.7
Crack	0.7 ^b	0.6 ^b	0.5 ^b	0.6 ^b	0.6 ^b	0.6 ^b	0.4 ^b	0.4 ^a	0.3	0.2	0.4	0.2	0.3
Heroin	0.2 ^b	0.1 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^a	0.2 ^b	0.3 ^a	0.3 ^a	0.3
Hallucinogens	2.0 ^b	1.7	1.6	1.6	1.6	1.5	1.5	1.8 ^a	1.8 ^a	1.6	1.7	1.7	1.6
LSD	0.4	0.2 ^b	0.2 ^b	0.2 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.4	0.4	0.5
PCP	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^a	0.1 ^a	0.0	0.0	0.0	0.0	0.1 ^a	0.0	0.0
Ecstasy	1.3 ^b	0.9	0.8	0.8	0.9	0.9	0.8	1.1 ^b	1.0 ^a	0.9	1.0	1.0	0.9
Inhalants	0.9 ^b	0.9 ^b	0.9 ^b	0.9 ^b	0.9 ^b	0.8 ^b	0.8 ^b	0.8 ^b	0.8 ^b	0.7 ^a	0.7	0.6	0.6
Nonmedical Use of Psychotherapeutics ^{2,3}	6.3 ^b	6.4 ^b	6.2 ^b	6.3 ^b	6.7 ^b	6.6 ^b	6.1 ^a	6.4 ^b	6.3 ^b	5.7	6.4 ^b	5.8	5.6
Pain Relievers	4.7 ^b	4.9 ^b	4.7 ^b	4.9 ^b	5.1 ^b	5.1 ^b	4.8 ^b	4.9 ^b	4.8 ^b	4.3 ^b	4.8 ^b	4.2 ^a	3.9
OxyContin [®]	--	--	0.5	0.5	0.5	0.6 ^a	0.6 ^a	0.7 ^b	0.7 ^b	0.6 ^b	0.6 ^a	0.5	0.5
Tranquilizers	2.1	2.1	2.1	2.2	2.1	2.1	2.0	2.2	2.2 ^a	2.0	2.3 ^b	2.0	2.0
Stimulants ³	1.4	1.3	1.4	1.3	1.5	1.2 ^a	1.1 ^b	1.2 ^a	1.1 ^b	1.0 ^b	1.3	1.3	1.4
Methamphetamine ³	0.7 ^b	0.7 ^b	0.8 ^b	0.7 ^b	0.8 ^b	0.5	0.3 ^b	0.5	0.4 ^a	0.4	0.4	0.5	0.5
Sedatives	0.4 ^a	0.3	0.3	0.3	0.4	0.3	0.3	0.3	0.4	0.2 ^a	0.2	0.2	0.3
ILLCIT DRUGS OTHER THAN MARIJUANA¹	8.7 ^b	8.5 ^b	8.2 ^b	8.3 ^b	8.6 ^b	8.6 ^b	8.0 ^b	8.4 ^b	8.1 ^b	7.4	8.2 ^b	7.6	7.4

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.3A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	19,522 ^b	19,470 ^b	19,071 ^b	19,720 ^b	20,387 ^b	19,892 ^b	20,123 ^b	21,930 ^b	22,648 ^b	22,454 ^b	23,863 ^b	24,573 ^b	26,983
Marijuana and Hashish	14,584 ^b	14,638 ^b	14,576 ^b	14,626 ^b	14,849 ^b	14,470 ^b	15,269 ^b	16,826 ^b	17,409 ^b	18,071 ^b	18,855 ^b	19,810 ^b	22,188
Cocaine	2,020 ^b	2,281 ^b	2,021 ^b	2,397 ^b	2,426 ^b	2,077 ^b	1,865 ^a	1,642	1,472	1,369	1,650	1,549	1,530
Crack	567 ^a	604 ^a	467	682 ^b	704 ^b	611 ^b	363	497	378	228	443	377	354
Heroin	166 ^b	119 ^b	166 ^b	136 ^b	339	161 ^b	213 ^b	193 ^b	239 ^b	281 ^a	335	289 ^a	435
Hallucinogens	1,196	1,042	929 ^a	1,088	1,008	998	1,059	1,263	1,207	972	1,134	1,333	1,173
LSD	112 ^b	133 ^b	141 ^b	104 ^b	130 ^b	144 ^b	157 ^b	159 ^b	155 ^b	152 ^b	184 ^a	229	287
PCP	58	56	49	48	31	43	25	54	36	26	32	33	*
Ecstasy	676	470	450 ^a	502	529	503	546	763	691	544	628	660	609
Inhalants	635	570	638	611	762 ^a	617	644	562	690	624	493	496	546
Nonmedical Use of Psychotherapeutics ^{2,3}	6,287	6,451	6,110	6,491	7,098	6,907	6,210	6,980	6,957	6,119	6,831	6,484	6,537
Pain Relievers	4,377	4,693	4,404	4,658	5,226 ^b	5,188 ^b	4,732	5,282 ^b	5,093 ^b	4,471	4,862	4,521	4,325
OxyContin [®]	--	--	325	334	275	371	436	510 ^a	566 ^a	434	358	492	356
Tranquilizers	1,804	1,830	1,616	1,817	1,761	1,838	1,796	2,030	2,159	1,840	2,139	1,705	1,875
Stimulants ³	1,303 ^a	1,310	1,312	1,188 ^b	1,385	1,051 ^b	908 ^b	1,299	1,082 ^b	970 ^b	1,171 ^b	1,365	1,595
Methamphetamine ³	683	726	706	628	731	530	314 ^b	502	353 ^a	439	440	595	569
Sedatives	436	294	265	272	385	345	237	367	375	231	270	251	330
ILLCIT DRUGS OTHER THAN MARIJUANA¹	8,777	8,849	8,247	8,963	9,620 ^a	9,295	8,556	9,195	9,016	8,020	8,883	8,665	8,719

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.3B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	8.3 ^b	8.2 ^b	7.9 ^b	8.1 ^b	8.3 ^b	8.0 ^b	8.1 ^b	8.7 ^b	8.9 ^b	8.7 ^b	9.2 ^b	9.4 ^b	10.2
Marijuana and Hashish	6.2 ^b	6.2 ^b	6.1 ^b	6.0 ^b	6.0 ^b	5.8 ^b	6.1 ^b	6.7 ^b	6.9 ^b	7.0 ^b	7.3 ^b	7.5 ^b	8.4
Cocaine	0.9 ^b	1.0 ^b	0.8 ^b	1.0 ^b	1.0 ^b	0.8 ^b	0.7 ^a	0.7	0.6	0.5	0.6	0.6	0.6
Crack	0.2 ^b	0.3 ^b	0.2	0.3 ^b	0.3 ^b	0.2 ^b	0.1	0.2	0.1	0.1	0.2	0.1	0.1
Heroin	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^a	0.1	0.1 ^a	0.2
Hallucinogens	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.4	0.4	0.5	0.4
LSD	0.0 ^b	0.1 ^b	0.1 ^b	0.0 ^b	0.1 ^b	0.1 ^a	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1	0.1	0.1
PCP	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
Ecstasy	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.3 ^a	0.3	0.2	0.2	0.3	0.2
Inhalants	0.3	0.2	0.3	0.3	0.3 ^b	0.2	0.3	0.2	0.3	0.2	0.2	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{2,3}	2.7	2.7	2.5	2.7	2.9 ^b	2.8 ^a	2.5	2.8 ^a	2.7 ^a	2.4	2.6	2.5	2.5
Pain Relievers	1.9 ^a	2.0 ^b	1.8 ^a	1.9 ^b	2.1 ^b	2.1 ^b	1.9 ^a	2.1 ^b	2.0 ^b	1.7	1.9 ^a	1.7	1.6
OxyContin [®]	--	--	0.1	0.1	0.1	0.1	0.2	0.2 ^a	0.2 ^b	0.2	0.1	0.2	0.1
Tranquilizers	0.8	0.8	0.7	0.7	0.7	0.7	0.7	0.8	0.9 ^a	0.7	0.8	0.6	0.7
Stimulants ³	0.6	0.6	0.5	0.5 ^a	0.6	0.4 ^b	0.4 ^b	0.5	0.4 ^b	0.4 ^b	0.5 ^b	0.5	0.6
Methamphetamine ³	0.3	0.3 ^a	0.3	0.3	0.3	0.2	0.1 ^a	0.2	0.1 ^a	0.2	0.2	0.2	0.2
Sedatives	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	3.7 ^b	3.7 ^b	3.4	3.7 ^b	3.9 ^b	3.8 ^b	3.4	3.7 ^a	3.6	3.1	3.4	3.3	3.3

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.4A Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	7,642 ^b	7,631 ^b	7,566 ^b	7,022 ^b	7,014 ^b	6,638 ^b	6,565 ^b	6,629 ^b	6,278 ^b	6,373 ^b	6,044	5,794	5,791
Marijuana and Hashish	5,104 ^b	4,888 ^b	4,788 ^b	4,402 ^a	4,400 ^a	4,096	4,138	4,204	4,157	4,378 ^a	4,232	4,093	4,091
Cocaine	670 ^b	648 ^b	611 ^b	582 ^b	571 ^b	544 ^b	473 ^b	405 ^b	363 ^b	337 ^b	265	215	222
Crack	163 ^b	157 ^b	125 ^b	108 ^b	129 ^b	107 ^b	80 ^b	58	60	68	34	23	35
Heroin	94 ^b	80 ^b	78 ^b	60	60	55	67 ^a	56	46	70 ^a	59	38	36
Hallucinogens	1,419 ^b	1,250 ^b	1,160 ^b	998 ^b	990 ^b	898 ^b	972 ^b	1,015 ^b	1,000 ^b	930 ^b	823 ^b	702	610
LSD	670 ^b	397 ^b	313	288	219 ^a	202 ^b	271	249	231	229	249	221 ^a	291
PCP	233 ^b	204 ^b	175 ^b	182 ^b	128 ^b	105 ^a	102 ^a	99 ^a	76	76	96 ^a	46	60
Ecstasy	820 ^b	603 ^b	523 ^b	399 ^b	476 ^b	446 ^b	524 ^b	572 ^b	618 ^b	605 ^b	488 ^b	369 ^a	297
Inhalants	2,605 ^b	2,670 ^b	2,762 ^b	2,658 ^b	2,559 ^b	2,428 ^b	2,326 ^b	2,283 ^b	2,015 ^b	1,877 ^b	1,613 ^b	1,327	1,307
Nonmedical Use of Psychotherapeutics ^{2,3}	3,402 ^b	3,372 ^b	3,403 ^b	3,019 ^b	3,175 ^b	2,954 ^b	2,783 ^b	2,810 ^b	2,591 ^b	2,522 ^a	2,482 ^a	2,188	2,290
Pain Relievers	2,775 ^b	2,798 ^b	2,879 ^b	2,517 ^b	2,650 ^b	2,457 ^b	2,353 ^b	2,393 ^b	2,245 ^b	2,150 ^b	2,069 ^b	1,805	1,803
OxyContin [®]	224	254	305	276	343 ^b	360 ^b	344 ^b	381 ^b	369 ^b	369 ^b	322 ^a	274	252
Tranquilizers	834 ^b	874 ^b	816 ^b	766	742	748	721	696	723	699	703	572	665
Stimulants ³	1,112 ^b	1,046 ^b	906 ^b	870 ^b	930 ^b	730 ^b	673 ^b	583	531	508	495	437	522
Methamphetamine ³	416 ^b	382 ^b	346 ^b	319 ^b	341 ^b	221	195	187	157	184	145	119 ^a	171
Sedatives	238	249	251	232	224	213	205	224	178	144 ^a	175	125 ^b	207
ILLCIT DRUGS OTHER THAN MARIJUANA¹	5,286 ^b	5,319 ^b	5,355 ^b	4,944 ^b	4,951 ^b	4,649 ^b	4,513 ^b	4,501 ^b	4,166 ^b	4,007 ^b	3,691 ^b	3,279	3,321

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.4B Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	30.9 ^b	30.5 ^b	30.0 ^b	27.7 ^b	27.6 ^b	26.3 ^b	26.4 ^b	26.9 ^b	25.8 ^b	25.5 ^b	24.2	23.3	23.3
Marijuana and Hashish	20.6 ^b	19.6 ^b	19.0 ^b	17.4	17.3	16.2	16.6	17.1	17.1	17.5 ^a	17.0	16.4	16.4
Cocaine	2.7 ^b	2.6 ^b	2.4 ^b	2.3 ^b	2.2 ^b	2.2 ^b	1.9 ^b	1.6 ^b	1.5 ^b	1.3 ^b	1.1	0.9	0.9
Crack	0.7 ^b	0.6 ^b	0.5 ^b	0.4 ^b	0.5 ^b	0.4 ^b	0.3 ^b	0.2	0.2	0.3	0.1	0.1	0.1
Heroin	0.4 ^b	0.3 ^b	0.3 ^b	0.2	0.2	0.2	0.3 ^a	0.2	0.2	0.3 ^a	0.2	0.2	0.1
Hallucinogens	5.7 ^b	5.0 ^b	4.6 ^b	3.9 ^b	3.9 ^b	3.6 ^b	3.9 ^b	4.1 ^b	4.1 ^b	3.7 ^b	3.3 ^b	2.8	2.5
LSD	2.7 ^b	1.6 ^b	1.2	1.1	0.9 ^a	0.8 ^b	1.1	1.0	0.9	0.9	1.0	0.9 ^a	1.2
PCP	0.9 ^b	0.8 ^b	0.7 ^b	0.7 ^b	0.5 ^b	0.4 ^a	0.4 ^a	0.4 ^a	0.3	0.3	0.4 ^a	0.2	0.2
Ecstasy	3.3 ^b	2.4 ^b	2.1 ^b	1.6 ^a	1.9 ^b	1.8 ^b	2.1 ^b	2.3 ^b	2.5 ^b	2.4 ^b	2.0 ^b	1.5 ^a	1.2
Inhalants	10.5 ^b	10.7 ^b	11.0 ^b	10.5 ^b	10.1 ^b	9.6 ^b	9.3 ^b	9.3 ^b	8.3 ^b	7.5 ^b	6.5 ^b	5.3	5.3
Nonmedical Use of Psychotherapeutics ^{2,3}	13.7 ^b	13.5 ^b	13.5 ^b	11.9 ^b	12.5 ^b	11.7 ^b	11.2 ^b	11.4 ^b	10.6 ^b	10.1 ^a	10.0 ^a	8.8	9.2
Pain Relievers	11.2 ^b	11.2 ^b	11.4 ^b	9.9 ^b	10.4 ^b	9.7 ^b	9.5 ^b	9.7 ^b	9.2 ^b	8.6 ^b	8.3 ^b	7.3	7.3
OxyContin [®]	0.9	1.0	1.2	1.1	1.4 ^a	1.4 ^b	1.4 ^b	1.5 ^b	1.5 ^b	1.5 ^b	1.3 ^a	1.1	1.0
Tranquilizers	3.4 ^b	3.5 ^b	3.2 ^a	3.0	2.9	3.0	2.9	2.8	3.0	2.8	2.8	2.3	2.7
Stimulants ³	4.5 ^b	4.2 ^b	3.6 ^b	3.4 ^b	3.7 ^b	2.9 ^b	2.7 ^b	2.4	2.2	2.0	2.0	1.8	2.1
Methamphetamine ³	1.7 ^b	1.5 ^b	1.4 ^b	1.3 ^b	1.3 ^b	0.9	0.8	0.8	0.6	0.7	0.6	0.5 ^a	0.7
Sedatives	1.0	1.0	1.0	0.9	0.9	0.8	0.8	0.9	0.7	0.6 ^a	0.7	0.5 ^b	0.8
ILLCIT DRUGS OTHER THAN MARIJUANA¹	21.4 ^b	21.3 ^b	21.2 ^b	19.5 ^b	19.5 ^b	18.4 ^b	18.1 ^b	18.3 ^b	17.1 ^b	16.0 ^b	14.8 ^b	13.2	13.4

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.5A Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	5,494 ^b	5,455 ^b	5,300 ^b	5,041 ^b	4,978 ^b	4,737 ^b	4,751 ^b	4,838 ^b	4,745 ^b	4,735 ^b	4,468	4,287	4,336
Marijuana and Hashish	3,905 ^b	3,745 ^b	3,656 ^b	3,383	3,355	3,158	3,258	3,363	3,414	3,553 ^a	3,363	3,346	3,262
Cocaine	508 ^b	438 ^b	411 ^b	423 ^b	411 ^b	385 ^b	306 ^b	242 ^a	239 ^a	229 ^a	180	131	168
Crack	100 ^b	95 ^b	63 ^b	54 ^b	74 ^b	65 ^b	34	29	18	29	19	11	20
Heroin	51	36	48	37	37	24	41	34	28	54	31	31	28
Hallucinogens	945 ^b	778 ^b	753 ^b	665 ^b	653 ^b	629 ^b	689 ^b	737 ^b	747 ^b	648 ^b	553 ^b	482	422
LSD	314 ^b	149 ^a	157	150 ^a	106 ^b	126 ^b	185	143 ^a	136 ^b	159	152 ^a	150 ^a	214
PCP	110 ^b	89 ^b	72 ^b	85 ^b	59	56	61 ^a	48	35	39	48	29	33
Ecstasy	547 ^b	329 ^b	310 ^b	256 ^a	308 ^b	328 ^b	346 ^b	416 ^b	465 ^b	419 ^b	288 ^b	232	182
Inhalants	1,078 ^b	1,119 ^b	1,168 ^b	1,144 ^b	1,109 ^b	995 ^b	986 ^b	960 ^b	885 ^b	820 ^b	647 ^b	480	512
Nonmedical Use of Psychotherapeutics ^{2,3}	2,285 ^b	2,295 ^b	2,228 ^b	2,101 ^b	2,184 ^b	2,037 ^b	1,919 ^b	1,895 ^b	1,812 ^b	1,749 ^a	1,642	1,441	1,554
Pain Relievers	1,870 ^b	1,917 ^b	1,865 ^b	1,741 ^b	1,821 ^b	1,684 ^b	1,613 ^b	1,628 ^b	1,522 ^b	1,484 ^b	1,333 ^a	1,157	1,165
OxyContin [®]	--	--	194	187	200	232 ^b	226 ^a	225 ^a	235 ^a	219 ^a	187	150	162
Tranquilizers	568 ^b	586 ^b	532 ^a	497	484	505	483	462	466	475	452	347	424
Stimulants ³	665 ^b	578 ^b	507 ^b	506 ^b	548 ^b	415	373	334	325	299	324	273 ^a	361
Methamphetamine ³	245 ^b	185 ^b	177 ^b	178 ^b	185 ^b	116	94	87	98	106	82	73	105
Sedatives	138	131	133	118	110	111	113	126	111	84	81	65 ^a	116
ILLCIT DRUGS OTHER THAN MARIJUANA¹	3,338 ^b	3,351 ^b	3,269 ^b	3,129 ^b	3,147 ^b	2,921 ^b	2,860 ^b	2,849 ^b	2,758 ^b	2,565 ^b	2,297 ^a	1,979	2,076

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.5B Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	22.2 ^b	21.8 ^b	21.0 ^b	19.9 ^b	19.6 ^b	18.8 ^a	19.1 ^b	19.7 ^b	19.5 ^b	19.0 ^b	17.9	17.2	17.4
Marijuana and Hashish	15.8 ^b	15.0 ^b	14.5 ^b	13.3	13.2	12.5	13.1	13.7	14.0	14.2 ^a	13.5	13.4	13.1
Cocaine	2.1 ^b	1.8 ^b	1.6 ^b	1.7 ^b	1.6 ^b	1.5 ^b	1.2 ^b	1.0 ^a	1.0 ^a	0.9 ^a	0.7	0.5	0.7
Crack	0.4 ^b	0.4 ^b	0.3 ^b	0.2 ^b	0.3 ^b	0.3 ^b	0.1	0.1	0.1	0.1	0.1	0.0	0.1
Heroin	0.2	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1
Hallucinogens	3.8 ^b	3.1 ^b	3.0 ^b	2.6 ^b	2.6 ^b	2.5 ^b	2.8 ^b	3.0 ^b	3.1 ^b	2.6 ^b	2.2 ^b	1.9	1.7
LSD	1.3 ^b	0.6 ^a	0.6 ^a	0.6 ^a	0.4 ^b	0.5 ^b	0.7	0.6 ^a	0.6 ^a	0.6	0.6 ^a	0.6 ^a	0.9
PCP	0.4 ^b	0.4 ^b	0.3 ^a	0.3 ^b	0.2	0.2	0.2 ^a	0.2	0.1	0.2	0.2	0.1	0.1
Ecstasy	2.2 ^b	1.3 ^b	1.2 ^b	1.0 ^a	1.2 ^b	1.3 ^b	1.4 ^b	1.7 ^b	1.9 ^b	1.7 ^b	1.2 ^b	0.9	0.7
Inhalants	4.4 ^b	4.5 ^b	4.6 ^b	4.5 ^b	4.4 ^b	3.9 ^b	4.0 ^b	3.9 ^b	3.6 ^b	3.3 ^b	2.6 ^b	1.9	2.1
Nonmedical Use of Psychotherapeutics ^{2,3}	9.2 ^b	9.2 ^b	8.8 ^b	8.3 ^b	8.6 ^b	8.1 ^b	7.7 ^b	7.7 ^b	7.4 ^b	7.0 ^a	6.6	5.8	6.2
Pain Relievers	7.6 ^b	7.7 ^b	7.4 ^b	6.9 ^b	7.2 ^b	6.7 ^b	6.5 ^b	6.6 ^b	6.3 ^b	5.9 ^b	5.3 ^a	4.6	4.7
OxyContin [®]	--	--	0.8	0.7	0.8	0.9 ^a	0.9 ^a	0.9 ^a	1.0 ^a	0.9 ^a	0.7	0.6	0.7
Tranquilizers	2.3 ^b	2.3 ^b	2.1 ^a	2.0	1.9	2.0	1.9	1.9	1.9	1.9	1.8	1.4	1.7
Stimulants ³	2.7 ^b	2.3 ^b	2.0 ^b	2.0 ^b	2.2 ^b	1.6	1.5	1.4	1.3	1.2	1.3	1.1 ^a	1.5
Methamphetamine ³	1.0 ^b	0.7 ^b	0.7 ^b	0.7 ^b	0.7 ^b	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.4
Sedatives	0.6	0.5	0.5	0.5	0.4	0.4	0.5	0.5	0.5	0.3	0.3	0.3 ^a	0.5
ILLCIT DRUGS OTHER THAN MARIJUANA¹	13.5 ^b	13.4 ^b	13.0 ^b	12.3 ^b	12.4 ^b	11.6 ^b	11.5 ^b	11.6 ^b	11.3 ^b	10.3 ^b	9.2 ^a	8.0	8.3

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.6A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	2,878 ^b	2,811 ^b	2,674 ^b	2,511	2,488	2,415	2,312	2,484	2,468	2,522	2,377	2,197	2,338
Marijuana and Hashish	2,023 ^a	1,971	1,909	1,730	1,695	1,693	1,668	1,813	1,802	1,967	1,800	1,762	1,830
Cocaine	146 ^b	153 ^b	126 ^b	142 ^b	110 ^b	104 ^b	98 ^b	70	60	67	30	43	39
Crack	26	32 ^b	16	25 ^a	12	16	7	8	4	3	*	3	8
Heroin	11	18	15	15	16	3 ^a	14	13	8	15	*	13	16
Hallucinogens	255 ^b	260 ^b	211 ^b	205 ^a	182	174	245 ^b	217 ^b	215 ^b	215 ^b	160	154	136
LSD	61	51	38	32 ^a	31 ^a	30 ^a	52	35	46	34	26 ^a	51	65
PCP	23 ^b	20 ^b	11	17 ^a	12	9	15 ^a	12	6	4	5	7	3
Ecstasy	124 ^b	91 ^b	77 ^a	74 ^a	83 ^b	65	105 ^b	115 ^b	123 ^b	108 ^b	73 ^a	48	39
Inhalants	301 ^b	320 ^b	312 ^b	313 ^b	340 ^b	294 ^b	270 ^b	244 ^b	262 ^b	223 ^b	188	121	149
Nonmedical Use of Psychotherapeutics ^{2,3}	986 ^b	997 ^b	914 ^b	838 ^b	846 ^b	830 ^b	711	774 ^a	721	688	693	549 ^a	655
Pain Relievers	794 ^b	801 ^b	751 ^b	686 ^b	692 ^b	671 ^b	566 ^a	658 ^b	607 ^b	563 ^a	558 ^a	425	467
OxyContin [®]	--	--	66 ^a	34	29	47	49	70 ^a	47	51	49	30	35
Tranquilizers	188 ^b	226 ^b	161 ^b	161 ^a	125	181 ^b	151 ^a	144	134	162 ^a	140	90	103
Stimulants ³	208	222	180	172	171	126	112 ^a	127	107 ^a	98 ^b	118	84 ^b	169
Methamphetamine ³	71	70	58	72	53	34	29	36	25	37	30	22	45
Sedatives	47	47	26	27	45	34	35	40	35	24	26	25	41
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1,421 ^b	1,417 ^b	1,328 ^b	1,253 ^b	1,252 ^b	1,182 ^b	1,092 ^b	1,122 ^b	1,092 ^b	1,026 ^a	918	757 ^a	884

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.6B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	11.6 ^b	11.2 ^b	10.6 ^b	9.9	9.8	9.6	9.3	10.1	10.1	10.1	9.5	8.8	9.4
Marijuana and Hashish	8.2 ^a	7.9	7.6	6.8	6.7 ^a	6.7	6.7	7.4	7.4	7.9	7.2	7.1	7.4
Cocaine	0.6 ^b	0.6 ^b	0.5 ^b	0.6 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.3	0.2	0.3	0.1	0.2	0.2
Crack	0.1	0.1 ^b	0.1	0.1 ^a	0.0	0.1	0.0	0.0	0.0	0.0	*	0.0	0.0
Heroin	0.0	0.1	0.1	0.1	0.1	0.0 ^a	0.1	0.1	0.0	0.1	*	0.1	0.1
Hallucinogens	1.0 ^b	1.0 ^b	0.8 ^b	0.8 ^a	0.7	0.7	1.0 ^b	0.9 ^b	0.9 ^b	0.9 ^b	0.6	0.6	0.5
LSD	0.2	0.2	0.2	0.1 ^a	0.1 ^a	0.1 ^a	0.2	0.1	0.2	0.1	0.1 ^a	0.2	0.3
PCP	0.1 ^b	0.1 ^b	0.0	0.1 ^a	0.0	0.0	0.1 ^a	0.1	0.0	0.0	0.0	0.0	0.0
Ecstasy	0.5 ^b	0.4 ^b	0.3 ^a	0.3 ^a	0.3 ^b	0.3	0.4 ^b	0.5 ^b	0.5 ^b	0.4 ^b	0.3 ^a	0.2	0.2
Inhalants	1.2 ^b	1.3 ^b	1.2 ^b	1.2 ^b	1.3 ^b	1.2 ^b	1.1 ^b	1.0 ^b	1.1 ^b	0.9 ^b	0.8	0.5	0.6
Nonmedical Use of Psychotherapeutics ^{2,3}	4.0 ^b	4.0 ^b	3.6 ^b	3.3 ^b	3.3 ^b	3.3 ^b	2.9	3.1 ^a	3.0	2.8	2.8	2.2 ^a	2.6
Pain Relievers	3.2 ^b	3.2 ^b	3.0 ^b	2.7 ^b	2.7 ^b	2.7 ^b	2.3 ^a	2.7 ^b	2.5 ^b	2.3 ^a	2.2	1.7	1.9
OxyContin [®]	--	--	0.3 ^a	0.1	0.1	0.2	0.2	0.3 ^a	0.2	0.2	0.2	0.1	0.1
Tranquilizers	0.8 ^b	0.9 ^b	0.6 ^a	0.6 ^a	0.5	0.7 ^b	0.6 ^a	0.6	0.5	0.6 ^a	0.6	0.4	0.4
Stimulants ³	0.8	0.9	0.7	0.7	0.7	0.5	0.5 ^a	0.5	0.4 ^a	0.4 ^b	0.5	0.3 ^b	0.7
Methamphetamine ³	0.3	0.3	0.2	0.3	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2
Sedatives	0.2	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA¹	5.7 ^b	5.7 ^b	5.3 ^b	4.9 ^b	4.9 ^b	4.7 ^b	4.4 ^b	4.6 ^b	4.5 ^b	4.1 ^a	3.7	3.0 ^a	3.6

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.7A Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	18,562 ^b	19,195 ^b	19,043 ^b	19,222 ^b	19,328 ^b	18,831 ^b	18,772 ^b	19,658 ^a	19,538 ^b	19,534 ^b	19,977	19,826	20,217
Marijuana and Hashish	16,680 ^b	17,103 ^b	16,983 ^b	17,032 ^b	17,180 ^b	16,652 ^b	16,718 ^b	17,677 ^b	17,529 ^b	17,796 ^a	18,052	18,051	18,390
Cocaine	4,786 ^b	4,755 ^b	4,893 ^b	4,905 ^b	5,139 ^b	4,925 ^b	4,772 ^b	4,997 ^b	4,568 ^b	4,253 ^a	4,267 ^a	4,041	3,881
Crack	1,182 ^b	1,193 ^b	1,135 ^b	1,145 ^b	1,176 ^b	985 ^b	916 ^b	965 ^b	886 ^b	714 ^b	654	563	546
Heroin	487 ^b	516 ^b	529 ^a	496 ^b	514 ^b	498 ^b	467 ^b	587	614	594	664	619	683
Hallucinogens	7,517 ^b	7,377 ^b	6,855 ^b	6,820 ^b	6,625 ^b	6,291 ^b	5,876	6,418 ^b	6,295 ^b	6,029	6,103	6,122	5,784
LSD	4,928 ^b	4,450 ^b	3,900 ^b	3,417 ^b	2,933 ^b	2,381	2,167 ^a	2,326	2,194 ^a	2,054 ^b	2,056 ^b	2,246	2,451
PCP	834 ^b	948 ^b	737 ^b	716 ^b	641 ^b	540 ^b	523 ^b	480 ^b	447 ^b	351	345	326	305
Ecstasy	4,674 ^b	4,683 ^b	4,439	4,464	4,392	4,186	4,008	4,194	4,230	4,222	4,452	4,436	4,200
Inhalants	4,867 ^b	4,729 ^b	4,516 ^b	4,333 ^b	4,097 ^b	3,703 ^b	3,445 ^b	3,620 ^b	3,396 ^b	3,106 ^b	2,918 ^b	2,613	2,444
Nonmedical Use of Psychotherapeutics ^{2,3}	8,778 ^a	9,360	9,564	9,990 ^b	10,067 ^b	9,807 ^b	9,694 ^a	9,896 ^b	9,832 ^b	9,373	9,709 ^a	9,239	9,197
Pain Relievers	6,854	7,513 ^a	7,813 ^b	8,279 ^b	8,353 ^b	8,143 ^b	8,096 ^b	8,221 ^b	8,143 ^b	7,600 ^b	7,759 ^b	7,251	7,049
OxyContin [®]	820 ^b	1,145 ^b	1,376 ^b	1,616	1,684	1,716	1,765	2,110 ^b	2,149 ^b	2,079 ^b	2,011 ^a	1,889	1,783
Tranquilizers	3,473 ^b	3,898 ^b	3,919 ^b	4,356	4,286	4,155	4,069	4,323	4,503	4,232	4,420	4,184	4,362
Stimulants ³	3,796 ^b	3,873 ^b	3,892 ^b	4,008 ^b	3,926 ^b	3,569 ^b	3,485 ^b	3,587 ^b	3,534 ^b	3,172	3,374	3,279	3,129
Methamphetamine ³	2,301 ^b	2,192 ^b	2,247 ^b	2,190 ^b	2,103 ^b	1,847 ^b	1,548 ^b	1,518 ^b	1,361 ^b	1,150	1,089	1,042	1,040
Sedatives	648 ^b	565	627 ^b	650 ^b	612 ^b	583 ^a	511	528	469	472	420	439	460
ILLCIT DRUGS OTHER THAN MARIJUANA¹	12,436 ^a	12,770 ^b	12,630 ^b	12,800 ^b	12,959 ^b	12,573 ^b	12,146	12,592 ^b	12,410 ^a	11,860	12,216	11,853	11,862

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.7B Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	59.8 ^b	60.5 ^b	59.2	59.2	59.0	57.5	57.0	58.5	57.3	56.9	57.8	57.0	57.9
Marijuana and Hashish	53.8	53.9	52.8	52.4	52.5	50.9 ^a	50.8 ^a	52.6	51.4	51.9	52.2	51.9	52.6
Cocaine	15.4 ^b	15.0 ^b	15.2 ^b	15.1 ^b	15.7 ^b	15.0 ^b	14.5 ^b	14.9 ^b	13.4 ^b	12.4 ^b	12.3 ^b	11.6	11.1
Crack	3.8 ^b	3.8 ^b	3.5 ^b	3.5 ^b	3.6 ^b	3.0 ^b	2.8 ^b	2.9 ^b	2.6 ^b	2.1 ^b	1.9 ^a	1.6	1.6
Heroin	1.6 ^a	1.6	1.6	1.5 ^a	1.6 ^a	1.5 ^a	1.4 ^b	1.7	1.8	1.7	1.9	1.8	2.0
Hallucinogens	24.2 ^b	23.3 ^b	21.3 ^b	21.0 ^b	20.2 ^b	19.2 ^b	17.8 ^a	19.1 ^b	18.5 ^b	17.6	17.6 ^a	17.6	16.6
LSD	15.9 ^b	14.0 ^b	12.1 ^b	10.5 ^b	9.0 ^b	7.3	6.6	6.9	6.4	6.0 ^b	5.9 ^b	6.5	7.0
PCP	2.7 ^b	3.0 ^b	2.3 ^b	2.2 ^b	2.0 ^b	1.6 ^b	1.6 ^b	1.4 ^b	1.3 ^b	1.0	1.0	0.9	0.9
Ecstasy	15.1 ^b	14.8 ^b	13.8 ^b	13.7 ^b	13.4 ^b	12.8	12.2	12.5	12.4	12.3	12.9	12.8	12.0
Inhalants	15.7 ^b	14.9 ^b	14.0 ^b	13.3 ^b	12.5 ^b	11.3 ^b	10.5 ^b	10.8 ^b	10.0 ^b	9.1 ^b	8.4 ^b	7.5	7.0
Nonmedical Use of Psychotherapeutics ^{2,3}	28.3 ^b	29.5 ^b	29.7 ^b	30.8 ^b	30.7 ^b	30.0 ^b	29.4 ^b	29.5 ^b	28.9 ^b	27.3	28.1 ^b	26.6	26.3
Pain Relievers	22.1 ^b	23.7 ^b	24.3 ^b	25.5 ^b	25.5 ^b	24.9 ^b	24.6 ^b	24.5 ^b	23.9 ^b	22.2 ^b	22.4 ^b	20.8	20.2
OxyContin [®]	2.6 ^b	3.6 ^b	4.3 ^b	5.0	5.1	5.2	5.4	6.3 ^b	6.3 ^b	6.1 ^b	5.8 ^a	5.4	5.1
Tranquilizers	11.2 ^b	12.3	12.2	13.4	13.1	12.7	12.4	12.9	13.2	12.3	12.8	12.0	12.5
Stimulants ³	12.2 ^b	12.2 ^b	12.1 ^b	12.3 ^b	12.0 ^b	10.9 ^b	10.6 ^b	10.7 ^b	10.4 ^b	9.2	9.8	9.4	9.0
Methamphetamine ³	7.4 ^b	6.9 ^b	7.0 ^b	6.7 ^b	6.4 ^b	5.6 ^b	4.7 ^b	4.5 ^b	4.0 ^b	3.4	3.1	3.0	3.0
Sedatives	2.1 ^b	1.8 ^b	1.9 ^b	2.0 ^b	1.9 ^b	1.8 ^b	1.6	1.6	1.4	1.4	1.2	1.3	1.3
ILLCIT DRUGS OTHER THAN MARIJUANA¹	40.1 ^b	40.2 ^b	39.2 ^b	39.4 ^b	39.6 ^b	38.4 ^b	36.9 ^b	37.5 ^b	36.4 ^b	34.6	35.3 ^a	34.1	34.0

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.8A Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	11,000 ^b	10,976 ^b	10,913 ^b	11,117 ^b	11,267 ^b	10,889 ^b	11,090 ^b	12,186	11,990 ^a	12,067 ^a	12,551	12,448	12,597
Marijuana and Hashish	9,238 ^b	9,044 ^b	8,963 ^b	9,111 ^b	9,188 ^b	9,008 ^b	9,154 ^b	10,343 ^b	10,215 ^b	10,554 ^a	10,887	11,000	11,154
Cocaine	2,087 ^b	2,098 ^b	2,134 ^b	2,247 ^b	2,263 ^b	2,087 ^b	1,844 ^a	1,790	1,593	1,563	1,600	1,543	1,604
Crack	266 ^b	275 ^b	263 ^b	328 ^b	293 ^b	249 ^b	217 ^b	172	180	107	140	95	127
Heroin	122 ^b	96 ^b	118 ^b	159 ^b	147 ^b	142 ^b	149 ^b	183 ^a	211	230	272	244	268
Hallucinogens	2,610 ^b	2,112	1,944 ^a	2,051	2,151	2,042	2,101	2,430	2,439	2,323	2,236	2,338	2,205
LSD	553 ^b	336 ^b	317 ^b	322 ^b	380 ^b	375 ^b	507 ^b	523 ^b	549 ^b	586 ^b	622 ^a	708	800
PCP	95 ^b	117 ^b	100 ^b	57	76 ^a	57	30	35	37	63	57	26	36
Ecstasy	1,784 ^b	1,164	999 ^a	998 ^a	1,235	1,146	1,270	1,456 ^a	1,500 ^b	1,413 ^a	1,412	1,379	1,219
Inhalants	685 ^b	662 ^b	674 ^b	685 ^b	578 ^b	528 ^b	538 ^b	635 ^b	597 ^b	500 ^a	473	478	379
Nonmedical Use of Psychotherapeutics ^{2,3}	4,452 ^a	4,637 ^b	4,807 ^b	4,915 ^b	5,130 ^b	4,920 ^b	4,891 ^b	5,061 ^b	4,876 ^b	4,366	4,729 ^b	4,245	4,133
Pain Relievers	3,540 ^b	3,803 ^b	3,834 ^b	4,030 ^b	4,077 ^b	3,981 ^b	3,956 ^b	4,027 ^b	3,779 ^b	3,353 ^b	3,507 ^b	3,061 ^a	2,740
OxyContin [®]	--	--	562 ^b	559 ^a	577 ^b	568 ^b	602 ^b	773 ^b	752 ^b	628 ^b	568 ^a	473	427
Tranquilizers	1,509	1,682 ^a	1,658 ^a	1,796 ^b	1,793 ^b	1,702 ^a	1,762 ^b	1,802 ^b	1,803 ^b	1,555	1,718 ^b	1,568	1,447
Stimulants ³	1,233	1,197	1,280	1,242	1,376	1,132 ^a	1,123 ^b	1,238	1,242	1,065 ^b	1,361	1,302	1,359
Methamphetamine ³	619 ^b	594 ^b	619 ^b	576 ^b	553 ^b	406	270	303	268	225 ^b	341	318	350
Sedatives	162	156	163	191	185	171	178	160	166	128	119	138	153
ILLCIT DRUGS OTHER THAN MARIJUANA¹	6,277 ^b	6,266 ^b	6,201 ^a	6,467 ^b	6,638 ^b	6,374 ^b	6,253 ^b	6,701 ^b	6,395 ^b	5,972	6,217 ^a	5,854	5,780

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.8B Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	35.5	34.6 ^a	33.9 ^b	34.2 ^b	34.4 ^a	33.3 ^b	33.7 ^b	36.3	35.2	35.2	36.3	35.8	36.1
Marijuana and Hashish	29.8 ^b	28.5 ^b	27.8 ^b	28.0 ^b	28.1 ^b	27.5 ^b	27.8 ^b	30.8	30.0 ^b	30.8	31.5	31.6	31.9
Cocaine	6.7 ^b	6.6 ^b	6.6 ^b	6.9 ^b	6.9 ^b	6.4 ^b	5.6 ^b	5.3 ^a	4.7	4.6	4.6	4.4	4.6
Crack	0.9 ^b	0.9 ^b	0.8 ^b	1.0 ^b	0.9 ^b	0.8 ^b	0.7 ^b	0.5	0.5	0.3	0.4	0.3	0.4
Heroin	0.4 ^b	0.3 ^b	0.4 ^b	0.5 ^a	0.4 ^b	0.4 ^b	0.5 ^b	0.5 ^a	0.6	0.7	0.8	0.7	0.8
Hallucinogens	8.4 ^b	6.7	6.0	6.3	6.6	6.2	6.4	7.2 ^b	7.2 ^a	6.8	6.5	6.7	6.3
LSD	1.8 ^a	1.1 ^b	1.0 ^b	1.0 ^b	1.2 ^b	1.1 ^b	1.5 ^b	1.6 ^b	1.6 ^b	1.7 ^b	1.8 ^a	2.0	2.3
PCP	0.3 ^b	0.4 ^b	0.3 ^b	0.2	0.2 ^b	0.2	0.1	0.1	0.1	0.2	0.2	0.1	0.1
Ecstasy	5.8 ^b	3.7	3.1	3.1	3.8	3.5	3.9	4.3 ^b	4.4 ^b	4.1 ^a	4.1 ^a	4.0	3.5
Inhalants	2.2 ^b	2.1 ^b	2.1 ^b	2.1 ^b	1.8 ^b	1.6 ^b	1.6 ^b	1.9 ^b	1.8 ^b	1.5 ^a	1.4	1.4	1.1
Nonmedical Use of Psychotherapeutics ^{2,3}	14.4 ^b	14.6 ^b	14.9 ^b	15.1 ^b	15.7 ^b	15.0 ^b	14.9 ^b	15.1 ^b	14.3 ^b	12.7	13.7 ^b	12.2	11.8
Pain Relievers	11.4 ^b	12.0 ^b	11.9 ^b	12.4 ^b	12.5 ^b	12.2 ^b	12.0 ^b	12.0 ^b	11.1 ^b	9.8 ^b	10.1 ^b	8.8 ^a	7.8
OxyContin [®]	--	--	1.7 ^b	1.7 ^b	1.8 ^b	1.7 ^b	1.8 ^b	2.3 ^b	2.2 ^b	1.8 ^b	1.6 ^a	1.4	1.2
Tranquilizers	4.9 ^b	5.3 ^b	5.2 ^b	5.5 ^b	5.5 ^b	5.2 ^b	5.4 ^b	5.4 ^b	5.3 ^b	4.5	5.0 ^b	4.5	4.1
Stimulants ³	4.0	3.8	4.0	3.8	4.2	3.5	3.4	3.7	3.6	3.1 ^b	3.9	3.7	3.9
Methamphetamine ³	2.0 ^b	1.9 ^b	1.9 ^b	1.8 ^b	1.7 ^b	1.2	0.8	0.9	0.8	0.7 ^b	1.0	0.9	1.0
Sedatives	0.5	0.5	0.5	0.6	0.6	0.5	0.5	0.5	0.5	0.4	0.3	0.4	0.4
ILLCIT DRUGS OTHER THAN MARIJUANA¹	20.2 ^b	19.7 ^b	19.3 ^b	19.9 ^b	20.3 ^b	19.5 ^b	19.0 ^b	20.0 ^b	18.8 ^b	17.4	18.0 ^b	16.8	16.5

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.9A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	6,264 ^b	6,435 ^b	6,239 ^b	6,543 ^b	6,498 ^b	6,475 ^b	6,490 ^b	7,176 ^a	7,345	7,352	7,381	7,490	7,676
Marijuana and Hashish	5,376 ^b	5,400 ^b	5,196 ^b	5,388 ^b	5,337 ^b	5,392 ^b	5,459 ^b	6,128 ^b	6,313 ^a	6,526	6,485	6,636	6,836
Cocaine	631 ^b	696 ^b	692 ^b	832 ^b	730 ^b	572	515	466	503	464	398	398	473
Crack	52	55	80 ^b	104 ^b	72 ^b	56	54	40	72 ^a	20	38	36	29
Heroin	39 ^a	25 ^b	43 ^a	59	55	49	76	67	90	107	138 ^a	92	82
Hallucinogens	589	545	477	503	564	475	545	620	672 ^a	557	600	617	502
LSD	36 ^b	67 ^a	83	60 ^b	58 ^b	69 ^a	98	97	105	92	118	105	118
PCP	15	30 ^a	17	11	14	15	4	5	6	12	12	4	5
Ecstasy	355	230	231	267	326	230	295	378 ^a	402 ^b	315	361	319	270
Inhalants	150 ^b	134 ^a	144 ^b	146 ^a	130 ^a	131 ^a	104	148 ^a	135 ^a	144	123	106	80
Nonmedical Use of Psychotherapeutics ^{2,3}	1,698	1,922 ^b	1,978 ^b	2,046 ^b	2,122 ^b	1,947 ^b	1,936 ^b	2,133 ^b	1,997 ^b	1,711	1,832 ^b	1,661	1,551
Pain Relievers	1,262 ^b	1,486 ^b	1,504 ^b	1,535 ^b	1,627 ^b	1,518 ^b	1,489 ^b	1,610 ^b	1,510 ^b	1,220 ^b	1,311 ^b	1,132	978
OxyContin [®]	--	--	142	146	140	157 ^a	130	179 ^b	239 ^b	158	145	107	102
Tranquilizers	487	525 ^a	566 ^b	606 ^b	651 ^b	573 ^b	570 ^b	622 ^b	570 ^b	546 ^a	537 ^a	435	416
Stimulants ³	409	412	479	468	457	350	351	424	414	344	400	462	406
Methamphetamine ³	183 ^b	197 ^b	220 ^b	225 ^b	183 ^b	116	74	84	83	79	97	120	86
Sedatives	49	58	49	74	54	71	67	84	59	42	52	34	56
ILLCIT DRUGS OTHER THAN MARIJUANA¹	2,459	2,665 ^b	2,619 ^b	2,850 ^b	2,912 ^b	2,642 ^b	2,579 ^b	2,811 ^b	2,709 ^b	2,385	2,435	2,338	2,248

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.9B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	20.2 ^b	20.3 ^b	19.4 ^b	20.1 ^b	19.8 ^b	19.8 ^b	19.7 ^b	21.4	21.6	21.4	21.3	21.5	22.0
Marijuana and Hashish	17.3 ^b	17.0 ^b	16.1 ^b	16.6 ^b	16.3 ^b	16.5 ^b	16.6 ^b	18.2 ^a	18.5	19.0	18.7	19.1	19.6
Cocaine	2.0 ^b	2.2 ^b	2.1 ^b	2.6 ^b	2.2 ^b	1.7 ^a	1.6	1.4	1.5	1.4	1.1	1.1	1.4
Crack	0.2 ^a	0.2	0.3 ^b	0.3 ^b	0.2 ^b	0.2 ^a	0.2	0.1	0.2 ^a	0.1	0.1	0.1	0.1
Heroin	0.1 ^a	0.1 ^b	0.1	0.2	0.2	0.1	0.2	0.2	0.3	0.3	0.4 ^a	0.3	0.2
Hallucinogens	1.9 ^a	1.7	1.5	1.5	1.7	1.5	1.7	1.8 ^a	2.0 ^b	1.6	1.7	1.8	1.4
LSD	0.1 ^b	0.2	0.3	0.2 ^a	0.2 ^a	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.3
PCP	0.0	0.1 ^a	0.1 ^a	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Ecstasy	1.1 ^b	0.7	0.7	0.8	1.0	0.7	0.9	1.1 ^a	1.2 ^b	0.9	1.0	0.9	0.8
Inhalants	0.5 ^b	0.4 ^a	0.4 ^b	0.5 ^b	0.4 ^a	0.4 ^a	0.3	0.4 ^b	0.4 ^a	0.4	0.4	0.3	0.2
Nonmedical Use of Psychotherapeutics ^{2,3}	5.5 ^b	6.1 ^b	6.1 ^b	6.3 ^b	6.5 ^b	5.9 ^b	5.9 ^b	6.4 ^b	5.9 ^b	5.0	5.3 ^b	4.8	4.4
Pain Relievers	4.1 ^b	4.7 ^b	4.7 ^b	4.7 ^b	5.0 ^b	4.6 ^b	4.5 ^b	4.8 ^b	4.4 ^b	3.6 ^b	3.8 ^b	3.3	2.8
OxyContin [®]	--	--	0.4	0.4	0.4	0.5 ^a	0.4	0.5 ^b	0.7 ^b	0.5 ^a	0.4	0.3	0.3
Tranquilizers	1.6 ^a	1.7 ^b	1.8 ^b	1.9 ^b	2.0 ^b	1.7 ^b	1.7 ^b	1.9 ^b	1.7 ^b	1.6 ^a	1.6 ^a	1.2	1.2
Stimulants ³	1.3	1.3	1.5 ^a	1.4	1.4	1.1	1.1	1.3	1.2	1.0	1.2	1.3	1.2
Methamphetamine ³	0.6 ^b	0.6 ^b	0.7 ^b	0.7 ^b	0.6 ^b	0.4	0.2	0.2	0.2	0.2	0.3	0.3	0.2
Sedatives	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA¹	7.9 ^b	8.4 ^b	8.1 ^b	8.8 ^b	8.9 ^b	8.1 ^b	7.8 ^b	8.4 ^b	8.0 ^b	7.0	7.0	6.7	6.4

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.10A Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	82,051 ^b	83,380 ^b	83,448 ^b	85,841 ^b	85,405 ^b	89,040 ^b	92,478 ^b	92,973 ^b	94,117 ^b	95,171 ^b	98,787 ^b	101,838	104,324
Marijuana and Hashish	73,162 ^b	74,620 ^b	75,001 ^b	76,111 ^b	76,213 ^b	79,988 ^b	82,001 ^b	83,070 ^b	84,927 ^b	85,668 ^b	88,955 ^b	92,568	94,732
Cocaine	28,453 ^b	29,488 ^b	28,649 ^b	28,185 ^b	29,593 ^b	30,477 ^b	31,671 ^b	31,339 ^b	32,430 ^b	32,332 ^b	33,156 ^a	33,378	35,097
Crack	7,057 ^b	6,599 ^b	6,580 ^b	6,675 ^b	7,254 ^b	7,524 ^b	7,458 ^b	7,368 ^b	8,262	7,433 ^b	8,327	8,284	8,843
Heroin	3,087 ^b	3,148 ^b	2,538 ^b	2,979 ^b	3,214 ^b	3,253 ^b	3,262 ^b	3,036 ^b	3,483	3,499	3,841	4,155	4,095
Hallucinogens	25,378 ^b	25,736 ^b	26,318 ^b	25,910 ^b	27,675 ^b	27,088 ^b	29,217 ^b	29,970 ^b	30,248 ^b	29,403 ^b	30,982 ^b	32,912	33,253
LSD	18,917 ^b	19,577 ^b	19,185 ^b	18,728 ^b	20,213 ^b	20,137 ^b	21,166	21,138	20,950	20,716 ^a	21,430	22,303	22,293
PCP	6,352	5,955	5,851	5,705	5,865	5,533	6,032	5,692	5,731	5,676	6,112	6,090	6,023
Ecstasy	4,656 ^b	5,619 ^b	6,168 ^b	6,632 ^b	7,414 ^b	7,786 ^b	8,431 ^b	9,513 ^b	11,082 ^b	9,743 ^b	11,222 ^b	13,009	13,051
Inhalants	15,398 ^b	15,596 ^b	15,521 ^b	15,754 ^b	16,200 ^a	16,426	16,659	16,649	16,367	15,540 ^b	16,563	17,129	17,543
Nonmedical Use of Psychotherapeutics ^{2,3}	35,779 ^b	36,269 ^b	36,190 ^b	36,562 ^b	37,741 ^b	37,781 ^b	39,677 ^b	39,285 ^b	39,409 ^b	39,348 ^b	42,198	41,745	42,909
Pain Relievers	19,983 ^b	20,896 ^b	21,076 ^b	21,896 ^b	22,475 ^b	22,533 ^b	24,543 ^b	24,583 ^b	24,521 ^b	24,497 ^b	27,218	26,417	27,211
OxyContin [®]	881 ^b	1,433 ^b	1,392 ^b	1,589 ^b	2,089 ^b	2,303 ^b	2,748 ^b	3,373 ^b	3,642 ^b	3,469 ^b	4,247 ^a	4,811	4,996
Tranquilizers	14,959 ^b	15,447 ^b	15,117 ^b	15,919 ^b	16,290 ^b	15,350 ^b	16,762 ^b	16,830 ^b	16,960 ^b	16,725 ^b	18,515	18,737	19,823
Stimulants ³	18,588	18,085	17,499 ^a	16,105 ^b	17,639	17,369 ^a	17,126 ^a	17,863	17,674	16,699 ^b	17,629	17,940	18,879
Methamphetamine ³	12,648	12,565	11,920	10,154 ^b	11,782	11,012	10,891	11,203	11,542	10,594 ^a	11,025	11,096	11,731
Sedatives	9,075 ^b	8,696 ^b	9,012 ^b	8,101 ^a	8,007	7,604	8,204 ^a	7,882	7,006	6,899	7,344	6,916	7,160
ILLCIT DRUGS OTHER THAN MARIJUANA¹	52,578 ^b	53,040 ^b	52,672 ^b	54,078 ^b	55,024 ^b	56,431 ^b	59,248 ^b	58,987 ^b	59,895 ^b	59,580 ^b	62,127 ^a	62,945	64,936

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.10B Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	45.7 ^b	46.1 ^b	45.6 ^b	46.3 ^b	45.5 ^b	46.9 ^b	48.2 ^b	48.0 ^b	48.2 ^b	48.0 ^b	49.3 ^a	50.2	50.8
Marijuana and Hashish	40.8 ^b	41.2 ^b	41.0 ^b	41.1 ^b	40.6 ^b	42.1 ^b	42.7 ^b	42.9 ^b	43.5 ^b	43.2 ^b	44.4 ^b	45.7	46.1
Cocaine	15.9 ^b	16.3	15.6 ^b	15.2 ^b	15.7 ^b	16.1 ^a	16.5	16.2 ^a	16.6	16.3	16.5	16.5	17.1
Crack	3.9	3.6 ^b	3.6 ^b	3.6 ^b	3.9	4.0	3.9	3.8 ^a	4.2	3.7 ^a	4.2	4.1	4.3
Heroin	1.7	1.7	1.4 ^b	1.6 ^a	1.7	1.7	1.7	1.6 ^b	1.8	1.8	1.9	2.0	2.0
Hallucinogens	14.1 ^b	14.2 ^b	14.4 ^b	14.0 ^b	14.7 ^b	14.3 ^b	15.2 ^a	15.5	15.5	14.8 ^b	15.4	16.2	16.2
LSD	10.5	10.8	10.5	10.1 ^a	10.8	10.6	11.0	10.9	10.7	10.4	10.7	11.0	10.9
PCP	3.5 ^b	3.3	3.2	3.1	3.1	2.9	3.1	2.9	2.9	2.9	3.0	3.0	2.9
Ecstasy	2.6 ^b	3.1 ^b	3.4 ^b	3.6 ^b	3.9 ^b	4.1 ^b	4.4 ^b	4.9 ^b	5.7 ^a	4.9 ^b	5.6 ^b	6.4	6.4
Inhalants	8.6	8.6	8.5	8.5	8.6	8.7	8.7	8.6	8.4	7.8 ^a	8.3	8.4	8.5
Nonmedical Use of Psychotherapeutics ^{2,3}	19.9	20.0	19.8 ^a	19.7 ^a	20.1	19.9 ^a	20.7	20.3	20.2	19.8 ^a	21.0	20.6	20.9
Pain Relievers	11.1 ^b	11.5 ^b	11.5 ^b	11.8 ^b	12.0 ^b	11.9 ^b	12.8	12.7	12.6	12.4 ^a	13.6	13.0	13.3
OxyContin [®]	0.5 ^b	0.8 ^b	0.8 ^b	0.9 ^b	1.1 ^b	1.2 ^b	1.4 ^b	1.7 ^b	1.9 ^b	1.7 ^b	2.1 ^a	2.4	2.4
Tranquilizers	8.3 ^b	8.5 ^b	8.3 ^b	8.6 ^b	8.7 ^b	8.1 ^b	8.7 ^b	8.7 ^b	8.7 ^b	8.4 ^b	9.2	9.2	9.7
Stimulants ³	10.4 ^b	10.0 ^a	9.6	8.7	9.4	9.1	8.9	9.2	9.1	8.4 ^a	8.8	8.8	9.2
Methamphetamine ³	7.1 ^b	6.9 ^b	6.5 ^b	5.5	6.3 ^a	5.8	5.7	5.8	5.9	5.3	5.5	5.5	5.7
Sedatives	5.1 ^b	4.8 ^b	4.9 ^b	4.4 ^b	4.3 ^b	4.0 ^a	4.3 ^b	4.1 ^a	3.6	3.5	3.7	3.4	3.5
ILLCIT DRUGS OTHER THAN MARIJUANA¹	29.3 ^b	29.3 ^b	28.8 ^b	29.2 ^b	29.3 ^b	29.7 ^b	30.9	30.5 ^a	30.7	30.0 ^b	31.0	31.1	31.6

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.11A Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	18,639 ^b	18,562 ^b	18,594 ^b	18,883 ^b	19,565 ^b	20,163 ^b	19,801 ^b	21,099 ^b	22,174 ^b	21,485 ^b	24,460 ^b	24,856 ^b	27,223
Marijuana and Hashish	12,612 ^b	12,441 ^b	12,833 ^b	12,880 ^b	12,889 ^b	12,970 ^b	13,479 ^b	14,983 ^b	15,672 ^b	15,632 ^b	17,263 ^b	18,606 ^b	20,707
Cocaine	3,307 ^a	3,372 ^a	3,114	2,852	3,400 ^a	3,275	3,133	2,773	2,700	2,065 ^b	2,890	2,508	2,782
Crack	1,188 ^b	1,036 ^b	978 ^a	1,000 ^b	1,111 ^b	1,138 ^b	862	807	686	490	762	526	627
Heroin	231 ^b	182 ^b	232 ^b	184 ^b	376 ^a	207 ^b	265 ^b	364 ^a	382 ^a	336 ^b	366 ^a	406 ^a	618
Hallucinogens	1,194 ^b	1,046 ^b	1,182 ^a	1,093 ^b	1,166 ^b	1,089 ^b	881 ^b	1,369	1,356	1,098 ^b	1,516	1,610	1,624
LSD	131 ^a	73 ^b	117 ^b	91 ^b	181	120 ^a	115 ^b	124 ^b	197	135 ^a	283	253	275
PCP	31	13	38	22	52	26	*	42	24	17	67	35	21
Ecstasy	836	626 ^a	606 ^a	705	595 ^b	647 ^a	499 ^b	932	687 ^a	590 ^b	910	977	942
Inhalants	320 ^b	295 ^b	413 ^b	358 ^b	532	562	540	514	566	541	573	575	725
Nonmedical Use of Psychotherapeutics ^{2,3}	8,057 ^b	8,231 ^a	7,814 ^b	8,331 ^a	9,168	9,377	8,380 ^a	9,108	9,362	8,542	10,296 ^a	9,662	9,280
Pain Relievers	5,582 ^a	5,950	5,556 ^a	6,044	6,752	6,851	6,347	6,795	6,941	6,306	7,649 ^b	6,865	6,432
OxyContin [®]	--	--	457 ^a	480	547	626	634	687	893	776	723	819	649
Tranquilizers	2,772 ^a	2,783 ^a	2,878	2,956	2,788 ^a	3,086	2,853	3,228	3,314	3,080	3,903 ^a	3,354	3,331
Stimulants ³	1,482 ^b	1,256 ^b	1,467 ^b	1,340 ^b	1,864	1,447 ^b	1,152 ^b	1,521 ^a	1,335 ^b	1,336 ^b	1,632	1,917	1,995
Methamphetamine ³	891	823	1,012	849	1,146	823	490 ^b	777	593	703	732	794	845
Sedatives	681	544	441	441	632	582	335 ^a	530	630	314 ^a	390	437	506
ILLCIT DRUGS OTHER THAN MARIJUANA¹	10,808 ^a	10,689 ^a	10,189 ^b	10,513 ^b	11,477	11,910	10,906	11,526	11,472	10,422 ^b	12,753	12,035	11,864

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.11B Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	10.4 ^b	10.3 ^b	10.2 ^b	10.2 ^b	10.4 ^b	10.6 ^b	10.3 ^b	10.9 ^b	11.4 ^b	10.8 ^b	12.2 ^b	12.3 ^a	13.3
Marijuana and Hashish	7.0 ^b	6.9 ^b	7.0 ^b	6.9 ^b	6.9 ^b	6.8 ^b	7.0 ^b	7.7 ^b	8.0 ^b	7.9 ^b	8.6 ^b	9.2 ^b	10.1
Cocaine	1.8 ^b	1.9 ^b	1.7 ^b	1.5	1.8 ^b	1.7 ^b	1.6 ^a	1.4	1.4	1.0 ^b	1.4	1.2	1.4
Crack	0.7 ^b	0.6 ^b	0.5 ^b	0.5 ^b	0.6 ^b	0.6 ^b	0.4 ^a	0.4	0.4	0.2	0.4	0.3	0.3
Heroin	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.2	0.1 ^b	0.1 ^b	0.2 ^a	0.2 ^a	0.2 ^b	0.2 ^a	0.2 ^a	0.3
Hallucinogens	0.7	0.6 ^a	0.6	0.6 ^a	0.6 ^a	0.6 ^b	0.5 ^b	0.7	0.7	0.6 ^b	0.8	0.8	0.8
LSD	0.1 ^a	0.0 ^b	0.1 ^b	0.0 ^b	0.1	0.1 ^a	0.1 ^b	0.1 ^b	0.1	0.1 ^a	0.1	0.1	0.1
PCP	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0
Ecstasy	0.5	0.3	0.3	0.4	0.3 ^a	0.3	0.3 ^b	0.5	0.4	0.3 ^b	0.5	0.5	0.5
Inhalants	0.2 ^b	0.2 ^b	0.2 ^a	0.2 ^b	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4
Nonmedical Use of Psychotherapeutics ^{2,3}	4.5	4.5	4.3	4.5	4.9	4.9	4.4	4.7	4.8	4.3	5.1 ^b	4.8	4.5
Pain Relievers	3.1	3.3	3.0	3.3	3.6 ^a	3.6 ^a	3.3	3.5 ^a	3.6 ^a	3.2	3.8 ^b	3.4	3.1
OxyContin [®]	--	--	0.2	0.3	0.3	0.3	0.3	0.4	0.5 ^a	0.4	0.4	0.4	0.3
Tranquilizers	1.5	1.5	1.6	1.6	1.5	1.6	1.5	1.7	1.7	1.6	1.9 ^a	1.7	1.6
Stimulants ³	0.8	0.7 ^b	0.8	0.7 ^b	1.0	0.8 ^a	0.6 ^b	0.8	0.7 ^b	0.7 ^b	0.8	0.9	1.0
Methamphetamine ³	0.5	0.5	0.6	0.5	0.6 ^a	0.4	0.3 ^b	0.4	0.3	0.4	0.4	0.4	0.4
Sedatives	0.4	0.3	0.2	0.2	0.3	0.3	0.2	0.3	0.3	0.2 ^a	0.2	0.2	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA¹	6.0	5.9	5.6	5.7	6.1	6.3	5.7	6.0	5.9	5.3 ^a	6.4 ^a	5.9	5.8

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.12A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	10,380 ^b	10,224 ^b	10,159 ^b	10,666 ^b	11,400 ^b	11,001 ^b	11,321 ^b	12,270 ^b	12,835 ^b	12,580 ^b	14,105 ^b	14,886 ^b	16,969
Marijuana and Hashish	7,185 ^b	7,267 ^b	7,471 ^b	7,508 ^b	7,818 ^b	7,385 ^b	8,142 ^b	8,885 ^b	9,294 ^b	9,578 ^b	10,571 ^b	11,412 ^b	13,521
Cocaine	1,243	1,431 ^a	1,204	1,423 ^a	1,586 ^b	1,400 ^a	1,252	1,106	909	838	1,222	1,108	1,018
Crack	488 ^a	517	370	553 ^a	619 ^b	538 ^a	302	448	303	205	405	338	317
Heroin	116 ^b	76 ^b	108 ^b	62 ^b	268	109 ^b	124 ^b	112 ^b	141 ^b	159 ^b	197 ^a	185 ^a	337
Hallucinogens	352	238 ^b	241 ^b	380	262 ^b	349	269 ^b	427	320 ^a	200 ^b	374	562	535
LSD	15 ^a	15 ^b	20 ^a	12 ^b	41	45	*	27 ^a	*	26 ^a	39	72	104
PCP	20	*	21	20	*	19	*	36	24	*	16	22	*
Ecstasy	197	149 ^a	142 ^a	161 ^a	120 ^b	209	146 ^a	269	165	120 ^b	193	292	300
Inhalants	183	117 ^b	181	152 ^a	292	191	270	170	293	257	182	269	316
Nonmedical Use of Psychotherapeutics ^{2,3}	3,602 ^a	3,532 ^a	3,218 ^b	3,607 ^a	4,130	4,130	3,563 ^b	4,072	4,239	3,720 ^a	4,306	4,274	4,331
Pain Relievers	2,321 ^a	2,406	2,149 ^b	2,437	2,908	2,998	2,676	3,014	2,975	2,688	2,993	2,964	2,879
OxyContin [®]	--	--	118	153	106 ^a	168	256	262	280	225	165	355	219
Tranquilizers	1,129	1,078	889 ^b	1,050 ^a	984 ^a	1,085	1,075	1,265	1,456	1,132	1,462	1,180	1,356
Stimulants ³	685 ^a	676 ^a	653 ^b	548 ^b	757	574 ^b	445 ^b	748	561 ^b	528 ^b	654 ^a	819	1,021
Methamphetamine ³	428	459	428	331	495	380	211 ^a	383	245 ^a	323	312	453	438
Sedatives	340	189	190	171	285	240	136	243	280	164	192	192	233
ILLCIT DRUGS OTHER THAN MARIJUANA¹	4,897	4,767 ^a	4,301 ^b	4,860 ^a	5,456	5,471	4,884 ^a	5,262	5,214	4,610 ^b	5,530	5,570	5,587

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.12B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	5.8 ^b	5.6 ^b	5.5 ^b	5.8 ^b	6.1 ^b	5.8 ^b	5.9 ^b	6.3 ^b	6.6 ^b	6.3 ^b	7.0 ^b	7.3 ^b	8.3
Marijuana and Hashish	4.0 ^b	4.0 ^b	4.1 ^b	4.1 ^b	4.2 ^b	3.9 ^b	4.2 ^b	4.6 ^b	4.8 ^b	4.8 ^b	5.3 ^b	5.6 ^b	6.6
Cocaine	0.7 ^a	0.8 ^b	0.7 ^a	0.8 ^b	0.8 ^b	0.7 ^b	0.7 ^a	0.6	0.5	0.4	0.6	0.5	0.5
Crack	0.3 ^b	0.3 ^a	0.2	0.3 ^b	0.3 ^b	0.3 ^b	0.2	0.2	0.2	0.1	0.2	0.2	0.2
Heroin	0.1 ^b	0.0 ^b	0.1 ^b	0.0 ^b	0.1	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^a	0.1	0.1 ^a	0.2
Hallucinogens	0.2	0.1 ^b	0.1 ^b	0.2	0.1 ^b	0.2	0.1 ^b	0.2	0.2 ^a	0.1 ^b	0.2	0.3	0.3
LSD	0.0 ^a	0.0 ^a	0.0 ^a	0.0 ^b	0.0	0.0	*	0.0 ^a	*	0.0 ^a	0.0	0.0	0.1
PCP	0.0	*	0.0	0.0	*	0.0	*	0.0	0.0	*	0.0	0.0	*
Ecstasy	0.1	0.1	0.1 ^a	0.1	0.1 ^a	0.1	0.1 ^a	0.1	0.1	0.1 ^b	0.1	0.1	0.1
Inhalants	0.1	0.1 ^b	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2
Nonmedical Use of Psychotherapeutics ^{2,3}	2.0	2.0	1.8 ^a	1.9	2.2	2.2	1.9	2.1	2.2	1.9	2.1	2.1	2.1
Pain Relievers	1.3	1.3	1.2 ^a	1.3	1.5	1.6	1.4	1.6	1.5	1.4	1.5	1.5	1.4
OxyContin [®]	--	--	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1
Tranquilizers	0.6	0.6	0.5 ^a	0.6	0.5	0.6	0.6	0.7	0.7	0.6	0.7	0.6	0.7
Stimulants ³	0.4	0.4	0.4 ^a	0.3 ^b	0.4	0.3 ^b	0.2 ^b	0.4	0.3 ^b	0.3 ^b	0.3 ^a	0.4	0.5
Methamphetamine ³	0.2	0.3	0.2	0.2	0.3	0.2	0.1 ^a	0.2	0.1 ^a	0.2	0.2	0.2	0.2
Sedatives	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	2.7	2.6	2.3 ^a	2.6	2.9	2.9	2.5	2.7	2.7	2.3 ^a	2.8	2.7	2.7

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.13A Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	100,612 ^b	102,574 ^b	102,491 ^b	105,063 ^b	104,733 ^b	107,871 ^b	111,250 ^b	112,631 ^b	113,655 ^b	114,705 ^b	118,764 ^b	121,664 ^a	124,541
Marijuana and Hashish	89,842 ^b	91,723 ^b	91,984 ^b	93,143 ^b	93,393 ^b	96,641 ^b	98,720 ^b	100,747 ^b	102,456 ^b	103,464 ^b	107,007 ^b	110,619	113,121
Cocaine	33,239 ^b	34,243 ^b	33,542 ^b	33,090 ^b	34,732 ^b	35,402 ^b	36,443 ^b	36,337 ^b	36,998 ^a	36,584 ^a	37,423	37,419	38,978
Crack	8,239 ^b	7,792 ^b	7,715 ^b	7,820 ^b	8,430 ^a	8,509	8,374 ^a	8,332 ^a	9,148	8,146 ^b	8,981	8,847	9,389
Heroin	3,574 ^b	3,664 ^b	3,067 ^b	3,475 ^b	3,728 ^b	3,751 ^b	3,729 ^b	3,623 ^b	4,097 ^a	4,093 ^a	4,505	4,774	4,777
Hallucinogens	32,895 ^b	33,113 ^b	33,173 ^b	32,730 ^b	34,300 ^b	33,380 ^b	35,093 ^b	36,388 ^b	36,543 ^b	35,432 ^b	37,085 ^a	39,034	39,037
LSD	23,845	24,027	23,085 ^a	22,145 ^b	23,146 ^a	22,518 ^b	23,333	23,464	23,143 ^a	22,771 ^b	23,487	24,549	24,744
PCP	7,185 ^a	6,903	6,587	6,421	6,505	6,073	6,554	6,173	6,179	6,027	6,456	6,416	6,328
Ecstasy	9,330 ^b	10,301 ^b	10,607 ^b	11,096 ^b	11,806 ^b	11,971 ^b	12,439 ^b	13,707 ^b	15,312 ^b	13,964 ^b	15,673 ^b	17,445	17,251
Inhalants	20,265	20,325	20,036	20,087	20,298	20,130	20,104	20,269	19,763	18,646 ^a	19,482	19,741	19,986
Nonmedical Use of Psychotherapeutics ^{2,3}	44,556 ^b	45,629 ^b	45,754 ^b	46,552 ^b	47,809 ^b	47,588 ^b	49,371 ^b	49,181 ^b	49,241 ^b	48,721 ^b	51,908	50,984	52,105
Pain Relievers	26,836 ^b	28,409 ^b	28,890 ^b	30,175 ^b	30,828 ^b	30,676 ^b	32,639 ^a	32,804	32,664	32,097 ^b	34,977	33,668	34,261
OxyContin [®]	1,700 ^b	2,579 ^b	2,767 ^b	3,205 ^b	3,773 ^b	4,019 ^b	4,513 ^b	5,483 ^b	5,791 ^b	5,547 ^b	6,257	6,699	6,779
Tranquilizers	18,433 ^b	19,346 ^b	19,036 ^b	20,275 ^b	20,576 ^b	19,505 ^b	20,831 ^b	21,152 ^b	21,463 ^b	20,956 ^b	22,936	22,921	24,185
Stimulants ³	22,384	21,958	21,391	20,113 ^b	21,565	20,939	20,612 ^a	21,450	21,209	19,871 ^b	21,003	21,219	22,008
Methamphetamine ³	14,949 ^b	14,756 ^b	14,166 ^b	12,344	13,885 ^a	12,859	12,438	12,721	12,903	11,743	12,114	12,138	12,771
Sedatives	9,723 ^b	9,261 ^b	9,640 ^b	8,751 ^a	8,619 ^a	8,188	8,715 ^a	8,409	7,475	7,371	7,764	7,355	7,620
ILLCIT DRUGS OTHER THAN MARIJUANA¹	65,014 ^b	65,809 ^b	65,302 ^b	66,879 ^b	67,984 ^b	69,005 ^b	71,394 ^b	71,579 ^b	72,306 ^b	71,440 ^b	74,343 ^a	74,797	76,798

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.13B Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	47.8 ^b	48.2 ^b	47.6 ^b	48.2 ^b	47.5 ^b	48.5 ^b	49.5 ^b	49.6 ^b	49.6 ^b	49.3 ^b	50.5 ^a	51.2	51.8
Marijuana and Hashish	42.7 ^b	43.1 ^b	42.7 ^b	42.8 ^b	42.3 ^b	43.4 ^b	43.9 ^b	44.3 ^b	44.7 ^b	44.5 ^b	45.5 ^b	46.6	47.1
Cocaine	15.8	16.1	15.6	15.2 ^b	15.7	15.9	16.2	16.0	16.1	15.7	15.9	15.8	16.2
Crack	3.9	3.7	3.6	3.6	3.8	3.8	3.7	3.7	4.0	3.5 ^a	3.8	3.7	3.9
Heroin	1.7 ^a	1.7 ^a	1.4 ^b	1.6 ^b	1.7 ^a	1.7 ^a	1.7 ^a	1.6 ^b	1.8	1.8	1.9	2.0	2.0
Hallucinogens	15.6	15.6	15.4 ^a	15.0 ^b	15.5	15.0 ^b	15.6	16.0	15.9	15.2 ^b	15.8	16.4	16.2
LSD	11.3 ^b	11.3 ^b	10.7	10.2	10.5	10.1	10.4	10.3	10.1	9.8	10.0	10.3	10.3
PCP	3.4 ^b	3.2 ^b	3.1 ^a	2.9	2.9	2.7	2.9	2.7	2.7	2.6	2.7	2.7	2.6
Ecstasy	4.4 ^b	4.8 ^b	4.9 ^b	5.1 ^b	5.4 ^b	5.4 ^b	5.5 ^b	6.0 ^b	6.7 ^a	6.0 ^b	6.7 ^a	7.3	7.2
Inhalants	9.6 ^b	9.6 ^b	9.3 ^b	9.2 ^b	9.2 ^b	9.0 ^a	8.9 ^a	8.9 ^a	8.6	8.0	8.3	8.3	8.3
Nonmedical Use of Psychotherapeutics ^{2,3}	21.2	21.5	21.3	21.4	21.7	21.4	22.0	21.6	21.5	20.9	22.1	21.5	21.7
Pain Relievers	12.8 ^b	13.4 ^b	13.4 ^a	13.9	14.0	13.8	14.5	14.4	14.2	13.8	14.9	14.2	14.3
OxyContin [®]	0.8 ^b	1.2 ^b	1.3 ^b	1.5 ^b	1.7 ^b	1.8 ^b	2.0 ^b	2.4 ^b	2.5 ^a	2.4 ^b	2.7	2.8	2.8
Tranquilizers	8.8 ^b	9.1 ^b	8.8 ^b	9.3 ^b	9.3 ^b	8.8 ^b	9.3 ^b	9.3 ^b	9.4 ^a	9.0 ^b	9.8	9.7	10.1
Stimulants ³	10.6 ^b	10.3 ^b	9.9 ^b	9.2	9.8 ^a	9.4	9.2	9.4	9.3	8.5 ^a	8.9	8.9	9.2
Methamphetamine ³	7.1 ^b	6.9 ^b	6.6 ^b	5.7	6.3 ^b	5.8 ^a	5.5	5.6	5.6	5.0	5.2	5.1	5.3
Sedatives	4.6 ^b	4.4 ^b	4.5 ^b	4.0 ^b	3.9 ^b	3.7 ^a	3.9 ^b	3.7 ^a	3.3	3.2	3.3	3.1	3.2
ILLCIT DRUGS OTHER THAN MARIJUANA¹	30.9 ^a	30.9 ^a	30.3 ^b	30.7 ^a	30.8 ^a	31.0	31.7	31.5	31.5	30.7 ^a	31.6	31.5	32.0

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.14A Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	29,638 ^b	29,538 ^b	29,506 ^b	29,999 ^b	30,831 ^b	31,052 ^b	30,892 ^b	33,285 ^b	34,163 ^b	33,552 ^b	37,011 ^b	37,304 ^b	39,821
Marijuana and Hashish	21,850 ^b	21,486 ^b	21,796 ^b	21,991 ^b	22,077 ^b	21,978 ^b	22,633 ^b	25,325 ^b	25,887 ^b	26,186 ^b	28,150 ^b	29,606 ^b	31,862
Cocaine	5,394 ^b	5,470 ^b	5,247 ^b	5,099 ^b	5,663 ^b	5,362 ^b	4,977 ^a	4,564	4,294	3,628 ^b	4,491	4,051	4,385
Crack	1,454 ^b	1,310 ^b	1,241 ^b	1,327 ^b	1,404 ^b	1,387 ^b	1,079 ^b	979	866	597	902	621	754
Heroin	353 ^b	278 ^b	350 ^b	343 ^b	524 ^b	349 ^b	415 ^b	547 ^b	593 ^b	566 ^b	638 ^a	650 ^a	885
Hallucinogens	3,804	3,158 ^b	3,125 ^b	3,144 ^b	3,317 ^a	3,130 ^b	2,982 ^b	3,799	3,795	3,421 ^a	3,753	3,948	3,829
LSD	684 ^b	409 ^b	434 ^b	413 ^b	561 ^b	495 ^b	623 ^b	647 ^b	746 ^b	721 ^b	905	962	1,075
PCP	126 ^b	130 ^b	138 ^b	79	129	83	37	77	61	80	124	61	57
Ecstasy	2,621 ^b	1,790 ^a	1,605 ^b	1,703 ^b	1,830 ^a	1,794 ^a	1,770 ^b	2,388	2,188	2,003	2,322	2,356	2,160
Inhalants	1,005	956	1,087	1,043	1,110	1,090	1,079	1,149	1,163	1,041	1,046	1,053	1,104
Nonmedical Use of Psychotherapeutics ^{2,3}	12,509	12,868	12,621	13,246	14,298	14,297	13,271	14,169	14,238	12,908	15,025 ^b	13,907	13,412
Pain Relievers	9,122	9,753	9,391	10,074 ^a	10,829 ^b	10,833 ^b	10,304 ^b	10,822 ^b	10,720 ^b	9,659	11,157 ^b	9,925	9,172
OxyContin [®]	--	--	1,019	1,039	1,124	1,195	1,236	1,460 ^b	1,645 ^b	1,404 ^a	1,290	1,292	1,075
Tranquilizers	4,281	4,465	4,536	4,752	4,581	4,788	4,616	5,031	5,117	4,635	5,621 ^b	4,922	4,778
Stimulants ³	2,716 ^b	2,453 ^b	2,747 ^b	2,581 ^b	3,241	2,579 ^b	2,275 ^b	2,759 ^b	2,577 ^b	2,401 ^b	2,992	3,219	3,354
Methamphetamine ³	1,510 ^a	1,417	1,631 ^b	1,425	1,700 ^b	1,229	760 ^b	1,080	861 ^a	928	1,073	1,113	1,195
Sedatives	843	700	604	632	818	754	513	690	796	442 ^a	509	575	659
ILLCIT DRUGS OTHER THAN MARIJUANA¹	17,086	16,954	16,389 ^a	16,980	18,114	18,283	17,159	18,228	17,866	16,395 ^a	18,970 ^a	17,889	17,643

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.14B Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	14.1 ^b	13.9 ^b	13.7 ^b	13.8 ^b	14.0 ^b	13.9 ^b	13.7 ^b	14.6 ^b	14.9 ^b	14.4 ^b	15.7 ^a	15.7 ^a	16.6
Marijuana and Hashish	10.4 ^b	10.1 ^b	10.1 ^b	10.1 ^b	10.0 ^b	9.9 ^b	10.1 ^b	11.1 ^b	11.3 ^b	11.3 ^b	12.0 ^b	12.5 ^a	13.3
Cocaine	2.6 ^b	2.6 ^b	2.4 ^b	2.3 ^b	2.6 ^b	2.4 ^b	2.2 ^b	2.0	1.9	1.6 ^a	1.9	1.7	1.8
Crack	0.7 ^b	0.6 ^b	0.6 ^b	0.6 ^b	0.6 ^b	0.6 ^b	0.5 ^b	0.4 ^a	0.4	0.3	0.4	0.3	0.3
Heroin	0.2 ^b	0.1 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.3 ^a	0.2 ^b	0.3 ^a	0.3 ^a	0.4
Hallucinogens	1.8 ^a	1.5	1.5	1.4	1.5	1.4 ^a	1.3 ^b	1.7	1.7	1.5	1.6	1.7	1.6
LSD	0.3 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.3 ^b	0.2 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.4	0.4	0.4
PCP	0.1 ^b	0.1 ^b	0.1 ^b	0.0	0.1 ^a	0.0	0.0	0.0	0.0	0.0	0.1 ^a	0.0	0.0
Ecstasy	1.2 ^b	0.8	0.7 ^a	0.8	0.8	0.8	0.8	1.1 ^a	1.0	0.9	1.0	1.0	0.9
Inhalants	0.5	0.4	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.5
Nonmedical Use of Psychotherapeutics ^{2,3}	5.9	6.1 ^a	5.9	6.1 ^a	6.5 ^b	6.4 ^b	5.9	6.2 ^b	6.2 ^b	5.5	6.4 ^b	5.9	5.6
Pain Relievers	4.3 ^b	4.6 ^b	4.4 ^b	4.6 ^b	4.9 ^b	4.9 ^b	4.6 ^b	4.8 ^b	4.7 ^b	4.2 ^a	4.7 ^b	4.2 ^a	3.8
OxyContin [®]	--	--	0.5	0.5	0.5	0.5	0.5 ^a	0.6 ^b	0.7 ^b	0.6 ^b	0.5	0.5	0.4
Tranquilizers	2.0	2.1	2.1	2.2	2.1	2.2	2.1	2.2	2.2 ^a	2.0	2.4 ^b	2.1	2.0
Stimulants ³	1.3	1.2 ^b	1.3	1.2 ^a	1.5	1.2 ^a	1.0 ^b	1.2 ^a	1.1 ^b	1.0 ^b	1.3	1.4	1.4
Methamphetamine ³	0.7 ^b	0.7 ^b	0.8 ^b	0.7 ^a	0.8 ^b	0.6	0.3 ^b	0.5	0.4	0.4	0.5	0.5	0.5
Sedatives	0.4 ^a	0.3	0.3	0.3	0.4	0.3	0.2	0.3	0.3	0.2 ^a	0.2	0.2	0.3
ILLCIT DRUGS OTHER THAN MARIJUANA¹	8.1 ^b	8.0 ^a	7.6	7.8	8.2 ^b	8.2 ^b	7.6	8.0 ^b	7.8	7.0	8.1 ^b	7.5	7.3

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.15A Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	16,644 ^b	16,659 ^b	16,397 ^b	17,209 ^b	17,898 ^b	17,477 ^b	17,811 ^b	19,446 ^b	20,180 ^b	19,932 ^b	21,486 ^b	22,376 ^b	24,646
Marijuana and Hashish	12,561 ^b	12,667 ^b	12,667 ^b	12,896 ^b	13,154 ^b	12,777 ^b	13,601 ^b	15,013 ^b	15,607 ^b	16,103 ^b	17,055 ^b	18,048 ^b	20,357
Cocaine	1,874 ^a	2,128 ^b	1,895 ^a	2,255 ^b	2,316 ^b	1,972 ^b	1,767	1,572	1,412	1,302	1,620	1,505	1,491
Crack	541 ^a	572 ^a	451	657 ^b	691 ^b	595 ^a	356	489	375	225	443	375	347
Heroin	155 ^b	101 ^b	151 ^b	121 ^b	323	157 ^b	200 ^b	180 ^b	231 ^b	267 ^a	335	277 ^a	419
Hallucinogens	941	783 ^a	719 ^b	883	826	824	814 ^a	1,047	992	757 ^b	974	1,179	1,037
LSD	51 ^b	82 ^b	103 ^b	72 ^b	99 ^b	114 ^a	105 ^b	124 ^a	109 ^b	118 ^a	158	177	222
PCP	35	35	37	31	19	34	*	42	30	22	27	26	*
Ecstasy	553	379 ^a	373 ^a	428	446	438	441	647	567	435	554	611	570
Inhalants	334	250 ^a	325	298	422	323	374	318	428	402	305	375	397
Nonmedical Use of Psychotherapeutics ^{2,3}	5,300	5,453	5,196 ^a	5,653	6,252	6,077	5,499	6,205	6,236	5,431	6,138	5,935	5,882
Pain Relievers	3,583	3,892	3,653	3,972	4,534 ^a	4,517 ^a	4,165	4,624 ^b	4,485 ^a	3,908	4,304	4,096	3,858
OxyContin [®]	--	--	260	299	246	324	387	440	519 ^a	383	310	462	321
Tranquilizers	1,617	1,604	1,455 ^a	1,657	1,636	1,657	1,646	1,887	2,026	1,678	1,999	1,615	1,772
Stimulants ³	1,094 ^a	1,088 ^a	1,132 ^a	1,016 ^b	1,214	924 ^b	796 ^b	1,172	975 ^b	872 ^b	1,053 ^a	1,281	1,427
Methamphetamine ³	612	655	648	556	678	496	285 ^a	466	328 ^a	402	409	573	524
Sedatives	389	247	239	245	340	311	203	327	340	206	244	226	289
ILLCIT DRUGS OTHER THAN MARIJUANA¹	7,356	7,432	6,920 ^a	7,710	8,368	8,113	7,463	8,073	7,924	6,994 ^a	7,965	7,908	7,835

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.15B Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	7.9 ^b	7.8 ^b	7.6 ^b	7.9 ^b	8.1 ^b	7.9 ^b	7.9 ^b	8.6 ^b	8.8 ^b	8.6 ^b	9.1 ^b	9.4 ^b	10.3
Marijuana and Hashish	6.0 ^b	6.0 ^b	5.9 ^b	5.9 ^b	6.0 ^b	5.7 ^b	6.0 ^b	6.6 ^b	6.8 ^b	6.9 ^b	7.3 ^b	7.6 ^b	8.5
Cocaine	0.9 ^b	1.0 ^b	0.9 ^b	1.0 ^b	1.0 ^b	0.9 ^b	0.8 ^a	0.7	0.6	0.6	0.7	0.6	0.6
Crack	0.3 ^b	0.3 ^a	0.2	0.3 ^b	0.3 ^b	0.3 ^b	0.2	0.2	0.2	0.1	0.2	0.2	0.1
Heroin	0.1 ^b	0.0 ^b	0.1 ^b	0.1 ^b	0.1	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^a	0.1	0.1	0.1 ^a	0.2
Hallucinogens	0.4	0.4	0.3 ^a	0.4	0.4	0.4	0.4	0.5	0.4	0.3 ^a	0.4	0.5	0.4
LSD	0.0 ^b	0.0 ^b	0.0 ^a	0.0 ^b	0.0 ^a	0.1	0.0 ^b	0.1 ^a	0.0 ^a	0.1 ^a	0.1	0.1	0.1
PCP	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	*
Ecstasy	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.3	0.2
Inhalants	0.2	0.1	0.2	0.1	0.2	0.1	0.2	0.1	0.2	0.2	0.1	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{2,3}	2.5	2.6	2.4	2.6	2.8 ^b	2.7	2.4	2.7 ^a	2.7	2.3	2.6	2.5	2.4
Pain Relievers	1.7	1.8	1.7	1.8	2.1 ^b	2.0 ^b	1.9 ^a	2.0 ^b	2.0 ^b	1.7	1.8 ^a	1.7	1.6
OxyContin [®]	--	--	0.1	0.1	0.1	0.1	0.2	0.2	0.2 ^b	0.2	0.1	0.2	0.1
Tranquilizers	0.8	0.8	0.7	0.8	0.7	0.7	0.7	0.8	0.9	0.7	0.9	0.7	0.7
Stimulants ³	0.5	0.5	0.5	0.5 ^a	0.6	0.4 ^b	0.4 ^b	0.5	0.4 ^b	0.4 ^b	0.4 ^a	0.5	0.6
Methamphetamine ³	0.3	0.3	0.3	0.3	0.3	0.2	0.1 ^a	0.2	0.1	0.2	0.2	0.2	0.2
Sedatives	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	3.5	3.5	3.2	3.5	3.8 ^b	3.6 ^a	3.3	3.6	3.5	3.0	3.4	3.3	3.3

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.16A Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	4,035 ^b	3,968 ^b	3,956 ^b	3,679 ^b	3,658 ^b	3,501 ^b	3,335 ^b	3,524 ^b	3,256 ^b	3,336 ^b	3,123 ^a	3,056	2,898
Marijuana and Hashish	2,714 ^b	2,569 ^b	2,552 ^b	2,348 ^a	2,315 ^a	2,233	2,203	2,352 ^a	2,269	2,387 ^b	2,266	2,188	2,117
Cocaine	337 ^b	319 ^b	339 ^b	283 ^b	257 ^b	258 ^b	218 ^b	220 ^b	190 ^b	175 ^b	129	121	105
Crack	61 ^b	89 ^b	60 ^b	48 ^b	51 ^b	43 ^a	32	28	34	36	15	15	17
Heroin	34	34	43 ^a	32	25	28	37	25	23	33	27	18	18
Hallucinogens	740 ^b	618 ^b	616 ^b	518 ^b	512 ^b	480 ^b	501 ^b	537 ^b	516 ^b	496 ^b	405	412 ^a	335
LSD	364 ^b	206	174	177	125	111 ^a	149	140	120	137	128	146	161
PCP	106 ^b	100 ^b	97 ^b	97 ^b	53 ^a	53 ^b	57 ^b	44 ^a	27	40	45 ^a	23	22
Ecstasy	376 ^b	257 ^b	243 ^b	181	202 ^a	215 ^a	224 ^b	277 ^b	277 ^b	298 ^b	214 ^a	201 ^a	150
Inhalants	1,426 ^b	1,390 ^b	1,429 ^b	1,360 ^b	1,302 ^b	1,225 ^b	1,081 ^b	1,109 ^b	985 ^b	909 ^b	749 ^b	683	595
Nonmedical Use of Psychotherapeutics ^{2,3}	1,643 ^b	1,639 ^b	1,616 ^b	1,444 ^b	1,561 ^b	1,441 ^b	1,263 ^b	1,352 ^b	1,198 ^a	1,215 ^a	1,209 ^a	1,105	1,062
Pain Relievers	1,362 ^b	1,391 ^b	1,378 ^b	1,211 ^b	1,315 ^b	1,200 ^b	1,068 ^b	1,178 ^b	1,053 ^b	1,027 ^b	1,029 ^b	902	856
OxyContin [®]	112	146	162	136	160	173	159	189 ^a	184	165	157	131	134
Tranquilizers	385 ^a	398 ^b	353	342	316	323	317	303	325	315	309	277	288
Stimulants ³	469 ^b	472 ^b	387 ^b	364 ^b	399 ^b	302 ^a	248	234	230	232	203	229	231
Methamphetamine ³	150 ^b	160 ^b	140 ^b	133 ^a	126 ^a	81	65	62	76	81	59	58	76
Sedatives	84	126	128 ^a	110	97	89	90	103	77	66	73	70	86
ILLCIT DRUGS OTHER THAN MARIJUANA¹	2,726 ^b	2,711 ^b	2,721 ^b	2,488 ^b	2,527 ^b	2,378 ^b	2,179 ^b	2,252 ^b	2,030 ^b	2,008 ^b	1,811 ^a	1,708	1,610

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.16B Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	31.9 ^b	31.1 ^b	30.7 ^b	28.4 ^b	28.2 ^b	27.2 ^b	26.2 ^b	28.0 ^b	26.1 ^b	26.1 ^b	24.5 ^a	24.1	22.8
Marijuana and Hashish	21.5 ^b	20.1 ^b	19.8 ^b	18.1 ^a	17.9	17.3	17.3	18.7 ^b	18.2 ^a	18.7 ^b	17.8	17.2	16.7
Cocaine	2.7 ^b	2.5 ^b	2.6 ^b	2.2 ^b	2.0 ^b	2.0 ^b	1.7 ^b	1.8 ^b	1.5 ^b	1.4 ^b	1.0	1.0	0.8
Crack	0.5 ^b	0.7 ^b	0.5 ^b	0.4 ^b	0.4 ^b	0.3 ^a	0.2	0.2	0.3	0.3	0.1	0.1	0.1
Heroin	0.3	0.3	0.3 ^a	0.2	0.2	0.2	0.3	0.2	0.2	0.3	0.2	0.1	0.1
Hallucinogens	5.8 ^b	4.8 ^b	4.8 ^b	4.0 ^b	4.0 ^b	3.7 ^b	3.9 ^b	4.3 ^b	4.1 ^b	3.9 ^b	3.2	3.2 ^a	2.6
LSD	2.9 ^b	1.6	1.4	1.4	1.0	0.9 ^a	1.2	1.1	1.0	1.1	1.0	1.1	1.3
PCP	0.8 ^b	0.8 ^b	0.8 ^b	0.7 ^b	0.4 ^a	0.4 ^b	0.5 ^b	0.3 ^a	0.2	0.3	0.4 ^a	0.2	0.2
Ecstasy	3.0 ^b	2.0 ^b	1.9 ^b	1.4	1.6	1.7 ^a	1.8 ^b	2.2 ^b	2.2 ^b	2.3 ^b	1.7 ^a	1.6	1.2
Inhalants	11.3 ^b	10.9 ^b	11.1 ^b	10.5 ^b	10.0 ^b	9.5 ^b	8.5 ^b	8.8 ^b	7.9 ^b	7.1 ^b	5.9 ^b	5.4	4.7
Nonmedical Use of Psychotherapeutics ^{2,3}	13.0 ^b	12.8 ^b	12.6 ^b	11.1 ^b	12.0 ^b	11.2 ^b	9.9 ^b	10.8 ^b	9.6 ^a	9.5 ^a	9.5 ^a	8.7	8.4
Pain Relievers	10.8 ^b	10.9 ^b	10.7 ^b	9.4 ^b	10.1 ^b	9.3 ^b	8.4 ^b	9.4 ^b	8.5 ^b	8.0 ^b	8.1 ^b	7.1	6.7
OxyContin [®]	0.9	1.1	1.3	1.0	1.2	1.3	1.3	1.5 ^a	1.5 ^a	1.3	1.2	1.0	1.1
Tranquilizers	3.0 ^a	3.1 ^b	2.7	2.6	2.4	2.5	2.5	2.4	2.6	2.5	2.4	2.2	2.3
Stimulants ³	3.7 ^b	3.7 ^b	3.0 ^b	2.8 ^b	3.1 ^b	2.3 ^a	2.0	1.9	1.8	1.8	1.6	1.8	1.8
Methamphetamine ³	1.2 ^b	1.3 ^b	1.1 ^b	1.0 ^a	1.0 ^a	0.6	0.5	0.5	0.6	0.6	0.5	0.5	0.6
Sedatives	0.7	1.0	1.0	0.8	0.7	0.7	0.7	0.8	0.6	0.5	0.6	0.6	0.7
ILLCIT DRUGS OTHER THAN MARIJUANA¹	21.6 ^b	21.2 ^b	21.1 ^b	19.2 ^b	19.5 ^b	18.4 ^b	17.1 ^b	17.9 ^b	16.3 ^b	15.7 ^b	14.2 ^a	13.4	12.7

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.17A Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	3,607 ^b	3,663 ^b	3,610 ^b	3,342 ^b	3,356 ^b	3,137 ^a	3,230 ^b	3,104 ^a	3,022	3,037	2,921	2,738	2,893
Marijuana and Hashish	2,391 ^b	2,318 ^b	2,236 ^b	2,055	2,084	1,863	1,935	1,852	1,888	1,991	1,966	1,905	1,975
Cocaine	333 ^b	330 ^b	272 ^b	300 ^b	313 ^b	287 ^b	254 ^b	185 ^b	173 ^a	162	136	94	117
Crack	102 ^b	68 ^b	66 ^b	60 ^b	78 ^b	64 ^b	48 ^a	30	26	31	19	7	17
Heroin	60 ^b	45 ^b	35	28	35	27	30	32	23	37	32	21	17
Hallucinogens	680 ^b	632 ^b	543 ^b	480 ^b	477 ^b	418 ^b	470 ^b	479 ^b	485 ^b	435 ^b	418 ^b	291	274
LSD	306 ^b	191 ^a	139	111	94	91	122	109	111	92	121	75 ^a	130
PCP	127 ^b	104 ^b	77 ^a	85 ^b	75 ^a	52	45	55	49	37	51	23	39
Ecstasy	444 ^b	346 ^b	280 ^b	218 ^a	274 ^b	231 ^b	301 ^b	295 ^b	340 ^b	307 ^b	274 ^b	168	146
Inhalants	1,179 ^b	1,281 ^b	1,333 ^b	1,298 ^b	1,257 ^b	1,203 ^b	1,245 ^b	1,174 ^b	1,030 ^b	968 ^b	864 ^b	644	712
Nonmedical Use of Psychotherapeutics ^{2,3}	1,759 ^b	1,733 ^b	1,787 ^b	1,575 ^b	1,613 ^b	1,513 ^b	1,521 ^b	1,458 ^b	1,392 ^a	1,307	1,273	1,083 ^a	1,227
Pain Relievers	1,412 ^b	1,407 ^b	1,500 ^b	1,305 ^b	1,335 ^b	1,257 ^b	1,286 ^b	1,215 ^b	1,191 ^b	1,123 ^b	1,040	904	948
OxyContin [®]	112	108	143	140	184 ^b	187 ^b	185 ^b	191 ^b	185 ^b	204 ^b	165	143	118
Tranquilizers	450	476 ^a	464 ^a	424	426	425	403	392	398	385	394	295 ^a	377
Stimulants ³	643 ^b	574 ^b	519 ^b	506 ^b	531 ^b	428 ^b	425 ^b	349	301	276	292	208 ^a	290
Methamphetamine ³	266 ^b	222 ^b	206 ^b	185 ^b	216 ^b	140 ^a	130	125	81	103	86	60	95
Sedatives	154	123	123	122	126	124	115	121	101	77 ^a	102	55 ^b	120
ILLCIT DRUGS OTHER THAN MARIJUANA¹	2,559 ^b	2,607 ^b	2,634 ^b	2,456 ^b	2,424 ^b	2,271 ^b	2,334 ^b	2,250 ^b	2,136 ^b	1,999 ^b	1,881 ^a	1,571	1,711

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.17B Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	29.8 ^b	30.0 ^b	29.3 ^b	26.9 ^b	27.0 ^b	25.4 ^a	26.5 ^b	25.8 ^a	25.4 ^a	24.9	24.0	22.5	23.7
Marijuana and Hashish	19.7 ^b	19.0 ^b	18.1 ^b	16.6	16.8	15.1	15.9	15.4	15.9	16.3	16.1	15.6	16.2
Cocaine	2.8 ^b	2.7 ^b	2.2 ^b	2.4 ^b	2.5 ^b	2.3 ^b	2.1 ^b	1.5 ^b	1.5 ^a	1.3	1.1	0.8	1.0
Crack	0.8 ^b	0.6 ^b	0.5 ^b	0.5 ^b	0.6 ^b	0.5 ^b	0.4 ^a	0.3	0.2	0.3	0.2	0.1	0.1
Heroin	0.5 ^b	0.4 ^b	0.3	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.3	0.2	0.1
Hallucinogens	5.6 ^b	5.2 ^b	4.4 ^b	3.9 ^b	3.8 ^b	3.4 ^b	3.9 ^b	4.0 ^b	4.1 ^b	3.6 ^b	3.4 ^b	2.4	2.3
LSD	2.5 ^b	1.6 ^a	1.1	0.9	0.8	0.7	1.0	0.9	0.9	0.8	1.0	0.6 ^a	1.1
PCP	1.0 ^b	0.8 ^b	0.6 ^a	0.7 ^b	0.6 ^a	0.4	0.4	0.5	0.4	0.3	0.4	0.2	0.3
Ecstasy	3.7 ^b	2.8 ^b	2.3 ^b	1.8 ^a	2.2 ^b	1.9 ^b	2.5 ^b	2.5 ^b	2.9 ^b	2.5 ^b	2.2 ^b	1.4	1.2
Inhalants	9.7 ^b	10.5 ^b	10.8 ^b	10.5 ^b	10.1 ^b	9.7 ^b	10.2 ^b	9.7 ^b	8.7 ^b	7.9 ^b	7.1 ^b	5.3	5.8
Nonmedical Use of Psychotherapeutics ^{2,3}	14.5 ^b	14.2 ^b	14.5 ^b	12.7 ^b	13.0 ^b	12.2 ^b	12.5 ^b	12.1 ^b	11.7 ^b	10.7	10.4	8.9 ^a	10.1
Pain Relievers	11.7 ^b	11.5 ^b	12.2 ^b	10.5 ^b	10.7 ^b	10.2 ^b	10.6 ^b	10.1 ^b	10.0 ^b	9.2 ^b	8.5	7.4	7.8
OxyContin [®]	0.9	0.9	1.2	1.1	1.5 ^b	1.5 ^b	1.5 ^b	1.6 ^b	1.6 ^b	1.7 ^b	1.4	1.2	1.0
Tranquilizers	3.7	3.9 ^a	3.8	3.4	3.4	3.4	3.3	3.3	3.4	3.2	3.2	2.4 ^a	3.1
Stimulants ³	5.3 ^b	4.7 ^b	4.2 ^b	4.1 ^b	4.3 ^b	3.5 ^b	3.5 ^b	2.9	2.5	2.3	2.4	1.7 ^a	2.4
Methamphetamine ³	2.2 ^b	1.8 ^b	1.7 ^b	1.5 ^b	1.7 ^b	1.1 ^a	1.1	1.0	0.7	0.8	0.7	0.5	0.8
Sedatives	1.3	1.0	1.0	1.0	1.0	1.0	0.9	1.0	0.9	0.6 ^a	0.8	0.5 ^b	1.0
ILLCIT DRUGS OTHER THAN MARIJUANA¹	21.1 ^b	21.3 ^b	21.4 ^b	19.8 ^b	19.5 ^b	18.4 ^b	19.2 ^b	18.7 ^b	18.0 ^b	16.4 ^b	15.4 ^a	12.9	14.0

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.18A Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	2,842 ^b	2,776 ^b	2,653 ^b	2,557 ^b	2,531 ^b	2,507 ^b	2,433 ^b	2,556 ^b	2,422 ^b	2,438 ^b	2,268	2,263	2,112
Marijuana and Hashish	2,064 ^b	1,955 ^b	1,918 ^b	1,796	1,752	1,754	1,738	1,898 ^b	1,861 ^a	1,933 ^b	1,799	1,793	1,675
Cocaine	263 ^b	222 ^b	223 ^b	208 ^b	179 ^b	174 ^b	161 ^b	146 ^a	116	115	88	74	87
Crack	36 ^b	47 ^b	31 ^a	23	32 ^a	25	19	16	8	16	8	7	12
Heroin	20	17	32	17	14	16	27	17	13	22	15	16	14
Hallucinogens	501 ^b	386 ^b	389 ^b	352 ^b	320 ^b	342 ^b	377 ^b	407 ^b	382 ^b	369 ^b	285	295 ^a	226
LSD	183 ^b	73	94	92	51 ^b	76	119	90	73	98	82	98	111
PCP	46 ^b	44 ^b	39 ^b	44 ^b	26	32 ^b	37 ^b	26	12	19	20	16	11
Ecstasy	262 ^b	142 ^a	141 ^a	125	130	165 ^b	160 ^b	212 ^b	206 ^b	218 ^b	123	140 ^a	96
Inhalants	586 ^b	554 ^b	587 ^b	538 ^b	520 ^b	501 ^b	470 ^b	458 ^b	381 ^b	396 ^b	271	237	217
Nonmedical Use of Psychotherapeutics ^{2,3}	1,091 ^b	1,090 ^b	974 ^b	962 ^b	1,055 ^b	973 ^b	867 ^b	878 ^b	806 ^a	829 ^b	777	728	686
Pain Relievers	913 ^b	930 ^b	837 ^b	818 ^b	885 ^b	819 ^b	730 ^b	755 ^b	688 ^b	693 ^b	653 ^b	575	508
OxyContin [®]	--	--	99	101	96	107	122 ^a	110	113	102	98	73	80
Tranquilizers	249	258 ^a	226	229	196	214	226	210	200	216	184	169	193
Stimulants ³	294 ^b	253 ^b	210	194	224 ^a	174	140	138	133	134	147	141	158
Methamphetamine ³	97 ^a	82	69	78	74	39	33	29	44	44	38	32	54
Sedatives	45	67	59	51	56	42	47	70	41	32	29	33	44
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1,683 ^b	1,637 ^b	1,537 ^b	1,479 ^b	1,543 ^b	1,450 ^b	1,383 ^b	1,376 ^b	1,265 ^b	1,263 ^b	1,087 ^a	1,024	950

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.18B Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	22.5 ^b	21.7 ^b	20.6 ^b	19.7 ^b	19.5 ^b	19.4 ^b	19.1 ^b	20.3 ^b	19.4 ^b	19.1 ^b	17.8	17.8	16.7
Marijuana and Hashish	16.3 ^b	15.3 ^b	14.9 ^b	13.9	13.5	13.6	13.7	15.1 ^b	14.9 ^a	15.1 ^b	14.1	14.1	13.2
Cocaine	2.1 ^b	1.7 ^b	1.7 ^b	1.6 ^b	1.4 ^b	1.4 ^b	1.3 ^b	1.2 ^a	0.9	0.9	0.7	0.6	0.7
Crack	0.3 ^b	0.4 ^b	0.2 ^a	0.2	0.2 ^a	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Heroin	0.2	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1
Hallucinogens	4.0 ^b	3.0 ^b	3.0 ^b	2.7 ^b	2.5 ^b	2.7 ^b	3.0 ^b	3.2 ^b	3.1 ^b	2.9 ^b	2.2	2.3 ^a	1.8
LSD	1.5 ^b	0.6 ^a	0.7	0.7	0.4 ^b	0.6	0.9	0.7	0.6	0.8	0.6	0.8	0.9
PCP	0.4 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.2	0.2 ^b	0.3 ^b	0.2	0.1	0.2	0.2	0.1	0.1
Ecstasy	2.1 ^b	1.1 ^a	1.1	1.0	1.0	1.3 ^b	1.3 ^b	1.7 ^b	1.7 ^b	1.7 ^b	1.0	1.1 ^a	0.8
Inhalants	4.6 ^b	4.3 ^b	4.6 ^b	4.2 ^b	4.0 ^b	3.9 ^b	3.7 ^b	3.6 ^b	3.1 ^b	3.1 ^b	2.1	1.9	1.7
Nonmedical Use of Psychotherapeutics ^{2,3}	8.6 ^b	8.5 ^b	7.6 ^b	7.4 ^b	8.1 ^b	7.5 ^b	6.8 ^b	7.0 ^b	6.5 ^a	6.5 ^a	6.1	5.7	5.4
Pain Relievers	7.2 ^b	7.3 ^b	6.5 ^b	6.3 ^b	6.8 ^b	6.4 ^b	5.7 ^b	6.0 ^b	5.5 ^b	5.4 ^b	5.1 ^b	4.5	4.0
OxyContin [®]	--	--	0.8	0.8	0.7	0.8	1.0 ^a	0.9	0.9	0.8	0.8	0.6	0.6
Tranquilizers	2.0	2.0 ^a	1.8	1.8	1.5	1.7	1.8	1.7	1.6	1.7	1.4	1.3	1.5
Stimulants ³	2.3 ^b	2.0 ^b	1.6	1.5	1.7 ^a	1.4	1.1	1.1	1.1	1.1	1.2	1.1	1.2
Methamphetamine ³	0.8 ^a	0.6	0.5	0.6	0.6	0.3	0.3	0.2	0.4	0.3	0.3	0.2	0.4
Sedatives	0.4	0.5	0.5	0.4	0.4	0.3	0.4	0.6	0.3	0.2	0.2	0.3	0.3
ILLCIT DRUGS OTHER THAN MARIJUANA¹	13.3 ^b	12.8 ^b	11.9 ^b	11.4 ^b	11.9 ^b	11.2 ^b	10.9 ^b	10.9 ^b	10.2 ^b	9.9 ^b	8.5 ^a	8.1	7.5

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.19A Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	2,652 ^b	2,679 ^b	2,648 ^b	2,484 ^b	2,447 ^a	2,231	2,318	2,282	2,322	2,297	2,201	2,023 ^a	2,224
Marijuana and Hashish	1,841 ^b	1,790 ^a	1,738	1,587	1,603	1,404 ^a	1,520	1,464	1,553	1,620	1,565	1,554	1,587
Cocaine	245 ^b	216 ^b	188 ^b	215 ^b	232 ^b	211 ^b	145 ^b	96	124	113	92	58	81
Crack	64 ^b	48 ^b	32 ^b	31 ^b	41 ^b	40 ^b	16	13	10	12	11	4	8
Heroin	31	19	16	20	23	8	14	17	15	32	16	15	14
Hallucinogens	444 ^b	392 ^b	364 ^b	313 ^b	333 ^b	287 ^b	312 ^b	330 ^b	365 ^b	279 ^b	269 ^a	186	195
LSD	131	76	63 ^a	59 ^a	55 ^a	50 ^b	66	53 ^b	62	61 ^a	70	52 ^b	103
PCP	64 ^b	44 ^a	33	41	33	24	25	22	23	20	28	13	23
Ecstasy	285 ^b	188 ^b	169 ^b	131 ^a	177 ^b	163 ^b	186 ^b	204 ^b	258 ^b	201 ^b	164 ^b	92	85
Inhalants	492 ^b	565 ^b	581 ^b	606 ^b	589 ^b	494 ^b	516 ^b	502 ^b	504 ^b	424 ^b	376 ^a	243	296
Nonmedical Use of Psychotherapeutics ^{2,3}	1,194 ^b	1,205 ^b	1,254 ^b	1,139 ^b	1,129 ^b	1,064 ^b	1,052 ^b	1,017 ^a	1,006 ^a	920	865	713 ^b	867
Pain Relievers	957 ^b	987 ^b	1,028 ^b	922 ^b	937 ^b	866 ^b	883 ^b	873 ^b	833 ^b	791 ^a	680	582	657
OxyContin [®]	--	--	95	87	105	125 ^a	104	115	121	117	88	78	81
Tranquilizers	319 ^a	328 ^b	307 ^a	268	288	291	257	252	266	259	268	178	232
Stimulants ³	371 ^b	325 ^b	297 ^b	312 ^b	324 ^b	240	233	196	191	165	177	132 ^a	203
Methamphetamine ³	148 ^b	103 ^b	108 ^b	100 ^b	111 ^b	77	62	58	54	62	45	41	51
Sedatives	93	64	75	67	54	68	67	56	70	52	53	32 ^a	72
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1,655 ^b	1,714 ^b	1,732 ^b	1,650 ^b	1,605 ^b	1,471 ^b	1,477 ^b	1,474 ^b	1,493 ^b	1,302 ^b	1,210	955 ^b	1,126

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.19B Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	21.9 ^b	21.9 ^b	21.5 ^b	20.0 ^a	19.7 ^a	18.1	19.0	18.9	19.5	18.8	18.0	16.6 ^a	18.2
Marijuana and Hashish	15.2 ^b	14.6 ^a	14.1	12.8	12.9	11.4 ^b	12.5	12.2	13.1	13.3	12.8	12.8	13.0
Cocaine	2.0 ^b	1.8 ^b	1.5 ^b	1.7 ^b	1.9 ^b	1.7 ^b	1.2 ^b	0.8	1.0 ^a	0.9	0.8	0.5	0.7
Crack	0.5 ^b	0.4 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.1	0.1	0.1	0.1	0.1	0.0	0.1
Heroin	0.3	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.3	0.1	0.1	0.1
Hallucinogens	3.7 ^b	3.2 ^b	3.0 ^b	2.5 ^b	2.7 ^b	2.3 ^b	2.6 ^b	2.7 ^b	3.1 ^b	2.3 ^b	2.2 ^a	1.5	1.6
LSD	1.1	0.6	0.5 ^a	0.5 ^a	0.4 ^a	0.4 ^b	0.5	0.4 ^b	0.5	0.5 ^a	0.6	0.4 ^b	0.8
PCP	0.5 ^b	0.4	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2
Ecstasy	2.4 ^b	1.5 ^b	1.4 ^b	1.1 ^a	1.4 ^b	1.3 ^b	1.5 ^b	1.7 ^b	2.2 ^b	1.7 ^b	1.3 ^b	0.8	0.7
Inhalants	4.1 ^b	4.6 ^b	4.7 ^b	4.9 ^b	4.7 ^b	4.0 ^b	4.2 ^b	4.2 ^b	4.2 ^b	3.5 ^b	3.1 ^a	2.0	2.4
Nonmedical Use of Psychotherapeutics ^{2,3}	9.9 ^b	9.9 ^b	10.2 ^b	9.2 ^b	9.1 ^b	8.6 ^b	8.6 ^b	8.4 ^b	8.5 ^b	7.5	7.1	5.9 ^b	7.1
Pain Relievers	7.9 ^b	8.1 ^b	8.3 ^b	7.4 ^b	7.5 ^b	7.0 ^b	7.2 ^b	7.2 ^b	7.0 ^b	6.5 ^a	5.6	4.8	5.4
OxyContin [®]	--	--	0.8	0.7	0.8	1.0 ^a	0.8	1.0	1.0 ^a	1.0	0.7	0.6	0.7
Tranquilizers	2.6 ^a	2.7 ^b	2.5 ^a	2.2	2.3	2.4	2.1	2.1	2.2	2.1	2.2	1.5	1.9
Stimulants ³	3.1 ^b	2.7 ^b	2.4 ^b	2.5 ^b	2.6 ^b	1.9	1.9	1.6	1.6	1.3	1.5	1.1 ^a	1.7
Methamphetamine ³	1.2 ^b	0.8 ^b	0.9 ^b	0.8 ^b	0.9 ^b	0.6	0.5	0.5	0.5	0.5	0.4	0.3	0.4
Sedatives	0.8	0.5	0.6	0.5	0.4	0.6	0.5	0.5	0.6	0.4	0.4	0.3 ^a	0.6
ILLCIT DRUGS OTHER THAN MARIJUANA¹	13.7 ^b	14.0 ^b	14.0 ^b	13.3 ^b	12.9 ^b	11.9 ^b	12.1 ^b	12.2 ^b	12.6 ^b	10.7 ^b	9.9	7.8 ^b	9.2

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.20A Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	1,560 ^b	1,457 ^b	1,360 ^a	1,313	1,277	1,296	1,205	1,344	1,299	1,382 ^a	1,217	1,220	1,222
Marijuana and Hashish	1,152 ^a	1,093	1,037	967	893	973	929	1,056	1,042	1,151 ^a	951	1,004	1,000
Cocaine	77 ^b	70 ^b	75 ^b	79 ^b	49 ^b	48 ^b	62 ^b	43 ^a	37	23	13	19	16
Crack	10	22	9	14	5	4	6	5	*	2	1	2	7
Heroin	6	8	11	11	3	3	8	8	2	8	*	4	8
Hallucinogens	130 ^a	138 ^a	120	98	86	103	148 ^b	128 ^a	116	134 ^a	92	98	85
LSD	37	27	23	24	10 ^a	24	36	21	30	26	10 ^a	36	38
PCP	11	14	7	8	6	9	10	8	4	3	4	2	3
Ecstasy	58 ^a	37	45	32	38	34	58 ^a	65 ^a	50	57 ^a	31	29	30
Inhalants	159 ^b	139 ^b	141 ^b	160 ^b	156 ^b	158 ^b	136 ^b	120 ^a	115 ^a	98	86	59	70
Nonmedical Use of Psychotherapeutics ^{2,3}	460 ^b	478 ^b	413 ^b	400 ^b	403 ^b	392 ^a	310	351	282	302	308	257	295
Pain Relievers	376 ^b	385 ^b	337 ^b	325 ^b	331 ^b	325 ^b	248	302 ^b	246	244	250	189	205
OxyContin [®]	--	--	29	17	19	25	33	40 ^a	27	29	29	11	17
Tranquilizers	70	111 ^b	76	84	46	84	63	58	61	74	52	42	52
Stimulants ³	106	107	89	79	81	48 ^a	54	53	49	43 ^a	51	44 ^a	83
Methamphetamine ³	40	32	24	39	24	10	16	18	8	13	9	13	22
Sedatives	6	27	4 ^a	9	21	15	8	25	14	10	7	16	18
ILLCIT DRUGS OTHER THAN MARIJUANA¹	705 ^b	682 ^b	626 ^b	616 ^b	604 ^b	584 ^b	536 ^b	539 ^b	466	484	436	373	417

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.20B Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	12.3 ^b	11.4 ^b	10.6	10.1	9.8	10.1	9.5	10.7	10.4	10.8 ^a	9.6	9.6	9.6
Marijuana and Hashish	9.1 ^a	8.6	8.1	7.5	6.9 ^a	7.5	7.3	8.4	8.4	9.0 ^a	7.5	7.9	7.9
Cocaine	0.6 ^b	0.5 ^b	0.6 ^b	0.6 ^b	0.4 ^b	0.4 ^b	0.5 ^b	0.3 ^a	0.3	0.2	0.1	0.2	0.1
Crack	0.1	0.2	0.1	0.1	0.0	0.0	0.0	0.0	*	0.0	0.0	0.0	0.1
Heroin	0.0	0.1	0.1	0.1	0.0	0.0	0.1	0.1	0.0	0.1	*	0.0	0.1
Hallucinogens	1.0 ^a	1.1 ^a	0.9	0.8	0.7	0.8	1.2 ^b	1.0 ^a	0.9	1.0 ^a	0.7	0.8	0.7
LSD	0.3	0.2	0.2	0.2	0.1 ^a	0.2	0.3	0.2	0.2	0.2	0.1 ^a	0.3	0.3
PCP	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0
Ecstasy	0.5 ^a	0.3	0.4	0.3	0.3	0.3	0.5 ^a	0.5 ^a	0.4	0.4 ^a	0.2	0.2	0.2
Inhalants	1.3 ^b	1.1 ^b	1.1 ^b	1.2 ^b	1.2 ^b	1.2 ^b	1.1 ^b	1.0 ^a	0.9 ^a	0.8	0.7	0.5	0.6
Nonmedical Use of Psychotherapeutics ^{2,3}	3.6 ^b	3.7 ^b	3.2 ^b	3.1 ^b	3.1 ^b	3.0 ^a	2.4	2.8	2.3	2.4	2.4	2.0	2.3
Pain Relievers	3.0 ^b	3.0 ^b	2.6 ^b	2.5 ^b	2.5 ^b	2.5 ^b	1.9	2.4 ^b	2.0	1.9	2.0	1.5	1.6
OxyContin [®]	--	--	0.2	0.1	0.1	0.2	0.3	0.3 ^a	0.2	0.2	0.2	0.1	0.1
Tranquilizers	0.6	0.9 ^b	0.6	0.7	0.4	0.6	0.5	0.5	0.5	0.6	0.4	0.3	0.4
Stimulants ³	0.8	0.8	0.7	0.6	0.6	0.4 ^a	0.4	0.4	0.4	0.3 ^a	0.4	0.3 ^a	0.7
Methamphetamine ³	0.3	0.3	0.2	0.3	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2
Sedatives	0.0	0.2	0.0 ^a	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	5.6 ^b	5.3 ^b	4.9 ^b	4.8 ^b	4.7 ^b	4.5 ^b	4.2 ^b	4.3 ^b	3.7	3.8	3.4	2.9	3.3

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.21A Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	1,318 ^b	1,355 ^b	1,314 ^b	1,198	1,211	1,119	1,107	1,141	1,169	1,140	1,160	977 ^a	1,116
Marijuana and Hashish	871	878	872	763	802	721 ^a	739	757	760	816	849	758	831
Cocaine	68 ^b	83 ^b	51 ^a	63 ^b	61 ^b	56 ^b	36	27	24	44	17	24	23
Crack	16 ^a	10 ^b	6	12 ^a	7	12 ^a	1	3	3	1	*	*	1
Heroin	5	10	4	4	13	*	5	5	6	6	*	8	8
Hallucinogens	126 ^b	122 ^b	91 ^a	107 ^b	96 ^a	71	97 ^a	89 ^a	99 ^b	82	68	57	51
LSD	23	23	15	8	21	6 ^a	16	14	16	8	16	16	27
PCP	11	6	4	9	6	*	5	4	3	1	*	5	*
Ecstasy	66 ^b	54 ^b	32 ^b	41 ^b	45 ^b	31 ^a	47 ^b	50 ^b	73 ^b	52 ^b	43 ^b	19	9
Inhalants	143 ^b	181 ^b	171 ^b	153 ^b	184 ^b	136 ^b	134 ^b	124 ^a	147 ^b	124 ^a	102	62	79
Nonmedical Use of Psychotherapeutics ^{2,3}	527 ^b	519 ^b	501 ^b	439	443 ^a	438	401	424	440	386	385	292	360
Pain Relievers	418 ^b	416 ^b	414 ^b	361 ^b	361 ^b	346 ^a	319	356 ^b	362 ^b	319	308	236	262
OxyContin [®]	--	--	37	17	10	22	17	29	20	21	20	19	18
Tranquilizers	117 ^b	115 ^b	86 ^a	76	79	97 ^a	88 ^a	86 ^a	73	88 ^a	87	49	52
Stimulants ³	102	115	91	93	90	78	58	74	58	55	67	40 ^a	86
Methamphetamine ³	31	38	34	32	29	24	13	18	16	23	22	9	23
Sedatives	41	20	22	18	24	19	27	15	21	14	20	9	22
ILLCIT DRUGS OTHER THAN MARIJUANA¹	715 ^b	735 ^b	702 ^b	637 ^b	648 ^b	598 ^b	557	583 ^a	626 ^b	541	482	383 ^a	467

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.21B Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	10.9 ^b	11.1 ^b	10.6 ^b	9.7	9.8	9.1	9.1	9.5	9.8	9.3	9.5	8.0 ^a	9.1
Marijuana and Hashish	7.2	7.2	7.1	6.2	6.5	5.8 ^a	6.1	6.3	6.4	6.7	7.0	6.2	6.8
Cocaine	0.6 ^b	0.7 ^b	0.4 ^a	0.5 ^b	0.5 ^b	0.5 ^b	0.3	0.2	0.2	0.4	0.1	0.2	0.2
Crack	0.1 ^a	0.1 ^b	0.1	0.1 ^a	0.1	0.1 ^a	0.0	0.0	0.0	0.0	*	*	0.0
Heroin	0.0	0.1	0.0	0.0	0.1	*	0.0	0.0	0.1	0.1	*	0.1	0.1
Hallucinogens	1.0 ^b	1.0 ^b	0.7 ^a	0.9 ^b	0.8 ^a	0.6	0.8 ^a	0.7 ^a	0.8 ^b	0.7	0.6	0.5	0.4
LSD	0.2	0.2	0.1	0.1	0.2	0.0 ^a	0.1	0.1	0.1	0.1	0.1	0.1	0.2
PCP	0.1	0.1	0.0	0.1	0.0	*	0.0	0.0	0.0	0.0	*	0.0	*
Ecstasy	0.5 ^b	0.4 ^b	0.3 ^b	0.3 ^b	0.4 ^b	0.3 ^a	0.4 ^b	0.4 ^b	0.6 ^b	0.4 ^b	0.3 ^b	0.2	0.1
Inhalants	1.2 ^b	1.5 ^b	1.4 ^b	1.2 ^b	1.5 ^b	1.1 ^a	1.1 ^b	1.0 ^a	1.2 ^b	1.0 ^a	0.8	0.5	0.6
Nonmedical Use of Psychotherapeutics ^{2,3}	4.4 ^b	4.2 ^b	4.1 ^b	3.5	3.6	3.5	3.3	3.5	3.7 ^a	3.2	3.2	2.4	3.0
Pain Relievers	3.5 ^b	3.4 ^b	3.4 ^b	2.9 ^b	2.9 ^b	2.8 ^a	2.6	3.0 ^b	3.0 ^b	2.6	2.5	1.9	2.2
OxyContin [®]	--	--	0.3	0.1	0.1	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.1
Tranquilizers	1.0 ^b	0.9 ^b	0.7 ^a	0.6	0.6	0.8 ^a	0.7 ^a	0.7 ^a	0.6	0.7 ^a	0.7	0.4	0.4
Stimulants ³	0.8	0.9	0.7	0.8	0.7	0.6	0.5	0.6	0.5	0.5	0.5	0.3 ^a	0.7
Methamphetamine ³	0.3	0.3	0.3	0.3	0.2	0.2	0.1	0.2	0.1	0.2	0.2	0.1	0.2
Sedatives	0.3	0.2	0.2	0.1	0.2	0.2	0.2	0.1	0.2	0.1	0.2	0.1	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA¹	5.9 ^b	6.0 ^b	5.7 ^b	5.1 ^b	5.2 ^b	4.8 ^b	4.6	4.8 ^a	5.3 ^b	4.4	4.0	3.1	3.8

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.22A GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	1,635	1,326	1,198	1,443	1,698	1,389	1,446	1,454	1,363
AGE GROUP									
12 to 17	40 ^b	21	21	30	21	17	20	6	14
18 to 25	475 ^b	307 ^b	202	255 ^b	217 ^a	142	150	138	143
26 or Older	1,120	998	976	1,158	1,459	1,230	1,276	1,310	1,206
GENDER									
Male	986	864	765	965	1,211 ^a	976	964	892	873
Female	650	462	433	479	486	413	482	562	490
PAST YEAR USE									
TOTAL	130	68 ^a	100	99	170	108	143	120	139
AGE GROUP									
12 to 17	16	7	8	14	3	11	15	3	7
18 to 25	46	42	22	31	30	23	33	18	24
26 or Older	68	19 ^b	70	55	137	74	96	98	108
GENDER									
Male	105	55	81	78	123	96	101	62	97
Female	26	14	19	21	47	12	42	58	42
PAST MONTH USE									
TOTAL	28	24	28	18	70	43	66	30	60
AGE GROUP									
12 to 17	5	3	1	5	*	1	11	*	4
18 to 25	6	21	5	4	6	3	12	3	*
26 or Older	17	*	22	*	63	39	43	26	56
GENDER									
Male	24	22	22	8	40	43	32	28	43
Female	*	*	*	9	30	*	34	*	18

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.22B GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.7 ^a	0.5	0.5	0.6	0.7 ^a	0.5	0.6	0.6	0.5
AGE GROUP									
12 to 17	0.2 ^b	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1
18 to 25	1.5 ^b	0.9 ^b	0.6 ^a	0.8 ^b	0.6 ^a	0.4	0.4	0.4	0.4
26 or Older	0.6	0.5	0.5	0.6	0.7	0.6	0.6	0.6	0.6
GENDER									
Male	0.8	0.7	0.6	0.8	1.0 ^a	0.8	0.8	0.7	0.7
Female	0.5 ^a	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.4
PAST YEAR USE									
TOTAL	0.1	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.1
AGE GROUP									
12 to 17	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.0
18 to 25	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
26 or Older	0.0	0.0 ^b	0.0	0.0	0.1	0.0	0.0	0.0	0.1
GENDER									
Male	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.0	0.1
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PAST MONTH USE									
TOTAL	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
AGE GROUP									
12 to 17	0.0	0.0	0.0	0.0	*	0.0	0.0	*	0.0
18 to 25	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	*
26 or Older	0.0	*	0.0	*	0.0	0.0	0.0	0.0	0.0
GENDER									
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	*	*	0.0	0.0	*	0.0	*	0.0

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.23A Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	4,704 ^b	4,951 ^b	5,366 ^b	6,751 ^b	7,936 ^b	8,198 ^b	9,624 ^b	10,721 ^a	11,717
AGE GROUP									
12 to 17	701	673	613	631	664	621	665	614	662
18 to 25	2,778 ^b	2,868 ^b	3,111 ^b	3,701 ^b	4,038 ^b	4,079 ^b	4,392 ^a	4,610	4,823
26 or Older	1,224 ^b	1,410 ^b	1,642 ^b	2,419 ^b	3,234 ^b	3,498 ^b	4,567 ^b	5,498 ^a	6,231
GENDER									
Male	2,662 ^b	2,798 ^b	2,959 ^b	4,012 ^b	4,707 ^b	4,327 ^b	5,382 ^b	5,829 ^a	6,510
Female	2,042 ^b	2,153 ^b	2,407 ^b	2,739 ^b	3,228 ^b	3,870 ^b	4,241 ^b	4,892	5,207
PAST YEAR USE									
TOTAL	1,991 ^b	2,003 ^b	2,099 ^b	2,779 ^b	2,904 ^b	3,076 ^b	3,509	3,772	3,892
AGE GROUP									
12 to 17	434	395	382	398	414	409	391	390	428
18 to 25	1,212 ^b	1,216 ^b	1,218 ^b	1,608 ^b	1,774 ^a	1,766 ^a	1,961	1,972	2,089
26 or Older	345 ^b	391 ^b	499 ^b	773 ^b	716 ^b	902 ^b	1,157	1,410	1,375
GENDER									
Male	1,055 ^b	1,157 ^b	1,155 ^b	1,649 ^b	1,661 ^b	1,684 ^b	1,934 ^a	2,059	2,252
Female	937 ^b	846 ^b	944 ^b	1,129 ^b	1,243 ^b	1,392	1,575	1,713	1,640
PAST MONTH USE									
TOTAL	551 ^b	529 ^b	613 ^b	858 ^b	921 ^b	836 ^b	948 ^b	1,100	1,278
AGE GROUP									
12 to 17	119	134	102	118	126	122	103	112	133
18 to 25	339 ^b	344 ^b	346 ^b	466 ^b	501 ^a	490 ^a	548	546	638
26 or Older	94 ^b	51 ^b	165 ^b	274 ^b	293 ^b	224 ^b	297 ^a	442	507
GENDER									
Male	299 ^b	300 ^b	344 ^b	512 ^a	547	502 ^a	507 ^a	511 ^a	694
Female	252 ^b	229 ^b	269 ^b	346 ^b	373 ^b	334 ^b	441 ^a	589	585

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.23B Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	1.9 ^b	2.0 ^b	2.1 ^b	2.7 ^b	3.1 ^b	3.2 ^b	3.7 ^b	4.1 ^a	4.4
AGE GROUP									
12 to 17	2.8	2.7	2.5	2.6	2.7	2.5	2.7	2.5	2.7
18 to 25	8.5 ^b	8.8 ^b	9.4 ^b	11.0 ^b	11.9 ^b	11.9 ^b	12.7 ^a	13.3	13.8
26 or Older	0.7 ^b	0.7 ^b	0.9 ^b	1.2 ^b	1.7 ^b	1.8 ^b	2.3 ^b	2.7	3.0
GENDER									
Male	2.2 ^b	2.3 ^b	2.4 ^b	3.3 ^b	3.8 ^b	3.5 ^b	4.3 ^b	4.6 ^a	5.1
Female	1.6 ^b	1.7 ^b	1.9 ^b	2.1 ^b	2.5 ^b	2.9 ^b	3.2 ^b	3.6	3.8
PAST YEAR USE									
TOTAL	0.8 ^b	0.8 ^b	0.8 ^b	1.1 ^b	1.1 ^b	1.2 ^b	1.3	1.4	1.5
AGE GROUP									
12 to 17	1.7	1.6	1.5	1.6	1.7	1.6	1.6	1.6	1.7
18 to 25	3.7 ^b	3.7 ^b	3.7 ^b	4.8 ^b	5.2 ^a	5.1 ^a	5.7	5.7	6.0
26 or Older	0.2 ^b	0.2 ^b	0.3 ^b	0.4 ^b	0.4 ^b	0.5 ^b	0.6	0.7	0.7
GENDER									
Male	0.9 ^b	1.0 ^b	1.0 ^b	1.3 ^b	1.3 ^b	1.4 ^b	1.5	1.6	1.8
Female	0.7 ^b	0.7 ^b	0.7 ^b	0.9 ^b	1.0 ^b	1.0	1.2	1.3	1.2
PAST MONTH USE									
TOTAL	0.2 ^b	0.2 ^b	0.2 ^b	0.3 ^b	0.4 ^b	0.3 ^b	0.4 ^b	0.4	0.5
AGE GROUP									
12 to 17	0.5	0.5	0.4	0.5	0.5	0.5	0.4	0.4	0.5
18 to 25	1.0 ^b	1.1 ^b	1.1 ^b	1.4 ^a	1.5	1.4 ^a	1.6	1.6	1.8
26 or Older	0.0 ^b	0.0 ^b	0.1 ^b	0.1 ^a	0.2 ^a	0.1 ^b	0.1 ^a	0.2	0.2
GENDER									
Male	0.3 ^b	0.2 ^b	0.3 ^b	0.4	0.4	0.4 ^a	0.4 ^a	0.4 ^a	0.5
Female	0.2 ^b	0.2 ^b	0.2 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.3	0.4	0.4

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.24A Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	4,440 ^b	5,197 ^b	5,525 ^a	6,231	6,444	6,123	6,537	6,514	6,345
AGE GROUP									
12 to 17	315 ^b	297 ^a	268	287	231	273	273	225	228
18 to 25	1,187	1,087	1,182	1,284 ^a	1,288 ^a	1,113	1,204	1,051	1,081
26 or Older	2,939 ^b	3,813 ^b	4,075 ^b	4,660	4,925	4,738	5,060	5,238	5,035
GENDER									
Male	2,141 ^b	2,366 ^b	2,345 ^b	2,932	3,227	2,703	3,029	3,172	3,084
Female	2,299 ^b	2,831	3,180	3,299	3,217	3,421	3,508	3,342	3,262
PAST YEAR USE									
TOTAL	2,034 ^b	2,048 ^b	2,065 ^b	2,163 ^b	2,264 ^b	1,904 ^a	1,844	1,747	1,498
AGE GROUP									
12 to 17	174 ^b	152 ^b	139 ^a	142 ^a	123	150 ^b	121	102	90
18 to 25	536 ^b	476 ^b	480 ^b	530 ^b	453 ^a	382	414	297	335
26 or Older	1,324	1,420 ^a	1,446 ^a	1,492 ^a	1,688 ^b	1,371	1,309	1,348	1,073
GENDER									
Male	997 ^a	995 ^a	940	1,089 ^b	1,160 ^b	789	814	819	718
Female	1,037 ^a	1,053 ^a	1,125 ^a	1,074 ^a	1,104 ^a	1,115 ^b	1,030	928	780
PAST MONTH USE									
TOTAL	544	377	442	613 ^a	815 ^b	486	527	317	387
AGE GROUP									
12 to 17	46	30	28	35	35	35	30	29	25
18 to 25	98	108	133 ^a	132 ^a	109	92	92	53	80
26 or Older	400	238	281	446	671 ^b	359	405	235	282
GENDER									
Male	329	135	206	308	460 ^a	205	191	201	192
Female	215	241	236	305	355	281	336	116	195

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.24B Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	1.8 ^b	2.1 ^a	2.2	2.5	2.5	2.4	2.5	2.5	2.4
AGE GROUP									
12 to 17	1.2 ^a	1.2 ^a	1.1	1.2	0.9	1.1	1.1	0.9	0.9
18 to 25	3.6 ^a	3.3	3.6 ^a	3.8 ^b	3.8 ^b	3.2	3.5	3.0	3.1
26 or Older	1.6 ^b	2.0 ^b	2.1 ^a	2.4	2.5	2.4	2.5	2.6	2.5
GENDER									
Male	1.8 ^b	2.0 ^a	1.9 ^b	2.4	2.6	2.2	2.4	2.5	2.4
Female	1.8 ^b	2.2	2.5	2.5	2.5	2.6	2.6	2.5	2.4
PAST YEAR USE									
TOTAL	0.8 ^b	0.8 ^b	0.8 ^b	0.9 ^b	0.9 ^b	0.7 ^a	0.7 ^a	0.7	0.6
AGE GROUP									
12 to 17	0.7 ^b	0.6 ^b	0.6 ^a	0.6 ^a	0.5	0.6 ^b	0.5	0.4	0.4
18 to 25	1.6 ^b	1.5 ^b	1.5 ^b	1.6 ^b	1.3 ^a	1.1	1.2	0.9	1.0
26 or Older	0.7 ^a	0.7 ^a	0.8 ^a	0.8 ^b	0.9 ^b	0.7 ^a	0.7	0.7	0.5
GENDER									
Male	0.8 ^b	0.8 ^b	0.8 ^a	0.9 ^b	0.9 ^b	0.6	0.6	0.6	0.6
Female	0.8 ^b	0.8 ^b	0.9 ^b	0.8 ^b	0.8 ^b	0.8 ^b	0.8	0.7	0.6
PAST MONTH USE									
TOTAL	0.2	0.2	0.2	0.2 ^a	0.3 ^b	0.2	0.2	0.1	0.1
AGE GROUP									
12 to 17	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
18 to 25	0.3	0.3	0.4 ^a	0.4 ^a	0.3	0.3	0.3	0.2	0.2
26 or Older	0.2	0.1	0.1	0.2 ^a	0.3 ^b	0.2	0.2	0.1	0.1
GENDER									
Male	0.3	0.1	0.2	0.3 ^a	0.4 ^a	0.2	0.2	0.2	0.1
Female	0.2	0.2	0.2	0.2	0.3 ^a	0.2	0.3	0.1	0.1

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.25A Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	7,770	7,108 ^b	7,532 ^a	7,419 ^a	8,044	7,678	8,311	8,652	8,278
AGE GROUP									
12 to 17	932 ^b	855 ^b	795 ^b	750	760 ^a	704	699	594	652
18 to 25	2,134	2,040	2,183	2,325	2,324	2,270	2,373	2,380	2,198
26 or Older	4,705 ^a	4,213 ^b	4,554 ^a	4,344 ^b	4,959	4,704 ^a	5,239	5,679	5,428
GENDER									
Male	4,879	4,211 ^b	4,428 ^a	4,262 ^b	5,030	4,696	5,150	5,412	5,142
Female	2,891	2,898	3,103	3,157	3,014	2,982	3,162	3,241	3,136
PAST YEAR USE									
TOTAL	1,582 ^a	1,354	1,434	1,470	1,781 ^b	1,577 ^a	1,308	1,557	1,282
AGE GROUP									
12 to 17	486 ^b	433 ^b	404 ^b	373 ^b	374 ^b	332	313	268	277
18 to 25	511 ^a	482	462	485 ^a	489 ^a	519 ^a	487	484	384
26 or Older	586	439	568	611	918	726	508	805	621
GENDER									
Male	887	690	779	727	1,102 ^a	807	631	919	725
Female	696	664	655	743	679	771 ^a	678	638	556
PAST MONTH USE									
TOTAL	423	403	404	412	467	395	410	448	351
AGE GROUP									
12 to 17	123 ^b	125 ^b	104 ^a	108 ^b	94 ^a	77	87	69	62
18 to 25	107	130	107	124	127	94	117	113	113
26 or Older	193	147	192	181	246	224	205	266	176
GENDER									
Male	263	205	220	195	248	189	157	246	180
Female	160	197	184	217	219	206	252	202	171

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.25B Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	3.2	2.9	3.0	2.9	3.2	3.0	3.2	3.3	3.1
AGE GROUP									
12 to 17	3.7 ^b	3.4 ^b	3.2 ^b	3.0 ^a	3.1 ^a	2.8	2.8	2.4	2.6
18 to 25	6.5	6.2	6.6	6.9	6.8	6.6	6.9	6.8	6.3
26 or Older	2.5	2.2 ^a	2.4	2.2 ^a	2.5	2.4	2.6	2.8	2.6
GENDER									
Male	4.1	3.5 ^a	3.7	3.5 ^a	4.1	3.8	4.1	4.3	4.0
Female	2.3	2.3	2.4	2.4	2.3	2.2	2.4	2.4	2.3
PAST YEAR USE									
TOTAL	0.6 ^b	0.5	0.6	0.6	0.7 ^b	0.6 ^a	0.5	0.6	0.5
AGE GROUP									
12 to 17	1.9 ^b	1.7 ^b	1.6 ^b	1.5 ^b	1.5 ^b	1.3	1.3	1.1	1.1
18 to 25	1.6 ^b	1.5 ^a	1.4 ^a	1.4 ^a	1.4 ^a	1.5 ^b	1.4 ^a	1.4 ^a	1.1
26 or Older	0.3	0.2	0.3	0.3	0.5	0.4	0.3	0.4	0.3
GENDER									
Male	0.7 ^a	0.6	0.6	0.6	0.9 ^a	0.6	0.5	0.7	0.6
Female	0.5 ^a	0.5	0.5	0.6 ^a	0.5	0.6 ^b	0.5	0.5	0.4
PAST MONTH USE									
TOTAL	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1
AGE GROUP									
12 to 17	0.5 ^b	0.5 ^b	0.4 ^a	0.4 ^b	0.4 ^a	0.3	0.4	0.3	0.2
18 to 25	0.3	0.4	0.3	0.4	0.4	0.3	0.3	0.3	0.3
26 or Older	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
GENDER									
Male	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.1
Female	0.1	0.2	0.1	0.2	0.2	0.2	0.2	0.1	0.1

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.26A Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	2,283 ^b	2,191 ^b	2,148 ^b	2,309 ^b	2,411 ^b	2,379 ^b	2,669	2,720	3,058
AGE GROUP									
12 to 17	83	71	74	78	80	88	67	41 ^a	73
18 to 25	919 ^b	796 ^b	636	595	555	503	496	539	542
26 or Older	1,281 ^b	1,324 ^b	1,439 ^b	1,637 ^b	1,776 ^b	1,788 ^b	2,106	2,141	2,443
GENDER									
Male	1,468 ^b	1,436 ^b	1,354 ^b	1,547 ^b	1,747	1,552 ^b	1,911	1,705 ^a	2,102
Female	814	755 ^a	794	762	663 ^b	827	758	1,015	955
PAST YEAR USE									
TOTAL	203	212	150 ^a	202	246	336	280	274	247
AGE GROUP									
12 to 17	33	32	34	35	51	54 ^a	36	26	25
18 to 25	74	122	70	92	136	152	98	137	104
26 or Older	95	58	46	75	59	130	146	111	117
GENDER									
Male	141	143	105	115	175	224	206	181	160
Female	62	69	45 ^a	88	71	112	74	93	87
PAST MONTH USE									
TOTAL	53	39	27	86 ^a	61	82	53	58	33
AGE GROUP									
12 to 17	11	12	10	17	14	25 ^a	8	5	5
18 to 25	14	17	17	26	22	29	22	21	15
26 or Older	28	10	*	43	24	28	23	32	14
GENDER									
Male	34	27	16	42	39	38	20	49	17
Female	19	12	10	43	21	44	33	9	16

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.26B Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.9 ^b	0.9 ^b	0.9 ^b	0.9 ^b	1.0 ^a	0.9 ^b	1.0	1.0	1.2
AGE GROUP									
12 to 17	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.2 ^a	0.3
18 to 25	2.8 ^b	2.4 ^b	1.9 ^a	1.8	1.6	1.5	1.4	1.5	1.6
26 or Older	0.7 ^b	0.7 ^b	0.7 ^b	0.8 ^b	0.9 ^b	0.9 ^b	1.1	1.1	1.2
GENDER									
Male	1.2 ^b	1.2 ^b	1.1 ^b	1.3 ^b	1.4	1.2 ^b	1.5	1.3 ^a	1.6
Female	0.6	0.6	0.6	0.6	0.5 ^b	0.6	0.6	0.8	0.7
PAST YEAR USE									
TOTAL	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
AGE GROUP									
12 to 17	0.1	0.1	0.1	0.1	0.2	0.2 ^a	0.1	0.1	0.1
18 to 25	0.2	0.4	0.2	0.3	0.4	0.4	0.3	0.4	0.3
26 or Older	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1
GENDER									
Male	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1	0.1
Female	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1
PAST MONTH USE									
TOTAL	0.0	0.0	0.0	0.0 ^a	0.0	0.0	0.0	0.0	0.0
AGE GROUP									
12 to 17	0.0	0.0	0.0	0.1	0.1	0.1 ^a	0.0	0.0	0.0
18 to 25	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0
26 or Older	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0
GENDER									
Male	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

**Table 7.27A DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Numbers in Thousands, 2006-2014**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	688 ^b	678 ^b	731 ^b	885 ^b	1,212 ^b	1,082 ^b	1,475 ^b	1,742	1,936
AGE GROUP									
12 to 17	45	44	49	58	61	65	98	83	77
18 to 25	307 ^b	270 ^b	280 ^b	367 ^b	444 ^b	483 ^b	619 ^a	797	767
26 or Older	336 ^b	363 ^b	402 ^b	460 ^b	707 ^b	535 ^b	757 ^a	862	1,092
GENDER									
Male	508 ^b	521 ^b	474 ^b	644 ^b	883 ^b	825 ^b	1,104 ^a	1,264	1,437
Female	180 ^b	157 ^b	258 ^b	241 ^b	329 ^a	257 ^b	371	478	498
PAST YEAR USE									
TOTAL	104 ^b	82 ^b	150 ^b	194 ^b	234 ^b	265 ^a	325	391	390
AGE GROUP									
12 to 17	16 ^a	20 ^a	29	38	38	36	52	53	45
18 to 25	64 ^b	52 ^b	73 ^b	111 ^b	118 ^b	183	198	243	191
26 or Older	24 ^b	10 ^b	47 ^b	46 ^b	79	47 ^b	75	95	153
GENDER									
Male	83 ^b	63 ^b	111 ^b	117 ^b	161 ^b	213 ^a	255	290	329
Female	21 ^b	19 ^b	39	77	73	53	70	102	60
PAST MONTH USE									
TOTAL	20 ^b	*	46	57	81	47	75	47	82
AGE GROUP									
12 to 17	*	*	8	11	17	7	17	14	12
18 to 25	15	10 ^a	19	12 ^a	21	25	46	30	35
26 or Older	*	*	19	34	43	14	12	*	35
GENDER									
Male	13 ^b	7 ^b	40	22 ^a	50	27 ^a	47	40	73
Female	7	*	*	35	31	20	29 ^a	8	9

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine. AMT is also known as alpha-methyltryptamine. The drug 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy."

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.27B DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.3 ^b	0.3 ^b	0.3 ^b	0.4 ^b	0.5 ^b	0.4 ^b	0.6 ^b	0.7	0.7
AGE GROUP									
12 to 17	0.2	0.2	0.2	0.2	0.3	0.3	0.4	0.3	0.3
18 to 25	0.9 ^b	0.8 ^b	0.8 ^b	1.1 ^b	1.3 ^b	1.4 ^b	1.8 ^a	2.3	2.2
26 or Older	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.4 ^a	0.3 ^b	0.4 ^a	0.4	0.5
GENDER									
Male	0.4 ^b	0.4 ^b	0.4 ^b	0.5 ^b	0.7 ^b	0.7 ^b	0.9 ^a	1.0	1.1
Female	0.1 ^b	0.1 ^b	0.2 ^b	0.2 ^b	0.3	0.2 ^b	0.3	0.4	0.4
PAST YEAR USE									
TOTAL	0.0 ^b	0.0 ^b	0.1 ^b	0.1 ^b	0.1 ^a	0.1 ^a	0.1	0.1	0.1
AGE GROUP									
12 to 17	0.1 ^a	0.1 ^a	0.1	0.2	0.2	0.1	0.2	0.2	0.2
18 to 25	0.2 ^b	0.2 ^b	0.2 ^b	0.3 ^a	0.3 ^a	0.5	0.6	0.7	0.5
26 or Older	0.0 ^b	0.0 ^b	0.0 ^a	0.0 ^b	0.0	0.0 ^b	0.0	0.0	0.1
GENDER									
Male	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.2 ^a	0.2	0.2	0.3
Female	0.0 ^b	0.0 ^b	0.0	0.1	0.1	0.0	0.1	0.1	0.0
PAST MONTH USE									
TOTAL	0.0 ^b	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0
AGE GROUP									
12 to 17	*	*	0.0	0.0	0.1	0.0	0.1	0.1	0.0
18 to 25	0.0	0.0 ^a	0.1	0.0 ^a	0.1	0.1	0.1	0.1	0.1
26 or Older	*	*	0.0	0.0	0.0	0.0	0.0	*	0.0
GENDER									
Male	0.0 ^b	0.0 ^b	0.0	0.0 ^a	0.0	0.0 ^a	0.0	0.0	0.1
Female	0.0	*	*	0.0	0.0	0.0	0.0 ^a	0.0	0.0

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine. AMT is also known as alpha-methyltryptamine. The drug 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy."

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.28A Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	1,802 ^b	2,522 ^b	3,201 ^b	4,406 ^b	5,235	5,285	5,223	5,417	5,218
AGE GROUP									
12 to 17	229 ^b	298 ^b	394 ^b	418 ^b	399 ^b	375 ^b	267 ^b	196	140
18 to 25	1,188 ^b	1,599 ^b	1,962	2,698 ^b	2,999 ^b	3,015 ^b	2,788 ^b	2,518 ^b	2,172
26 or Older	386 ^b	624 ^b	846 ^b	1,289 ^b	1,837 ^b	1,895 ^b	2,168 ^b	2,704	2,907
GENDER									
Male	1,436 ^b	1,873 ^b	2,296 ^b	3,391 ^a	3,907	3,733	3,891	3,898	3,849
Female	366 ^b	649 ^b	905 ^b	1,015 ^b	1,329	1,551	1,332	1,520	1,369
PAST YEAR USE									
TOTAL	761 ^b	1,007 ^b	1,213 ^b	1,283 ^b	1,191 ^b	871 ^b	648 ^b	518	354
AGE GROUP									
12 to 17	149 ^b	192 ^b	264 ^b	244 ^b	229 ^b	208 ^b	129 ^b	81	69
18 to 25	552 ^b	656 ^b	738 ^b	784 ^b	651 ^b	522 ^b	378 ^b	226	167
26 or Older	60	158	211	255 ^a	311 ^b	141	140	212	118
GENDER									
Male	600 ^b	737 ^b	877 ^b	965 ^b	929 ^b	629 ^b	463 ^b	379	238
Female	161	270 ^b	337 ^b	318 ^b	262 ^b	242 ^b	184 ^a	140	116
PAST MONTH USE									
TOTAL	136 ^b	200 ^b	198 ^b	194 ^b	162 ^b	136 ^b	122 ^b	145	44
AGE GROUP									
12 to 17	28	42 ^b	59 ^b	50 ^b	44 ^b	37 ^b	24	15	10
18 to 25	104 ^b	134 ^b	129 ^b	95 ^b	93 ^b	85 ^b	48	36	24
26 or Older	*	24	10	49	25	13	50	94	*
GENDER									
Male	112 ^b	151 ^b	160 ^b	134 ^b	134 ^b	95 ^b	88 ^a	118	28
Female	24	49 ^a	37	60 ^a	28	41	33	27	16

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.28B Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.7 ^b	1.0 ^b	1.3 ^b	1.7 ^a	2.1	2.1	2.0	2.1	2.0
AGE GROUP									
12 to 17	0.9 ^b	1.2 ^b	1.6 ^b	1.7 ^b	1.6 ^b	1.5 ^b	1.1 ^b	0.8	0.6
18 to 25	3.6 ^b	4.9 ^b	6.0	8.0 ^b	8.8 ^b	8.8 ^b	8.1 ^b	7.2 ^b	6.2
26 or Older	0.2 ^b	0.3 ^b	0.4 ^b	0.7 ^b	0.9 ^b	1.0 ^b	1.1 ^b	1.3	1.4
GENDER									
Male	1.2 ^b	1.6 ^b	1.9 ^b	2.8	3.2	3.0	3.1	3.1	3.0
Female	0.3 ^b	0.5 ^b	0.7 ^b	0.8 ^b	1.0	1.2	1.0	1.1	1.0
PAST YEAR USE									
TOTAL	0.3 ^b	0.4 ^b	0.5 ^b	0.5 ^b	0.5 ^b	0.3 ^b	0.2 ^b	0.2	0.1
AGE GROUP									
12 to 17	0.6 ^b	0.8 ^b	1.1 ^b	1.0 ^b	0.9 ^b	0.8 ^b	0.5 ^b	0.3	0.3
18 to 25	1.7 ^b	2.0 ^b	2.2 ^b	2.3 ^b	1.9 ^b	1.5 ^b	1.1 ^b	0.6	0.5
26 or Older	0.0	0.1	0.1	0.1 ^b	0.2 ^b	0.1	0.1	0.1	0.1
GENDER									
Male	0.5 ^b	0.6 ^b	0.7 ^b	0.8 ^b	0.8 ^b	0.5 ^b	0.4 ^b	0.3	0.2
Female	0.1	0.2 ^b	0.3 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.1 ^a	0.1	0.1
PAST MONTH USE									
TOTAL	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.0 ^b	0.1	0.0
AGE GROUP									
12 to 17	0.1	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.1 ^b	0.1	0.1	0.0
18 to 25	0.3 ^b	0.4 ^b	0.4 ^b	0.3 ^b	0.3 ^b	0.2 ^b	0.1	0.1	0.1
26 or Older	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*
GENDER									
Male	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^a	0.1	0.0
Female	0.0	0.0 ^a	0.0 ^a	0.0 ^a	0.0	0.0 ^a	0.0	0.0	0.0

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.29A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Numbers in Thousands, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	71,499 ^b	70,757 ^b	70,257 ^b	71,519 ^b	72,878 ^b	71,073 ^b	70,955 ^b	69,825 ^b	69,663 ^a	68,225	69,497 ^a	66,879	66,899
Cigarettes	61,136 ^b	60,434 ^b	59,896 ^b	60,532 ^b	61,568 ^b	60,183 ^b	59,833 ^b	58,723 ^b	58,336 ^b	56,819	57,525 ^a	55,778	55,241
Smokeless Tobacco	7,787 ^a	7,725 ^a	7,154 ^b	7,682 ^b	8,226	8,059	8,706	8,615	8,879	8,243	9,036	8,813	8,662
Cigars	12,751	12,837 ^a	13,727 ^b	13,640 ^b	13,701 ^b	13,300 ^b	13,225 ^b	13,345 ^b	13,273 ^b	12,865	13,417 ^b	12,438	11,964
Pipe Tobacco	1,816 ^a	1,619 ^b	1,835 ^a	2,190	2,315	2,044	1,885	2,104	2,155	2,137	2,530	2,271	2,247
ALCOHOL	119,820 ^b	118,965 ^b	120,934 ^b	126,028 ^b	125,360 ^b	126,865 ^b	128,884 ^b	130,642 ^b	131,374 ^b	133,385 ^b	135,502 ^b	136,868 ^a	139,677
Binge Alcohol Use ²	53,787 ^b	53,770 ^b	54,725 ^b	55,090 ^b	56,680 ^b	57,858 ^b	58,368 ^a	59,690	58,535 ^a	58,341 ^a	59,685	60,064	60,941
Heavy Alcohol Use ²	15,860	16,144	16,689	16,035	16,958	17,023	17,373	17,175	16,899	15,865	17,021	16,541	16,342
MALE													
TOBACCO PRODUCTS¹	41,991 ^b	41,288	41,569 ^a	42,175 ^b	43,427 ^b	42,413 ^b	41,884 ^a	40,930	41,622 ^a	40,232	41,522	39,549	39,952
Cigarettes	32,636 ^b	32,263 ^b	32,278 ^b	32,312 ^b	33,249 ^b	32,632 ^b	31,913 ^b	30,916	31,378 ^a	30,331	31,021	30,041	29,822
Smokeless Tobacco	7,242 ^b	7,096 ^b	6,730 ^b	7,174 ^b	7,838	7,598	8,248	8,203	8,367	7,676	8,496	8,211	8,205
Cigars	10,669 ^b	10,372 ^a	11,375 ^b	11,355 ^b	11,102 ^b	10,971 ^b	10,980 ^b	10,739 ^b	10,583 ^a	10,179	10,756 ^b	9,755	9,580
Pipe Tobacco	1,487	1,400 ^a	1,579	1,877	2,026	1,794	1,488	1,787	1,739	1,728	2,038	1,878	1,774
ALCOHOL	65,210 ^b	65,927 ^b	66,317 ^b	68,497 ^b	68,049 ^b	68,129 ^b	69,908 ^b	70,371 ^b	70,765 ^b	70,736 ^b	71,187 ^b	72,605	73,559
Binge Alcohol Use ²	35,456 ^b	35,565 ^b	36,195 ^b	36,025 ^b	37,379	38,151	38,384	38,691	38,116	37,375	38,277	38,379	38,586
Heavy Alcohol Use ²	12,216	11,958	12,388	12,172	12,781	12,790	12,933 ^a	12,636	12,460	11,663	12,469	12,093	11,938
FEMALE													
TOBACCO PRODUCTS¹	29,509 ^b	29,469 ^b	28,688 ^b	29,344 ^b	29,452 ^b	28,660 ^a	29,071 ^b	28,895 ^b	28,041	27,993	27,976	27,331	26,947
Cigarettes	28,500 ^b	28,171 ^b	27,618 ^b	28,220 ^b	28,319 ^b	27,551 ^b	27,920 ^b	27,807 ^b	26,957 ^a	26,488	26,504	25,737	25,418
Smokeless Tobacco	545	628	424	508	388	461	458	412	511	568	540	603	457
Cigars	2,082	2,465	2,352	2,285	2,599	2,329	2,246	2,606	2,691	2,686	2,661	2,683	2,384
Pipe Tobacco	330	219 ^b	256 ^b	313 ^b	289 ^b	250 ^b	397	317 ^b	416	410	492	392	473
ALCOHOL	54,610 ^b	53,038 ^b	54,616 ^b	57,531 ^b	57,311 ^b	58,736 ^b	58,977 ^b	60,271 ^b	60,609 ^b	62,649 ^b	64,315	64,263	66,118
Binge Alcohol Use ²	18,331 ^b	18,205 ^b	18,530 ^b	19,065 ^b	19,301 ^b	19,707 ^b	19,983 ^b	20,998 ^a	20,419 ^b	20,966 ^a	21,409	21,685	22,355
Heavy Alcohol Use ²	3,645 ^b	4,186	4,301	3,863 ^a	4,177	4,233	4,441	4,539	4,439	4,202	4,553	4,448	4,404

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.29B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Percentages, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	30.4 ^b	29.8 ^b	29.2 ^b	29.4 ^b	29.6 ^b	28.7 ^b	28.4 ^b	27.7 ^b	27.5 ^b	26.5 ^b	26.7 ^b	25.5	25.2
Cigarettes	26.0 ^b	25.4 ^b	24.9 ^b	24.9 ^b	25.0 ^b	24.3 ^b	24.0 ^b	23.3 ^b	23.0 ^b	22.1 ^b	22.1 ^b	21.3	20.8
Smokeless Tobacco	3.3	3.3	3.0 ^a	3.2	3.3	3.3	3.5	3.4	3.5	3.2	3.5	3.4	3.3
Cigars	5.4 ^b	5.4 ^b	5.7 ^b	5.6 ^b	5.6 ^b	5.4 ^b	5.3 ^b	5.3 ^b	5.2 ^b	5.0 ^b	5.2 ^b	4.7	4.5
Pipe Tobacco	0.8	0.7 ^a	0.8	0.9	0.9	0.8	0.8	0.8	0.8	0.8	1.0	0.9	0.8
ALCOHOL	51.0 ^b	50.1 ^b	50.3 ^b	51.8	51.0 ^b	51.2 ^b	51.6 ^a	51.9	51.8	51.8	52.1	52.2	52.7
Binge Alcohol Use ²	22.9	22.6	22.8	22.7	23.0	23.3	23.4	23.7	23.1	22.6	23.0	22.9	23.0
Heavy Alcohol Use ²	6.7 ^b	6.8 ^b	6.9 ^b	6.6 ^a	6.9 ^b	6.9 ^b	7.0 ^b	6.8 ^b	6.7 ^a	6.2	6.5	6.3	6.2
MALE													
TOBACCO PRODUCTS¹	37.0 ^b	35.9 ^b	35.7 ^b	35.8 ^b	36.4 ^b	35.3 ^b	34.5 ^b	33.5 ^b	33.7 ^b	32.3	33.0 ^b	31.1	31.1
Cigarettes	28.7 ^b	28.1 ^b	27.7 ^b	27.4 ^b	27.9 ^b	27.1 ^b	26.3 ^b	25.3 ^b	25.4 ^b	24.3	24.6 ^a	23.6	23.2
Smokeless Tobacco	6.4	6.2	5.8 ^a	6.1	6.6	6.3	6.8	6.7	6.8	6.2	6.7	6.5	6.4
Cigars	9.4 ^b	9.0 ^b	9.8 ^b	9.6 ^b	9.3 ^b	9.1 ^b	9.1 ^b	8.8 ^b	8.6 ^b	8.2 ^a	8.5 ^b	7.7	7.5
Pipe Tobacco	1.3	1.2	1.4	1.6	1.7	1.5	1.2	1.5	1.4	1.4	1.6	1.5	1.4
ALCOHOL	57.4	57.3	56.9	58.1	57.0	56.6	57.7	57.5	57.3	56.8	56.5	57.1	57.3
Binge Alcohol Use ²	31.2	30.9	31.1	30.5	31.3 ^a	31.7 ^b	31.7 ^a	31.6 ^a	30.9	30.0	30.4	30.2	30.0
Heavy Alcohol Use ²	10.8 ^b	10.4 ^b	10.6 ^b	10.3 ^b	10.7 ^b	10.6 ^b	10.7 ^b	10.3 ^b	10.1 ^a	9.4	9.9	9.5	9.3
FEMALE													
TOBACCO PRODUCTS¹	24.3 ^b	24.0 ^b	23.1 ^b	23.4 ^b	23.3 ^b	22.5 ^b	22.6 ^b	22.3 ^b	21.5 ^b	21.1 ^a	20.9 ^a	20.2	19.7
Cigarettes	23.4 ^b	23.0 ^b	22.3 ^b	22.5 ^b	22.4 ^b	21.6 ^b	21.7 ^b	21.5 ^b	20.7 ^b	19.9 ^b	19.8 ^a	19.0	18.6
Smokeless Tobacco	0.4	0.5	0.3	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.4	0.4	0.3
Cigars	1.7	2.0 ^a	1.9	1.8	2.1 ^a	1.8	1.7	2.0 ^a	2.1 ^a	2.0 ^a	2.0	2.0	1.7
Pipe Tobacco	0.3	0.2 ^b	0.2 ^b	0.3 ^a	0.2 ^a	0.2 ^b	0.3	0.2 ^a	0.3	0.3	0.4	0.3	0.3
ALCOHOL	44.9 ^b	43.2 ^b	44.0 ^b	45.9 ^b	45.2 ^b	46.0 ^b	45.9 ^b	46.5 ^b	46.6 ^b	47.1	47.9	47.5	48.4
Binge Alcohol Use ²	15.1 ^b	14.8 ^b	14.9 ^b	15.2 ^b	15.2 ^b	15.4 ^a	15.5	16.2	15.7	15.8	16.0	16.0	16.4
Heavy Alcohol Use ²	3.0	3.4	3.5	3.1	3.3	3.3	3.5	3.5	3.4	3.2	3.4	3.3	3.2

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.30A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Numbers in Thousands, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	3,767 ^b	3,610 ^b	3,635 ^b	3,324 ^b	3,282 ^b	3,141 ^b	2,870 ^b	2,894 ^b	2,612 ^b	2,490 ^b	2,152 ^b	1,951 ^a	1,742
Cigarettes	3,209 ^b	3,044 ^b	2,998 ^b	2,733 ^b	2,642 ^b	2,495 ^b	2,282 ^b	2,216 ^b	2,040 ^b	1,939 ^b	1,634 ^b	1,397 ^a	1,213
Smokeless Tobacco	487	511	571	523	601 ^a	620 ^b	555	583 ^a	570	517	515	487	490
Cigars	1,117 ^b	1,115 ^b	1,201 ^b	1,064 ^b	1,049 ^b	1,073 ^b	951 ^b	990 ^b	788 ^b	858 ^b	645 ^b	576	521
Pipe Tobacco	137	160	178	158	168	181	178	231	155	182	165	160	179
ALCOHOL	4,365 ^b	4,425 ^b	4,436 ^b	4,191 ^b	4,228 ^b	4,040 ^b	3,671 ^b	3,649 ^b	3,319 ^b	3,326 ^b	3,217 ^b	2,877	2,872
Binge Alcohol Use ²	2,641 ^b	2,658 ^b	2,794 ^b	2,513 ^b	2,621 ^b	2,457 ^b	2,208 ^b	2,189 ^b	1,912 ^b	1,849 ^b	1,799 ^b	1,556	1,512
Heavy Alcohol Use ²	630 ^b	652 ^b	671 ^b	611 ^b	604 ^b	592 ^b	496 ^b	520 ^b	409 ^b	379 ^b	326	293	257
MALE													
TOBACCO PRODUCTS¹	2,019 ^b	1,988 ^b	1,972 ^b	1,843 ^b	1,815 ^b	1,825 ^b	1,614 ^b	1,733 ^b	1,529 ^b	1,475 ^b	1,270 ^b	1,158	1,036
Cigarettes	1,556 ^b	1,514 ^b	1,453 ^b	1,387 ^b	1,307 ^b	1,293 ^b	1,153 ^b	1,170 ^b	1,072 ^b	1,048 ^b	865 ^b	722	649
Smokeless Tobacco	436	479	521 ^a	476	547 ^b	569 ^b	499	523 ^a	498	461	470	438	417
Cigars	790 ^b	789 ^b	852 ^b	748 ^b	719 ^b	781 ^b	683 ^b	666 ^b	539 ^b	562 ^b	445 ^b	407	337
Pipe Tobacco	89	121	110	109	114	116	99	162 ^b	98	101	87	88	97
ALCOHOL	2,200 ^b	2,185 ^b	2,216 ^b	2,057 ^b	2,114 ^b	2,057 ^b	1,827 ^b	1,921 ^b	1,710 ^b	1,697 ^b	1,607 ^b	1,429	1,369
Binge Alcohol Use ²	1,441 ^b	1,421 ^b	1,496 ^b	1,341 ^b	1,394 ^b	1,372 ^b	1,140 ^b	1,221 ^b	1,038 ^b	1,004 ^b	941 ^a	845	813
Heavy Alcohol Use ²	395 ^b	374 ^b	407 ^b	383 ^b	369 ^b	367 ^b	296 ^b	297 ^b	257 ^b	235 ^b	177	173	147
FEMALE													
TOBACCO PRODUCTS¹	1,748 ^b	1,622 ^b	1,663 ^b	1,481 ^b	1,467 ^b	1,316 ^b	1,256 ^b	1,161 ^b	1,083 ^b	1,014 ^b	882 ^b	793	706
Cigarettes	1,653 ^b	1,530 ^b	1,545 ^b	1,345 ^b	1,336 ^b	1,202 ^b	1,130 ^b	1,046 ^b	969 ^b	890 ^b	769 ^b	675 ^a	564
Smokeless Tobacco	51	32 ^b	50	47	55	50	56	60	72	56	45	49	73
Cigars	327 ^b	326 ^b	349 ^b	316 ^b	330 ^b	292 ^b	268 ^b	324 ^b	250 ^a	296 ^b	200	169	184
Pipe Tobacco	49 ^a	40 ^b	68	49	55	65	79	68	57	82	78	72	82
ALCOHOL	2,165 ^b	2,241 ^b	2,220 ^b	2,134 ^b	2,113 ^b	1,984 ^b	1,844 ^b	1,728 ^b	1,609	1,629	1,609	1,448	1,503
Binge Alcohol Use ²	1,200 ^b	1,237 ^b	1,298 ^b	1,172 ^b	1,228 ^b	1,086 ^b	1,068 ^b	968 ^b	874 ^b	845 ^b	859 ^b	711	699
Heavy Alcohol Use ²	235 ^b	278 ^b	264 ^b	229 ^b	235 ^b	225 ^b	200 ^b	223 ^b	152	144	149	119	111

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.30B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Percentages, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	15.2 ^b	14.4 ^b	14.4 ^b	13.1 ^b	12.9 ^b	12.4 ^b	11.5 ^b	11.8 ^b	10.7 ^b	10.0 ^b	8.6 ^b	7.8 ^a	7.0
Cigarettes	13.0 ^b	12.2 ^b	11.9 ^b	10.8 ^b	10.4 ^b	9.9 ^b	9.2 ^b	9.0 ^b	8.4 ^b	7.8 ^b	6.6 ^b	5.6 ^a	4.9
Smokeless Tobacco	2.0	2.0	2.3	2.1	2.4 ^a	2.5 ^a	2.2	2.4 ^a	2.3 ^a	2.1	2.1	2.0	2.0
Cigars	4.5 ^b	4.5 ^b	4.8 ^b	4.2 ^b	4.1 ^b	4.3 ^b	3.8 ^b	4.0 ^b	3.2 ^b	3.4 ^b	2.6 ^b	2.3	2.1
Pipe Tobacco	0.6	0.6	0.7	0.6	0.7	0.7	0.7	0.9	0.6	0.7	0.7	0.6	0.7
ALCOHOL	17.6 ^b	17.7 ^b	17.6 ^b	16.5 ^b	16.7 ^b	16.0 ^b	14.7 ^b	14.8 ^b	13.6 ^b	13.3 ^b	12.9 ^b	11.6	11.5
Binge Alcohol Use ²	10.7 ^b	10.6 ^b	11.1 ^b	9.9 ^b	10.3 ^b	9.7 ^b	8.9 ^b	8.9 ^b	7.9 ^b	7.4 ^b	7.2 ^b	6.2	6.1
Heavy Alcohol Use ²	2.5 ^b	2.6 ^b	2.7 ^b	2.4 ^b	2.4 ^b	2.3 ^b	2.0 ^b	2.1 ^b	1.7 ^b	1.5 ^b	1.3	1.2	1.0
MALE													
TOBACCO PRODUCTS¹	16.0 ^b	15.6 ^b	15.3 ^b	14.2 ^b	14.0 ^b	14.2 ^b	12.7 ^b	13.8 ^b	12.3 ^b	11.6 ^b	10.0 ^b	9.1	8.2
Cigarettes	12.3 ^b	11.9 ^b	11.3 ^b	10.7 ^b	10.1 ^b	10.0 ^b	9.1 ^b	9.3 ^b	8.6 ^b	8.2 ^b	6.8 ^b	5.7	5.1
Smokeless Tobacco	3.4	3.7	4.0 ^a	3.7	4.2 ^b	4.4 ^b	3.9	4.2 ^a	4.0 ^a	3.6	3.7	3.4	3.3
Cigars	6.2 ^b	6.2 ^b	6.6 ^b	5.8 ^b	5.5 ^b	6.1 ^b	5.4 ^b	5.3 ^b	4.3 ^b	4.4 ^b	3.5 ^b	3.2	2.7
Pipe Tobacco	0.7	0.9	0.9	0.8	0.9	0.9	0.8	1.3 ^b	0.8	0.8	0.7	0.7	0.8
ALCOHOL	17.4 ^b	17.1 ^b	17.2 ^b	15.9 ^b	16.3 ^b	16.0 ^b	14.4 ^b	15.3 ^b	13.7 ^b	13.3 ^b	12.6 ^b	11.2	10.8
Binge Alcohol Use ²	11.4 ^b	11.1 ^b	11.6 ^b	10.4 ^b	10.7 ^b	10.6 ^b	9.0 ^b	9.7 ^b	8.3 ^b	7.9 ^b	7.4 ^a	6.6	6.4
Heavy Alcohol Use ²	3.1 ^b	2.9 ^b	3.2 ^b	3.0 ^b	2.8 ^b	2.8 ^b	2.3 ^b	2.4 ^b	2.1 ^b	1.8 ^b	1.4	1.4	1.2
FEMALE													
TOBACCO PRODUCTS¹	14.4 ^b	13.3 ^b	13.5 ^b	11.9 ^b	11.8 ^b	10.7 ^b	10.3 ^b	9.6 ^b	9.1 ^b	8.3 ^b	7.2 ^b	6.5	5.8
Cigarettes	13.6 ^b	12.5 ^b	12.5 ^b	10.8 ^b	10.8 ^b	9.7 ^b	9.3 ^b	8.7 ^b	8.2 ^b	7.3 ^b	6.3 ^b	5.5 ^a	4.6
Smokeless Tobacco	0.4	0.3 ^b	0.4	0.4	0.4	0.4	0.5	0.5	0.6	0.5	0.4	0.4	0.6
Cigars	2.7 ^b	2.7 ^b	2.8 ^b	2.5 ^b	2.7 ^b	2.4 ^b	2.2 ^b	2.7 ^b	2.1 ^a	2.4 ^b	1.6	1.4	1.5
Pipe Tobacco	0.4 ^a	0.3 ^b	0.5	0.4	0.4	0.5	0.6	0.6	0.5	0.7	0.6	0.6	0.7
ALCOHOL	17.9 ^b	18.3 ^b	18.0 ^b	17.2 ^b	17.0 ^b	16.1 ^b	15.1 ^b	14.4 ^b	13.5	13.3	13.2	11.9	12.3
Binge Alcohol Use ²	9.9 ^b	10.1 ^b	10.5 ^b	9.4 ^b	9.9 ^b	8.8 ^b	8.8 ^b	8.0 ^b	7.4 ^b	6.9 ^b	7.0 ^b	5.8	5.7
Heavy Alcohol Use ²	1.9 ^b	2.3 ^b	2.1 ^b	1.8 ^b	1.9 ^b	1.8 ^b	1.6 ^b	1.8 ^b	1.3	1.2	1.2	1.0	0.9

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.31A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Numbers in Thousands, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	14,049 ^b	14,201 ^b	14,359 ^b	14,382 ^b	14,396 ^b	13,700 ^b	13,638 ^b	13,986 ^b	13,934 ^b	13,532 ^b	13,167 ^b	12,882 ^a	12,244
Cigarettes	12,650 ^b	12,764 ^b	12,731 ^b	12,667 ^b	12,595 ^b	11,845 ^b	11,774 ^b	12,038 ^b	11,702 ^b	11,488 ^b	11,008 ^b	10,657 ^b	9,905
Smokeless Tobacco	1,502 ^b	1,486 ^b	1,589 ^b	1,664 ^b	1,715 ^a	1,718 ^a	1,762	2,056	2,182	1,867	1,911	2,018	1,964
Cigars	3,417	3,623	4,103 ^b	3,895 ^b	3,972 ^b	3,887 ^b	3,770 ^b	3,871 ^b	3,858 ^b	3,724 ^a	3,716 ^a	3,472	3,392
Pipe Tobacco	332 ^b	292 ^b	396 ^b	471 ^b	430 ^b	395 ^b	461 ^b	592	619	635	625	758	666
ALCOHOL	18,759 ^b	19,466 ^b	19,474 ^b	19,783 ^b	20,298 ^a	20,049 ^b	20,125 ^b	20,752	20,917	20,814	20,828	20,738	20,815
Binge Alcohol Use ²	12,699	13,188	13,271	13,597	13,838 ^a	13,709 ^a	13,573	14,045 ^b	13,796 ^a	13,653	13,662	13,187	13,184
Heavy Alcohol Use ²	4,625 ^b	4,788 ^b	4,848 ^b	4,967 ^b	5,105 ^b	4,837 ^b	4,807 ^b	4,627 ^b	4,617 ^b	4,165 ^a	4,386 ^b	3,939	3,782
MALE													
TOBACCO PRODUCTS¹	8,097 ^a	8,226 ^b	8,368 ^b	8,425 ^b	8,414 ^b	8,229 ^b	8,068 ^a	8,411 ^b	8,445 ^b	8,155 ^b	8,091 ^a	8,080 ^a	7,666
Cigarettes	6,897 ^b	7,040 ^b	7,041 ^b	7,007 ^b	6,915 ^b	6,669 ^b	6,532 ^b	6,817 ^b	6,588 ^b	6,582 ^b	6,343 ^b	6,346 ^b	5,883
Smokeless Tobacco	1,455 ^b	1,418 ^b	1,532 ^b	1,588 ^b	1,644 ^a	1,633 ^a	1,687	1,926	2,063	1,769	1,820	1,888	1,877
Cigars	2,605	2,758 ^a	3,179 ^b	2,997 ^b	3,078 ^b	3,051 ^b	2,892 ^b	2,967 ^b	2,906 ^b	2,742 ^a	2,836 ^b	2,602	2,471
Pipe Tobacco	262 ^b	228 ^b	335 ^b	372	353 ^a	314 ^b	339 ^a	466	418	449	465	555	456
ALCOHOL	10,129 ^b	10,652	10,501	10,830	10,871	10,756	10,630	11,126	11,360 ^b	10,867	10,908	10,875	10,793
Binge Alcohol Use ²	7,787	8,163 ^b	8,101 ^a	8,442 ^b	8,295 ^b	8,214 ^b	8,013 ^a	8,402 ^b	8,322 ^b	7,961	7,941	7,750	7,650
Heavy Alcohol Use ²	3,282 ^b	3,372 ^b	3,434 ^b	3,545 ^b	3,469 ^b	3,284 ^b	3,305 ^b	3,224 ^b	3,173 ^b	2,814 ^a	3,016 ^b	2,659	2,497
FEMALE													
TOBACCO PRODUCTS¹	5,952 ^b	5,975 ^b	5,990 ^b	5,957 ^b	5,982 ^b	5,471 ^b	5,570 ^b	5,575 ^b	5,489 ^b	5,377 ^b	5,076 ^b	4,802	4,578
Cigarettes	5,753 ^b	5,724 ^b	5,690 ^b	5,660 ^b	5,680 ^b	5,176 ^b	5,243 ^b	5,221 ^b	5,114 ^b	4,906 ^b	4,665 ^b	4,310 ^a	4,021
Smokeless Tobacco	47 ^a	67	56	75	71	86	75	131	119	98	91	130	87
Cigars	813	865	924	898	894	836	878	905	952	982	880	870	921
Pipe Tobacco	69 ^b	65 ^b	61 ^b	100 ^b	77 ^b	82 ^b	123 ^b	126 ^b	201	186	160	203	210
ALCOHOL	8,630 ^b	8,814 ^b	8,974 ^b	8,953 ^b	9,427 ^b	9,292 ^b	9,494 ^b	9,626 ^a	9,558 ^b	9,947	9,920	9,864	10,021
Binge Alcohol Use ²	4,911 ^b	5,025 ^b	5,170 ^a	5,155 ^a	5,543	5,496	5,561	5,643	5,474	5,692	5,721	5,437	5,534
Heavy Alcohol Use ²	1,343	1,417	1,414	1,422	1,636 ^b	1,553 ^b	1,502 ^a	1,402	1,444	1,351	1,370	1,280	1,286

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.31B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Percentages, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	45.3 ^b	44.8 ^b	44.6 ^b	44.3 ^b	44.0 ^b	41.9 ^b	41.4 ^b	41.6 ^b	40.9 ^b	39.5 ^b	38.1 ^b	37.0 ^b	35.0
Cigarettes	40.8 ^b	40.2 ^b	39.5 ^b	39.0 ^b	38.5 ^b	36.2 ^b	35.7 ^b	35.8 ^b	34.3 ^b	33.5 ^b	31.8 ^b	30.6 ^b	28.4
Smokeless Tobacco	4.8 ^b	4.7 ^b	4.9 ^a	5.1	5.2	5.2	5.3	6.1	6.4 ^a	5.4	5.5	5.8	5.6
Cigars	11.0 ^b	11.4 ^b	12.7 ^b	12.0 ^b	12.1 ^b	11.9 ^b	11.4 ^b	11.5 ^b	11.3 ^b	10.9 ^b	10.7 ^a	10.0	9.7
Pipe Tobacco	1.1 ^b	0.9 ^b	1.2 ^b	1.5 ^b	1.3 ^b	1.2 ^b	1.4 ^b	1.8	1.8	1.9	1.8	2.2	1.9
ALCOHOL	60.5	61.4 ^a	60.5	60.9	62.0 ^b	61.3 ^a	61.1 ^a	61.8 ^b	61.4 ^a	60.7	60.2	59.6	59.6
Binge Alcohol Use ²	40.9 ^b	41.6 ^b	41.2 ^b	41.9 ^b	42.3 ^b	41.9 ^b	41.2 ^b	41.8 ^b	40.5 ^b	39.8 ^b	39.5 ^a	37.9	37.7
Heavy Alcohol Use ²	14.9 ^b	15.1 ^b	15.1 ^b	15.3 ^b	15.6 ^b	14.8 ^b	14.6 ^b	13.8 ^b	13.5 ^b	12.1 ^a	12.7 ^b	11.3	10.8
MALE													
TOBACCO PRODUCTS¹	52.1 ^b	51.7 ^b	51.7 ^b	51.6 ^b	51.0 ^b	49.9 ^b	48.7 ^b	49.8 ^b	48.9 ^b	47.5 ^b	46.6 ^b	46.3 ^a	43.7
Cigarettes	44.4 ^b	44.2 ^b	43.5 ^b	42.9 ^b	41.9 ^b	40.5 ^b	39.4 ^b	40.4 ^b	38.1 ^b	38.3 ^b	36.6 ^b	36.3 ^b	33.6
Smokeless Tobacco	9.4 ^a	8.9 ^b	9.5 ^a	9.7	10.0	9.9	10.2	11.4	11.9 ^a	10.3	10.5	10.8	10.7
Cigars	16.8 ^b	17.3 ^b	19.7 ^b	18.3 ^b	18.7 ^b	18.5 ^b	17.5 ^b	17.6 ^b	16.8 ^b	16.0 ^b	16.4 ^b	14.9	14.1
Pipe Tobacco	1.7 ^b	1.4 ^b	2.1	2.3	2.1	1.9 ^a	2.0 ^a	2.8	2.4	2.6	2.7	3.2	2.6
ALCOHOL	65.2 ^b	66.9 ^b	64.9 ^b	66.3 ^b	65.9 ^b	65.3 ^b	64.2 ^b	65.9 ^b	65.7 ^b	63.3	62.9	62.3	61.6
Binge Alcohol Use ²	50.2 ^b	51.3 ^b	50.1 ^b	51.7 ^b	50.3 ^b	49.8 ^b	48.4 ^b	49.8 ^b	48.1 ^b	46.3 ^a	45.8 ^a	44.4	43.6
Heavy Alcohol Use ²	21.1 ^b	21.2 ^b	21.2 ^b	21.7 ^b	21.0 ^b	19.9 ^b	19.9 ^b	19.1 ^b	18.4 ^b	16.4 ^b	17.4 ^b	15.2	14.2
FEMALE													
TOBACCO PRODUCTS¹	38.4 ^b	37.8 ^b	37.4 ^b	36.9 ^b	36.8 ^b	33.7 ^b	34.0 ^b	33.4 ^b	32.7 ^b	31.4 ^b	29.4 ^b	27.7	26.3
Cigarettes	37.1 ^b	36.2 ^b	35.5 ^b	35.0 ^b	35.0 ^b	31.9 ^b	32.0 ^b	31.3 ^b	30.5 ^b	28.7 ^b	27.1 ^b	24.9 ^a	23.1
Smokeless Tobacco	0.3	0.4	0.4	0.5	0.4	0.5	0.5	0.8 ^a	0.7	0.6	0.5	0.8	0.5
Cigars	5.2	5.5	5.8	5.6	5.5	5.1	5.4	5.4	5.7	5.7	5.1	5.0	5.3
Pipe Tobacco	0.4 ^b	0.4 ^b	0.4 ^b	0.6 ^b	0.5 ^b	0.5 ^b	0.7 ^b	0.8 ^b	1.2	1.1	0.9	1.2	1.2
ALCOHOL	55.7	55.8	56.0	55.4 ^a	58.0	57.2	58.0	57.7	56.9	58.1	57.5	56.9	57.6
Binge Alcohol Use ²	31.7	31.8	32.3	31.9	34.1 ^a	33.8 ^a	34.0 ^a	33.8 ^a	32.6	33.2	33.2	31.4	31.8
Heavy Alcohol Use ²	8.7 ^a	9.0 ^b	8.8 ^b	8.8 ^b	10.1 ^b	9.6 ^b	9.2 ^b	8.4	8.6 ^a	7.9	7.9	7.4	7.4

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.32A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Numbers in Thousands, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	53,683	52,946	52,263	53,813	55,200 ^a	54,232	54,446	52,946	53,117	52,203	54,178	52,046	52,913
Cigarettes	45,278	44,626	44,166	45,133	46,331 ^a	45,843	45,777	44,469	44,593	43,392	44,882	43,724	44,123
Smokeless Tobacco	5,797	5,728	4,994 ^b	5,495 ^a	5,910	5,722	6,388	5,976	6,127	5,859	6,609	6,308	6,209
Cigars	8,217	8,099	8,423	8,681	8,680	8,340	8,504	8,484	8,627	8,283	9,055 ^a	8,389	8,051
Pipe Tobacco	1,347	1,166	1,261	1,561	1,717	1,467	1,246	1,280	1,381	1,320	1,740	1,353	1,402
ALCOHOL	96,696 ^b	95,073 ^b	97,023 ^b	102,054 ^b	100,835 ^b	102,777 ^b	105,089 ^b	106,242 ^b	107,138 ^b	109,245 ^b	111,457 ^b	113,253 ^a	115,990
Binge Alcohol Use ²	38,447 ^b	37,924 ^b	38,661 ^b	38,980 ^b	40,220 ^b	41,692 ^b	42,586 ^b	43,456 ^b	42,827 ^b	42,839 ^b	44,224 ^a	45,322	46,245
Heavy Alcohol Use ²	10,605 ^b	10,704 ^b	11,170 ^a	10,457 ^b	11,250 ^a	11,594	12,070	12,028	11,874	11,321	12,309	12,310	12,303
MALE													
TOBACCO PRODUCTS¹	31,874	31,074	31,229	31,907	33,198 ^b	32,359	32,202	30,787	31,647	30,602	32,161	30,310	31,250
Cigarettes	24,184	23,709	23,784	23,917	25,027 ^a	24,669	24,229	22,929	23,718	22,701	23,813	22,973	23,290
Smokeless Tobacco	5,351	5,199 ^a	4,677 ^b	5,110 ^a	5,648	5,396	6,062	5,754	5,807	5,446	6,205	5,885	5,911
Cigars	7,274	6,825	7,344	7,610 ^a	7,305	7,139	7,404	7,107	7,138	6,875	7,475	6,745	6,772
Pipe Tobacco	1,136	1,051	1,133	1,396	1,559	1,364	1,051	1,158	1,223	1,178	1,487	1,235	1,220
ALCOHOL	52,880 ^b	53,090 ^b	53,601 ^b	55,610 ^b	55,064 ^b	55,316 ^b	57,451 ^b	57,324 ^b	57,695 ^b	58,172 ^b	58,672 ^b	60,302	61,396
Binge Alcohol Use ²	26,227 ^b	25,981 ^b	26,598 ^b	26,242 ^b	27,691 ^b	28,566 ^a	29,231	29,069	28,757	28,410 ^a	29,395	29,784	30,123
Heavy Alcohol Use ²	8,539	8,212 ^a	8,548	8,245 ^a	8,943	9,139	9,332	9,114	9,030	8,614	9,275	9,261	9,294
FEMALE													
TOBACCO PRODUCTS¹	21,809	21,872	21,034	21,906	22,002	21,873	22,245	22,159	21,470	21,601	22,017	21,736	21,663
Cigarettes	21,094	20,918	20,382	21,215	21,304	21,173	21,548	21,539	20,875	20,691	21,069	20,752	20,833
Smokeless Tobacco	447	529	318	385	262	325	327	221	320	413	404	423	297
Cigars	942 ^a	1,274	1,079	1,071	1,374	1,201	1,100	1,377	1,489	1,408	1,581	1,644 ^a	1,279
Pipe Tobacco	212	115	128	165	158	103	196	123	158	142	253	118	181
ALCOHOL	43,816 ^b	41,983 ^b	43,423 ^b	46,444 ^b	45,771 ^b	47,460 ^b	47,638 ^b	48,918 ^b	49,443 ^b	51,073 ^b	52,785 ^a	52,951	54,594
Binge Alcohol Use ²	12,220 ^b	11,943 ^b	12,063 ^b	12,738 ^b	12,530 ^b	13,126 ^b	13,355 ^b	14,387 ^b	14,070 ^b	14,429 ^b	14,829 ^a	15,538	16,122
Heavy Alcohol Use ²	2,067 ^b	2,492 ^a	2,623	2,212 ^b	2,306 ^b	2,455 ^a	2,739	2,914	2,843	2,707	3,034	3,049	3,008

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.32B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Percentages, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	29.9 ^b	29.3 ^b	28.5 ^b	29.0 ^b	29.4 ^b	28.6 ^b	28.4 ^b	27.3 ^b	27.2 ^b	26.3	27.0 ^a	25.7	25.8
Cigarettes	25.2 ^b	24.7 ^b	24.1 ^b	24.3 ^b	24.7 ^b	24.1 ^b	23.8 ^b	23.0 ^b	22.8 ^b	21.9	22.4	21.6	21.5
Smokeless Tobacco	3.2	3.2	2.7	3.0	3.1	3.0	3.3	3.1	3.1	3.0	3.3	3.1	3.0
Cigars	4.6 ^b	4.5 ^a	4.6 ^b	4.7 ^b	4.6 ^b	4.4 ^a	4.4 ^a	4.4 ^a	4.4 ^a	4.2	4.5 ^b	4.1	3.9
Pipe Tobacco	0.8	0.6	0.7	0.8	0.9 ^a	0.8	0.6	0.7	0.7	0.7	0.9	0.7	0.7
ALCOHOL	53.9 ^b	52.5 ^b	53.0 ^b	55.1 ^a	53.7 ^b	54.1 ^b	54.7 ^b	54.9 ^b	54.9 ^a	55.1 ^a	55.6	55.9	56.5
Binge Alcohol Use ²	21.4 ^a	21.0 ^b	21.1 ^b	21.0 ^b	21.4 ^a	22.0	22.2	22.4	21.9	21.6 ^a	22.1	22.4	22.5
Heavy Alcohol Use ²	5.9	5.9	6.1	5.6	6.0	6.1	6.3	6.2	6.1	5.7	6.1	6.1	6.0
MALE													
TOBACCO PRODUCTS¹	37.3 ^b	36.0 ^b	35.7 ^b	36.0 ^b	36.9 ^b	35.6 ^b	35.0 ^b	33.2	33.8 ^a	32.3	33.6 ^a	31.3	31.8
Cigarettes	28.3 ^b	27.5 ^b	27.2 ^b	27.0 ^b	27.8 ^b	27.1 ^b	26.3 ^b	24.7	25.3 ^a	24.0	24.9	23.7	23.7
Smokeless Tobacco	6.3	6.0	5.3	5.8	6.3	5.9	6.6	6.2	6.2	5.8	6.5	6.1	6.0
Cigars	8.5 ^b	7.9 ^a	8.4 ^b	8.6 ^b	8.1 ^b	7.9 ^a	8.0 ^b	7.7	7.6	7.3	7.8 ^a	7.0	6.9
Pipe Tobacco	1.3	1.2	1.3	1.6	1.7 ^a	1.5	1.1	1.2	1.3	1.2	1.6	1.3	1.2
ALCOHOL	61.9	61.5	61.3	62.7	61.3	60.9	62.5	61.7	61.6	61.4	61.2	62.2	62.5
Binge Alcohol Use ²	30.7	30.1	30.4	29.6	30.8	31.4	31.8	31.3	30.7	30.0	30.7	30.7	30.7
Heavy Alcohol Use ²	10.0	9.5	9.8	9.3	9.9	10.1	10.1	9.8	9.6	9.1	9.7	9.6	9.5
FEMALE													
TOBACCO PRODUCTS¹	23.2 ^b	23.1 ^b	22.0 ^b	22.6 ^b	22.5 ^b	22.1 ^b	22.2 ^b	22.0 ^b	21.1	20.8	21.0	20.6	20.2
Cigarettes	22.5 ^b	22.1 ^b	21.3 ^b	21.9 ^b	21.7 ^b	21.4 ^b	21.5 ^b	21.4 ^b	20.6	20.0	20.1	19.6	19.5
Smokeless Tobacco	0.5	0.6 ^a	0.3	0.4	0.3	0.3	0.3	0.2	0.3	0.4	0.4	0.4	0.3
Cigars	1.0	1.3	1.1	1.1	1.4	1.2	1.1	1.4	1.5	1.4	1.5 ^a	1.6 ^a	1.2
Pipe Tobacco	0.2	0.1	0.1	0.2	0.2	0.1	0.2	0.1	0.2	0.1	0.2	0.1	0.2
ALCOHOL	46.6 ^b	44.3 ^b	45.4 ^b	48.0 ^b	46.7 ^b	47.9 ^b	47.6 ^b	48.5 ^b	48.7 ^b	49.3 ^a	50.4	50.1	51.0
Binge Alcohol Use ²	13.0 ^b	12.6 ^b	12.6 ^b	13.2 ^b	12.8 ^b	13.3 ^b	13.4 ^b	14.3	13.9 ^a	13.9 ^a	14.2	14.7	15.1
Heavy Alcohol Use ²	2.2 ^b	2.6	2.7	2.3 ^a	2.4 ^a	2.5	2.7	2.9	2.8	2.6	2.9	2.9	2.8

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.33A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Numbers in Thousands, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	67,732 ^a	67,147	66,622	68,195 ^b	69,596 ^b	67,932 ^a	68,085 ^a	66,932	67,051	65,735	67,345	64,928	65,157
Cigarettes	57,928 ^b	57,391 ^b	56,897 ^b	57,799 ^b	58,926 ^b	57,688 ^b	57,551 ^b	56,507 ^a	56,295 ^a	54,880	55,890	54,381	54,028
Smokeless Tobacco	7,299 ^a	7,214 ^a	6,583 ^b	7,158 ^b	7,625	7,440 ^a	8,150	8,032	8,309	7,726	8,521	8,326	8,173
Cigars	11,634	11,722	12,526 ^a	12,576 ^a	12,652 ^b	12,227	12,274	12,355 ^a	12,485 ^a	12,007	12,772 ^b	11,862	11,443
Pipe Tobacco	1,679	1,458 ^b	1,657 ^a	2,032	2,147	1,863	1,707	1,873	2,000	1,955	2,365	2,111	2,067
ALCOHOL	115,455 ^b	114,539 ^b	116,498 ^b	121,837 ^b	121,133 ^b	122,825 ^b	125,213 ^b	126,994 ^b	128,055 ^b	130,059 ^b	132,285 ^b	133,991 ^a	136,804
Binge Alcohol Use ²	51,146 ^b	51,112 ^b	51,931 ^b	52,577 ^b	54,058 ^b	55,401 ^b	56,159 ^b	57,501	56,623 ^b	56,491 ^b	57,886	58,508	59,429
Heavy Alcohol Use ²	15,231	15,492	16,018	15,424	16,354	16,431	16,877	16,655	16,490	15,486	16,695	16,248	16,085
MALE													
TOBACCO PRODUCTS¹	39,972	39,300	39,597	40,332	41,612 ^b	40,587 ^a	40,270	39,198	40,093	38,757	40,251	38,390	38,916
Cigarettes	31,080 ^b	30,749 ^a	30,825 ^a	30,924 ^a	31,942 ^b	31,338 ^b	30,760 ^a	29,746	30,307	29,282	30,156	29,319	29,173
Smokeless Tobacco	6,806 ^b	6,618 ^b	6,209 ^b	6,698 ^b	7,291	7,029 ^a	7,749	7,680	7,870	7,215	8,026	7,773	7,788
Cigars	9,879	9,583	10,523 ^b	10,607 ^b	10,383 ^b	10,190 ^a	10,296 ^a	10,073 ^a	10,044	9,617	10,311 ^a	9,348	9,243
Pipe Tobacco	1,398	1,279 ^a	1,469	1,768	1,912	1,678	1,389	1,624	1,641	1,627	1,952	1,790	1,676
ALCOHOL	63,009 ^b	63,742 ^b	64,101 ^b	66,440 ^b	65,935 ^b	66,073 ^b	68,081 ^b	68,450 ^b	69,055 ^b	69,039 ^b	69,580 ^b	71,176	72,189
Binge Alcohol Use ²	34,014 ^b	34,144 ^b	34,699 ^b	34,684 ^b	35,985 ^a	36,780	37,244	37,470	37,078	36,371	37,336	37,534	37,773
Heavy Alcohol Use ²	11,821	11,584	11,981	11,790	12,412	12,423	12,637	12,338	12,203	11,428	12,291	11,920	11,791
FEMALE													
TOBACCO PRODUCTS¹	27,761 ^a	27,847 ^a	27,025	27,863 ^a	27,984 ^b	27,345	27,815 ^a	27,734 ^a	26,958	26,979	27,093	26,538	26,241
Cigarettes	26,847 ^b	26,641 ^b	26,072	26,875 ^b	26,984 ^b	26,349 ^a	26,791 ^b	26,761 ^b	25,989	25,598	25,734	25,062	24,855
Smokeless Tobacco	493	596	374	460	333	411	402	352	439	511	495	553	384
Cigars	1,755 ^b	2,139	2,003	1,969	2,268	2,037	1,978	2,282	2,441	2,390	2,461	2,514	2,200
Pipe Tobacco	281	179 ^b	189 ^b	264 ^a	234 ^a	185 ^b	318	249 ^b	359	328	414	320	391
ALCOHOL	52,446 ^b	50,797 ^b	52,396 ^b	55,397 ^b	55,198 ^b	56,752 ^b	57,133 ^b	58,543 ^b	59,000 ^b	61,020 ^b	62,706 ^a	62,815	64,615
Binge Alcohol Use ²	17,132 ^b	16,968 ^b	17,232 ^b	17,893 ^b	18,073 ^b	18,621 ^b	18,915 ^b	20,030 ^b	19,544 ^b	20,121 ^b	20,550	20,975	21,656
Heavy Alcohol Use ²	3,409 ^b	3,908	4,037	3,634 ^b	3,942	4,008	4,241	4,316	4,287	4,058	4,404	4,328	4,294

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.33B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Percentages, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	32.2 ^b	31.6 ^b	30.9 ^b	31.3 ^b	31.5 ^b	30.5 ^b	30.3 ^b	29.5 ^b	29.2 ^b	28.3 ^a	28.6 ^b	27.3	27.1
Cigarettes	27.5 ^b	27.0 ^b	26.4 ^b	26.5 ^b	26.7 ^b	25.9 ^b	25.6 ^b	24.9 ^b	24.6 ^b	23.6 ^a	23.8 ^b	22.9	22.5
Smokeless Tobacco	3.5	3.4	3.1 ^a	3.3	3.5	3.3	3.6	3.5	3.6	3.3	3.6	3.5	3.4
Cigars	5.5 ^b	5.5 ^b	5.8 ^b	5.8 ^b	5.7 ^b	5.5 ^b	5.5 ^b	5.4 ^b	5.4 ^b	5.2 ^a	5.4 ^b	5.0	4.8
Pipe Tobacco	0.8	0.7 ^a	0.8	0.9	1.0	0.8	0.8	0.8	0.9	0.8	1.0	0.9	0.9
ALCOHOL	54.9 ^b	53.9 ^b	54.1 ^b	55.9	54.9 ^b	55.2 ^b	55.7 ^a	55.9	55.9 ^a	55.9	56.3	56.4	56.9
Binge Alcohol Use ²	24.3	24.0	24.1	24.1	24.5	24.9	25.0	25.3	24.7	24.3	24.6	24.6	24.7
Heavy Alcohol Use ²	7.2 ^a	7.3 ^a	7.4 ^b	7.1	7.4 ^b	7.4 ^b	7.5 ^b	7.3 ^b	7.2 ^a	6.7	7.1	6.8	6.7
MALE													
TOBACCO PRODUCTS¹	39.6 ^b	38.4 ^b	38.2 ^b	38.4 ^b	39.1 ^b	37.8 ^b	37.1 ^b	35.7 ^b	36.1 ^b	34.6	35.6 ^b	33.6	33.6
Cigarettes	30.8 ^b	30.1 ^b	29.8 ^b	29.5 ^b	30.0 ^b	29.2 ^b	28.3 ^b	27.1 ^b	27.3 ^b	26.2	26.7 ^a	25.6	25.2
Smokeless Tobacco	6.7	6.5	6.0 ^a	6.4	6.9	6.5	7.1	7.0	7.1	6.4	7.1	6.8	6.7
Cigars	9.8 ^b	9.4 ^b	10.2 ^b	10.1 ^b	9.8 ^b	9.5 ^b	9.5 ^b	9.2 ^b	9.1 ^b	8.6	9.1 ^b	8.2	8.0
Pipe Tobacco	1.4	1.3	1.4	1.7	1.8	1.6	1.3	1.5	1.5	1.5	1.7	1.6	1.4
ALCOHOL	62.4	62.4	61.9	63.3	62.0	61.5	62.7	62.4	62.2	61.7	61.5	62.2	62.4
Binge Alcohol Use ²	33.7	33.4	33.5	33.0	33.8	34.3 ^a	34.3 ^a	34.1 ^a	33.4	32.5	33.0	32.8	32.6
Heavy Alcohol Use ²	11.7 ^b	11.3 ^b	11.6 ^b	11.2 ^a	11.7 ^b	11.6 ^b	11.6 ^b	11.2 ^a	11.0	10.2	10.9	10.4	10.2
FEMALE													
TOBACCO PRODUCTS¹	25.4 ^b	25.2 ^b	24.2 ^b	24.7 ^b	24.5 ^b	23.7 ^b	23.9 ^b	23.6 ^b	22.8 ^b	22.3 ^a	22.2 ^a	21.6	21.1
Cigarettes	24.5 ^b	24.1 ^b	23.3 ^b	23.8 ^b	23.6 ^b	22.9 ^b	23.0 ^b	22.8 ^b	22.0 ^b	21.2 ^a	21.1 ^a	20.4	20.0
Smokeless Tobacco	0.5	0.5	0.3	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.3
Cigars	1.6	1.9	1.8	1.7	2.0	1.8	1.7	1.9	2.1	2.0	2.0	2.0	1.8
Pipe Tobacco	0.3	0.2 ^b	0.2 ^b	0.2	0.2 ^a	0.2 ^b	0.3	0.2 ^a	0.3	0.3	0.3	0.3	0.3
ALCOHOL	47.9 ^b	46.0 ^b	46.9 ^b	49.1 ^b	48.3 ^b	49.3 ^b	49.1 ^b	49.8 ^b	49.9 ^b	50.5	51.4	51.0	51.9
Binge Alcohol Use ²	15.7 ^b	15.4 ^b	15.4 ^b	15.8 ^b	15.8 ^b	16.2 ^b	16.3 ^a	17.0	16.5	16.7	16.8	17.0	17.4
Heavy Alcohol Use ²	3.1	3.5	3.6	3.2	3.5	3.5	3.6	3.7	3.6	3.4	3.6	3.5	3.4

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.34A Daily Cigarette Use in the Past Month among Past Month Users Aged 12 or Older and Smoked One or More Packs of Cigarettes Per Day among Daily Cigarette Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
DAILY CIGARETTE USE AMONG PAST MONTH USERS	38,738 ^b	37,989 ^b	37,343 ^b	38,131 ^b	38,331 ^b	36,901 ^b	36,788 ^b	35,834 ^b	34,718 ^a	34,488 ^a	34,936 ^b	33,247	32,471
12-17	1,021 ^b	904 ^b	828 ^b	704 ^b	701 ^b	659 ^b	510 ^b	511 ^b	459 ^b	440 ^b	360	271	292
18-25	6,549 ^b	6,726 ^b	6,574 ^b	6,349 ^b	6,149 ^b	5,829 ^b	5,623 ^b	5,458 ^b	5,358 ^b	5,207 ^b	4,967 ^b	4,591	4,263
26 or Older	31,168 ^b	30,359 ^b	29,942 ^a	31,079 ^b	31,481 ^b	30,414 ^b	30,655 ^b	29,865 ^a	28,901	28,841	29,610	28,385	27,916
SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS¹	20,520 ^b	20,284 ^b	20,160 ^b	19,573 ^b	19,394 ^b	18,752 ^b	18,066 ^b	16,433 ^b	15,619 ^b	15,093 ^b	14,650 ^b	13,721	13,058
12-17	221 ^b	199 ^b	160 ^b	142 ^b	125 ^b	123 ^b	93 ^b	91 ^b	76 ^b	65 ^a	38	32	35
18-25	2,556 ^b	2,494 ^b	2,296 ^b	2,342 ^b	2,116 ^b	1,915 ^b	1,774 ^b	1,609 ^b	1,461 ^b	1,361 ^b	1,247 ^b	1,021	959
26 or Older	17,743 ^b	17,591 ^b	17,704 ^b	17,089 ^b	17,153 ^b	16,715 ^b	16,199 ^b	14,733 ^b	14,082 ^b	13,668 ^b	13,365 ^a	12,667	12,065

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Daily Cigarette Use is defined as smoking on each of the past 30 days.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.34B Daily Cigarette Use in the Past Month among Past Month Users Aged 12 or Older and Smoked One or More Packs of Cigarettes Per Day among Daily Cigarette Users Aged 12 or Older, by Age Group: Percentages, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
DAILY CIGARETTE USE AMONG PAST MONTH USERS	63.4 ^b	62.9 ^b	62.3 ^b	63.0 ^b	62.3 ^b	61.3 ^b	61.5 ^b	61.0 ^a	59.5	60.7 ^a	60.7 ^a	59.6	58.8
12-17	31.8 ^b	29.7 ^a	27.6	25.8	26.5	26.4	22.3	23.0	22.5	22.7	22.0	19.4 ^a	24.1
18-25	51.8 ^b	52.7 ^b	51.6 ^b	50.1 ^b	48.8 ^b	49.2 ^b	47.8 ^b	45.3	45.8 ^a	45.3	45.1	43.1	43.0
26 or Older	68.8 ^b	68.0 ^b	67.8 ^b	68.9 ^b	67.9 ^b	66.3 ^b	67.0 ^b	67.2 ^b	64.8	66.5 ^b	66.0 ^a	64.9	63.3
SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS¹	53.1 ^b	53.5 ^b	54.0 ^b	51.4 ^b	50.6 ^b	50.9 ^b	49.2 ^b	45.9 ^b	45.1 ^b	43.8 ^b	42.0	41.3	40.3
12-17	21.8 ^b	22.0 ^b	19.4 ^a	20.1 ^b	17.9	18.7 ^a	18.4 ^a	17.9	16.7	14.8	10.8	11.9	11.9
18-25	39.1 ^b	37.1 ^b	34.9 ^b	36.9 ^b	34.4 ^b	32.9 ^b	31.6 ^b	29.5 ^b	27.3 ^b	26.1 ^a	25.1	22.3	22.5
26 or Older	57.1 ^b	58.0 ^b	59.2 ^b	55.1 ^b	54.5 ^b	55.1 ^b	53.0 ^b	49.4 ^b	48.8 ^b	47.4 ^b	45.2	44.7	43.3

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Daily Cigarette Use is defined as smoking on each of the past 30 days.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.35A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Gender: Numbers in Thousands, 2002-2014

Gender/Alcohol Use	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
Lifetime	20,911 ^b	20,936 ^b	20,709 ^b	20,671 ^b	20,584 ^b	20,380 ^b	19,929 ^b	20,145 ^b	19,196 ^b	18,669 ^b	17,917 ^b	17,251	16,749
Past Year	17,492 ^b	17,554 ^b	17,568 ^b	17,729 ^b	17,605 ^b	17,366 ^b	16,904 ^b	17,081 ^b	16,434 ^b	15,938 ^b	15,372 ^b	14,692	14,305
Past Month	10,713 ^b	10,876 ^b	10,838 ^b	10,819 ^b	10,831 ^b	10,750 ^b	10,083 ^b	10,392 ^b	9,934 ^b	9,654 ^b	9,294 ^a	8,651	8,661
Binge Alcohol Use ¹	7,175 ^b	7,190 ^b	7,397 ^b	7,197 ^b	7,248 ^b	7,198 ^b	6,680 ^b	6,945 ^b	6,423 ^b	6,075 ^b	5,852 ^a	5,399	5,251
Heavy Alcohol Use ¹	2,301 ^b	2,297 ^b	2,375 ^b	2,296 ^b	2,361 ^b	2,321 ^b	2,136 ^b	2,063 ^b	1,940 ^b	1,698 ^b	1,651 ^b	1,412	1,301
MALE													
Lifetime	10,719 ^b	10,537 ^b	10,624 ^b	10,500 ^b	10,644 ^b	10,471 ^b	10,180 ^b	10,576 ^b	10,048 ^b	9,484 ^b	9,114 ^b	8,791	8,334
Past Year	8,847 ^b	8,735 ^b	8,946 ^b	8,927 ^b	9,056 ^b	8,903 ^b	8,520 ^b	8,949 ^b	8,638 ^b	8,004 ^b	7,821 ^b	7,425	7,041
Past Month	5,619 ^b	5,720 ^b	5,723 ^b	5,658 ^b	5,749 ^b	5,618 ^b	5,290 ^b	5,591 ^b	5,517 ^b	5,002 ^b	4,813	4,482	4,424
Binge Alcohol Use ¹	4,140 ^b	4,162 ^b	4,271 ^b	4,175 ^b	4,193 ^b	4,172 ^b	3,765 ^b	4,046 ^b	3,856 ^b	3,416 ^b	3,229	3,094	2,999
Heavy Alcohol Use ¹	1,544 ^b	1,506 ^b	1,582 ^b	1,492 ^b	1,559 ^b	1,537 ^b	1,370 ^b	1,374 ^b	1,307 ^b	1,088 ^b	1,017 ^a	908	829
FEMALE													
Lifetime	10,192 ^b	10,399 ^b	10,085 ^b	10,171 ^b	9,940 ^b	9,909 ^b	9,750 ^b	9,570 ^b	9,148 ^b	9,185 ^b	8,803	8,459	8,415
Past Year	8,645 ^b	8,819 ^b	8,622 ^b	8,802 ^b	8,549 ^b	8,463 ^b	8,384 ^b	8,132 ^b	7,797 ^a	7,934 ^a	7,551	7,267	7,263
Past Month	5,094 ^b	5,156 ^b	5,115 ^b	5,160 ^b	5,082 ^b	5,132 ^b	4,793 ^b	4,801 ^b	4,417	4,652	4,481	4,170	4,236
Binge Alcohol Use ¹	3,035 ^b	3,027 ^b	3,127 ^b	3,022 ^b	3,055 ^b	3,026 ^b	2,915 ^b	2,900 ^b	2,566 ^a	2,659 ^b	2,624 ^b	2,305	2,252
Heavy Alcohol Use ¹	756 ^b	791 ^b	793 ^b	803 ^b	802 ^b	783 ^b	765 ^b	690 ^b	634 ^b	610	634 ^b	505	471

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.35B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Gender: Percentages, 2002-2014

Gender/Alcohol Use	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
Lifetime	56.2 ^b	55.8 ^b	54.9 ^b	53.9 ^b	53.9 ^b	53.0 ^b	52.3 ^b	52.7 ^b	50.5 ^b	48.5 ^b	46.9 ^b	45.3	44.1
Past Year	47.0 ^b	46.8 ^b	46.6 ^b	46.3 ^b	46.1 ^b	45.2 ^b	44.4 ^b	44.7 ^b	43.3 ^b	41.4 ^b	40.2 ^b	38.6	37.7
Past Month	28.8 ^b	29.0 ^b	28.7 ^b	28.2 ^b	28.4 ^b	28.0 ^b	26.5 ^b	27.2 ^b	26.2 ^b	25.1 ^b	24.3 ^a	22.7	22.8
Binge Alcohol Use ¹	19.3 ^b	19.2 ^b	19.6 ^b	18.8 ^b	19.0 ^b	18.7 ^b	17.5 ^b	18.2 ^b	16.9 ^b	15.8 ^b	15.3 ^b	14.2	13.8
Heavy Alcohol Use ¹	6.2 ^b	6.1 ^b	6.3 ^b	6.0 ^b	6.2 ^b	6.0 ^b	5.6 ^b	5.4 ^b	5.1 ^b	4.4 ^b	4.3 ^b	3.7	3.4
MALE													
Lifetime	56.5 ^b	55.0 ^b	54.9 ^b	53.7 ^b	54.1 ^b	53.1 ^b	52.1 ^b	53.8 ^b	51.3 ^b	48.5 ^b	46.7 ^b	45.0 ^a	42.9
Past Year	46.6 ^b	45.6 ^b	46.3 ^b	45.6 ^b	46.0 ^b	45.2 ^b	43.6 ^b	45.6 ^b	44.1 ^b	40.9 ^b	40.1 ^b	38.0 ^a	36.2
Past Month	29.6 ^b	29.9 ^b	29.6 ^b	28.9 ^b	29.2 ^b	28.5 ^b	27.1 ^b	28.5 ^b	28.1 ^b	25.6 ^b	24.7 ^a	23.0	22.8
Binge Alcohol Use ¹	21.8 ^b	21.7 ^b	22.1 ^b	21.3 ^b	21.3 ^b	21.2 ^b	19.3 ^b	20.6 ^b	19.7 ^b	17.5 ^b	16.5	15.8	15.4
Heavy Alcohol Use ¹	8.1 ^b	7.9 ^b	8.2 ^b	7.6 ^b	7.9 ^b	7.8 ^b	7.0 ^b	7.0 ^b	6.7 ^b	5.6 ^b	5.2 ^a	4.6	4.3
FEMALE													
Lifetime	56.0 ^b	56.6 ^b	54.8 ^b	54.2 ^b	53.8 ^b	52.9 ^b	52.5 ^b	51.4 ^b	49.8 ^b	48.5 ^b	47.1	45.6	45.4
Past Year	47.5 ^b	48.0 ^b	46.9 ^b	46.9 ^b	46.2 ^b	45.2 ^b	45.2 ^b	43.7 ^b	42.4 ^b	41.9 ^b	40.4	39.2	39.2
Past Month	28.0 ^b	28.1 ^b	27.8 ^b	27.5 ^b	27.5 ^b	27.4 ^b	25.8 ^b	25.8 ^b	24.0	24.6	24.0	22.5	22.9
Binge Alcohol Use ¹	16.7 ^b	16.5 ^b	17.0 ^b	16.1 ^b	16.5 ^b	16.2 ^b	15.7 ^b	15.6 ^b	14.0 ^b	14.0 ^b	14.0 ^b	12.4	12.1
Heavy Alcohol Use ¹	4.2 ^b	4.3 ^b	4.3 ^b	4.3 ^b	4.3 ^b	4.2 ^b	4.1 ^b	3.7 ^b	3.4 ^b	3.2	3.4 ^b	2.7	2.5

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.36A Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	166,124 ^b	168,717 ^b	177,663 ^b	179,990 ^b	180,376 ^b	181,872 ^b	182,579 ^b	177,436 ^b	179,806 ^b	182,546 ^b	185,136 ^a	184,585 ^b	187,503
Marijuana													
Smoke Once a Month	88,596 ^b	92,711 ^b	94,048 ^b	91,661 ^b	94,948 ^b	93,056 ^b	90,738 ^b	85,263 ^b	80,919 ^b	80,875 ^b	78,874 ^b	72,547 ^b	68,922
Smoke Once or Twice a Week	118,697 ^b	123,443 ^b	124,258 ^b	123,054 ^b	123,833 ^b	123,760 ^b	119,654 ^b	113,048 ^b	107,617 ^b	107,133 ^b	103,277 ^b	95,410 ^b	89,132
Cocaine													
Use Once a Month	165,536 ^b	166,317 ^b	167,920 ^b	170,419 ^b	172,009 ^b	174,409 ^b	172,419 ^b	174,166 ^b	175,021 ^b	175,835 ^a	180,470	178,108	178,477
Use Once or Twice a Week	207,016 ^b	208,765 ^b	211,346 ^b	214,452 ^b	215,075 ^b	217,547 ^b	217,984 ^b	217,776 ^b	219,569 ^b	221,840 ^b	225,252	224,860	224,747
Heroin													
Try Once or Twice	190,861 ^b	192,487 ^b	192,541 ^b	196,045 ^b	198,665 ^b	199,700 ^b	200,985 ^b	202,626 ^b	203,667 ^b	207,554 ^b	211,215 ^b	211,257 ^b	216,026
Use Once or Twice a Week	217,455 ^b	220,004 ^b	222,286 ^b	224,922 ^b	226,790 ^b	229,046 ^b	230,150 ^b	231,856 ^b	232,912 ^b	236,448 ^b	239,203 ^b	240,078 ^b	242,829
LSD													
Try Once or Twice	169,629 ^b	170,754 ^b	169,691 ^b	172,935 ^a	173,271	174,476	172,464 ^a	174,144	172,951	176,281	177,840 ^a	174,818	175,144
Use Once or Twice a Week	204,413 ^b	206,339 ^b	209,139 ^b	211,266 ^b	212,215 ^b	214,118 ^b	213,556 ^b	213,572 ^b	213,198 ^b	215,905	218,024 ^a	216,427	216,174
Alcohol													
Have Four or Five Drinks Nearly Every Day	161,928 ^b	162,769 ^b	163,535 ^b	166,577 ^b	169,277 ^b	171,609	171,141 ^a	170,912 ^a	173,467	174,163	176,417 ^a	174,033	173,739
Have Five or More Drinks Once or Twice a Week	98,785 ^b	98,548 ^b	97,537 ^b	100,148 ^b	101,846 ^b	103,961	102,342 ^b	105,312	108,077	107,573	109,323 ^b	107,350	105,792
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	130,549 ^b	134,014 ^b	132,553 ^b	133,697 ^b	134,287 ^b	135,977 ^b	134,071 ^b	135,688 ^b	138,341 ^b	141,832 ^b	142,880 ^b	146,638 ^b	152,532
Cocaine	70,675 ^b	71,461 ^b	71,128 ^b	70,672 ^b	72,161 ^b	71,396 ^b	68,996 ^b	67,181 ^b	63,475 ^a	64,021 ^b	62,263	58,931	60,741
Crack	65,373 ^b	67,179 ^b	66,302 ^b	66,430 ^b	66,938 ^b	66,824 ^b	62,722 ^b	60,699 ^b	57,920 ^b	57,296 ^b	55,482	51,867	53,766
Heroin	40,380	41,125	40,518	40,101	41,261	40,583	39,718 ^a	39,086 ^b	36,992 ^b	38,213 ^b	38,332 ^b	37,097 ^b	41,567
LSD	44,659 ^b	44,053 ^b	41,970 ^b	40,919 ^b	40,501 ^b	39,407 ^b	38,013 ^b	36,510 ^a	35,762	35,635	35,264	33,951	34,568
Approached in the Past Month by Someone Selling Drugs	18,152	18,803 ^b	18,635 ^a	18,309 ^a	19,135 ^b	18,873 ^b	17,696	18,333	18,854 ^b	18,240	18,542 ^a	17,650	17,335

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.² Respondents with unknown Perceived Availability data were excluded from the respective analyses.³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.36B Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 or Older: Percentages, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	71.1	71.4	74.3 ^b	74.4 ^b	73.8 ^b	73.8 ^b	73.5 ^b	71.0	71.4	71.3	71.6	70.9	71.2
Marijuana													
Smoke Once a Month	38.3 ^b	39.6 ^b	39.7 ^b	38.3 ^b	39.3 ^b	38.3 ^b	36.9 ^b	34.4 ^b	32.5 ^b	31.9 ^b	30.8 ^b	28.2 ^b	26.5
Smoke Once or Twice a Week	51.3 ^b	52.8 ^b	52.4 ^b	51.5 ^b	51.2 ^b	50.9 ^b	48.7 ^b	45.7 ^b	43.2 ^b	42.3 ^b	40.3 ^b	37.1 ^b	34.3
Cocaine													
Use Once a Month	71.5 ^b	71.0 ^b	70.8 ^b	71.1 ^b	71.1 ^b	71.5 ^b	70.1 ^b	70.4 ^b	70.2 ^b	69.4	70.5 ^b	69.2	68.6
Use Once or Twice a Week	89.4 ^b	89.0 ^b	89.0 ^b	89.4 ^b	88.8 ^b	89.1 ^b	88.6 ^b	87.9 ^b	88.0 ^b	87.5 ^b	87.9 ^b	87.3 ^b	86.3
Heroin													
Try Once or Twice	82.4 ^a	82.2 ^b	81.3 ^b	81.8 ^b	82.2 ^b	81.9 ^b	81.8 ^b	81.9 ^b	81.8 ^b	82.0 ^b	82.6	82.2 ^a	83.1
Use Once or Twice a Week	93.9 ^b	93.9 ^b	93.7 ^a	93.8 ^b	93.8 ^a	93.9 ^b	93.6	93.6	93.4	93.4	93.4	93.3	93.3
LSD													
Try Once or Twice	73.7 ^b	73.4 ^b	72.1 ^b	72.8 ^b	72.3 ^b	72.2 ^b	70.8 ^b	71.0 ^b	70.1 ^b	70.3 ^b	70.1 ^b	68.7	68.0
Use Once or Twice a Week	88.8 ^b	88.7 ^b	88.8 ^b	88.8 ^b	88.5 ^b	88.5 ^b	87.7 ^b	86.9 ^b	86.3 ^b	86.1 ^b	85.9 ^b	85.0 ^b	83.9
Alcohol													
Have Four or Five Drinks Nearly Every Day	69.4 ^b	68.9 ^b	68.5 ^b	69.0 ^b	69.4 ^b	69.7 ^b	69.0 ^b	68.4 ^b	69.0 ^b	68.2 ^b	68.3 ^b	66.9	66.1
Have Five or More Drinks Once or Twice a Week	42.3 ^b	41.7 ^b	40.9	41.5 ^a	41.8 ^b	42.3 ^b	41.3 ^a	42.1 ^b	43.0 ^b	42.1 ^b	42.4 ^b	41.3 ^a	40.3
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	58.0 ^b	58.8 ^b	57.5 ^b	57.3 ^b	57.1 ^b	57.4 ^b	56.0 ^b	56.2 ^b	57.1 ^b	57.5 ^b	57.5 ^b	58.6 ^b	60.2
Cocaine	31.8 ^b	31.8 ^b	31.4 ^b	30.8 ^b	31.2 ^b	30.6 ^b	29.3 ^b	28.3 ^b	26.7 ^b	26.5 ^b	25.5 ^a	24.0	24.4
Crack	29.5 ^b	30.0 ^b	29.3 ^b	29.0 ^b	29.0 ^b	28.7 ^b	26.8 ^b	25.6 ^b	24.4 ^b	23.7 ^b	22.7 ^a	21.2	21.7
Heroin	18.3 ^b	18.4 ^b	18.0 ^b	17.6 ^a	17.9 ^b	17.5	17.0	16.6	15.6 ^b	15.9 ^b	15.8 ^b	15.2 ^b	16.8
LSD	20.3 ^b	19.8 ^b	18.7 ^b	18.0 ^b	17.6 ^b	17.1 ^b	16.3 ^b	15.5 ^b	15.2 ^b	14.9 ^b	14.6	14.0	14.0
Approached in the Past Month by Someone Selling Drugs	7.7 ^b	7.9 ^b	7.8 ^b	7.5 ^b	7.8 ^b	7.6 ^b	7.1 ^b	7.3 ^b	7.4 ^b	7.1 ^b	7.1 ^b	6.7	6.5

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.37A Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	15,493 ^b	15,899 ^a	16,871 ^b	17,158 ^b	17,292 ^b	17,177 ^b	17,148 ^b	15,929 ^a	15,706 ^b	16,327	16,173	15,759 ^b	16,258
Marijuana													
Smoke Once a Month	7,918 ^b	8,614 ^b	8,699 ^b	8,487 ^b	8,664 ^b	8,549 ^b	8,206 ^b	7,329 ^b	7,078 ^b	6,758 ^b	6,471 ^b	5,899 ^a	5,572
Smoke Once or Twice a Week	12,614 ^b	13,449 ^b	13,637 ^b	13,777 ^b	13,598 ^b	13,613 ^b	12,963 ^b	11,880 ^b	11,314 ^b	10,986 ^b	10,682 ^b	9,645 ^b	9,139
Cocaine													
Use Once a Month	12,296	12,628 ^b	12,320	12,151	12,193	12,234	12,061	11,817	11,679 ^a	11,712 ^a	12,235	11,901	12,064
Use Once or Twice a Week	19,463 ^b	19,835 ^b	19,842 ^b	19,894 ^b	19,754 ^b	19,530 ^b	19,348 ^b	18,894	18,649	19,035 ^a	19,215 ^b	19,005	18,745
Heroin													
Try Once or Twice	14,211	14,400	14,077	13,994	14,163	13,984	13,946 ^a	13,565 ^b	13,559 ^b	13,729 ^b	13,826 ^b	13,984	14,282
Use Once or Twice a Week	20,073 ^b	20,244 ^b	20,150 ^b	20,276 ^b	20,128 ^b	19,942 ^b	19,752 ^b	19,386	19,035 ^a	19,357	19,336	19,185	19,304
LSD													
Try Once or Twice	12,499 ^b	12,805 ^b	12,692 ^b	12,445 ^b	12,418 ^b	12,093 ^b	11,840 ^b	11,200 ^b	10,897	11,060	11,026	10,928	10,753
Use Once or Twice a Week	18,156 ^b	18,475 ^b	18,470 ^b	18,361 ^b	18,018 ^b	17,618 ^b	17,447 ^b	16,746 ^b	16,274 ^a	16,577 ^b	16,532 ^b	16,199	15,929
Alcohol													
Have Four or Five Drinks Nearly Every Day	15,288	15,277	15,468	16,028 ^b	16,259 ^b	16,265 ^b	16,203 ^b	15,611	15,584	15,973 ^b	15,762 ^b	15,318	15,307
Have Five or More Drinks Once or Twice a Week	9,379	9,528	9,514	9,636	9,868	9,797	9,879	9,620	9,717	10,026 ^a	9,777	9,546	9,615
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	13,299 ^b	13,084 ^b	12,843 ^b	12,601 ^b	12,404 ^b	12,053 ^b	11,953 ^b	11,941 ^b	11,516	11,544	11,463	11,659	11,453
Cocaine	5,981 ^b	6,010 ^b	5,917 ^b	6,073 ^b	6,325 ^b	5,909 ^b	5,269 ^b	4,909 ^b	4,420 ^b	4,178 ^b	3,782 ^b	3,397	3,385
Crack	6,324 ^b	6,286 ^b	6,069 ^b	6,149 ^b	6,376 ^b	6,111 ^b	5,533 ^b	5,205 ^b	4,653 ^b	4,331 ^b	3,938 ^b	3,517	3,372
Heroin	3,765 ^b	3,649 ^b	3,373 ^b	3,384 ^b	3,477 ^b	3,395 ^b	3,055 ^b	2,991 ^b	2,679 ^b	2,542 ^b	2,320	2,132	2,210
LSD	4,562 ^b	4,154 ^b	4,035 ^b	3,732 ^b	3,310 ^b	3,377 ^b	3,199 ^b	3,079 ^b	2,925 ^a	2,851	2,646	2,603	2,685
Approached in the Past Month by Someone Selling Drugs	4,121 ^b	4,013 ^b	4,103 ^b	3,926 ^b	3,885 ^b	3,667 ^b	3,418 ^b	3,531 ^b	3,474 ^b	3,438 ^b	3,272 ^a	3,087	3,013

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.37B Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 to 17: Percentages, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	63.1 ^b	64.2 ^b	67.5 ^a	68.3 ^b	68.7 ^b	68.8 ^b	69.5 ^b	65.5	65.3	66.2	65.7	64.3 ^b	66.3
Marijuana													
Smoke Once a Month	32.4 ^b	34.9 ^b	35.0 ^b	34.0 ^b	34.6 ^b	34.4 ^b	33.4 ^b	30.3 ^b	29.6 ^b	27.6 ^b	26.5 ^b	24.2 ^a	22.9
Smoke Once or Twice a Week	51.5 ^b	54.4 ^b	54.7 ^b	55.0 ^b	54.2 ^b	54.6 ^b	52.8 ^b	49.0 ^b	47.2 ^b	44.8 ^b	43.6 ^b	39.5 ^b	37.4
Cocaine													
Use Once a Month	50.5	51.4 ^a	49.6	48.8	49.0	49.6	49.4	49.2	49.1	48.1 ^a	50.4	49.3	49.8
Use Once or Twice a Week	79.8 ^b	80.7 ^b	79.8 ^b	79.9 ^b	79.2 ^b	78.9 ^b	79.1 ^b	78.4 ^a	78.3 ^a	78.1	78.9 ^b	78.4 ^a	77.2
Heroin													
Try Once or Twice	58.5	58.8	57.0 ^b	56.5 ^b	57.2 ^b	56.9 ^b	57.4 ^b	56.7 ^b	57.4 ^b	56.6 ^b	57.3 ^b	58.2	59.2
Use Once or Twice a Week	82.5 ^b	82.6 ^b	81.4 ^b	81.8 ^b	81.2 ^a	81.0	81.3 ^a	80.9	80.4	79.7	80.0	79.8	79.9
LSD													
Try Once or Twice	52.6 ^b	53.4 ^b	52.6 ^b	51.7 ^b	51.6 ^b	51.0 ^b	50.2 ^b	48.1 ^b	47.8 ^a	47.0	47.2	47.1	46.2
Use Once or Twice a Week	76.2 ^b	76.9 ^b	76.4 ^b	76.1 ^b	74.7 ^b	74.1 ^b	73.8 ^b	71.7 ^b	71.3 ^b	70.4 ^b	70.6 ^b	69.7 ^a	68.4
Alcohol													
Have Four or Five Drinks Nearly Every Day	62.2	61.6	61.8	63.8 ^a	64.5 ^b	65.1 ^b	65.6 ^b	64.1 ^b	64.7 ^b	64.8 ^b	63.9 ^a	62.5	62.4
Have Five or More Drinks Once or Twice a Week	38.2	38.5	38.1	38.4	39.3	39.3	40.0	39.6	40.4	40.7 ^a	39.7	39.0	39.2
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	55.0 ^b	53.6 ^b	52.2 ^b	51.0 ^b	50.1 ^b	49.2 ^a	49.4 ^a	50.0 ^b	48.9	47.7	47.8	48.6	47.8
Cocaine	25.0 ^b	25.0 ^b	24.4 ^b	24.9 ^b	25.9 ^b	24.5 ^b	22.1 ^b	20.9 ^b	19.1 ^b	17.5 ^b	16.0 ^b	14.4	14.4
Crack	26.5 ^b	26.2 ^b	25.0 ^b	25.3 ^b	26.2 ^b	25.3 ^b	23.2 ^b	22.1 ^b	20.1 ^b	18.2 ^b	16.7 ^b	14.9	14.3
Heroin	15.8 ^b	15.3 ^b	14.0 ^b	14.0 ^b	14.4 ^b	14.2 ^b	12.9 ^b	12.8 ^b	11.6 ^b	10.7 ^b	9.9	9.1	9.4
LSD	19.4 ^b	17.6 ^b	16.9 ^b	15.7 ^b	14.0 ^b	14.4 ^b	13.7 ^b	13.4 ^b	13.0 ^b	12.2	11.5	11.3	11.6
Approached in the Past Month by Someone Selling Drugs	16.7 ^b	16.1 ^b	16.3 ^b	15.5 ^b	15.3 ^b	14.5 ^b	13.8 ^b	14.4 ^b	14.3 ^b	13.8 ^b	13.2 ^a	12.4	12.1

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.38A Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	20,152 ^b	20,743 ^b	22,378 ^b	22,595 ^b	23,016	23,036	22,588 ^b	21,650 ^b	22,245 ^b	22,501 ^b	22,958	22,859	23,253
Marijuana													
Smoke Once a Month	7,252 ^b	7,812 ^b	8,169 ^b	7,874 ^b	7,938 ^b	8,018 ^b	7,414 ^b	6,649 ^b	6,259 ^b	6,142 ^b	5,741 ^b	5,147 ^b	4,689
Smoke Once or Twice a Week	10,947 ^b	11,600 ^b	12,177 ^b	11,920 ^b	11,659 ^b	11,651 ^b	10,829 ^b	9,560 ^b	9,140 ^b	8,736 ^b	8,062 ^b	7,257 ^b	6,357
Cocaine													
Use Once a Month	19,683 ^b	19,971 ^b	19,928 ^b	20,060 ^b	20,268 ^b	20,473 ^b	20,183 ^b	20,239 ^b	20,847 ^b	21,041 ^b	21,727	21,609	21,671
Use Once or Twice a Week	26,794 ^b	27,246 ^b	27,354 ^b	27,639 ^b	27,789 ^b	27,817 ^b	28,012 ^b	28,021 ^b	28,475 ^b	28,774	29,259	28,990	29,096
Heroin													
Try Once or Twice	23,917 ^b	24,296 ^b	24,466 ^b	24,676 ^b	25,194 ^b	25,268 ^b	25,175 ^b	25,491 ^b	25,860 ^b	26,458 ^b	27,007 ^b	27,206 ^a	27,640
Use Once or Twice a Week	28,723 ^b	29,367 ^b	29,700 ^b	30,105 ^b	30,297 ^b	30,281 ^b	30,335 ^b	30,805 ^b	31,176 ^b	31,427 ^b	31,893 ^b	31,930 ^b	32,251
LSD													
Try Once or Twice	19,046	19,606	19,790 ^a	19,821 ^b	20,309 ^b	20,411 ^b	19,767 ^a	19,702 ^a	19,737 ^a	20,106 ^b	20,007 ^b	19,690	19,207
Use Once or Twice a Week	25,890 ^a	26,587	27,118 ^b	27,298 ^b	27,237 ^b	27,260 ^b	26,915 ^b	26,822 ^a	26,877 ^b	26,942 ^b	27,105 ^b	26,380	26,322
Alcohol													
Have Four or Five Drinks Nearly Every Day	19,201 ^b	19,284 ^b	19,795 ^b	20,005 ^b	20,277 ^a	20,346 ^a	20,634	20,676	21,026	21,165	21,135	20,690	20,847
Have Five or More Drinks Once or Twice a Week	10,248 ^b	10,078 ^b	10,086 ^b	10,566 ^b	10,626 ^b	10,817 ^b	10,810 ^b	11,089 ^a	11,422	11,801	11,874	11,494	11,651
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	23,718 ^b	24,409 ^b	24,122 ^b	24,524 ^b	24,715 ^b	24,494 ^b	24,431 ^b	25,221 ^a	25,425	25,353	25,674	25,751	25,703
Cocaine	12,176 ^b	12,622 ^b	12,753 ^b	13,199 ^b	13,212 ^b	12,860 ^b	12,147 ^b	11,690 ^b	11,230 ^b	10,513 ^b	10,071 ^b	9,324	9,290
Crack	9,805 ^b	9,909 ^b	9,960 ^b	10,263 ^b	10,251 ^b	10,133 ^b	9,191 ^b	8,850 ^b	8,260 ^b	7,722 ^b	7,314 ^b	6,571	6,371
Heroin	6,062 ^b	6,117 ^b	5,914 ^b	5,964 ^b	5,980 ^b	5,910 ^b	5,647 ^b	5,452 ^a	5,528 ^b	5,042	5,211	5,003	5,036
LSD	8,634 ^b	8,113 ^b	7,308 ^b	7,075 ^b	6,646 ^b	6,317 ^b	6,163 ^a	5,978	6,072	5,851	5,795	5,621	5,734
Approached in the Past Month by Someone Selling Drugs	5,894	6,223 ^b	6,063 ^a	6,134 ^b	6,020 ^a	6,096 ^a	6,077 ^a	6,211 ^b	6,490 ^b	6,182 ^b	6,535 ^b	5,656	5,644

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.38B Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 18 to 25: Percentages, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	65.2 ^b	65.7	69.8 ^b	69.8 ^b	70.6 ^b	70.7 ^b	68.8 ^b	64.8 ^b	65.5 ^a	65.9	66.6	66.0	66.9
Marijuana													
Smoke Once a Month	23.5 ^b	24.8 ^b	25.6 ^b	24.4 ^b	24.4 ^b	24.7 ^b	22.7 ^b	19.9 ^b	18.5 ^b	18.0 ^b	16.7 ^b	14.9 ^b	13.5
Smoke Once or Twice a Week	35.5 ^b	36.8 ^b	38.1 ^b	37.0 ^b	35.9 ^b	35.8 ^b	33.1 ^b	28.6 ^b	27.0 ^b	25.6 ^b	23.5 ^b	21.0 ^b	18.3
Cocaine													
Use Once a Month	64.1	63.6	62.6	62.4	62.5	63.2	61.8	60.9 ^b	61.9	62.0	63.5	62.8	62.9
Use Once or Twice a Week	87.2 ^b	86.6 ^b	85.9 ^b	85.8 ^b	85.7 ^b	85.8 ^b	85.7 ^b	84.2	84.5	84.8	85.5 ^a	84.3	84.4
Heroin													
Try Once or Twice	78.0 ^b	77.5 ^b	76.9 ^b	76.9 ^b	77.9 ^b	78.1 ^b	77.2 ^b	76.8 ^b	76.8 ^b	78.0 ^b	79.0 ^a	79.2	80.2
Use Once or Twice a Week	93.6	93.5	93.3	93.7	93.6	93.5	92.9	92.8 ^a	92.6 ^a	92.6 ^a	93.2	92.9	93.5
LSD													
Try Once or Twice	62.4 ^b	63.0 ^b	62.6 ^b	62.1 ^b	63.3 ^b	63.5 ^b	61.2 ^b	60.0 ^b	59.3 ^b	60.1 ^b	59.1 ^b	57.9 ^a	56.4
Use Once or Twice a Week	84.8 ^b	85.3 ^b	85.7 ^b	85.5 ^b	84.9 ^b	84.9 ^b	83.3 ^b	81.5 ^b	80.6 ^b	80.4 ^b	80.0 ^b	77.5	77.3
Alcohol													
Have Four or Five Drinks Nearly Every Day	62.1 ^b	61.1	61.7 ^b	61.8 ^b	62.2 ^b	62.4 ^b	62.9 ^b	61.8 ^b	62.0 ^b	62.0 ^b	61.4 ^a	59.8	60.0
Have Five or More Drinks Once or Twice a Week	33.2	31.9 ^a	31.5 ^b	32.7	32.6	33.2	33.0	33.2	33.7	34.5	34.5	33.2	33.5
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	77.4 ^b	77.9 ^b	76.1	76.5 ^b	76.6 ^b	76.0	75.3	76.1 ^a	75.7	75.0	75.2	75.3	74.9
Cocaine	40.5 ^b	41.2 ^b	41.0 ^b	42.0 ^b	41.7 ^b	40.7 ^b	38.1 ^b	36.0 ^b	34.0 ^b	31.7 ^b	30.1 ^b	27.9	27.5
Crack	32.7 ^b	32.5 ^b	32.2 ^b	32.8 ^b	32.5 ^b	32.1 ^b	29.0 ^b	27.4 ^b	25.2 ^b	23.4 ^b	22.0 ^b	19.7	19.0
Heroin	20.3 ^b	20.2 ^b	19.2 ^b	19.1 ^b	19.1 ^b	18.9 ^b	17.9 ^b	16.9 ^b	16.9 ^b	15.3	15.7	15.1	15.0
LSD	28.9 ^b	26.7 ^b	23.7 ^b	22.7 ^b	21.2 ^b	20.2 ^b	19.5 ^b	18.6 ^b	18.6 ^b	17.9	17.5	16.9	17.1
Approached in the Past Month by Someone Selling Drugs	19.0 ^b	19.6 ^b	18.9 ^b	18.9 ^b	18.4 ^b	18.6 ^b	18.5 ^b	18.5 ^b	19.1 ^b	18.0 ^b	18.9 ^b	16.3	16.2

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.39A Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	130,478 ^b	132,075 ^b	138,413 ^b	140,236 ^b	140,067 ^b	141,658 ^b	142,843 ^b	139,858 ^b	141,855 ^b	143,718 ^b	146,005	145,967	147,993
Marijuana													
Smoke Once a Month	73,427 ^b	76,285 ^b	77,180 ^b	75,300 ^b	78,346 ^b	76,489 ^b	75,118 ^b	71,284 ^b	67,581 ^b	67,975 ^b	66,662 ^b	61,502 ^a	58,661
Smoke Once or Twice a Week	95,135 ^b	98,394 ^b	98,443 ^b	97,357 ^b	98,575 ^b	98,496 ^b	95,862 ^b	91,608 ^b	87,164 ^b	87,410 ^b	84,534 ^b	78,509 ^b	73,636
Cocaine													
Use Once a Month	133,558 ^b	133,718 ^b	135,672 ^b	138,207 ^b	139,549 ^b	141,702 ^b	140,176 ^b	142,110 ^a	142,495 ^a	143,082	146,508	144,598	144,742
Use Once or Twice a Week	160,759 ^b	161,684 ^b	164,149 ^b	166,919 ^b	167,532 ^b	170,200 ^b	170,623 ^b	170,862 ^b	172,445 ^b	174,031 ^b	176,778	176,865	176,905
Heroin													
Try Once or Twice	152,733 ^b	153,791 ^b	153,998 ^b	157,375 ^b	159,308 ^b	160,447 ^b	161,865 ^b	163,570 ^b	164,249 ^b	167,368 ^b	170,382 ^b	170,067 ^b	174,105
Use Once or Twice a Week	168,658 ^b	170,393 ^b	172,436 ^b	174,541 ^b	176,365 ^b	178,823 ^b	180,063 ^b	181,665 ^b	182,701 ^b	185,663 ^b	187,974 ^b	188,963 ^b	191,273
LSD													
Try Once or Twice	138,085 ^b	138,342 ^b	137,209 ^b	140,669 ^b	140,545 ^b	141,972 ^b	140,857 ^b	143,242	142,317 ^b	145,114	146,807	144,200	145,185
Use Once or Twice a Week	160,367 ^b	161,277 ^b	163,552 ^b	165,607 ^b	166,960 ^b	169,240 ^b	169,194 ^b	170,004 ^b	170,048 ^b	172,386 ^a	174,387	173,848	173,923
Alcohol													
Have Four or Five Drinks Nearly Every Day	127,439 ^b	128,208 ^b	128,272 ^b	130,544 ^b	132,741 ^b	134,997 ^a	134,303 ^b	134,625 ^b	136,857	137,025	139,520	138,025	137,585
Have Five or More Drinks Once or Twice a Week	79,158 ^b	78,943 ^b	77,936 ^b	79,945 ^b	81,352 ^b	83,348	81,653 ^a	84,603	86,938 ^a	85,746	87,672 ^b	86,310	84,527
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	93,532 ^b	96,521 ^b	95,589 ^b	96,571 ^b	97,167 ^b	99,430 ^b	97,686 ^b	98,526 ^b	101,400 ^b	104,935 ^b	105,743 ^b	109,228 ^b	115,376
Cocaine	52,518 ^b	52,829 ^b	52,457 ^b	51,401 ^b	52,624 ^b	52,627 ^b	51,580 ^b	50,581 ^a	47,825	49,330	48,410	46,210	48,066
Crack	49,244 ^b	50,984 ^b	50,272 ^b	50,019 ^b	50,311 ^b	50,580 ^b	47,998 ^b	46,644 ^b	45,007	45,243	44,231	41,779 ^a	44,023
Heroin	30,554 ^b	31,359 ^b	31,232 ^b	30,753 ^b	31,805 ^b	31,278 ^b	31,016 ^b	30,644 ^b	28,785 ^b	30,629 ^b	30,801 ^b	29,962 ^b	34,321
LSD	31,463 ^b	31,785 ^b	30,627 ^b	30,111 ^b	30,544 ^b	29,713 ^b	28,651 ^b	27,453	26,766	26,932	26,822	25,726	26,150
Approached in the Past Month by Someone Selling Drugs	8,137	8,567	8,469	8,249	9,231	9,110	8,201	8,591	8,889	8,621	8,735	8,906	8,678

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.39B Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 26 or Older: Percentages, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	73.3	73.4	75.9 ^b	76.1 ^b	75.0 ^b	75.0 ^b	74.8 ^b	72.8	73.2	72.8	73.2	72.5	72.6
Marijuana													
Smoke Once a Month	41.7 ^b	42.9 ^b	42.9 ^b	41.3 ^b	42.5 ^b	41.1 ^b	39.8 ^b	37.5 ^b	35.4 ^b	34.9 ^b	33.8 ^b	31.0 ^b	29.2
Smoke Once or Twice a Week	54.1 ^b	55.4 ^b	54.7 ^b	53.6 ^b	53.5 ^b	53.0 ^b	50.9 ^b	48.2 ^b	45.6 ^b	44.9 ^b	42.9 ^b	39.6 ^b	36.6
Cocaine													
Use Once a Month	75.7 ^b	75.0 ^b	75.2 ^b	75.7 ^b	75.6 ^b	75.8 ^b	74.2 ^b	74.7 ^b	74.3 ^b	73.4 ^b	74.2 ^b	72.8	71.9
Use Once or Twice a Week	91.1 ^b	90.6 ^b	90.8 ^b	91.3 ^b	90.7 ^b	91.0 ^b	90.3 ^b	89.7 ^b	89.8 ^b	89.2 ^b	89.4 ^b	89.0 ^b	87.8
Heroin													
Try Once or Twice	86.5	86.2	85.3 ^b	86.1	86.3	85.9	85.8	85.9	85.7	85.9	86.3	85.7	86.5
Use Once or Twice a Week	95.5 ^b	95.5 ^a	95.5 ^a	95.5 ^a	95.5 ^a	95.6 ^b	95.3	95.4 ^a	95.2	95.2	95.1	95.1	94.8
LSD													
Try Once or Twice	78.5 ^b	77.9 ^b	76.4 ^b	77.5 ^b	76.6 ^b	76.4 ^b	75.0 ^b	75.7 ^b	74.6 ^b	74.9 ^b	74.8 ^b	73.1	72.5
Use Once or Twice a Week	91.2 ^b	90.9 ^b	91.0 ^b	91.1 ^b	91.0 ^b	91.0 ^b	90.2 ^b	89.7 ^b	89.1 ^b	89.0 ^b	88.7 ^b	88.1 ^b	86.8
Alcohol													
Have Four or Five Drinks Nearly Every Day	71.7 ^b	71.2 ^b	70.6 ^b	71.0 ^b	71.4 ^b	71.6 ^b	70.5 ^b	70.1 ^b	70.7 ^b	69.7 ^b	70.1 ^b	68.7 ^a	67.6
Have Five or More Drinks Once or Twice a Week	44.5 ^b	43.9 ^b	42.9 ^a	43.5 ^b	43.8 ^b	44.2 ^b	42.9 ^a	44.0 ^b	44.9 ^b	43.6 ^b	44.1 ^b	43.0 ^a	41.6
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	54.9 ^b	56.0 ^b	54.9 ^b	54.7 ^b	54.6 ^b	55.2 ^b	53.5 ^b	53.4 ^b	54.8 ^b	55.6 ^b	55.5 ^b	56.9 ^b	59.2
Cocaine	31.3 ^b	31.1 ^b	30.6 ^b	29.6 ^b	30.0 ^b	29.6 ^b	28.7 ^b	27.8 ^b	26.3 ^a	26.7 ^b	25.9	24.5	25.2
Crack	29.4 ^b	30.1 ^b	29.4 ^b	28.8 ^b	28.8 ^b	28.5 ^b	26.8 ^b	25.7 ^b	24.8 ^b	24.5 ^b	23.6	22.2	23.1
Heroin	18.3	18.6	18.3	17.8	18.2	17.7	17.4	17.0 ^a	15.9 ^b	16.6 ^b	16.5 ^b	15.9 ^b	18.0
LSD	18.9 ^b	18.8 ^b	18.0 ^b	17.4 ^b	17.5 ^b	16.9 ^b	16.1 ^b	15.2 ^b	14.8 ^a	14.7 ^a	14.4	13.7	13.8
Approached in the Past Month by Someone Selling Drugs	4.5	4.7 ^a	4.6	4.5	4.9 ^b	4.8 ^a	4.3	4.4	4.6	4.4	4.4	4.4	4.2

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.40A Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Youths Aged 12 to 17: Numbers in Thousands, 2002-2014

Behavior	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Smoking One or More Packs of Cigarettes Per Day	21,890 ^b	22,084 ^b	22,500 ^b	22,726 ^b	22,839	22,846	22,661 ^b	22,479 ^b	22,255 ^b	22,926	22,871	22,920	22,965
Trying Marijuana or Hashish Once or Twice	21,834 ^b	22,029 ^b	22,358 ^b	22,512 ^b	22,652 ^b	22,592 ^b	22,271 ^b	22,014 ^b	21,570	21,972 ^b	21,959 ^b	21,677 ^a	21,423
Using Marijuana or Hashish Once a Month or More	22,536 ^b	22,747 ^b	23,127 ^b	23,203 ^b	23,319 ^b	23,177 ^b	22,859 ^b	22,602 ^b	22,119	22,574 ^b	22,448 ^b	22,209	22,028
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	21,810 ^b	21,840 ^b	22,158	22,209	22,442 ^a	22,272	22,044	21,946 ^b	21,773 ^b	22,274	22,273	22,232	22,210

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Response categories for the questions on Youths' Perceptions of Parents' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove." Respondents with unknown Perceptions of Parents' Feelings data were excluded.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.40B Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Youths Aged 12 to 17: Percentages, 2002-2014

Behavior	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Smoking One or More Packs of Cigarettes Per Day	89.5 ^b	89.8 ^b	90.6 ^b	91.1 ^b	91.4 ^b	92.1 ^b	92.4 ^b	92.6 ^b	92.6 ^b	93.2	93.1	93.5	93.8
Trying Marijuana or Hashish Once or Twice	89.1 ^b	89.4 ^b	89.8 ^b	90.2 ^b	90.4 ^b	91.0 ^b	90.7 ^b	90.5 ^b	89.6 ^b	89.3 ^b	89.3 ^b	88.4 ^a	87.5
Using Marijuana or Hashish Once a Month or More	92.0 ^b	92.2 ^b	93.0 ^b	92.9 ^b	93.1 ^b	93.3 ^b	93.1 ^b	93.0 ^b	91.9 ^b	91.6 ^b	91.3 ^b	90.6	90.0
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	89.0 ^b	88.5 ^b	89.0 ^b	88.9 ^b	89.6 ^a	89.6 ^a	89.7 ^a	90.3	90.5	90.5	90.5	90.7	90.6

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Response categories for the questions on Youths' Perceptions of Parents' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove." Respondents with unknown Perceptions of Parents' Feelings data were excluded.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.41A Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Youths Aged 12 to 17: Numbers in Thousands, 2002-2014

Behavior	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Smoking One or More Packs of Cigarettes Per Day	21,396 ^b	21,642 ^b	21,967 ^b	22,054 ^b	22,337 ^b	22,321 ^b	22,065 ^b	21,956 ^b	21,793 ^b	22,405 ^b	22,510 ^a	22,424 ^b	22,736
Trying Marijuana or Hashish Once or Twice	19,525	19,762	20,079 ^b	20,225 ^b	20,474 ^b	20,521 ^b	20,211 ^b	19,873 ^b	19,553	19,724	19,816 ^a	19,509	19,516
Using Marijuana or Hashish Once a Month or More	19,749 ^a	19,908 ^b	20,350 ^b	20,386 ^b	20,734 ^b	20,641 ^b	20,354 ^b	19,970 ^b	19,630	19,787 ^b	19,783 ^b	19,436	19,425
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	20,798 ^b	20,875 ^b	21,215 ^b	21,447 ^b	21,664 ^b	21,577 ^b	21,451 ^b	21,278 ^b	21,212 ^b	21,708 ^b	21,854	21,778 ^a	22,015

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Response categories for the questions on Youths' Feelings about Peers include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Feelings about Peers data were excluded.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.41B Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Youths Aged 12 to 17: Percentages, 2002-2014

Behavior	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Smoking One or More Packs of Cigarettes Per Day	87.1 ^b	87.6 ^b	88.1 ^b	88.2 ^b	89.1 ^b	89.8 ^b	89.7 ^b	90.3 ^b	90.5 ^b	91.0 ^b	91.4 ^b	91.4 ^b	92.5
Trying Marijuana or Hashish Once or Twice	79.5	80.0	80.6 ^a	80.8 ^a	81.7 ^b	82.5 ^b	82.2 ^b	81.7 ^b	81.3 ^b	80.0	80.5	79.5	79.5
Using Marijuana or Hashish Once a Month or More	80.4 ^a	80.6 ^b	81.6 ^b	81.4 ^b	82.7 ^b	82.9 ^b	82.7 ^b	82.1 ^b	81.5 ^b	80.3 ^a	80.3 ^a	79.2	79.2
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	84.7 ^b	84.4 ^b	85.0 ^b	85.6 ^b	86.4 ^b	86.6 ^b	87.1 ^b	87.5 ^b	88.1 ^b	88.1 ^b	88.7 ^a	88.7 ^a	89.7

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Response categories for the questions on Youths' Feelings about Peers include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Feelings about Peers data were excluded.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.42A Exposure to Substance Use Prevention Program or Message in the Past Year among Youths Aged 12 to 17: Numbers in Thousands, 2002-2014

Program/Message	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School	3,134 ^b	3,455 ^b	3,070 ^b	2,946 ^a	2,885	2,841	2,742	2,944 ^a	2,774	2,894	2,949 ^a	2,840	2,735
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School	20,483 ^b	20,757 ^b	20,762 ^b	20,365 ^b	19,928 ^b	19,407 ^b	19,179 ^b	18,711 ^b	18,243 ^b	18,489 ^b	18,712 ^b	17,792	17,849
Drug or Alcohol Prevention Messages Were Seen or Heard in School among Youths Enrolled in School ^{1,2}	17,599 ^a	17,424	17,794 ^b	18,191 ^b	17,884 ^b	17,604 ^a	17,534 ^a	17,126	17,170	17,294	17,425	16,909	16,857

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents with unknown Substance Use Program Participation or Prevention Message Exposure data were excluded.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

² Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.42B Exposure to Substance Use Prevention Program or Message in the Past Year among Youths Aged 12 to 17: Percentages, 2002-2014

Program/Message	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School	12.7 ^b	13.9 ^b	12.2 ^b	11.7	11.4	11.4	11.1	12.1 ^a	11.5	11.7	11.9 ^a	11.5	11.1
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School	83.2 ^b	83.6 ^b	83.0 ^b	81.1 ^b	79.3 ^b	77.8 ^b	77.9 ^b	76.9 ^b	75.9 ^b	75.1 ^b	75.9 ^b	72.6	72.9
Drug or Alcohol Prevention Messages Were Seen or Heard in School among Youths Enrolled in School ^{1,2}	78.8 ^b	78.1 ^b	78.2 ^b	77.9 ^b	76.9 ^b	75.8 ^b	76.1 ^b	75.2 ^b	75.7 ^b	74.6 ^b	75.0 ^b	73.5	73.0

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents with unknown Substance Use Program Participation or Prevention Message Exposure data were excluded.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

² Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.43A Substance Use in the Past Month among Youths Aged 12 to 17, by Exposure to Substance Use Prevention Program or Message in the Past Year: Numbers in Thousands, 2002-2014

Program/Message	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School													
Illicit Drugs ¹	347 ^b	376 ^b	339 ^a	256	256	268	241	306	264	314	281	254	257
Marijuana	224	234	243	171	176	176	155	201	191	216	199	198	193
Cigarettes	404 ^b	435 ^b	355 ^b	308 ^b	258 ^b	259 ^b	221 ^b	261 ^b	225 ^b	195 ^b	213 ^b	184 ^a	131
Binge Alcohol ²	273 ^b	361 ^b	293 ^b	251 ^b	283 ^b	224 ^b	206 ^a	236 ^b	203 ^a	199 ^a	231 ^b	149	147
Did Not Participate in Alcohol, Tobacco, or Drug Prevention Program Outside of School													
Illicit Drugs ¹	2,515 ^b	2,418 ^b	2,310 ^b	2,233 ^a	2,213	2,116	2,030	2,153	2,152	2,168	2,073	1,904	2,037
Marijuana	1,793 ^a	1,728	1,649	1,539	1,507	1,493	1,482	1,594	1,578	1,718	1,583	1,533	1,608
Cigarettes	2,788 ^b	2,597 ^b	2,611 ^b	2,401 ^b	2,368 ^b	2,209 ^b	2,029 ^b	1,940 ^b	1,782 ^b	1,707 ^b	1,397 ^b	1,180	1,049
Binge Alcohol ²	2,357 ^b	2,286 ^b	2,485 ^b	2,248 ^b	2,322 ^b	2,217 ^b	1,978 ^b	1,934 ^b	1,680 ^b	1,632 ^b	1,555 ^b	1,379	1,341
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School													
Illicit Drugs ¹	2,316 ^b	2,236 ^b	2,147 ^b	1,988 ^b	1,846 ^a	1,785	1,726	1,825 ^a	1,850 ^a	1,850 ^a	1,752	1,498	1,624
Marijuana	1,647 ^b	1,553 ^b	1,539 ^b	1,378	1,245	1,241	1,284	1,332	1,345	1,457 ^a	1,328	1,222	1,273
Cigarettes	2,565 ^b	2,383 ^b	2,397 ^b	2,101 ^b	1,907 ^b	1,861 ^b	1,697 ^b	1,640 ^b	1,430 ^b	1,330 ^b	1,155 ^b	922 ^a	799
Binge Alcohol ²	2,115 ^b	2,144 ^b	2,268 ^b	2,006 ^b	2,009 ^b	1,873 ^b	1,663 ^b	1,676 ^b	1,423 ^b	1,394 ^b	1,314 ^b	1,073	1,059
Did Not See or Hear Alcohol or Drug Prevention Message from Sources Outside School													
Illicit Drugs ¹	546 ^a	561	502 ^b	495 ^b	624	589	551 ^a	632	577	625	593	671	660
Marijuana	363 ^b	410 ^a	351 ^b	335 ^b	444	414 ^a	355 ^b	465	427	472	449	517	520
Cigarettes	620 ^b	644 ^b	569 ^b	596 ^b	717 ^b	595 ^b	547 ^b	557 ^b	577 ^b	564 ^b	449	450	379
Binge Alcohol ²	517 ^a	509	511 ^a	489	596 ^b	552 ^b	522 ^a	491	458	429	472	460	424
Saw or Heard Drug or Alcohol Prevention Messages in School among Youths Enrolled in School ^{3,4}													
Illicit Drugs ¹	1,927 ^b	1,817 ^b	1,785 ^b	1,665 ^b	1,593 ^a	1,540	1,485	1,575	1,600 ^a	1,597 ^a	1,543	1,414	1,426
Marijuana	1,407 ^b	1,285 ^a	1,256	1,173	1,072	1,066	1,070	1,155	1,177	1,244	1,170	1,128	1,117
Cigarettes	2,090 ^b	1,945 ^b	1,947 ^b	1,813 ^b	1,675 ^b	1,570 ^b	1,471 ^b	1,396 ^b	1,247 ^b	1,180 ^b	1,004 ^b	852 ^b	660
Binge Alcohol ²	1,756 ^b	1,764 ^b	1,876 ^b	1,740 ^b	1,751 ^b	1,584 ^b	1,470 ^b	1,439 ^b	1,224 ^b	1,278 ^b	1,214 ^b	977	916
Did Not See or Hear Drug or Alcohol Prevention Messages in School among Youths Enrolled in School ^{3,4}													
Illicit Drugs ¹	688	726	675	682	713	677	678	730	696	776	715	621 ^b	768
Marijuana	477 ^b	545	527	467 ^b	535	515 ^a	507 ^a	562	528	636	563	528	619
Cigarettes	831 ^b	808 ^b	855 ^b	763 ^b	817 ^b	754 ^b	675 ^b	710 ^b	658 ^b	635 ^b	541	463	465
Binge Alcohol ²	672 ^b	688 ^b	730 ^b	653 ^b	715 ^b	718 ^b	638 ^b	637 ^b	594 ^a	491	502	486	496

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents with unknown Substance Use Prevention Program or Message Exposure data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

⁴ Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.43B Substance Use in the Past Month among Youths Aged 12 to 17, by Exposure to Substance Use Prevention Program or Message in the Past Year: Percentages, 2002-2014

Program/Message	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School													
Illicit Drugs ¹	11.1	10.9	11.0	8.7	8.9	9.4	8.8	10.4	9.5	10.8	9.5	8.9	9.4
Marijuana	7.1	6.8	7.9	5.8	6.1	6.2	5.7	6.8	6.9	7.5	6.7	7.0	7.0
Cigarettes	12.9 ^b	12.6 ^b	11.6 ^b	10.4 ^b	9.0 ^b	9.1 ^b	8.0 ^b	8.9 ^b	8.1 ^b	6.8 ^a	7.2 ^b	6.5 ^a	4.8
Binge Alcohol ²	8.7 ^b	10.4 ^b	9.5 ^b	8.5 ^b	9.8 ^b	7.9 ^b	7.5 ^a	8.0 ^b	7.3 ^a	6.9	7.8 ^b	5.2	5.4
Did Not Participate in Alcohol, Tobacco, or Drug Prevention Program Outside of School													
Illicit Drugs ¹	11.7 ^b	11.3 ^b	10.5 ^b	10.0	9.9	9.5	9.2	10.0	10.1	9.9	9.5	8.7	9.3
Marijuana	8.3 ^a	8.1	7.5	6.9	6.7	6.7	6.7	7.4	7.4	7.9	7.3	7.0	7.3
Cigarettes	13.0 ^b	12.1 ^b	11.9 ^b	10.8 ^b	10.6 ^b	10.0 ^b	9.2 ^b	9.0 ^b	8.4 ^b	7.8 ^b	6.4 ^b	5.4 ^a	4.8
Binge Alcohol ²	11.0 ^b	10.7 ^b	11.3 ^b	10.1 ^b	10.4 ^b	10.0 ^b	9.0 ^b	9.0 ^b	7.9 ^b	7.5 ^b	7.1 ^b	6.3	6.1
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School													
Illicit Drugs ¹	11.3 ^b	10.8 ^b	10.3 ^b	9.8	9.3	9.2	9.0	9.8	10.1 ^a	10.0	9.4	8.4	9.1
Marijuana	8.0 ^a	7.5	7.4	6.8	6.2 ^a	6.4	6.7	7.1	7.4	7.9	7.1	6.9	7.1
Cigarettes	12.5 ^b	11.5 ^b	11.5 ^b	10.3 ^b	9.6 ^b	9.6 ^b	8.9 ^b	8.8 ^b	7.8 ^b	7.2 ^b	6.2 ^b	5.2 ^a	4.5
Binge Alcohol ²	10.3 ^b	10.3 ^b	10.9 ^b	9.9 ^b	10.1 ^b	9.7 ^b	8.7 ^b	9.0 ^b	7.8 ^b	7.5 ^b	7.0 ^b	6.0	5.9
Did Not See or Hear Alcohol or Drug Prevention Message from Sources Outside School													
Illicit Drugs ¹	13.2 ^b	13.7 ^b	11.8 ^a	10.4	12.0 ^a	10.7	10.2	11.3	9.9	10.2	10.0	10.0	9.9
Marijuana	8.8	10.0 ^b	8.2	7.1	8.5	7.5	6.5	8.3	7.3	7.7	7.6	7.7	7.8
Cigarettes	15.0 ^b	15.8 ^b	13.4 ^b	12.6 ^b	13.8 ^b	10.8 ^b	10.1 ^b	9.9 ^b	9.9 ^b	9.2 ^b	7.6 ^b	6.7	5.7
Binge Alcohol ²	12.5 ^b	12.5 ^b	12.0 ^b	10.3 ^b	11.5 ^b	10.0 ^b	9.6 ^b	8.7 ^b	7.9 ^a	7.0	7.9 ^a	6.8	6.4
Saw or Heard Drug or Alcohol Prevention Messages in School among Youths Enrolled in School ^{3,4}													
Illicit Drugs ¹	10.9 ^b	10.4 ^b	10.0 ^b	9.2	8.9	8.7	8.5	9.2	9.3	9.2	8.9	8.4	8.5
Marijuana	8.0 ^b	7.4	7.1	6.4	6.0	6.1	6.1	6.7	6.9	7.2	6.7	6.7	6.6
Cigarettes	11.9 ^b	11.2 ^b	10.9 ^b	10.0 ^b	9.4 ^b	8.9 ^b	8.4 ^b	8.1 ^b	7.3 ^b	6.8 ^b	5.8 ^b	5.0 ^b	3.9
Binge Alcohol ²	10.0 ^b	10.1 ^b	10.5 ^b	9.6 ^b	9.8 ^b	9.0 ^b	8.4 ^b	8.4 ^b	7.1 ^b	7.4 ^b	7.0 ^b	5.8	5.4
Did Not See or Hear Drug or Alcohol Prevention Messages in School among Youths Enrolled in School ^{3,4}													
Illicit Drugs ¹	14.6 ^a	14.8 ^b	13.6	13.2	13.3	12.1	12.3	12.9	12.6	13.2	12.3	10.2 ^a	12.3
Marijuana	10.1	11.1	10.6	9.1	10.0	9.2	9.2	9.9	9.6	10.8	9.7	8.7	9.9
Cigarettes	17.6 ^b	16.5 ^b	17.3 ^b	14.8 ^b	15.2 ^b	13.4 ^b	12.3 ^b	12.5 ^b	11.9 ^b	10.8 ^b	9.3 ^b	7.6	7.4
Binge Alcohol ²	14.2 ^b	14.0 ^b	14.7 ^b	12.7 ^b	13.3 ^b	12.8 ^b	11.6 ^b	11.3 ^b	10.8 ^b	8.4	8.7	8.0	7.9

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents with unknown Substance Use Prevention Program or Message Exposure data were excluded from the respective analyses.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

⁴ Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.44A Past Year Initiation of Substance Use among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	2,656 ^a	2,627 ^b	2,784	2,908	2,785	2,672 ^a	2,905	3,136	2,982	3,083	2,883	2,848	3,011
Marijuana and Hashish	2,196 ^b	1,973 ^b	2,142 ^b	2,114 ^b	2,061 ^b	2,089 ^b	2,224 ^b	2,379	2,439	2,617	2,398	2,427	2,568
Cocaine	1,032 ^b	986 ^b	998 ^b	872	977 ^a	906	724	623	642	670	639	601 ^a	766
Crack	337 ^b	269 ^b	215 ^b	230 ^b	243 ^b	353 ^b	209 ^a	95	83	76	84	58	109
Heroin	117 ^a	92 ^b	118 ^a	108 ^a	90 ^b	106 ^a	116 ^a	187	142	178	156	169	212
Hallucinogens	1,152 ^b	886	934	953	1,118 ^a	1,061	1,132 ^a	1,276 ^b	1,240 ^b	1,137 ^a	1,073	1,065	936
LSD	338 ^b	200 ^b	235 ^b	243 ^b	265 ^b	271 ^b	400 ^b	341 ^b	381 ^b	358 ^b	421 ^b	482	586
PCP	123 ^b	105 ^b	106 ^b	77 ^a	70	58	53	45	46	48	90 ^a	32	41
Ecstasy	1,206 ^b	642	607	615	863 ^a	777	892 ^b	1,118 ^b	949 ^b	922 ^b	869 ^a	751	676
Inhalants	849 ^b	871 ^b	857 ^b	877 ^b	783 ^b	777 ^b	738 ^b	822 ^b	800 ^b	719 ^b	584	563	512
Nonmedical Use of Psychotherapeutics ^{2,3}	2,552 ^b	2,583 ^b	2,836 ^b	2,526 ^b	2,584 ^b	2,542 ^b	2,529 ^b	2,583 ^b	2,428 ^a	2,346	2,449 ^a	2,006	2,111
Pain Relievers	2,320 ^b	2,456 ^b	2,422 ^b	2,193 ^b	2,155 ^b	2,159 ^b	2,189 ^b	2,193 ^b	2,013 ^b	1,888 ^b	1,880 ^b	1,539	1,425
OxyContin [®]	--	--	615 ^b	526 ^b	535 ^b	557 ^b	483 ^b	588 ^b	600 ^b	483 ^b	372	436	300
Tranquilizers	1,184	1,071	1,180	1,286	1,118	1,231	1,134	1,234	1,244	1,204	1,427 ^a	1,180	1,133
Stimulants ²	783	715	793	647	846	640	602	710	626	670	676	603	690
Methamphetamine ²	299 ^b	260	318 ^b	192	259	157	97 ^a	155	107 ^a	133	133	144	183
Sedatives	209	194	240	247	267	198	183	186	253	159	166	128	173
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	2,569	2,523	2,664	2,768 ^a	2,722 ^a	2,567	2,711 ^a	2,825 ^b	2,667	2,553	2,538	2,220	2,413
CIGARETTES	1,940	1,983	2,122	2,282	2,456 ^a	2,231	2,453 ^a	2,545 ^b	2,403 ^a	2,394	2,336	2,071	2,164
Daily Cigarette Use ⁴	1,016 ^b	1,064 ^b	1,101 ^b	965 ^b	1,049 ^b	983 ^b	945 ^a	1,136 ^b	962 ^b	878	778	813	756
SMOKELESS TOBACCO⁵	951	928	999	1,134	1,335 ^b	1,299 ^b	1,413 ^b	1,468 ^b	1,425 ^b	1,297 ^b	997	1,116	1,016
CIGARS	2,858	2,736	3,058 ^b	3,349 ^b	3,061 ^b	3,078 ^b	2,918 ^a	3,146 ^b	2,950 ^a	2,800	2,664	2,770	2,597
ALCOHOL	3,942 ^b	4,082 ^b	4,396	4,274 ^a	4,378	4,551	4,466	4,561	4,675	4,699	4,589	4,559	4,655

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.44B Past Year Initiation of Substance Use among Persons Aged 12 or Older: Percentages, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	1.1	1.1	1.2	1.2	1.1	1.1	1.2	1.2	1.2	1.2	1.1	1.1	1.1
Marijuana and Hashish	0.9	0.8 ^b	0.9	0.9	0.8 ^b	0.8 ^b	0.9	0.9	1.0	1.0	0.9	0.9	1.0
Cocaine	0.4 ^b	0.4 ^b	0.4 ^b	0.4 ^a	0.4 ^b	0.4 ^a	0.3	0.2	0.3	0.3	0.2	0.2 ^a	0.3
Crack	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.0	0.0	0.0	0.0	0.0	0.0
Heroin	0.0	0.0 ^b	0.0	0.0 ^a	0.0 ^b	0.0 ^a	0.0 ^a	0.1	0.1	0.1	0.1	0.1	0.1
Hallucinogens	0.5 ^b	0.4	0.4	0.4	0.5 ^b	0.4 ^a	0.5 ^b	0.5 ^b	0.5 ^b	0.4 ^b	0.4	0.4	0.4
LSD	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.2 ^b	0.1 ^b	0.2 ^b	0.1 ^b	0.2 ^a	0.2	0.2
PCP	0.1 ^b	0.0 ^b	0.0 ^b	0.0 ^a	0.0 ^a	0.0	0.0	0.0	0.0	0.0	0.0 ^a	0.0	0.0
Ecstasy	0.5 ^b	0.3	0.3	0.3	0.4 ^b	0.3 ^a	0.4 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.3 ^b	0.3	0.3
Inhalants	0.4 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.2	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{2,3}	1.1 ^b	1.1 ^b	1.2 ^b	1.0 ^b	1.1 ^b	1.0 ^b	1.0 ^b	1.0 ^b	1.0 ^b	0.9	0.9 ^a	0.8	0.8
Pain Relievers	1.0 ^b	1.0 ^b	1.0 ^b	0.9 ^b	0.9 ^b	0.9 ^b	0.9 ^b	0.9 ^b	0.8 ^b	0.7 ^b	0.7 ^b	0.6	0.5
OxyContin [®]	--	--	0.3 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.1	0.2	0.1
Tranquilizers	0.5	0.5	0.5	0.5 ^a	0.5	0.5	0.5	0.5	0.5	0.5	0.5 ^a	0.4	0.4
Stimulants ²	0.3 ^a	0.3	0.3	0.3	0.3 ^a	0.3	0.2	0.3	0.2	0.3	0.3	0.2	0.3
Methamphetamine ²	0.1 ^b	0.1 ^a	0.1 ^b	0.1	0.1	0.1	0.0 ^a	0.1	0.0 ^a	0.1	0.1	0.1	0.1
Sedatives	0.1	0.1	0.1	0.1	0.1 ^a	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	1.1 ^b	1.1 ^b	1.1 ^b	1.1 ^b	1.1 ^b	1.0 ^a	1.1 ^b	1.1 ^b	1.1 ^a	1.0	1.0	0.8	0.9
CIGARETTES	0.8	0.8	0.9	0.9 ^a	1.0 ^b	0.9	1.0 ^b	1.0 ^b	0.9 ^b	0.9 ^a	0.9	0.8	0.8
Daily Cigarette Use ⁴	0.4 ^b	0.4 ^b	0.5 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.5 ^b	0.4 ^b	0.3	0.3	0.3	0.3
SMOKELESS TOBACCO⁵	0.4	0.4	0.4	0.5 ^a	0.5 ^b	0.5 ^b	0.6 ^b	0.6 ^b	0.6 ^b	0.5 ^b	0.4	0.4	0.4
CIGARS	1.2 ^b	1.2 ^b	1.3 ^b	1.4 ^b	1.2 ^b	1.2 ^b	1.2 ^b	1.2 ^b	1.2 ^b	1.1	1.0	1.1	1.0
ALCOHOL	1.7	1.7	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.7	1.8

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.45A Past Year Initiation of Substance Use among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLICIT DRUGS^{1,2}	1,596 ^a	1,562 ^a	1,487	1,501	1,467	1,474	1,502	1,613 ^b	1,530	1,577 ^a	1,427	1,353	1,410
Marijuana and Hashish	1,373 ^a	1,219	1,252	1,139	1,194	1,168	1,248	1,343 ^a	1,274	1,375 ^a	1,255	1,200	1,203
Cocaine	310 ^b	282 ^b	274 ^b	286 ^b	260 ^b	254 ^b	196 ^b	145	156	146	120	94	117
Crack	86 ^b	76 ^b	42 ^b	32 ^a	41 ^b	52 ^b	17	18	14	19	18	10	11
Heroin	39 ^a	25	31	18	24	16	29	19	23	38 ^a	21	21	13
Hallucinogens	474 ^b	417 ^b	437 ^b	398 ^b	397 ^b	397 ^b	469 ^b	462 ^b	498 ^b	419 ^b	358 ^b	306	258
LSD	180	96 ^b	99 ^b	105 ^a	76 ^b	97 ^b	147	106 ^a	100 ^a	123	125	122	165
PCP	77 ^b	59 ^b	43 ^a	55 ^b	43 ^a	38 ^a	37 ^a	26	22	29	45 ^a	19	17
Ecstasy	350 ^b	219 ^b	225 ^b	183 ^a	207 ^b	242 ^b	246 ^b	311 ^b	324 ^b	300 ^b	201 ^b	182 ^a	125
Inhalants	591 ^b	600 ^b	603 ^b	605 ^b	571 ^b	500 ^b	494 ^b	535 ^b	516 ^b	463 ^b	331	254	271
Nonmedical Use of													
Psychotherapeutics ^{2,3}	985 ^b	913 ^b	980 ^b	850 ^b	908 ^b	854 ^b	892 ^b	851 ^b	783 ^a	753 ^a	727	601	641
Pain Relievers	908 ^b	890 ^b	908 ^b	782 ^b	803 ^b	750 ^b	822 ^b	790 ^b	748 ^b	671 ^b	611 ^a	508	489
OxyContin [®]	--	--	141 ^a	127	136 ^a	152 ^b	156 ^b	157 ^b	159 ^b	145 ^a	122	99	95
Tranquilizers	314	319	310	274	241	267	283	280	237	298	280	219	254
Stimulants ²	341 ^b	292 ^b	271 ^b	264 ^a	296 ^b	224	196	196	190	169	183	138	183
Methamphetamine ²	131 ^b	107 ^b	99 ^a	84	85	70	45	49	45	76	31	40	51
Sedatives	63	66	66	59	63	67	69	76	65	54	44	26 ^a	59
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	1,211 ^b	1,205 ^b	1,171 ^b	1,172 ^b	1,143 ^b	1,067 ^b	1,153 ^b	1,121 ^b	1,102 ^b	1,016 ^b	914	788	814
CIGARETTES	1,187 ^b	1,226 ^b	1,294 ^b	1,303 ^b	1,333 ^b	1,198 ^b	1,288 ^b	1,273 ^b	1,205 ^b	1,165 ^b	1,032 ^b	932	838
Daily Cigarette Use ⁴	403 ^b	439 ^b	417 ^b	334 ^b	386 ^b	333 ^b	277 ^b	313 ^b	286 ^b	268 ^b	197	209	165
SMOKELESS TOBACCO⁵	459	487	447	545 ^a	594 ^b	622 ^b	612 ^b	653 ^b	572 ^b	508	415	484	431
CIGARS	1,113 ^b	1,163 ^b	1,246 ^b	1,270 ^b	1,217 ^b	1,145 ^b	1,120 ^b	1,085 ^b	940 ^a	969 ^b	849	730	797
ALCOHOL	2,588 ^b	2,593 ^b	2,743 ^b	2,749 ^b	2,706 ^b	2,698 ^b	2,568 ^a	2,662 ^b	2,476	2,622 ^b	2,448	2,417	2,335

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.45B Past Year Initiation of Substance Use among Persons Aged 12 to 17: Percentages, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	6.4 ^a	6.2	5.9	5.9	5.8	5.8	6.0	6.6 ^b	6.3	6.3 ^a	5.7	5.4	5.7
Marijuana and Hashish	5.5 ^a	4.9	5.0	4.5	4.7	4.6	5.0	5.5 ^a	5.2	5.5 ^a	5.0	4.8	4.8
Cocaine	1.3 ^b	1.1 ^b	1.1 ^b	1.1 ^b	1.0 ^b	1.0 ^b	0.8 ^b	0.6	0.6	0.6	0.5	0.4	0.5
Crack	0.3 ^b	0.3 ^b	0.2 ^b	0.1 ^a	0.2 ^b	0.2 ^b	0.1	0.1	0.1	0.1	0.1	0.0	0.0
Heroin	0.2 ^a	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2 ^a	0.1	0.1	0.1
Hallucinogens	1.9 ^b	1.7 ^b	1.7 ^b	1.6 ^b	1.6 ^b	1.6 ^b	1.9 ^b	1.9 ^b	2.0 ^b	1.7 ^b	1.4 ^b	1.2	1.0
LSD	0.7	0.4 ^b	0.4 ^b	0.4 ^a	0.3 ^b	0.4 ^b	0.6	0.4 ^a	0.4 ^a	0.5	0.5	0.5	0.7
PCP	0.3 ^b	0.2 ^b	0.2 ^a	0.2 ^b	0.2 ^a	0.2 ^a	0.1 ^a	0.1	0.1	0.1	0.2 ^a	0.1	0.1
Ecstasy	1.4 ^b	0.9 ^b	0.9 ^b	0.7 ^a	0.8 ^b	1.0 ^b	1.0 ^b	1.3 ^b	1.3 ^b	1.2 ^b	0.8 ^b	0.7 ^a	0.5
Inhalants	2.4 ^b	2.4 ^b	2.4 ^b	2.4 ^b	2.2 ^b	2.0 ^b	2.0 ^b	2.2 ^b	2.1 ^b	1.9 ^b	1.3	1.0	1.1
Nonmedical Use of Psychotherapeutics ^{2,3}	4.0 ^b	3.7 ^b	3.9 ^b	3.4 ^b	3.6 ^b	3.4 ^b	3.6 ^b	3.5 ^b	3.2 ^b	3.0 ^a	2.9	2.4	2.6
Pain Relievers	3.7 ^b	3.6 ^b	3.6 ^b	3.1 ^b	3.2 ^b	3.0 ^b	3.3 ^b	3.2 ^b	3.1 ^b	2.7 ^b	2.4 ^a	2.0	2.0
OxyContin [®]	--	--	0.6 ^a	0.5	0.5	0.6 ^b	0.6 ^b	0.6 ^b	0.7 ^b	0.6 ^a	0.5	0.4	0.4
Tranquilizers	1.3	1.3	1.2	1.1	0.9	1.1	1.1	1.1	1.0	1.2	1.1	0.9	1.0
Stimulants ²	1.4 ^b	1.2 ^b	1.1 ^b	1.0 ^a	1.2 ^b	0.9	0.8	0.8	0.8	0.7	0.7	0.6	0.7
Methamphetamine ²	0.5 ^b	0.4 ^b	0.4 ^a	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.1	0.2	0.2
Sedatives	0.3	0.3	0.3	0.2	0.2	0.3	0.3	0.3	0.3	0.2	0.2	0.1 ^a	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	4.9 ^b	4.8 ^b	4.6 ^b	4.6 ^b	4.5 ^b	4.2 ^b	4.6 ^b	4.6 ^b	4.5 ^b	4.1 ^b	3.7	3.2	3.3
CIGARETTES	4.8 ^b	4.9 ^b	5.1 ^b	5.1 ^b	5.2 ^b	4.7 ^b	5.2 ^b	5.2 ^b	4.9 ^b	4.7 ^b	4.1 ^b	3.7	3.4
Daily Cigarette Use ⁴	1.6 ^b	1.8 ^b	1.7 ^b	1.3 ^b	1.5 ^b	1.3 ^b	1.1 ^b	1.3 ^b	1.2 ^b	1.1 ^b	0.8	0.8	0.7
SMOKELESS TOBACCO⁵	1.9	1.9	1.8	2.1 ^a	2.3 ^b	2.5 ^b	2.5 ^b	2.7 ^b	2.3 ^b	2.0	1.7	1.9	1.7
CIGARS	4.5 ^b	4.7 ^b	4.9 ^b	5.0 ^b	4.8 ^b	4.5 ^b	4.5 ^b	4.4 ^b	3.9 ^b	3.9 ^b	3.4	2.9	3.2
ALCOHOL	10.5 ^b	10.4 ^b	10.9 ^b	10.8 ^b	10.7 ^b	10.7 ^b	10.3 ^a	10.8 ^b	10.2 ^a	10.5 ^b	9.8	9.7	9.4

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.46A Past Year Initiation of Substance Use among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	793 ^b	742 ^b	818 ^b	828 ^b	903 ^b	872 ^b	984	1,091	995	1,137	1,117	1,106	1,131
Marijuana and Hashish	733 ^b	666 ^b	714 ^b	723 ^b	742 ^b	787 ^b	817 ^b	988	918 ^a	1,060	966	1,017	1,094
Cocaine	594	576	592	498	570	541	426	397 ^a	372 ^a	467	443	432	501
Crack	100 ^a	109 ^b	120 ^b	142 ^b	132 ^b	88	91	62	39	40	49	25	54
Heroin	66	42	46	57	56	70	58	83	83	100	95	66	75
Hallucinogens	584	425 ^b	412 ^b	475 ^a	569	536	579	691	691	644	625	611	608
LSD	142 ^b	98 ^b	112 ^b	114 ^b	162 ^b	171 ^b	235 ^b	228 ^b	261 ^a	222 ^b	264 ^a	312	371
PCP	46	41	49	22	27	19	16	17	24	18	28	13	24
Ecstasy	613 ^b	337 ^a	303 ^b	322 ^a	495	410	523	611 ^b	543	501	543	494	448
Inhalants	247 ^b	247 ^b	225 ^a	240 ^a	191	201	221	235 ^a	225 ^a	220	218	188	157
Nonmedical Use of Psychotherapeutics ^{2,3}	937	962	945	847	900	934	959	995	882	919	890	847	884
Pain Relievers	885 ^b	930 ^b	830 ^b	803 ^b	795 ^b	857 ^b	843 ^b	846 ^b	736 ^b	701 ^a	710 ^a	637	574
OxyContin [®]	--	--	290 ^b	214 ^a	225 ^b	216 ^b	229 ^b	269 ^b	241 ^b	190	148	144	139
Tranquilizers	414	457	422	456	497	448	458	496	507	442	494	453	440
Stimulants ²	337	240 ^b	344	276	310	276	282	351	327	315	332	345	356
Methamphetamine ²	132 ^b	92	139 ^b	88	100	71	37	86	52	41	85	93	67
Sedatives	53	42	53	67	60	51	57	63	80	49	41	35	64
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	899 ^a	823 ^b	885 ^a	904	1,035	951	984	1,096	1,018	1,040	1,002	952	1,044
CIGARETTES	641 ^b	659 ^b	765 ^b	848 ^b	1,041	989 ^a	1,076	1,147	1,120	1,156	1,204	1,031	1,181
Daily Cigarette Use ⁴	447	474	566	493	554	566	549	618 ^a	599 ^a	525	488	505	479
SMOKELESS TOBACCO⁵	396	353 ^a	432	506	568	597 ^a	583	664 ^b	652 ^b	599 ^a	496	514	468
CIGARS	1,031 ^b	1,055 ^b	1,199	1,332	1,275	1,379	1,277	1,417	1,388	1,238	1,291	1,334	1,311
ALCOHOL	1,230 ^b	1,430 ^b	1,484 ^b	1,421 ^b	1,612 ^b	1,741 ^b	1,706 ^b	1,775 ^b	2,008	1,971 ^a	1,945 ^a	2,056	2,225

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.46B Past Year Initiation of Substance Use among Persons Aged 18 to 25: Percentages, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	2.6 ^b	2.3 ^b	2.5 ^b	2.5 ^b	2.8 ^a	2.7 ^a	3.0	3.2	2.9	3.3	3.2	3.2	3.2
Marijuana and Hashish	2.4 ^b	2.1 ^b	2.2 ^b	2.2 ^b	2.3 ^b	2.4 ^b	2.5 ^b	2.9	2.7	3.1	2.8	2.9	3.1
Cocaine	1.9 ^b	1.8 ^a	1.8 ^a	1.5	1.7	1.7	1.3	1.2	1.1 ^a	1.4	1.3	1.2	1.4
Crack	0.3 ^b	0.3 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.3	0.3	0.2	0.1	0.1	0.1	0.1	0.2
Heroin	0.2	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2
Hallucinogens	1.9	1.3 ^a	1.3 ^b	1.5	1.7	1.6	1.8	2.1	2.0	1.9	1.8	1.8	1.7
LSD	0.5 ^b	0.3 ^b	0.3 ^b	0.4 ^b	0.5 ^b	0.5 ^b	0.7 ^b	0.7 ^b	0.8 ^a	0.6 ^b	0.8 ^a	0.9	1.1
PCP	0.1	0.1	0.2	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.0	0.1
Ecstasy	2.0 ^b	1.1	0.9 ^a	1.0 ^a	1.5	1.3	1.6	1.8 ^b	1.6	1.5	1.6	1.4	1.3
Inhalants	0.8 ^b	0.8 ^b	0.7 ^a	0.7 ^b	0.6	0.6	0.7 ^a	0.7 ^a	0.7 ^a	0.6	0.6	0.5	0.4
Nonmedical Use of Psychotherapeutics ^{2,3}	3.0 ^a	3.0 ^a	2.9	2.6	2.7	2.9	2.9	3.0	2.6	2.7	2.6	2.4	2.5
Pain Relievers	2.9 ^b	2.9 ^b	2.6 ^b	2.5 ^b	2.4 ^b	2.6 ^b	2.6 ^b	2.5 ^b	2.2 ^b	2.0 ^a	2.1 ^a	1.8	1.6
OxyContin [®]	--	--	0.9 ^b	0.7 ^b	0.7 ^b	0.7 ^b	0.7 ^b	0.8 ^b	0.7 ^b	0.6	0.4	0.4	0.4
Tranquilizers	1.3	1.4	1.3	1.4	1.5	1.4	1.4	1.5	1.5	1.3	1.4	1.3	1.3
Stimulants ²	1.1	0.8 ^a	1.1	0.8	0.9	0.8	0.9	1.0	1.0	0.9	1.0	1.0	1.0
Methamphetamine ²	0.4 ^b	0.3	0.4 ^b	0.3	0.3	0.2	0.1	0.3	0.2	0.1	0.2	0.3	0.2
Sedatives	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	2.9	2.6	2.7	2.8	3.2	2.9	3.0	3.3	3.0	3.0	2.9	2.7	3.0
CIGARETTES	2.1 ^b	2.1 ^b	2.4 ^b	2.6 ^b	3.2	3.0	3.3	3.4	3.3	3.4	3.5	3.0	3.4
Daily Cigarette Use ⁴	1.4	1.5	1.8 ^a	1.5	1.7 ^a	1.7 ^a	1.7	1.8 ^b	1.8 ^a	1.5	1.4	1.5	1.4
SMOKELESS TOBACCO⁵	1.3	1.1	1.3	1.6	1.7 ^a	1.8 ^b	1.8 ^a	2.0 ^b	1.9 ^b	1.7 ^a	1.4	1.5	1.3
CIGARS	3.3	3.3	3.7	4.1	3.9	4.2	3.9	4.2	4.1	3.6	3.7	3.8	3.8
ALCOHOL	4.0 ^b	4.5 ^b	4.6 ^b	4.4 ^b	4.9 ^b	5.3 ^b	5.2 ^b	5.3 ^b	5.9	5.7	5.6 ^a	5.9	6.4

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.47A Past Year Initiation of Substance Use among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLICIT DRUGS^{1,2}	268 ^a	324	479	579	415	326	419	433	457	368	339	389	469
Marijuana and Hashish	90 ^b	88 ^b	176	252	126 ^a	134 ^a	159	49 ^b	247	182	177	210	271
Cocaine	127	128	133	87	147	112	102	81	114	56 ^a	76	75	148
Crack	151 ^a	83	53	55	70	212 ^a	101	15	30	17	17	23	44
Heroin	12 ^b	25 ^b	40 ^a	33 ^b	9 ^b	20 ^b	28 ^b	85	37 ^a	40 ^a	40 ^a	82	124
Hallucinogens	93	44	85	80	152	128	85	123	51	75	90	147	70
LSD	16	*	24	24	28	*	18	*	20	13	33	48	50
PCP	*	*	14	*	*	*	*	*	*	*	17	*	*
Ecstasy	243 ^a	86	78	110	161	124	124	196	82	120	124	75	104
Inhalants	11 ^a	24	29	32	22	76	23	53	58	37	36	121	84
Nonmedical Use of Psychotherapeutics ^{2,3}	630	708	911 ^b	829 ^a	775	754	679	737	763	674	833 ^a	558	585
Pain Relievers	527	636 ^a	684 ^b	608 ^a	557 ^a	552	524	557 ^a	529	516	559 ^a	394	362
OxyContin [®]	--	--	183 ^a	185 ^a	173 ^a	189 ^b	98	162	200 ^b	147	101	193	66
Tranquilizers	456	295	448	557	380	517	393	457	499	464	653	508	439
Stimulants ²	106	183	178	108	239	140	124	163	108	185	160	121	152
Methamphetamine ²	36	61	80	21	74	16 ^a	15 ^a	20	10 ^a	17 ^a	16	11 ^a	65
Sedatives	93	86	121	121	145 ^a	80	56	48	108	56	82	68	50
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	459	496	608	691	544	550	575	608	547	497	622	480	555
CIGARETTES	111	98	63 ^a	131	83	45 ^b	89	124	78	73	101	108	144
Daily Cigarette Use ⁴	166	150	118	137	109	84	119	204	77	85	92	99	113
SMOKELESS TOBACCO⁵	95	88	121	83	174	80	219	152	202	190	85	118	117
CIGARS	714 ^a	518	614	747 ^a	570	555	521	644	622	593	524	706	489
ALCOHOL	124	60	169	105	60	112	193	124	191	106	196	85	95

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.47B Past Year Initiation of Substance Use among Persons Aged 26 or Older: Percentages, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLICIT DRUGS^{1,2}	0.1	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Marijuana and Hashish	0.1 ^b	0.0 ^b	0.1	0.1	0.1 ^a	0.1 ^a	0.1	0.0 ^b	0.1	0.1	0.1	0.1	0.1
Cocaine	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.1	0.0 ^a	0.0	0.0	0.1
Crack	0.1 ^b	0.0	0.0	0.0	0.0	0.1 ^a	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Heroin	0.0 ^b	0.0 ^b	0.0	0.0 ^a	0.0 ^b	0.0 ^b	0.0 ^a	0.0	0.0 ^a	0.0 ^a	0.0 ^a	0.0	0.1
Hallucinogens	0.1	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.1	0.0
LSD	0.0	*	0.0	0.0	0.0	*	0.0	*	0.0	0.0	0.0	0.0	0.0
PCP	*	*	0.0	*	*	*	*	*	*	*	0.0	*	*
Ecstasy	0.1 ^b	0.0	0.0	0.1	0.1	0.1	0.1	0.1 ^a	0.0	0.1	0.1	0.0	0.1
Inhalants	0.0 ^a	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Nonmedical Use of Psychotherapeutics ^{2,3}	0.4	0.4	0.5 ^b	0.4 ^b	0.4 ^a	0.4	0.4	0.4	0.4	0.3	0.4 ^a	0.3	0.3
Pain Relievers	0.3 ^a	0.4 ^b	0.4 ^b	0.3 ^b	0.3 ^a	0.3 ^a	0.3 ^a	0.3 ^a	0.3 ^a	0.3	0.3 ^a	0.2	0.2
OxyContin [®]	--	--	0.1 ^b	0.1 ^a	0.1 ^b	0.1 ^b	0.1	0.1 ^a	0.1 ^b	0.1	0.1	0.1	0.0
Tranquilizers	0.3	0.2	0.2	0.3	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.3	0.2
Stimulants ²	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Methamphetamine ²	0.0	0.0	0.0	0.0	0.0	0.0 ^a	0.0 ^a	0.0	0.0 ^a	0.0	0.0	0.0 ^a	0.0
Sedatives	0.1	0.0	0.1	0.1	0.1 ^a	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.3	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3
CIGARETTES	0.1	0.1	0.0 ^a	0.1	0.0	0.0 ^b	0.0	0.1	0.0	0.0	0.1	0.1	0.1
Daily Cigarette Use ⁴	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.1
SMOKELESS TOBACCO⁵	0.1	0.0	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.0	0.1	0.1
CIGARS	0.4 ^b	0.3	0.3	0.4 ^b	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2
ALCOHOL	0.1	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.48A Past Year Initiation of Illicit Drugs and First Drug Initiated in the Past Year among Persons Aged 12 or Older Who Initiated Illicit Drug Use in the Past Year: Numbers in Thousands, 2002-2014

	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PAST YEAR INITIATION OF ILLICIT DRUGS^{1,2}	2,656 ^a	2,627 ^b	2,784	2,908	2,785	2,672 ^a	2,905	3,136	2,982	3,083	2,883	2,848	3,011
FIRST DRUG INITIATED³													
Marijuana and Hashish	1,555 ^b	1,463 ^b	1,516 ^b	1,586 ^b	1,508 ^b	1,502 ^b	1,649 ^b	1,856	1,848	2,080	1,890	2,003	2,035
Cocaine	29	21	43	29	34	16	24	8	4	6	4	3	16
Crack	*	*	*	1	*	*	*	*	*	*	*	*	*
Heroin	1	1	1	*	0	*	4	3	2	2	2	*	*
Hallucinogens	88	63	56	67	92	55	92	66	90	86	57	74	64
LSD	1	5	1	2	6	4	2	1	2	3	2	6	10
PCP	2	1	1	1	9	*	1	0	2	3	5	1	*
Ecstasy	52	37	26	23	46	15	28	30	41	42	20	26	29
Inhalants	246 ^a	340 ^b	305 ^b	331 ^b	276 ^b	288 ^b	282 ^b	309 ^b	268 ^b	232	182	181	177
Nonmedical Use of Psychotherapeutics ^{2,4}	742	742	869	900	878	817	857	895	777	679	749	588	719
Pain Relievers	525 ^a	520	635 ^b	596 ^b	618 ^b	508 ^a	652 ^b	535 ^a	514 ^a	432	489	357	382
OxyContin [®]	--	--	1	8	1 ^a	2	6	24	15	29	10	*	12
Tranquilizers	64 ^b	87 ^a	109	179	112	173	93	268	136	130	118	148	165
Stimulants ²	92	102	99	73	114	108	88	61 ^a	73	80	104	76	135
Methamphetamine ²	13	8	18	8	7	16	11	8	2	3	*	3	8
Sedatives	61	34	27	52	35	28	24	30	57	36	38	6 ^b	37
Illicit Drugs Other Than Marijuana ^{1,2}	1,104	1,164	1,269 ^b	1,328 ^b	1,282 ^b	1,176 ^a	1,258 ^b	1,281 ^b	1,141	1,004	993	845	976

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ First Drug Initiated is based on imputation-revised month and year and imputed day of first use for each drug. Respondents may be counted in more than one drug category if they reported initiating multiple drugs on the same day the first time they used drugs. Also, several drugs listed are subcategories of other drugs, and will have the same start date. Therefore, the sum of the estimates for the first drug or drugs initiated may not equal the total estimates for past year initiates.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.48B Past Year Initiation of Illicit Drugs and First Drug Initiated in the Past Year among Persons Aged 12 or Older Who Initiated Illicit Drug Use in the Past Year: Percentages, 2002-2014

	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PAST YEAR INITIATION OF ILLICIT DRUGS^{1,2}	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
FIRST DRUG INITIATED³													
Marijuana and Hashish	58.6 ^b	55.7 ^b	54.5 ^b	54.5 ^b	54.1 ^b	56.2 ^b	56.7 ^b	59.2 ^b	62.0 ^a	67.5	65.6	70.3	67.6
Cocaine	1.1	0.8	1.5	1.0	1.2	0.6	0.8	0.3	0.1	0.2	0.1	0.1	0.5
Crack	*	*	*	0.0	*	*	*	*	*	*	*	*	*
Heroin	0.0	0.0	0.0	*	0.0	*	0.1	0.1	0.1	0.1	0.1	*	*
Hallucinogens	3.3	2.4	2.0	2.3	3.3	2.1	3.2	2.1	3.0	2.8	2.0	2.6	2.1
LSD	0.0	0.2	0.0	0.1	0.2	0.2	0.1	0.0	0.1	0.1	0.1	0.2	0.3
PCP	0.1	0.0	0.0	0.0	0.3	*	0.0	0.0	0.1	0.1	0.2	0.0	*
Ecstasy	1.9	1.4	0.9	0.8	1.7	0.6	1.0	0.9	1.4	1.4	0.7	0.9	1.0
Inhalants	9.3 ^b	12.9 ^b	10.9 ^b	11.4 ^b	9.9 ^b	10.8 ^b	9.7 ^b	9.9 ^b	9.0 ^b	7.5	6.3	6.3	5.9
Nonmedical Use of Psychotherapeutics ^{2,4}													
Pain Relievers	19.8 ^b	19.8 ^b	22.8 ^b	20.5 ^b	22.2 ^b	19.0 ^b	22.5 ^b	17.1 ^a	17.2 ^a	14.0	17.0 ^a	12.5	12.7
OxyContin [®]	--	--	0.0	0.3	0.0	0.1	0.2	0.8	0.5	1.0	0.3	*	0.4
Tranquilizers	2.4 ^b	3.3	3.9	6.2	4.0	6.5	3.2	8.5	4.6	4.2	4.1	5.2	5.5
Stimulants ²	3.5	3.9	3.5	2.5	4.1	4.0	3.0	2.0 ^a	2.5	2.6	3.6	2.7	4.5
Methamphetamine ²	0.5	0.3	0.7	0.3	0.3	0.6	0.4	0.3	0.1	0.1	*	0.1	0.3
Sedatives	2.3	1.3	1.0	1.8	1.2	1.1	0.8	0.9	1.9	1.2	1.3	0.2 ^b	1.2
Illicit Drugs Other Than Marijuana ^{1,2}	41.6 ^b	44.3 ^b	45.6 ^b	45.7 ^b	46.0 ^b	44.0 ^b	43.3 ^b	40.9 ^b	38.3 ^a	32.6	34.5	29.7	32.4

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ First Drug Initiated is based on imputation-revised month and year and imputed day of first use for each drug. Respondents may be counted in more than one drug category if they reported initiating multiple drugs on the same day the first time they used drugs. Also, several drugs listed are subcategories of other drugs, and will have the same start date. Therefore, the sum of the estimates for the first drug or drugs initiated may not equal the total estimates for past year initiates.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.49B Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49: 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	17.9 ^a	17.5 ^b	18.1	18.7	19.0	18.0 ^a	18.8	17.6 ^b	19.1	18.1	18.7	19.0	19.0
Marijuana and Hashish	17.0 ^b	16.8 ^b	17.1 ^b	17.4 ^b	17.4 ^b	17.6 ^a	17.8	17.0 ^b	18.4	17.5 ^b	17.9	18.0	18.5
Cocaine	19.8 ^b	19.8 ^b	20.0 ^b	19.7 ^b	20.3 ^a	20.2 ^a	19.8 ^b	19.9 ^b	21.2	20.1 ^a	20.0 ^b	20.4 ^a	21.8
Crack	25.0	21.8	21.9	23.4	22.8	29.6	27.1	20.6 ^a	24.8	20.8	20.5	24.4	26.4
Heroin	21.0 ^a	20.9 ^b	24.4	22.2 ^b	20.7 ^b	21.8 ^b	23.5	25.3	21.4 ^b	22.1 ^b	23.0 ^b	24.5	28.0
Hallucinogens	18.5	17.9 ^b	18.7	18.7	19.7	19.1	18.6	18.9	18.3 ^b	18.7	19.1	19.9	19.3
LSD	17.4 ^b	17.2 ^b	18.4	18.3	19.4	18.2 ^b	18.4 ^b	18.3 ^b	19.0	18.6 ^a	19.0	19.7	19.7
PCP	16.0 ^a	17.4	18.9	16.5	16.3	16.4	15.9 ^a	16.8	17.6	17.8	16.6	17.1	17.7
Ecstasy	21.2	19.7	19.5	20.7	20.6	20.3	20.3	20.2	19.4 ^a	19.6 ^a	20.3	20.5	21.0
Inhalants	15.9 ^b	16.0 ^b	16.0 ^b	16.1 ^a	15.7 ^b	17.1	15.9 ^b	16.9	16.3 ^a	16.4 ^a	16.9	19.2	18.2
Nonmedical Use of Psychotherapeutics ^{2,3}	21.3	21.6	21.7	22.2	22.9	21.8	22.0	21.0	22.4	22.4	22.9	22.4	21.8
Pain Relievers	20.8	21.1	20.9	21.2	22.0	21.2	21.2	20.8	21.0	21.8	22.3	21.7	21.2
OxyContin [®]	--	--	22.9	23.2	22.6	24.0	21.7	22.3	22.8	22.8	22.0	23.6	21.2
Tranquilizers	23.8	22.7	24.4	25.7 ^a	24.0	24.5	24.4	22.4	24.6	24.6	23.6	25.4	23.4
Stimulants ²	19.2 ^a	21.5	20.0	20.1	23.0	21.9	21.3	21.5	21.2	22.2	22.1	21.6	21.6
Methamphetamine ²	18.9 ^a	20.4	20.6	18.6 ^a	22.2	19.4	19.3	19.3	18.8 ^a	17.8 ^b	19.7	18.9 ^a	22.0
Sedatives	26.6	21.2	23.5	22.9	26.5 ^a	24.3	21.6	19.7	23.5	22.0	26.2 ^a	25.0	21.4
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	19.7	19.5 ^a	19.9	20.5	20.7	20.0	20.4	19.3 ^b	20.2	20.1	21.2	20.6	20.6
CIGARETTES	16.9 ^b	16.9 ^b	16.7 ^b	17.3 ^b	17.1 ^b	16.9 ^b	17.4 ^b	17.5 ^b	17.3 ^b	17.2 ^b	17.8 ^a	17.8 ^a	18.6
Daily Cigarette Use ⁴	19.9	19.8	18.8 ^b	19.7	18.9 ^a	19.2 ^a	20.1	20.7	19.1 ^a	19.1 ^a	19.9	19.8	20.6
SMOKELESS TOBACCO⁵	19.1	18.1	18.8	18.3	19.0	18.0 ^a	18.9	18.9	19.3	19.8	18.8	18.4	19.0
CIGARS	21.8 ^a	20.3	20.2	21.2	19.9	20.5	20.0	20.7	20.5	19.6	20.5	21.6	20.4
ALCOHOL	16.6 ^b	16.4 ^b	16.4 ^b	16.4 ^b	16.6 ^b	16.8 ^b	17.0	16.9 ^b	17.1	17.1	17.4	17.3	17.3

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.50A Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	7,116	6,835	7,298	6,833	7,024	6,866	7,012	7,114	7,144	6,531	7,312	6,852	7,077
Marijuana and Hashish	4,294	4,198	4,469	4,090	4,184	3,941	4,228	4,322	4,505	4,165	4,304	4,206	4,176
Cocaine	1,488 ^b	1,515 ^b	1,571 ^b	1,549 ^b	1,665 ^b	1,604 ^b	1,412 ^b	1,108	1,012	821	1,119	855	913
Heroin	214 ^b	189 ^b	270 ^b	227 ^b	324 ^b	214 ^b	283 ^b	369 ^a	361 ^b	426	467	517	586
Hallucinogens	426 ^b	321	449 ^b	371 ^a	380 ^a	369 ^a	362 ^a	373 ^a	402 ^b	342	331	277	246
Inhalants	180 ^a	169 ^a	233 ^b	221 ^b	176 ^a	164 ^a	175 ^a	164 ^a	169 ^a	141	164	132	96
Nonmedical Use of Psychotherapeutics ^{2,3}	2,018 ^a	1,923 ^b	2,048 ^a	1,959 ^b	2,036 ^a	2,167	2,177	2,297	2,378	2,139	2,597	2,281	2,417
Pain Relievers	1,509 ^a	1,424 ^b	1,388 ^b	1,546 ^a	1,636	1,715	1,715	1,878	1,923	1,768	2,056	1,879	1,918
Tranquilizers	509	435	573	419	403	443	453	476	522	400	629	423	472
Stimulants ²	436	378	470	409	388	405	351	380	358	329 ^a	535	469	476
Sedatives	154	158	128	97	121	154	127	147	162	78	135	99	143
ALCOHOL	18,100 ^a	17,805	18,654 ^b	18,658 ^b	18,852 ^b	18,687 ^b	18,478 ^b	18,763 ^b	17,967	16,672	17,714	17,298	16,994
BOTH ILLICIT DRUGS AND ALCOHOL¹	3,210 ^b	3,054 ^a	3,445 ^b	3,273 ^b	3,215 ^b	3,184 ^b	3,102 ^b	3,243 ^b	2,889	2,598	2,840	2,589	2,592
ILLCIT DRUGS OR ALCOHOL¹	22,006	21,586	22,506	22,218	22,661 ^a	22,369	22,388	22,634	22,221	20,605	22,187	21,561	21,480

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.50B Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	3.0 ^b	2.9	3.0 ^b	2.8	2.9	2.8	2.8	2.8	2.8	2.5	2.8	2.6	2.7
Marijuana and Hashish	1.8 ^b	1.8 ^a	1.9 ^b	1.7	1.7	1.6	1.7	1.7	1.8 ^a	1.6	1.7	1.6	1.6
Cocaine	0.6 ^b	0.6 ^b	0.7 ^b	0.6 ^b	0.7 ^b	0.6 ^b	0.6 ^b	0.4	0.4	0.3	0.4	0.3	0.3
Heroin	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^a	0.1 ^b	0.1 ^b	0.1 ^a	0.1 ^a	0.2	0.2	0.2	0.2
Hallucinogens	0.2 ^b	0.1 ^a	0.2 ^b	0.2 ^b	0.2 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.2 ^b	0.1	0.1	0.1	0.1
Inhalants	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^a	0.1 ^b	0.1 ^a	0.1	0.1	0.1	0.0
Nonmedical Use of Psychotherapeutics ^{2,3}	0.9	0.8	0.9	0.8	0.8	0.9	0.9	0.9	0.9	0.8	1.0	0.9	0.9
Pain Relievers	0.6	0.6 ^a	0.6 ^b	0.6	0.7	0.7	0.7	0.7	0.8	0.7	0.8	0.7	0.7
Tranquilizers	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Stimulants ²	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.1	0.1 ^a	0.2	0.2	0.2
Sedatives	0.1	0.1	0.1	0.0	0.0	0.1	0.1	0.1	0.1	0.0	0.1	0.0	0.1
ALCOHOL	7.7 ^b	7.5 ^b	7.8 ^b	7.7 ^b	7.7 ^b	7.5 ^b	7.4 ^b	7.5 ^b	7.1 ^b	6.5	6.8	6.6	6.4
BOTH ILLICIT DRUGS AND ALCOHOL¹	1.4 ^b	1.3 ^b	1.4 ^b	1.3 ^b	1.3 ^b	1.3 ^b	1.2 ^b	1.3 ^b	1.1 ^a	1.0	1.1	1.0	1.0
ILLCIT DRUGS OR ALCOHOL¹	9.4 ^b	9.1 ^b	9.4 ^b	9.1 ^b	9.2 ^b	9.0 ^b	9.0 ^b	9.0 ^b	8.8 ^b	8.0	8.5	8.2	8.1

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.51A Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	1,376 ^b	1,287 ^b	1,347 ^b	1,186 ^b	1,175 ^b	1,092 ^b	1,155 ^b	1,058 ^b	1,154 ^b	1,157 ^b	996 ^a	875	867
Marijuana and Hashish	1,055 ^b	955 ^b	979 ^b	907 ^b	869 ^b	786 ^a	846 ^b	834 ^b	868 ^b	879 ^b	806 ^a	715	667
Cocaine	105 ^b	86 ^b	91 ^b	93 ^b	94 ^b	96 ^b	83 ^b	45	33	42	39	22	27
Heroin	13	12	21	9	12	8	17	18	7	30	20	10	18
Hallucinogens	138 ^b	106 ^b	121 ^b	128 ^b	121 ^b	117 ^b	96 ^a	109 ^b	138 ^b	85	87	63	55
Inhalants	101 ^b	104 ^b	134 ^b	108 ^b	108 ^b	99 ^b	96 ^b	92 ^b	88 ^b	79 ^b	59	41	39
Nonmedical Use of Psychotherapeutics ^{2,3}	333 ^b	361 ^b	388 ^b	329 ^a	350 ^b	316 ^a	337 ^b	264	293	301	203	177 ^a	241
Pain Relievers	237 ^a	281 ^b	309 ^b	275 ^b	265 ^b	238 ^a	254 ^b	218	238 ^a	250 ^b	150	134	168
Tranquilizers	87 ^a	96 ^a	75	71	62	61	73	52	68	62	54	40	52
Stimulants ²	98 ^a	98 ^a	86	73	95	73	68	58	53	68	50	42	60
Sedatives	28	40	22	34	23	23	23	19	16	12	18	11	29
ALCOHOL	1,453 ^b	1,471 ^b	1,517 ^b	1,407 ^b	1,362 ^b	1,367 ^b	1,220 ^b	1,136 ^b	1,109 ^b	947 ^b	855 ^b	697	679
BOTH ILLICIT DRUGS AND ALCOHOL¹	620 ^b	544 ^b	635 ^b	555 ^b	492 ^b	517 ^b	466 ^b	456 ^b	477 ^b	386 ^a	334	270	296
ILLCIT DRUGS OR ALCOHOL¹	2,209 ^b	2,214 ^b	2,228 ^b	2,038 ^b	2,045 ^b	1,942 ^b	1,908 ^b	1,738 ^b	1,786 ^b	1,718 ^b	1,518 ^b	1,302	1,251

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.51B Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	5.6 ^b	5.1 ^b	5.3 ^b	4.7 ^b	4.6 ^b	4.3 ^b	4.6 ^b	4.3 ^b	4.7 ^b	4.6 ^b	4.0 ^a	3.5	3.5
Marijuana and Hashish	4.3 ^b	3.8 ^b	3.9 ^b	3.6 ^b	3.4 ^b	3.1 ^a	3.4 ^b	3.4 ^b	3.6 ^b	3.5 ^b	3.2 ^a	2.9	2.7
Cocaine	0.4 ^b	0.3 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.3 ^b	0.2	0.1	0.2	0.2	0.1	0.1
Heroin	0.1	0.0	0.1	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.1	0.0	0.1
Hallucinogens	0.6 ^b	0.4 ^b	0.5 ^b	0.5 ^b	0.5 ^b	0.5 ^b	0.4 ^a	0.4 ^b	0.6 ^b	0.3	0.3	0.3	0.2
Inhalants	0.4 ^b	0.4 ^b	0.5 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.3 ^a	0.2	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{2,3}	1.3 ^b	1.4 ^b	1.5 ^b	1.3 ^a	1.4 ^b	1.3 ^a	1.4 ^b	1.1	1.2	1.2	0.8	0.7 ^a	1.0
Pain Relievers	1.0 ^a	1.1 ^b	1.2 ^b	1.1 ^b	1.0 ^b	0.9 ^a	1.0 ^b	0.9 ^a	1.0 ^b	1.0 ^b	0.6	0.5	0.7
Tranquilizers	0.4 ^a	0.4 ^a	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.2	0.2	0.2	0.2
Stimulants ²	0.4 ^a	0.4 ^a	0.3	0.3	0.4	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.2
Sedatives	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.0	0.1
ALCOHOL	5.9 ^b	5.9 ^b	6.0 ^b	5.5 ^b	5.4 ^b	5.4 ^b	4.9 ^b	4.6 ^b	4.6 ^b	3.8 ^b	3.4 ^b	2.8	2.7
BOTH ILLICIT DRUGS AND ALCOHOL¹	2.5 ^b	2.2 ^b	2.5 ^b	2.2 ^b	1.9 ^b	2.1 ^b	1.9 ^b	1.9 ^b	2.0 ^b	1.5 ^a	1.3	1.1	1.2
ILLCIT DRUGS OR ALCOHOL¹	8.9 ^b	8.9 ^b	8.8 ^b	8.0 ^b	8.1 ^b	7.7 ^b	7.7 ^b	7.1 ^b	7.3 ^b	6.9 ^b	6.1 ^b	5.2	5.0

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.52A Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	2,548	2,486	2,684 ^b	2,728 ^b	2,591 ^a	2,599 ^a	2,591 ^a	2,596 ^a	2,677 ^b	2,574 ^a	2,706 ^b	2,557	2,320
Marijuana and Hashish	1,860	1,886	1,936 ^a	1,911	1,862	1,840	1,861	1,865	1,955 ^a	1,948 ^a	1,900	1,878	1,706
Cocaine	377 ^b	393 ^b	462 ^b	484 ^b	422 ^b	451 ^b	393 ^b	307 ^b	230	204	217	232	185
Heroin	47 ^b	42 ^b	70 ^b	89 ^b	66 ^b	77 ^b	99 ^a	104 ^a	119	139	173	182	168
Hallucinogens	242 ^b	152	214 ^b	188 ^a	192 ^a	173	198 ^a	186 ^a	193 ^a	156	123	147	120
Inhalants	29	41 ^a	52 ^b	49 ^a	28	28	26	54 ^b	44 ^a	37	14	47 ^a	14
Nonmedical Use of Psychotherapeutics ^{2,3}	587	516	652	702 ^a	659	697 ^a	735 ^b	736 ^b	760 ^b	687 ^a	843 ^b	666	559
Pain Relievers	419	350	454	541 ^a	486	559 ^a	580 ^b	565 ^a	609 ^b	580 ^b	649 ^b	485	430
Tranquilizers	144	147	154 ^a	176 ^b	154	134	139	156 ^a	144	133	183 ^a	147	104
Stimulants ²	137	142	172	170	140	117	117	146	139	90	170	171	134
Sedatives	26	20	24	20	31	20	33	38	26	18	16	30	20
ALCOHOL	5,477 ^b	5,462 ^b	5,609 ^b	5,692 ^b	5,778 ^b	5,526 ^b	5,726 ^b	5,422 ^b	5,342 ^b	4,936 ^b	4,956 ^b	4,529	4,284
BOTH ILLICIT DRUGS AND ALCOHOL¹	1,292 ^b	1,270 ^b	1,453 ^b	1,351 ^b	1,371 ^b	1,336 ^b	1,405 ^b	1,266 ^b	1,216 ^b	1,141 ^b	1,127 ^b	1,073 ^a	908
ILLCIT DRUGS OR ALCOHOL¹	6,733 ^b	6,678 ^b	6,840 ^b	7,069 ^b	6,998 ^b	6,790 ^b	6,912 ^b	6,753 ^b	6,803 ^b	6,369 ^b	6,535 ^b	6,014	5,696

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.52B Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	8.2 ^b	7.8 ^b	8.3 ^b	8.4 ^b	7.9 ^b	7.9 ^b	7.9 ^b	7.7 ^b	7.9 ^b	7.5 ^a	7.8 ^b	7.4	6.6
Marijuana and Hashish	6.0 ^b	5.9 ^b	6.0 ^b	5.9 ^b	5.7 ^b	5.6 ^a	5.6 ^a	5.6 ^a	5.7 ^b	5.7 ^a	5.5	5.4	4.9
Cocaine	1.2 ^b	1.2 ^b	1.4 ^b	1.5 ^b	1.3 ^b	1.4 ^b	1.2 ^b	0.9 ^b	0.7	0.6	0.6	0.7	0.5
Heroin	0.2 ^b	0.1 ^b	0.2 ^b	0.3 ^a	0.2 ^b	0.2 ^b	0.3 ^a	0.3 ^a	0.3	0.4	0.5	0.5	0.5
Hallucinogens	0.8 ^b	0.5	0.7 ^b	0.6 ^b	0.6 ^b	0.5 ^a	0.6 ^b	0.6 ^a	0.6 ^a	0.5	0.4	0.4	0.3
Inhalants	0.1	0.1 ^a	0.2 ^b	0.2 ^a	0.1	0.1	0.1	0.2 ^b	0.1 ^a	0.1 ^a	0.0	0.1 ^a	0.0
Nonmedical Use of Psychotherapeutics ^{2,3}	1.9	1.6	2.0 ^a	2.2 ^b	2.0 ^a	2.1 ^b	2.2 ^b	2.2 ^b	2.2 ^b	2.0 ^a	2.4 ^b	1.9	1.6
Pain Relievers	1.4	1.1	1.4	1.7 ^b	1.5	1.7 ^b	1.8 ^b	1.7 ^b	1.8 ^b	1.7 ^b	1.9 ^b	1.4	1.2
Tranquilizers	0.5 ^a	0.5 ^a	0.5 ^a	0.5 ^b	0.5 ^a	0.4	0.4	0.5 ^a	0.4	0.4	0.5 ^a	0.4	0.3
Stimulants ²	0.4	0.4	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.5	0.5	0.4
Sedatives	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1
ALCOHOL	17.7 ^b	17.2 ^b	17.4 ^b	17.5 ^b	17.6 ^b	16.9 ^b	17.4 ^b	16.1 ^b	15.7 ^b	14.4 ^b	14.3 ^b	13.0	12.3
BOTH ILLICIT DRUGS AND ALCOHOL¹	4.2 ^b	4.0 ^b	4.5 ^b	4.2 ^b	4.2 ^b	4.1 ^b	4.3 ^b	3.8 ^b	3.6 ^b	3.3 ^b	3.3 ^b	3.1 ^a	2.6
ILLCIT DRUGS OR ALCOHOL¹	21.7 ^b	21.0 ^b	21.2 ^b	21.8 ^b	21.4 ^b	20.7 ^b	21.0 ^b	20.1 ^b	20.0 ^b	18.6 ^b	18.9 ^b	17.3	16.3

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.53A Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	3,192 ^a	3,062 ^b	3,267 ^a	2,918 ^b	3,258 ^a	3,175 ^a	3,266 ^a	3,460	3,313 ^a	2,801 ^b	3,610	3,419	3,890
Marijuana and Hashish	1,378 ^a	1,357 ^b	1,554	1,271 ^b	1,453	1,316 ^b	1,521	1,623	1,683	1,338 ^b	1,598	1,612	1,804
Cocaine	1,006 ^a	1,036 ^a	1,018 ^a	972 ^a	1,150 ^b	1,057 ^a	937	756	749	575	862	601	702
Heroin	153 ^b	135 ^b	179 ^b	129 ^b	245	129 ^b	167 ^b	246	236 ^a	256	274	325	400
Hallucinogens	46	63	114	55	67	79	67	77	71	101	121	67	71
Inhalants	50	25	47	64	40	36	52	17	37	25	92	45	42
Nonmedical Use of Psychotherapeutics ^{2,3}	1,098 ^b	1,046 ^b	1,008 ^b	928 ^b	1,027 ^b	1,153 ^b	1,105 ^b	1,298	1,325	1,150 ^b	1,552	1,438	1,617
Pain Relievers	853 ^b	793 ^b	624 ^b	730 ^b	885 ^b	919 ^b	881 ^b	1,095	1,075	938 ^b	1,258	1,260	1,320
Tranquilizers	278	192	344	173 ^a	187	247	241	268	311	205	392	236	316
Stimulants ²	202	138 ^a	212	166	153 ^a	215	166	176	167	172	315	256	281
Sedatives	100	99	82	44	67	110	72	91	121	48	101	58	94
ALCOHOL	11,169	10,872 ^a	11,528	11,560	11,711	11,793	11,532	12,204	11,516	10,788 ^b	11,903	12,072	12,032
BOTH ILLICIT DRUGS AND ALCOHOL¹	1,298	1,240	1,357	1,367	1,352	1,331	1,231	1,522	1,197	1,071 ^a	1,379	1,246	1,388
ILLCIT DRUGS OR ALCOHOL¹	13,064 ^b	12,694 ^b	13,438 ^a	13,111 ^b	13,618	13,638	13,568	14,142	13,631	12,518 ^b	14,134	14,245	14,534

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.53B Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	1.8	1.7	1.8	1.6 ^a	1.7	1.7	1.7	1.8	1.7	1.4 ^b	1.8	1.7	1.9
Marijuana and Hashish	0.8	0.7	0.8	0.7 ^a	0.8	0.7 ^a	0.8	0.8	0.9	0.7 ^a	0.8	0.8	0.9
Cocaine	0.6 ^b	0.6 ^b	0.6 ^b	0.5 ^b	0.6 ^b	0.6 ^b	0.5 ^a	0.4	0.4	0.3	0.4	0.3	0.3
Heroin	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1	0.1 ^b	0.1 ^b	0.1	0.1	0.1	0.1	0.2	0.2
Hallucinogens	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0
Inhalants	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Nonmedical Use of Psychotherapeutics ^{2,3}	0.6	0.6 ^b	0.6 ^b	0.5 ^b	0.5 ^b	0.6 ^a	0.6 ^b	0.7	0.7	0.6 ^b	0.8	0.7	0.8
Pain Relievers	0.5 ^a	0.4 ^b	0.3 ^b	0.4 ^b	0.5 ^a	0.5 ^a	0.5 ^b	0.6	0.6	0.5 ^a	0.6	0.6	0.6
Tranquilizers	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.2	0.1	0.2
Stimulants ²	0.1	0.1 ^a	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1
Sedatives	0.1	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.0	0.0
ALCOHOL	6.2	6.0	6.3	6.2	6.2	6.2	6.0	6.3	5.9	5.4	5.9	6.0	5.9
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.8	0.6	0.5	0.7	0.6	0.7
ILLCIT DRUGS OR ALCOHOL¹	7.3	7.0	7.3	7.1	7.2	7.2	7.1	7.3	7.0	6.3 ^b	7.0	7.0	7.1

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.54A Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	4,636	4,374 ^a	4,852	4,718	4,840	4,696	4,879	4,888	4,848	4,566	5,056	4,917	4,991
Marijuana and Hashish	2,614	2,520	2,670	2,619	2,625	2,508	2,701	2,644	2,813	2,641	2,690	2,804	2,656
Cocaine	1,025 ^b	961 ^b	1,175 ^b	1,144 ^b	1,205 ^b	1,166 ^b	959 ^b	741	689	581	815 ^a	624	571
Heroin	198 ^b	169 ^b	226 ^b	199 ^b	298 ^b	179 ^b	254 ^b	324 ^a	321 ^b	369 ^a	438	434	539
Hallucinogens	136	135	180 ^b	143	119	107	126	185 ^a	158 ^a	143	124	102	94
Inhalants	47	37	90	46	48	45	83	64	65	47	34	39	43
Nonmedical Use of Psychotherapeutics ^{2,3}	1,242 ^b	1,275 ^b	1,395 ^b	1,366 ^b	1,312 ^b	1,556	1,525 ^a	1,731	1,758	1,625	1,929	1,758	1,846
Pain Relievers	936 ^b	943 ^b	970 ^b	1,102 ^b	1,073 ^b	1,270	1,240 ^a	1,407	1,400	1,370	1,569	1,488	1,507
Tranquilizers	251	196	252	243	178	201	233	261	313	178	315	237	284
Stimulants ²	259	260	359	273	276	303	234	299	251	252	379	367	335
Sedatives	131	112	89	54	53	94	92	115	130	67	110	61	115
ALCOHOL	8,222	7,563	8,355	7,974	8,423	8,468	8,825 ^a	8,945 ^a	8,340	7,756	8,627	7,998	8,063
BOTH ILLICIT DRUGS AND ALCOHOL¹	1,334	1,066	1,414 ^a	1,366 ^a	1,330	1,372 ^a	1,323	1,379 ^a	1,171	1,159	1,267	1,098	1,086
ILLCIT DRUGS OR ALCOHOL¹	11,524	10,871 ^b	11,794	11,326	11,933	11,792	12,381	12,453	12,017	11,163	12,416	11,817	11,968

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.54B Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	2.0	1.8	2.0	1.9	2.0	1.9	2.0	1.9	1.9	1.8	1.9	1.9	1.9
Marijuana and Hashish	1.1	1.1	1.1	1.1	1.1	1.0	1.1	1.0	1.1	1.0	1.0	1.1	1.0
Cocaine	0.4 ^b	0.4 ^b	0.5 ^b	0.5 ^b	0.5 ^b	0.5 ^b	0.4 ^b	0.3	0.3	0.2	0.3 ^a	0.2	0.2
Heroin	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^a	0.1 ^b	0.1 ^b	0.1 ^a	0.1 ^b	0.1	0.2	0.2	0.2
Hallucinogens	0.1 ^a	0.1	0.1 ^b	0.1 ^a	0.0	0.0	0.1	0.1 ^a	0.1 ^a	0.1	0.0	0.0	0.0
Inhalants	0.0	0.0	0.0 ^a	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Nonmedical Use of Psychotherapeutics ^{2,3}	0.5 ^b	0.5 ^b	0.6 ^a	0.6 ^a	0.5 ^b	0.6	0.6	0.7	0.7	0.6	0.7	0.7	0.7
Pain Relievers	0.4 ^b	0.4 ^b	0.4 ^b	0.5 ^a	0.4 ^b	0.5	0.5	0.6	0.6	0.5	0.6	0.6	0.6
Tranquilizers	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Stimulants ²	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Sedatives	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
ALCOHOL	3.5 ^b	3.2	3.5 ^b	3.3	3.4 ^b	3.4 ^a	3.5 ^b	3.6 ^b	3.3	3.0	3.3	3.0	3.0
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.6 ^b	0.4	0.6 ^b	0.6 ^b	0.5 ^a	0.6 ^b	0.5 ^a	0.5 ^a	0.5	0.4	0.5	0.4	0.4
ILLCIT DRUGS OR ALCOHOL¹	4.9 ^a	4.6	4.9 ^a	4.7	4.9	4.8	5.0 ^a	4.9 ^a	4.7	4.3	4.8	4.5	4.5

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.55A Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	781 ^b	694 ^b	737 ^b	678 ^b	638 ^b	576 ^b	644 ^b	573 ^a	620 ^b	631 ^b	499	480	455
Marijuana and Hashish	608 ^b	502 ^b	508 ^b	506 ^b	448 ^b	383 ^a	454 ^b	444 ^b	462 ^b	470 ^b	396 ^a	383 ^a	304
Cocaine	49 ^b	52 ^b	72 ^b	57 ^b	49 ^b	62 ^b	40 ^b	23	20	24	11	11	11
Heroin	7	8	11	4	6	7	10	11	7	21	8	3	12
Hallucinogens	38 ^a	44 ^a	52 ^b	56 ^b	38 ^a	35	35	47 ^a	62 ^b	34	26	22	17
Inhalants	35 ^a	29	45 ^a	33	34 ^a	28	33	30	24	21	18	15	15
Nonmedical Use of Psychotherapeutics ^{2,3}	189	222 ^b	213 ^a	187	188	184	178	141	176	163	114	100 ^a	148
Pain Relievers	133	164 ^b	169 ^b	162 ^a	135	129	139	117	146	130	89	86	103
Tranquilizers	31	36	23	30	22	32	24	20	31	17	25	17	21
Stimulants ²	52	57	47	35	58	41	37	43	30	41	31	23	31
Sedatives	14	30	15	13	13	14	7	7	13	11	9	4	21
ALCOHOL	527 ^b	512 ^b	535 ^b	549 ^b	473 ^b	554 ^b	470 ^b	478 ^b	407 ^b	359 ^b	311 ^a	261	239
BOTH ILLICIT DRUGS AND ALCOHOL¹	210 ^b	168 ^b	197 ^b	192 ^b	153 ^b	165 ^b	147 ^b	140 ^b	138 ^b	117 ^a	89	91	74
ILLCIT DRUGS OR ALCOHOL¹	1,099 ^b	1,038 ^b	1,075 ^b	1,036 ^b	958 ^b	964 ^b	967 ^b	911 ^b	889 ^b	874 ^b	721	650	620

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.55B Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	3.2 ^b	2.8 ^b	2.9 ^b	2.7 ^b	2.5 ^b	2.3 ^a	2.6 ^b	2.3 ^b	2.5 ^b	2.5 ^b	2.0	1.9	1.8
Marijuana and Hashish	2.5 ^b	2.0 ^b	2.0 ^b	2.0 ^b	1.8 ^b	1.5 ^a	1.8 ^b	1.8 ^b	1.9 ^b	1.9 ^b	1.6 ^a	1.5 ^a	1.2
Cocaine	0.2 ^b	0.2 ^b	0.3 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.1	0.1	0.1	0.0	0.0	0.0
Heroin	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
Hallucinogens	0.2 ^a	0.2 ^a	0.2 ^b	0.2 ^b	0.1	0.1	0.1	0.2 ^a	0.3 ^b	0.1	0.1	0.1	0.1
Inhalants	0.1 ^a	0.1	0.2 ^a	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Nonmedical Use of Psychotherapeutics ^{2,3}	0.8	0.9 ^b	0.8 ^a	0.7	0.7	0.7	0.7	0.6	0.7	0.7	0.5	0.4 ^a	0.6
Pain Relievers	0.5	0.7 ^b	0.7 ^b	0.6 ^a	0.5	0.5	0.6	0.5	0.6 ^a	0.5	0.4	0.3	0.4
Tranquilizers	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Stimulants ²	0.2	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.1	0.2	0.1	0.1	0.1
Sedatives	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.1	0.0	0.0	0.0	0.1
ALCOHOL	2.1 ^b	2.0 ^b	2.1 ^b	2.2 ^b	1.9 ^b	2.2 ^b	1.9 ^b	1.9 ^b	1.7 ^b	1.4 ^b	1.2 ^a	1.0	1.0
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.8 ^b	0.7 ^b	0.8 ^b	0.8 ^b	0.6 ^b	0.7 ^b	0.6 ^b	0.6 ^b	0.6 ^b	0.5 ^a	0.4	0.4	0.3
ILLCIT DRUGS OR ALCOHOL¹	4.4 ^b	4.2 ^b	4.3 ^b	4.1 ^b	3.8 ^b	3.8 ^b	3.9 ^b	3.7 ^b	3.6 ^b	3.5 ^b	2.9	2.6	2.5

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.56A Substance Dependence for Specific Substances in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	1,711	1,654	1,786	1,899 ^a	1,784	1,786	1,874 ^a	1,827	1,845	1,818	1,922 ^a	1,808	1,661
Marijuana and Hashish	1,199	1,260	1,237	1,259	1,219	1,224	1,284	1,259	1,288	1,276	1,258	1,251	1,148
Cocaine	242 ^b	228 ^b	287 ^b	326 ^b	297 ^b	300 ^b	244 ^b	181 ^a	138	138	137	141	114
Heroin	42 ^b	37 ^b	53 ^b	82 ^b	56 ^b	61 ^b	87 ^a	86 ^a	110	131	156	155	152
Hallucinogens	88 ^a	61	91 ^a	80	69	43	64	83	78	65	47	50	50
Inhalants	6	8	18	7	14	6	8	23 ^a	15	3	6	16	5
Nonmedical Use of Psychotherapeutics ^{2,3}	358	310 ^b	445	476	436	473	525	548 ^a	571 ^a	534	635 ^b	513	437
Pain Relievers	257 ^a	210 ^b	314	353	334	393	420	455 ^a	465 ^a	472 ^a	508 ^b	385	346
Tranquilizers	51	47	74	90	65	40	59	65	76	57	91	83	61
Stimulants ²	90	107	126	125	109	79	94	110	107	67	112	131	106
Sedatives	17	13	16	9	17	14	19	27	18	15	14	22	17
ALCOHOL	2,171	2,139	2,313 ^b	2,350 ^b	2,498 ^b	2,380 ^b	2,401 ^b	2,228 ^a	2,224 ^a	2,057	2,081	1,922	1,989
BOTH ILLICIT DRUGS AND ALCOHOL¹	458	432	553 ^b	514 ^b	549 ^b	505 ^a	560 ^b	517 ^b	472	450	428	444	377
ILLCIT DRUGS OR ALCOHOL¹	3,424	3,362	3,546 ^a	3,735 ^b	3,733 ^b	3,661 ^b	3,715 ^b	3,538	3,598 ^a	3,425	3,574 ^a	3,285	3,274

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.56B Substance Dependence for Specific Substances in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	5.5 ^b	5.2	5.5 ^a	5.8 ^b	5.5 ^a	5.5 ^a	5.7 ^b	5.4 ^a	5.4 ^a	5.3	5.6 ^a	5.2	4.8
Marijuana and Hashish	3.9 ^a	4.0 ^b	3.8 ^a	3.9 ^a	3.7	3.7	3.9 ^a	3.7	3.8	3.7	3.6	3.6	3.3
Cocaine	0.8 ^b	0.7 ^b	0.9 ^b	1.0 ^b	0.9 ^b	0.9 ^b	0.7 ^b	0.5 ^a	0.4	0.4	0.4	0.4	0.3
Heroin	0.1 ^b	0.1 ^b	0.2 ^b	0.3 ^a	0.2 ^b	0.2 ^b	0.3 ^a	0.3 ^a	0.3	0.4	0.5	0.4	0.4
Hallucinogens	0.3 ^a	0.2	0.3 ^a	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.1	0.1
Inhalants	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.1 ^a	0.0	0.0	0.0	0.0	0.0
Nonmedical Use of Psychotherapeutics ^{2,3}	1.2	1.0	1.4	1.5	1.3	1.4	1.6 ^a	1.6 ^a	1.7 ^b	1.6	1.8 ^b	1.5	1.3
Pain Relievers	0.8	0.7 ^b	1.0	1.1	1.0	1.2	1.3 ^a	1.4 ^a	1.4 ^a	1.4 ^b	1.5 ^b	1.1	1.0
Tranquilizers	0.2	0.1	0.2	0.3	0.2	0.1	0.2	0.2	0.2	0.2	0.3	0.2	0.2
Stimulants ²	0.3	0.3	0.4	0.4	0.3	0.2	0.3	0.3	0.3	0.2	0.3	0.4	0.3
Sedatives	0.1	0.0	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.0	0.0	0.1	0.0
ALCOHOL	7.0 ^b	6.7 ^b	7.2 ^b	7.2 ^b	7.6 ^b	7.3 ^b	7.3 ^b	6.6 ^b	6.5 ^a	6.0	6.0	5.5	5.7
BOTH ILLICIT DRUGS AND ALCOHOL¹	1.5 ^b	1.4	1.7 ^b	1.6 ^b	1.7 ^b	1.5 ^b	1.7 ^b	1.5 ^b	1.4	1.3	1.2	1.3	1.1
ILLCIT DRUGS OR ALCOHOL¹	11.0 ^b	10.6 ^b	11.0 ^b	11.5 ^b	11.4 ^b	11.2 ^b	11.3 ^b	10.5 ^b	10.6 ^b	10.0	10.3 ^a	9.4	9.4

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.57A Substance Dependence for Specific Substances in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	2,143 ^b	2,025 ^b	2,329 ^a	2,141 ^b	2,418 ^a	2,335 ^a	2,361 ^a	2,489	2,383 ^a	2,117 ^b	2,635	2,629	2,875
Marijuana and Hashish	806 ^b	757 ^b	925 ^a	854 ^a	958	901 ^a	962	941	1,062	894 ^a	1,037	1,171	1,204
Cocaine	733 ^a	681 ^a	816 ^b	760 ^b	859 ^b	804 ^b	675 ^a	538	531	419	667	472	446
Heroin	149 ^b	124 ^b	162 ^b	113 ^b	236	111 ^b	157 ^b	227	204 ^a	217 ^a	274	276	374
Hallucinogens	11	29	37	*	12	29	27	55	18	44	51	30	27
Inhalants	*	*	27	*	*	10	43	11	26	22	*	*	24
Nonmedical Use of Psychotherapeutics ^{2,3}	695 ^b	743 ^b	736 ^b	703 ^b	687 ^b	899 ^a	823 ^b	1,041	1,011	928 ^a	1,180	1,145	1,261
Pain Relievers	546 ^b	569 ^b	488 ^b	586 ^b	604 ^b	748 ^a	681 ^b	835	789 ^a	768 ^a	972	1,017	1,058
Tranquilizers	169	113	155	123	92 ^a	128	150	176	206	103	198	137	202
Stimulants ²	117	96 ^a	186	113	108	184	103	146	114	145	235	213	199
Sedatives	100	69	58	32	23	65	66	81	99	42	87	35	78
ALCOHOL	5,524	4,912 ^b	5,508	5,075 ^a	5,452	5,534	5,955	6,238	5,708	5,340	6,235	5,815	5,835
BOTH ILLICIT DRUGS AND ALCOHOL¹	667	467	664	660	627	702	616	722	561	592	750	562	635
ILLCIT DRUGS OR ALCOHOL¹	7,000 ^b	6,470 ^b	7,173 ^a	6,555 ^b	7,242 ^a	7,167 ^a	7,700	8,005	7,531	6,865 ^b	8,120	7,882	8,074

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.57B Substance Dependence for Specific Substances in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	1.2	1.1 ^a	1.3	1.2 ^a	1.3	1.2	1.2	1.3	1.2	1.1 ^b	1.3	1.3	1.4
Marijuana and Hashish	0.4 ^a	0.4 ^a	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5 ^a	0.5	0.6	0.6
Cocaine	0.4 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.5 ^b	0.4 ^b	0.4 ^b	0.3	0.3	0.2	0.3 ^a	0.2	0.2
Heroin	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1	0.1 ^b	0.1 ^b	0.1	0.1 ^a	0.1 ^a	0.1	0.1	0.2
Hallucinogens	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Inhalants	*	*	0.0	*	*	0.0	0.0	0.0	0.0	0.0	*	*	0.0
Nonmedical Use of Psychotherapeutics ^{2,3}	0.4 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.4 ^b	0.5	0.4 ^b	0.5	0.5	0.5 ^a	0.6	0.6	0.6
Pain Relievers	0.3 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.3 ^b	0.4	0.4 ^b	0.4	0.4	0.4 ^a	0.5	0.5	0.5
Tranquilizers	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Stimulants ²	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Sedatives	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
ALCOHOL	3.1	2.7	3.0	2.7	2.9	2.9	3.1	3.2 ^a	2.9	2.7	3.1	2.9	2.8
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.4	0.3	0.4	0.4	0.3	0.4	0.3	0.4	0.3	0.3	0.4	0.3	0.3
ILLCIT DRUGS OR ALCOHOL¹	3.9	3.6	3.9	3.5 ^a	3.9	3.8	4.0	4.1	3.9	3.5 ^a	4.0	3.9	3.9

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.58A Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	2,013 ^a	1,802 ^b	2,192	2,172	2,460	2,182	2,093	2,306	2,201	2,017 ^a	2,235	2,214	2,440
Alcohol	2,405	2,359	2,658	2,843	2,764	2,744	2,911	3,081	2,868	2,644	2,611	2,731	2,683
Illicit Drugs or Alcohol ^{1,2}	3,483 ^b	3,327 ^b	3,791	3,930	4,031	3,927	4,061	4,276	4,152	3,764	4,004	4,051	4,149
Both Illicit Drugs and Alcohol ¹	1,319	1,255	1,467	1,522	1,571	1,416	1,324	1,535	1,313	1,211	1,213	1,320	1,427
Illicit Drugs but not Alcohol ¹	693 ^b	545 ^b	724 ^a	649 ^b	885	766 ^a	768 ^a	767 ^a	887	794	1,012	857	1,008
Alcohol but not Illicit Drugs ¹	1,086	1,103	1,191	1,318	1,192	1,326	1,583	1,545	1,553	1,432	1,377	1,404	1,256
SPECIALTY FACILITY													
Illicit Drugs ¹	1,412	1,103 ^b	1,427	1,280 ^a	1,578	1,362	1,211 ^b	1,484	1,518	1,354	1,532	1,483	1,604
Alcohol	1,549	1,298	1,535	1,626	1,553	1,575	1,564	1,709	1,591	1,472	1,492	1,421	1,572
Illicit Drugs or Alcohol ^{1,2}	2,346	1,874 ^b	2,327	2,308	2,533	2,430	2,293	2,627	2,594	2,325	2,496	2,466	2,606
Both Illicit Drugs and Alcohol ¹	709	595	718	748	734	624	580	750	629	574	633	547	684
Illicit Drugs but not Alcohol ¹	645 ^a	442 ^b	609 ^a	481 ^b	809	678	585 ^b	653	787	701	824	786	861
Alcohol but not Illicit Drugs ¹	783	646	727	818	725	898	890	843	891	828	801	840	764

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.58B Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	0.9	0.8 ^a	0.9	0.9	1.0	0.9	0.8	0.9	0.9	0.8	0.9	0.8	0.9
Alcohol	1.0	1.0	1.1	1.2	1.1	1.1	1.2	1.2 ^a	1.1	1.0	1.0	1.0	1.0
Illicit Drugs or Alcohol ^{1,2}	1.5	1.4	1.6	1.6	1.6	1.6	1.6	1.7	1.6	1.5	1.5	1.5	1.6
Both Illicit Drugs and Alcohol ¹	0.6	0.5	0.6	0.6	0.6	0.6	0.5	0.6	0.5	0.5	0.5	0.5	0.5
Illicit Drugs but not Alcohol ¹	0.3	0.2 ^b	0.3	0.3 ^b	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.4
Alcohol but not Illicit Drugs ¹	0.5	0.5	0.5	0.5	0.5	0.5	0.6 ^a	0.6 ^a	0.6 ^a	0.6	0.5	0.5	0.5
SPECIALTY FACILITY													
Illicit Drugs ¹	0.6	0.5 ^a	0.6	0.5	0.6	0.5	0.5 ^a	0.6	0.6	0.5	0.6	0.6	0.6
Alcohol	0.7	0.5	0.6	0.7	0.6	0.6	0.6	0.7	0.6	0.6	0.6	0.5	0.6
Illicit Drugs or Alcohol ^{1,2}	1.0	0.8 ^b	1.0	0.9	1.0	1.0	0.9	1.0	1.0	0.9	1.0	0.9	1.0
Both Illicit Drugs and Alcohol ¹	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.3
Illicit Drugs but not Alcohol ¹	0.3	0.2 ^b	0.3	0.2 ^b	0.3	0.3	0.2 ^a	0.3	0.3	0.3	0.3	0.3	0.3
Alcohol but not Illicit Drugs ¹	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.3	0.4	0.3	0.3	0.3	0.3

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.59A Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	255 ^b	242 ^b	276 ^b	252 ^b	258 ^b	230 ^a	220	209	205	228 ^a	223	175	166
Alcohol	227 ^b	216 ^b	236 ^b	215 ^b	197 ^b	192 ^b	185 ^b	179 ^b	147	144	164 ^a	137	107
Illicit Drugs or Alcohol ^{1,2}	369 ^b	362 ^b	406 ^b	348 ^b	353 ^b	341 ^b	331 ^b	286	290	301 ^a	285	232	227
Both Illicit Drugs and Alcohol ¹	172 ^b	161 ^b	180 ^b	167 ^b	152 ^b	153 ^b	145 ^a	141 ^a	112	113	135	113	91
Illicit Drugs but not Alcohol ¹	83	81	95	83	106	77	75	67	91	113	88	60	75
Alcohol but not Illicit Drugs ¹	55 ^b	55 ^b	55 ^b	47 ^b	45 ^b	37 ^b	40 ^a	37 ^a	35 ^a	30	29	24	16
SPECIALTY FACILITY													
Illicit Drugs ¹	142 ^b	113 ^a	134 ^b	142 ^b	137 ^b	111 ^a	112 ^a	117 ^a	99	125 ^a	121 ^a	90	74
Alcohol	121 ^b	95 ^a	126 ^b	119 ^b	101 ^a	83	78	97 ^a	70	63	76	73	55
Illicit Drugs or Alcohol ^{1,2}	186 ^b	168 ^a	185 ^b	181 ^b	181 ^b	151	145	152	139	146	157 ^a	122	109
Both Illicit Drugs and Alcohol ¹	81 ^b	55 ^b	81 ^b	84 ^b	63 ^b	53 ^a	54 ^b	68 ^b	39	43	52 ^a	49 ^a	25
Illicit Drugs but not Alcohol ¹	56	52	47	54	72	46	52	42	52	75	67	38	47
Alcohol but not Illicit Drugs ¹	36	30	37	27	28	22	17	20	19	16	19	13	18

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.59B Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	1.0 ^b	1.0 ^b	1.1 ^b	1.0 ^b	1.0 ^b	0.9 ^a	0.9	0.8	0.8	0.9 ^a	0.9	0.7	0.7
Alcohol	0.9 ^b	0.9 ^b	0.9 ^b	0.8 ^b	0.8 ^b	0.8 ^b	0.7 ^b	0.7 ^b	0.6	0.6	0.7 ^a	0.6	0.4
Illicit Drugs or Alcohol ^{1,2}	1.5 ^b	1.4 ^b	1.6 ^b	1.4 ^b	1.4 ^b	1.4 ^b	1.3 ^b	1.2	1.2 ^a	1.2 ^a	1.1	0.9	0.9
Both Illicit Drugs and Alcohol ¹	0.7 ^b	0.6 ^b	0.7 ^b	0.7 ^b	0.6 ^a	0.6 ^b	0.6 ^a	0.6 ^a	0.5	0.5	0.5	0.5	0.4
Illicit Drugs but not Alcohol ¹	0.3	0.3	0.4	0.3	0.4	0.3	0.3	0.3	0.4	0.5	0.4	0.2	0.3
Alcohol but not Illicit Drugs ¹	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.2 ^b	0.1 ^a	0.2 ^a	0.2 ^a	0.1 ^a	0.1	0.1	0.1	0.1
SPECIALTY FACILITY													
Illicit Drugs ¹	0.6 ^b	0.5 ^a	0.5 ^b	0.6 ^b	0.5 ^b	0.4	0.4 ^a	0.5 ^a	0.4	0.5 ^a	0.5 ^a	0.4	0.3
Alcohol	0.5 ^b	0.4 ^a	0.5 ^b	0.5 ^b	0.4 ^a	0.3	0.3	0.4 ^a	0.3	0.3	0.3	0.3	0.2
Illicit Drugs or Alcohol ^{1,2}	0.8 ^b	0.7 ^a	0.7 ^b	0.7 ^b	0.7 ^b	0.6	0.6	0.6 ^a	0.6	0.6	0.6 ^a	0.5	0.4
Both Illicit Drugs and Alcohol ¹	0.3 ^b	0.2 ^b	0.3 ^b	0.3 ^b	0.2 ^b	0.2 ^a	0.2 ^b	0.3 ^b	0.2	0.2	0.2 ^a	0.2 ^a	0.1
Illicit Drugs but not Alcohol ¹	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2
Alcohol but not Illicit Drugs ¹	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.60A Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	418	543	543	564	553	478	514	599	570	581	552	509	496
Alcohol	453	598 ^b	575 ^b	592 ^b	626 ^b	627 ^b	610 ^b	672 ^b	616 ^b	544 ^a	411	466	425
Illicit Drugs or Alcohol ^{1,2}	686	891	874	962 ^b	875	864	864	984 ^b	969 ^b	872	841	764	764
Both Illicit Drugs and Alcohol ¹	256	352 ^b	317	317	375 ^b	301	325 ^a	383 ^b	317	326	221	304	244
Illicit Drugs but not Alcohol ¹	161 ^b	190	226	247	178 ^a	177 ^a	188	215	253	249	331	205	252
Alcohol but not Illicit Drugs ¹	197	247 ^a	258 ^a	273 ^b	251 ^a	325 ^b	285 ^b	288 ^b	299 ^b	219	189	162	181
SPECIALTY FACILITY													
Illicit Drugs ¹	287	340	352	350	342	307	305	377	355	401	365	341	328
Alcohol	270	305	323 ^a	295	317	293	345 ^a	353 ^b	312	315	180	233	244
Illicit Drugs or Alcohol ^{1,2}	435	486	548	522	505	489	502	586 ^a	537	567	467	458	470
Both Illicit Drugs and Alcohol ¹	148	178	154	138	185	139	157	169	146	169	92	137	122
Illicit Drugs but not Alcohol ¹	133 ^a	147	182	194	148	156	137 ^a	180	202	221	259	179	205
Alcohol but not Illicit Drugs ¹	118	114	156	151	118	145	169 ^a	154	147	140	80	89	112

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.60B Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	1.3	1.7	1.7	1.7	1.7	1.5	1.6	1.8 ^a	1.7	1.7	1.6	1.5	1.4
Alcohol	1.5	1.9 ^b	1.8 ^b	1.8 ^b	1.9 ^b	1.9 ^b	1.9 ^b	2.0 ^b	1.8 ^b	1.6 ^a	1.2	1.3	1.2
Illicit Drugs or Alcohol ^{1,2}	2.2	2.8 ^b	2.7 ^a	3.0 ^b	2.7 ^a	2.6 ^a	2.6 ^a	2.9 ^b	2.8 ^b	2.5	2.4	2.2	2.2
Both Illicit Drugs and Alcohol ¹	0.8	1.1 ^b	1.0 ^a	1.0 ^a	1.1 ^b	0.9	1.0 ^a	1.1 ^b	0.9	0.9	0.6	0.9	0.7
Illicit Drugs but not Alcohol ¹	0.5 ^a	0.6	0.7	0.8	0.5	0.5	0.6	0.6	0.7	0.7	1.0	0.6	0.7
Alcohol but not Illicit Drugs ¹	0.6	0.8 ^b	0.8 ^b	0.8 ^b	0.8 ^a	1.0 ^b	0.9 ^b	0.9 ^b	0.9 ^b	0.6	0.5	0.5	0.5
SPECIALTY FACILITY													
Illicit Drugs ¹	0.9	1.1	1.1	1.1	1.0	0.9	0.9	1.1	1.0	1.2	1.1	1.0	0.9
Alcohol	0.9	1.0 ^a	1.0 ^b	0.9	1.0 ^a	0.9	1.0 ^b	1.1 ^b	0.9	0.9	0.5	0.7	0.7
Illicit Drugs or Alcohol ^{1,2}	1.4	1.5	1.7 ^a	1.6	1.5	1.5	1.5	1.7 ^a	1.6	1.7	1.3	1.3	1.3
Both Illicit Drugs and Alcohol ¹	0.5	0.6 ^a	0.5	0.4	0.6 ^a	0.4	0.5	0.5	0.4	0.5	0.3	0.4	0.3
Illicit Drugs but not Alcohol ¹	0.4	0.5	0.6	0.6	0.5	0.5	0.4	0.5	0.6	0.6	0.7	0.5	0.6
Alcohol but not Illicit Drugs ¹	0.4	0.4	0.5 ^a	0.5	0.4	0.4	0.5 ^a	0.5	0.4	0.4	0.2	0.3	0.3

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.61A Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	1,341 ^a	1,017 ^b	1,373 ^a	1,357 ^a	1,650	1,474	1,359 ^a	1,498	1,427	1,208 ^b	1,461	1,530	1,778
Alcohol	1,725 ^a	1,544 ^b	1,848	2,036	1,940	1,925	2,115	2,231	2,105	1,956	2,036	2,128	2,151
Illicit Drugs or Alcohol ^{1,2}	2,428 ^b	2,074 ^b	2,511 ^a	2,620 ^a	2,803	2,721	2,867	3,006	2,893	2,591 ^a	2,878	3,055	3,158
Both Illicit Drugs and Alcohol ¹	891	742 ^a	970	1,038	1,044	962	854	1,011	885	772 ^a	857	903	1,092
Illicit Drugs but not Alcohol ¹	450 ^a	275 ^b	404 ^b	319 ^b	602	513	505	484	542	432 ^a	593	592	681
Alcohol but not Illicit Drugs ¹	834	802	877	999	896	964	1,257	1,220	1,219	1,183	1,159	1,218	1,059
SPECIALTY FACILITY													
Illicit Drugs ¹	983	649 ^b	941	789 ^b	1,099	944	794 ^b	990	1,064	828 ^b	1,046	1,051	1,201
Alcohol	1,157	898 ^a	1,086	1,213	1,135	1,199	1,142	1,260	1,209	1,093	1,236	1,115	1,272
Illicit Drugs or Alcohol ^{1,2}	1,724	1,221 ^b	1,594 ^a	1,605 ^a	1,846	1,790	1,646	1,888	1,919	1,613 ^a	1,872	1,886	2,028
Both Illicit Drugs and Alcohol ¹	480	363	483	527	486	432	368	512	445	361	489	361	536
Illicit Drugs but not Alcohol ¹	456	243 ^b	380 ^a	233 ^b	589	476	395 ^a	431	533	405 ^a	497	569	609
Alcohol but not Illicit Drugs ¹	629	502	534	640	579	731	704	669	725	672	702	738	634

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.61B Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	0.7	0.6 ^b	0.8	0.7	0.9	0.8	0.7	0.8	0.7	0.6 ^b	0.7	0.8	0.9
Alcohol	1.0	0.9	1.0	1.1	1.0	1.0	1.1	1.2	1.1	1.0	1.0	1.0	1.0
Illicit Drugs or Alcohol ^{1,2}	1.4	1.1 ^b	1.4	1.4	1.5	1.4	1.5	1.6	1.5	1.3	1.4	1.5	1.5
Both Illicit Drugs and Alcohol ¹	0.5	0.4	0.5	0.6	0.6	0.5	0.4	0.5	0.5	0.4	0.4	0.4	0.5
Illicit Drugs but not Alcohol ¹	0.3	0.2 ^b	0.2 ^a	0.2 ^b	0.3	0.3	0.3	0.3	0.3	0.2 ^a	0.3	0.3	0.3
Alcohol but not Illicit Drugs ¹	0.5	0.4	0.5	0.5	0.5	0.5	0.7	0.6	0.6	0.6	0.6	0.6	0.5
SPECIALTY FACILITY													
Illicit Drugs ¹	0.5	0.4 ^b	0.5	0.4 ^a	0.6	0.5	0.4 ^a	0.5	0.5	0.4 ^a	0.5	0.5	0.6
Alcohol	0.6	0.5	0.6	0.7	0.6	0.6	0.6	0.7	0.6	0.6	0.6	0.5	0.6
Illicit Drugs or Alcohol ^{1,2}	1.0	0.7 ^b	0.9	0.9	1.0	0.9	0.9	1.0	1.0	0.8	0.9	0.9	1.0
Both Illicit Drugs and Alcohol ¹	0.3	0.2	0.3	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.3
Illicit Drugs but not Alcohol ¹	0.3	0.1 ^b	0.2	0.1 ^b	0.3	0.3	0.2	0.2	0.3	0.2 ^a	0.2	0.3	0.3
Alcohol but not Illicit Drugs ¹	0.4	0.3	0.3	0.3	0.3	0.4	0.4	0.3	0.4	0.3	0.4	0.4	0.3

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.62A Substances for Which Last or Current Treatment Was Received among Persons Aged 12 or Older Who Received Substance Use Treatment in the Past Year: Numbers in Thousands, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	3,483 ^a	3,327 ^b	3,791	3,930	4,031	3,927	4,061	4,276	4,152	3,764	4,004	4,051	4,149
Marijuana	974	975	1,017	1,062	1,233	942	955	1,234	1,028	872	957	845	1,034
Cocaine	796	557	884	797	930	818	664	777	706	511 ^a	658	584	781
Heroin	277 ^b	281 ^b	283 ^b	326 ^b	467	343 ^b	343 ^b	486	422	430	450	526	618
Hallucinogens	275	278	348	358	443	306	291	445	335	293	366	303	339
Inhalants	139	135	213	167	254	187	186	248	222	152	153	187	189
Pain Relievers	360 ^b	415 ^b	424 ^b	466 ^b	547 ^a	565 ^a	604	736	761	726	973	746	772
Tranquilizers	197 ^b	252 ^b	276 ^a	278 ^b	387	269 ^b	327	413	353	318 ^a	458	376	480
Stimulants ²	268 ^a	344	393	351	537	314	339	517	345	309	357	461	425
Sedatives	163	107	117	98	190	62 ^a	130	185	188	120	140	154	152
Alcohol	2,222	2,163	2,413	2,463	2,545	2,471	2,678	2,881 ^a	2,607	2,364	2,395	2,513	2,406

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.62B Substances for Which Last or Current Treatment Was Received among Persons Aged 12 or Older Who Received Substance Use Treatment in the Past Year: Percentages, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marijuana	28.0	29.3	26.8	27.0	30.6 ^a	24.0	23.5	28.9	24.8	23.2	23.9	20.9	24.9
Cocaine	22.8	16.7	23.3	20.3	23.1	20.8	16.3	18.2	17.0	13.6 ^a	16.4	14.4	18.8
Heroin	8.0 ^b	8.4 ^b	7.5 ^b	8.3 ^b	11.6	8.7 ^a	8.4 ^b	11.4	10.2 ^a	11.4	11.2	13.0	14.9
Hallucinogens	7.9	8.4	9.2	9.1	11.0	7.8	7.2	10.4	8.1	7.8	9.1	7.5	8.2
Inhalants	4.0	4.1	5.6	4.2	6.3	4.8	4.6	5.8	5.3	4.0	3.8	4.6	4.6
Pain Relievers	10.3 ^b	12.5 ^b	11.2 ^b	11.9 ^b	13.6 ^a	14.4	14.9	17.2	18.3	19.3	24.3 ^a	18.4	18.6
Tranquilizers	5.6 ^b	7.6 ^a	7.3 ^a	7.1 ^a	9.6	6.9 ^a	8.1	9.7	8.5	8.5	11.4	9.3	11.6
Stimulants ²	7.7	10.3	10.4	8.9	13.3	8.0	8.3	12.1	8.3	8.2	8.9	11.4	10.2
Sedatives	4.7	3.2	3.1	2.5	4.7	1.6 ^a	3.2	4.3	4.5	3.2	3.5	3.8	3.7
Alcohol	63.8	65.0 ^a	63.6	62.7	63.1	62.9	65.9 ^a	67.4 ^b	62.8	62.8	59.8	62.0	58.0

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.63A Substances for Which Last or Current Treatment Was Received among Persons Aged 12 to 17 Who Received Substance Use Treatment in the Past Year: Numbers in Thousands, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	369 ^b	362 ^b	406 ^b	348 ^b	353 ^b	341 ^b	331 ^b	286	290	301 ^a	285	232	227
Marijuana	203 ^b	189 ^b	219 ^b	185 ^b	189 ^b	172 ^a	162	159	155	165	187 ^b	127	117
Cocaine	45 ^b	33 ^b	41 ^b	32 ^b	48 ^b	34 ^b	23 ^a	28 ^a	27 ^a	18	24	20	8
Heroin	6	8	10	10	5	8	9	7	3	14	9	*	12
Hallucinogens	45 ^a	39 ^b	35 ^b	39 ^b	47 ^b	30 ^a	34 ^a	35 ^a	29	38 ^a	32	25	13
Inhalants	26	27 ^a	29 ^a	25 ^a	20	12	17	26	35 ^a	12	19	12	11
Pain Relievers	37	40	60 ^b	47 ^a	57 ^b	41	44 ^a	44 ^a	50 ^a	51 ^a	44 ^a	29	22
Tranquilizers	27	16	29	23	35	24	21	22	24	32	44	19	23
Stimulants ²	34	23	36	23	44 ^a	22	14	19	27	25	17	21	19
Sedatives	10	9	13	6	10	5	3	5	*	8	8	*	11
Alcohol	187 ^b	171 ^b	197 ^b	171 ^b	149 ^b	162 ^b	145 ^b	135 ^b	122 ^a	103	122 ^a	105	79

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.63B Substances for Which Last or Current Treatment Was Received among Persons Aged 12 to 17 Who Received Substance Use Treatment in the Past Year: Percentages, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marijuana	55.1	52.1	54.0	53.1	53.7	50.4	48.8	55.5	53.3	54.8	65.5 ^a	54.6	51.5
Cocaine	12.1 ^b	9.1 ^a	10.1 ^a	9.3 ^a	13.6 ^b	10.1 ^b	6.9	9.9 ^a	9.4 ^a	5.8	8.6	8.5	3.6
Heroin	1.6	2.1	2.5	3.0	1.3	2.5	2.8	2.3	1.1	4.7	3.0	*	5.3
Hallucinogens	12.3	10.7	8.6	11.1	13.2 ^a	8.8	10.2	12.3 ^a	9.9	12.8 ^a	11.1	10.6	5.8
Inhalants	7.1	7.5	7.0	7.2	5.7	3.6	5.3	9.0	12.2 ^a	3.9	6.7	5.2	4.7
Pain Relievers	10.1	11.1	14.7	13.6	16.2	12.1	13.4	15.5	17.1	17.0	15.6	12.5	9.9
Tranquilizers	7.2	4.3	7.2	6.6	9.8	7.2	6.2	7.9	8.4	10.7	15.4	8.3	10.3
Stimulants ²	9.2	6.4	8.8	6.6	12.5	6.6	4.1	6.6	9.4	8.3	5.9	9.0	8.5
Sedatives	2.7	2.6	3.3	1.9	2.8	1.6	0.8	1.6	*	2.7	3.0	*	4.7
Alcohol	50.8 ^b	47.3 ^a	48.5 ^a	49.1 ^a	42.2	47.6 ^a	43.8	47.2 ^a	42.1	34.3	42.9	45.0	34.9

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.64A Substances for Which Last or Current Treatment Was Received among Persons Aged 18 to 25 Who Received Substance Use Treatment in the Past Year: Numbers in Thousands, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	686	891	874	962 ^b	875	864	864	984 ^b	969 ^b	872	841	764	764
Marijuana	264	356 ^b	346 ^b	332 ^a	379 ^b	281	310	362 ^b	335 ^a	298	279	268	238
Cocaine	114	138	161 ^b	140 ^a	167 ^b	134	144 ^a	142 ^a	143 ^a	121	106	116	88
Heroin	39 ^b	50 ^b	60 ^b	87	67 ^b	74 ^a	67 ^b	95	86	112	104	128	133
Hallucinogens	77	106	116	90	114	99	97	125 ^a	83	90	83	90	72
Inhalants	30	39	62	48	61	60	61	65	30	42	38	46	35
Pain Relievers	75 ^b	136	140	137	159	131	171	207	182	228	247	190	183
Tranquilizers	59	78	77	79	107	77	79	99	78	110	109	104	98
Stimulants ²	60	130	116	98	122	103	78	104	76	85	113	94	97
Sedatives	18	22	19	22	28	21	29	29	18	14	13	25	21
Alcohol	387	524 ^b	493 ^a	531 ^b	556 ^b	570 ^b	553 ^b	610 ^b	537 ^b	472	358	398	377

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.64B Substances for Which Last or Current Treatment Was Received among Persons Aged 18 to 25 Who Received Substance Use Treatment in the Past Year: Percentages, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marijuana	38.5	39.9 ^a	39.6 ^a	34.5	43.3 ^b	32.5	35.9	36.7	34.6	34.1	33.1	35.1	31.2
Cocaine	16.7	15.5	18.4 ^a	14.6	19.1 ^a	15.4	16.7	14.4	14.8	13.9	12.6	15.2	11.5
Heroin	5.7 ^b	5.6 ^b	6.9 ^b	9.1 ^b	7.6 ^b	8.6 ^b	7.7 ^b	9.6 ^b	8.9 ^b	12.9	12.4	16.8	17.4
Hallucinogens	11.2	11.9	13.2	9.4	13.1	11.5	11.3	12.7	8.6	10.3	9.9	11.8	9.4
Inhalants	4.3	4.3	7.1	5.0	6.9	6.9	7.0	6.6	3.1	4.8	4.6	6.0	4.6
Pain Relievers	10.9 ^b	15.2 ^b	16.0 ^a	14.2 ^b	18.2	15.1 ^b	19.8	21.0	18.8	26.2	29.3	24.9	23.9
Tranquilizers	8.6	8.8	8.9	8.2	12.2	8.9	9.1	10.0	8.0	12.6	12.9	13.6	12.8
Stimulants ²	8.7	14.6	13.2	10.2	13.9	11.9	9.0	10.6	7.9	9.7	13.5	12.3	12.7
Sedatives	2.6	2.5	2.1	2.3	3.2	2.5	3.4	3.0	1.8	1.6	1.5	3.3	2.7
Alcohol	56.4	58.8 ^a	56.4	55.2	63.5 ^b	66.0 ^b	64.1 ^b	62.0 ^b	55.4	54.2	42.6	52.1	49.3

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.65A Substances for Which Last or Current Treatment Was Received among Persons Aged 26 or Older Who Received Substance Use Treatment in the Past Year: Numbers in Thousands, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	2,428 ^b	2,074 ^b	2,511 ^a	2,620 ^a	2,803	2,721	2,867	3,006	2,893	2,591 ^a	2,878	3,055	3,158
Marijuana	507	431 ^a	451 ^a	546	665	490	483	714	539	409 ^b	491	450 ^a	679
Cocaine	637	386 ^b	683	624	715	650	497	607	535	372 ^b	528	448 ^a	685
Heroin	232 ^b	224 ^b	213 ^b	228 ^b	396	261 ^a	267 ^a	385	333	303	338	392	473
Hallucinogens	152	134 ^a	197	229	282	176	160	285	223	165	252	189	254
Inhalants	83	69	122	94	173	115	108	157	156	98	95	129	143
Pain Relievers	247 ^b	239 ^b	224 ^b	282 ^b	330 ^b	393	388 ^a	485	529	447	682	527	567
Tranquilizers	111 ^b	158 ^b	170 ^a	176 ^a	245	168 ^b	228	292	251	176 ^a	305	253	359
Stimulants ²	174 ^a	190	241	230	370	189	247	394	242	199	227	346	308
Sedatives	136	76	85	70	152	35 ^a	99	151	164	98	118	123	121
Alcohol	1,647	1,469 ^a	1,723	1,761	1,840	1,739	1,980	2,136	1,948	1,789	1,915	2,010	1,950

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.65B Substances for Which Last or Current Treatment Was Received among Persons Aged 26 or Older Who Received Substance Use Treatment in the Past Year: Percentages, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marijuana	20.9	20.8	18.0	20.8	23.7	18.0	16.9	23.7	18.6	15.8	17.1	14.7 ^a	21.5
Cocaine	26.2	18.6	27.2	23.8	25.5	23.9	17.3	20.2	18.5	14.4 ^a	18.3	14.7 ^a	21.7
Heroin	9.6	10.8	8.5 ^a	8.7 ^a	14.1	9.6	9.3 ^a	12.8	11.5	11.7	11.7	12.8	15.0
Hallucinogens	6.3	6.4	7.9	8.7	10.1	6.5	5.6	9.5	7.7	6.4	8.7	6.2	8.0
Inhalants	3.4	3.3	4.8	3.6	6.2	4.2	3.8	5.2	5.4	3.8	3.3	4.2	4.5
Pain Relievers	10.2 ^b	11.5 ^a	8.9 ^b	10.8 ^b	11.8 ^a	14.4	13.5	16.1	18.3	17.2	23.7	17.2	18.0
Tranquilizers	4.6 ^b	7.6	6.8	6.7	8.8	6.2 ^a	8.0	9.7	8.7	6.8	10.6	8.3	11.4
Stimulants ²	7.2	9.2	9.6	8.8	13.2	6.9	8.6	13.1	8.3	7.7	7.9	11.3	9.8
Sedatives	5.6	3.6	3.4	2.7	5.4	1.3 ^a	3.5	5.0	5.7	3.8	4.1	4.0	3.8
Alcohol	67.8	70.8 ^a	68.6	67.2	65.6	63.9	69.1	71.1 ^a	67.3	69.0	66.5	65.8	61.8

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.66A Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Numbers in Thousands, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	3,483 ^a	3,327 ^b	3,791	3,930	4,031	3,927	4,061	4,276	4,152	3,764	4,004	4,051	4,149
Hospital - Inpatient	859	587 ^b	775	773	815	784	681 ^a	855	737	871	861	879	921
Rehabilitation Facility - Inpatient	1,094	752 ^b	947	1,109	934	1,055	744 ^b	1,194	992	1,001	1,010	1,042	1,076
Rehabilitation Facility - Outpatient	1,536	1,243 ^b	1,666	1,546	1,616	1,723	1,460	1,993	1,702	1,531	1,505	1,753	1,731
Mental Health Center - Outpatient	1,016	729 ^b	982	1,046	1,107	890 ^a	1,055	1,092	1,003	1,031	1,000	1,176	1,157
Emergency Room	469	251 ^b	453	399	395	525	374	492	472	574	597	603	521
Private Doctor's Office	523 ^a	377 ^b	490 ^b	460 ^b	614	599	688	721	655	700	735	770	780
Self-Help Group	2,021	1,911	2,098	2,102	2,183	2,188	2,199	2,434	2,343	2,139	2,119	2,292	2,210
Prison/Jail	259	206 ^a	310	344	421	302	344	373	344	435	388	263	366

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.66B Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Percentages, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Hospital - Inpatient	24.7	17.7	20.4	19.7	20.2	20.0	16.8 ^a	20.0	17.8	23.2	21.5	21.7	22.2
Rehabilitation Facility - Inpatient	31.4	22.6	25.0	28.2	23.2	26.9	18.3 ^b	27.9	23.9	26.6	25.2	25.7	25.9
Rehabilitation Facility - Outpatient	44.1	37.3	43.9	39.3	40.1	43.9	35.9	46.6	41.0	40.7	37.6	43.3	41.7
Mental Health Center - Outpatient	29.2	21.9 ^a	25.9	26.6	27.5	22.7	26.0	25.5	24.2	27.4	25.0	29.0	27.9
Emergency Room	13.5	7.5 ^b	11.9	10.2	9.8	13.4	9.2	11.5	11.4	15.2	14.9	14.9	12.6
Private Doctor's Office	15.0	11.3 ^b	12.9 ^b	11.7 ^b	15.2	15.3	16.9	16.9	15.8	18.6	18.4	19.0	18.8
Self-Help Group	58.0	57.4	55.3	53.5	54.2	55.7	54.1	56.9	56.4	56.8	52.9	56.6	53.3
Prison/Jail	7.4	6.2	8.2	8.8	10.5	7.7	8.5	8.7	8.3	11.6	9.7	6.5	8.8

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.67A Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 to 17 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Numbers in Thousands, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	369 ^b	362 ^b	406 ^b	348 ^b	353 ^b	341 ^b	331 ^b	286	290	301 ^a	285	232	227
Hospital - Inpatient	72	57	63	73	66	60	54	52	46	47	37	43	51
Rehabilitation Facility - Inpatient	83	65	82 ^a	71	54	70	59	70	46	66	59	49	47
Rehabilitation Facility - Outpatient	127 ^b	111 ^b	122 ^b	115 ^b	106 ^a	111 ^b	88	93 ^a	68	71	119 ^b	89	59
Mental Health Center - Outpatient	79	73	85 ^a	73	98 ^a	57	59	53	55	70	53	62	51
Emergency Room	28	50	44	46	53	45	34	32	33	38	28	40	45
Private Doctor's Office	49	56	67	47	53	51	30	27	37	35	34	30	45
Self-Help Group	173 ^b	132 ^b	138 ^b	121 ^b	123 ^a	151 ^b	87	90	95	105	102	80	71
Prison/Jail	39	30	35	21	36	22	24	25	28	21	26	22	21

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.67B Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 to 17 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Percentages, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Hospital - Inpatient	19.4	15.9	15.4	21.0	18.6	17.7	16.4	18.2	15.7	15.5	13.1	18.3	22.4
Rehabilitation Facility - Inpatient	22.4	18.0	20.2	20.3	15.4	20.6	17.8	24.4	15.9	22.0	20.8	21.1	20.5
Rehabilitation Facility - Outpatient	34.5	30.7	30.0	33.2	30.1	32.7	26.7	32.4	23.6	23.7	41.6 ^a	38.4 ^a	26.0
Mental Health Center - Outpatient	21.5	20.3	21.0	21.1	27.7	16.7	18.0	18.6	19.0	23.3	18.7	26.7	22.5
Emergency Room	7.7 ^b	13.9	10.7	13.1	14.9	13.2	10.2 ^a	11.0	11.2	12.8	9.7 ^a	17.3	20.0
Private Doctor's Office	13.2	15.6	16.5	13.6	15.2	15.0	8.9 ^a	9.4 ^a	12.8	11.6	11.9	13.0	19.8
Self-Help Group	47.0 ^b	36.4	33.9	34.8	34.9	44.4 ^a	26.3	31.4	32.9	34.8	35.7	34.6	31.1
Prison/Jail	10.5	8.4	8.5	6.0	10.1	6.5	7.1	8.8	9.8	7.1	9.2	9.3	9.3

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.68A Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 18 to 25 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Numbers in Thousands, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	686	891	874	962 ^b	875	864	864	984 ^b	969 ^b	872	841	764	764
Hospital - Inpatient	117	123	130	149	157	138	142	174	122	192	146	148	154
Rehabilitation Facility - Inpatient	160	185	193	174	193	190	166	220	208	234	201	202	212
Rehabilitation Facility - Outpatient	269	316	381	359	356	353	375	447 ^a	378	395	310	349	326
Mental Health Center - Outpatient	169	159	196	224	216	162	195	240	196	236	171	181	206
Emergency Room	61	67	79	76	99	94	89	114	75	121	102	91	85
Private Doctor's Office	80	81	90	116	148	135	152	147	120	158	154	126	109
Self-Help Group	351	471	459	449	516 ^a	401	449	538 ^a	533 ^a	471	421	369	402
Prison/Jail	57	62	70	107 ^b	102 ^a	107 ^b	61	104 ^b	99 ^a	127 ^b	54	57	48

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.68B Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 18 to 25 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Percentages, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Hospital - Inpatient	17.0	13.8 ^a	14.9	15.5	17.9	16.0	16.5	17.7	12.6 ^a	22.0	17.3	19.4	20.2
Rehabilitation Facility - Inpatient	23.3	20.8	22.1	18.1 ^b	22.0	22.0	19.2 ^a	22.4	21.5	26.9	23.9	26.4	27.8
Rehabilitation Facility - Outpatient	39.2	35.5	43.5	37.3	40.6	40.8	43.4	45.4	39.0	45.3	36.9	45.7	42.7
Mental Health Center - Outpatient	24.7	17.9 ^b	22.4	23.3	24.7	18.8 ^a	22.6	24.4	20.2 ^a	27.1	20.3	23.7	27.0
Emergency Room	8.9	7.5	9.0	7.9	11.3	10.9	10.4	11.6	7.8	13.9	12.1	11.9	11.1
Private Doctor's Office	11.7	9.1 ^a	10.3	12.0	16.9	15.6	17.6	14.9	12.4	18.1	18.3	16.5	14.3
Self-Help Group	51.1	52.8	52.5	46.7	59.0	46.4	52.0	54.7	55.1	54.0	50.1	48.3	52.5
Prison/Jail	8.3	6.9	8.0	11.1 ^a	11.7 ^a	12.3 ^a	7.1	10.5	10.3	14.6 ^b	6.5	7.5	6.3

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.69A Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 26 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Numbers in Thousands, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	2,428 ^b	2,074 ^b	2,511 ^a	2,620 ^a	2,803	2,721	2,867	3,006	2,893	2,591 ^a	2,878	3,055	3,158
Hospital - Inpatient	671	407 ^b	582	550	593	586	484 ^a	629	569	632	678	688	716
Rehabilitation Facility - Inpatient	852	502 ^b	672	864	687	794	519 ^b	904	737	700	750	791	817
Rehabilitation Facility - Outpatient	1,140	815 ^b	1,163	1,072	1,154	1,258	996 ^a	1,453	1,256	1,065	1,076	1,314	1,346
Mental Health Center - Outpatient	768	496 ^b	701	748	793	671	801	800	751	725	776	933	900
Emergency Room	380	134 ^b	330	277	244	386	251	347	364	414	467	472	391
Private Doctor's Office	394 ^a	240 ^b	333 ^b	297 ^b	412 ^a	413 ^a	506	547	498	507	548	613	626
Self-Help Group	1,497	1,309 ^a	1,501	1,532	1,544	1,635	1,663	1,806	1,715	1,563	1,596	1,843	1,738
Prison/Jail	163 ^a	114 ^b	206	216	284	173	259	245	216	286	308	184	296

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.69B Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 26 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Percentages, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Hospital - Inpatient	27.6	19.6	23.2	21.0	21.2	21.5	16.9	20.9	19.7	24.4	23.6	22.5	22.7
Rehabilitation Facility - Inpatient	35.1 ^a	24.2	26.8	33.0	24.5	29.2	18.1 ^a	30.1	25.5	27.0	26.1	25.9	25.9
Rehabilitation Facility - Outpatient	46.9	39.3	46.3	40.9	41.2	46.2	34.7 ^a	48.3	43.4	41.1	37.4	43.0	42.6
Mental Health Center - Outpatient	31.6	23.9	27.9	28.5	28.3	24.6	27.9	26.6	26.0	28.0	27.0	30.6	28.5
Emergency Room	15.6	6.5 ^b	13.2	10.6	8.7	14.2	8.8	11.5	12.6	16.0	16.2	15.4	12.4
Private Doctor's Office	16.2	11.6 ^b	13.3 ^a	11.3 ^b	14.7	15.2	17.7	18.2	17.2	19.6	19.0	20.1	19.8
Self-Help Group	61.7	63.1	59.8	58.4	55.1	60.1	58.0	60.1	59.3	60.3	55.4	60.3	55.0
Prison/Jail	6.7	5.5	8.2	8.3	10.1	6.3	9.0	8.1	7.5	11.1	10.7	6.0	9.4

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.70A Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	7,748	7,333	8,053	7,550	7,762	7,558	7,580	7,863	7,936	7,186 ^a	8,040	7,608	7,881
Received Treatment at a Specialty Facility	1,412	1,103 ^b	1,427	1,280 ^a	1,578	1,362	1,211 ^b	1,484	1,518	1,354	1,532	1,483	1,604
Did Not Receive Treatment at a Specialty Facility	6,335	6,230	6,626	6,269	6,184	6,196	6,369	6,379	6,418	5,832	6,508	6,126	6,278
NEEDED TREATMENT FOR ALCOHOL	18,638	18,215	19,360 ^b	19,378 ^b	19,568 ^b	19,350 ^b	19,102 ^b	19,420 ^b	18,633	17,389	18,291	18,049	17,645
Received Treatment at a Specialty Facility	1,549	1,298	1,535	1,626	1,553	1,575	1,564	1,709	1,591	1,472	1,492	1,421	1,572
Did Not Receive Treatment at a Specialty Facility	17,089	16,917	17,824 ^b	17,752 ^b	18,015 ^b	17,775 ^b	17,537 ^b	17,711 ^b	17,042	15,917	16,800	16,628	16,073
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	22,811	22,165	23,476	23,172	23,635	23,265	23,208	23,632	23,209	21,579	23,061	22,687	22,478
Received Treatment at a Specialty Facility ²	2,346	1,874 ^b	2,327	2,308	2,533	2,430	2,293	2,627	2,594	2,325	2,496	2,466	2,606
Did Not Receive Treatment at a Specialty Facility	20,465	20,290	21,149 ^a	20,864	21,102 ^a	20,835	20,914	21,005	20,615	19,254	20,565	20,221	19,872

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.70B Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	3.3 ^a	3.1	3.3 ^b	3.1	3.2	3.0	3.0	3.1	3.1	2.8	3.1	2.9	3.0
Received Treatment at a Specialty Facility	0.6	0.5 ^a	0.6	0.5	0.6	0.5	0.5 ^a	0.6	0.6	0.5	0.6	0.6	0.6
Did Not Receive Treatment at a Specialty Facility	2.7 ^b	2.6 ^a	2.8 ^b	2.6	2.5	2.5	2.5	2.5	2.5	2.3	2.5	2.3	2.4
NEEDED TREATMENT FOR ALCOHOL	7.9 ^b	7.7 ^b	8.0 ^b	8.0 ^b	8.0 ^b	7.8 ^b	7.6 ^b	7.7 ^b	7.3 ^b	6.8	7.0	6.9	6.7
Received Treatment at a Specialty Facility	0.7	0.5	0.6	0.7	0.6	0.6	0.6	0.7	0.6	0.6	0.6	0.5	0.6
Did Not Receive Treatment at a Specialty Facility	7.3 ^b	7.1 ^b	7.4 ^b	7.3 ^b	7.3 ^b	7.2 ^b	7.0 ^b	7.0 ^b	6.7 ^b	6.2	6.5	6.3	6.1
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	9.7 ^b	9.3 ^b	9.8 ^b	9.5 ^b	9.6 ^b	9.4 ^b	9.3 ^b	9.4 ^b	9.2 ^b	8.4	8.9	8.6	8.5
Received Treatment at a Specialty Facility ²	1.0	0.8 ^b	1.0	0.9	1.0	1.0	0.9	1.0	1.0	0.9	1.0	0.9	1.0
Did Not Receive Treatment at a Specialty Facility	8.7 ^b	8.5 ^b	8.8 ^b	8.6 ^b	8.6 ^b	8.4 ^b	8.4 ^b	8.3 ^b	8.1 ^b	7.5	7.9	7.7	7.5

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.71A Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	1,414 ^b	1,327 ^b	1,397 ^b	1,254 ^b	1,218 ^b	1,124 ^b	1,197 ^b	1,100 ^b	1,179 ^b	1,184 ^b	1,041 ^a	908	894
Received Treatment at a Specialty Facility	142 ^b	113 ^a	134 ^b	142 ^b	137 ^b	111 ^a	112 ^a	117 ^a	99	125 ^a	121 ^a	90	74
Did Not Receive Treatment at a Specialty Facility	1,272 ^b	1,214 ^b	1,262 ^b	1,112 ^b	1,081 ^b	1,013 ^b	1,085 ^b	983 ^b	1,080 ^b	1,060 ^b	920	817	820
NEEDED TREATMENT FOR ALCOHOL	1,495 ^b	1,501 ^b	1,570 ^b	1,460 ^b	1,409 ^b	1,400 ^b	1,255 ^b	1,182 ^b	1,144 ^b	978 ^b	889 ^b	735	699
Received Treatment at a Specialty Facility	121 ^b	95 ^a	126 ^b	119 ^b	101 ^a	83	78	97 ^a	70	63	76	73	55
Did Not Receive Treatment at a Specialty Facility	1,374 ^b	1,406 ^b	1,444 ^b	1,341 ^b	1,308 ^b	1,318 ^b	1,177 ^b	1,086 ^b	1,074 ^b	915 ^b	814 ^b	662	644
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	2,256 ^b	2,253 ^b	2,288 ^b	2,096 ^b	2,090 ^b	1,984 ^b	1,949 ^b	1,784 ^b	1,833 ^b	1,744 ^b	1,563 ^b	1,340	1,284
Received Treatment at a Specialty Facility ²	186 ^b	168 ^a	185 ^b	181 ^b	181 ^b	151	145	152	139	146	157 ^a	122	109
Did Not Receive Treatment at a Specialty Facility	2,071 ^b	2,085 ^b	2,103 ^b	1,915 ^b	1,908 ^b	1,833 ^b	1,804 ^b	1,632 ^b	1,694 ^b	1,598 ^b	1,407 ^b	1,218	1,175

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.71B Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	5.7 ^b	5.3 ^b	5.5 ^b	4.9 ^b	4.8 ^b	4.5 ^b	4.8 ^b	4.5 ^b	4.8 ^b	4.7 ^b	4.2 ^a	3.6	3.6
Received Treatment at a Specialty Facility	0.6 ^b	0.5 ^a	0.5 ^b	0.6 ^b	0.5 ^b	0.4	0.4 ^a	0.5 ^a	0.4	0.5 ^a	0.5 ^a	0.4	0.3
Did Not Receive Treatment at a Specialty Facility	5.1 ^b	4.9 ^b	5.0 ^b	4.4 ^b	4.3 ^b	4.0 ^b	4.4 ^b	4.0 ^b	4.4 ^b	4.2 ^b	3.7	3.3	3.3
NEEDED TREATMENT FOR ALCOHOL	6.0 ^b	6.0 ^b	6.2 ^b	5.8 ^b	5.5 ^b	5.5 ^b	5.0 ^b	4.8 ^b	4.7 ^b	3.9 ^b	3.6 ^b	3.0	2.8
Received Treatment at a Specialty Facility	0.5 ^b	0.4 ^a	0.5 ^b	0.5 ^b	0.4 ^a	0.3	0.3	0.4 ^a	0.3	0.3	0.3	0.3	0.2
Did Not Receive Treatment at a Specialty Facility	5.6 ^b	5.6 ^b	5.7 ^b	5.3 ^b	5.2 ^b	5.2 ^b	4.7 ^b	4.4 ^b	4.4 ^b	3.7 ^b	3.3 ^b	2.7	2.6
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	9.1 ^b	9.0 ^b	9.1 ^b	8.3 ^b	8.2 ^b	7.9 ^b	7.8 ^b	7.3 ^b	7.5 ^b	7.0 ^b	6.3 ^b	5.4	5.2
Received Treatment at a Specialty Facility ²	0.8 ^b	0.7 ^a	0.7 ^b	0.7 ^b	0.7 ^b	0.6	0.6	0.6 ^a	0.6	0.6	0.6 ^a	0.5	0.4
Did Not Receive Treatment at a Specialty Facility	8.4 ^b	8.3 ^b	8.3 ^b	7.6 ^b	7.5 ^b	7.3 ^b	7.2 ^b	6.6 ^b	7.0 ^b	6.4 ^b	5.6 ^b	4.9	4.7

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.72A Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	2,680	2,624	2,856 ^b	2,854 ^b	2,734 ^a	2,754 ^a	2,697	2,747 ^a	2,815 ^b	2,761 ^a	2,849 ^b	2,671	2,460
Received Treatment at a Specialty Facility	287	340	352	350	342	307	305	377	355	401	365	341	328
Did Not Receive Treatment at a Specialty Facility	2,393 ^a	2,284	2,505 ^b	2,504 ^b	2,392 ^a	2,447 ^b	2,392 ^a	2,370 ^a	2,460 ^b	2,360	2,483 ^b	2,330	2,132
NEEDED TREATMENT FOR ALCOHOL	5,584 ^b	5,589 ^b	5,744 ^b	5,816 ^b	5,909 ^b	5,641 ^b	5,847 ^b	5,538 ^b	5,455 ^b	5,062 ^b	5,038 ^b	4,633	4,361
Received Treatment at a Specialty Facility	270	305	323 ^a	295	317	293	345 ^a	353 ^b	312	315	180	233	244
Did Not Receive Treatment at a Specialty Facility	5,313 ^b	5,284 ^b	5,421 ^b	5,522 ^b	5,592 ^b	5,348 ^b	5,502 ^b	5,184 ^b	5,143 ^b	4,747 ^b	4,858 ^b	4,400	4,117
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	6,874 ^b	6,824 ^b	7,047 ^b	7,220 ^b	7,155 ^b	6,939 ^b	7,054 ^b	6,929 ^b	6,956 ^b	6,560 ^b	6,681 ^b	6,146	5,845
Received Treatment at a Specialty Facility ²	435	486	548	522	505	489	502	586 ^a	537	567	467	458	470
Did Not Receive Treatment at a Specialty Facility	6,439 ^b	6,338 ^b	6,499 ^b	6,699 ^b	6,650 ^b	6,450 ^b	6,552 ^b	6,343 ^b	6,419 ^b	5,994 ^b	6,215 ^b	5,689	5,375

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.72B Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	8.6 ^b	8.3 ^b	8.9 ^b	8.8 ^b	8.4 ^b	8.4 ^b	8.2 ^b	8.2 ^b	8.3 ^b	8.0 ^b	8.2 ^b	7.7	7.0
Received Treatment at a Specialty Facility	0.9	1.1	1.1	1.1	1.0	0.9	0.9	1.1	1.0	1.2	1.1	1.0	0.9
Did Not Receive Treatment at a Specialty Facility	7.7 ^b	7.2 ^b	7.8 ^b	7.7 ^b	7.3 ^b	7.5 ^b	7.3 ^b	7.1 ^b	7.2 ^b	6.9 ^a	7.2 ^b	6.7	6.1
NEEDED TREATMENT FOR ALCOHOL	18.0 ^b	17.6 ^b	17.8 ^b	17.9 ^b	18.0 ^b	17.2 ^b	17.8 ^b	16.5 ^b	16.0 ^b	14.8 ^b	14.6 ^b	13.3	12.5
Received Treatment at a Specialty Facility	0.9	1.0 ^a	1.0 ^b	0.9	1.0 ^a	0.9	1.0 ^b	1.1 ^b	0.9	0.9	0.5	0.7	0.7
Did Not Receive Treatment at a Specialty Facility	17.1 ^b	16.7 ^b	16.8 ^b	17.0 ^b	17.1 ^b	16.3 ^b	16.7 ^b	15.4 ^b	15.1 ^b	13.8 ^b	14.0 ^b	12.6	11.8
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	22.2 ^b	21.5 ^b	21.9 ^b	22.2 ^b	21.9 ^b	21.2 ^b	21.4 ^b	20.6 ^b	20.4 ^b	19.1 ^b	19.3 ^b	17.7	16.7
Received Treatment at a Specialty Facility ²	1.4	1.5	1.7 ^a	1.6	1.5	1.5	1.5	1.7 ^a	1.6	1.7	1.3	1.3	1.3
Did Not Receive Treatment at a Specialty Facility	20.8 ^b	20.0 ^b	20.2 ^b	20.6 ^b	20.3 ^b	19.7 ^b	19.9 ^b	18.9 ^b	18.8 ^b	17.5 ^b	18.0 ^b	16.4	15.4

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.73A Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	3,654 ^b	3,382 ^b	3,801 ^a	3,442 ^b	3,810 ^a	3,680 ^b	3,686 ^b	4,016	3,943	3,240 ^b	4,150	4,030	4,527
Received Treatment at a Specialty Facility	983	649 ^b	941	789 ^b	1,099	944	794 ^b	990	1,064	828 ^b	1,046	1,051	1,201
Did Not Receive Treatment at a Specialty Facility	2,670 ^b	2,732 ^a	2,860	2,653 ^b	2,711 ^a	2,735 ^a	2,892	3,026	2,878	2,412 ^b	3,104	2,978	3,326
NEEDED TREATMENT FOR ALCOHOL	11,559 ^a	11,125 ^b	12,045	12,102	12,251	12,309	12,000	12,701	12,034	11,349 ^a	12,364	12,681	12,584
Received Treatment at a Specialty Facility	1,157	898 ^a	1,086	1,213	1,135	1,199	1,142	1,260	1,209	1,093	1,236	1,115	1,272
Did Not Receive Treatment at a Specialty Facility	10,402	10,228 ^a	10,960	10,889	11,116	11,110	10,858	11,441	10,825	10,255 ^a	11,128	11,567	11,312
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	13,680 ^b	13,088 ^b	14,142 ^a	13,856 ^b	14,390	14,342	14,204 ^a	14,918	14,420	13,274 ^b	14,817	15,200	15,349
Received Treatment at a Specialty Facility ²	1,724	1,221 ^b	1,594 ^a	1,605 ^a	1,846	1,790	1,646	1,888	1,919	1,613 ^a	1,872	1,886	2,028
Did Not Receive Treatment at a Specialty Facility	11,956 ^b	11,867 ^b	12,547	12,251 ^a	12,543	12,552	12,558	13,030	12,502	11,661 ^b	12,944	13,314	13,321

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.73B Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	2.0	1.9 ^a	2.1	1.9 ^a	2.0	1.9	1.9 ^a	2.1	2.0	1.6 ^b	2.1	2.0	2.2
Received Treatment at a Specialty Facility	0.5	0.4 ^b	0.5	0.4 ^a	0.6	0.5	0.4 ^a	0.5	0.5	0.4 ^a	0.5	0.5	0.6
Did Not Receive Treatment at a Specialty Facility	1.5	1.5	1.6	1.4	1.4	1.4	1.5	1.6	1.5	1.2 ^b	1.5	1.5	1.6
NEEDED TREATMENT FOR ALCOHOL	6.4	6.1	6.6	6.5	6.5	6.5	6.3	6.6	6.2	5.7	6.2	6.3	6.1
Received Treatment at a Specialty Facility	0.6	0.5	0.6	0.7	0.6	0.6	0.6	0.7	0.6	0.6	0.6	0.5	0.6
Did Not Receive Treatment at a Specialty Facility	5.8	5.7	6.0	5.9	5.9	5.9	5.7	5.9	5.5	5.2	5.5	5.7	5.5
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	7.6	7.2	7.7	7.5	7.7	7.6	7.4	7.7	7.4	6.7 ^b	7.4	7.5	7.5
Received Treatment at a Specialty Facility ²	1.0	0.7 ^b	0.9	0.9	1.0	0.9	0.9	1.0	1.0	0.8	0.9	0.9	1.0
Did Not Receive Treatment at a Specialty Facility	6.7	6.6	6.9	6.6	6.7	6.6	6.5	6.7	6.4	5.9 ^a	6.5	6.6	6.5

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.74A Type of Substance Treatment Received at Any Location in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use in the Past Year, by Age Group: Numbers in Thousands, 2002-2014

Substance for Which Treatment Was Received/Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS BUT NOT ALCOHOL													
12-17	83	81	95	83	106	77	75	67	91	113	88	60	75
18-25	161 ^b	190	226	247	178 ^a	177 ^a	188	215	253	249	331	205	252
26 or Older	450 ^a	275 ^b	404 ^b	319 ^b	602	513	505	484	542	432 ^a	593	592	681
ALCOHOL BUT NOT ILLICIT DRUGS													
12-17	55 ^b	55 ^b	55 ^b	47 ^b	45 ^b	37 ^b	40 ^a	37 ^a	35 ^a	30	29	24	16
18-25	197	247 ^a	258 ^a	273 ^b	251 ^a	325 ^b	285 ^b	288 ^b	299 ^b	219	189	162	181
26 or Older	834	802	877	999	896	964	1,257	1,220	1,219	1,183	1,159	1,218	1,059
ILLCIT DRUGS AND ALCOHOL													
12-17	172 ^b	161 ^b	180 ^b	167 ^b	152 ^b	153 ^b	145 ^a	141 ^a	112	113	135	113	91
18-25	256	352 ^b	317	317	375 ^b	301	325 ^a	383 ^b	317	326	221	304	244
26 or Older	891	742 ^a	970	1,038	1,044	962	854	1,011	885	772 ^a	857	903	1,092

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.74B Type of Substance Treatment Received at Any Location in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use in the Past Year, by Age Group: Percentages, 2002-2014

Substance for Which Treatment Was Received/Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS BUT NOT ALCOHOL													
12-17	19.9	16.4 ^b	19.1 ^a	16.5 ^b	22.0	19.5	18.9 ^a	17.9 ^a	21.4	21.1	25.3	21.2	24.3
18-25	22.4	22.3	23.4	24.0	30.0	22.5	22.7	23.6	31.5	37.5	31.0	26.0	33.0
26 or Older	23.4 ^a	21.3 ^b	25.8	25.6 ^a	20.3 ^b	20.4 ^b	21.8 ^b	21.9 ^b	26.1	28.6	39.3	26.8	33.0
	18.5	13.3 ^b	16.1	12.2 ^b	21.5	18.8	17.6	16.1	18.7	16.7	20.6	19.4	21.6
ALCOHOL BUT NOT ILLICIT DRUGS													
12-17	31.2	33.2	31.4	33.5	29.6	33.8	39.0 ^b	36.1	37.4 ^a	38.1 ^a	34.4	34.7	30.3
18-25	14.9 ^a	15.1 ^a	13.6 ^a	13.5 ^a	12.9 ^a	11.0	12.2	13.0	12.1	10.1	10.1	10.5	6.9
26 or Older	28.7	27.7	29.5	28.4	28.7	37.6 ^b	33.0 ^b	29.3	30.9 ^a	25.1	22.5	21.2	23.7
	34.4	38.7	34.9	38.1	31.9	35.4	43.9 ^a	40.6	42.1 ^a	45.7 ^b	40.3	39.9	33.5
ILLCIT DRUGS AND ALCOHOL													
12-17	37.9	37.7	38.7	38.7	39.0	36.1	32.6	35.9	31.6	32.2	30.3	32.6	34.4
18-25	46.5	44.6	44.5	47.9	43.1	45.0	43.8	49.3	38.6	37.6	47.4	48.7	40.3
26 or Older	37.3	39.5	36.3	33.0	42.9 ^b	34.9	37.6	38.9	32.7	37.4	26.3	39.8	31.9
	36.7	35.8	38.6	39.6	37.2	35.3	29.8	33.6	30.6	29.8	29.8	29.6	34.6

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.75A Type of Substance Treatment Received at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2002-2014

Substance for Which Treatment Was Received/Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS BUT NOT ALCOHOL	645 ^a	442 ^b	609 ^a	481 ^b	809	678	585 ^b	653	787	701	824	786	861
12-17	56	52	47	54	72	46	52	42	52	75	67	38	*
18-25	133 ^a	147	182	194	148	156	137 ^a	180	202	221	259	179	205
26 or Older	456	243 ^b	380 ^a	233 ^b	589	476	395 ^a	431	533	405 ^a	497	569	609
ALCOHOL BUT NOT ILLICIT DRUGS	783	646	727	818	725	898	890	843	891	828	801	840	764
12-17	36	30	37	27	28	22	17	20	19	16	19	13	*
18-25	118	114	156	151	118	145	169 ^a	154	147	140	80	89	112
26 or Older	629	502	534	640	579	731	704	669	725	672	702	738	634
ILLCIT DRUGS AND ALCOHOL	709	595	718	748	734	624	580	750	629	574	633	547	684
12-17	81	55	81	84	63	53	54	68	39	43	52	49	*
18-25	148	178	154	138	185	139	157	169	146	169	92	137	122
26 or Older	480	363	483	527	486	432	368	512	445	361	489	361	536

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.75B Type of Substance Treatment Received at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use at a Specialty Facility in the Past Year, by Age Group: Percentages, 2002-2014

Substance for Which Treatment Was Received/Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS BUT NOT ALCOHOL													
12-17	27.5	23.6 ^b	26.2	20.9 ^b	31.9	27.9	25.5 ^a	24.9 ^a	30.4	30.1	33.0	31.9	33.0
18-25	30.3	31.1	25.7	30.0	39.8	30.5	36.1	27.5	37.9	51.2	43.0	30.7	*
26 or Older	30.5 ^b	30.2 ^b	33.1 ^a	37.1	29.2 ^b	31.9 ^a	27.3 ^b	30.8 ^a	37.7	39.0	55.6 ^a	39.2	43.6
	26.4	19.9 ^a	23.9	14.5 ^b	31.9	26.6	24.0	22.8	27.8	25.1	26.6	30.2	30.0
ALCOHOL BUT NOT ILLICIT DRUGS													
12-17	33.4	34.5	31.2	35.5	28.6	37.0	38.8 ^a	32.1	34.3	35.6	32.1	34.1	29.3
18-25	19.1	18.0	20.1	14.8	15.2	14.4	11.4	13.1	13.4	11.0	11.9	10.6	*
26 or Older	27.2	23.5	28.4	29.0	23.4	29.8	33.7 ^a	26.2	27.4	24.7	17.2	19.5	23.8
	36.5	41.1	33.5	39.9	31.4	40.8	42.8 ^a	35.4	37.8	41.7	37.5	39.1	31.3
ILLCIT DRUGS AND ALCOHOL													
12-17	30.2	31.8	30.9	32.4	29.0	25.7	25.3	28.5	24.3	24.7	25.4	22.2	26.2
18-25	43.9	32.8	43.8	46.2	34.5	35.2	37.5	44.7	28.3	29.7	33.5	39.8	*
26 or Older	34.0	36.5 ^a	28.1	26.4	36.6 ^a	28.4	31.3	28.9	27.1	29.9	19.6	30.0	26.0
	27.8	29.7	30.3	32.8	26.3	24.1	22.3	27.1	23.2	22.4	26.1	19.1	26.5

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.76A Receipt of Illicit Drug Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug Treatment, by Age Group: Numbers in Thousands, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	1,412	1,103 ^b	1,427	1,280 ^a	1,578	1,362	1,211 ^b	1,484	1,518	1,354	1,532	1,483	1,604
12-17	142 ^b	113 ^a	134 ^b	142 ^b	137 ^b	111 ^a	112 ^a	117 ^a	99	125 ^a	121 ^a	90	74
18-25	287	340	352	350	342	307	305	377	355	401	365	341	328
26 or Older	983	649 ^b	941	789 ^b	1,099	944	794 ^b	990	1,064	828 ^b	1,046	1,051	1,201
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	6,335	6,230	6,626	6,269	6,184	6,196	6,369	6,379	6,418	5,832	6,508	6,126	6,278
12-17	1,272 ^b	1,214 ^b	1,262 ^b	1,112 ^b	1,081 ^b	1,013 ^b	1,085 ^b	983 ^a	1,080 ^b	1,060 ^b	920	817	820
18-25	2,393 ^a	2,284	2,505 ^b	2,504 ^b	2,392 ^a	2,447 ^a	2,392 ^a	2,370	2,460 ^a	2,360	2,483 ^a	2,330	2,132
26 or Older	2,670 ^a	2,732 ^a	2,860	2,653 ^b	2,711 ^a	2,735 ^a	2,892	3,026	2,878	2,412 ^b	3,104	2,978	3,326

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.76B Receipt of Illicit Drug Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug Treatment, by Age Group: Percentages, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	18.2	15.0 ^b	17.7	17.0	20.3	18.0	16.0 ^b	18.9	19.1	18.8	19.1	19.5	20.3
12-17	10.1	8.5	9.6	11.3	11.2	9.9	9.3	10.7	8.4	10.5	11.6	10.0	8.3
18-25	10.7	13.0	12.3	12.3	12.5	11.1	11.3	13.7	12.6	14.5	12.8	12.8	13.3
26 or Older	26.9	19.2 ^a	24.8	22.9	28.9	25.7	21.6	24.7	27.0	25.5	25.2	26.1	26.5
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	81.8	85.0 ^b	82.3	83.0	79.7	82.0	84.0 ^b	81.1	80.9	81.2	80.9	80.5	79.7
12-17	89.9	91.5	90.4	88.7	88.8	90.1	90.7	89.3	91.6	89.5	88.4	90.0	91.7
18-25	89.3	87.0	87.7	87.7	87.5	88.9	88.7	86.3	87.4	85.5	87.2	87.2	86.7
26 or Older	73.1	80.8 ^a	75.2	77.1	71.1	74.3	78.4	75.3	73.0	74.5	74.8	73.9	73.5

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.77A Receipt of Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Alcohol Treatment, by Age Group: Numbers in Thousands, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	1,549	1,298	1,535	1,626	1,553	1,575	1,564	1,709	1,591	1,472	1,492	1,421	1,572
12-17	121 ^b	95 ^a	126 ^b	119 ^b	101 ^a	83	78	97 ^a	70	63	76	73	55
18-25	270	305	323 ^a	295	317	293	345 ^a	353 ^b	312	315	180	233	244
26 or Older	1,157	898 ^a	1,086	1,213	1,135	1,199	1,142	1,260	1,209	1,093	1,236	1,115	1,272
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	17,089	16,917	17,824 ^b	17,752 ^b	18,015 ^b	17,775 ^b	17,537 ^b	17,711 ^b	17,042	15,917	16,800	16,628	16,073
12-17	1,374 ^b	1,406 ^b	1,444 ^b	1,341 ^b	1,308 ^b	1,318 ^b	1,177 ^b	1,086 ^b	1,074 ^b	915 ^b	814 ^b	662	644
18-25	5,313 ^b	5,284 ^b	5,421 ^b	5,522 ^b	5,592 ^b	5,348 ^b	5,502 ^b	5,184 ^b	5,143 ^b	4,747 ^b	4,858 ^b	4,400	4,117
26 or Older	10,402	10,228 ^a	10,960	10,889	11,116	11,110	10,858	11,441	10,825	10,255 ^a	11,128	11,567	11,312

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.77B Receipt of Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Alcohol Treatment, by Age Group: Percentages, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	8.3	7.1 ^a	7.9	8.4	7.9	8.1	8.2	8.8	8.5	8.5	8.2	7.9	8.9
12-17	8.1	6.3	8.0	8.1	7.2	5.9	6.2	8.2	6.1	6.4	8.5	10.0	7.9
18-25	4.8	5.5	5.6	5.1	5.4	5.2	5.9	6.4	5.7	6.2	3.6 ^b	5.0	5.6
26 or Older	10.0	8.1	9.0	10.0	9.3	9.7	9.5	9.9	10.0	9.6	10.0	8.8	10.1
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	91.7	92.9 ^a	92.1	91.6	92.1	91.9	91.8	91.2	91.5	91.5	91.8	92.1	91.1
12-17	91.9	93.7	92.0	91.9	92.8	94.1	93.8	91.8	93.9	93.6	91.5	90.0	92.1
18-25	95.2	94.5	94.4	94.9	94.6	94.8	94.1	93.6	94.3	93.8	96.4 ^b	95.0	94.4
26 or Older	90.0	91.9	91.0	90.0	90.7	90.3	90.5	90.1	90.0	90.4	90.0	91.2	89.9

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.78A Receipt of Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment, by Age Group: Numbers in Thousands, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	2,346	1,874 ^b	2,327	2,308	2,533	2,430	2,293	2,627	2,594	2,325	2,496	2,466	2,606
12-17	186 ^b	168 ^a	185 ^b	181 ^b	181 ^b	151	145	152	139	146	157 ^a	122	109
18-25	435	486	548	522	505	489	502	586 ^a	537	567	467	458	470
26 or Older	1,724	1,221 ^b	1,594 ^a	1,605 ^a	1,846	1,790	1,646	1,888	1,919	1,613 ^a	1,872	1,886	2,028
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	20,465	20,290	21,149 ^a	20,864	21,102 ^a	20,835	20,914	21,005	20,615	19,254	20,565	20,221	19,872
12-17	2,071 ^b	2,085 ^b	2,103 ^b	1,915 ^b	1,908 ^b	1,833 ^b	1,804 ^b	1,632 ^b	1,694 ^b	1,598 ^b	1,407 ^b	1,218	1,175
18-25	6,439 ^b	6,338 ^b	6,499 ^b	6,699 ^b	6,650 ^b	6,450 ^b	6,552 ^b	6,343 ^b	6,419 ^b	5,994 ^b	6,215 ^b	5,689	5,375
26 or Older	11,956 ^a	11,867 ^b	12,547	12,251 ^a	12,543	12,552	12,558	13,030	12,502	11,661 ^b	12,944	13,314	13,321

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.78B Receipt of Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment, by Age Group: Percentages, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	10.3	8.5 ^b	9.9	10.0	10.7	10.4	9.9	11.1	11.2	10.8	10.8	10.9	11.6
12-17	8.2	7.4	8.1	8.6	8.7	7.6	7.4	8.5	7.6	8.4	10.0	9.1	8.5
18-25	6.3 ^a	7.1	7.8	7.2	7.1	7.0	7.1	8.5	7.7	8.6	7.0	7.4	8.0
26 or Older	12.6	9.3 ^b	11.3	11.6	12.8	12.5	11.6	12.7	13.3	12.1	12.6	12.4	13.2
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	89.7	91.5 ^b	90.1	90.0	89.3	89.6	90.1	88.9	88.8	89.2	89.2	89.1	88.4
12-17	91.8	92.6	91.9	91.4	91.3	92.4	92.6	91.5	92.4	91.6	90.0	90.9	91.5
18-25	93.7 ^a	92.9	92.2	92.8	92.9	93.0	92.9	91.5	92.3	91.4	93.0	92.6	92.0
26 or Older	87.4	90.7 ^b	88.7	88.4	87.2	87.5	88.4	87.3	86.7	87.9	87.4	87.6	86.8

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2014 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2014 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.79B Marijuana Use in the Past Month, by Age Group: Percentages, 1971-2014

Survey Year	Aged 12-17	Aged 18-25	Aged 26-34
1971	5.1	17.3	5.7
1972	6.0	27.8	10.3
1974	10.2	25.2	9.1
1976	10.5	25.0	12.5
1977	14.1	27.4	13.7
1979	14.2	35.6	19.7
1982	9.9	27.2	19.0
1985	10.2	21.7	19.0
1988	5.4	15.3	12.3
1990	4.4	12.7	9.5
1991	3.6	12.9	7.7
1992	3.4	10.9	9.3
1993	4.0	11.1	7.5
1994	6.0	12.1	6.9
1995	8.2	12.0	6.7
1996	7.1	13.2	6.3
1997	9.4	12.8	6.0
1998	8.3	13.8	5.5
1999	7.2	14.2	5.4
2000	7.2	13.6	5.9
2001	8.0	16.0	6.8
2002	8.2	17.3	7.7
2003	7.9	17.0	8.4
2004	7.6	16.1	8.3
2005	6.8	16.6	8.6
2006	6.7	16.3	8.5
2007	6.7	16.5	7.9
2008	6.7	16.6	8.8
2009	7.4	18.2	9.6
2010	7.4	18.5	10.6
2011	7.9	19.0	10.2
2012	7.2	18.7	11.3
2013	7.1	19.1	12.6
2014	7.4	19.6	12.7

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for 1971 through 1977 for youths aged 12 to 17 and for adults aged 26 to 34 may differ from estimates for these age groups for these survey years that were published in other NSDUH reports. The estimates shown here for these survey years have been adjusted to improve their comparability with estimates based on later versions of the NSDUH instrument. Estimates were adjusted by applying the ratio of the adjusted and unadjusted rate from 1979 for each age group to the original estimate.

NOTE: Estimates for 1979 through 1993 have been adjusted for comparability with estimates for 1994 through 1998. Estimates were adjusted by formally modeling the effect of the new methodology, relative to the old methodology, using 1994 data. See Appendix E of the *National Household Survey on Drug Abuse: Main Findings 1998* (<http://oas.samhsa.gov/nsduh/reports.htm>) for information on the adjustment procedures.

NOTE: Estimates for 1999 through 2014 have not been adjusted for comparability. See Appendix C of the *Results from the 2004 National Survey on Drug Use and Health: National Findings* for information on the comparability of the 1999 and later estimates with pre-1999 estimates.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 1971-2014.

Table 7.80B Cocaine Use in the Past Year, by Age Group: Percentages, 1971-2014

Survey Year	Aged 12-17	Aged 18-25	Aged 26-34
1971	--	--	--
1972	1.3	--	--
1974	2.3	7.0	--
1976	2.0	6.1	--
1977	2.2	8.8	--
1979	3.6	17.0	5.7
1982	3.7	15.9	9.3
1985	3.4	13.6	10.5
1988	2.5	10.5	7.0
1990	1.9	6.5	5.9
1991	1.3	6.7	4.4
1992	1.0	5.5	4.3
1993	0.7	4.4	3.8
1994	1.1	3.6	3.5
1995	1.7	4.3	3.1
1996	1.4	4.7	3.5
1997	2.2	3.9	3.1
1998	1.7	4.7	2.7
1999	1.6	5.2	2.4
2000	1.7	4.4	2.1
2001	1.5	5.7	2.7
2002	2.1	6.7	3.6
2003	1.8	6.6	4.1
2004	1.6	6.6	4.0
2005	1.7	6.9	3.1
2006	1.6	6.9	4.4
2007	1.5	6.4	3.6
2008	1.2	5.6	4.0
2009	1.0	5.3	3.4
2010	1.0	4.7	3.6
2011	0.9	4.6	2.3
2012	0.7	4.6	3.1
2013	0.5	4.4	3.4
2014	0.7	4.6	3.0

*Low precision; no estimate reported.

-- Not available.

NOTE: Cocaine use includes the use of crack.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for 1971 through 1977 for youths aged 12 to 17 may differ from estimates for these age groups for these survey years that were published in other NSDUH reports. The estimates shown here for these survey years have been adjusted to improve their comparability with estimates based on later versions of the NSDUH instrument. Estimates have been adjusted by applying the ratio of the adjusted and unadjusted rate from 1979 for each age group to the original estimate.

NOTE: Estimates for 1979 through 1993 have been adjusted for comparability with estimates for 1994 through 1998. Estimates were adjusted using a simple ratio correction factor that measured the effect of the new methodology, relative to the old methodology, estimated at the total population level using pooled 1993 and 1994 data. See Appendix E of the *National Household Survey on Drug Abuse: Main Findings 1998* (<http://oas.samhsa.gov/nsduh/reports.htm>) for further information on the adjustment procedures.

NOTE: Estimates for 1999 through 2014 have not been adjusted for comparability. See Appendix C of the *Results from the 2004 National Survey on Drug Use and Health: National Findings* for information on the comparability of the 1999 and later estimates with pre-1999 estimates.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 1971-2014.

Table 7.1C Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	963	909	969	940	899	974	927	966	1,023	1,036	956	1,015	873
Marijuana and Hashish	912	868	912	925	909	964	917	949	1,019	1,013	1,005	1,017	872
Cocaine	583	565	636	595	655	660	672	655	682	701	703	693	632
Crack	306	264	280	292	329	339	323	316	379	327	346	364	317
Heroin	223	205	193	226	238	240	224	217	249	227	245	281	213
Hallucinogens	602	589	618	608	644	640	647	666	687	628	674	707	636
LSD	514	494	506	521	520	550	548	564	560	530	561	580	508
PCP	284	271	298	289	294	298	339	298	321	307	311	346	274
Ecstasy	297	285	329	335	355	346	344	371	430	367	405	464	384
Inhalants	474	441	480	460	471	503	490	473	473	491	489	519	428
Nonmedical Use of Psychotherapeutics ^{2,3}	743	653	736	719	701	698	757	740	784	752	765	785	658
Pain Relievers	569	517	564	562	540	563	597	584	610	598	622	640	553
OxyContin [®]	116	140	134	142	168	181	187	222	246	211	245	280	220
Tranquilizers	497	435	490	480	477	469	506	494	494	494	563	576	443
Stimulants ³	515	475	498	491	508	472	546	509	557	500	556	501	441
Methamphetamine ³	417	403	401	375	413	368	400	404	452	410	426	393	363
Sedatives	370	325	403	347	353	360	401	354	355	353	377	377	324
ILLCIT DRUGS OTHER THAN MARIJUANA¹	815	756	827	835	809	858	855	877	900	889	873	923	803

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.1D Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.41	0.38	0.40	0.39	0.37	0.39	0.37	0.38	0.40	0.40	0.37	0.39	0.33
Marijuana and Hashish	0.39	0.37	0.38	0.38	0.37	0.39	0.37	0.38	0.40	0.39	0.39	0.39	0.33
Cocaine	0.25	0.24	0.26	0.24	0.27	0.27	0.27	0.26	0.27	0.27	0.27	0.26	0.24
Crack	0.13	0.11	0.12	0.12	0.13	0.14	0.13	0.13	0.15	0.13	0.13	0.14	0.12
Heroin	0.10	0.09	0.08	0.09	0.10	0.10	0.09	0.09	0.10	0.09	0.09	0.11	0.08
Hallucinogens	0.26	0.25	0.26	0.25	0.26	0.26	0.26	0.26	0.27	0.24	0.26	0.27	0.24
LSD	0.22	0.21	0.21	0.21	0.21	0.22	0.22	0.22	0.22	0.21	0.22	0.22	0.19
PCP	0.12	0.11	0.12	0.12	0.12	0.12	0.14	0.12	0.13	0.12	0.12	0.13	0.10
Ecstasy	0.13	0.12	0.14	0.14	0.14	0.14	0.14	0.15	0.17	0.14	0.16	0.18	0.14
Inhalants	0.20	0.19	0.20	0.19	0.19	0.20	0.20	0.19	0.19	0.19	0.19	0.20	0.16
Nonmedical Use of Psychotherapeutics ^{2,3}	0.32	0.27	0.31	0.30	0.28	0.28	0.30	0.29	0.31	0.29	0.29	0.30	0.25
Pain Relievers	0.24	0.22	0.23	0.23	0.22	0.23	0.24	0.23	0.24	0.23	0.24	0.24	0.21
OxyContin [®]	0.05	0.06	0.06	0.06	0.07	0.07	0.07	0.09	0.10	0.08	0.09	0.11	0.08
Tranquilizers	0.21	0.18	0.20	0.20	0.19	0.19	0.20	0.20	0.19	0.19	0.22	0.22	0.17
Stimulants ³	0.22	0.20	0.21	0.20	0.21	0.19	0.22	0.20	0.22	0.19	0.21	0.19	0.17
Methamphetamine ³	0.18	0.17	0.17	0.15	0.17	0.15	0.16	0.16	0.18	0.16	0.16	0.15	0.14
Sedatives	0.16	0.14	0.17	0.14	0.14	0.15	0.16	0.14	0.14	0.14	0.14	0.14	0.12
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.35	0.32	0.34	0.34	0.33	0.35	0.34	0.35	0.35	0.35	0.34	0.35	0.30

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.1P Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0004	0.0004	0.0000	0.0184	0.2511	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0048	0.3327	
Cocaine	0.2891	0.7531	0.1003	0.0060	0.2225	0.4303	0.9815	0.5808	0.8799	0.2109	0.4157	0.2134	
Crack	0.9156	0.1976	0.0774	0.0820	0.6744	0.6664	0.3334	0.1993	0.6904	0.0362	0.6235	0.3420	
Heroin	0.0404	0.0418	0.0000	0.0033	0.0286	0.0264	0.0141	0.0027	0.1525	0.0946	0.6279	0.8887	
Hallucinogens	0.3032	0.1501	0.0536	0.0018	0.0862	0.0015	0.1430	0.7785	0.6765	0.0144	0.2857	0.5990	
LSD	0.0008	0.0033	0.3160	0.4451	0.8486	0.3469	0.9841	0.9306	0.4394	0.0679	0.2742	0.9928	
PCP	0.0000	0.0002	0.0130	0.0528	0.0701	0.5990	0.1351	0.6068	0.7287	0.7981	0.4872	0.7509	
Ecstasy	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.1300	0.0000	0.0580	0.4555	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0002	0.0247	0.7973	0.7467	0.9936	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.7626	0.7886	0.8417	0.7254	0.5856	0.7401	0.3583	0.7361	0.8403	0.1037	0.3023	0.5162	
Pain Relievers	0.0016	0.1166	0.2092	0.6033	0.9873	0.4467	0.2026	0.2303	0.6126	0.3235	0.0433	0.7945	
OxyContin [®]	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0077	0.0803	0.0024	0.3307	0.9676	
Tranquilizers	0.0000	0.0005	0.0000	0.0054	0.0058	0.0000	0.0046	0.0070	0.0151	0.0002	0.3003	0.1285	
Stimulants ³	0.0000	0.0000	0.0037	0.6220	0.0151	0.3330	0.9359	0.3364	0.7892	0.0219	0.3925	0.3336	
Methamphetamine ³	0.0000	0.0000	0.0000	0.1157	0.0000	0.0500	0.4052	0.2470	0.2332	0.2314	0.4322	0.2996	
Sedatives	0.0000	0.0000	0.0000	0.0001	0.0007	0.0216	0.0022	0.0108	0.7252	0.8507	0.5954	0.5922	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.4831	0.5036	0.0664	0.1322	0.1993	0.2754	0.7171	0.9882	0.8853	0.0429	0.6378	0.3178	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.2C Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	588	558	535	550	566	596	553	598	646	603	648	677	595
Marijuana and Hashish	471	455	451	473	478	493	474	541	559	514	556	577	538
Cocaine	222	229	205	205	230	242	207	209	205	168	217	189	186
Crack	124	124	117	114	134	123	94	105	92	69	111	77	81
Heroin	55	48	56	43	86	54	59	79	75	72	78	70	75
Hallucinogens	151	147	153	146	149	140	130	158	178	143	175	204	159
LSD	63	44	47	48	56	55	50	55	72	60	85	81	76
PCP	25	24	28	21	38	19	14	22	24	20	34	21	16
Ecstasy	126	105	112	114	106	108	96	126	128	107	134	153	121
Inhalants	95	83	90	95	107	106	115	111	115	107	120	105	95
Nonmedical Use of Psychotherapeutics ^{2,3}	379	358	352	346	360	381	352	366	397	363	406	418	321
Pain Relievers	300	299	300	304	306	316	294	322	336	302	355	343	253
OxyContin [®]	--	--	74	81	93	96	88	105	116	104	105	119	83
Tranquilizers	225	195	208	207	197	222	200	222	207	211	252	230	190
Stimulants ³	154	136	157	134	180	152	123	149	158	151	161	178	159
Methamphetamine ³	116	100	128	99	137	115	75	100	105	118	113	114	104
Sedatives	116	85	72	74	99	94	61	80	103	64	66	84	75
ILLCIT DRUGS OTHER THAN MARIJUANA¹	432	420	410	398	418	449	407	420	446	404	459	469	371

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.2D Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.25	0.23	0.22	0.23	0.23	0.24	0.22	0.24	0.25	0.23	0.25	0.26	0.22
Marijuana and Hashish	0.20	0.19	0.19	0.19	0.19	0.20	0.19	0.21	0.22	0.20	0.21	0.22	0.20
Cocaine	0.09	0.10	0.09	0.08	0.09	0.10	0.08	0.08	0.08	0.07	0.08	0.07	0.07
Crack	0.05	0.05	0.05	0.05	0.05	0.05	0.04	0.04	0.04	0.03	0.04	0.03	0.03
Heroin	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03
Hallucinogens	0.06	0.06	0.06	0.06	0.06	0.06	0.05	0.06	0.07	0.06	0.07	0.08	0.06
LSD	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.03	0.03	0.03
PCP	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Ecstasy	0.05	0.04	0.05	0.05	0.04	0.04	0.04	0.05	0.05	0.04	0.05	0.06	0.05
Inhalants	0.04	0.03	0.04	0.04	0.04	0.04	0.05	0.04	0.05	0.04	0.05	0.04	0.04
Nonmedical Use of Psychotherapeutics ^{2,3}	0.16	0.15	0.15	0.14	0.15	0.15	0.14	0.15	0.16	0.14	0.16	0.16	0.12
Pain Relievers	0.13	0.13	0.12	0.12	0.12	0.13	0.12	0.13	0.13	0.12	0.14	0.13	0.10
OxyContin [®]	--	--	0.03	0.03	0.04	0.04	0.04	0.04	0.05	0.04	0.04	0.05	0.03
Tranquilizers	0.10	0.08	0.09	0.08	0.08	0.09	0.08	0.09	0.08	0.08	0.10	0.09	0.07
Stimulants ³	0.07	0.06	0.07	0.06	0.07	0.06	0.05	0.06	0.06	0.06	0.06	0.07	0.06
Methamphetamine ³	0.05	0.04	0.05	0.04	0.06	0.05	0.03	0.04	0.04	0.05	0.04	0.04	0.04
Sedatives	0.05	0.04	0.03	0.03	0.04	0.04	0.02	0.03	0.04	0.02	0.03	0.03	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.18	0.18	0.17	0.16	0.17	0.18	0.16	0.17	0.18	0.16	0.18	0.18	0.14

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.2P Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0358	0.0189	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0214	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0789	0.5120	0.0216	0.4703	0.2191	
Crack	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0016	0.0365	0.2306	0.2274	0.2349	0.2313	
Heroin	0.0000	0.0000	0.0000	0.0000	0.0096	0.0000	0.0000	0.0072	0.0153	0.0094	0.0356	0.0295	
Hallucinogens	0.0000	0.5403	0.9159	0.6639	0.9014	0.2954	0.0920	0.0229	0.0419	0.7723	0.5598	0.3865	
LSD	0.1178	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0001	0.0662	0.1356	
PCP	0.0000	0.0000	0.0001	0.0018	0.0119	0.0256	0.5032	0.1453	0.7162	0.2139	0.0241	0.9760	
Ecstasy	0.0000	0.8947	0.1806	0.2360	0.8179	0.6673	0.5431	0.0007	0.0170	0.3559	0.0813	0.1639	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0001	0.0007	0.0402	0.4825	0.6320	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0014	0.0002	0.0055	0.0004	0.0000	0.0000	0.0193	0.0001	0.0006	0.8090	0.0001	0.3072	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0048	0.0000	0.0451	
OxyContin [®]	--	--	0.3922	0.4147	0.1472	0.0294	0.0126	0.0001	0.0000	0.0014	0.0486	0.1321	
Tranquilizers	0.4026	0.1343	0.1971	0.0782	0.3668	0.1298	0.4633	0.0539	0.0280	0.8448	0.0020	0.6857	
Stimulants ³	0.6838	0.1291	0.5867	0.1067	0.1441	0.0245	0.0000	0.0402	0.0030	0.0000	0.1445	0.4362	
Methamphetamine ³	0.0001	0.0015	0.0001	0.0031	0.0001	0.3905	0.0028	0.6283	0.0487	0.1395	0.4296	0.5088	
Sedatives	0.0286	0.2063	0.7272	0.6959	0.0854	0.2315	0.2638	0.4520	0.1883	0.0187	0.0843	0.2542	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0009	0.0001	0.0000	0.0000	0.0075	0.0000	0.0021	0.7120	0.0011	0.5555	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.3C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	388	404	415	432	427	445	427	457	488	460	492	509	476
Marijuana and Hashish	320	327	360	360	357	359	377	395	407	409	431	442	424
Cocaine	128	147	119	142	153	149	131	136	108	106	142	133	105
Crack	68	91	66	87	101	80	55	77	60	42	94	64	55
Heroin	42	28	37	19	77	38	39	37	43	48	51	43	55
Hallucinogens	80	72	58	83	70	75	68	92	84	60	90	132	88
LSD	18	18	26	16	24	33	19	22	20	23	31	37	40
PCP	15	13	17	13	9	13	8	18	21	10	13	15	*
Ecstasy	64	55	47	44	51	56	51	69	61	45	72	91	65
Inhalants	64	45	64	57	73	67	71	58	83	68	65	64	65
Nonmedical Use of Psychotherapeutics ^{2,3}	230	256	212	232	244	251	225	250	268	231	265	256	222
Pain Relievers	186	207	170	194	217	214	196	214	215	189	211	206	174
OxyContin [®]	--	--	36	38	35	53	58	59	67	54	46	81	50
Tranquilizers	136	118	103	118	115	129	118	137	138	133	162	121	115
Stimulants ³	94	102	99	81	105	94	75	99	86	89	101	116	116
Methamphetamine ³	78	81	76	62	79	80	53	67	53	72	74	85	78
Sedatives	64	55	48	47	69	68	37	55	72	49	50	67	54
ILLCIT DRUGS OTHER THAN MARIJUANA¹	276	289	255	279	284	306	261	292	300	256	309	306	261

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.3D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.17	0.17	0.17	0.18	0.17	0.18	0.17	0.18	0.19	0.18	0.19	0.19	0.18
Marijuana and Hashish	0.14	0.14	0.15	0.15	0.15	0.14	0.15	0.16	0.16	0.16	0.17	0.17	0.16
Cocaine	0.05	0.06	0.05	0.06	0.06	0.06	0.05	0.05	0.04	0.04	0.05	0.05	0.04
Crack	0.03	0.04	0.03	0.04	0.04	0.03	0.02	0.03	0.02	0.02	0.04	0.02	0.02
Heroin	0.02	0.01	0.02	0.01	0.03	0.02	0.02	0.01	0.02	0.02	0.02	0.02	0.02
Hallucinogens	0.03	0.03	0.02	0.03	0.03	0.03	0.03	0.04	0.03	0.02	0.03	0.05	0.03
LSD	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02
PCP	0.01	0.01	0.01	0.01	0.00	0.01	0.00	0.01	0.01	0.00	0.00	0.01	*
Ecstasy	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.03	0.03	0.02
Inhalants	0.03	0.02	0.03	0.02	0.03	0.03	0.03	0.02	0.03	0.03	0.02	0.02	0.02
Nonmedical Use of Psychotherapeutics ^{2,3}	0.10	0.11	0.09	0.10	0.10	0.10	0.09	0.10	0.11	0.09	0.10	0.10	0.08
Pain Relievers	0.08	0.09	0.07	0.08	0.09	0.09	0.08	0.09	0.08	0.07	0.08	0.08	0.07
OxyContin [®]	--	--	0.02	0.02	0.01	0.02	0.02	0.02	0.03	0.02	0.02	0.03	0.02
Tranquilizers	0.06	0.05	0.04	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.06	0.05	0.04
Stimulants ³	0.04	0.04	0.04	0.03	0.04	0.04	0.03	0.04	0.03	0.03	0.04	0.04	0.04
Methamphetamine ³	0.03	0.03	0.03	0.03	0.03	0.03	0.02	0.03	0.02	0.03	0.03	0.03	0.03
Sedatives	0.03	0.02	0.02	0.02	0.03	0.03	0.01	0.02	0.03	0.02	0.02	0.03	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.12	0.12	0.11	0.11	0.12	0.12	0.10	0.12	0.12	0.10	0.12	0.12	0.10

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.3P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0022	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0101	0.2625	0.9546	0.4276	0.3968	0.8370	
Crack	0.0025	0.0059	0.0798	0.0004	0.0010	0.0034	0.7009	0.0863	0.6220	0.0895	0.3778	0.7509	
Heroin	0.0006	0.0000	0.0003	0.0000	0.4844	0.0001	0.0026	0.0006	0.0096	0.0492	0.2145	0.0413	
Hallucinogens	0.1645	0.9291	0.1739	0.9214	0.4553	0.3753	0.6660	0.2306	0.4743	0.1096	0.8909	0.2769	
LSD	0.0004	0.0023	0.0075	0.0001	0.0022	0.0131	0.0078	0.0099	0.0059	0.0050	0.0530	0.3098	
PCP	*	*	*	*	*	*	*	*	*	*	*	*	
Ecstasy	0.1161	0.3403	0.1741	0.4427	0.6475	0.4224	0.7272	0.0474	0.2151	0.5364	0.7533	0.6122	
Inhalants	0.0804	0.2695	0.1000	0.1809	0.0069	0.2387	0.1672	0.6057	0.1055	0.3102	0.6384	0.6223	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.1071	0.0693	0.5398	0.1100	0.0013	0.0148	0.8703	0.0187	0.0401	0.4634	0.2226	0.9662	
Pain Relievers	0.0256	0.0017	0.0388	0.0062	0.0000	0.0000	0.0104	0.0000	0.0005	0.2908	0.0230	0.3709	
OxyContin [®]	--	--	0.9711	0.9048	0.3381	0.5869	0.1788	0.0243	0.0065	0.2289	0.8912	0.1404	
Tranquilizers	0.4075	0.3445	0.5621	0.5402	0.8954	0.6126	0.8556	0.1550	0.0384	0.9168	0.1304	0.3646	
Stimulants ³	0.4202	0.4085	0.3503	0.0398	0.5276	0.0022	0.0000	0.1442	0.0016	0.0001	0.0097	0.1908	
Methamphetamine ³	0.0885	0.0439	0.0686	0.2623	0.0596	0.9818	0.0145	0.7039	0.0368	0.2747	0.2676	0.7826	
Sedatives	0.0724	0.9888	0.6248	0.6531	0.3578	0.6642	0.2453	0.4738	0.5057	0.2115	0.4635	0.3799	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0038	0.0056	0.3329	0.0089	0.0000	0.0036	0.3442	0.0172	0.0841	0.2105	0.4100	0.9303	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.4C Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	104	107	96	100	104	103	98	103	102	103	92	97	108
Marijuana and Hashish	88	91	87	85	87	88	82	86	91	89	82	86	95
Cocaine	36	36	33	34	32	32	32	28	26	26	23	19	26
Crack	17	17	14	14	15	13	12	10	11	16	10	5	9
Heroin	14	11	12	10	11	10	13	9	9	12	12	9	9
Hallucinogens	49	48	45	44	42	41	41	43	44	43	39	35	37
LSD	34	26	24	23	21	18	21	20	21	19	21	19	27
PCP	20	20	18	19	16	14	12	13	11	10	13	8	11
Ecstasy	40	32	31	26	28	30	33	34	35	37	30	24	27
Inhalants	67	68	67	67	66	68	65	66	61	58	51	52	56
Nonmedical Use of Psychotherapeutics ^{2,3}	75	76	70	72	75	74	69	70	68	64	64	63	69
Pain Relievers	69	70	67	65	68	69	66	64	64	59	58	58	63
OxyContin [®]	21	20	24	21	24	25	23	27	28	25	23	22	24
Tranquilizers	38	40	36	38	36	37	35	33	36	37	38	33	42
Stimulants ³	46	41	38	41	43	34	39	31	31	31	28	28	37
Methamphetamine ³	29	24	25	26	28	19	22	20	18	21	15	14	22
Sedatives	22	22	23	22	19	20	18	21	17	16	17	14	23
ILLCIT DRUGS OTHER THAN MARIJUANA¹	89	97	84	87	87	92	83	88	87	82	77	77	84

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.4D Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.42	0.43	0.38	0.39	0.41	0.41	0.39	0.42	0.42	0.41	0.37	0.39	0.43
Marijuana and Hashish	0.36	0.37	0.34	0.33	0.34	0.35	0.33	0.35	0.37	0.35	0.33	0.35	0.38
Cocaine	0.15	0.15	0.13	0.14	0.13	0.13	0.13	0.11	0.11	0.11	0.09	0.08	0.10
Crack	0.07	0.07	0.06	0.05	0.06	0.05	0.05	0.04	0.04	0.06	0.04	0.02	0.03
Heroin	0.05	0.04	0.05	0.04	0.04	0.04	0.05	0.04	0.04	0.05	0.05	0.04	0.03
Hallucinogens	0.20	0.19	0.18	0.17	0.17	0.16	0.16	0.18	0.18	0.17	0.15	0.14	0.15
LSD	0.14	0.10	0.09	0.09	0.08	0.07	0.08	0.08	0.09	0.08	0.09	0.08	0.11
PCP	0.08	0.08	0.07	0.08	0.06	0.05	0.05	0.05	0.05	0.04	0.05	0.03	0.05
Ecstasy	0.16	0.13	0.12	0.10	0.11	0.12	0.13	0.14	0.14	0.15	0.12	0.10	0.11
Inhalants	0.27	0.27	0.27	0.26	0.26	0.27	0.26	0.27	0.25	0.23	0.21	0.21	0.23
Nonmedical Use of Psychotherapeutics ^{2,3}	0.30	0.30	0.28	0.28	0.30	0.29	0.28	0.28	0.28	0.26	0.26	0.25	0.28
Pain Relievers	0.28	0.28	0.27	0.26	0.27	0.27	0.26	0.26	0.26	0.23	0.23	0.23	0.25
OxyContin [®]	0.09	0.08	0.09	0.08	0.09	0.10	0.09	0.11	0.12	0.10	0.09	0.09	0.10
Tranquilizers	0.16	0.16	0.14	0.15	0.14	0.15	0.14	0.13	0.15	0.15	0.15	0.13	0.17
Stimulants ³	0.19	0.16	0.15	0.16	0.17	0.13	0.16	0.12	0.13	0.12	0.11	0.11	0.15
Methamphetamine ³	0.12	0.10	0.10	0.10	0.11	0.07	0.09	0.08	0.08	0.08	0.06	0.06	0.09
Sedatives	0.09	0.09	0.09	0.09	0.08	0.08	0.07	0.09	0.07	0.06	0.07	0.06	0.09
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.36	0.39	0.33	0.34	0.34	0.36	0.33	0.36	0.36	0.33	0.31	0.31	0.34

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.4P Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0910	0.9952	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0720	0.0866	0.6691	0.7307	0.2227	0.2425	0.0383	0.2975	0.9930	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0023	0.2277	0.8252	
Crack	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0028	0.0620	0.0587	0.0643	0.9772	0.2332	
Heroin	0.0003	0.0019	0.0053	0.0705	0.0961	0.1616	0.0406	0.0879	0.3580	0.0227	0.1056	0.8141	
Hallucinogens	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0681	
LSD	0.0000	0.0051	0.6146	0.8126	0.0239	0.0046	0.5600	0.2524	0.1147	0.0598	0.2151	0.0332	
PCP	0.0000	0.0000	0.0000	0.0000	0.0008	0.0133	0.0114	0.0264	0.2909	0.3017	0.0373	0.2942	
Ecstasy	0.0000	0.0000	0.0000	0.0118	0.0000	0.0003	0.0000	0.0000	0.0000	0.0000	0.0000	0.0462	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.7986	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0186	0.0485	0.2713	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0024	0.9936	
OxyContin [®]	0.3967	1.0000	0.1534	0.5662	0.0131	0.0030	0.0062	0.0003	0.0010	0.0010	0.0413	0.5099	
Tranquilizers	0.0026	0.0005	0.0116	0.1260	0.2637	0.1969	0.3168	0.4818	0.1924	0.5761	0.5239	0.0814	
Stimulants ³	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0051	0.1615	0.6770	0.7469	0.5473	0.0659	
Methamphetamine ³	0.0000	0.0000	0.0000	0.0000	0.0000	0.1041	0.4507	0.5533	0.7158	0.6903	0.3173	0.0428	
Sedatives	0.3092	0.1904	0.1964	0.5135	0.6753	0.9027	0.9556	0.5306	0.3943	0.0226	0.2605	0.0028	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0015	0.6935	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.5C Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	95	91	87	88	95	88	87	90	94	93	83	86	95
Marijuana and Hashish	79	79	77	75	78	75	73	78	84	83	75	78	82
Cocaine	32	28	27	28	28	27	24	22	22	20	19	15	23
Crack	14	13	10	10	12	11	8	7	5	7	8	4	7
Heroin	11	8	10	8	8	6	11	7	8	11	9	8	8
Hallucinogens	40	37	36	36	34	34	35	38	40	35	32	29	31
LSD	25	15	17	17	14	15	18	14	17	18	16	15	25
PCP	13	12	12	14	11	9	10	10	8	7	11	7	9
Ecstasy	32	23	23	21	23	26	27	29	32	29	23	19	23
Inhalants	44	48	45	47	45	43	44	42	42	40	34	29	36
Nonmedical Use of Psychotherapeutics ^{2,3}	64	64	59	61	62	63	58	59	58	53	53	50	60
Pain Relievers	59	59	56	56	56	59	54	54	54	48	47	46	53
OxyContin [®]	--	--	18	18	18	19	17	20	24	18	17	16	19
Tranquilizers	34	33	29	29	30	31	29	26	30	31	31	24	35
Stimulants ³	35	30	28	31	33	27	28	23	24	23	23	21	33
Methamphetamine ³	22	17	17	20	21	14	14	13	15	16	12	10	18
Sedatives	16	16	15	16	13	15	14	16	14	12	11	11	18
ILLCIT DRUGS OTHER THAN MARIJUANA¹	76	79	71	71	73	73	69	72	74	65	64	59	70

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.5D Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.38	0.36	0.34	0.35	0.37	0.35	0.35	0.36	0.38	0.37	0.33	0.35	0.38
Marijuana and Hashish	0.32	0.31	0.31	0.30	0.31	0.30	0.29	0.32	0.34	0.33	0.30	0.31	0.33
Cocaine	0.13	0.11	0.11	0.11	0.11	0.11	0.10	0.09	0.09	0.08	0.08	0.06	0.09
Crack	0.06	0.05	0.04	0.04	0.05	0.04	0.03	0.03	0.02	0.03	0.03	0.02	0.03
Heroin	0.04	0.03	0.04	0.03	0.03	0.02	0.04	0.03	0.03	0.05	0.04	0.03	0.03
Hallucinogens	0.16	0.15	0.14	0.14	0.13	0.14	0.14	0.16	0.16	0.14	0.13	0.12	0.13
LSD	0.10	0.06	0.07	0.07	0.05	0.06	0.07	0.06	0.07	0.07	0.07	0.06	0.10
PCP	0.05	0.05	0.05	0.06	0.04	0.04	0.04	0.04	0.03	0.03	0.04	0.03	0.03
Ecstasy	0.13	0.09	0.09	0.08	0.09	0.10	0.11	0.12	0.13	0.12	0.09	0.08	0.09
Inhalants	0.18	0.19	0.18	0.19	0.18	0.17	0.18	0.17	0.17	0.16	0.14	0.12	0.14
Nonmedical Use of Psychotherapeutics ^{2,3}	0.26	0.26	0.23	0.24	0.24	0.25	0.23	0.24	0.24	0.21	0.21	0.20	0.24
Pain Relievers	0.24	0.24	0.22	0.22	0.22	0.23	0.22	0.22	0.22	0.19	0.19	0.18	0.21
OxyContin [®]	--	--	0.07	0.07	0.07	0.07	0.07	0.08	0.10	0.07	0.07	0.06	0.08
Tranquilizers	0.14	0.13	0.12	0.11	0.12	0.12	0.12	0.11	0.12	0.12	0.13	0.09	0.14
Stimulants ³	0.14	0.12	0.11	0.12	0.13	0.11	0.11	0.09	0.10	0.09	0.09	0.09	0.13
Methamphetamine ³	0.09	0.07	0.07	0.08	0.08	0.06	0.06	0.05	0.06	0.06	0.05	0.04	0.07
Sedatives	0.07	0.06	0.06	0.06	0.05	0.06	0.06	0.06	0.06	0.05	0.05	0.04	0.07
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.31	0.32	0.28	0.28	0.29	0.29	0.28	0.29	0.30	0.26	0.26	0.24	0.28

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.5P Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0001	0.0100	0.0015	0.0000	0.0002	0.0043	0.3352	0.6826	
Marijuana and Hashish	0.0000	0.0000	0.0022	0.6025	0.8243	0.1753	0.9536	0.2281	0.0569	0.0174	0.4007	0.4690	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0169	0.0167	0.0458	0.6862	0.1860	
Crack	0.0000	0.0000	0.0005	0.0045	0.0001	0.0005	0.1475	0.3201	0.9114	0.3495	0.9361	0.2519	
Heroin	0.0982	0.4801	0.1368	0.4984	0.4930	0.6106	0.3711	0.5571	0.9830	0.0740	0.8245	0.7859	
Hallucinogens	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0035	0.1625	
LSD	0.0041	0.0226	0.0474	0.0247	0.0001	0.0018	0.3236	0.0131	0.0123	0.0663	0.0341	0.0256	
PCP	0.0000	0.0002	0.0108	0.0028	0.0849	0.0909	0.0324	0.2749	0.8257	0.6101	0.2849	0.6833	
Ecstasy	0.0000	0.0000	0.0001	0.0241	0.0002	0.0000	0.0000	0.0000	0.0000	0.0000	0.0012	0.0878	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0070	0.4803	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0187	0.2940	0.1489	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0201	0.9004	
OxyContin [®]	--	--	0.2498	0.3914	0.1786	0.0111	0.0122	0.0177	0.0114	0.0300	0.3389	0.6238	
Tranquilizers	0.0028	0.0009	0.0258	0.1579	0.2678	0.1121	0.1966	0.3289	0.2568	0.2988	0.5646	0.0642	
Stimulants ³	0.0000	0.0000	0.0011	0.0024	0.0002	0.2593	0.7810	0.5498	0.4702	0.1106	0.3447	0.0237	
Methamphetamine ³	0.0000	0.0017	0.0059	0.0087	0.0053	0.6794	0.6313	0.4342	0.8180	1.0000	0.2853	0.1223	
Sedatives	0.3598	0.5592	0.5119	0.9915	0.7015	0.7647	0.9066	0.6342	0.9099	0.1365	0.1060	0.0150	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0231	0.2851	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.6C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	71	67	67	64	67	68	61	66	70	68	63	62	73
Marijuana and Hashish	58	60	57	56	55	56	54	59	62	61	55	56	67
Cocaine	18	16	15	16	13	13	13	13	11	12	7	9	10
Crack	9	8	5	7	4	5	3	4	2	2	*	1	4
Heroin	5	6	5	5	5	2	6	5	3	8	*	5	6
Hallucinogens	20	20	19	19	18	17	21	20	21	22	17	17	19
LSD	12	10	8	8	7	8	10	7	10	8	6	10	14
PCP	6	6	5	6	5	5	4	5	3	2	3	3	2
Ecstasy	16	12	13	11	11	11	15	16	15	15	11	9	10
Inhalants	22	25	26	23	26	26	24	22	23	20	19	15	19
Nonmedical Use of Psychotherapeutics ^{2,3}	43	42	40	37	40	40	33	36	38	35	37	32	41
Pain Relievers	39	39	36	33	36	36	29	34	35	31	32	29	33
OxyContin [®]	--	--	10	7	6	8	8	11	10	10	10	7	9
Tranquilizers	18	22	15	17	14	19	16	15	15	17	19	12	16
Stimulants ³	22	19	19	18	17	14	14	14	15	12	14	12	23
Methamphetamine ³	14	11	12	13	10	8	6	8	8	8	6	6	12
Sedatives	9	10	6	6	8	9	8	9	8	6	6	6	10
ILLCIT DRUGS OTHER THAN MARIJUANA¹	52	48	49	44	48	49	42	45	47	42	43	37	46

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.6D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.29	0.27	0.27	0.25	0.27	0.27	0.25	0.27	0.29	0.27	0.25	0.25	0.30
Marijuana and Hashish	0.24	0.24	0.23	0.22	0.21	0.22	0.22	0.24	0.25	0.24	0.22	0.23	0.27
Cocaine	0.07	0.06	0.06	0.06	0.05	0.05	0.05	0.05	0.05	0.05	0.03	0.04	0.04
Crack	0.03	0.03	0.02	0.03	0.02	0.02	0.01	0.02	0.01	0.01	*	0.01	0.02
Heroin	0.02	0.02	0.02	0.02	0.02	0.01	0.03	0.02	0.01	0.03	*	0.02	0.02
Hallucinogens	0.08	0.08	0.07	0.07	0.07	0.07	0.08	0.08	0.09	0.09	0.07	0.07	0.08
LSD	0.05	0.04	0.03	0.03	0.03	0.03	0.04	0.03	0.04	0.03	0.02	0.04	0.06
PCP	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01
Ecstasy	0.06	0.05	0.05	0.04	0.04	0.04	0.06	0.06	0.06	0.06	0.04	0.04	0.04
Inhalants	0.09	0.10	0.10	0.09	0.10	0.10	0.10	0.09	0.10	0.08	0.08	0.06	0.07
Nonmedical Use of Psychotherapeutics ^{2,3}	0.17	0.17	0.16	0.14	0.16	0.16	0.13	0.15	0.16	0.14	0.15	0.13	0.16
Pain Relievers	0.16	0.15	0.14	0.13	0.14	0.14	0.12	0.14	0.14	0.12	0.13	0.11	0.13
OxyContin [®]	--	--	0.04	0.03	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.03	0.04
Tranquilizers	0.07	0.09	0.06	0.07	0.06	0.07	0.07	0.06	0.06	0.07	0.08	0.05	0.06
Stimulants ³	0.09	0.08	0.08	0.07	0.07	0.06	0.05	0.06	0.06	0.05	0.06	0.05	0.09
Methamphetamine ³	0.06	0.04	0.05	0.05	0.04	0.03	0.02	0.03	0.03	0.03	0.03	0.02	0.05
Sedatives	0.03	0.04	0.02	0.03	0.03	0.04	0.03	0.04	0.03	0.02	0.03	0.03	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.21	0.19	0.20	0.17	0.19	0.19	0.17	0.18	0.19	0.17	0.17	0.15	0.19

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.6P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0025	0.1943	0.3121	0.6701	0.7768	0.0811	0.0741	0.0805	0.7306	0.1397	
Marijuana and Hashish	0.0229	0.1447	0.5459	0.1251	0.0478	0.0624	0.0575	0.9779	0.9087	0.1523	0.6864	0.4269	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0004	0.0527	0.1445	0.0739	0.4394	0.7663	
Crack	0.0565	0.0052	0.2403	0.0428	0.4626	0.2157	0.9229	0.9231	0.3706	0.3297	*	0.2562	
Heroin	0.5650	0.8256	0.8972	0.8728	0.9875	0.0462	0.7850	0.7246	0.2808	0.8834	*	0.6692	
Hallucinogens	0.0000	0.0000	0.0072	0.0146	0.1057	0.1769	0.0001	0.0028	0.0043	0.0068	0.3670	0.4918	
LSD	0.8213	0.4051	0.0932	0.0412	0.0316	0.0286	0.4486	0.0601	0.2975	0.0550	0.0127	0.4291	
PCP	0.0017	0.0069	0.1482	0.0247	0.1004	0.2425	0.0154	0.0785	0.3500	0.7534	0.7140	0.3177	
Ecstasy	0.0000	0.0014	0.0227	0.0279	0.0058	0.0974	0.0003	0.0000	0.0000	0.0002	0.0259	0.5023	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0009	0.0001	0.0071	0.1456	0.2319	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0000	0.0000	0.0000	0.0021	0.0022	0.0043	0.2916	0.0201	0.1456	0.5659	0.5095	0.0396	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0255	0.0000	0.0017	0.0392	0.0509	0.3296	
OxyContin [®]	--	--	0.0304	0.9090	0.5108	0.3440	0.2409	0.0137	0.3449	0.2356	0.3149	0.6153	
Tranquilizers	0.0004	0.0000	0.0122	0.0205	0.3694	0.0024	0.0410	0.0575	0.1405	0.0136	0.1542	0.5092	
Stimulants ³	0.1942	0.0770	0.7644	0.9897	0.9677	0.0965	0.0340	0.1336	0.0305	0.0059	0.0569	0.0010	
Methamphetamine ³	0.1616	0.1239	0.4568	0.1440	0.6626	0.3946	0.2193	0.5367	0.1647	0.5453	0.2638	0.0868	
Sedatives	0.5967	0.6569	0.2139	0.2309	0.7852	0.6193	0.6603	0.9963	0.7035	0.1658	0.2367	0.1931	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0010	0.0001	0.0006	0.0273	0.6160	0.0305	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.7C Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	156	155	165	161	166	162	155	163	167	174	175	168	190
Marijuana and Hashish	157	157	176	168	175	166	160	169	167	177	178	170	197
Cocaine	114	107	109	115	116	117	112	113	112	108	113	108	120
Crack	51	54	50	54	55	49	48	45	47	42	40	40	40
Heroin	35	37	36	36	38	34	32	37	41	43	43	44	48
Hallucinogens	140	134	130	132	140	130	118	135	134	121	131	133	139
LSD	115	108	98	96	89	78	74	80	79	74	87	81	92
PCP	45	49	42	41	38	35	36	37	34	32	30	30	31
Ecstasy	112	109	108	113	112	111	100	104	115	102	117	114	120
Inhalants	111	104	111	102	101	96	92	95	91	95	84	85	93
Nonmedical Use of Psychotherapeutics ^{2,3}	135	136	143	143	154	145	142	153	143	145	151	149	165
Pain Relievers	119	130	129	131	145	136	138	139	134	130	138	133	146
OxyContin [®]	40	56	54	64	65	62	68	71	75	74	77	74	78
Tranquilizers	94	98	98	107	112	104	103	104	106	104	109	108	121
Stimulants ³	93	101	104	97	104	97	91	97	97	93	100	98	100
Methamphetamine ³	71	76	72	77	79	71	63	62	62	56	62	57	58
Sedatives	39	37	40	40	38	40	37	40	34	36	36	37	40
ILLCIT DRUGS OTHER THAN MARIJUANA¹	157	151	157	154	166	158	156	163	158	157	160	161	174

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.7D Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.50	0.49	0.51	0.49	0.51	0.49	0.47	0.49	0.49	0.51	0.51	0.48	0.54
Marijuana and Hashish	0.51	0.50	0.55	0.52	0.53	0.51	0.49	0.50	0.49	0.51	0.51	0.49	0.56
Cocaine	0.37	0.34	0.34	0.35	0.36	0.36	0.34	0.34	0.33	0.32	0.33	0.31	0.34
Crack	0.16	0.17	0.16	0.17	0.17	0.15	0.15	0.13	0.14	0.12	0.12	0.11	0.11
Heroin	0.11	0.12	0.11	0.11	0.12	0.11	0.10	0.11	0.12	0.13	0.12	0.13	0.14
Hallucinogens	0.45	0.42	0.40	0.41	0.43	0.40	0.36	0.40	0.39	0.35	0.38	0.38	0.40
LSD	0.37	0.34	0.31	0.30	0.27	0.24	0.22	0.24	0.23	0.22	0.25	0.23	0.26
PCP	0.15	0.16	0.13	0.13	0.12	0.11	0.11	0.11	0.10	0.09	0.09	0.09	0.09
Ecstasy	0.36	0.34	0.34	0.35	0.34	0.34	0.30	0.31	0.34	0.30	0.34	0.33	0.34
Inhalants	0.36	0.33	0.34	0.31	0.31	0.29	0.28	0.28	0.27	0.28	0.24	0.24	0.27
Nonmedical Use of Psychotherapeutics ^{2,3}	0.44	0.43	0.44	0.44	0.47	0.44	0.43	0.46	0.42	0.42	0.44	0.43	0.47
Pain Relievers	0.39	0.41	0.40	0.40	0.44	0.42	0.42	0.42	0.39	0.38	0.40	0.38	0.42
OxyContin [®]	0.13	0.18	0.17	0.20	0.20	0.19	0.21	0.21	0.22	0.22	0.22	0.21	0.22
Tranquilizers	0.30	0.31	0.31	0.33	0.34	0.32	0.31	0.31	0.31	0.30	0.32	0.31	0.35
Stimulants ³	0.30	0.32	0.32	0.30	0.32	0.30	0.28	0.29	0.28	0.27	0.29	0.28	0.29
Methamphetamine ³	0.23	0.24	0.23	0.24	0.24	0.22	0.19	0.18	0.18	0.16	0.18	0.17	0.17
Sedatives	0.13	0.12	0.12	0.12	0.12	0.12	0.11	0.12	0.10	0.11	0.10	0.11	0.11
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.51	0.47	0.49	0.48	0.51	0.48	0.47	0.49	0.47	0.46	0.46	0.46	0.50

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.7P Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0083	0.0003	0.0877	0.0773	0.1177	0.6453	0.2215	0.3591	0.4714	0.2150	0.8737	0.2296	
Marijuana and Hashish	0.1388	0.0927	0.8871	0.7830	0.8303	0.0204	0.0116	1.0000	0.1114	0.3203	0.5543	0.3159	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0058	0.0096	0.2725	
Crack	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0022	0.0451	0.7324	
Heroin	0.0310	0.0684	0.0814	0.0149	0.0328	0.0128	0.0015	0.2443	0.4098	0.2327	0.8524	0.3514	
Hallucinogens	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0168	0.0000	0.0006	0.0551	0.0486	0.0590	
LSD	0.0000	0.0000	0.0000	0.0000	0.0000	0.4698	0.2068	0.8002	0.1006	0.0027	0.0034	0.1116	
PCP	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0010	0.2430	0.3173	0.5934	
Ecstasy	0.0000	0.0000	0.0003	0.0005	0.0043	0.1131	0.7500	0.3139	0.4152	0.5311	0.0795	0.1242	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.1541	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0023	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.1156	0.0068	0.7131	
Pain Relievers	0.0008	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0001	0.2409	
OxyContin [®]	0.0000	0.0000	0.0031	0.6642	0.8932	0.6355	0.4048	0.0001	0.0001	0.0022	0.0255	0.2930	
Tranquilizers	0.0052	0.6674	0.4963	0.0545	0.2141	0.6578	0.7767	0.4062	0.1171	0.7447	0.5330	0.3268	
Stimulants ³	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.4658	0.0515	0.2446	
Methamphetamine ³	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.1067	0.4834	0.9414	
Sedatives	0.0000	0.0047	0.0002	0.0001	0.0008	0.0051	0.1429	0.1219	0.6876	0.6957	0.5185	0.7336	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.3573	0.0453	0.8604	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.8C Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	143	153	154	152	157	156	154	160	167	172	174	166	185
Marijuana and Hashish	135	146	152	145	150	150	150	156	166	167	171	163	186
Cocaine	74	72	80	74	80	78	70	69	69	72	74	72	81
Crack	23	23	24	29	27	24	23	20	22	16	18	16	22
Heroin	17	14	15	21	18	20	19	20	25	26	29	27	31
Hallucinogens	81	77	72	74	79	77	75	82	90	81	89	89	90
LSD	44	30	27	30	32	30	36	40	45	43	49	50	56
PCP	14	19	17	11	14	11	8	8	10	16	11	7	9
Ecstasy	67	50	48	52	61	57	60	65	70	65	69	66	73
Inhalants	41	39	41	42	40	38	36	45	45	47	38	41	37
Nonmedical Use of Psychotherapeutics ^{2,3}	97	104	110	109	118	110	112	119	114	110	111	110	115
Pain Relievers	86	97	94	98	103	97	101	103	100	94	97	97	90
OxyContin [®]	--	--	35	37	39	36	37	43	44	43	47	42	37
Tranquilizers	57	66	66	68	75	69	70	70	72	63	71	70	71
Stimulants ³	57	56	66	58	61	61	54	61	59	55	72	62	71
Methamphetamine ³	41	39	41	41	38	35	26	29	28	25	43	31	35
Sedatives	18	18	18	22	22	21	23	22	21	18	18	21	22
ILLCIT DRUGS OTHER THAN MARIJUANA¹	116	127	126	121	130	126	123	131	132	129	127	128	135

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.8D Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.46	0.48	0.48	0.47	0.48	0.48	0.47	0.48	0.49	0.50	0.50	0.48	0.53
Marijuana and Hashish	0.43	0.46	0.47	0.45	0.46	0.46	0.46	0.46	0.49	0.49	0.49	0.47	0.53
Cocaine	0.24	0.23	0.25	0.23	0.24	0.24	0.21	0.20	0.20	0.21	0.21	0.21	0.23
Crack	0.08	0.07	0.07	0.09	0.08	0.07	0.07	0.06	0.06	0.05	0.05	0.05	0.06
Heroin	0.05	0.04	0.05	0.06	0.06	0.06	0.06	0.06	0.07	0.07	0.08	0.08	0.09
Hallucinogens	0.26	0.24	0.22	0.23	0.24	0.24	0.23	0.24	0.27	0.23	0.26	0.25	0.26
LSD	0.14	0.09	0.08	0.09	0.10	0.09	0.11	0.12	0.13	0.13	0.14	0.14	0.16
PCP	0.04	0.06	0.05	0.03	0.04	0.03	0.02	0.02	0.03	0.05	0.03	0.02	0.02
Ecstasy	0.22	0.16	0.15	0.16	0.19	0.18	0.18	0.19	0.20	0.19	0.20	0.19	0.21
Inhalants	0.13	0.12	0.13	0.13	0.12	0.12	0.11	0.14	0.13	0.14	0.11	0.12	0.11
Nonmedical Use of Psychotherapeutics ^{2,3}	0.31	0.33	0.34	0.33	0.36	0.34	0.34	0.35	0.33	0.32	0.32	0.32	0.33
Pain Relievers	0.28	0.30	0.29	0.30	0.31	0.30	0.31	0.31	0.29	0.27	0.28	0.28	0.26
OxyContin [®]	--	--	0.11	0.11	0.12	0.11	0.11	0.13	0.13	0.13	0.14	0.12	0.11
Tranquilizers	0.18	0.21	0.20	0.21	0.23	0.21	0.21	0.21	0.21	0.18	0.21	0.20	0.20
Stimulants ³	0.18	0.18	0.20	0.18	0.19	0.18	0.16	0.18	0.17	0.16	0.21	0.18	0.20
Methamphetamine ³	0.13	0.12	0.13	0.13	0.12	0.11	0.08	0.09	0.08	0.07	0.12	0.09	0.10
Sedatives	0.06	0.06	0.06	0.07	0.07	0.06	0.07	0.06	0.06	0.05	0.05	0.06	0.06
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.37	0.40	0.39	0.37	0.40	0.38	0.37	0.39	0.39	0.38	0.37	0.37	0.39

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.8P Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.3890	0.0407	0.0025	0.0093	0.0212	0.0001	0.0008	0.7475	0.2278	0.2272	0.7579	0.7006	
Marijuana and Hashish	0.0018	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.1101	0.0070	0.1082	0.5319	0.6654	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0015	0.0171	0.7810	0.9159	0.9105	0.6188	
Crack	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0016	0.0893	0.0675	0.5119	0.5915	0.2483	
Heroin	0.0003	0.0000	0.0001	0.0104	0.0023	0.0019	0.0029	0.0363	0.1958	0.4136	0.8635	0.5789	
Hallucinogens	0.0000	0.3348	0.4225	0.9938	0.4656	0.8321	0.8458	0.0097	0.0225	0.1888	0.6760	0.2608	
LSD	0.0175	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0002	0.0011	0.0044	0.0215	0.2399	
PCP	0.0001	0.0001	0.0003	0.0774	0.0076	0.0910	0.7491	0.9505	0.8603	0.1296	0.1248	0.3729	
Ecstasy	0.0000	0.4915	0.1320	0.1124	0.3103	0.9584	0.1828	0.0029	0.0018	0.0261	0.0397	0.0926	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0008	0.0003	0.0000	0.0001	0.0331	0.0662	0.0698	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0513	0.0001	0.4133	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0124	
OxyContin [®]	--	--	0.0006	0.0013	0.0008	0.0008	0.0001	0.0000	0.0000	0.0002	0.0150	0.3930	
Tranquilizers	0.0083	0.0001	0.0005	0.0000	0.0000	0.0003	0.0000	0.0000	0.0001	0.1536	0.0044	0.1980	
Stimulants ³	0.7557	0.6652	0.7680	0.8028	0.2581	0.1189	0.0669	0.4582	0.3629	0.0027	0.8813	0.5864	
Methamphetamine ³	0.0000	0.0000	0.0000	0.0000	0.0000	0.1042	0.1535	0.4531	0.0994	0.0055	0.9251	0.5185	
Sedatives	0.3234	0.5253	0.4320	0.1038	0.1669	0.3447	0.2747	0.6696	0.5910	0.4121	0.2402	0.6307	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.1082	0.0076	0.5950	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.9C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	116	128	130	130	122	127	129	135	136	142	138	143	160
Marijuana and Hashish	110	116	119	121	115	120	123	126	130	135	133	135	157
Cocaine	37	42	43	47	41	38	39	36	39	41	33	34	40
Crack	9	13	13	16	13	11	11	11	16	6	9	12	9
Heroin	10	7	9	10	12	11	14	12	17	20	22	16	16
Hallucinogens	38	37	33	36	43	35	39	46	48	40	46	48	47
LSD	8	13	14	12	12	12	15	16	17	14	18	18	19
PCP	6	10	5	5	7	6	3	3	3	6	4	2	3
Ecstasy	31	23	23	26	34	24	31	33	37	30	38	31	35
Inhalants	18	19	18	21	17	19	18	21	21	36	22	18	16
Nonmedical Use of Psychotherapeutics ^{2,3}	60	71	69	72	77	69	71	74	72	69	74	69	71
Pain Relievers	51	61	60	61	66	60	61	62	64	58	65	59	57
OxyContin [®]	--	--	19	19	17	18	16	19	24	21	24	19	20
Tranquilizers	35	36	38	40	43	39	40	40	39	40	42	34	36
Stimulants ³	31	30	34	37	34	31	29	34	35	33	31	37	36
Methamphetamine ³	21	21	24	27	21	18	14	15	17	16	16	18	15
Sedatives	9	11	10	15	13	13	15	17	12	10	13	9	15
ILLCIT DRUGS OTHER THAN MARIJUANA¹	75	82	82	87	89	81	82	88	84	87	85	82	91

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.9D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.37	0.40	0.40	0.40	0.37	0.39	0.39	0.40	0.40	0.42	0.40	0.41	0.46
Marijuana and Hashish	0.36	0.37	0.37	0.37	0.35	0.37	0.37	0.38	0.38	0.39	0.39	0.39	0.45
Cocaine	0.12	0.13	0.13	0.15	0.13	0.12	0.12	0.11	0.11	0.12	0.09	0.10	0.11
Crack	0.03	0.04	0.04	0.05	0.04	0.03	0.03	0.03	0.05	0.02	0.03	0.03	0.03
Heroin	0.03	0.02	0.03	0.03	0.04	0.03	0.04	0.04	0.05	0.06	0.06	0.05	0.05
Hallucinogens	0.12	0.12	0.10	0.11	0.13	0.11	0.12	0.14	0.14	0.12	0.13	0.14	0.13
LSD	0.03	0.04	0.04	0.04	0.04	0.04	0.05	0.05	0.05	0.04	0.05	0.05	0.05
PCP	0.02	0.03	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.02	0.01	0.01	0.01
Ecstasy	0.10	0.07	0.07	0.08	0.10	0.07	0.09	0.10	0.11	0.09	0.11	0.09	0.10
Inhalants	0.06	0.06	0.06	0.06	0.05	0.06	0.05	0.06	0.06	0.10	0.06	0.05	0.05
Nonmedical Use of Psychotherapeutics ^{2,3}	0.19	0.22	0.21	0.22	0.23	0.21	0.22	0.22	0.21	0.20	0.21	0.20	0.20
Pain Relievers	0.16	0.19	0.19	0.19	0.20	0.18	0.19	0.19	0.19	0.17	0.19	0.17	0.16
OxyContin [®]	--	--	0.06	0.06	0.05	0.06	0.05	0.06	0.07	0.06	0.07	0.05	0.06
Tranquilizers	0.11	0.11	0.12	0.12	0.13	0.12	0.12	0.12	0.11	0.12	0.12	0.10	0.10
Stimulants ³	0.10	0.10	0.11	0.11	0.10	0.09	0.09	0.10	0.10	0.10	0.09	0.11	0.10
Methamphetamine ³	0.07	0.07	0.07	0.08	0.06	0.06	0.04	0.05	0.05	0.05	0.05	0.05	0.04
Sedatives	0.03	0.04	0.03	0.05	0.04	0.04	0.05	0.05	0.04	0.03	0.04	0.03	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.24	0.26	0.26	0.27	0.27	0.25	0.25	0.26	0.25	0.25	0.25	0.24	0.26

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.9P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0027	0.0056	0.0000	0.0027	0.0003	0.0003	0.0002	0.3225	0.4928	0.3817	0.2964	0.4747	
Marijuana and Hashish	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0248	0.0774	0.3621	0.1662	0.4081	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0161	0.2065	0.8293	0.4503	1.0000	0.1718	0.1629	
Crack	0.0335	0.0724	0.0007	0.0000	0.0034	0.0360	0.0604	0.3914	0.0184	0.4429	0.4875	0.6272	
Heroin	0.0497	0.0020	0.0555	0.3195	0.2487	0.1253	0.9267	0.5364	0.6764	0.2995	0.0375	0.6588	
Hallucinogens	0.0111	0.1157	0.7836	0.5213	0.1265	0.9238	0.2228	0.0321	0.0059	0.2914	0.1150	0.0780	
LSD	0.0002	0.0649	0.2420	0.0197	0.0144	0.0543	0.5636	0.4939	0.6813	0.2960	0.9653	0.6211	
PCP	0.0967	0.0161	0.0424	0.2519	0.1526	0.1058	0.9550	0.8775	0.7911	0.2579	0.1742	0.8628	
Ecstasy	0.0077	0.6974	0.6665	0.6922	0.1181	0.5657	0.3620	0.0111	0.0053	0.2673	0.0673	0.2725	
Inhalants	0.0007	0.0124	0.0033	0.0057	0.0187	0.0214	0.2378	0.0064	0.0314	0.0952	0.1150	0.2937	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0003	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0558	0.0039	0.2389	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0014	0.0001	0.0545	
OxyContin [®]	--	--	0.0755	0.0539	0.0805	0.0193	0.1674	0.0033	0.0000	0.0446	0.1556	0.8414	
Tranquilizers	0.0138	0.0026	0.0003	0.0000	0.0000	0.0005	0.0008	0.0000	0.0019	0.0102	0.0241	0.6807	
Stimulants ³	0.2690	0.3314	0.0265	0.0660	0.1056	0.5150	0.4843	0.4833	0.7138	0.2581	0.9632	0.2592	
Methamphetamine ³	0.0000	0.0000	0.0000	0.0000	0.0000	0.1238	0.7392	0.9511	0.9815	0.8207	0.5771	0.1459	
Sedatives	0.9469	0.6792	0.8488	0.2865	0.9382	0.3399	0.5063	0.1745	0.8094	0.4564	0.8340	0.2070	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0000	0.1566	0.0932	0.4192	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.10C Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	935	874	928	903	854	935	898	932	988	1,011	925	978	818
Marijuana and Hashish	888	831	875	887	863	921	887	907	986	993	967	986	811
Cocaine	574	552	621	580	635	655	660	646	663	704	683	674	611
Crack	303	257	274	283	322	337	317	314	371	324	344	359	313
Heroin	220	201	188	223	235	236	221	214	243	223	241	278	207
Hallucinogens	572	549	583	573	613	615	631	646	651	618	641	678	601
LSD	496	469	494	499	508	540	542	557	549	526	538	567	491
PCP	282	262	292	284	290	294	337	292	317	303	309	342	273
Ecstasy	247	255	286	294	322	317	319	352	396	344	374	442	345
Inhalants	457	407	448	436	445	488	468	463	458	479	471	505	412
Nonmedical Use of Psychotherapeutics ^{2,3}	715	629	706	693	663	662	730	705	752	740	740	751	622
Pain Relievers	539	477	529	527	497	526	567	544	579	589	589	616	520
OxyContin [®]	103	125	120	120	152	165	171	203	224	196	228	268	201
Tranquilizers	481	414	471	460	451	453	484	472	479	483	540	558	424
Stimulants ³	495	456	480	474	482	459	536	496	538	486	539	486	425
Methamphetamine ³	402	392	394	361	398	359	397	395	439	404	418	386	356
Sedatives	366	316	396	344	347	357	395	351	353	349	374	376	316
ILLCIT DRUGS OTHER THAN MARIJUANA¹	789	728	788	797	772	830	828	845	863	879	842	874	769

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.10D Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.52	0.48	0.51	0.49	0.45	0.49	0.47	0.48	0.51	0.51	0.46	0.48	0.40
Marijuana and Hashish	0.50	0.46	0.48	0.48	0.46	0.48	0.46	0.47	0.50	0.50	0.48	0.49	0.39
Cocaine	0.32	0.30	0.34	0.31	0.34	0.35	0.34	0.33	0.34	0.35	0.34	0.33	0.30
Crack	0.17	0.14	0.15	0.15	0.17	0.18	0.17	0.16	0.19	0.16	0.17	0.18	0.15
Heroin	0.12	0.11	0.10	0.12	0.12	0.12	0.12	0.11	0.12	0.11	0.12	0.14	0.10
Hallucinogens	0.32	0.30	0.32	0.31	0.33	0.32	0.33	0.33	0.33	0.31	0.32	0.33	0.29
LSD	0.28	0.26	0.27	0.27	0.27	0.28	0.28	0.29	0.28	0.27	0.27	0.28	0.24
PCP	0.16	0.14	0.16	0.15	0.15	0.15	0.18	0.15	0.16	0.15	0.15	0.17	0.13
Ecstasy	0.14	0.14	0.16	0.16	0.17	0.17	0.17	0.18	0.20	0.17	0.19	0.22	0.17
Inhalants	0.25	0.22	0.24	0.24	0.24	0.26	0.24	0.24	0.23	0.24	0.23	0.25	0.20
Nonmedical Use of Psychotherapeutics ^{2,3}	0.40	0.35	0.39	0.37	0.35	0.35	0.38	0.36	0.39	0.37	0.37	0.37	0.30
Pain Relievers	0.30	0.26	0.29	0.28	0.26	0.28	0.30	0.28	0.30	0.30	0.29	0.30	0.25
OxyContin [®]	0.06	0.07	0.07	0.06	0.08	0.09	0.09	0.11	0.11	0.10	0.11	0.13	0.10
Tranquilizers	0.27	0.23	0.26	0.25	0.24	0.24	0.25	0.24	0.25	0.24	0.27	0.28	0.21
Stimulants ³	0.28	0.25	0.26	0.26	0.26	0.24	0.28	0.26	0.28	0.24	0.27	0.24	0.21
Methamphetamine ³	0.22	0.22	0.22	0.19	0.21	0.19	0.21	0.20	0.22	0.20	0.21	0.19	0.17
Sedatives	0.20	0.17	0.22	0.19	0.18	0.19	0.21	0.18	0.18	0.18	0.19	0.19	0.15
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.44	0.40	0.43	0.43	0.41	0.44	0.43	0.44	0.44	0.44	0.42	0.43	0.37

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.10P Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0112	0.3587	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0044	0.4483	
Cocaine	0.0050	0.0611	0.0014	0.0000	0.0029	0.0223	0.1891	0.0424	0.2874	0.0877	0.2155	0.1591	
Crack	0.1019	0.0016	0.0009	0.0011	0.0523	0.1418	0.0610	0.0246	0.7608	0.0126	0.5004	0.3458	
Heroin	0.0852	0.0898	0.0000	0.0136	0.0773	0.0793	0.0540	0.0045	0.1901	0.1281	0.6155	0.7454	
Hallucinogens	0.0000	0.0000	0.0000	0.0000	0.0009	0.0000	0.0266	0.1063	0.1149	0.0014	0.0854	0.9291	
LSD	0.3951	0.9115	0.2920	0.0363	0.7824	0.4975	0.6524	0.8744	0.7340	0.2488	0.6337	0.6946	
PCP	0.0032	0.0693	0.2078	0.4792	0.3561	0.9242	0.3446	0.9753	0.9901	0.7228	0.5751	0.7424	
Ecstasy	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0101	0.0000	0.0025	0.8254	
Inhalants	0.9012	0.8050	0.8300	0.8819	0.8015	0.7429	0.6734	0.8619	0.6050	0.0244	0.3570	0.7675	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0577	0.0636	0.0210	0.0148	0.0812	0.0304	0.6326	0.1981	0.1476	0.0279	0.7631	0.5229	
Pain Relievers	0.0000	0.0000	0.0000	0.0002	0.0004	0.0002	0.2277	0.1409	0.0767	0.0211	0.4108	0.5751	
OxyContin [®]	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0000	0.0357	0.7152	
Tranquilizers	0.0001	0.0003	0.0000	0.0010	0.0019	0.0000	0.0047	0.0026	0.0027	0.0001	0.2138	0.2313	
Stimulants ³	0.0007	0.0146	0.2796	0.1228	0.5592	0.8817	0.4299	0.9262	0.6833	0.0159	0.2338	0.2760	
Methamphetamine ³	0.0000	0.0000	0.0041	0.3655	0.0422	0.7372	0.8793	0.7877	0.4827	0.1646	0.4262	0.3517	
Sedatives	0.0000	0.0000	0.0000	0.0003	0.0013	0.0337	0.0023	0.0145	0.6694	0.9707	0.4711	0.7537	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0001	0.0000	0.0000	0.0000	0.0000	0.0010	0.1797	0.0434	0.1037	0.0064	0.2506	0.3131	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.11C Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	514	489	473	493	494	529	497	526	572	550	583	622	515
Marijuana and Hashish	407	387	395	418	410	418	422	470	488	462	488	519	451
Cocaine	200	207	180	180	210	223	192	189	183	154	200	172	165
Crack	121	120	112	109	132	120	90	103	89	67	110	75	79
Heroin	52	45	52	36	83	50	53	76	70	66	72	62	68
Hallucinogens	109	109	132	112	113	105	95	122	129	106	138	177	120
LSD	35	28	29	31	43	42	28	32	47	35	65	65	44
PCP	16	6	18	10	34	12	*	18	20	9	30	19	10
Ecstasy	93	86	97	98	81	85	69	103	90	78	108	135	90
Inhalants	66	54	69	65	86	88	102	89	100	87	107	90	85
Nonmedical Use of Psychotherapeutics ^{2,3}	345	335	313	315	323	346	310	329	363	338	380	386	286
Pain Relievers	275	275	265	274	272	290	259	290	305	280	332	315	228
OxyContin [®]	--	--	61	69	82	85	79	90	103	91	90	107	72
Tranquilizers	210	175	189	189	180	208	184	203	194	197	234	215	167
Stimulants ³	129	115	136	114	163	129	103	132	132	135	141	165	140
Methamphetamine ³	99	91	116	89	131	102	69	96	93	115	101	106	96
Sedatives	113	81	69	69	94	88	54	73	101	59	63	81	67
ILLCIT DRUGS OTHER THAN MARIJUANA¹	384	375	364	354	367	407	364	375	397	370	423	431	332

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.11D Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.29	0.27	0.26	0.27	0.26	0.28	0.26	0.27	0.29	0.28	0.29	0.31	0.25
Marijuana and Hashish	0.23	0.21	0.22	0.23	0.22	0.22	0.22	0.24	0.25	0.23	0.24	0.26	0.22
Cocaine	0.11	0.11	0.10	0.10	0.11	0.12	0.10	0.10	0.09	0.08	0.10	0.08	0.08
Crack	0.07	0.07	0.06	0.06	0.07	0.06	0.05	0.05	0.05	0.03	0.05	0.04	0.04
Heroin	0.03	0.02	0.03	0.02	0.04	0.03	0.03	0.04	0.04	0.03	0.04	0.03	0.03
Hallucinogens	0.06	0.06	0.07	0.06	0.06	0.06	0.05	0.06	0.07	0.05	0.07	0.09	0.06
LSD	0.02	0.02	0.02	0.02	0.02	0.02	0.01	0.02	0.02	0.02	0.03	0.03	0.02
PCP	0.01	0.00	0.01	0.01	0.02	0.01	*	0.01	0.01	0.00	0.01	0.01	0.01
Ecstasy	0.05	0.05	0.05	0.05	0.04	0.04	0.04	0.05	0.05	0.04	0.05	0.07	0.04
Inhalants	0.04	0.03	0.04	0.03	0.05	0.05	0.05	0.05	0.05	0.04	0.05	0.04	0.04
Nonmedical Use of Psychotherapeutics ^{2,3}	0.19	0.18	0.17	0.17	0.17	0.18	0.16	0.17	0.19	0.17	0.19	0.19	0.14
Pain Relievers	0.15	0.15	0.14	0.15	0.14	0.15	0.13	0.15	0.16	0.14	0.17	0.16	0.11
OxyContin [®]	--	--	0.03	0.04	0.04	0.04	0.04	0.05	0.05	0.05	0.05	0.05	0.04
Tranquilizers	0.12	0.10	0.10	0.10	0.10	0.11	0.10	0.11	0.10	0.10	0.12	0.11	0.08
Stimulants ³	0.07	0.06	0.07	0.06	0.09	0.07	0.05	0.07	0.07	0.07	0.07	0.08	0.07
Methamphetamine ³	0.06	0.05	0.06	0.05	0.07	0.05	0.04	0.05	0.05	0.06	0.05	0.05	0.05
Sedatives	0.06	0.04	0.04	0.04	0.05	0.05	0.03	0.04	0.05	0.03	0.03	0.04	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.21	0.21	0.20	0.19	0.20	0.21	0.19	0.19	0.20	0.19	0.21	0.21	0.16

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.11P Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0059	0.0120	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0073	
Cocaine	0.0004	0.0003	0.0067	0.1456	0.0010	0.0094	0.0309	0.5394	0.8179	0.0051	0.4989	0.3133	
Crack	0.0000	0.0005	0.0016	0.0009	0.0004	0.0001	0.0185	0.0884	0.4398	0.2546	0.2645	0.3901	
Heroin	0.0001	0.0000	0.0001	0.0000	0.0693	0.0000	0.0002	0.0281	0.0318	0.0053	0.0160	0.0262	
Hallucinogens	0.1379	0.0116	0.1165	0.0170	0.0428	0.0072	0.0000	0.3288	0.2759	0.0028	0.7016	0.9728	
LSD	0.0357	0.0004	0.0087	0.0016	0.2261	0.0216	0.0042	0.0097	0.3048	0.0170	0.8566	0.8116	
PCP	0.4757	0.6219	0.3385	0.8419	0.3491	0.6558	*	0.2727	0.8496	0.8218	0.1391	0.5008	
Ecstasy	0.9111	0.0815	0.0639	0.2540	0.0209	0.0610	0.0005	0.7413	0.0943	0.0064	0.9457	0.7672	
Inhalants	0.0017	0.0002	0.0230	0.0032	0.2538	0.3576	0.2856	0.1576	0.3380	0.1827	0.3181	0.2517	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.9071	0.9013	0.2535	0.9067	0.1048	0.0685	0.4679	0.4023	0.2343	0.3342	0.0092	0.2962	
Pain Relievers	0.9122	0.4092	0.5891	0.4906	0.0118	0.0119	0.3215	0.0441	0.0277	0.7955	0.0007	0.1846	
OxyContin [®]	--	--	0.1703	0.2667	0.6555	0.8090	0.7897	0.5054	0.0265	0.1938	0.4383	0.1649	
Tranquilizers	0.5889	0.5045	0.6984	0.8308	0.2709	0.9830	0.2788	0.7354	0.5580	0.5879	0.0231	0.8107	
Stimulants ³	0.1432	0.0030	0.0911	0.0067	0.8531	0.0294	0.0000	0.0542	0.0028	0.0021	0.1075	0.8058	
Methamphetamine ³	0.2417	0.5316	0.0741	0.4927	0.0184	0.7596	0.0083	0.8786	0.1078	0.4434	0.4967	0.7781	
Sedatives	0.0617	0.3286	0.9171	0.8661	0.1320	0.2878	0.0963	0.5863	0.2121	0.0483	0.2549	0.5522	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.3566	0.6249	0.4038	0.6689	0.1934	0.0662	0.6956	0.4889	0.7042	0.0344	0.0291	0.5526	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.12C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	353	363	366	395	376	395	393	401	441	420	452	472	415
Marijuana and Hashish	279	284	312	323	311	309	342	340	362	374	392	407	361
Cocaine	119	137	110	129	147	143	122	127	99	96	138	129	98
Crack	66	90	64	85	101	78	55	76	58	41	94	63	54
Heroin	40	26	35	15	76	37	36	35	40	43	46	39	52
Hallucinogens	67	54	44	73	51	63	53	77	61	38	77	123	74
LSD	11	8	13	7	20	30	*	13	*	16	26	31	33
PCP	13	*	15	10	*	11	*	17	20	*	12	15	*
Ecstasy	53	46	37	34	36	49	38	61	44	30	61	85	56
Inhalants	56	32	55	48	64	59	65	48	77	54	56	59	59
Nonmedical Use of Psychotherapeutics ^{2,3}	216	243	190	218	222	237	204	228	252	215	247	244	207
Pain Relievers	172	193	152	179	199	200	177	198	196	178	199	196	162
OxyContin [®]	--	--	28	31	29	50	56	54	62	49	37	77	45
Tranquilizers	127	112	91	109	105	124	108	128	130	123	155	114	107
Stimulants ³	85	93	87	69	97	86	66	92	76	81	96	106	109
Methamphetamine ³	71	77	67	56	77	76	49	65	49	69	72	81	75
Sedatives	63	53	47	44	67	67	33	51	71	47	47	67	51
ILLCIT DRUGS OTHER THAN MARIJUANA¹	259	274	226	257	257	288	242	264	278	232	284	293	242

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.12D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.20	0.20	0.20	0.21	0.20	0.21	0.20	0.21	0.23	0.21	0.23	0.23	0.20
Marijuana and Hashish	0.16	0.16	0.17	0.17	0.17	0.16	0.18	0.18	0.19	0.19	0.20	0.20	0.18
Cocaine	0.07	0.08	0.06	0.07	0.08	0.08	0.06	0.07	0.05	0.05	0.07	0.06	0.05
Crack	0.04	0.05	0.03	0.05	0.05	0.04	0.03	0.04	0.03	0.02	0.05	0.03	0.03
Heroin	0.02	0.01	0.02	0.01	0.04	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03
Hallucinogens	0.04	0.03	0.02	0.04	0.03	0.03	0.03	0.04	0.03	0.02	0.04	0.06	0.04
LSD	0.01	0.00	0.01	0.00	0.01	0.02	*	0.01	*	0.01	0.01	0.02	0.02
PCP	0.01	*	0.01	0.01	*	0.01	*	0.01	0.01	*	0.01	0.01	*
Ecstasy	0.03	0.03	0.02	0.02	0.02	0.03	0.02	0.03	0.02	0.01	0.03	0.04	0.03
Inhalants	0.03	0.02	0.03	0.03	0.03	0.03	0.03	0.02	0.04	0.03	0.03	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{2,3}	0.12	0.13	0.10	0.12	0.12	0.12	0.11	0.12	0.13	0.11	0.12	0.12	0.10
Pain Relievers	0.10	0.11	0.08	0.10	0.11	0.11	0.09	0.10	0.10	0.09	0.10	0.10	0.08
OxyContin [®]	--	--	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.02	0.02	0.04	0.02
Tranquilizers	0.07	0.06	0.05	0.06	0.06	0.07	0.06	0.07	0.07	0.06	0.08	0.06	0.05
Stimulants ³	0.05	0.05	0.05	0.04	0.05	0.05	0.03	0.05	0.04	0.04	0.05	0.05	0.05
Methamphetamine ³	0.04	0.04	0.04	0.03	0.04	0.04	0.03	0.03	0.03	0.04	0.04	0.04	0.04
Sedatives	0.04	0.03	0.03	0.02	0.04	0.04	0.02	0.03	0.04	0.02	0.02	0.03	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.14	0.15	0.12	0.14	0.14	0.15	0.13	0.14	0.14	0.12	0.14	0.14	0.12

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.12P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0029	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	
Cocaine	0.0159	0.0010	0.0351	0.0013	0.0002	0.0067	0.0487	0.3518	0.6643	0.2816	0.1744	0.5229	
Crack	0.0092	0.0201	0.2754	0.0069	0.0035	0.0087	0.9401	0.1049	0.9850	0.1276	0.3778	0.7624	
Heroin	0.0034	0.0000	0.0010	0.0000	0.6523	0.0009	0.0017	0.0007	0.0050	0.0127	0.0531	0.0228	
Hallucinogens	0.2139	0.0058	0.0030	0.2943	0.0071	0.1154	0.0081	0.4517	0.0429	0.0001	0.1571	0.8146	
LSD	0.0142	0.0114	0.0246	0.0078	0.1332	0.2308	*	0.0362	*	0.0379	0.1342	0.5079	
PCP	*	*	*	*	*	*	*	*	*	*	*	*	
Ecstasy	0.3661	0.0864	0.0414	0.0704	0.0134	0.3303	0.0369	0.8639	0.0816	0.0057	0.2229	0.9620	
Inhalants	0.2226	0.0082	0.1878	0.0629	0.9776	0.2067	0.7629	0.0810	0.9343	0.5393	0.1160	0.6026	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.5205	0.3489	0.0152	0.2903	0.5683	0.6812	0.0842	0.9684	0.7036	0.1153	0.8127	0.9959	
Pain Relievers	0.3829	0.5829	0.0462	0.4815	0.2723	0.1801	0.9445	0.2332	0.3422	0.6935	0.4786	0.6333	
OxyContin [®]	--	--	0.1122	0.3820	0.0617	0.5903	0.4580	0.4227	0.3358	0.8345	0.3915	0.1185	
Tranquilizers	0.7261	0.4244	0.0157	0.2335	0.0747	0.2859	0.1908	0.9317	0.3130	0.2707	0.4629	0.3089	
Stimulants ³	0.1064	0.0947	0.0488	0.0020	0.2026	0.0054	0.0000	0.1195	0.0015	0.0006	0.0168	0.2122	
Methamphetamine ³	0.6394	0.4739	0.6978	0.4616	0.3671	0.8040	0.0213	0.7509	0.0479	0.3198	0.2605	0.8571	
Sedatives	0.0774	0.8179	0.7908	0.5366	0.3769	0.7633	0.1584	0.7341	0.4941	0.3759	0.6063	0.6503	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.9609	0.6512	0.0295	0.5851	0.3108	0.4048	0.3049	0.9843	0.7876	0.0171	0.8416	0.8856	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.13C Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	963	902	965	937	896	966	924	961	1,018	1,035	953	999	857
Marijuana and Hashish	915	865	918	924	903	952	915	940	1,016	1,019	1,002	1,008	860
Cocaine	584	564	635	597	657	660	674	654	680	704	699	692	627
Crack	307	263	280	293	329	339	323	316	378	328	346	364	316
Heroin	222	204	193	227	238	240	223	217	248	226	244	280	213
Hallucinogens	599	584	613	608	642	638	644	665	681	628	670	706	630
LSD	513	494	504	521	520	549	548	565	558	532	557	578	506
PCP	281	270	297	288	295	298	339	297	320	306	311	346	274
Ecstasy	296	281	326	334	355	344	342	371	427	363	405	463	381
Inhalants	472	435	471	456	466	496	483	470	468	487	486	515	424
Nonmedical Use of Psychotherapeutics ^{2,3}	737	648	731	721	696	692	753	733	777	752	759	778	657
Pain Relievers	565	510	559	560	533	558	593	577	606	597	614	636	554
OxyContin [®]	114	140	133	139	165	178	186	219	242	209	243	278	218
Tranquilizers	493	433	484	478	476	467	503	490	491	493	557	572	441
Stimulants ³	508	472	497	492	504	471	545	508	554	500	554	495	439
Methamphetamine ³	414	403	400	375	413	366	401	405	449	412	426	393	362
Sedatives	370	322	399	346	352	360	400	354	355	353	377	378	323
ILLCIT DRUGS OTHER THAN MARIJUANA¹	813	748	820	831	805	857	853	869	890	887	866	913	798

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.13D Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.46	0.42	0.45	0.43	0.41	0.43	0.41	0.42	0.44	0.44	0.41	0.42	0.36
Marijuana and Hashish	0.44	0.41	0.43	0.42	0.41	0.43	0.41	0.41	0.44	0.44	0.43	0.42	0.36
Cocaine	0.28	0.27	0.30	0.27	0.30	0.30	0.30	0.29	0.30	0.30	0.30	0.29	0.26
Crack	0.15	0.12	0.13	0.13	0.15	0.15	0.14	0.14	0.16	0.14	0.15	0.15	0.13
Heroin	0.11	0.10	0.09	0.10	0.11	0.11	0.10	0.10	0.11	0.10	0.10	0.12	0.09
Hallucinogens	0.28	0.27	0.28	0.28	0.29	0.29	0.29	0.29	0.30	0.27	0.28	0.30	0.26
LSD	0.24	0.23	0.23	0.24	0.24	0.25	0.24	0.25	0.24	0.23	0.24	0.24	0.21
PCP	0.13	0.13	0.14	0.13	0.13	0.13	0.15	0.13	0.14	0.13	0.13	0.15	0.11
Ecstasy	0.14	0.13	0.15	0.15	0.16	0.15	0.15	0.16	0.19	0.16	0.17	0.20	0.16
Inhalants	0.22	0.20	0.22	0.21	0.21	0.22	0.21	0.21	0.20	0.21	0.21	0.22	0.18
Nonmedical Use of Psychotherapeutics ^{2,3}	0.35	0.30	0.34	0.33	0.32	0.31	0.33	0.32	0.34	0.32	0.32	0.33	0.27
Pain Relievers	0.27	0.24	0.26	0.26	0.24	0.25	0.26	0.25	0.26	0.26	0.26	0.27	0.23
OxyContin [®]	0.05	0.07	0.06	0.06	0.07	0.08	0.08	0.10	0.11	0.09	0.10	0.12	0.09
Tranquilizers	0.23	0.20	0.22	0.22	0.22	0.21	0.22	0.22	0.21	0.21	0.24	0.24	0.18
Stimulants ³	0.24	0.22	0.23	0.23	0.23	0.21	0.24	0.22	0.24	0.22	0.24	0.21	0.18
Methamphetamine ³	0.20	0.19	0.19	0.17	0.19	0.16	0.18	0.18	0.20	0.18	0.18	0.17	0.15
Sedatives	0.18	0.15	0.19	0.16	0.16	0.16	0.18	0.16	0.15	0.15	0.16	0.16	0.13
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.39	0.35	0.38	0.38	0.37	0.38	0.38	0.38	0.39	0.38	0.37	0.38	0.33

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.13P Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0142	0.2681	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0048	0.3598	
Cocaine	0.2648	0.7393	0.1020	0.0063	0.2240	0.4177	0.9569	0.5519	0.8262	0.2139	0.4371	0.2311	
Crack	0.9679	0.1760	0.0791	0.0905	0.6617	0.6702	0.3421	0.2086	0.6974	0.0354	0.6542	0.3655	
Heroin	0.0358	0.0422	0.0000	0.0041	0.0326	0.0298	0.0131	0.0026	0.1506	0.0816	0.5966	0.8837	
Hallucinogens	0.1134	0.0738	0.0302	0.0014	0.0735	0.0013	0.0963	0.5533	0.4344	0.0070	0.2192	0.6374	
LSD	0.0014	0.0015	0.1797	0.6722	0.5445	0.5715	0.8176	0.9316	0.5243	0.1007	0.3280	0.9079	
PCP	0.0000	0.0004	0.0176	0.0729	0.0735	0.5914	0.1385	0.6328	0.7346	0.8041	0.5213	0.7140	
Ecstasy	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0406	0.0000	0.0283	0.5123	
Inhalants	0.0000	0.0000	0.0005	0.0011	0.0014	0.0111	0.0263	0.0271	0.2654	0.2681	0.9025	0.9802	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.2514	0.5673	0.3166	0.4552	0.9638	0.4537	0.5447	0.9208	0.6281	0.0792	0.3587	0.6047	
Pain Relievers	0.0000	0.0068	0.0155	0.2349	0.3887	0.1590	0.4745	0.6054	0.9683	0.1800	0.0778	0.8108	
OxyContin [®]	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0021	0.0339	0.0007	0.2442	0.9953	
Tranquilizers	0.0000	0.0004	0.0000	0.0080	0.0091	0.0000	0.0055	0.0077	0.0126	0.0002	0.2981	0.1701	
Stimulants ³	0.0000	0.0001	0.0087	0.8068	0.0363	0.3804	0.9915	0.3323	0.7671	0.0287	0.4448	0.4148	
Methamphetamine ³	0.0000	0.0000	0.0000	0.1263	0.0001	0.0390	0.3584	0.2252	0.2072	0.2499	0.4880	0.3594	
Sedatives	0.0000	0.0000	0.0000	0.0001	0.0005	0.0163	0.0017	0.0103	0.6661	0.9890	0.5333	0.7200	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0371	0.0346	0.0013	0.0123	0.0198	0.0574	0.6561	0.3619	0.4012	0.0132	0.4837	0.3529	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.14C Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	565	540	519	541	548	592	545	583	631	589	639	666	581
Marijuana and Hashish	459	437	437	463	466	484	466	527	548	503	549	565	528
Cocaine	217	225	202	202	227	239	207	208	202	167	217	189	184
Crack	124	122	115	114	133	123	94	105	92	68	111	77	81
Heroin	54	47	55	42	85	54	58	78	75	71	78	70	75
Hallucinogens	144	140	147	140	143	134	124	151	170	138	172	200	154
LSD	58	41	41	45	55	53	47	53	69	57	83	79	71
PCP	21	20	25	15	37	17	10	19	23	19	32	20	14
Ecstasy	122	101	108	111	102	104	92	123	121	103	132	152	117
Inhalants	80	66	78	79	94	96	108	101	109	99	115	101	88
Nonmedical Use of Psychotherapeutics ^{2,3}	368	356	341	345	354	377	345	360	389	357	398	415	314
Pain Relievers	293	298	289	300	298	312	288	318	329	297	349	339	246
OxyContin [®]	--	--	72	80	90	94	87	102	114	102	103	116	81
Tranquilizers	219	192	203	206	193	219	199	220	205	207	249	229	185
Stimulants ³	147	133	153	130	177	150	120	146	155	149	160	176	158
Methamphetamine ³	110	99	125	97	136	113	75	100	103	118	113	113	104
Sedatives	115	83	70	73	98	92	59	78	103	63	65	83	72
ILLCIT DRUGS OTHER THAN MARIJUANA¹	415	408	399	393	405	444	400	410	433	397	450	463	362

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.14D Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.27	0.25	0.24	0.25	0.25	0.27	0.24	0.26	0.28	0.25	0.27	0.28	0.24
Marijuana and Hashish	0.22	0.21	0.20	0.21	0.21	0.22	0.21	0.23	0.24	0.22	0.23	0.24	0.22
Cocaine	0.10	0.11	0.09	0.09	0.10	0.11	0.09	0.09	0.09	0.07	0.09	0.08	0.08
Crack	0.06	0.06	0.05	0.05	0.06	0.06	0.04	0.05	0.04	0.03	0.05	0.03	0.03
Heroin	0.03	0.02	0.03	0.02	0.04	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03
Hallucinogens	0.07	0.07	0.07	0.06	0.06	0.06	0.06	0.07	0.07	0.06	0.07	0.08	0.06
LSD	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.04	0.03	0.03
PCP	0.01	0.01	0.01	0.01	0.02	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01
Ecstasy	0.06	0.05	0.05	0.05	0.05	0.05	0.04	0.05	0.05	0.04	0.06	0.06	0.05
Inhalants	0.04	0.03	0.04	0.04	0.04	0.04	0.05	0.04	0.05	0.04	0.05	0.04	0.04
Nonmedical Use of Psychotherapeutics ^{2,3}	0.17	0.17	0.16	0.16	0.16	0.17	0.15	0.16	0.17	0.15	0.17	0.17	0.13
Pain Relievers	0.14	0.14	0.13	0.14	0.14	0.14	0.13	0.14	0.14	0.13	0.15	0.14	0.10
OxyContin [®]	--	--	0.03	0.04	0.04	0.04	0.04	0.04	0.05	0.04	0.04	0.05	0.03
Tranquilizers	0.10	0.09	0.09	0.09	0.09	0.10	0.09	0.10	0.09	0.09	0.11	0.10	0.08
Stimulants ³	0.07	0.06	0.07	0.06	0.08	0.07	0.05	0.06	0.07	0.06	0.07	0.07	0.07
Methamphetamine ³	0.05	0.05	0.06	0.04	0.06	0.05	0.03	0.04	0.04	0.05	0.05	0.05	0.04
Sedatives	0.05	0.04	0.03	0.03	0.04	0.04	0.03	0.03	0.04	0.03	0.03	0.03	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.20	0.19	0.19	0.18	0.18	0.20	0.18	0.18	0.19	0.17	0.19	0.19	0.15

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.14P Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0222	0.0193	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0141	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0013	0.1251	0.6854	0.0117	0.4811	0.2783	
Crack	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0020	0.0405	0.2223	0.2002	0.2277	0.2640	
Heroin	0.0000	0.0000	0.0000	0.0000	0.0084	0.0000	0.0000	0.0062	0.0150	0.0044	0.0334	0.0274	
Hallucinogens	0.0225	0.2350	0.1297	0.0973	0.3228	0.0331	0.0016	0.3974	0.5287	0.1587	0.9810	0.5166	
LSD	0.0026	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0037	0.0003	0.1722	0.3385	
PCP	0.0017	0.0008	0.0017	0.1566	0.0485	0.1413	0.3330	0.3157	0.7807	0.2729	0.0463	0.8354	
Ecstasy	0.0000	0.3975	0.0285	0.0977	0.2998	0.1683	0.0786	0.0376	0.4444	0.5624	0.2351	0.2493	
Inhalants	0.7305	0.8341	0.3808	0.7119	0.4380	0.5966	0.7415	0.4240	0.4292	0.8278	0.8065	0.7694	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0965	0.0279	0.1738	0.0156	0.0000	0.0001	0.1153	0.0015	0.0035	0.8665	0.0002	0.2110	
Pain Relievers	0.0028	0.0000	0.0014	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0414	0.0000	0.0403	
OxyContin [®]	--	--	0.5901	0.5570	0.2446	0.0990	0.0473	0.0005	0.0000	0.0049	0.0683	0.1048	
Tranquilizers	0.7214	0.3513	0.3325	0.1153	0.4519	0.1944	0.5889	0.0686	0.0398	0.9749	0.0022	0.4975	
Stimulants ³	0.2714	0.0076	0.2145	0.0175	0.4828	0.0117	0.0000	0.0482	0.0040	0.0001	0.1913	0.6808	
Methamphetamine ³	0.0012	0.0080	0.0003	0.0119	0.0003	0.4146	0.0035	0.7196	0.0506	0.1393	0.5235	0.6521	
Sedatives	0.0434	0.2625	0.8883	0.7264	0.0720	0.2103	0.2466	0.5182	0.1773	0.0379	0.1584	0.4854	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0018	0.0103	0.2613	0.0562	0.0003	0.0005	0.2217	0.0040	0.0637	0.1940	0.0031	0.4448	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.15C Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	379	396	404	430	416	440	422	447	482	453	485	503	462
Marijuana and Hashish	313	317	349	357	350	353	374	384	401	403	426	437	414
Cocaine	126	146	119	141	152	149	130	135	106	106	142	132	105
Crack	67	91	66	87	101	80	55	77	60	42	94	64	55
Heroin	41	27	36	18	77	38	39	37	43	48	51	42	55
Hallucinogens	78	68	55	82	68	72	65	89	80	56	89	131	84
LSD	13	15	22	14	23	32	16	21	17	21	31	36	38
PCP	14	11	16	11	7	13	*	17	21	10	12	15	*
Ecstasy	62	53	45	43	50	54	50	67	58	43	71	91	63
Inhalants	59	37	58	52	66	62	67	52	80	65	62	61	61
Nonmedical Use of Psychotherapeutics ^{2,3}	227	255	206	232	236	248	221	246	265	228	263	255	217
Pain Relievers	182	205	164	191	211	211	192	212	212	187	210	205	170
OxyContin [®]	--	--	35	38	34	53	57	58	67	54	45	80	49
Tranquilizers	133	116	102	116	114	129	117	135	137	132	161	120	113
Stimulants ³	92	100	96	79	103	93	74	98	84	88	101	116	114
Methamphetamine ³	75	80	75	60	79	79	52	67	53	71	75	85	77
Sedatives	64	54	48	46	68	68	36	54	72	48	49	66	53
ILLCIT DRUGS OTHER THAN MARIJUANA¹	270	286	247	278	275	303	258	287	296	251	305	305	255

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.15D Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.18	0.19	0.19	0.20	0.19	0.20	0.19	0.20	0.21	0.19	0.21	0.21	0.19
Marijuana and Hashish	0.15	0.15	0.16	0.16	0.16	0.16	0.17	0.17	0.18	0.17	0.18	0.18	0.17
Cocaine	0.06	0.07	0.06	0.06	0.07	0.07	0.06	0.06	0.05	0.05	0.06	0.06	0.04
Crack	0.03	0.04	0.03	0.04	0.05	0.04	0.02	0.03	0.03	0.02	0.04	0.03	0.02
Heroin	0.02	0.01	0.02	0.01	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Hallucinogens	0.04	0.03	0.03	0.04	0.03	0.03	0.03	0.04	0.03	0.02	0.04	0.06	0.04
LSD	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02
PCP	0.01	0.01	0.01	0.01	0.00	0.01	*	0.01	0.01	0.00	0.01	0.01	*
Ecstasy	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.02	0.03	0.04	0.03
Inhalants	0.03	0.02	0.03	0.02	0.03	0.03	0.03	0.02	0.04	0.03	0.03	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{2,3}	0.11	0.12	0.10	0.11	0.11	0.11	0.10	0.11	0.12	0.10	0.11	0.11	0.09
Pain Relievers	0.09	0.10	0.08	0.09	0.10	0.09	0.09	0.09	0.09	0.08	0.09	0.09	0.07
OxyContin [®]	--	--	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.02	0.02	0.03	0.02
Tranquilizers	0.06	0.05	0.05	0.05	0.05	0.06	0.05	0.06	0.06	0.06	0.07	0.05	0.05
Stimulants ³	0.04	0.05	0.04	0.04	0.05	0.04	0.03	0.04	0.04	0.04	0.04	0.05	0.05
Methamphetamine ³	0.04	0.04	0.03	0.03	0.04	0.04	0.02	0.03	0.02	0.03	0.03	0.04	0.03
Sedatives	0.03	0.03	0.02	0.02	0.03	0.03	0.02	0.02	0.03	0.02	0.02	0.03	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.13	0.13	0.11	0.13	0.12	0.14	0.11	0.13	0.13	0.11	0.13	0.13	0.11

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.15P Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0036	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	
Cocaine	0.0003	0.0000	0.0002	0.0000	0.0000	0.0009	0.0226	0.3328	0.9408	0.3359	0.3590	0.8494	
Crack	0.0041	0.0102	0.0878	0.0007	0.0010	0.0041	0.6788	0.0844	0.5799	0.1045	0.3404	0.7016	
Heroin	0.0008	0.0000	0.0002	0.0000	0.4996	0.0003	0.0028	0.0007	0.0127	0.0505	0.3063	0.0454	
Hallucinogens	0.7575	0.1818	0.0247	0.6060	0.2211	0.1994	0.1253	0.5813	0.9795	0.0129	0.7336	0.3199	
LSD	0.0001	0.0021	0.0183	0.0006	0.0122	0.0549	0.0088	0.0394	0.0103	0.0227	0.2176	0.4220	
PCP	*	*	*	*	*	*	*	*	*	*	*	*	
Ecstasy	0.5198	0.1046	0.0582	0.2167	0.3155	0.2633	0.2341	0.2279	0.7783	0.1203	0.9728	0.6653	
Inhalants	0.8615	0.1264	0.7066	0.4165	0.5103	0.5922	0.9792	0.4645	0.6202	0.8429	0.3313	0.8427	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.6144	0.4414	0.7909	0.2953	0.0061	0.0501	0.9791	0.0455	0.0648	0.3942	0.2593	0.7175	
Pain Relievers	0.3843	0.0611	0.3818	0.0545	0.0002	0.0004	0.0266	0.0003	0.0027	0.4887	0.0490	0.2866	
OxyContin [®]	--	--	0.6166	0.8868	0.3929	0.6991	0.2415	0.0672	0.0092	0.3137	0.9477	0.1227	
Tranquilizers	0.6951	0.8200	0.3550	0.7476	0.9566	0.9256	0.9340	0.2218	0.0547	0.8274	0.1756	0.4051	
Stimulants ³	0.2543	0.2163	0.2957	0.0326	0.5142	0.0047	0.0000	0.2237	0.0050	0.0003	0.0226	0.4230	
Methamphetamine ³	0.1317	0.0681	0.0802	0.3793	0.0638	0.9225	0.0217	0.7693	0.0572	0.3098	0.3304	0.6325	
Sedatives	0.0859	0.9062	0.7662	0.7979	0.3782	0.6085	0.2683	0.4664	0.4686	0.2983	0.5836	0.4819	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.1584	0.1739	0.7622	0.0952	0.0012	0.0267	0.7160	0.0772	0.2440	0.0931	0.4515	0.6814	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.16C Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	75	71	67	75	74	74	68	73	73	77	68	72	76
Marijuana and Hashish	64	63	61	64	63	65	59	64	67	64	62	62	66
Cocaine	26	24	24	24	21	23	19	21	21	20	16	15	17
Crack	11	13	10	9	10	8	7	7	9	13	7	4	6
Heroin	7	8	9	7	7	8	10	6	7	8	8	6	6
Hallucinogens	35	33	32	31	30	31	30	32	32	32	27	28	26
LSD	25	19	17	18	16	13	15	16	15	15	15	16	19
PCP	13	13	13	14	11	10	10	9	7	7	10	6	6
Ecstasy	27	21	20	17	19	21	20	25	24	27	20	19	18
Inhalants	46	49	47	48	47	46	44	47	42	42	36	40	37
Nonmedical Use of Psychotherapeutics ^{2,3}	51	49	48	51	52	52	45	50	48	46	46	47	48
Pain Relievers	47	47	46	47	48	47	41	48	46	41	42	43	42
OxyContin [®]	14	16	17	14	16	17	16	21	19	16	16	15	16
Tranquilizers	27	26	22	25	23	23	24	22	25	25	25	24	27
Stimulants ³	30	27	25	27	29	23	20	21	21	21	19	20	25
Methamphetamine ³	18	17	16	18	17	12	10	11	13	13	11	10	15
Sedatives	13	16	15	16	13	13	12	14	11	11	10	12	14
ILLCIT DRUGS OTHER THAN MARIJUANA¹	63	63	61	63	62	66	57	63	60	62	55	59	59

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.16D Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.59	0.56	0.52	0.58	0.57	0.57	0.54	0.58	0.59	0.60	0.53	0.57	0.60
Marijuana and Hashish	0.50	0.49	0.47	0.49	0.48	0.50	0.46	0.51	0.53	0.50	0.49	0.48	0.52
Cocaine	0.21	0.18	0.18	0.18	0.16	0.18	0.15	0.17	0.17	0.16	0.13	0.12	0.13
Crack	0.09	0.10	0.08	0.07	0.08	0.06	0.06	0.06	0.07	0.10	0.05	0.03	0.05
Heroin	0.05	0.06	0.07	0.06	0.05	0.06	0.08	0.05	0.06	0.07	0.07	0.04	0.05
Hallucinogens	0.27	0.26	0.25	0.24	0.23	0.24	0.23	0.26	0.25	0.25	0.21	0.22	0.20
LSD	0.20	0.15	0.13	0.14	0.12	0.10	0.12	0.13	0.12	0.12	0.12	0.13	0.15
PCP	0.10	0.10	0.10	0.11	0.08	0.08	0.08	0.07	0.05	0.06	0.08	0.05	0.04
Ecstasy	0.21	0.16	0.16	0.13	0.15	0.16	0.16	0.20	0.19	0.21	0.16	0.15	0.14
Inhalants	0.37	0.39	0.37	0.37	0.37	0.36	0.34	0.37	0.33	0.33	0.28	0.31	0.29
Nonmedical Use of Psychotherapeutics ^{2,3}	0.41	0.38	0.37	0.39	0.40	0.40	0.36	0.40	0.39	0.36	0.36	0.37	0.38
Pain Relievers	0.37	0.37	0.36	0.36	0.37	0.36	0.33	0.38	0.37	0.32	0.33	0.34	0.33
OxyContin [®]	0.11	0.12	0.13	0.11	0.12	0.13	0.12	0.16	0.16	0.12	0.13	0.12	0.13
Tranquilizers	0.21	0.20	0.17	0.19	0.18	0.18	0.19	0.17	0.20	0.20	0.20	0.19	0.21
Stimulants ³	0.24	0.21	0.19	0.21	0.22	0.18	0.16	0.16	0.17	0.17	0.15	0.16	0.19
Methamphetamine ³	0.14	0.13	0.13	0.14	0.13	0.09	0.08	0.09	0.10	0.10	0.09	0.07	0.12
Sedatives	0.10	0.13	0.12	0.12	0.10	0.10	0.10	0.11	0.09	0.09	0.08	0.09	0.11
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.50	0.50	0.47	0.49	0.47	0.51	0.45	0.50	0.48	0.49	0.43	0.46	0.47

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.16P Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0378	0.1455	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0452	0.1015	0.3804	0.3570	0.0055	0.0422	0.0057	0.1235	0.4576	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0011	0.0084	0.3187	0.4863	
Crack	0.0005	0.0000	0.0002	0.0054	0.0035	0.0123	0.1110	0.2420	0.1077	0.1667	0.8397	0.7764	
Heroin	0.0826	0.1030	0.0257	0.1607	0.4788	0.3186	0.1071	0.4444	0.5995	0.1597	0.3847	0.9234	
Hallucinogens	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0000	0.0000	0.0000	0.0001	0.0708	0.0485	
LSD	0.0000	0.1043	0.6795	0.6282	0.1157	0.0270	0.6298	0.4492	0.1180	0.3233	0.1683	0.5534	
PCP	0.0000	0.0000	0.0000	0.0000	0.0133	0.0067	0.0014	0.0323	0.5419	0.0566	0.0459	0.9274	
Ecstasy	0.0000	0.0001	0.0008	0.2674	0.0684	0.0238	0.0069	0.0000	0.0000	0.0000	0.0186	0.0503	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0037	0.1107	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0028	0.0000	0.0227	0.0293	0.0344	0.5446	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.0000	0.0006	0.0051	0.0048	0.4647	
OxyContin [®]	0.2973	0.6521	0.2861	0.9348	0.3402	0.1369	0.2799	0.0326	0.0399	0.1992	0.3368	0.8811	
Tranquilizers	0.0108	0.0043	0.0929	0.2016	0.5583	0.4077	0.4326	0.6156	0.2575	0.5130	0.6065	0.7315	
Stimulants ³	0.0000	0.0000	0.0000	0.0005	0.0000	0.0498	0.6166	0.8821	0.9349	0.9792	0.3391	0.9201	
Methamphetamine ³	0.0018	0.0003	0.0057	0.0198	0.0418	0.8622	0.5506	0.4870	0.9481	0.8272	0.3603	0.3185	
Sedatives	0.9108	0.0733	0.0545	0.3205	0.6580	0.9379	0.8498	0.3925	0.6356	0.2504	0.4368	0.3673	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0174	0.2573	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.17C Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	66	73	66	67	68	67	71	70	69	69	66	68	71
Marijuana and Hashish	59	62	59	55	57	56	60	57	56	60	58	60	62
Cocaine	24	26	24	23	24	23	25	18	18	18	16	13	19
Crack	13	10	12	10	12	10	10	6	6	9	7	3	6
Heroin	12	8	8	6	9	7	8	6	6	9	8	7	6
Hallucinogens	34	35	31	29	29	28	29	30	31	30	27	22	26
LSD	22	18	16	14	13	12	14	12	15	12	15	11	19
PCP	14	14	12	12	12	9	7	10	9	7	8	6	10
Ecstasy	28	24	22	20	22	21	25	23	28	26	23	17	20
Inhalants	45	45	50	49	45	47	48	45	44	43	38	34	41
Nonmedical Use of Psychotherapeutics ^{2,3}	52	54	49	50	50	51	51	48	48	46	46	44	51
Pain Relievers	48	50	47	46	45	49	47	44	45	42	43	39	44
OxyContin [®]	16	13	16	16	18	17	17	17	19	20	18	15	16
Tranquilizers	27	28	29	27	28	27	26	25	26	27	30	23	31
Stimulants ³	35	31	29	31	30	26	31	23	25	22	23	19	27
Methamphetamine ³	22	19	20	18	21	15	20	16	13	15	12	10	16
Sedatives	17	15	15	15	14	15	14	16	13	11	13	9	19
ILLCIT DRUGS OTHER THAN MARIJUANA¹	59	66	58	61	57	60	61	59	61	58	53	52	59

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.17D Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.54	0.60	0.54	0.54	0.55	0.54	0.58	0.58	0.58	0.57	0.54	0.56	0.58
Marijuana and Hashish	0.49	0.51	0.48	0.45	0.46	0.45	0.49	0.47	0.47	0.49	0.47	0.49	0.51
Cocaine	0.20	0.21	0.19	0.18	0.19	0.18	0.21	0.15	0.15	0.15	0.13	0.10	0.15
Crack	0.11	0.08	0.09	0.08	0.10	0.08	0.08	0.05	0.05	0.07	0.06	0.03	0.05
Heroin	0.10	0.07	0.06	0.05	0.07	0.06	0.06	0.05	0.05	0.07	0.07	0.05	0.05
Hallucinogens	0.28	0.28	0.25	0.23	0.23	0.22	0.24	0.25	0.26	0.24	0.22	0.18	0.21
LSD	0.18	0.15	0.13	0.11	0.10	0.10	0.12	0.10	0.12	0.10	0.12	0.09	0.15
PCP	0.12	0.12	0.10	0.10	0.09	0.08	0.06	0.08	0.08	0.06	0.06	0.05	0.08
Ecstasy	0.23	0.20	0.18	0.16	0.18	0.17	0.21	0.19	0.23	0.21	0.19	0.14	0.16
Inhalants	0.37	0.36	0.41	0.39	0.36	0.38	0.39	0.38	0.37	0.35	0.31	0.28	0.34
Nonmedical Use of Psychotherapeutics ^{2,3}	0.43	0.44	0.40	0.40	0.40	0.41	0.42	0.40	0.40	0.37	0.38	0.36	0.42
Pain Relievers	0.40	0.41	0.38	0.37	0.36	0.39	0.39	0.36	0.38	0.35	0.35	0.32	0.36
OxyContin [®]	0.13	0.10	0.13	0.13	0.14	0.14	0.14	0.14	0.16	0.16	0.15	0.13	0.13
Tranquilizers	0.23	0.23	0.23	0.22	0.22	0.22	0.22	0.21	0.22	0.22	0.24	0.19	0.25
Stimulants ³	0.29	0.25	0.24	0.25	0.24	0.21	0.26	0.19	0.21	0.18	0.18	0.16	0.22
Methamphetamine ³	0.18	0.16	0.16	0.15	0.17	0.12	0.16	0.14	0.11	0.13	0.10	0.08	0.13
Sedatives	0.14	0.12	0.12	0.12	0.11	0.12	0.11	0.13	0.11	0.09	0.10	0.07	0.15
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.49	0.54	0.47	0.49	0.46	0.48	0.50	0.49	0.51	0.47	0.44	0.43	0.48

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.17P Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0001	0.0000	0.0351	0.0008	0.0125	0.0390	0.1561	0.7735	0.1172	
Marijuana and Hashish	0.0000	0.0001	0.0061	0.5867	0.3943	0.1029	0.6585	0.2406	0.6568	0.8696	0.9141	0.4300	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0069	0.0220	0.0881	0.4424	0.3129	
Crack	0.0000	0.0000	0.0003	0.0003	0.0000	0.0001	0.0107	0.1392	0.3093	0.2140	0.8817	0.1571	
Heroin	0.0019	0.0060	0.0907	0.2485	0.1136	0.3256	0.2046	0.0995	0.4314	0.0707	0.1563	0.6844	
Hallucinogens	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0000	0.0000	0.0000	0.0001	0.0001	0.6255	
LSD	0.0000	0.0204	0.7634	0.3566	0.0977	0.0687	0.7300	0.3750	0.5001	0.0860	0.7050	0.0103	
PCP	0.0000	0.0002	0.0139	0.0048	0.0200	0.3370	0.6091	0.2357	0.3931	0.8668	0.3327	0.1838	
Ecstasy	0.0000	0.0000	0.0000	0.0152	0.0000	0.0050	0.0000	0.0000	0.0000	0.0000	0.0000	0.4007	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0066	0.2061	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0000	0.0004	0.0045	0.2523	0.5092	0.0316	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0046	0.1354	0.4613	
OxyContin [®]	0.8250	0.6177	0.3129	0.3701	0.0097	0.0050	0.0048	0.0017	0.0056	0.0009	0.0534	0.2696	
Tranquilizers	0.0640	0.0189	0.0511	0.3248	0.3139	0.2964	0.5058	0.6024	0.4295	0.8538	0.6804	0.0330	
Stimulants ³	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.0011	0.0776	0.6267	0.6759	0.9624	0.0130	
Methamphetamine ³	0.0000	0.0000	0.0000	0.0003	0.0000	0.0452	0.1745	0.1787	0.5684	0.7274	0.6375	0.0643	
Sedatives	0.1660	0.8987	0.9462	0.9927	0.8618	0.9237	0.8240	0.9224	0.4889	0.0463	0.4160	0.0015	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0332	0.0750	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.18C Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	65	64	62	66	66	67	58	65	66	67	60	65	66
Marijuana and Hashish	55	53	53	56	54	57	52	59	60	59	57	57	59
Cocaine	23	19	20	19	18	18	17	17	16	14	14	11	16
Crack	8	9	7	7	8	7	6	5	3	5	5	3	5
Heroin	5	6	8	6	6	5	9	6	6	8	7	6	6
Hallucinogens	29	25	26	25	22	26	26	28	28	25	23	24	21
LSD	18	11	13	14	10	11	14	12	12	14	12	13	16
PCP	8	9	9	11	8	7	8	7	4	5	7	5	4
Ecstasy	22	15	16	14	14	19	18	21	22	20	15	15	15
Inhalants	31	33	34	33	32	31	31	29	27	29	21	21	23
Nonmedical Use of Psychotherapeutics ^{2,3}	42	43	39	41	43	45	38	41	41	37	38	39	40
Pain Relievers	38	40	37	38	40	40	35	38	40	34	35	34	34
OxyContin [®]	--	--	13	13	12	13	14	15	16	12	12	11	13
Tranquilizers	22	22	19	21	18	19	20	18	21	21	20	18	23
Stimulants ³	24	20	18	19	21	18	15	16	16	16	16	16	21
Methamphetamine ³	16	12	12	13	14	8	6	7	10	10	9	7	13
Sedatives	10	13	10	11	11	9	9	12	8	7	7	8	10
ILLCIT DRUGS OTHER THAN MARIJUANA¹	52	53	49	52	51	55	46	48	50	47	44	45	47

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.18D Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.52	0.50	0.48	0.51	0.51	0.52	0.46	0.51	0.53	0.53	0.47	0.51	0.52
Marijuana and Hashish	0.43	0.42	0.41	0.43	0.42	0.44	0.41	0.47	0.49	0.46	0.45	0.45	0.46
Cocaine	0.18	0.15	0.15	0.15	0.14	0.14	0.13	0.14	0.13	0.11	0.11	0.09	0.13
Crack	0.06	0.07	0.05	0.05	0.06	0.05	0.05	0.04	0.02	0.04	0.04	0.02	0.04
Heroin	0.04	0.05	0.06	0.04	0.04	0.04	0.07	0.04	0.05	0.06	0.05	0.04	0.04
Hallucinogens	0.23	0.20	0.20	0.19	0.17	0.20	0.20	0.22	0.22	0.20	0.18	0.19	0.17
LSD	0.15	0.09	0.10	0.10	0.07	0.09	0.11	0.09	0.10	0.11	0.09	0.10	0.13
PCP	0.06	0.07	0.07	0.09	0.06	0.06	0.06	0.06	0.03	0.04	0.06	0.04	0.03
Ecstasy	0.18	0.12	0.12	0.11	0.11	0.14	0.14	0.17	0.18	0.16	0.12	0.12	0.12
Inhalants	0.25	0.26	0.26	0.26	0.25	0.24	0.24	0.23	0.22	0.23	0.17	0.17	0.18
Nonmedical Use of Psychotherapeutics ^{2,3}	0.33	0.33	0.30	0.32	0.33	0.35	0.30	0.33	0.33	0.29	0.29	0.30	0.32
Pain Relievers	0.30	0.31	0.29	0.30	0.31	0.31	0.27	0.30	0.32	0.26	0.27	0.27	0.27
OxyContin [®]	--	--	0.10	0.10	0.09	0.10	0.11	0.12	0.13	0.09	0.09	0.09	0.11
Tranquilizers	0.17	0.17	0.14	0.16	0.14	0.15	0.16	0.14	0.17	0.17	0.16	0.14	0.18
Stimulants ³	0.19	0.15	0.14	0.14	0.16	0.14	0.12	0.13	0.13	0.12	0.13	0.13	0.16
Methamphetamine ³	0.12	0.09	0.09	0.10	0.11	0.06	0.05	0.06	0.08	0.08	0.07	0.05	0.10
Sedatives	0.08	0.10	0.08	0.08	0.08	0.07	0.07	0.10	0.06	0.06	0.05	0.07	0.08
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.41	0.41	0.38	0.40	0.39	0.42	0.36	0.38	0.40	0.37	0.35	0.36	0.37

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.18P Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0001	0.0002	0.0003	0.0000	0.0002	0.0010	0.1036	0.1124	
Marijuana and Hashish	0.0000	0.0008	0.0068	0.2979	0.6300	0.5376	0.4501	0.0043	0.0105	0.0032	0.1576	0.1657	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0002	0.0005	0.0017	0.0121	0.1768	0.1920	0.9765	0.4983	
Crack	0.0087	0.0008	0.0240	0.2119	0.0396	0.1223	0.4040	0.5520	0.5436	0.5503	0.6160	0.3857	
Heroin	0.4450	0.6947	0.0755	0.7712	1.0000	0.8529	0.2390	0.6491	0.9240	0.4036	0.9093	0.7773	
Hallucinogens	0.0000	0.0000	0.0000	0.0003	0.0042	0.0009	0.0000	0.0000	0.0000	0.0000	0.0654	0.0328	
LSD	0.0034	0.0483	0.3754	0.3102	0.0013	0.0698	0.7404	0.3060	0.0751	0.5340	0.1526	0.5121	
PCP	0.0001	0.0004	0.0043	0.0054	0.0868	0.0100	0.0034	0.0685	0.7224	0.1967	0.2374	0.3966	
Ecstasy	0.0000	0.0364	0.0505	0.2083	0.1335	0.0052	0.0061	0.0000	0.0000	0.0000	0.2165	0.0411	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0892	0.5242	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0013	0.0006	0.0213	0.0120	0.1120	0.4700	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0002	0.0036	0.1775	
OxyContin [®]	--	--	0.3539	0.3260	0.4622	0.1723	0.0293	0.1279	0.1008	0.2276	0.3214	0.6424	
Tranquilizers	0.0733	0.0437	0.3159	0.3110	0.9640	0.5437	0.2868	0.5057	0.7249	0.4814	0.7531	0.4047	
Stimulants ³	0.0000	0.0012	0.0761	0.2476	0.0388	0.6340	0.4711	0.4722	0.3928	0.3309	0.6558	0.4992	
Methamphetamine ³	0.0366	0.1256	0.4505	0.2420	0.3452	0.3009	0.1357	0.0968	0.5470	0.5045	0.2963	0.1282	
Sedatives	0.9224	0.1673	0.3205	0.6509	0.4470	0.8693	0.8320	0.0896	0.8938	0.3159	0.1996	0.3815	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0400	0.2691	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.19C Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	63	64	63	60	61	58	63	61	64	61	60	60	65
Marijuana and Hashish	53	55	54	50	51	48	52	48	53	55	53	54	56
Cocaine	23	19	19	19	20	20	17	14	16	15	14	10	15
Crack	11	9	8	7	8	9	5	4	4	4	6	3	4
Heroin	10	5	6	5	6	4	6	5	5	9	6	5	6
Hallucinogens	27	26	26	24	25	23	25	26	29	23	22	18	22
LSD	16	11	10	10	10	9	11	8	12	11	11	9	17
PCP	10	8	7	9	8	6	5	7	7	5	8	4	8
Ecstasy	22	17	17	15	18	18	22	20	26	21	18	11	16
Inhalants	30	31	34	33	32	30	31	30	32	29	27	20	27
Nonmedical Use of Psychotherapeutics ^{2,3}	43	46	45	44	43	42	42	42	42	38	38	32	45
Pain Relievers	40	42	43	39	39	40	39	40	40	36	33	30	38
OxyContin [®]	--	--	13	12	13	14	12	13	17	14	13	11	13
Tranquilizers	24	24	24	21	24	24	21	20	21	23	24	16	25
Stimulants ³	25	23	21	24	25	20	23	18	20	17	16	14	24
Methamphetamine ³	17	13	14	14	15	12	13	11	11	12	8	8	12
Sedatives	13	11	11	12	8	12	11	10	12	9	9	6	15
ILLCIT DRUGS OTHER THAN MARIJUANA¹	51	56	52	50	49	48	52	51	55	45	45	39	50

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.19D Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.52	0.52	0.51	0.48	0.49	0.47	0.52	0.51	0.54	0.50	0.50	0.50	0.53
Marijuana and Hashish	0.44	0.45	0.44	0.40	0.41	0.39	0.43	0.40	0.45	0.45	0.44	0.44	0.46
Cocaine	0.19	0.16	0.15	0.15	0.16	0.16	0.14	0.11	0.14	0.12	0.11	0.09	0.13
Crack	0.09	0.07	0.06	0.06	0.07	0.07	0.04	0.03	0.03	0.04	0.05	0.02	0.03
Heroin	0.08	0.04	0.05	0.04	0.05	0.03	0.05	0.04	0.04	0.07	0.05	0.04	0.05
Hallucinogens	0.23	0.22	0.21	0.19	0.20	0.18	0.21	0.21	0.24	0.19	0.18	0.15	0.18
LSD	0.13	0.09	0.08	0.08	0.08	0.07	0.09	0.07	0.10	0.09	0.09	0.08	0.14
PCP	0.08	0.06	0.06	0.07	0.06	0.05	0.04	0.06	0.06	0.04	0.06	0.04	0.06
Ecstasy	0.18	0.14	0.14	0.12	0.14	0.14	0.18	0.16	0.22	0.17	0.15	0.09	0.13
Inhalants	0.25	0.25	0.27	0.27	0.26	0.24	0.26	0.25	0.27	0.23	0.22	0.17	0.22
Nonmedical Use of Psychotherapeutics ^{2,3}	0.36	0.38	0.36	0.35	0.35	0.34	0.34	0.35	0.35	0.31	0.31	0.27	0.37
Pain Relievers	0.33	0.34	0.35	0.32	0.31	0.33	0.32	0.33	0.33	0.29	0.27	0.25	0.31
OxyContin [®]	--	--	0.10	0.09	0.11	0.12	0.10	0.11	0.14	0.11	0.10	0.09	0.11
Tranquilizers	0.20	0.19	0.19	0.17	0.19	0.19	0.17	0.16	0.18	0.18	0.20	0.13	0.20
Stimulants ³	0.21	0.19	0.17	0.20	0.20	0.16	0.19	0.15	0.17	0.14	0.13	0.11	0.20
Methamphetamine ³	0.14	0.10	0.11	0.11	0.12	0.09	0.10	0.09	0.09	0.10	0.07	0.06	0.10
Sedatives	0.10	0.09	0.09	0.10	0.06	0.10	0.09	0.08	0.10	0.08	0.08	0.05	0.13
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.42	0.46	0.43	0.40	0.39	0.39	0.42	0.42	0.46	0.37	0.37	0.32	0.41

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.19P Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0130	0.0451	0.7986	0.2945	0.3383	0.0883	0.4371	0.7859	0.0249	
Marijuana and Hashish	0.0006	0.0120	0.0914	0.7210	0.8591	0.0067	0.3897	0.1623	0.9319	0.6913	0.7721	0.6811	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0055	0.4241	0.0411	0.1266	0.5780	0.2206	
Crack	0.0000	0.0000	0.0072	0.0046	0.0003	0.0012	0.1935	0.3804	0.6070	0.4548	0.6898	0.4527	
Heroin	0.1407	0.5387	0.8518	0.5007	0.3503	0.3319	0.9465	0.7089	0.8884	0.0980	0.8378	0.9228	
Hallucinogens	0.0000	0.0000	0.0000	0.0005	0.0001	0.0052	0.0005	0.0001	0.0000	0.0085	0.0188	0.7579	
LSD	0.2269	0.1922	0.0439	0.0238	0.0134	0.0061	0.0707	0.0100	0.0668	0.0390	0.1059	0.0097	
PCP	0.0009	0.0509	0.3623	0.1436	0.3784	0.9495	0.8417	0.9482	0.9486	0.7789	0.6455	0.2704	
Ecstasy	0.0000	0.0000	0.0004	0.0422	0.0002	0.0013	0.0002	0.0000	0.0000	0.0000	0.0010	0.7397	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0013	0.0371	0.1202	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0000	0.0000	0.0000	0.0001	0.0001	0.0030	0.0027	0.0086	0.0084	0.3783	0.9607	0.0058	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.0000	0.0000	0.0004	0.0110	0.6518	0.1270	
OxyContin [®]	--	--	0.4838	0.8289	0.2505	0.0303	0.2194	0.0620	0.0458	0.0658	0.7181	0.8210	
Tranquilizers	0.0100	0.0055	0.0349	0.3213	0.1275	0.1046	0.4206	0.4647	0.2092	0.4244	0.2860	0.0663	
Stimulants ³	0.0000	0.0003	0.0046	0.0023	0.0009	0.2714	0.3583	0.8729	0.8344	0.1915	0.3758	0.0113	
Methamphetamine ³	0.0000	0.0028	0.0027	0.0084	0.0021	0.1225	0.5302	0.6337	0.7818	0.5125	0.6729	0.5023	
Sedatives	0.2837	0.6593	0.9359	0.7481	0.2559	0.8050	0.7695	0.3959	0.9670	0.2706	0.2807	0.0166	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0099	0.2106	0.0076	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.20C Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	52	47	44	49	46	54	43	48	52	50	47	46	52
Marijuana and Hashish	44	43	39	43	39	44	38	45	47	46	41	42	48
Cocaine	12	10	12	13	9	9	10	10	9	5	4	6	6
Crack	4	7	4	6	2	2	2	4	*	2	1	1	4
Heroin	3	5	4	4	2	2	4	4	1	6	*	2	4
Hallucinogens	14	15	15	13	11	13	16	16	16	16	13	14	15
LSD	8	7	7	7	4	7	8	5	8	7	3	8	11
PCP	4	5	4	4	3	5	4	4	2	2	3	1	2
Ecstasy	10	8	10	7	8	8	10	12	10	10	7	7	9
Inhalants	16	16	16	18	18	19	17	16	15	13	13	10	13
Nonmedical Use of Psychotherapeutics ^{2,3}	30	28	26	26	27	29	22	24	24	24	26	23	26
Pain Relievers	26	26	24	24	25	26	20	22	23	21	23	20	22
OxyContin [®]	--	--	7	6	5	6	7	9	8	8	8	4	6
Tranquilizers	11	14	12	13	9	12	10	9	11	12	12	9	11
Stimulants ³	17	13	13	14	12	9	9	10	9	8	10	8	15
Methamphetamine ³	13	8	8	10	7	4	5	5	4	4	3	5	8
Sedatives	3	8	2	4	7	6	3	8	5	4	3	6	6
ILLCIT DRUGS OTHER THAN MARIJUANA¹	36	32	32	33	33	37	30	30	30	30	31	27	31

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.20D Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.41	0.37	0.34	0.38	0.36	0.42	0.34	0.38	0.42	0.39	0.37	0.36	0.41
Marijuana and Hashish	0.35	0.34	0.30	0.33	0.30	0.34	0.30	0.36	0.37	0.36	0.32	0.33	0.38
Cocaine	0.10	0.08	0.09	0.10	0.07	0.07	0.08	0.08	0.07	0.04	0.03	0.05	0.05
Crack	0.03	0.05	0.03	0.04	0.02	0.02	0.02	0.03	*	0.01	0.01	0.01	0.03
Heroin	0.02	0.04	0.03	0.03	0.01	0.01	0.03	0.03	0.01	0.05	*	0.02	0.03
Hallucinogens	0.11	0.11	0.11	0.10	0.09	0.10	0.13	0.12	0.12	0.13	0.10	0.11	0.12
LSD	0.07	0.06	0.05	0.06	0.03	0.06	0.06	0.04	0.07	0.05	0.03	0.06	0.08
PCP	0.03	0.04	0.03	0.03	0.02	0.04	0.03	0.03	0.01	0.01	0.02	0.01	0.02
Ecstasy	0.08	0.06	0.08	0.05	0.06	0.06	0.08	0.09	0.08	0.08	0.05	0.06	0.07
Inhalants	0.13	0.13	0.12	0.14	0.14	0.15	0.14	0.13	0.12	0.10	0.10	0.08	0.10
Nonmedical Use of Psychotherapeutics ^{2,3}	0.23	0.22	0.20	0.20	0.21	0.22	0.17	0.19	0.19	0.18	0.20	0.18	0.21
Pain Relievers	0.21	0.20	0.18	0.18	0.20	0.20	0.15	0.18	0.19	0.17	0.18	0.16	0.17
OxyContin [®]	--	--	0.05	0.04	0.04	0.05	0.05	0.07	0.06	0.06	0.06	0.03	0.04
Tranquilizers	0.08	0.11	0.09	0.10	0.07	0.09	0.08	0.07	0.09	0.09	0.09	0.07	0.09
Stimulants ³	0.13	0.11	0.10	0.10	0.09	0.07	0.07	0.08	0.08	0.06	0.08	0.07	0.12
Methamphetamine ³	0.10	0.06	0.06	0.08	0.06	0.03	0.04	0.04	0.03	0.03	0.02	0.04	0.06
Sedatives	0.02	0.06	0.01	0.03	0.05	0.04	0.02	0.06	0.04	0.03	0.02	0.05	0.05
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.28	0.25	0.25	0.26	0.25	0.28	0.23	0.24	0.24	0.23	0.24	0.21	0.25

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.20P Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0015	0.0826	0.3659	0.7025	0.4770	0.7793	0.0615	0.1791	0.0368	0.8834	0.9520	
Marijuana and Hashish	0.0175	0.1839	0.7277	0.4068	0.0392	0.5075	0.2333	0.3179	0.3693	0.0294	0.4003	0.9710	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0034	0.0034	0.0002	0.0228	0.0566	0.3984	0.6560	0.6893	
Crack	0.5561	0.0626	0.6749	0.3603	0.6674	0.5867	0.8212	0.7772	*	0.2712	0.1496	0.2950	
Heroin	0.6874	0.9312	0.5877	0.6746	0.1558	0.2275	0.9631	0.9963	0.1133	0.9696	*	0.3684	
Hallucinogens	0.0275	0.0122	0.1097	0.5712	0.9581	0.4183	0.0046	0.0409	0.1298	0.0278	0.7385	0.5388	
LSD	0.9580	0.3919	0.2223	0.2533	0.0115	0.2442	0.8396	0.1476	0.5469	0.3244	0.0120	0.8540	
PCP	0.0848	0.0575	0.3558	0.3284	0.4477	0.2428	0.1402	0.2855	0.8845	0.8793	0.8248	0.6070	
Ecstasy	0.0355	0.5484	0.2902	0.8728	0.5483	0.7830	0.0427	0.0179	0.1293	0.0463	0.9670	0.9460	
Inhalants	0.0000	0.0011	0.0008	0.0001	0.0002	0.0002	0.0029	0.0154	0.0234	0.1413	0.4095	0.4913	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0000	0.0000	0.0025	0.0086	0.0082	0.0184	0.6657	0.0959	0.8226	0.8741	0.7475	0.2751	
Pain Relievers	0.0000	0.0000	0.0001	0.0004	0.0004	0.0006	0.1519	0.0016	0.1634	0.2192	0.1669	0.5876	
OxyContin [®]	--	--	0.1835	0.9827	0.8646	0.3578	0.0727	0.0239	0.2715	0.2008	0.2282	0.3671	
Tranquilizers	0.2385	0.0014	0.1627	0.0773	0.6385	0.0599	0.4861	0.6563	0.5263	0.1924	0.9897	0.4740	
Stimulants ³	0.2977	0.2459	0.8096	0.7810	0.8255	0.0424	0.0985	0.1031	0.0648	0.0163	0.0702	0.0236	
Methamphetamine ³	0.2375	0.3882	0.8985	0.2037	0.9177	0.1585	0.4802	0.6322	0.1191	0.3035	0.1038	0.3360	
Sedatives	0.0752	0.4150	0.0366	0.2274	0.8201	0.6623	0.1351	0.5333	0.6416	0.2737	0.0966	0.7476	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0000	0.0001	0.0010	0.0068	0.0039	0.1907	0.1370	0.7035	0.2756	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.21C Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	44	45	50	43	45	44	44	45	49	44	46	41	50
Marijuana and Hashish	35	39	38	35	37	35	39	36	40	38	39	37	43
Cocaine	13	12	9	10	10	10	7	7	7	10	6	7	8
Crack	7	3	3	5	3	4	1	2	2	1	*	*	1
Heroin	5	4	2	2	5	*	5	2	3	4	*	4	5
Hallucinogens	15	13	12	14	14	11	14	12	14	14	12	10	12
LSD	9	7	4	3	6	3	7	5	5	4	5	6	10
PCP	4	3	3	4	3	*	2	3	2	1	*	3	*
Ecstasy	12	10	7	9	9	8	11	10	12	12	9	5	5
Inhalants	16	19	20	16	18	17	16	15	17	15	14	11	13
Nonmedical Use of Psychotherapeutics ^{2,3}	30	31	31	26	28	28	26	28	30	24	27	21	31
Pain Relievers	28	28	28	24	25	25	22	26	26	22	23	19	25
OxyContin [®]	--	--	8	5	3	5	5	6	6	6	6	5	7
Tranquilizers	14	16	12	10	11	15	13	12	11	13	16	8	11
Stimulants ³	13	14	14	12	12	11	9	11	12	10	10	8	16
Methamphetamine ³	7	8	9	8	8	7	4	6	7	7	5	3	9
Sedatives	8	6	6	5	5	7	8	5	6	4	6	3	8
ILLCIT DRUGS OTHER THAN MARIJUANA¹	35	37	38	31	33	33	31	34	37	29	31	25	35

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.21D Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.37	0.37	0.40	0.35	0.36	0.35	0.36	0.38	0.41	0.36	0.38	0.34	0.41
Marijuana and Hashish	0.29	0.32	0.31	0.29	0.29	0.29	0.32	0.30	0.34	0.31	0.32	0.30	0.35
Cocaine	0.11	0.10	0.07	0.08	0.08	0.08	0.06	0.06	0.05	0.08	0.05	0.06	0.06
Crack	0.06	0.03	0.03	0.04	0.03	0.03	0.01	0.01	0.02	0.01	*	*	0.01
Heroin	0.04	0.03	0.02	0.01	0.04	*	0.04	0.02	0.02	0.03	*	0.03	0.04
Hallucinogens	0.12	0.11	0.10	0.11	0.11	0.09	0.11	0.10	0.12	0.11	0.09	0.08	0.10
LSD	0.07	0.06	0.04	0.03	0.05	0.02	0.05	0.04	0.04	0.03	0.04	0.05	0.08
PCP	0.03	0.02	0.02	0.03	0.03	*	0.02	0.02	0.02	0.01	*	0.03	*
Ecstasy	0.10	0.08	0.06	0.07	0.07	0.06	0.09	0.08	0.10	0.10	0.08	0.04	0.04
Inhalants	0.13	0.15	0.16	0.13	0.15	0.14	0.13	0.13	0.15	0.12	0.11	0.09	0.11
Nonmedical Use of Psychotherapeutics ^{2,3}	0.25	0.25	0.25	0.21	0.22	0.23	0.21	0.23	0.25	0.20	0.22	0.17	0.25
Pain Relievers	0.23	0.23	0.23	0.19	0.20	0.20	0.18	0.21	0.22	0.18	0.19	0.15	0.20
OxyContin [®]	--	--	0.07	0.04	0.03	0.04	0.04	0.05	0.05	0.05	0.05	0.04	0.06
Tranquilizers	0.11	0.13	0.09	0.08	0.09	0.12	0.10	0.10	0.10	0.10	0.13	0.07	0.09
Stimulants ³	0.10	0.11	0.11	0.10	0.10	0.09	0.07	0.09	0.10	0.08	0.08	0.06	0.13
Methamphetamine ³	0.05	0.06	0.07	0.06	0.06	0.05	0.03	0.05	0.06	0.06	0.05	0.03	0.07
Sedatives	0.07	0.05	0.04	0.04	0.04	0.06	0.06	0.04	0.05	0.04	0.05	0.02	0.06
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.29	0.30	0.31	0.25	0.27	0.26	0.25	0.28	0.31	0.24	0.25	0.20	0.28

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.21P Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0016	0.0005	0.0092	0.3483	0.2699	0.8711	0.9015	0.5626	0.2379	0.7240	0.5207	0.0328	
Marijuana and Hashish	0.4014	0.4334	0.5848	0.1452	0.4350	0.0311	0.1145	0.2582	0.3914	0.7830	0.7595	0.2033	
Cocaine	0.0038	0.0000	0.0256	0.0022	0.0035	0.0094	0.2433	0.7190	0.9488	0.1111	0.5348	0.9565	
Crack	0.0491	0.0054	0.1125	0.0230	0.0593	0.0135	0.7161	0.2713	0.2243	0.8004	*	*	
Heroin	0.6709	0.7084	0.4750	0.4925	0.4412	*	0.7090	0.6262	0.8007	0.7953	*	0.9369	
Hallucinogens	0.0001	0.0001	0.0267	0.0031	0.0193	0.2546	0.0138	0.0302	0.0072	0.1016	0.3242	0.7446	
LSD	0.7864	0.7639	0.2459	0.0645	0.6049	0.0379	0.3611	0.2265	0.3600	0.0677	0.3161	0.3138	
PCP	*	*	*	*	*	*	*	*	*	*	*	*	
Ecstasy	0.0000	0.0000	0.0086	0.0016	0.0004	0.0162	0.0026	0.0002	0.0000	0.0008	0.0013	0.1704	
Inhalants	0.0017	0.0000	0.0002	0.0004	0.0000	0.0105	0.0077	0.0209	0.0011	0.0218	0.2175	0.3155	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0001	0.0003	0.0021	0.0795	0.0723	0.0857	0.3113	0.1012	0.0376	0.5216	0.5507	0.0681	
Pain Relievers	0.0000	0.0000	0.0001	0.0073	0.0087	0.0229	0.0916	0.0066	0.0033	0.0932	0.1805	0.4038	
OxyContin [®]	--	--	0.0878	0.8906	0.2654	0.6698	0.8520	0.2364	0.8332	0.7157	0.8641	0.9380	
Tranquilizers	0.0002	0.0014	0.0382	0.1178	0.0991	0.0151	0.0315	0.0295	0.1553	0.0306	0.0641	0.8305	
Stimulants ³	0.4084	0.1765	0.8446	0.7674	0.8806	0.6664	0.1459	0.5729	0.2024	0.1117	0.3319	0.0130	
Methamphetamine ³	0.4469	0.1871	0.3726	0.4367	0.6194	0.9606	0.3047	0.6894	0.5953	0.9593	0.9161	0.1425	
Sedatives	0.0863	0.8465	0.9553	0.5886	0.8729	0.7757	0.6520	0.4752	0.9248	0.3755	0.7925	0.1260	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0000	0.0000	0.0000	0.0006	0.0004	0.0095	0.0523	0.0118	0.0006	0.1031	0.7431	0.0507	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.22C GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	118	106	106	119	155	121	121	122	93
AGE GROUP									
12 to 17	8	7	5	8	6	5	7	3	5
18 to 25	37	30	22	25	27	20	22	22	22
26 or Older	111	100	100	117	150	120	117	120	89
GENDER									
Male	88	87	87	98	130	101	106	95	75
Female	70	59	55	63	68	63	66	77	51
PAST YEAR USE									
TOTAL	31	13	31	26	47	37	49	39	32
AGE GROUP									
12 to 17	5	3	3	5	2	4	7	2	4
18 to 25	10	10	8	10	9	8	10	7	11
26 or Older	28	11	30	24	46	35	45	38	28
GENDER									
Male	29	12	30	24	43	36	41	29	29
Female	7	4	9	9	19	6	28	26	14
PAST MONTH USE									
TOTAL	18	9	18	9	29	28	34	20	21
AGE GROUP									
12 to 17	3	3	1	3	*	1	6	*	3
18 to 25	4	8	3	3	3	3	7	3	*
26 or Older	17	*	17	*	29	28	29	20	20
GENDER									
Male	18	9	17	4	23	28	20	20	18
Female	*	*	*	8	18	*	28	*	9

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.22D GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.05	0.04	0.04	0.05	0.06	0.05	0.05	0.05	0.04
AGE GROUP									
12 to 17	0.03	0.03	0.02	0.03	0.02	0.02	0.03	0.01	0.02
18 to 25	0.11	0.09	0.07	0.08	0.08	0.06	0.06	0.06	0.06
26 or Older	0.06	0.05	0.05	0.06	0.08	0.06	0.06	0.06	0.04
GENDER									
Male	0.07	0.07	0.07	0.08	0.11	0.08	0.08	0.08	0.06
Female	0.06	0.05	0.04	0.05	0.05	0.05	0.05	0.06	0.04
PAST YEAR USE									
TOTAL	0.01	0.01	0.01	0.01	0.02	0.01	0.02	0.01	0.01
AGE GROUP									
12 to 17	0.02	0.01	0.01	0.02	0.01	0.02	0.03	0.01	0.02
18 to 25	0.03	0.03	0.02	0.03	0.03	0.02	0.03	0.02	0.03
26 or Older	0.01	0.01	0.02	0.01	0.02	0.02	0.02	0.02	0.01
GENDER									
Male	0.02	0.01	0.02	0.02	0.03	0.03	0.03	0.02	0.02
Female	0.01	0.00	0.01	0.01	0.01	0.00	0.02	0.02	0.01
PAST MONTH USE									
TOTAL	0.01	0.00	0.01	0.00	0.01	0.01	0.01	0.01	0.01
AGE GROUP									
12 to 17	0.01	0.01	0.00	0.01	*	0.00	0.02	*	0.01
18 to 25	0.01	0.03	0.01	0.01	0.01	0.01	0.02	0.01	*
26 or Older	0.01	*	0.01	*	0.01	0.01	0.01	0.01	0.01
GENDER									
Male	0.01	0.01	0.01	0.00	0.02	0.02	0.02	0.02	0.01
Female	*	*	*	0.01	0.01	*	0.02	*	0.01

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.22P GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2006-2013

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.0114	0.7021	0.5315	0.3145	0.0273	0.6672	0.4710	0.4913	
AGE GROUP									
12 to 17	0.0098	0.4166	0.3949	0.0801	0.3266	0.6560	0.5159	0.1721	
18 to 25	0.0000	0.0000	0.0302	0.0005	0.0245	0.9640	0.8009	0.8816	
26 or Older	0.9034	0.3666	0.2452	0.8831	0.0695	0.6580	0.4981	0.4197	
GENDER									
Male	0.1226	0.6790	0.5950	0.2735	0.0126	0.3019	0.4026	0.8180	
Female	0.0205	0.9443	0.7114	0.8521	0.8089	0.4276	0.9898	0.4005	
PAST YEAR USE									
TOTAL	0.9745	0.0613	0.4735	0.4180	0.5097	0.5774	0.9060	0.7234	
AGE GROUP									
12 to 17	0.1558	0.9520	0.8032	0.2259	0.4186	0.4534	0.3156	0.4410	
18 to 25	0.1077	0.1624	0.9772	0.5870	0.6196	0.9661	0.5305	0.6729	
26 or Older	0.4187	0.0044	0.4259	0.1803	0.5253	0.4949	0.8483	0.8484	
GENDER									
Male	0.7208	0.2160	0.7844	0.6932	0.5605	0.9790	0.9136	0.3957	
Female	0.3952	0.0775	0.2107	0.2628	0.7612	0.0665	0.9603	0.5678	
PAST MONTH USE									
TOTAL	0.2891	0.1266	0.2741	0.0637	0.7297	0.6533	0.8656	0.3006	
AGE GROUP									
12 to 17	0.7578	0.9390	0.4012	0.6950	*	0.4356	0.2846	*	
18 to 25	*	*	*	*	*	*	*	*	
26 or Older	0.1870	*	0.2512	*	0.7666	0.6651	0.7429	0.3116	
GENDER									
Male	0.5283	0.3471	0.4372	0.0698	0.9610	0.9732	0.7082	0.5873	
Female	*	*	*	0.5375	0.4981	*	0.5598	*	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.23C Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	165	170	180	220	257	232	286	321	284
AGE GROUP									
12 to 17	34	37	31	32	37	34	37	36	40
18 to 25	98	97	95	112	117	113	122	126	142
26 or Older	112	116	139	164	204	194	233	285	216
GENDER									
Male	123	125	127	174	197	161	214	213	202
Female	96	102	118	113	139	158	172	209	173
PAST YEAR USE									
TOTAL	92	99	102	135	128	134	152	172	146
AGE GROUP									
12 to 17	27	28	25	25	29	27	26	29	33
18 to 25	66	69	65	82	84	86	88	88	93
26 or Older	53	58	71	90	76	91	112	142	96
GENDER									
Male	62	72	74	107	97	93	106	112	112
Female	60	61	65	70	67	92	96	119	89
PAST MONTH USE									
TOTAL	43	45	55	65	73	65	70	93	88
AGE GROUP									
12 to 17	14	17	13	13	16	12	14	16	19
18 to 25	32	37	31	36	46	41	43	45	49
26 or Older	24	15	42	53	49	47	55	76	66
GENDER									
Male	33	32	41	57	57	45	53	47	65
Female	27	30	33	35	37	42	45	75	57

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.23D Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.07	0.07	0.07	0.09	0.10	0.09	0.11	0.12	0.11
AGE GROUP									
12 to 17	0.14	0.15	0.12	0.13	0.15	0.13	0.15	0.14	0.16
18 to 25	0.30	0.30	0.29	0.33	0.34	0.33	0.35	0.36	0.41
26 or Older	0.06	0.06	0.07	0.08	0.10	0.10	0.12	0.14	0.11
GENDER									
Male	0.10	0.10	0.10	0.14	0.16	0.13	0.17	0.17	0.16
Female	0.08	0.08	0.09	0.09	0.11	0.12	0.13	0.15	0.13
PAST YEAR USE									
TOTAL	0.04	0.04	0.04	0.05	0.05	0.05	0.06	0.07	0.06
AGE GROUP									
12 to 17	0.11	0.11	0.10	0.10	0.12	0.11	0.10	0.12	0.13
18 to 25	0.20	0.21	0.20	0.24	0.25	0.25	0.25	0.25	0.27
26 or Older	0.03	0.03	0.04	0.05	0.04	0.05	0.06	0.07	0.05
GENDER									
Male	0.05	0.06	0.06	0.09	0.08	0.07	0.08	0.09	0.09
Female	0.05	0.05	0.05	0.05	0.05	0.07	0.07	0.09	0.06
PAST MONTH USE									
TOTAL	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.04	0.03
AGE GROUP									
12 to 17	0.05	0.07	0.05	0.05	0.06	0.05	0.06	0.07	0.08
18 to 25	0.10	0.11	0.09	0.11	0.13	0.12	0.12	0.13	0.14
26 or Older	0.01	0.01	0.02	0.03	0.03	0.02	0.03	0.04	0.03
GENDER									
Male	0.03	0.03	0.03	0.05	0.05	0.04	0.04	0.04	0.05
Female	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.06	0.04

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.23P Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2006-2013

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0406	
AGE GROUP									
12 to 17	0.6365	0.9831	0.3239	0.6294	0.7712	0.4006	0.9852	0.3625	
18 to 25	0.0000	0.0000	0.0000	0.0000	0.0003	0.0003	0.0391	0.3073	
26 or Older	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0661	
GENDER									
Male	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0357	
Female	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.3335	
PAST YEAR USE									
TOTAL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.1406	0.7223	
AGE GROUP									
12 to 17	0.9494	0.3802	0.2660	0.5369	0.9133	0.6286	0.3750	0.3933	
18 to 25	0.0000	0.0000	0.0000	0.0011	0.0341	0.0237	0.3993	0.3974	
26 or Older	0.0000	0.0000	0.0000	0.0000	0.0000	0.0011	0.2019	0.7612	
GENDER									
Male	0.0000	0.0000	0.0000	0.0011	0.0006	0.0005	0.0728	0.2799	
Female	0.0000	0.0000	0.0000	0.0001	0.0032	0.1089	0.7891	0.5410	
PAST MONTH USE									
TOTAL	0.0000	0.0000	0.0000	0.0008	0.0071	0.0002	0.0061	0.1950	
AGE GROUP									
12 to 17	0.4684	0.9536	0.1749	0.5625	0.8685	0.6142	0.1935	0.3977	
18 to 25	0.0000	0.0000	0.0000	0.0131	0.0663	0.0299	0.1965	0.1764	
26 or Older	0.0000	0.0000	0.0000	0.0127	0.0177	0.0008	0.0191	0.5586	
GENDER									
Male	0.0000	0.0000	0.0000	0.0774	0.1586	0.0280	0.0369	0.0274	
Female	0.0000	0.0000	0.0000	0.0014	0.0057	0.0008	0.0659	0.9082	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.24C Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	206	216	233	265	289	264	277	297	235
AGE GROUP									
12 to 17	24	24	21	25	21	22	24	19	22
18 to 25	61	54	54	55	62	54	63	56	62
26 or Older	190	207	221	254	279	258	265	291	227
GENDER									
Male	143	149	149	185	213	166	188	191	159
Female	140	162	181	178	193	204	202	225	163
PAST YEAR USE									
TOTAL	138	140	151	149	184	138	152	175	116
AGE GROUP									
12 to 17	18	16	16	18	15	17	15	12	14
18 to 25	42	38	35	37	36	32	35	28	36
26 or Older	130	133	144	141	177	133	147	171	108
GENDER									
Male	102	98	95	106	142	86	97	99	82
Female	93	98	112	103	114	104	115	130	75
PAST MONTH USE									
TOTAL	85	54	58	83	133	72	88	61	51
AGE GROUP									
12 to 17	10	7	6	7	8	7	7	6	7
18 to 25	17	16	18	18	16	16	18	11	17
26 or Older	81	51	55	79	132	71	88	59	48
GENDER									
Male	70	27	38	49	113	53	48	54	34
Female	41	45	44	66	73	49	74	32	39

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.24D Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.08	0.09	0.09	0.11	0.11	0.10	0.11	0.11	0.09
AGE GROUP									
12 to 17	0.10	0.09	0.08	0.10	0.09	0.09	0.10	0.08	0.09
18 to 25	0.19	0.17	0.16	0.16	0.18	0.16	0.18	0.16	0.18
26 or Older	0.10	0.11	0.12	0.13	0.14	0.13	0.13	0.14	0.11
GENDER									
Male	0.12	0.12	0.12	0.15	0.17	0.13	0.15	0.15	0.12
Female	0.11	0.13	0.14	0.14	0.15	0.15	0.15	0.17	0.12
PAST YEAR USE									
TOTAL	0.06	0.06	0.06	0.06	0.07	0.05	0.06	0.07	0.04
AGE GROUP									
12 to 17	0.07	0.06	0.06	0.07	0.06	0.07	0.06	0.05	0.06
18 to 25	0.13	0.12	0.11	0.11	0.11	0.09	0.10	0.08	0.10
26 or Older	0.07	0.07	0.07	0.07	0.09	0.07	0.07	0.08	0.05
GENDER									
Male	0.09	0.08	0.08	0.09	0.12	0.07	0.08	0.08	0.06
Female	0.07	0.08	0.09	0.08	0.09	0.08	0.09	0.10	0.06
PAST MONTH USE									
TOTAL	0.03	0.02	0.02	0.03	0.05	0.03	0.03	0.02	0.02
AGE GROUP									
12 to 17	0.04	0.03	0.02	0.03	0.03	0.03	0.03	0.02	0.03
18 to 25	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.03	0.05
26 or Older	0.04	0.03	0.03	0.04	0.07	0.04	0.04	0.03	0.02
GENDER									
Male	0.06	0.02	0.03	0.04	0.09	0.04	0.04	0.04	0.03
Female	0.03	0.04	0.03	0.05	0.06	0.04	0.06	0.02	0.03

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.24P Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2006-2013

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.0000	0.0173	0.1585	0.5551	0.3077	0.9049	0.3850	0.5343	
AGE GROUP									
12 to 17	0.0148	0.0480	0.2070	0.0675	0.8098	0.1734	0.1902	0.9014	
18 to 25	0.0395	0.3509	0.0402	0.0026	0.0072	0.5267	0.1270	0.7614	
26 or Older	0.0000	0.0042	0.0391	0.7880	0.6970	0.7087	0.6816	0.4678	
GENDER									
Male	0.0005	0.0137	0.0077	0.9864	0.3164	0.2016	0.9793	0.6282	
Female	0.0005	0.3383	0.6344	0.3771	0.6558	0.3367	0.2349	0.6791	
PAST YEAR USE									
TOTAL	0.0003	0.0003	0.0005	0.0001	0.0001	0.0119	0.0490	0.2050	
AGE GROUP									
12 to 17	0.0004	0.0053	0.0211	0.0199	0.0935	0.0070	0.1446	0.5486	
18 to 25	0.0000	0.0016	0.0008	0.0000	0.0128	0.2595	0.1015	0.4358	
26 or Older	0.0362	0.0101	0.0120	0.0058	0.0011	0.0482	0.1492	0.1521	
GENDER									
Male	0.0097	0.0098	0.0328	0.0022	0.0040	0.4299	0.3843	0.3955	
Female	0.0070	0.0071	0.0032	0.0075	0.0075	0.0051	0.0525	0.2960	
PAST MONTH USE									
TOTAL	0.0578	0.8378	0.3061	0.0110	0.0018	0.2079	0.1450	0.4038	
AGE GROUP									
12 to 17	0.1050	0.6319	0.7243	0.3254	0.3230	0.3437	0.6169	0.7032	
18 to 25	0.3139	0.1386	0.0151	0.0243	0.1761	0.5385	0.5809	0.1941	
26 or Older	0.1231	0.7319	0.8151	0.0486	0.0041	0.3050	0.1955	0.5633	
GENDER									
Male	0.0514	0.2904	0.6219	0.0326	0.0193	0.7583	0.9607	0.8657	
Female	0.5299	0.3076	0.3575	0.1138	0.0387	0.1414	0.0834	0.1257	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.25C Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	283	260	274	258	288	254	286	316	239
AGE GROUP									
12 to 17	40	41	36	36	39	37	38	33	37
18 to 25	77	74	71	83	80	79	86	85	83
26 or Older	264	244	260	235	269	242	271	301	221
GENDER									
Male	229	203	223	193	236	215	244	247	195
Female	151	163	169	163	157	145	168	188	137
PAST YEAR USE									
TOTAL	99	85	98	111	162	111	99	133	92
AGE GROUP									
12 to 17	30	29	26	25	27	25	27	21	25
18 to 25	38	37	31	35	37	40	40	37	35
26 or Older	87	72	91	100	158	100	85	125	79
GENDER									
Male	76	55	81	75	147	86	59	104	73
Female	64	66	56	80	70	71	78	83	55
PAST MONTH USE									
TOTAL	51	46	53	60	67	60	70	77	53
AGE GROUP									
12 to 17	13	15	14	13	12	11	16	12	11
18 to 25	15	19	16	17	21	15	24	19	20
26 or Older	47	39	49	57	63	56	64	74	47
GENDER									
Male	45	34	43	37	52	43	31	68	41
Female	26	31	31	47	43	42	63	38	32

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.25D Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.11	0.10	0.11	0.10	0.11	0.10	0.11	0.12	0.09
AGE GROUP									
12 to 17	0.16	0.16	0.15	0.15	0.16	0.15	0.15	0.13	0.15
18 to 25	0.24	0.23	0.22	0.25	0.23	0.23	0.25	0.24	0.24
26 or Older	0.14	0.13	0.14	0.12	0.14	0.12	0.14	0.15	0.11
GENDER									
Male	0.19	0.17	0.18	0.16	0.19	0.17	0.19	0.19	0.15
Female	0.12	0.13	0.13	0.13	0.12	0.11	0.13	0.14	0.10
PAST YEAR USE									
TOTAL	0.04	0.03	0.04	0.04	0.06	0.04	0.04	0.05	0.03
AGE GROUP									
12 to 17	0.12	0.12	0.10	0.10	0.11	0.10	0.11	0.08	0.10
18 to 25	0.12	0.11	0.10	0.11	0.11	0.12	0.12	0.11	0.10
26 or Older	0.05	0.04	0.05	0.05	0.08	0.05	0.04	0.06	0.04
GENDER									
Male	0.06	0.05	0.07	0.06	0.12	0.07	0.05	0.08	0.06
Female	0.05	0.05	0.04	0.06	0.05	0.05	0.06	0.06	0.04
PAST MONTH USE									
TOTAL	0.02	0.02	0.02	0.02	0.03	0.02	0.03	0.03	0.02
AGE GROUP									
12 to 17	0.05	0.06	0.05	0.05	0.05	0.05	0.06	0.05	0.04
18 to 25	0.05	0.06	0.05	0.05	0.06	0.04	0.07	0.05	0.06
26 or Older	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.04	0.02
GENDER									
Male	0.04	0.03	0.04	0.03	0.04	0.03	0.02	0.05	0.03
Female	0.02	0.02	0.02	0.04	0.03	0.03	0.05	0.03	0.02

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.25P Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2006-2013

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.8047	0.0667	0.4498	0.1978	0.7343	0.2894	0.6045	0.2441	
AGE GROUP									
12 to 17	0.0000	0.0005	0.0061	0.0408	0.0212	0.3393	0.3890	0.2350	
18 to 25	0.5016	0.8601	0.2986	0.0655	0.1120	0.3273	0.0981	0.1066	
26 or Older	0.4291	0.0115	0.1163	0.0139	0.5541	0.0952	0.8564	0.3912	
GENDER									
Male	0.7314	0.0271	0.1406	0.0181	0.7694	0.3055	0.7241	0.3044	
Female	0.9408	0.8869	0.4701	0.3743	0.8961	0.7263	0.6984	0.5546	
PAST YEAR USE									
TOTAL	0.0028	0.1982	0.0851	0.0753	0.0027	0.0203	0.7020	0.0739	
AGE GROUP									
12 to 17	0.0000	0.0001	0.0004	0.0044	0.0041	0.1189	0.3238	0.7720	
18 to 25	0.0030	0.0147	0.0293	0.0184	0.0228	0.0075	0.0453	0.0467	
26 or Older	0.8785	0.1875	0.9147	0.8357	0.0616	0.3168	0.3897	0.1929	
GENDER									
Male	0.0380	0.9036	0.3745	0.7263	0.0130	0.3562	0.3862	0.1123	
Female	0.0274	0.0831	0.0840	0.0239	0.0880	0.0099	0.1649	0.3780	
PAST MONTH USE									
TOTAL	0.1725	0.2707	0.3142	0.3150	0.1185	0.4902	0.4534	0.2801	
AGE GROUP									
12 to 17	0.0006	0.0010	0.0154	0.0063	0.0391	0.3485	0.1932	0.6624	
18 to 25	0.9710	0.3581	0.9830	0.5673	0.5552	0.4956	0.8632	0.9952	
26 or Older	0.6086	0.7953	0.6672	0.8365	0.3085	0.4522	0.6697	0.2870	
GENDER									
Male	0.1021	0.4685	0.3922	0.6570	0.2477	0.8057	0.7029	0.3858	
Female	0.9661	0.3837	0.5820	0.3257	0.2828	0.4434	0.2270	0.5131	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.26C Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	145	140	136	151	166	152	161	177	148
AGE GROUP									
12 to 17	14	12	12	12	15	13	10	8	13
18 to 25	50	47	40	38	39	41	40	41	42
26 or Older	133	129	126	145	160	147	156	172	139
GENDER									
Male	123	115	108	124	152	120	138	139	127
Female	69	68	73	76	64	90	75	103	68
PAST YEAR USE									
TOTAL	38	31	28	32	36	49	42	41	35
AGE GROUP									
12 to 17	8	7	9	8	11	11	7	6	8
18 to 25	14	19	14	16	22	26	17	19	18
26 or Older	34	24	23	27	26	40	40	36	29
GENDER									
Male	35	27	25	23	33	39	37	37	29
Female	14	16	11	23	14	28	21	18	18
PAST MONTH USE									
TOTAL	18	11	7	24	22	26	14	24	10
AGE GROUP									
12 to 17	4	5	5	6	5	9	4	2	3
18 to 25	6	6	6	9	8	9	6	7	6
26 or Older	16	8	*	21	20	22	12	23	7
GENDER									
Male	14	10	6	13	21	12	6	24	8
Female	10	4	4	20	8	23	12	5	5

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.26D Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.06	0.06	0.05	0.06	0.07	0.06	0.06	0.07	0.06
AGE GROUP									
12 to 17	0.05	0.05	0.05	0.05	0.06	0.05	0.04	0.03	0.05
18 to 25	0.15	0.14	0.12	0.11	0.11	0.12	0.11	0.12	0.12
26 or Older	0.07	0.07	0.07	0.07	0.08	0.07	0.08	0.08	0.07
GENDER									
Male	0.10	0.10	0.09	0.10	0.12	0.10	0.11	0.11	0.10
Female	0.05	0.05	0.06	0.06	0.05	0.07	0.06	0.08	0.05
PAST YEAR USE									
TOTAL	0.02	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.01
AGE GROUP									
12 to 17	0.03	0.03	0.04	0.03	0.05	0.04	0.03	0.03	0.03
18 to 25	0.04	0.06	0.04	0.05	0.07	0.07	0.05	0.06	0.05
26 or Older	0.02	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.01
GENDER									
Male	0.03	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.02
Female	0.01	0.01	0.01	0.02	0.01	0.02	0.02	0.01	0.01
PAST MONTH USE									
TOTAL	0.01	0.00	0.00	0.01	0.01	0.01	0.01	0.01	0.00
AGE GROUP									
12 to 17	0.02	0.02	0.02	0.03	0.02	0.04	0.02	0.01	0.01
18 to 25	0.02	0.02	0.02	0.03	0.02	0.03	0.02	0.02	0.02
26 or Older	0.01	0.00	*	0.01	0.01	0.01	0.01	0.01	0.00
GENDER									
Male	0.01	0.01	0.00	0.01	0.02	0.01	0.01	0.02	0.01
Female	0.01	0.00	0.00	0.02	0.01	0.02	0.01	0.00	0.00

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.26P Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2006-2013

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.0055	0.0007	0.0002	0.0039	0.0184	0.0048	0.1282	0.1822	
AGE GROUP									
12 to 17	0.6371	0.8706	0.9648	0.7400	0.6604	0.4182	0.6913	0.0307	
18 to 25	0.0000	0.0000	0.0282	0.1836	0.6409	0.6185	0.4807	0.9867	
26 or Older	0.0000	0.0000	0.0000	0.0007	0.0085	0.0042	0.1768	0.2179	
GENDER									
Male	0.0046	0.0013	0.0001	0.0088	0.1611	0.0047	0.4205	0.0451	
Female	0.4467	0.1388	0.2782	0.1499	0.0069	0.3547	0.0735	0.5706	
PAST YEAR USE									
TOTAL	0.5977	0.6750	0.0587	0.4893	0.8406	0.1093	0.4848	0.5885	
AGE GROUP									
12 to 17	0.5148	0.5572	0.4349	0.3328	0.0543	0.0313	0.3038	0.9702	
18 to 25	0.2873	0.3414	0.1895	0.7302	0.2304	0.1149	0.8334	0.2171	
26 or Older	0.7858	0.1631	0.0728	0.3522	0.1711	0.7249	0.5228	0.9255	
GENDER									
Male	0.8610	0.8559	0.2179	0.2997	0.6272	0.1486	0.2877	0.6271	
Female	0.3865	0.5900	0.0671	0.8592	0.5933	0.4099	0.6854	0.7961	
PAST MONTH USE									
TOTAL	0.2590	0.5734	0.7150	0.0317	0.2282	0.0698	0.2144	0.3280	
AGE GROUP									
12 to 17	0.1979	0.1806	0.3008	0.0756	0.1032	0.0264	0.4822	0.8827	
18 to 25	0.9686	0.7059	0.7330	0.2564	0.3961	0.2017	0.3716	0.4791	
26 or Older	0.3860	0.7674	*	0.1731	0.5973	0.5259	0.4876	0.4521	
GENDER									
Male	0.2555	0.3853	0.9836	0.0756	0.3048	0.1441	0.7634	0.1985	
Female	0.7187	0.7025	0.5086	0.1668	0.4974	0.2120	0.1833	0.3987	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

**Table 7.27C DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Standard Errors of Numbers in Thousands, 2006-2014**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	62	71	74	79	101	84	108	127	118
AGE GROUP									
12 to 17	9	10	10	10	10	10	12	11	14
18 to 25	29	24	28	33	36	36	43	51	52
26 or Older	54	65	68	73	94	74	98	122	102
GENDER									
Male	55	66	60	71	87	70	92	117	97
Female	29	23	42	33	53	50	53	58	63
PAST YEAR USE									
TOTAL	21	16	27	29	35	32	36	40	43
AGE GROUP									
12 to 17	4	6	8	8	8	7	9	9	11
18 to 25	15	12	14	19	16	24	25	28	23
26 or Older	14	7	21	21	31	19	29	27	33
GENDER									
Male	21	14	23	19	28	27	33	35	40
Female	5	6	14	22	21	16	13	21	14
PAST MONTH USE									
TOTAL	7	*	18	20	28	17	16	12	21
AGE GROUP									
12 to 17	*	*	4	4	7	3	6	5	5
18 to 25	5	5	9	4	7	8	11	11	10
26 or Older	*	*	15	19	27	14	9	*	16
GENDER									
Male	6	4	17	8	22	8	13	12	20
Female	4	*	*	18	18	15	9	4	5

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine. AMT is also known as alpha-methyltryptamine. The drug 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

**Table 7.27D DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Standard Errors of Percentages, 2006-2014**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.03	0.03	0.03	0.03	0.04	0.03	0.04	0.05	0.04
AGE GROUP									
12 to 17	0.04	0.04	0.04	0.04	0.04	0.04	0.05	0.04	0.06
18 to 25	0.09	0.07	0.09	0.10	0.10	0.10	0.13	0.15	0.15
26 or Older	0.03	0.03	0.04	0.04	0.05	0.04	0.05	0.06	0.05
GENDER									
Male	0.05	0.05	0.05	0.06	0.07	0.06	0.07	0.09	0.08
Female	0.02	0.02	0.03	0.03	0.04	0.04	0.04	0.04	0.05
PAST YEAR USE									
TOTAL	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02
AGE GROUP									
12 to 17	0.02	0.02	0.03	0.03	0.03	0.03	0.04	0.04	0.05
18 to 25	0.05	0.04	0.04	0.06	0.05	0.07	0.07	0.08	0.07
26 or Older	0.01	0.00	0.01	0.01	0.02	0.01	0.01	0.01	0.02
GENDER									
Male	0.02	0.01	0.02	0.02	0.02	0.02	0.03	0.03	0.03
Female	0.00	0.00	0.01	0.02	0.02	0.01	0.01	0.02	0.01
PAST MONTH USE									
TOTAL	0.00	*	0.01	0.01	0.01	0.01	0.01	0.00	0.01
AGE GROUP									
12 to 17	*	*	0.02	0.02	0.03	0.01	0.02	0.02	0.02
18 to 25	0.02	0.01	0.03	0.01	0.02	0.02	0.03	0.03	0.03
26 or Older	*	*	0.01	0.01	0.01	0.01	0.00	*	0.01
GENDER									
Male	0.00	0.00	0.01	0.01	0.02	0.01	0.01	0.01	0.02
Female	0.00	*	*	0.01	0.01	0.01	0.01	0.00	0.00

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine. AMT is also known as alpha-methyltryptamine. The drug 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

**Table 7.27P DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
P Values from Tests of Differences of Percentages, 2014 Versus 2006-2013**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0076	0.3153	
AGE GROUP									
12 to 17	0.0533	0.0521	0.1089	0.2992	0.4050	0.4592	0.2620	0.7252	
18 to 25	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0367	0.6490	
26 or Older	0.0000	0.0000	0.0000	0.0000	0.0141	0.0000	0.0273	0.1733	
GENDER									
Male	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0209	0.2937	
Female	0.0000	0.0000	0.0036	0.0007	0.0682	0.0041	0.1454	0.8612	
PAST YEAR USE									
TOTAL	0.0000	0.0000	0.0000	0.0005	0.0106	0.0306	0.3054	0.9203	
AGE GROUP									
12 to 17	0.0135	0.0428	0.2497	0.6099	0.6272	0.4713	0.6298	0.5869	
18 to 25	0.0000	0.0000	0.0000	0.0132	0.0141	0.8911	0.7931	0.1516	
26 or Older	0.0004	0.0000	0.0101	0.0080	0.1248	0.0061	0.0802	0.1849	
GENDER									
Male	0.0000	0.0000	0.0000	0.0000	0.0012	0.0234	0.1860	0.4939	
Female	0.0097	0.0076	0.3516	0.4292	0.5286	0.7788	0.5419	0.0937	
PAST MONTH USE									
TOTAL	0.0072	*	0.2482	0.4738	0.9327	0.2166	0.8514	0.1696	
AGE GROUP									
12 to 17	*	*	0.5427	0.9399	0.5211	0.4366	0.4926	0.7907	
18 to 25	0.1060	0.0346	0.3036	0.0479	0.2870	0.4857	0.4249	0.7724	
26 or Older	*	*	0.5143	0.9761	0.7567	0.3552	0.2101	*	
GENDER									
Male	0.0039	0.0011	0.2599	0.0194	0.4926	0.0339	0.2807	0.1527	
Female	0.9042	*	*	0.1583	0.2211	0.4556	0.0424	0.8812	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine. AMT is also known as alpha-methyltryptamine. The drug 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.28C Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	92	110	129	159	189	175	182	223	178
AGE GROUP									
12 to 17	20	23	27	28	28	25	23	22	20
18 to 25	62	68	75	89	95	92	94	88	91
26 or Older	59	78	98	120	155	139	152	203	150
GENDER									
Male	81	91	104	142	159	142	167	191	150
Female	34	54	62	63	99	97	77	102	81
PAST YEAR USE									
TOTAL	48	68	70	72	77	52	51	86	44
AGE GROUP									
12 to 17	16	18	23	22	22	19	16	13	15
18 to 25	38	45	47	47	42	39	33	25	26
26 or Older	22	43	43	46	59	29	34	80	33
GENDER									
Male	41	55	55	58	73	43	46	81	35
Female	21	34	37	37	28	28	23	27	26
PAST MONTH USE									
TOTAL	18	26	24	28	28	20	26	74	12
AGE GROUP									
12 to 17	7	9	10	9	9	8	7	5	6
18 to 25	16	21	21	15	17	18	11	11	10
26 or Older	*	11	8	22	20	7	22	73	*
GENDER									
Male	17	22	22	19	27	17	24	72	9
Female	7	11	8	20	9	10	10	14	8

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.28D Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2014

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.04	0.04	0.05	0.06	0.07	0.07	0.07	0.09	0.07
AGE GROUP									
12 to 17	0.08	0.09	0.11	0.12	0.12	0.10	0.09	0.09	0.08
18 to 25	0.19	0.21	0.23	0.26	0.28	0.27	0.27	0.25	0.26
26 or Older	0.03	0.04	0.05	0.06	0.08	0.07	0.08	0.10	0.07
GENDER									
Male	0.07	0.08	0.09	0.12	0.13	0.11	0.13	0.15	0.12
Female	0.03	0.04	0.05	0.05	0.08	0.07	0.06	0.08	0.06
PAST YEAR USE									
TOTAL	0.02	0.03	0.03	0.03	0.03	0.02	0.02	0.03	0.02
AGE GROUP									
12 to 17	0.06	0.07	0.09	0.09	0.09	0.07	0.06	0.05	0.06
18 to 25	0.12	0.14	0.14	0.14	0.12	0.11	0.10	0.07	0.07
26 or Older	0.01	0.02	0.02	0.02	0.03	0.01	0.02	0.04	0.02
GENDER									
Male	0.03	0.05	0.05	0.05	0.06	0.03	0.04	0.06	0.03
Female	0.02	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02
PAST MONTH USE									
TOTAL	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.03	0.00
AGE GROUP									
12 to 17	0.03	0.03	0.04	0.04	0.04	0.03	0.03	0.02	0.02
18 to 25	0.05	0.06	0.06	0.04	0.05	0.05	0.03	0.03	0.03
26 or Older	*	0.01	0.00	0.01	0.01	0.00	0.01	0.04	*
GENDER									
Male	0.01	0.02	0.02	0.02	0.02	0.01	0.02	0.06	0.01
Female	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.28P Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2006-2013

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011	2012	2013	2014
LIFETIME USE									
TOTAL	0.0000	0.0000	0.0000	0.0179	0.3390	0.3828	0.6787	0.3738	
AGE GROUP									
12 to 17	0.0028	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0583	
18 to 25	0.0000	0.0001	0.4485	0.0000	0.0000	0.0000	0.0000	0.0049	
26 or Older	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0015	0.5078	
GENDER									
Male	0.0000	0.0000	0.0000	0.1748	0.3344	0.9918	0.5970	0.7176	
Female	0.0000	0.0000	0.0001	0.0046	0.8453	0.0801	0.9160	0.2053	
PAST YEAR USE									
TOTAL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0802	
AGE GROUP									
12 to 17	0.0004	0.0000	0.0000	0.0000	0.0000	0.0000	0.0063	0.5577	
18 to 25	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0981	
26 or Older	0.1964	0.3473	0.0578	0.0095	0.0029	0.5173	0.5884	0.2699	
GENDER									
Male	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.1043	
Female	0.0925	0.0001	0.0000	0.0000	0.0001	0.0006	0.0401	0.5049	
PAST MONTH USE									
TOTAL	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0056	0.1745	
AGE GROUP									
12 to 17	0.0609	0.0028	0.0000	0.0002	0.0018	0.0044	0.1113	0.5740	
18 to 25	0.0000	0.0000	0.0000	0.0000	0.0003	0.0022	0.0997	0.4013	
26 or Older	*	*	*	*	*	*	*	*	
GENDER									
Male	0.0000	0.0000	0.0000	0.0000	0.0002	0.0007	0.0167	0.2149	
Female	0.3598	0.0104	0.0413	0.0351	0.2469	0.0475	0.1591	0.4920	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2014.

Table 7.29C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	832	803	794	845	850	844	871	831	852	846	882	835	736
Cigarettes	795	789	768	777	807	809	810	797	780	816	839	798	700
Smokeless Tobacco	279	284	245	267	284	283	312	291	314	278	298	300	249
Cigars	350	331	323	354	352	337	366	361	363	373	388	354	282
Pipe Tobacco	166	131	147	146	167	170	147	151	147	144	184	151	139
ALCOHOL	989	919	957	975	968	1,019	983	957	990	1,006	1,016	1,076	869
Binge Alcohol Use ²	732	682	698	701	746	770	735	783	752	734	797	804	690
Heavy Alcohol Use ²	392	380	382	379	427	429	414	406	422	423	448	438	368
MALE													
TOBACCO PRODUCTS¹	572	573	570	576	585	598	628	602	626	581	618	612	513
Cigarettes	527	545	537	515	545	577	565	565	536	537	556	549	477
Smokeless Tobacco	264	256	234	241	273	252	296	283	302	264	281	282	240
Cigars	327	305	290	324	320	312	333	336	321	336	351	312	256
Pipe Tobacco	139	127	139	141	160	168	131	144	137	135	170	146	131
ALCOHOL	656	599	626	621	610	666	632	651	675	674	675	693	553
Binge Alcohol Use ²	540	517	513	529	559	585	586	605	577	567	625	607	519
Heavy Alcohol Use ²	346	321	319	321	354	374	357	349	353	355	382	367	311
FEMALE													
TOBACCO PRODUCTS¹	513	507	497	543	511	511	558	505	535	542	554	524	444
Cigarettes	501	488	491	538	503	493	553	498	526	532	539	513	438
Smokeless Tobacco	100	126	65	87	76	113	75	51	78	82	103	94	51
Cigars	114	123	122	112	131	116	114	130	153	142	146	161	111
Pipe Tobacco	90	35	43	42	46	30	55	37	54	46	61	35	43
ALCOHOL	644	600	664	637	645	676	673	687	635	684	686	736	620
Binge Alcohol Use ²	415	382	390	402	399	395	398	441	441	441	427	463	380
Heavy Alcohol Use ²	177	174	174	163	169	171	192	193	201	207	203	232	165

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.29D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Percentages, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	0.35	0.34	0.33	0.35	0.35	0.34	0.35	0.33	0.34	0.33	0.34	0.32	0.28
Cigarettes	0.34	0.33	0.32	0.32	0.33	0.33	0.32	0.32	0.31	0.32	0.32	0.30	0.26
Smokeless Tobacco	0.12	0.12	0.10	0.11	0.12	0.11	0.12	0.12	0.12	0.11	0.11	0.11	0.09
Cigars	0.15	0.14	0.13	0.15	0.14	0.14	0.15	0.14	0.14	0.14	0.15	0.14	0.11
Pipe Tobacco	0.07	0.06	0.06	0.06	0.07	0.07	0.06	0.06	0.06	0.06	0.07	0.06	0.05
ALCOHOL	0.42	0.39	0.40	0.40	0.39	0.41	0.39	0.38	0.39	0.39	0.39	0.41	0.33
Binge Alcohol Use ²	0.31	0.29	0.29	0.29	0.30	0.31	0.29	0.31	0.30	0.29	0.31	0.31	0.26
Heavy Alcohol Use ²	0.17	0.16	0.16	0.16	0.17	0.17	0.17	0.16	0.17	0.16	0.17	0.17	0.14
MALE													
TOBACCO PRODUCTS¹	0.50	0.50	0.49	0.49	0.49	0.50	0.52	0.49	0.51	0.47	0.49	0.48	0.40
Cigarettes	0.46	0.47	0.46	0.44	0.46	0.48	0.47	0.46	0.43	0.43	0.44	0.43	0.37
Smokeless Tobacco	0.23	0.22	0.20	0.20	0.23	0.21	0.24	0.23	0.25	0.21	0.22	0.22	0.19
Cigars	0.29	0.27	0.25	0.27	0.27	0.26	0.27	0.27	0.26	0.27	0.28	0.25	0.20
Pipe Tobacco	0.12	0.11	0.12	0.12	0.13	0.14	0.11	0.12	0.11	0.11	0.14	0.12	0.10
ALCOHOL	0.58	0.52	0.54	0.53	0.51	0.55	0.52	0.53	0.55	0.54	0.54	0.55	0.43
Binge Alcohol Use ²	0.48	0.45	0.44	0.45	0.47	0.49	0.48	0.49	0.47	0.45	0.50	0.48	0.40
Heavy Alcohol Use ²	0.30	0.28	0.27	0.27	0.30	0.31	0.29	0.29	0.29	0.28	0.30	0.29	0.24
FEMALE													
TOBACCO PRODUCTS¹	0.42	0.41	0.40	0.43	0.40	0.40	0.43	0.39	0.41	0.41	0.41	0.39	0.32
Cigarettes	0.41	0.40	0.40	0.43	0.40	0.39	0.43	0.38	0.40	0.40	0.40	0.38	0.32
Smokeless Tobacco	0.08	0.10	0.05	0.07	0.06	0.09	0.06	0.04	0.06	0.06	0.08	0.07	0.04
Cigars	0.09	0.10	0.10	0.09	0.10	0.09	0.09	0.10	0.12	0.11	0.11	0.12	0.08
Pipe Tobacco	0.07	0.03	0.03	0.03	0.04	0.02	0.04	0.03	0.04	0.03	0.05	0.03	0.03
ALCOHOL	0.53	0.49	0.54	0.51	0.51	0.53	0.52	0.53	0.49	0.51	0.51	0.54	0.45
Binge Alcohol Use ²	0.34	0.31	0.31	0.32	0.32	0.31	0.31	0.34	0.34	0.33	0.32	0.34	0.28
Heavy Alcohol Use ²	0.15	0.14	0.14	0.13	0.13	0.13	0.15	0.15	0.15	0.16	0.15	0.17	0.12

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.29P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0037	0.0007	0.5459	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0032	0.0021	0.2954	
Smokeless Tobacco	0.7700	0.9096	0.0348	0.4502	0.6079	0.9164	0.1640	0.3010	0.1336	0.6383	0.1621	0.5359	
Cigars	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0075	0.0004	0.1856	
Pipe Tobacco	0.3941	0.0293	0.2960	0.5068	0.2757	0.7934	0.2394	0.8804	0.9777	0.8176	0.1555	0.8172	
ALCOHOL	0.0012	0.0000	0.0000	0.0943	0.0008	0.0045	0.0333	0.1097	0.0832	0.0767	0.2565	0.3203	
Binge Alcohol Use ²	0.7829	0.3487	0.5506	0.3874	0.8954	0.3765	0.3357	0.0770	0.8122	0.3812	0.9304	0.8131	
Heavy Alcohol Use ²	0.0076	0.0031	0.0003	0.0403	0.0011	0.0016	0.0003	0.0021	0.0216	0.9797	0.0857	0.5200	
MALE													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0001	0.0565	0.0032	1.0000	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0001	0.0502	0.0142	0.4719	
Smokeless Tobacco	0.9627	0.4547	0.0263	0.2704	0.5481	0.7999	0.1798	0.2844	0.2059	0.4159	0.2171	0.8091	
Cigars	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0007	0.0352	0.0016	0.4990	
Pipe Tobacco	0.6480	0.2766	0.8703	0.1808	0.0607	0.5222	0.3008	0.6077	0.8537	0.9723	0.1599	0.5303	
ALCOHOL	0.8640	0.9333	0.6165	0.2354	0.6887	0.3701	0.5822	0.6986	0.9389	0.4532	0.2870	0.8148	
Binge Alcohol Use ²	0.0625	0.1441	0.0861	0.4043	0.0404	0.0082	0.0109	0.0129	0.1770	0.9270	0.5758	0.8169	
Heavy Alcohol Use ²	0.0002	0.0029	0.0003	0.0050	0.0002	0.0007	0.0003	0.0058	0.0333	0.8668	0.1179	0.5640	
FEMALE													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0103	0.0304	0.3309	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0098	0.0241	0.3843	
Smokeless Tobacco	0.2086	0.1044	0.9071	0.3713	0.6894	0.7786	0.7569	0.7641	0.4109	0.1997	0.4248	0.1602	
Cigars	0.8055	0.0399	0.2312	0.5064	0.0192	0.5016	0.9779	0.0380	0.0239	0.0401	0.0781	0.0969	
Pipe Tobacco	0.3554	0.0001	0.0028	0.0363	0.0141	0.0001	0.4832	0.0167	0.6072	0.4159	0.7093	0.1678	
ALCOHOL	0.0000	0.0000	0.0000	0.0003	0.0000	0.0009	0.0003	0.0087	0.0067	0.0680	0.5298	0.2246	
Binge Alcohol Use ²	0.0040	0.0003	0.0008	0.0075	0.0081	0.0296	0.0523	0.7474	0.1270	0.1756	0.3481	0.4647	
Heavy Alcohol Use ²	0.2390	0.3076	0.1853	0.4342	0.6722	0.5943	0.2273	0.1409	0.3398	0.7518	0.3761	0.7538	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.30C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Numbers in Thousands, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	81	80	80	78	72	76	69	71	69	68	61	59	61
Cigarettes	74	73	76	71	66	67	61	64	63	60	55	51	52
Smokeless Tobacco	27	30	32	29	31	36	30	32	30	30	31	31	34
Cigars	47	44	44	46	41	45	41	39	36	41	32	32	34
Pipe Tobacco	14	18	19	17	18	19	17	21	17	18	18	17	20
ALCOHOL	80	82	81	81	82	87	79	79	80	77	78	73	81
Binge Alcohol Use ²	67	67	73	65	69	69	57	62	62	55	56	54	60
Heavy Alcohol Use ²	31	33	36	32	33	37	30	30	28	25	25	23	24
MALE													
TOBACCO PRODUCTS¹	59	56	55	59	54	57	50	52	52	53	47	45	47
Cigarettes	53	47	52	53	46	48	42	44	45	46	41	36	36
Smokeless Tobacco	26	29	30	28	29	34	28	30	28	28	29	28	32
Cigars	37	36	37	38	34	37	34	33	31	31	26	28	26
Pipe Tobacco	13	17	15	14	15	16	12	18	14	14	12	12	14
ALCOHOL	55	57	59	57	57	61	53	58	57	53	55	52	54
Binge Alcohol Use ²	46	46	52	48	47	51	41	48	44	42	41	40	43
Heavy Alcohol Use ²	24	23	27	27	26	28	23	23	23	22	19	18	18
FEMALE													
TOBACCO PRODUCTS¹	51	52	53	50	47	48	46	46	43	43	37	40	37
Cigarettes	49	50	52	48	46	46	44	44	40	40	34	37	34
Smokeless Tobacco	9	6	11	8	10	8	10	9	12	10	8	9	12
Cigars	26	23	25	24	24	22	21	25	20	24	19	16	20
Pipe Tobacco	9	7	13	11	10	11	12	11	10	12	12	12	14
ALCOHOL	58	60	59	58	59	61	56	52	54	53	54	52	57
Binge Alcohol Use ²	46	46	50	43	49	45	42	40	43	37	38	37	41
Heavy Alcohol Use ²	19	23	23	20	21	23	20	19	17	15	16	16	16

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.30D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Percentages, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	0.33	0.32	0.32	0.31	0.29	0.30	0.28	0.29	0.28	0.27	0.25	0.24	0.25
Cigarettes	0.30	0.29	0.30	0.28	0.26	0.27	0.25	0.26	0.26	0.24	0.22	0.20	0.21
Smokeless Tobacco	0.11	0.12	0.13	0.12	0.12	0.14	0.12	0.13	0.12	0.12	0.12	0.12	0.14
Cigars	0.19	0.17	0.18	0.18	0.16	0.18	0.16	0.16	0.15	0.16	0.13	0.13	0.13
Pipe Tobacco	0.06	0.07	0.08	0.07	0.07	0.08	0.07	0.09	0.07	0.07	0.07	0.07	0.08
ALCOHOL	0.32	0.33	0.32	0.32	0.32	0.34	0.32	0.32	0.33	0.31	0.31	0.29	0.33
Binge Alcohol Use ²	0.27	0.27	0.29	0.26	0.27	0.27	0.23	0.25	0.25	0.22	0.22	0.22	0.24
Heavy Alcohol Use ²	0.12	0.13	0.14	0.13	0.13	0.15	0.12	0.12	0.11	0.10	0.10	0.09	0.10
MALE													
TOBACCO PRODUCTS¹	0.47	0.44	0.43	0.46	0.41	0.45	0.39	0.42	0.42	0.41	0.37	0.35	0.37
Cigarettes	0.42	0.37	0.40	0.41	0.36	0.37	0.33	0.35	0.36	0.36	0.32	0.28	0.28
Smokeless Tobacco	0.21	0.23	0.23	0.22	0.22	0.26	0.22	0.24	0.23	0.22	0.23	0.22	0.25
Cigars	0.30	0.28	0.28	0.29	0.26	0.29	0.27	0.27	0.25	0.25	0.20	0.22	0.21
Pipe Tobacco	0.10	0.13	0.11	0.11	0.12	0.12	0.10	0.14	0.11	0.11	0.09	0.10	0.11
ALCOHOL	0.43	0.45	0.45	0.44	0.44	0.47	0.41	0.46	0.46	0.42	0.43	0.41	0.42
Binge Alcohol Use ²	0.37	0.36	0.40	0.37	0.36	0.40	0.32	0.38	0.35	0.33	0.32	0.32	0.34
Heavy Alcohol Use ²	0.19	0.18	0.21	0.21	0.20	0.22	0.18	0.18	0.18	0.17	0.15	0.14	0.14
FEMALE													
TOBACCO PRODUCTS¹	0.42	0.42	0.43	0.40	0.38	0.39	0.38	0.38	0.36	0.35	0.30	0.33	0.30
Cigarettes	0.40	0.41	0.42	0.39	0.37	0.37	0.36	0.37	0.34	0.33	0.28	0.30	0.28
Smokeless Tobacco	0.08	0.05	0.09	0.07	0.08	0.07	0.08	0.08	0.11	0.08	0.06	0.07	0.10
Cigars	0.21	0.19	0.20	0.19	0.19	0.18	0.17	0.21	0.17	0.20	0.16	0.13	0.16
Pipe Tobacco	0.07	0.06	0.10	0.09	0.08	0.09	0.10	0.09	0.09	0.10	0.10	0.10	0.11
ALCOHOL	0.48	0.49	0.48	0.47	0.47	0.49	0.46	0.43	0.45	0.43	0.44	0.43	0.47
Binge Alcohol Use ²	0.38	0.38	0.41	0.35	0.39	0.37	0.35	0.33	0.36	0.30	0.31	0.30	0.33
Heavy Alcohol Use ²	0.16	0.19	0.19	0.16	0.17	0.18	0.16	0.15	0.14	0.12	0.13	0.13	0.13

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.30P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0151	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0115	
Smokeless Tobacco	1.0000	0.6797	0.1101	0.5953	0.0293	0.0133	0.1499	0.0331	0.0434	0.5697	0.5905	0.9525	
Cigars	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0081	0.2399	
Pipe Tobacco	0.0899	0.4591	0.8955	0.3545	0.5857	0.9801	0.9422	0.0639	0.4304	0.9327	0.5713	0.4508	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0028	0.9818	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0006	0.5992	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0526	0.3006	
MALE													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0007	0.0652	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.1600	
Smokeless Tobacco	0.6125	0.1692	0.0251	0.2386	0.0057	0.0019	0.0534	0.0117	0.0358	0.3273	0.2330	0.6309	
Cigars	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0039	0.0726	
Pipe Tobacco	0.6498	0.2941	0.5702	0.6270	0.5013	0.4115	0.9485	0.0039	0.8943	0.8910	0.5465	0.5982	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0027	0.4485	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0020	0.0359	0.6050	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0026	0.2659	0.3055	
FEMALE													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0008	0.1099	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0259	
Smokeless Tobacco	0.1660	0.0024	0.1507	0.0739	0.2123	0.1149	0.2741	0.4271	0.9638	0.2807	0.0582	0.1203	
Cigars	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.0036	0.0000	0.0116	0.0004	0.5509	0.5698	
Pipe Tobacco	0.0441	0.0057	0.4223	0.0550	0.0923	0.3001	0.8657	0.4621	0.1694	0.9803	0.8290	0.5858	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0015	0.0626	0.1087	0.1756	0.4865	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0011	0.0089	0.0045	0.8255	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0000	0.0562	0.1234	0.0916	0.6926	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.31C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Numbers in Thousands, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	150	151	161	155	160	162	153	169	166	167	164	170	189
Cigarettes	150	151	159	152	156	159	149	162	160	162	161	162	185
Smokeless Tobacco	59	56	60	64	65	63	68	78	81	70	70	71	82
Cigars	83	84	97	92	94	92	96	99	101	100	93	100	105
Pipe Tobacco	24	25	30	37	33	32	34	40	41	48	39	49	45
ALCOHOL	164	160	164	166	168	170	163	174	169	184	170	183	197
Binge Alcohol Use ²	161	155	166	171	169	169	167	182	168	189	175	182	197
Heavy Alcohol Use ²	112	114	120	120	125	114	124	128	127	125	121	107	125
MALE													
TOBACCO PRODUCTS¹	100	106	106	110	109	116	109	121	121	113	118	121	137
Cigarettes	99	109	105	107	108	116	106	121	113	111	113	117	132
Smokeless Tobacco	56	55	58	60	62	61	63	73	75	65	65	67	78
Cigars	73	72	83	79	78	81	80	81	84	89	80	86	85
Pipe Tobacco	22	21	27	33	30	30	29	35	33	39	34	41	37
ALCOHOL	111	100	110	101	108	110	109	117	112	119	118	121	130
Binge Alcohol Use ²	110	105	113	111	115	116	118	130	121	126	124	121	133
Heavy Alcohol Use ²	87	91	88	96	95	94	95	99	101	87	97	85	95
FEMALE													
TOBACCO PRODUCTS¹	101	93	102	101	102	97	101	98	96	105	104	101	114
Cigarettes	102	92	101	101	98	95	100	96	97	102	102	97	109
Smokeless Tobacco	8	11	10	12	13	13	14	18	20	14	14	18	16
Cigars	38	43	44	41	44	43	47	45	49	50	46	46	54
Pipe Tobacco	12	12	12	16	11	14	17	15	23	26	19	22	23
ALCOHOL	102	114	104	112	112	114	107	110	109	123	118	117	129
Binge Alcohol Use ²	99	100	96	104	103	110	102	105	101	116	110	112	126
Heavy Alcohol Use ²	55	59	60	56	63	60	65	61	59	72	63	65	66

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.31D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Percentages, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	0.48	0.48	0.50	0.48	0.49	0.50	0.47	0.50	0.49	0.49	0.47	0.49	0.54
Cigarettes	0.48	0.47	0.49	0.47	0.48	0.49	0.45	0.48	0.47	0.47	0.47	0.46	0.53
Smokeless Tobacco	0.19	0.18	0.19	0.20	0.20	0.19	0.20	0.23	0.24	0.20	0.20	0.20	0.23
Cigars	0.27	0.26	0.30	0.28	0.29	0.28	0.29	0.29	0.30	0.29	0.27	0.29	0.30
Pipe Tobacco	0.08	0.08	0.09	0.11	0.10	0.10	0.10	0.12	0.12	0.14	0.11	0.14	0.13
ALCOHOL	0.53	0.50	0.51	0.51	0.51	0.52	0.49	0.52	0.50	0.54	0.49	0.53	0.56
Binge Alcohol Use ²	0.52	0.49	0.52	0.53	0.52	0.52	0.51	0.54	0.49	0.55	0.51	0.52	0.57
Heavy Alcohol Use ²	0.36	0.36	0.37	0.37	0.38	0.35	0.38	0.38	0.37	0.36	0.35	0.31	0.36
MALE													
TOBACCO PRODUCTS¹	0.64	0.67	0.66	0.67	0.66	0.70	0.66	0.72	0.70	0.66	0.68	0.70	0.78
Cigarettes	0.64	0.69	0.65	0.66	0.66	0.70	0.64	0.71	0.65	0.65	0.65	0.67	0.75
Smokeless Tobacco	0.36	0.34	0.36	0.37	0.38	0.37	0.38	0.43	0.43	0.38	0.38	0.38	0.44
Cigars	0.47	0.45	0.51	0.48	0.48	0.49	0.48	0.48	0.49	0.52	0.46	0.50	0.49
Pipe Tobacco	0.14	0.13	0.17	0.20	0.18	0.18	0.17	0.21	0.19	0.23	0.20	0.24	0.21
ALCOHOL	0.71	0.63	0.68	0.62	0.66	0.67	0.66	0.69	0.65	0.69	0.68	0.69	0.74
Binge Alcohol Use ²	0.71	0.66	0.70	0.68	0.70	0.70	0.71	0.77	0.70	0.73	0.71	0.69	0.76
Heavy Alcohol Use ²	0.56	0.57	0.54	0.59	0.58	0.57	0.57	0.59	0.59	0.51	0.56	0.49	0.54
FEMALE													
TOBACCO PRODUCTS¹	0.65	0.59	0.64	0.63	0.63	0.59	0.62	0.59	0.57	0.61	0.60	0.58	0.66
Cigarettes	0.66	0.58	0.63	0.63	0.60	0.58	0.61	0.58	0.58	0.60	0.59	0.56	0.62
Smokeless Tobacco	0.05	0.07	0.06	0.08	0.08	0.08	0.08	0.11	0.12	0.08	0.08	0.11	0.09
Cigars	0.25	0.27	0.27	0.26	0.27	0.26	0.29	0.27	0.29	0.29	0.27	0.27	0.31
Pipe Tobacco	0.08	0.08	0.08	0.10	0.07	0.08	0.10	0.09	0.14	0.15	0.11	0.13	0.13
ALCOHOL	0.66	0.72	0.65	0.69	0.69	0.70	0.65	0.66	0.65	0.72	0.68	0.68	0.74
Binge Alcohol Use ²	0.64	0.63	0.60	0.65	0.64	0.68	0.62	0.63	0.60	0.68	0.64	0.64	0.72
Heavy Alcohol Use ²	0.35	0.37	0.37	0.35	0.39	0.37	0.39	0.36	0.35	0.42	0.37	0.38	0.38

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.31P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0067	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0013	
Smokeless Tobacco	0.0097	0.0014	0.0218	0.1017	0.2122	0.2189	0.3824	0.1268	0.0192	0.5664	0.7554	0.5588	
Cigars	0.0013	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0063	0.0105	0.5116	
Pipe Tobacco	0.0000	0.0000	0.0000	0.0086	0.0004	0.0000	0.0023	0.4209	0.6145	0.7762	0.5628	0.1539	
ALCOHOL	0.2514	0.0193	0.2312	0.0839	0.0016	0.0295	0.0432	0.0039	0.0162	0.1579	0.3979	0.9625	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0092	0.0209	0.8263	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0101	0.0002	0.2930	
MALE													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0049	0.0149	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0025	0.0059	
Smokeless Tobacco	0.0201	0.0014	0.0316	0.0886	0.2033	0.1649	0.3725	0.2620	0.0472	0.4853	0.7166	0.8554	
Cigars	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0091	0.0008	0.2462	
Pipe Tobacco	0.0003	0.0000	0.0521	0.2684	0.1037	0.0126	0.0426	0.5895	0.5202	0.9667	0.7873	0.0681	
ALCOHOL	0.0004	0.0000	0.0009	0.0000	0.0000	0.0002	0.0092	0.0000	0.0000	0.0971	0.1931	0.4838	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0107	0.0407	0.4702	
Heavy Alcohol Use ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0039	0.0001	0.1759	
FEMALE													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.1078	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0345	
Smokeless Tobacco	0.0538	0.5227	0.1727	0.7768	0.6068	0.8223	0.7314	0.0423	0.1646	0.5421	0.8223	0.0697	
Cigars	0.9102	0.6598	0.2486	0.5044	0.6043	0.7190	0.8639	0.7531	0.3738	0.2929	0.6512	0.5174	
Pipe Tobacco	0.0000	0.0000	0.0000	0.0004	0.0000	0.0000	0.0072	0.0058	0.9756	0.5553	0.1107	0.8462	
ALCOHOL	0.0570	0.0783	0.1108	0.0349	0.6591	0.7004	0.6759	0.9312	0.5093	0.6191	0.9595	0.5205	
Binge Alcohol Use ²	0.9150	0.9947	0.6163	0.9016	0.0158	0.0407	0.0232	0.0363	0.3895	0.1437	0.1518	0.6690	
Heavy Alcohol Use ²	0.0134	0.0030	0.0068	0.0057	0.0000	0.0000	0.0011	0.0536	0.0187	0.3737	0.2903	1.0000	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.32C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	797	748	752	798	799	801	849	784	813	805	841	803	679
Cigarettes	754	738	718	726	760	763	785	753	747	771	796	770	651
Smokeless Tobacco	270	274	236	256	267	268	303	274	302	266	286	288	234
Cigars	331	332	311	335	332	312	352	341	340	352	372	335	256
Pipe Tobacco	160	126	142	141	162	166	141	141	141	135	176	143	129
ALCOHOL	953	887	933	936	914	986	958	931	946	978	972	1,021	802
Binge Alcohol Use ²	702	639	654	650	697	714	703	751	715	699	749	766	623
Heavy Alcohol Use ²	355	349	356	347	392	402	391	377	398	396	420	420	334
MALE													
TOBACCO PRODUCTS¹	553	546	552	554	566	575	613	574	599	560	605	590	491
Cigarettes	506	521	512	493	527	551	546	538	514	514	545	530	453
Smokeless Tobacco	252	247	226	232	258	238	286	267	290	252	267	270	228
Cigars	311	302	280	309	305	290	325	318	306	321	334	293	242
Pipe Tobacco	132	124	135	136	156	164	128	136	134	129	165	140	125
ALCOHOL	630	577	610	598	583	640	616	631	645	665	652	669	524
Binge Alcohol Use ²	527	492	492	507	541	562	565	585	552	552	594	583	490
Heavy Alcohol Use ²	323	306	307	302	333	355	340	326	337	343	363	350	292
FEMALE													
TOBACCO PRODUCTS¹	488	477	469	515	477	486	543	485	516	520	525	511	414
Cigarettes	475	460	466	508	472	469	539	481	509	510	512	500	412
Smokeless Tobacco	99	125	63	85	74	111	73	49	72	80	102	91	47
Cigars	99	117	107	98	116	104	101	116	140	130	135	152	91
Pipe Tobacco	89	32	39	37	43	26	50	32	47	36	57	26	32
ALCOHOL	628	582	654	615	614	658	656	668	620	663	662	701	581
Binge Alcohol Use ²	393	347	361	370	356	364	382	422	427	406	406	435	338
Heavy Alcohol Use ²	158	155	160	146	147	157	181	181	190	184	192	219	151

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.32D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Percentages, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	0.44	0.41	0.41	0.43	0.43	0.42	0.44	0.40	0.42	0.41	0.42	0.40	0.33
Cigarettes	0.42	0.41	0.39	0.39	0.40	0.40	0.41	0.39	0.38	0.39	0.40	0.38	0.32
Smokeless Tobacco	0.15	0.15	0.13	0.14	0.14	0.14	0.16	0.14	0.15	0.13	0.14	0.14	0.11
Cigars	0.18	0.18	0.17	0.18	0.18	0.16	0.18	0.18	0.17	0.18	0.19	0.17	0.12
Pipe Tobacco	0.09	0.07	0.08	0.08	0.09	0.09	0.07	0.07	0.07	0.07	0.09	0.07	0.06
ALCOHOL	0.53	0.49	0.51	0.51	0.49	0.52	0.50	0.48	0.48	0.49	0.48	0.50	0.39
Binge Alcohol Use ²	0.39	0.35	0.36	0.35	0.37	0.38	0.37	0.39	0.37	0.35	0.37	0.38	0.30
Heavy Alcohol Use ²	0.20	0.19	0.19	0.19	0.21	0.21	0.20	0.19	0.20	0.20	0.21	0.21	0.16
MALE													
TOBACCO PRODUCTS¹	0.65	0.63	0.63	0.62	0.63	0.63	0.67	0.62	0.64	0.59	0.63	0.61	0.50
Cigarettes	0.59	0.60	0.59	0.56	0.59	0.61	0.59	0.58	0.55	0.54	0.57	0.55	0.46
Smokeless Tobacco	0.29	0.29	0.26	0.26	0.29	0.26	0.31	0.29	0.31	0.27	0.28	0.28	0.23
Cigars	0.36	0.35	0.32	0.35	0.34	0.32	0.35	0.34	0.33	0.34	0.35	0.30	0.25
Pipe Tobacco	0.15	0.14	0.15	0.15	0.17	0.18	0.14	0.15	0.14	0.14	0.17	0.14	0.13
ALCOHOL	0.74	0.67	0.70	0.67	0.65	0.70	0.67	0.68	0.69	0.70	0.68	0.69	0.53
Binge Alcohol Use ²	0.62	0.57	0.56	0.57	0.60	0.62	0.61	0.63	0.59	0.58	0.62	0.60	0.50
Heavy Alcohol Use ²	0.38	0.35	0.35	0.34	0.37	0.39	0.37	0.35	0.36	0.36	0.38	0.36	0.30
FEMALE													
TOBACCO PRODUCTS¹	0.52	0.50	0.49	0.53	0.49	0.49	0.54	0.48	0.51	0.50	0.50	0.48	0.39
Cigarettes	0.51	0.49	0.49	0.53	0.48	0.47	0.54	0.48	0.50	0.49	0.49	0.47	0.38
Smokeless Tobacco	0.11	0.13	0.07	0.09	0.08	0.11	0.07	0.05	0.07	0.08	0.10	0.09	0.04
Cigars	0.11	0.12	0.11	0.10	0.12	0.11	0.10	0.12	0.14	0.13	0.13	0.14	0.08
Pipe Tobacco	0.10	0.03	0.04	0.04	0.04	0.03	0.05	0.03	0.05	0.03	0.05	0.02	0.03
ALCOHOL	0.67	0.61	0.68	0.64	0.63	0.66	0.66	0.66	0.61	0.64	0.63	0.66	0.54
Binge Alcohol Use ²	0.42	0.37	0.38	0.38	0.36	0.37	0.38	0.42	0.42	0.39	0.39	0.41	0.32
Heavy Alcohol Use ²	0.17	0.16	0.17	0.15	0.15	0.16	0.18	0.18	0.19	0.18	0.18	0.21	0.14

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.32P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0027	0.0069	0.2936	0.0201	0.8507	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0034	0.0066	0.4385	0.0800	0.8733	
Smokeless Tobacco	0.2710	0.4559	0.0855	0.7378	0.5057	0.9535	0.1195	0.7326	0.5505	0.6911	0.1369	0.6310	
Cigars	0.0032	0.0127	0.0013	0.0006	0.0013	0.0227	0.0221	0.0332	0.0202	0.2403	0.0080	0.2945	
Pipe Tobacco	0.5300	0.6839	0.9524	0.1073	0.0307	0.4033	0.7298	0.8244	0.7978	0.8542	0.0876	0.8729	
ALCOHOL	0.0001	0.0000	0.0000	0.0242	0.0000	0.0003	0.0057	0.0089	0.0100	0.0253	0.1427	0.3269	
Binge Alcohol Use ²	0.0282	0.0008	0.0027	0.0013	0.0199	0.2417	0.4720	0.8695	0.2199	0.0474	0.3280	0.7311	
Heavy Alcohol Use ²	0.7571	0.7598	0.6701	0.1578	0.9854	0.6692	0.2583	0.3861	0.7286	0.2711	0.5823	0.7608	
MALE													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0913	0.0159	0.5170	0.0302	0.4805	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.1840	0.0257	0.7125	0.1197	0.9789	
Smokeless Tobacco	0.5152	0.9867	0.0541	0.4666	0.4764	0.8147	0.1421	0.6273	0.6440	0.4487	0.2072	0.8887	
Cigars	0.0002	0.0180	0.0002	0.0001	0.0034	0.0175	0.0075	0.0719	0.0772	0.3820	0.0341	0.8732	
Pipe Tobacco	0.6651	0.8989	0.7897	0.0957	0.0226	0.2422	0.5943	0.9818	0.7423	0.9935	0.1486	0.8701	
ALCOHOL	0.5004	0.2461	0.1688	0.7944	0.1329	0.0614	0.9452	0.3757	0.2857	0.2239	0.1404	0.7170	
Binge Alcohol Use ²	0.9704	0.4570	0.7391	0.1606	0.8674	0.3412	0.1626	0.4260	0.9758	0.3861	0.9903	0.9501	
Heavy Alcohol Use ²	0.2692	0.9097	0.4975	0.7206	0.3077	0.2289	0.1516	0.4420	0.7072	0.4353	0.6526	0.8500	
FEMALE													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0050	0.0003	0.0004	0.0028	0.0025	0.0045	0.1487	0.3313	0.2079	0.5993	
Cigarettes	0.0000	0.0000	0.0029	0.0002	0.0002	0.0016	0.0016	0.0018	0.0788	0.4124	0.2839	0.7813	
Smokeless Tobacco	0.0835	0.0439	0.4893	0.2221	0.9106	0.6722	0.5629	0.3746	0.6526	0.1750	0.3109	0.2080	
Cigars	0.1581	0.3124	0.6354	0.5109	0.1544	0.8897	0.4731	0.2305	0.0938	0.2798	0.0422	0.0317	
Pipe Tobacco	0.5719	0.2930	0.4806	0.9842	0.8763	0.1068	0.6535	0.2754	0.8000	0.4848	0.2441	0.1360	
ALCOHOL	0.0000	0.0000	0.0000	0.0004	0.0000	0.0004	0.0001	0.0045	0.0056	0.0435	0.4942	0.2897	
Binge Alcohol Use ²	0.0001	0.0000	0.0000	0.0002	0.0000	0.0002	0.0006	0.1377	0.0235	0.0246	0.0744	0.4855	
Heavy Alcohol Use ²	0.0056	0.4136	0.7567	0.0118	0.0274	0.1221	0.7602	0.7177	0.9729	0.3850	0.7022	0.7685	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.33C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	829	784	793	839	838	838	872	822	840	845	874	834	729
Cigarettes	790	774	765	772	799	803	809	790	772	814	832	796	691
Smokeless Tobacco	276	283	243	264	281	280	311	288	310	275	298	299	246
Cigars	346	333	323	351	348	331	363	358	358	369	387	354	280
Pipe Tobacco	163	130	145	145	165	169	146	149	147	143	183	150	137
ALCOHOL	972	909	950	969	954	1,011	977	945	974	1,006	1,004	1,052	843
Binge Alcohol Use ²	730	675	699	701	742	766	737	784	748	736	792	799	686
Heavy Alcohol Use ²	389	376	383	378	425	426	414	404	421	423	448	438	366
MALE													
TOBACCO PRODUCTS¹	565	561	568	576	581	594	630	597	616	580	613	615	514
Cigarettes	521	537	532	514	542	573	565	564	529	535	552	549	476
Smokeless Tobacco	260	255	233	239	271	249	295	281	299	261	280	279	238
Cigars	322	306	291	324	316	306	332	332	318	337	349	312	255
Pipe Tobacco	136	127	138	141	159	168	131	143	136	135	169	146	129
ALCOHOL	643	590	616	616	599	658	626	644	666	678	666	687	540
Binge Alcohol Use ²	540	517	511	529	561	585	588	606	573	568	621	606	521
Heavy Alcohol Use ²	346	320	318	322	352	373	358	349	352	355	382	367	311
FEMALE													
TOBACCO PRODUCTS¹	509	498	494	541	506	505	553	501	532	539	552	522	440
Cigarettes	497	481	488	537	497	487	548	494	524	529	538	511	433
Smokeless Tobacco	99	126	64	87	75	112	74	51	76	82	103	94	50
Cigars	110	121	119	109	128	112	113	127	150	140	145	161	109
Pipe Tobacco	90	35	41	40	45	28	54	35	53	45	60	33	41
ALCOHOL	640	593	662	635	641	673	667	679	634	677	682	717	611
Binge Alcohol Use ²	411	375	389	401	395	392	399	442	444	443	426	461	376
Heavy Alcohol Use ²	175	170	173	162	168	169	192	192	201	207	202	231	165

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.33D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Percentages, 2002-2014

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	0.39	0.37	0.37	0.39	0.38	0.38	0.39	0.36	0.37	0.36	0.37	0.35	0.30
Cigarettes	0.38	0.36	0.36	0.35	0.36	0.36	0.36	0.35	0.34	0.35	0.35	0.34	0.29
Smokeless Tobacco	0.13	0.13	0.11	0.12	0.13	0.13	0.14	0.13	0.14	0.12	0.13	0.13	0.10
Cigars	0.16	0.16	0.15	0.16	0.16	0.15	0.16	0.16	0.16	0.16	0.16	0.15	0.12
Pipe Tobacco	0.08	0.06	0.07	0.07	0.07	0.08	0.06	0.07	0.06	0.06	0.08	0.06	0.06
ALCOHOL	0.46	0.43	0.44	0.44	0.43	0.45	0.43	0.42	0.42	0.43	0.43	0.44	0.35
Binge Alcohol Use ²	0.35	0.32	0.32	0.32	0.34	0.34	0.33	0.34	0.33	0.32	0.34	0.34	0.29
Heavy Alcohol Use ²	0.19	0.18	0.18	0.17	0.19	0.19	0.18	0.18	0.18	0.18	0.19	0.18	0.15
MALE													
TOBACCO PRODUCTS¹	0.56	0.55	0.55	0.55	0.55	0.55	0.58	0.54	0.56	0.52	0.54	0.54	0.44
Cigarettes	0.52	0.53	0.51	0.49	0.51	0.53	0.52	0.51	0.48	0.48	0.49	0.48	0.41
Smokeless Tobacco	0.26	0.25	0.22	0.23	0.25	0.23	0.27	0.26	0.27	0.23	0.25	0.24	0.21
Cigars	0.32	0.30	0.28	0.31	0.30	0.29	0.31	0.30	0.29	0.30	0.31	0.27	0.22
Pipe Tobacco	0.13	0.12	0.13	0.13	0.15	0.16	0.12	0.13	0.12	0.12	0.15	0.13	0.11
ALCOHOL	0.64	0.58	0.59	0.59	0.56	0.61	0.58	0.59	0.60	0.61	0.59	0.60	0.47
Binge Alcohol Use ²	0.53	0.51	0.49	0.50	0.53	0.55	0.54	0.55	0.52	0.51	0.55	0.53	0.45
Heavy Alcohol Use ²	0.34	0.31	0.31	0.31	0.33	0.35	0.33	0.32	0.32	0.32	0.34	0.32	0.27
FEMALE													
TOBACCO PRODUCTS¹	0.47	0.45	0.44	0.48	0.44	0.44	0.47	0.43	0.45	0.45	0.45	0.42	0.35
Cigarettes	0.45	0.44	0.44	0.48	0.44	0.42	0.47	0.42	0.44	0.44	0.44	0.42	0.35
Smokeless Tobacco	0.09	0.11	0.06	0.08	0.07	0.10	0.06	0.04	0.06	0.07	0.08	0.08	0.04
Cigars	0.10	0.11	0.11	0.10	0.11	0.10	0.10	0.11	0.13	0.12	0.12	0.13	0.09
Pipe Tobacco	0.08	0.03	0.04	0.04	0.04	0.02	0.05	0.03	0.04	0.04	0.05	0.03	0.03
ALCOHOL	0.59	0.54	0.59	0.56	0.56	0.58	0.57	0.58	0.54	0.56	0.56	0.58	0.49
Binge Alcohol Use ²	0.38	0.34	0.35	0.36	0.35	0.34	0.34	0.38	0.37	0.37	0.35	0.37	0.30
Heavy Alcohol Use ²	0.16	0.15	0.16	0.14	0.15	0.15	0.16	0.16	0.17	0.17	0.17	0.19	0.13

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.33P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Gender/Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0165	0.0016	0.6390	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0151	0.0051	0.3546	
Smokeless Tobacco	0.6840	0.9531	0.0241	0.4643	0.7403	0.7138	0.1971	0.4133	0.1905	0.6071	0.1733	0.5218	
Cigars	0.0002	0.0001	0.0000	0.0000	0.0000	0.0001	0.0005	0.0006	0.0005	0.0433	0.0010	0.2215	
Pipe Tobacco	0.5181	0.0368	0.3050	0.4122	0.2325	0.8034	0.2416	0.6779	0.8918	0.8110	0.1323	0.7390	
ALCOHOL	0.0004	0.0000	0.0000	0.0724	0.0003	0.0022	0.0229	0.0543	0.0483	0.0639	0.2183	0.3529	
Binge Alcohol Use ²	0.3427	0.0991	0.1544	0.1608	0.5947	0.7354	0.5940	0.2026	0.9267	0.2889	0.7907	0.8183	
Heavy Alcohol Use ²	0.0235	0.0119	0.0015	0.0962	0.0036	0.0052	0.0008	0.0068	0.0374	0.8721	0.0968	0.5408	
MALE													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0029	0.0004	0.1339	0.0055	0.9215	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0039	0.0009	0.1237	0.0239	0.5064	
Smokeless Tobacco	0.9699	0.4316	0.0159	0.2566	0.7044	0.5556	0.2301	0.4112	0.2846	0.3699	0.2593	0.8396	
Cigars	0.0000	0.0002	0.0000	0.0000	0.0000	0.0000	0.0001	0.0015	0.0034	0.1018	0.0031	0.5997	
Pipe Tobacco	0.7168	0.2381	0.8593	0.1774	0.0621	0.5516	0.3069	0.8527	0.8557	0.9700	0.1389	0.4914	
ALCOHOL	0.9574	0.9890	0.5080	0.2195	0.5862	0.2780	0.6417	0.9858	0.8516	0.3950	0.2413	0.8328	
Binge Alcohol Use ²	0.1309	0.2569	0.2004	0.5490	0.0871	0.0224	0.0177	0.0339	0.2566	0.8585	0.6125	0.8071	
Heavy Alcohol Use ²	0.0005	0.0057	0.0008	0.0106	0.0006	0.0017	0.0007	0.0113	0.0519	0.9442	0.1183	0.5813	
FEMALE													
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0029	0.0267	0.0480	0.3803	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.0277	0.0433	0.4616	
Smokeless Tobacco	0.1514	0.0559	0.7043	0.2525	0.8264	0.6483	0.6273	0.8797	0.4103	0.1445	0.2980	0.1015	
Cigars	0.2223	0.2268	0.8468	0.8621	0.1243	0.9971	0.6059	0.2068	0.0550	0.1433	0.0905	0.0805	
Pipe Tobacco	0.5196	0.0009	0.0033	0.1005	0.0342	0.0002	0.4737	0.0222	0.8480	0.3913	0.6727	0.2049	
ALCOHOL	0.0000	0.0000	0.0000	0.0002	0.0000	0.0006	0.0002	0.0066	0.0055	0.0666	0.5142	0.2573	
Binge Alcohol Use ²	0.0003	0.0000	0.0000	0.0010	0.0007	0.0069	0.0129	0.4762	0.0702	0.1236	0.2385	0.4634	
Heavy Alcohol Use ²	0.1083	0.6606	0.4173	0.2400	0.9912	0.8809	0.3549	0.2841	0.4161	0.6820	0.4452	0.7684	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.34C Daily Cigarette Use in the Past Month among Past Month Users Aged 12 or Older and Smoked One or More Packs of Cigarettes Per Day among Daily Cigarette Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
DAILY CIGARETTE USE AMONG PAST MONTH USERS	774	714	693	664	690	727	730	726	737	709	740	707	578
12-17	41	39	43	34	35	35	29	30	30	28	25	21	27
18-25	135	144	146	135	134	135	129	130	120	134	129	115	127
26 or Older	738	678	649	632	672	696	705	696	709	681	709	691	545
SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS¹	538	525	505	489	482	509	512	503	486	441	455	466	365
12-17	19	17	17	15	15	16	12	12	11	10	7	7	8
18-25	79	81	76	77	74	71	67	64	60	62	54	49	57
26 or Older	528	508	491	476	473	500	501	495	479	435	447	458	355

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Daily Cigarette Use is defined as smoking on each of the past 30 days.

¹ Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.34D Daily Cigarette Use in the Past Month among Past Month Users Aged 12 or Older and Smoked One or More Packs of Cigarettes Per Day among Daily Cigarette Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
DAILY CIGARETTE USE AMONG PAST MONTH USERS	0.66	0.67	0.63	0.62	0.59	0.65	0.70	0.68	0.71	0.71	0.71	0.73	0.59
12-17	1.03	1.06	1.13	1.12	1.19	1.16	1.11	1.17	1.29	1.28	1.33	1.35	1.89
18-25	0.72	0.69	0.72	0.73	0.77	0.76	0.81	0.80	0.80	0.86	0.88	0.83	0.91
26 or Older	0.87	0.86	0.80	0.79	0.74	0.83	0.86	0.84	0.86	0.88	0.85	0.88	0.72
SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS¹	0.91	0.82	0.87	0.86	0.85	0.88	0.94	0.98	0.94	0.90	0.94	1.00	0.83
12-17	1.61	1.68	1.80	1.87	1.94	2.14	2.08	2.12	2.24	1.97	1.88	2.47	2.52
18-25	0.93	0.88	0.86	0.93	0.93	0.92	0.91	0.92	0.94	0.97	0.90	0.90	1.16
26 or Older	1.12	0.99	1.05	1.02	1.00	1.06	1.10	1.16	1.09	1.05	1.09	1.15	0.93

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Daily Cigarette Use is defined as smoking on each of the past 30 days.

¹ Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.34P Daily Cigarette Use in the Past Month among Past Month Users Aged 12 or Older and Smoked One or More Packs of Cigarettes Per Day among Daily Cigarette Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
DAILY CIGARETTE USE AMONG PAST MONTH USERS	0.0000	0.0000	0.0000	0.0000	0.0000	0.0042	0.0034	0.0130	0.4280	0.0379	0.0359	0.3802	
12-17	0.0004	0.0100	0.1135	0.4553	0.2832	0.3035	0.4212	0.6321	0.4772	0.5304	0.3624	0.0432	
18-25	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0575	0.0237	0.0676	0.1006	0.9722	
26 or Older	0.0000	0.0000	0.0000	0.0000	0.0000	0.0050	0.0010	0.0005	0.1692	0.0050	0.0153	0.1467	
SMOKED 1+ PACKS OF CIGARETTES PER DAY AMONG DAILY CIGARETTE USERS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0047	0.1802	0.4435	
12-17	0.0010	0.0009	0.0162	0.0087	0.0602	0.0398	0.0487	0.0699	0.1577	0.3679	0.7327	0.9950	
18-25	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0014	0.0168	0.0752	0.8538	
26 or Older	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0039	0.1940	0.3640	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Daily Cigarette Use is defined as smoking on each of the past 30 days.

¹ Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.35C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Gender: Standard Errors of Numbers in Thousands, 2002-2014

Gender/Alcohol Use	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
Lifetime	286	286	272	286	282	312	288	302	291	332	324	282	276
Past Year	250	253	246	268	259	293	264	273	266	307	308	255	261
Past Month	188	196	191	208	208	231	201	212	204	239	244	194	211
Binge Alcohol Use ¹	150	158	160	170	163	182	158	174	161	181	188	153	159
Heavy Alcohol Use ¹	76	79	98	86	93	93	85	91	84	95	88	68	73
MALE													
Lifetime	179	183	177	184	185	192	184	203	197	188	200	177	181
Past Year	158	166	162	171	170	179	165	187	180	173	187	162	169
Past Month	124	134	130	137	137	142	131	151	141	131	154	124	138
Binge Alcohol Use ¹	107	114	115	123	117	122	109	128	118	107	121	102	114
Heavy Alcohol Use ¹	59	59	70	67	73	72	66	76	66	59	66	53	56
FEMALE													
Lifetime	180	170	170	180	169	190	179	175	170	213	189	174	184
Past Year	160	159	155	171	156	181	169	157	156	199	179	159	172
Past Month	118	123	119	128	125	141	124	124	118	159	140	122	140
Binge Alcohol Use ¹	85	89	93	92	92	103	96	93	91	117	106	88	97
Heavy Alcohol Use ¹	41	43	50	44	46	49	47	39	45	60	51	40	38

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.35D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Gender: Standard Errors of Percentages, 2002-2014

Gender/Alcohol Use	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
Lifetime	0.40	0.43	0.41	0.41	0.42	0.45	0.43	0.45	0.44	0.49	0.48	0.47	0.48
Past Year	0.41	0.44	0.41	0.43	0.43	0.47	0.43	0.44	0.44	0.50	0.51	0.46	0.49
Past Month	0.39	0.41	0.39	0.41	0.42	0.46	0.40	0.43	0.41	0.47	0.48	0.40	0.46
Binge Alcohol Use ¹	0.34	0.36	0.36	0.37	0.36	0.39	0.34	0.38	0.35	0.39	0.40	0.34	0.37
Heavy Alcohol Use ¹	0.19	0.20	0.24	0.21	0.23	0.22	0.21	0.22	0.21	0.23	0.21	0.17	0.19
MALE													
Lifetime	0.55	0.53	0.56	0.55	0.54	0.59	0.55	0.57	0.60	0.61	0.61	0.58	0.66
Past Year	0.56	0.57	0.56	0.56	0.57	0.60	0.54	0.58	0.60	0.60	0.62	0.58	0.66
Past Month	0.52	0.53	0.55	0.53	0.54	0.57	0.53	0.58	0.55	0.54	0.61	0.51	0.60
Binge Alcohol Use ¹	0.48	0.49	0.52	0.51	0.49	0.51	0.47	0.53	0.50	0.47	0.52	0.45	0.53
Heavy Alcohol Use ¹	0.29	0.29	0.34	0.32	0.35	0.34	0.31	0.36	0.32	0.29	0.31	0.25	0.28
FEMALE													
Lifetime	0.51	0.59	0.55	0.56	0.58	0.61	0.58	0.60	0.59	0.67	0.63	0.62	0.65
Past Year	0.53	0.62	0.55	0.58	0.58	0.64	0.61	0.59	0.58	0.68	0.64	0.61	0.65
Past Month	0.50	0.57	0.51	0.54	0.56	0.59	0.53	0.55	0.53	0.65	0.59	0.54	0.61
Binge Alcohol Use ¹	0.43	0.44	0.44	0.43	0.45	0.48	0.45	0.45	0.44	0.53	0.49	0.42	0.47
Heavy Alcohol Use ¹	0.22	0.23	0.26	0.22	0.24	0.25	0.24	0.21	0.23	0.30	0.26	0.21	0.20

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.35P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Gender: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Gender/Alcohol Use	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL													
Lifetime	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0745	
Past Year	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.1733	
Past Month	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0214	0.8849	
Binge Alcohol Use ¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0069	0.4869	
Heavy Alcohol Use ¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0010	0.0016	0.2596	
MALE													
Lifetime	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0133	
Past Year	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0389	
Past Month	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0258	0.7981	
Binge Alcohol Use ¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0041	0.1319	0.5386	
Heavy Alcohol Use ¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0014	0.0238	0.3108	
FEMALE													
Lifetime	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0009	0.0569	0.8431	
Past Year	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0038	0.1767	0.9697	
Past Month	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0003	0.1447	0.0556	0.1846	0.6337	
Binge Alcohol Use ¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0047	0.0075	0.0052	0.6693	
Heavy Alcohol Use ¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0033	0.0598	0.0096	0.5385	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.36C Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	750	829	720	762	798	807	799	804	772	844	844	857	671
Marijuana													
Smoke Once a Month	912	897	897	927	930	941	941	964	918	982	928	993	737
Smoke Once or Twice a Week	881	860	905	873	883	921	937	1,003	978	954	984	1,050	820
Cocaine													
Use Once a Month	757	759	735	748	747	778	823	791	809	832	801	845	762
Use Once or Twice a Week	484	539	488	508	529	501	580	544	566	583	587	593	514
Heroin													
Try Once or Twice	563	595	619	608	602	628	657	629	675	667	605	692	548
Use Once or Twice a Week	363	378	385	384	386	369	412	386	448	425	407	431	357
LSD													
Try Once or Twice	696	732	765	753	735	792	783	808	848	853	839	867	754
Use Once or Twice a Week	474	497	483	545	488	526	536	587	614	623	590	637	547
Alcohol													
Have Four or Five Drinks Nearly Every Day	775	770	792	799	785	807	855	860	809	873	887	889	733
Have Five or More Drinks Once or Twice a Week	863	831	855	903	888	907	948	887	973	950	922	967	810
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	898	890	880	938	870	894	896	1,013	1,016	908	934	1,015	834
Cocaine	816	768	738	807	819	837	812	809	853	747	782	833	658
Crack	821	750	717	820	778	810	814	805	843	769	798	795	630
Heroin	669	651	614	665	641	697	693	691	667	622	670	703	575
LSD	661	612	604	655	607	654	668	622	640	582	612	636	514
Approached in the Past Month by Someone Selling Drugs	359	376	380	359	404	413	380	400	422	371	382	415	336

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.² Respondents with unknown Perceived Availability data were excluded from the respective analyses.³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.36D Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	0.32	0.35	0.30	0.32	0.33	0.33	0.32	0.32	0.31	0.33	0.33	0.33	0.25
Marijuana													
Smoke Once a Month	0.39	0.38	0.38	0.39	0.38	0.39	0.38	0.39	0.37	0.39	0.36	0.39	0.28
Smoke Once or Twice a Week	0.38	0.37	0.38	0.37	0.37	0.38	0.38	0.40	0.39	0.38	0.38	0.41	0.32
Cocaine													
Use Once a Month	0.33	0.32	0.31	0.31	0.31	0.32	0.33	0.32	0.32	0.33	0.31	0.33	0.29
Use Once or Twice a Week	0.21	0.23	0.21	0.21	0.22	0.21	0.24	0.22	0.23	0.23	0.23	0.23	0.20
Heroin													
Try Once or Twice	0.24	0.25	0.26	0.25	0.25	0.26	0.27	0.25	0.27	0.26	0.24	0.27	0.21
Use Once or Twice a Week	0.16	0.16	0.16	0.16	0.16	0.15	0.17	0.16	0.18	0.17	0.16	0.17	0.14
LSD													
Try Once or Twice	0.30	0.31	0.33	0.32	0.31	0.33	0.32	0.33	0.34	0.34	0.33	0.34	0.29
Use Once or Twice a Week	0.21	0.21	0.21	0.23	0.20	0.22	0.22	0.24	0.25	0.25	0.23	0.25	0.21
Alcohol													
Have Four or Five Drinks Nearly Every Day	0.33	0.33	0.33	0.33	0.32	0.33	0.34	0.34	0.32	0.34	0.34	0.34	0.28
Have Five or More Drinks Once or Twice a Week	0.37	0.35	0.36	0.37	0.36	0.37	0.38	0.36	0.39	0.37	0.36	0.37	0.31
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	0.40	0.39	0.38	0.40	0.37	0.38	0.37	0.42	0.42	0.37	0.38	0.41	0.33
Cocaine	0.37	0.34	0.33	0.35	0.35	0.36	0.35	0.34	0.36	0.31	0.32	0.34	0.26
Crack	0.37	0.33	0.32	0.36	0.34	0.35	0.35	0.34	0.35	0.32	0.33	0.32	0.25
Heroin	0.30	0.29	0.27	0.29	0.28	0.30	0.30	0.29	0.28	0.26	0.28	0.29	0.23
LSD	0.30	0.27	0.27	0.29	0.26	0.28	0.29	0.26	0.27	0.24	0.25	0.26	0.21
Approached in the Past Month by Someone Selling Drugs	0.15	0.16	0.16	0.15	0.16	0.17	0.15	0.16	0.17	0.14	0.15	0.16	0.13

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.36P Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	0.8112	0.6741	0.0000	0.0000	0.0000	0.0000	0.0000	0.5567	0.6711	0.9290	0.3821	0.4131	
Marijuana													
Smoke Once a Month	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	
Smoke Once or Twice a Week	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
Cocaine													
Use Once a Month	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0009	0.0001	0.0003	0.0695	0.0000	0.1640	
Use Once or Twice a Week	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0008	
Heroin													
Try Once or Twice	0.0373	0.0053	0.0000	0.0001	0.0043	0.0004	0.0002	0.0001	0.0001	0.0014	0.0872	0.0109	
Use Once or Twice a Week	0.0021	0.0059	0.0367	0.0098	0.0192	0.0042	0.1181	0.0705	0.5117	0.6811	0.4718	0.7837	
LSD													
Try Once or Twice	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.1181	
Use Once or Twice a Week	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	
Alcohol													
Have Four or Five Drinks Nearly Every Day	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0691	
Have Five or More Drinks Once or Twice a Week	0.0000	0.0019	0.2148	0.0119	0.0014	0.0000	0.0384	0.0001	0.0000	0.0001	0.0000	0.0327	
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	0.0000	0.0047	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0020	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0122	0.2750	
Crack	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0125	0.1915	
Heroin	0.0001	0.0000	0.0010	0.0362	0.0015	0.0598	0.5508	0.5335	0.0014	0.0083	0.0042	0.0000	
LSD	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0009	0.0075	0.0964	0.8577	
Approached in the Past Month by Someone Selling Drugs	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0056	0.0003	0.0000	0.0047	0.0024	0.3506	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.37C Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	108	104	104	110	106	108	101	103	107	107	106	109	113
Marijuana													
Smoke Once a Month	97	101	109	107	109	107	112	101	101	98	101	97	98
Smoke Once or Twice a Week	105	109	112	117	114	117	110	111	109	120	113	110	120
Cocaine													
Use Once a Month	108	112	106	114	107	110	108	110	107	115	109	108	115
Use Once or Twice a Week	86	88	92	95	90	95	95	93	91	92	94	97	95
Heroin													
Try Once or Twice	108	108	103	110	109	114	106	110	108	107	106	111	115
Use Once or Twice a Week	82	85	86	86	90	90	91	88	87	91	88	89	97
LSD													
Try Once or Twice	109	111	105	115	106	111	105	108	111	112	106	109	116
Use Once or Twice a Week	93	99	88	95	94	96	100	100	98	104	99	104	110
Alcohol													
Have Four or Five Drinks Nearly Every Day	118	112	112	108	109	108	108	107	104	114	103	106	121
Have Five or More Drinks Once or Twice a Week	106	103	101	118	108	112	111	101	109	120	111	109	116
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	109	106	108	110	110	109	110	111	119	116	109	111	118
Cocaine	86	90	102	99	97	97	94	89	85	80	88	79	83
Crack	90	91	100	96	99	94	96	94	89	81	82	80	82
Heroin	73	79	80	74	78	75	73	74	68	65	71	64	68
LSD	79	84	84	79	75	74	74	71	71	69	68	70	76
Approached in the Past Month by Someone Selling Drugs	82	83	85	86	76	79	79	79	81	80	77	79	87

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.37D Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	0.44	0.42	0.42	0.44	0.42	0.43	0.41	0.42	0.44	0.43	0.43	0.44	0.46
Marijuana													
Smoke Once a Month	0.40	0.41	0.44	0.43	0.44	0.43	0.46	0.42	0.42	0.40	0.41	0.40	0.40
Smoke Once or Twice a Week	0.43	0.44	0.45	0.47	0.45	0.47	0.45	0.46	0.46	0.49	0.46	0.45	0.49
Cocaine													
Use Once a Month	0.44	0.46	0.43	0.46	0.43	0.44	0.44	0.46	0.45	0.47	0.45	0.45	0.47
Use Once or Twice a Week	0.35	0.36	0.37	0.38	0.36	0.39	0.39	0.38	0.38	0.38	0.39	0.40	0.39
Heroin													
Try Once or Twice	0.45	0.44	0.42	0.44	0.44	0.46	0.44	0.46	0.46	0.44	0.44	0.46	0.48
Use Once or Twice a Week	0.34	0.35	0.35	0.35	0.36	0.36	0.38	0.37	0.37	0.37	0.36	0.37	0.40
LSD													
Try Once or Twice	0.46	0.46	0.44	0.48	0.44	0.47	0.45	0.46	0.49	0.47	0.45	0.47	0.50
Use Once or Twice a Week	0.39	0.41	0.36	0.39	0.39	0.41	0.42	0.43	0.43	0.44	0.42	0.45	0.47
Alcohol													
Have Four or Five Drinks Nearly Every Day	0.48	0.45	0.45	0.43	0.43	0.43	0.44	0.44	0.43	0.46	0.42	0.43	0.49
Have Five or More Drinks Once or Twice a Week	0.43	0.42	0.40	0.47	0.43	0.45	0.45	0.42	0.45	0.49	0.45	0.45	0.47
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	0.45	0.44	0.44	0.44	0.44	0.45	0.45	0.47	0.50	0.48	0.45	0.46	0.49
Cocaine	0.36	0.38	0.42	0.41	0.40	0.40	0.39	0.38	0.37	0.33	0.37	0.33	0.35
Crack	0.37	0.38	0.41	0.39	0.41	0.39	0.40	0.40	0.38	0.34	0.35	0.34	0.35
Heroin	0.31	0.33	0.33	0.31	0.32	0.31	0.31	0.32	0.30	0.28	0.30	0.28	0.29
LSD	0.34	0.36	0.35	0.33	0.32	0.31	0.32	0.31	0.32	0.29	0.29	0.30	0.33
Approached in the Past Month by Someone Selling Drugs	0.33	0.33	0.34	0.34	0.30	0.31	0.32	0.32	0.33	0.32	0.31	0.32	0.35

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.37P Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	0.0000	0.0007	0.0469	0.0015	0.0001	0.0001	0.0000	0.2258	0.1220	0.8528	0.3068	0.0014	
Marijuana													
Smoke Once a Month	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0179	
Smoke Once or Twice a Week	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0016	
Cocaine													
Use Once a Month	0.2621	0.0120	0.8453	0.1621	0.2206	0.7630	0.5561	0.3836	0.3478	0.0135	0.3543	0.4543	
Use Once or Twice a Week	0.0000	0.0000	0.0000	0.0000	0.0002	0.0021	0.0006	0.0254	0.0386	0.0909	0.0020	0.0279	
Heroin													
Try Once or Twice	0.2601	0.4843	0.0005	0.0000	0.0020	0.0006	0.0054	0.0001	0.0066	0.0000	0.0026	0.1391	
Use Once or Twice a Week	0.0000	0.0000	0.0050	0.0006	0.0204	0.0532	0.0146	0.0886	0.3653	0.6500	0.9388	0.7477	
LSD													
Try Once or Twice	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0053	0.0209	0.2309	0.1390	0.1810	
Use Once or Twice a Week	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0014	0.0005	0.0337	
Alcohol													
Have Four or Five Drinks Nearly Every Day	0.7582	0.2284	0.3673	0.0309	0.0011	0.0000	0.0000	0.0083	0.0004	0.0003	0.0207	0.9015	
Have Five or More Drinks Once or Twice a Week	0.1196	0.2368	0.0655	0.2223	0.9120	0.8948	0.2144	0.5452	0.0665	0.0278	0.4240	0.7098	
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	0.0000	0.0000	0.0000	0.0000	0.0004	0.0325	0.0161	0.0009	0.1157	0.9469	0.9910	0.2041	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0011	0.8839	
Crack	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.2148	
Heroin	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0011	0.2806	0.4668	
LSD	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0040	0.1854	0.7385	0.5077	
Approached in the Past Month by Someone Selling Drugs	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0000	0.0000	0.0005	0.0291	0.5288	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.38C Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	138	138	134	134	136	135	146	146	145	152	149	149	160
Marijuana													
Smoke Once a Month	130	124	137	133	127	143	129	135	128	134	127	122	122
Smoke Once or Twice a Week	148	150	150	156	152	154	145	149	146	155	145	143	137
Cocaine													
Use Once a Month	140	133	138	145	153	145	139	158	159	158	155	160	180
Use Once or Twice a Week	86	98	107	106	101	103	103	114	117	112	109	112	128
Heroin													
Try Once or Twice	111	118	117	120	116	123	118	134	133	126	123	129	146
Use Once or Twice a Week	73	74	75	77	74	72	77	81	82	82	80	80	87
LSD													
Try Once or Twice	140	136	140	145	148	152	154	153	161	158	163	167	188
Use Once or Twice a Week	92	97	101	105	104	109	110	117	124	125	130	133	155
Alcohol													
Have Four or Five Drinks Nearly Every Day	139	144	143	148	149	148	146	151	154	164	155	156	166
Have Five or More Drinks Once or Twice a Week	145	134	149	155	144	147	152	153	157	161	167	151	165
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	131	119	139	132	136	138	140	137	136	144	146	153	161
Cocaine	136	144	147	146	155	151	139	149	154	141	147	149	158
Crack	133	142	134	141	145	145	138	137	140	131	131	126	139
Heroin	110	109	112	115	122	122	115	113	116	112	115	113	120
LSD	117	126	113	116	121	113	115	116	114	112	123	122	131
Approached in the Past Month by Someone Selling Drugs	110	116	122	116	121	119	123	121	127	118	130	121	142

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.38D Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	0.45	0.44	0.42	0.41	0.42	0.41	0.44	0.44	0.43	0.45	0.43	0.43	0.46
Marijuana													
Smoke Once a Month	0.42	0.39	0.43	0.41	0.39	0.44	0.39	0.40	0.38	0.39	0.37	0.35	0.35
Smoke Once or Twice a Week	0.48	0.48	0.47	0.48	0.47	0.47	0.44	0.45	0.43	0.46	0.42	0.41	0.40
Cocaine													
Use Once a Month	0.45	0.42	0.43	0.45	0.47	0.45	0.43	0.47	0.47	0.47	0.45	0.47	0.52
Use Once or Twice a Week	0.28	0.31	0.33	0.33	0.31	0.32	0.31	0.34	0.35	0.33	0.32	0.33	0.37
Heroin													
Try Once or Twice	0.36	0.38	0.37	0.38	0.36	0.38	0.36	0.40	0.39	0.37	0.36	0.37	0.42
Use Once or Twice a Week	0.24	0.24	0.24	0.24	0.23	0.22	0.24	0.24	0.24	0.24	0.23	0.23	0.25
LSD													
Try Once or Twice	0.46	0.44	0.44	0.45	0.46	0.47	0.48	0.47	0.48	0.47	0.48	0.49	0.55
Use Once or Twice a Week	0.30	0.31	0.32	0.33	0.33	0.34	0.34	0.36	0.37	0.37	0.38	0.39	0.45
Alcohol													
Have Four or Five Drinks Nearly Every Day	0.45	0.45	0.45	0.46	0.46	0.45	0.44	0.45	0.46	0.48	0.45	0.45	0.48
Have Five or More Drinks Once or Twice a Week	0.47	0.42	0.46	0.48	0.44	0.45	0.46	0.46	0.46	0.47	0.48	0.44	0.48
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	0.43	0.38	0.44	0.41	0.42	0.43	0.43	0.41	0.41	0.43	0.43	0.45	0.47
Cocaine	0.45	0.47	0.47	0.46	0.49	0.48	0.44	0.46	0.47	0.42	0.44	0.44	0.47
Crack	0.44	0.47	0.43	0.45	0.46	0.46	0.44	0.42	0.43	0.40	0.39	0.38	0.41
Heroin	0.37	0.36	0.36	0.37	0.39	0.39	0.36	0.35	0.35	0.34	0.35	0.34	0.36
LSD	0.39	0.41	0.37	0.37	0.39	0.36	0.37	0.36	0.35	0.34	0.37	0.37	0.39
Approached in the Past Month by Someone Selling Drugs	0.35	0.37	0.38	0.36	0.37	0.36	0.37	0.36	0.37	0.34	0.38	0.35	0.41

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.38P Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 18 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	0.0092	0.0724	0.0000	0.0000	0.0000	0.0000	0.0019	0.0010	0.0376	0.1262	0.7355	0.1784	
Marijuana													
Smoke Once a Month	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0066	
Smoke Once or Twice a Week	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
Cocaine													
Use Once a Month	0.1037	0.3410	0.6421	0.4281	0.5922	0.6916	0.1009	0.0034	0.1294	0.1952	0.4110	0.9133	
Use Once or Twice a Week	0.0000	0.0000	0.0027	0.0027	0.0046	0.0032	0.0042	0.8222	0.8333	0.3680	0.0246	0.8420	
Heroin													
Try Once or Twice	0.0001	0.0000	0.0000	0.0000	0.0000	0.0003	0.0000	0.0000	0.0000	0.0001	0.0362	0.0877	
Use Once or Twice a Week	0.7661	0.9236	0.5431	0.5011	0.6776	0.9530	0.1030	0.0383	0.0127	0.0117	0.4642	0.0712	
LSD													
Try Once or Twice	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0003	0.0406	
Use Once or Twice a Week	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.6864	
Alcohol													
Have Four or Five Drinks Nearly Every Day	0.0011	0.1023	0.0071	0.0054	0.0007	0.0002	0.0000	0.0054	0.0021	0.0036	0.0370	0.7369	
Have Five or More Drinks Once or Twice a Week	0.5761	0.0116	0.0022	0.2048	0.1648	0.5866	0.3924	0.5676	0.8167	0.1353	0.1674	0.5966	
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	0.0001	0.0000	0.0702	0.0095	0.0084	0.0736	0.4909	0.0481	0.1978	0.8480	0.6281	0.5722	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.5734	
Crack	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.1821	
Heroin	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0002	0.5219	0.1904	0.9254	
LSD	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0056	0.0042	0.1469	0.4902	0.7395	
Approached in the Past Month by Someone Selling Drugs	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0000	0.0000	0.0000	0.0005	0.0000	0.8556	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.39C Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	732	795	694	733	774	787	770	778	758	836	812	827	634
Marijuana													
Smoke Once a Month	872	861	863	896	901	905	914	934	896	950	910	961	702
Smoke Once or Twice a Week	828	817	858	834	849	897	901	957	957	922	947	1,028	782
Cocaine													
Use Once a Month	726	721	702	708	718	752	802	758	778	788	774	816	707
Use Once or Twice a Week	477	507	474	475	510	482	557	525	539	563	556	570	480
Heroin													
Try Once or Twice	536	565	589	567	567	594	632	589	642	633	578	668	499
Use Once or Twice a Week	342	353	368	358	356	348	396	369	426	397	382	403	330
LSD													
Try Once or Twice	670	700	726	700	692	753	758	760	805	810	792	832	684
Use Once or Twice a Week	464	465	472	513	455	493	508	553	583	591	557	608	509
Alcohol													
Have Four or Five Drinks Nearly Every Day	764	753	773	757	759	776	831	839	784	841	842	859	702
Have Five or More Drinks Once or Twice a Week	834	818	834	868	854	876	916	858	938	915	889	939	764
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	873	861	855	908	837	857	854	968	982	873	891	975	793
Cocaine	792	736	718	776	780	809	796	777	819	734	748	804	627
Crack	793	733	703	791	744	778	787	774	813	746	769	780	604
Heroin	649	639	591	639	617	668	683	671	641	603	647	691	552
LSD	638	598	586	635	584	629	661	611	623	566	592	626	488
Approached in the Past Month by Someone Selling Drugs	330	328	341	306	355	374	330	343	383	329	316	370	276

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.39D Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	0.41	0.44	0.38	0.40	0.41	0.42	0.40	0.40	0.39	0.42	0.41	0.41	0.31
Marijuana													
Smoke Once a Month	0.50	0.48	0.48	0.49	0.49	0.49	0.48	0.49	0.47	0.49	0.46	0.48	0.35
Smoke Once or Twice a Week	0.47	0.46	0.48	0.46	0.46	0.48	0.48	0.50	0.50	0.47	0.48	0.52	0.39
Cocaine													
Use Once a Month	0.41	0.40	0.39	0.39	0.39	0.40	0.42	0.40	0.41	0.40	0.39	0.41	0.35
Use Once or Twice a Week	0.27	0.28	0.26	0.26	0.28	0.26	0.30	0.28	0.28	0.29	0.28	0.29	0.24
Heroin													
Try Once or Twice	0.30	0.32	0.33	0.31	0.31	0.32	0.33	0.31	0.33	0.33	0.29	0.34	0.25
Use Once or Twice a Week	0.19	0.20	0.20	0.20	0.19	0.19	0.21	0.19	0.22	0.20	0.19	0.20	0.16
LSD													
Try Once or Twice	0.38	0.39	0.40	0.39	0.38	0.41	0.40	0.40	0.42	0.42	0.40	0.42	0.34
Use Once or Twice a Week	0.26	0.26	0.26	0.28	0.25	0.27	0.27	0.29	0.31	0.31	0.28	0.31	0.25
Alcohol													
Have Four or Five Drinks Nearly Every Day	0.43	0.42	0.43	0.41	0.41	0.41	0.44	0.44	0.41	0.43	0.42	0.43	0.34
Have Five or More Drinks Once or Twice a Week	0.47	0.45	0.46	0.47	0.46	0.46	0.48	0.45	0.48	0.47	0.45	0.47	0.38
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	0.51	0.50	0.49	0.51	0.47	0.48	0.47	0.52	0.53	0.46	0.47	0.51	0.41
Cocaine	0.47	0.43	0.42	0.45	0.44	0.46	0.44	0.43	0.45	0.40	0.40	0.43	0.33
Crack	0.47	0.43	0.41	0.46	0.43	0.44	0.44	0.43	0.45	0.40	0.41	0.41	0.32
Heroin	0.39	0.38	0.35	0.37	0.35	0.38	0.38	0.37	0.35	0.33	0.35	0.37	0.29
LSD	0.38	0.35	0.34	0.37	0.33	0.36	0.37	0.34	0.34	0.31	0.32	0.33	0.26
Approached in the Past Month by Someone Selling Drugs	0.18	0.18	0.19	0.17	0.19	0.20	0.17	0.18	0.20	0.17	0.16	0.18	0.13

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.39P Perceived Risk of Harm Associated with Substance Use and Perceived Availability of Substances among Persons Aged 26 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Risk/Availability	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PERCEPTIONS OF GREAT RISK¹													
Cigarettes													
Smoke One or More Packs Per Day	0.1768	0.1213	0.0000	0.0000	0.0000	0.0000	0.0000	0.7046	0.2217	0.6109	0.2331	0.9522	
Marijuana													
Smoke Once a Month	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0021	
Smoke Once or Twice a Week	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	
Cocaine													
Use Once a Month	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0050	0.0000	0.0961	
Use Once or Twice a Week	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0000	0.0013	
Heroin													
Try Once or Twice	0.9323	0.5579	0.0057	0.3654	0.6077	0.1419	0.0937	0.1399	0.0521	0.1641	0.6064	0.0527	
Use Once or Twice a Week	0.0052	0.0132	0.0136	0.0123	0.0109	0.0016	0.0694	0.0223	0.2295	0.1642	0.2894	0.3953	
LSD													
Try Once or Twice	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0000	0.2747	
Use Once or Twice a Week	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0014	
Alcohol													
Have Four or Five Drinks Nearly Every Day	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0000	0.0491	
Have Five or More Drinks Once or Twice a Week	0.0000	0.0001	0.0239	0.0015	0.0002	0.0000	0.0284	0.0000	0.0000	0.0006	0.0000	0.0166	
PERCEIVED AVAILABILITY²													
Fairly or Very Easy to Obtain ³													
Marijuana	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0411	0.0036	0.1741	0.2111	
Crack	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0019	0.0064	0.2875	0.0865	
Heroin	0.5694	0.2429	0.4784	0.6411	0.6249	0.5386	0.2050	0.0282	0.0000	0.0016	0.0010	0.0000	
LSD	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0157	0.0225	0.1060	0.9809	
Approached in the Past Month by Someone Selling Drugs	0.1770	0.0248	0.0840	0.3018	0.0031	0.0168	0.8223	0.3372	0.1671	0.5681	0.5336	0.4586	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

¹ Response categories for the Perception of Risk questions include "No risk," "Slight risk," "Moderate risk," and "Great risk." The estimates in this table correspond to persons reporting "Great risk." Respondents with unknown Perception of Risk data were excluded from the respective analyses.

² Respondents with unknown Perceived Availability data were excluded from the respective analyses.

³ Response categories for the Perceived Availability questions pertaining to the listed illicit drugs include "Probably impossible," "Very difficult," "Fairly difficult," "Fairly easy," and "Very easy." The estimates in this table correspond to persons reporting "Fairly easy" or "Very easy."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.40C Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Youths Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Behavior	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Smoking One or More Packs of Cigarettes Per Day	64	68	70	64	61	61	57	55	58	54	54	55	60
Trying Marijuana or Hashish Once or Twice	70	69	69	64	64	64	64	66	71	68	67	68	83
Using Marijuana or Hashish Once a Month or More	59	55	58	59	55	56	56	56	64	59	61	64	77
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	66	71	63	69	69	67	69	64	68	61	68	66	70

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Response categories for the questions on Youths' Perceptions of Parents' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove." Respondents with unknown Perceptions of Parents' Feelings data were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.40D Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Youths Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Behavior	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Smoking One or More Packs of Cigarettes Per Day	0.26	0.28	0.28	0.26	0.24	0.25	0.23	0.23	0.24	0.22	0.22	0.23	0.25
Trying Marijuana or Hashish Once or Twice	0.29	0.28	0.28	0.26	0.26	0.26	0.26	0.27	0.29	0.28	0.27	0.28	0.34
Using Marijuana or Hashish Once a Month or More	0.24	0.22	0.23	0.23	0.22	0.22	0.23	0.23	0.26	0.24	0.25	0.26	0.31
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	0.27	0.29	0.25	0.27	0.28	0.27	0.28	0.26	0.28	0.25	0.27	0.27	0.28

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Response categories for the questions on Youths' Perceptions of Parents' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove." Respondents with unknown Perceptions of Parents' Feelings data were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.40P Youths Felt That Parents Would Strongly Disapprove of Youth's Substance Use Behaviors among Youths Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Behavior	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Smoking One or More Packs of Cigarettes Per Day	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0005	0.0008	0.0875	0.0527	0.4371	
Trying Marijuana or Hashish Once or Twice	0.0002	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0461	
Using Marijuana or Hashish Once a Month or More	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0977	
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	0.0001	0.0000	0.0001	0.0000	0.0202	0.0151	0.0315	0.4272	0.8364	0.8200	0.8457	0.6800	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Response categories for the questions on Youths' Perceptions of Parents' Feelings include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove." Respondents with unknown Perceptions of Parents' Feelings data were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.41C Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Youths Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Behavior	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Smoking One or More Packs of Cigarettes Per Day	73	68	71	73	70	68	68	64	68	64	63	65	61
Trying Marijuana or Hashish Once or Twice	87	86	87	93	87	85	85	86	92	89	88	92	102
Using Marijuana or Hashish Once a Month or More	86	87	84	90	82	86	84	87	88	90	86	92	103
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	81	76	74	76	77	78	77	76	73	77	72	74	76

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Response categories for the questions on Youths' Feelings about Peers include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Feelings about Peers data were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.41D Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Youths Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Behavior	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Smoking One or More Packs of Cigarettes Per Day	0.30	0.27	0.29	0.29	0.28	0.27	0.28	0.26	0.28	0.26	0.26	0.27	0.25
Trying Marijuana or Hashish Once or Twice	0.35	0.35	0.35	0.37	0.35	0.34	0.35	0.35	0.38	0.36	0.36	0.38	0.41
Using Marijuana or Hashish Once a Month or More	0.35	0.35	0.34	0.36	0.33	0.34	0.34	0.36	0.37	0.37	0.35	0.37	0.42
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	0.33	0.31	0.30	0.31	0.31	0.31	0.31	0.31	0.30	0.31	0.29	0.30	0.31

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Response categories for the questions on Youths' Feelings about Peers include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Feelings about Peers data were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.41P Youths Strongly Disapproved or Somewhat Disapproved of Peers' Substance Use Behaviors among Youths Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Behavior	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Smoking One or More Packs of Cigarettes Per Day	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0010	0.0021	
Trying Marijuana or Hashish Once or Twice	0.9726	0.3743	0.0483	0.0173	0.0001	0.0000	0.0000	0.0001	0.0019	0.3249	0.0602	0.9704	
Using Marijuana or Hashish Once a Month or More	0.0271	0.0091	0.0000	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0376	0.0337	0.9141	
Having One or Two Drinks of an Alcoholic Beverage Nearly Every Day	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0004	0.0202	0.0275	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Response categories for the questions on Youths' Feelings about Peers include "Neither approve nor disapprove," "Somewhat disapprove," and "Strongly disapprove." The estimates in this table correspond to persons reporting "Strongly disapprove" or "Somewhat disapprove." Respondents with unknown Feelings about Peers data were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.42C Exposure to Substance Use Prevention Program or Message in the Past Year among Youths Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Program/Message	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School	70	79	76	72	73	74	72	71	66	72	71	71	74
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School	89	81	87	91	90	99	88	98	96	93	98	104	109
Drug or Alcohol Prevention Messages Were Seen or Heard in School among Youths Enrolled in School ^{1,2}	248	249	239	249	226	251	248	255	251	258	263	256	235

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents with unknown Substance Use Program Participation or Prevention Message Exposure data were excluded.

¹ Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

² Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.42D Exposure to Substance Use Prevention Program or Message in the Past Year among Youths Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Program/Message	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School	0.28	0.32	0.30	0.29	0.29	0.30	0.29	0.29	0.28	0.29	0.29	0.29	0.30
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School	0.36	0.33	0.35	0.36	0.36	0.40	0.36	0.40	0.40	0.38	0.40	0.42	0.44
Drug or Alcohol Prevention Messages Were Seen or Heard in School among Youths Enrolled in School ^{1,2}	0.38	0.37	0.40	0.37	0.38	0.41	0.41	0.43	0.45	0.43	0.44	0.44	0.47

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents with unknown Substance Use Program Participation or Prevention Message Exposure data were excluded.

¹ Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

² Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.42P Exposure to Substance Use Prevention Program or Message in the Past Year among Youths Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Program/Message	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School	0.0001	0.0000	0.0075	0.1648	0.4306	0.5594	0.9922	0.0226	0.3213	0.1406	0.0493	0.2917	
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.6126	
Drug or Alcohol Prevention Messages Were Seen or Heard in School among Youths Enrolled in School ^{1,2}	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0000	0.0094	0.0014	0.3838	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents with unknown Substance Use Program Participation or Prevention Message Exposure data were excluded.

¹ Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

² Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.43C Substance Use in the Past Month among Youths Aged 12 to 17, by Exposure to Substance Use Prevention Program or Message in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014

Program/Message	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School													
Illicit Drugs ¹	25	26	27	20	21	24	21	23	22	24	26	21	22
Marijuana	20	20	21	18	17	20	16	19	19	20	22	19	20
Cigarettes	28	27	26	24	22	21	20	20	20	18	21	18	16
Binge Alcohol ²	19	24	24	22	26	22	20	20	19	20	22	15	17
Did Not Participate in Alcohol, Tobacco, or Drug Prevention Program Outside of School													
Illicit Drugs ¹	69	70	70	66	66	68	63	68	71	67	65	62	74
Marijuana	57	60	58	55	52	54	55	59	62	60	56	55	65
Cigarettes	74	73	76	70	66	68	59	64	61	59	52	49	49
Binge Alcohol ²	68	68	75	65	66	70	58	63	62	55	57	56	60
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School													
Illicit Drugs ¹	66	64	67	62	61	63	58	60	64	61	59	55	64
Marijuana	55	58	54	54	47	49	51	52	54	54	50	48	59
Cigarettes	71	67	71	65	56	63	54	59	54	50	48	41	43
Binge Alcohol ²	64	65	71	63	61	63	53	58	55	51	50	47	53
Did Not See or Hear Alcohol or Drug Prevention Message from Sources Outside School													
Illicit Drugs ¹	31	33	32	31	35	37	33	38	36	36	35	35	39
Marijuana	25	30	26	25	30	30	25	32	32	31	31	30	35
Cigarettes	33	35	33	34	38	33	31	32	34	35	29	30	28
Binge Alcohol ²	31	32	31	29	34	33	32	31	30	26	32	31	30
Saw or Heard Drug or Alcohol Prevention Messages in School among Youths Enrolled in School ^{3,4}													
Illicit Drugs ¹	58	60	64	57	57	55	53	57	59	57	58	53	61
Marijuana	49	51	52	51	45	44	46	50	49	49	47	47	55
Cigarettes	61	58	68	61	56	55	49	53	51	48	43	40	38
Binge Alcohol ²	58	59	65	58	58	57	51	54	50	51	47	45	48
Did Not See or Hear Drug or Alcohol Prevention Messages in School among Youths Enrolled in School ^{3,4}													
Illicit Drugs ¹	35	37	35	36	36	40	37	37	38	43	38	35	43
Marijuana	28	34	33	29	31	34	32	32	33	39	33	32	38
Cigarettes	38	39	44	36	37	40	35	36	35	36	32	29	33
Binge Alcohol ²	32	34	40	33	37	41	35	34	36	28	33	33	34

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents with unknown Substance Use Prevention Program or Message Exposure data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

⁴ Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.43D Substance Use in the Past Month among Youths Aged 12 to 17, by Exposure to Substance Use Prevention Program or Message in the Past Year: Standard Errors of Percentages, 2002-2014

Program/Message	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School													
Illicit Drugs ¹	0.76	0.71	0.81	0.66	0.70	0.80	0.73	0.74	0.76	0.79	0.81	0.71	0.79
Marijuana	0.60	0.57	0.65	0.58	0.59	0.67	0.57	0.62	0.65	0.66	0.70	0.63	0.72
Cigarettes	0.79	0.70	0.76	0.79	0.73	0.70	0.70	0.67	0.68	0.62	0.69	0.62	0.57
Binge Alcohol ²	0.58	0.64	0.72	0.72	0.84	0.72	0.70	0.67	0.66	0.66	0.70	0.53	0.61
Did Not Participate in Alcohol, Tobacco, or Drug Prevention Program Outside of School													
Illicit Drugs ¹	0.31	0.30	0.29	0.27	0.28	0.29	0.26	0.29	0.31	0.29	0.27	0.26	0.32
Marijuana	0.26	0.26	0.25	0.23	0.22	0.24	0.24	0.25	0.27	0.26	0.24	0.24	0.28
Cigarettes	0.32	0.32	0.32	0.30	0.28	0.28	0.26	0.28	0.27	0.26	0.23	0.21	0.22
Binge Alcohol ²	0.30	0.29	0.32	0.27	0.28	0.29	0.25	0.27	0.27	0.24	0.24	0.24	0.26
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School													
Illicit Drugs ¹	0.31	0.27	0.30	0.28	0.29	0.30	0.28	0.30	0.32	0.31	0.29	0.29	0.35
Marijuana	0.26	0.26	0.24	0.25	0.23	0.24	0.25	0.26	0.27	0.28	0.25	0.26	0.32
Cigarettes	0.32	0.30	0.31	0.30	0.27	0.30	0.27	0.30	0.28	0.26	0.24	0.23	0.23
Binge Alcohol ²	0.28	0.29	0.32	0.29	0.29	0.30	0.26	0.29	0.28	0.26	0.25	0.25	0.29
Did Not See or Hear Alcohol or Drug Prevention Message from Sources Outside School													
Illicit Drugs ¹	0.71	0.71	0.66	0.60	0.63	0.62	0.57	0.61	0.59	0.53	0.54	0.50	0.55
Marijuana	0.58	0.69	0.58	0.51	0.55	0.52	0.44	0.54	0.53	0.48	0.49	0.44	0.51
Cigarettes	0.71	0.76	0.73	0.68	0.67	0.56	0.54	0.54	0.55	0.53	0.46	0.42	0.42
Binge Alcohol ²	0.70	0.71	0.70	0.59	0.62	0.56	0.56	0.53	0.49	0.41	0.50	0.44	0.44
Saw or Heard Drug or Alcohol Prevention Messages in School among Youths Enrolled in School ^{3,4}													
Illicit Drugs ¹	0.31	0.31	0.33	0.29	0.30	0.30	0.28	0.31	0.32	0.31	0.30	0.29	0.34
Marijuana	0.27	0.28	0.27	0.26	0.25	0.24	0.25	0.28	0.27	0.28	0.26	0.26	0.31
Cigarettes	0.32	0.32	0.34	0.32	0.29	0.30	0.27	0.30	0.29	0.27	0.24	0.23	0.22
Binge Alcohol ²	0.30	0.31	0.33	0.31	0.30	0.31	0.27	0.29	0.28	0.28	0.26	0.25	0.27
Did Not See or Hear Drug or Alcohol Prevention Messages in School among Youths Enrolled in School ^{3,4}													
Illicit Drugs ¹	0.69	0.68	0.65	0.65	0.61	0.65	0.65	0.58	0.64	0.64	0.59	0.53	0.64
Marijuana	0.56	0.63	0.62	0.55	0.55	0.57	0.56	0.52	0.57	0.60	0.53	0.50	0.58
Cigarettes	0.76	0.73	0.79	0.65	0.64	0.66	0.61	0.59	0.61	0.56	0.51	0.45	0.51
Binge Alcohol ²	0.64	0.62	0.73	0.60	0.64	0.68	0.60	0.56	0.62	0.47	0.52	0.50	0.52

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents with unknown Substance Use Prevention Program or Message Exposure data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

⁴ Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.43P Substance Use in the Past Month among Youths Aged 12 to 17, by Exposure to Substance Use Prevention Program or Message in the Past Year: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Program/Message	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School													
Illicit Drugs ¹	0.1240	0.1530	0.1419	0.5026	0.6292	0.9707	0.5769	0.3474	0.9097	0.1892	0.8990	0.6793	
Marijuana	0.9240	0.7619	0.3753	0.1845	0.3056	0.3813	0.1314	0.8253	0.8578	0.6616	0.7573	0.9234	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0003	0.0000	0.0002	0.0207	0.0065	0.0456	
Binge Alcohol ²	0.0001	0.0000	0.0000	0.0009	0.0000	0.0073	0.0200	0.0035	0.0306	0.0931	0.0072	0.8814	
Did Not Participate in Alcohol, Tobacco, or Drug Prevention Program Outside of School													
Illicit Drugs ¹	0.0000	0.0000	0.0055	0.0842	0.1563	0.5832	0.8863	0.0905	0.0742	0.1349	0.6098	0.1763	
Marijuana	0.0104	0.0617	0.6951	0.2338	0.0950	0.0946	0.1055	0.8313	0.8937	0.1652	0.8314	0.4148	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0405	
Binge Alcohol ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0045	0.5544	
Saw or Heard Alcohol or Drug Prevention Message from Sources Outside School													
Illicit Drugs ¹	0.0000	0.0002	0.0065	0.1378	0.7175	0.8328	0.8181	0.1518	0.0271	0.0510	0.5590	0.1313	
Marijuana	0.0272	0.3903	0.4792	0.3702	0.0255	0.0668	0.2855	0.9804	0.5629	0.0773	0.9325	0.5192	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0308	
Binge Alcohol ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0040	0.7985	
Did Not See or Hear Alcohol or Drug Prevention Message from Sources Outside School													
Illicit Drugs ¹	0.0003	0.0000	0.0315	0.5496	0.0137	0.3816	0.7873	0.1089	1.0000	0.7275	0.9567	0.9568	
Marijuana	0.2104	0.0098	0.5866	0.2904	0.3411	0.6522	0.0588	0.5398	0.5174	0.8584	0.7104	0.8370	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0030	0.0984	
Binge Alcohol ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0224	0.3061	0.0197	0.4704	
Saw or Heard Drug or Alcohol Prevention Messages in School among Youths Enrolled in School ^{3,4}													
Illicit Drugs ¹	0.0000	0.0000	0.0008	0.1218	0.3244	0.5208	0.9736	0.1081	0.0638	0.0920	0.3841	0.8307	
Marijuana	0.0009	0.0729	0.2928	0.6603	0.1123	0.1484	0.1887	0.7714	0.5731	0.1734	0.8306	0.9132	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	
Binge Alcohol ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.3536	
Did Not See or Hear Drug or Alcohol Prevention Messages in School among Youths Enrolled in School ^{3,4}													
Illicit Drugs ¹	0.0167	0.0071	0.1473	0.3143	0.2600	0.7890	0.9882	0.4914	0.7013	0.3261	0.9600	0.0118	
Marijuana	0.8123	0.1573	0.3844	0.2877	0.9222	0.3570	0.3951	0.9727	0.6916	0.2726	0.8130	0.1058	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0089	0.8044	
Binge Alcohol ²	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.5502	0.3271	0.9433	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents with unknown Substance Use Prevention Program or Message Exposure data were excluded from the respective analyses.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ Youths who did not report their school enrollment status or reported not being enrolled in school in the past 12 months were excluded from this analysis. Youths reporting that they were "home-schooled" in the past 12 months were considered to be enrolled in school.

⁴ Drug or Alcohol Prevention Messages Seen or Heard in School includes respondents who reported having a special class about drugs or alcohol in school or having had films, lectures, discussion, or printed information about drugs or alcohol in a regular school class, such as health or physical education, or outside of a regular school class, such as in a special assembly.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.44C Past Year Initiation of Substance Use among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	90	99	100	139	102	94	107	119	107	115	106	102	107
Marijuana and Hashish	70	70	81	121	79	77	89	79	93	97	96	86	95
Cocaine	61	56	65	50	60	57	52	47	57	48	48	47	57
Crack	44	36	29	30	31	72	34	15	20	14	16	13	24
Heroin	20	20	28	20	15	21	23	30	24	26	23	36	35
Hallucinogens	54	46	55	51	72	60	59	66	61	58	58	72	58
LSD	30	20	25	29	32	23	31	28	39	30	41	40	48
PCP	15	14	20	13	13	11	10	9	11	10	21	7	10
Ecstasy	68	44	47	46	68	53	58	68	56	59	60	50	52
Inhalants	43	44	43	47	41	47	44	48	51	47	42	51	49
Nonmedical Use of Psychotherapeutics ^{2,3}	108	110	120	111	110	117	105	121	113	118	124	100	99
Pain Relievers	107	109	100	102	93	105	94	100	91	102	99	82	76
OxyContin [®]	--	--	54	63	50	50	36	55	53	49	39	75	31
Tranquilizers	80	64	74	88	68	82	73	95	86	90	115	89	73
Stimulants ²	49	57	70	48	69	50	48	56	52	59	62	48	58
Methamphetamine ²	31	35	35	21	42	20	16	26	16	20	27	21	31
Sedatives	33	38	43	40	46	29	32	29	46	29	29	27	25
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	97	103	100	112	102	107	104	119	109	109	104	108	103
CIGARETTES	75	72	72	86	79	71	90	89	81	86	89	81	90
Daily Cigarette Use ⁴	64	58	55	58	54	52	57	66	57	55	53	52	51
SMOKELESS TOBACCO⁵	52	50	53	58	71	61	83	67	72	75	53	65	59
CIGARS	103	99	112	113	104	107	105	121	120	143	108	144	104
ALCOHOL	101	104	127	108	107	111	116	112	131	124	130	113	127

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.44D Past Year Initiation of Substance Use among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	0.04	0.04	0.04	0.06	0.04	0.04	0.04	0.05	0.04	0.04	0.04	0.04	0.04
Marijuana and Hashish	0.03	0.03	0.03	0.05	0.03	0.03	0.04	0.03	0.04	0.04	0.04	0.03	0.04
Cocaine	0.03	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Crack	0.02	0.02	0.01	0.01	0.01	0.03	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Heroin	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Hallucinogens	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.03	0.02	0.02	0.02	0.03	0.02
LSD	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.02	0.02	0.02
PCP	0.01	0.01	0.01	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00
Ecstasy	0.03	0.02	0.02	0.02	0.03	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.02
Inhalants	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Nonmedical Use of Psychotherapeutics ^{2,3}	0.05	0.05	0.05	0.05	0.04	0.05	0.04	0.05	0.04	0.05	0.05	0.04	0.04
Pain Relievers	0.05	0.05	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.03	0.03
OxyContin®	--	--	0.02	0.03	0.02	0.02	0.01	0.02	0.02	0.02	0.01	0.03	0.01
Tranquilizers	0.03	0.03	0.03	0.04	0.03	0.03	0.03	0.04	0.03	0.03	0.04	0.03	0.03
Stimulants ²	0.02	0.02	0.03	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Methamphetamine ²	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Sedatives	0.01	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.04	0.04	0.04	0.05	0.04	0.04	0.04	0.05	0.04	0.04	0.04	0.04	0.04
CIGARETTES	0.03	0.03	0.03	0.04	0.03	0.03	0.04	0.04	0.03	0.03	0.03	0.03	0.03
Daily Cigarette Use ⁴	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.02
SMOKELESS TOBACCO⁵	0.02	0.02	0.02	0.02	0.03	0.02	0.03	0.03	0.03	0.03	0.02	0.02	0.02
CIGARS	0.04	0.04	0.05	0.05	0.04	0.04	0.04	0.05	0.05	0.06	0.04	0.06	0.04
ALCOHOL	0.04	0.04	0.05	0.04	0.04	0.04	0.05	0.04	0.05	0.05	0.05	0.04	0.05

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.44P Past Year Initiation of Substance Use among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	0.9147	0.6025	0.7070	0.3898	0.9526	0.2995	0.6410	0.0770	0.4922	0.3095	0.6395	0.3699	
Marijuana and Hashish	0.4555	0.0028	0.1124	0.1061	0.0066	0.0082	0.1203	0.6163	0.8929	0.3622	0.3672	0.3679	
Cocaine	0.0000	0.0001	0.0003	0.0194	0.0009	0.0146	0.9769	0.1464	0.2494	0.3074	0.1275	0.0317	
Crack	0.0000	0.0000	0.0013	0.0005	0.0002	0.0009	0.0097	0.7432	0.4715	0.2546	0.4069	0.0605	
Heroin	0.0568	0.0091	0.0804	0.0242	0.0030	0.0195	0.0388	0.7508	0.1447	0.5207	0.2149	0.4241	
Hallucinogens	0.0000	0.4984	0.2663	0.1991	0.0055	0.0221	0.0019	0.0000	0.0000	0.0049	0.0547	0.1303	
LSD	0.0005	0.0000	0.0000	0.0000	0.0000	0.0000	0.0055	0.0001	0.0030	0.0001	0.0142	0.1129	
PCP	0.0000	0.0001	0.0016	0.0123	0.0423	0.1757	0.2882	0.6349	0.6310	0.5782	0.0311	0.4850	
Ecstasy	0.0000	0.5763	0.9176	0.9297	0.0049	0.0459	0.0008	0.0000	0.0001	0.0007	0.0093	0.2557	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0000	0.0010	0.1955	0.4202	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0002	0.0058	0.0526	0.0161	0.5508	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.2508	
OxyContin [®]	--	--	0.0000	0.0003	0.0000	0.0000	0.0000	0.0000	0.0000	0.0010	0.1157	0.0868	
Tranquilizers	0.0811	0.5451	0.1267	0.0265	0.4895	0.1062	0.5055	0.1822	0.1488	0.3681	0.0203	0.6086	
Stimulants ²	0.0162	0.2129	0.0571	0.8428	0.0188	0.9450	0.5032	0.4865	0.6510	0.9885	0.9861	0.2824	
Methamphetamine ²	0.0010	0.0351	0.0008	0.5037	0.0843	0.6804	0.0229	0.6201	0.0422	0.2138	0.2477	0.3114	
Sedatives	0.1654	0.3782	0.0872	0.0570	0.0393	0.3322	0.6224	0.5602	0.0922	0.8120	0.9239	0.2476	
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.0014	0.0095	0.0006	0.0002	0.0006	0.0310	0.0022	0.0006	0.0153	0.1621	0.2414	0.2588	
CIGARETTES	0.8505	0.6892	0.1438	0.0126	0.0001	0.0592	0.0009	0.0001	0.0051	0.0179	0.0890	0.5617	
Daily Cigarette Use ⁴	0.0000	0.0000	0.0000	0.0003	0.0000	0.0001	0.0020	0.0000	0.0016	0.0531	0.6241	0.3732	
SMOKELESS TOBACCO⁵	0.4997	0.8102	0.3012	0.0110	0.0000	0.0000	0.0000	0.0000	0.0000	0.0010	0.9941	0.2074	
CIGARS	0.0001	0.0028	0.0000	0.0000	0.0000	0.0000	0.0011	0.0000	0.0029	0.1145	0.4322	0.2601	
ALCOHOL	0.2158	0.5527	0.3132	0.9827	0.7140	0.2209	0.6326	0.3968	0.2152	0.3149	0.8988	0.7716	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.45C Past Year Initiation of Substance Use among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	52	51	51	50	52	50	54	53	53	52	52	50	58
Marijuana and Hashish	50	45	45	44	48	45	49	49	51	53	49	46	52
Cocaine	24	22	23	23	22	22	20	17	18	16	16	13	20
Crack	13	11	9	7	8	10	5	5	4	5	8	4	5
Heroin	10	7	8	5	7	5	10	5	7	10	7	6	7
Hallucinogens	27	27	27	26	27	28	30	30	33	28	26	23	25
LSD	18	13	13	14	11	13	16	12	15	16	15	14	22
PCP	11	10	9	11	10	8	7	7	6	7	11	6	6
Ecstasy	25	19	20	18	18	23	23	25	27	24	20	18	18
Inhalants	34	36	32	36	33	30	31	32	33	30	25	22	28
Nonmedical Use of Psychotherapeutics ^{2,3}	42	39	42	37	40	39	42	41	39	35	35	34	40
Pain Relievers	42	41	42	36	37	36	40	39	39	32	32	31	34
OxyContin [®]	--	--	15	14	15	15	14	16	19	15	13	13	15
Tranquilizers	25	24	22	21	20	23	22	20	21	25	26	19	25
Stimulants ²	24	21	21	21	25	20	21	18	21	19	18	16	23
Methamphetamine ²	16	14	13	12	13	12	11	10	9	14	7	8	13
Sedatives	11	11	10	10	9	12	11	12	12	10	8	7	12
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	44	48	47	44	44	44	49	46	49	41	43	39	46
CIGARETTES	44	47	50	50	48	48	50	50	47	46	43	41	44
Daily Cigarette Use ⁴	27	27	32	24	27	23	23	24	24	22	22	19	19
SMOKELESS TOBACCO⁵	27	27	29	34	31	36	34	33	33	31	27	34	30
CIGARS	40	46	48	47	42	44	43	43	40	41	38	36	41
ALCOHOL	64	65	73	69	68	69	64	69	62	69	72	67	67

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.45D Past Year Initiation of Substance Use among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	0.21	0.20	0.20	0.20	0.20	0.20	0.22	0.21	0.22	0.21	0.21	0.20	0.23
Marijuana and Hashish	0.20	0.18	0.18	0.17	0.19	0.18	0.20	0.20	0.21	0.21	0.20	0.18	0.21
Cocaine	0.10	0.09	0.09	0.09	0.08	0.09	0.08	0.07	0.07	0.07	0.06	0.05	0.08
Crack	0.05	0.04	0.03	0.03	0.03	0.04	0.02	0.02	0.02	0.02	0.03	0.01	0.02
Heroin	0.04	0.03	0.03	0.02	0.03	0.02	0.04	0.02	0.03	0.04	0.03	0.02	0.03
Hallucinogens	0.11	0.11	0.11	0.10	0.11	0.11	0.12	0.12	0.14	0.11	0.10	0.09	0.10
LSD	0.07	0.05	0.05	0.05	0.04	0.05	0.07	0.05	0.06	0.07	0.06	0.06	0.09
PCP	0.04	0.04	0.04	0.04	0.04	0.03	0.03	0.03	0.03	0.03	0.04	0.02	0.02
Ecstasy	0.10	0.08	0.08	0.07	0.07	0.09	0.09	0.10	0.11	0.10	0.08	0.07	0.07
Inhalants	0.14	0.14	0.13	0.14	0.13	0.12	0.12	0.13	0.14	0.12	0.10	0.09	0.11
Nonmedical Use of Psychotherapeutics ^{2,3}	0.17	0.16	0.17	0.15	0.16	0.15	0.17	0.17	0.16	0.14	0.14	0.13	0.16
Pain Relievers	0.17	0.16	0.17	0.14	0.14	0.14	0.16	0.16	0.16	0.13	0.13	0.12	0.14
OxyContin [®]	--	--	0.06	0.06	0.06	0.06	0.06	0.07	0.08	0.06	0.05	0.05	0.06
Tranquilizers	0.10	0.10	0.09	0.08	0.08	0.09	0.09	0.08	0.09	0.10	0.10	0.08	0.10
Stimulants ²	0.10	0.09	0.09	0.08	0.10	0.08	0.08	0.07	0.08	0.07	0.07	0.07	0.09
Methamphetamine ²	0.06	0.05	0.05	0.05	0.05	0.05	0.04	0.04	0.04	0.06	0.03	0.03	0.05
Sedatives	0.04	0.04	0.04	0.04	0.04	0.05	0.05	0.05	0.05	0.04	0.03	0.03	0.05
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.18	0.19	0.19	0.18	0.17	0.17	0.20	0.19	0.20	0.16	0.17	0.16	0.19
CIGARETTES	0.18	0.19	0.20	0.20	0.19	0.19	0.20	0.20	0.19	0.18	0.17	0.16	0.18
Daily Cigarette Use ⁴	0.11	0.11	0.13	0.10	0.11	0.09	0.09	0.10	0.10	0.09	0.09	0.08	0.08
SMOKELESS TOBACCO⁵	0.11	0.11	0.11	0.13	0.12	0.14	0.14	0.13	0.14	0.13	0.11	0.14	0.12
CIGARS	0.16	0.18	0.19	0.19	0.17	0.18	0.17	0.17	0.16	0.17	0.15	0.14	0.17
ALCOHOL	0.26	0.26	0.29	0.27	0.27	0.27	0.26	0.28	0.25	0.28	0.29	0.27	0.27

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.45P Past Year Initiation of Substance Use among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLICIT DRUGS^{1,2}	0.0127	0.0611	0.4553	0.4088	0.7208	0.5735	0.2492	0.0053	0.0527	0.0372	0.8632	0.4479	
Marijuana and Hashish	0.0147	0.8874	0.6438	0.2026	0.6274	0.4400	0.5425	0.0313	0.1796	0.0256	0.4915	0.9544	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0051	0.2517	0.1157	0.2581	0.9002	0.3438	
Crack	0.0000	0.0000	0.0030	0.0143	0.0021	0.0002	0.4070	0.3108	0.6838	0.3161	0.4915	0.8464	
Heroin	0.0312	0.2352	0.1050	0.5843	0.2729	0.7520	0.1978	0.4875	0.3034	0.0416	0.4654	0.4007	
Hallucinogens	0.0000	0.0000	0.0000	0.0003	0.0003	0.0004	0.0000	0.0000	0.0000	0.0000	0.0058	0.1579	
LSD	0.5921	0.0057	0.0082	0.0149	0.0002	0.0061	0.4940	0.0204	0.0189	0.1187	0.1258	0.0970	
PCP	0.0000	0.0003	0.0190	0.0033	0.0281	0.0358	0.0423	0.3084	0.5771	0.2032	0.0240	0.8243	
Ecstasy	0.0000	0.0004	0.0003	0.0299	0.0022	0.0001	0.0000	0.0000	0.0000	0.0000	0.0043	0.0236	
Inhalants	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.1138	0.6196	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0000	0.0000	0.0000	0.0004	0.0000	0.0003	0.0000	0.0002	0.0052	0.0418	0.1159	0.4359	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0109	0.6845	
OxyContin [®]	--	--	0.0309	0.1481	0.0628	0.0084	0.0031	0.0040	0.0050	0.0176	0.1697	0.8350	
Tranquilizers	0.0775	0.0626	0.1180	0.6533	0.5628	0.7888	0.3817	0.3598	0.7307	0.2207	0.4676	0.2661	
Stimulants ²	0.0000	0.0006	0.0075	0.0151	0.0015	0.2101	0.6793	0.6039	0.7085	0.6266	0.9922	0.1150	
Methamphetamine ²	0.0001	0.0031	0.0119	0.0733	0.0781	0.3045	0.6935	0.9329	0.7145	0.2002	0.1649	0.4614	
Sedatives	0.7933	0.6674	0.6695	0.9754	0.8566	0.6703	0.5138	0.3019	0.6638	0.7290	0.2834	0.0157	
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0000	0.0000	0.0000	0.0015	0.1211	0.6605	
CIGARETTES	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0019	0.1219	
Daily Cigarette Use ⁴	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0002	0.0000	0.0000	0.0005	0.2652	0.1008	
SMOKELESS TOBACCO⁵	0.4577	0.1848	0.8173	0.0212	0.0004	0.0001	0.0001	0.0000	0.0008	0.0842	0.6738	0.2490	
CIGARS	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0048	0.0040	0.3666	0.2174	
ALCOHOL	0.0043	0.0088	0.0002	0.0002	0.0009	0.0007	0.0130	0.0003	0.0354	0.0043	0.2746	0.3988	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.46C Past Year Initiation of Substance Use among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	41	39	41	47	53	47	53	58	51	60	59	57	63
Marijuana and Hashish	37	37	45	45	46	45	49	55	51	61	57	54	62
Cocaine	42	36	41	35	40	38	33	32	32	38	37	37	40
Crack	15	15	17	21	18	15	15	11	8	9	11	6	14
Heroin	13	9	10	13	12	14	11	13	15	17	16	13	15
Hallucinogens	35	33	32	32	40	35	39	46	45	42	42	44	47
LSD	18	14	16	16	22	18	23	25	33	23	33	31	37
PCP	11	9	14	6	8	7	6	6	9	8	8	5	8
Ecstasy	37	28	25	28	41	31	37	42	40	37	41	39	42
Inhalants	24	25	23	24	21	23	24	29	27	32	26	24	22
Nonmedical Use of Psychotherapeutics ^{2,3}	42	47	46	43	48	50	50	52	48	50	53	53	55
Pain Relievers	45	46	43	43	46	50	46	48	43	44	43	48	45
OxyContin [®]	--	--	26	23	23	21	22	23	22	22	21	23	20
Tranquilizers	33	32	31	35	32	34	31	37	40	33	39	36	36
Stimulants ²	30	22	30	29	30	30	28	31	28	29	39	31	38
Methamphetamine ²	20	14	19	14	17	14	10	16	11	10	22	16	14
Sedatives	10	8	11	14	11	10	14	14	14	13	10	9	13
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	43	42	45	44	55	51	49	55	54	54	52	57	60
CIGARETTES	40	45	46	46	52	48	58	60	54	59	65	57	72
Daily Cigarette Use ⁴	31	35	36	33	36	38	35	39	44	37	39	36	40
SMOKELESS TOBACCO⁵	31	27	28	34	40	40	40	43	40	43	36	40	44
CIGARS	46	48	54	58	54	58	54	61	66	58	61	61	67
ALCOHOL	51	64	62	61	68	70	68	66	79	80	77	76	86

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.46D Past Year Initiation of Substance Use among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	0.13	0.12	0.13	0.14	0.16	0.14	0.16	0.17	0.15	0.18	0.17	0.16	0.18
Marijuana and Hashish	0.12	0.12	0.14	0.14	0.14	0.14	0.15	0.16	0.15	0.18	0.17	0.16	0.18
Cocaine	0.14	0.12	0.13	0.11	0.12	0.12	0.10	0.10	0.09	0.11	0.11	0.11	0.11
Crack	0.05	0.05	0.05	0.06	0.05	0.05	0.05	0.03	0.02	0.03	0.03	0.02	0.04
Heroin	0.04	0.03	0.03	0.04	0.04	0.04	0.03	0.04	0.04	0.05	0.05	0.04	0.04
Hallucinogens	0.11	0.10	0.10	0.10	0.12	0.11	0.12	0.14	0.13	0.12	0.12	0.13	0.14
LSD	0.06	0.05	0.05	0.05	0.07	0.06	0.07	0.07	0.10	0.07	0.10	0.09	0.10
PCP	0.03	0.03	0.04	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.01	0.02
Ecstasy	0.12	0.09	0.08	0.08	0.12	0.10	0.11	0.13	0.12	0.11	0.12	0.11	0.12
Inhalants	0.08	0.08	0.07	0.08	0.06	0.07	0.07	0.09	0.08	0.09	0.07	0.07	0.06
Nonmedical Use of Psychotherapeutics ^{2,3}	0.14	0.15	0.14	0.13	0.15	0.15	0.15	0.16	0.14	0.15	0.15	0.15	0.16
Pain Relievers	0.14	0.14	0.13	0.13	0.14	0.15	0.14	0.14	0.13	0.13	0.12	0.14	0.13
OxyContin [®]	--	--	0.08	0.07	0.07	0.07	0.07	0.07	0.07	0.06	0.06	0.07	0.06
Tranquilizers	0.10	0.10	0.10	0.11	0.10	0.10	0.09	0.11	0.12	0.10	0.11	0.10	0.10
Stimulants ²	0.10	0.07	0.09	0.09	0.09	0.09	0.09	0.09	0.08	0.08	0.11	0.09	0.11
Methamphetamine ²	0.06	0.04	0.06	0.04	0.05	0.04	0.03	0.05	0.03	0.03	0.06	0.05	0.04
Sedatives	0.03	0.03	0.03	0.04	0.03	0.03	0.04	0.04	0.04	0.04	0.03	0.03	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.14	0.13	0.14	0.14	0.17	0.16	0.15	0.16	0.16	0.16	0.15	0.16	0.17
CIGARETTES	0.13	0.14	0.14	0.14	0.16	0.15	0.18	0.18	0.16	0.17	0.19	0.17	0.21
Daily Cigarette Use ⁴	0.10	0.11	0.11	0.10	0.11	0.11	0.11	0.12	0.13	0.11	0.11	0.10	0.12
SMOKELESS TOBACCO⁵	0.10	0.08	0.09	0.11	0.12	0.12	0.12	0.13	0.12	0.12	0.10	0.12	0.13
CIGARS	0.15	0.15	0.17	0.18	0.17	0.18	0.16	0.18	0.19	0.17	0.18	0.18	0.19
ALCOHOL	0.16	0.20	0.19	0.19	0.21	0.21	0.21	0.20	0.23	0.23	0.22	0.22	0.25

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.46P Past Year Initiation of Substance Use among Persons Aged 18 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLICIT DRUGS^{1,2}	0.0023	0.0000	0.0016	0.0030	0.0478	0.0130	0.2994	0.9658	0.1749	0.7586	0.9730	0.8066	
Marijuana and Hashish	0.0003	0.0000	0.0001	0.0001	0.0001	0.0012	0.0051	0.4278	0.0580	0.8695	0.1631	0.3737	
Cocaine	0.0073	0.0195	0.0188	0.5312	0.0685	0.1866	0.3452	0.0902	0.0206	0.6409	0.3274	0.2130	
Crack	0.0063	0.0024	0.0012	0.0002	0.0003	0.0591	0.0502	0.5694	0.3761	0.4244	0.8165	0.0598	
Heroin	0.9969	0.1177	0.1931	0.5223	0.4569	1.0000	0.5174	0.5677	0.6400	0.2278	0.3317	0.6850	
Hallucinogens	0.4208	0.0188	0.0061	0.0945	0.9797	0.5478	0.9334	0.0991	0.1301	0.4613	0.7185	0.9327	
LSD	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0062	0.0031	0.0378	0.0009	0.0367	0.2396	
PCP	0.0502	0.1064	0.0801	0.9909	0.6772	0.7508	0.5219	0.5432	0.9700	0.6317	0.6791	0.2419	
Ecstasy	0.0000	0.1382	0.0173	0.0456	0.1823	0.8536	0.0632	0.0020	0.0630	0.2648	0.0866	0.3917	
Inhalants	0.0005	0.0010	0.0101	0.0033	0.1359	0.0805	0.0220	0.0201	0.0359	0.0880	0.0655	0.3363	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0191	0.0202	0.0573	0.7107	0.3073	0.1412	0.0843	0.0503	0.7905	0.4869	0.8513	0.6601	
Pain Relievers	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0040	0.0275	0.0214	0.3189	
OxyContin [®]	--	--	0.0000	0.0050	0.0018	0.0029	0.0008	0.0000	0.0004	0.0697	0.7100	0.8458	
Tranquilizers	0.6219	0.2202	0.7171	0.3368	0.0752	0.4614	0.3601	0.1512	0.1485	0.8504	0.2782	0.7805	
Stimulants ²	0.6524	0.0426	0.7282	0.2304	0.6155	0.2221	0.2431	0.8584	0.6693	0.4690	0.7131	0.8423	
Methamphetamine ²	0.0019	0.1010	0.0008	0.1927	0.0842	0.6580	0.1091	0.3110	0.4644	0.1488	0.4666	0.2239	
Sedatives	0.8105	0.2488	0.6929	0.7111	0.9662	0.5616	0.8436	0.9597	0.3889	0.4431	0.1635	0.0672	
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.6825	0.0679	0.2795	0.3487	0.4721	0.7202	0.9906	0.2472	0.9943	0.8495	0.6892	0.2924	
CIGARETTES	0.0000	0.0000	0.0001	0.0021	0.4375	0.1551	0.6753	0.8954	0.7189	0.9677	0.7237	0.1155	
Daily Cigarette Use ⁴	0.6502	0.4389	0.0166	0.3368	0.0440	0.0276	0.0621	0.0041	0.0263	0.3135	0.7972	0.5995	
SMOKELESS TOBACCO⁵	0.7067	0.1364	0.9912	0.1878	0.0259	0.0062	0.0139	0.0004	0.0010	0.0225	0.5550	0.4192	
CIGARS	0.0784	0.0809	0.9067	0.1865	0.5812	0.0808	0.6284	0.0796	0.2405	0.5755	0.9400	0.7502	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0013	0.0002	0.0006	0.1577	0.0666	0.0251	0.1639	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.47C Past Year Initiation of Substance Use among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	53	69	77	115	68	61	69	85	77	73	63	63	64
Marijuana and Hashish	25	30	47	97	33	37	45	15	60	49	53	44	45
Cocaine	33	36	39	23	38	32	34	28	41	22	26	29	35
Crack	38	31	21	19	24	69	30	10	18	9	8	11	18
Heroin	11	16	25	15	6	14	17	27	17	17	15	32	31
Hallucinogens	28	17	31	30	49	38	32	34	18	28	27	49	19
LSD	13	*	15	19	19	*	12	*	15	8	16	25	19
PCP	*	*	10	*	*	*	*	*	*	*	16	*	*
Ecstasy	50	27	29	33	48	35	36	43	24	40	37	26	23
Inhalants	8	13	13	15	12	26	11	24	27	17	18	39	34
Nonmedical Use of Psychotherapeutics ^{2,3}	84	91	100	96	91	99	81	100	91	92	103	78	69
Pain Relievers	82	92	79	84	73	85	71	77	69	84	77	61	53
OxyContin [®]	--	--	45	56	40	44	25	47	44	41	29	68	17
Tranquilizers	67	52	62	78	56	71	60	86	73	76	102	80	59
Stimulants ²	27	48	58	30	58	35	32	41	39	46	44	31	35
Methamphetamine ²	16	29	26	11	36	9	8	17	8	10	15	11	22
Sedatives	30	35	39	37	44	24	26	22	42	24	26	25	18
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	70	79	75	89	75	79	75	93	77	79	78	78	67
CIGARETTES	36	31	20	40	30	15	31	35	24	25	28	32	29
Daily Cigarette Use ⁴	48	39	29	42	33	23	37	49	23	32	27	31	25
SMOKELESS TOBACCO⁵	27	25	35	26	50	23	64	37	48	48	27	31	26
CIGARS	82	74	79	86	77	73	77	90	87	119	72	126	62
ALCOHOL	40	21	74	31	22	32	50	34	76	32	56	26	37

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.47D Past Year Initiation of Substance Use among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	0.03	0.04	0.04	0.06	0.04	0.03	0.04	0.04	0.04	0.04	0.03	0.03	0.03
Marijuana and Hashish	0.01	0.02	0.03	0.05	0.02	0.02	0.02	0.01	0.03	0.02	0.03	0.02	0.02
Cocaine	0.02	0.02	0.02	0.01	0.02	0.02	0.02	0.01	0.02	0.01	0.01	0.01	0.02
Crack	0.02	0.02	0.01	0.01	0.01	0.04	0.02	0.01	0.01	0.00	0.00	0.01	0.01
Heroin	0.01	0.01	0.01	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02
Hallucinogens	0.02	0.01	0.02	0.02	0.03	0.02	0.02	0.02	0.01	0.01	0.01	0.02	0.01
LSD	0.01	*	0.01	0.01	0.01	*	0.01	*	0.01	0.00	0.01	0.01	0.01
PCP	*	*	0.01	*	*	*	*	*	*	*	0.01	*	*
Ecstasy	0.03	0.01	0.02	0.02	0.03	0.02	0.02	0.02	0.01	0.02	0.02	0.01	0.01
Inhalants	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02
Nonmedical Use of Psychotherapeutics ^{2,3}	0.05	0.05	0.05	0.05	0.05	0.05	0.04	0.05	0.05	0.05	0.05	0.04	0.03
Pain Relievers	0.05	0.05	0.04	0.05	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.03	0.03
OxyContin [®]	--	--	0.02	0.03	0.02	0.02	0.01	0.02	0.02	0.02	0.01	0.03	0.01
Tranquilizers	0.04	0.03	0.03	0.04	0.03	0.04	0.03	0.04	0.04	0.04	0.05	0.04	0.03
Stimulants ²	0.01	0.03	0.03	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Methamphetamine ²	0.01	0.02	0.01	0.01	0.02	0.00	0.00	0.01	0.00	0.01	0.01	0.01	0.01
Sedatives	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.04	0.04	0.04	0.05	0.04	0.04	0.04	0.05	0.04	0.04	0.04	0.04	0.03
CIGARETTES	0.02	0.02	0.01	0.02	0.02	0.01	0.02	0.02	0.01	0.01	0.01	0.02	0.01
Daily Cigarette Use ⁴	0.03	0.02	0.02	0.02	0.02	0.01	0.02	0.03	0.01	0.02	0.01	0.02	0.01
SMOKELESS TOBACCO⁵	0.01	0.01	0.02	0.01	0.03	0.01	0.03	0.02	0.02	0.02	0.01	0.02	0.01
CIGARS	0.05	0.04	0.04	0.05	0.04	0.04	0.04	0.05	0.04	0.06	0.04	0.06	0.03
ALCOHOL	0.02	0.01	0.04	0.02	0.01	0.02	0.03	0.02	0.04	0.02	0.03	0.01	0.02

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.47P Past Year Initiation of Substance Use among Persons Aged 26 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	0.0660	0.3148	0.5313	0.2292	0.8677	0.2027	0.8298	0.9227	0.9149	0.3752	0.1793	0.4062	
Marijuana and Hashish	0.0019	0.0027	0.2957	0.9399	0.0222	0.0382	0.1265	0.0000	0.8925	0.2249	0.2042	0.3621	
Cocaine	0.9616	0.9545	0.9906	0.2387	0.8070	0.5769	0.4363	0.1793	0.6097	0.0305	0.1113	0.1165	
Crack	0.0070	0.2016	0.6061	0.5497	0.3096	0.0165	0.0872	0.1739	0.6404	0.1899	0.1756	0.3166	
Heroin	0.0013	0.0097	0.0633	0.0145	0.0004	0.0039	0.0101	0.4270	0.0190	0.0233	0.0181	0.3772	
Hallucinogens	0.3218	0.4560	0.5184	0.6215	0.0875	0.1269	0.5931	0.1375	0.5381	0.8283	0.5094	0.1310	
LSD	0.1913	*	0.3645	0.4205	0.4812	*	0.1772	*	0.2338	0.0808	0.5094	0.9504	
PCP	*	*	*	*	*	*	*	*	*	*	*	*	
Ecstasy	0.0043	0.8782	0.6896	0.6647	0.2077	0.4856	0.5201	0.0393	0.6068	0.6561	0.5907	0.4234	
Inhalants	0.0389	0.1166	0.1610	0.1936	0.0953	0.9706	0.0939	0.5057	0.6053	0.2224	0.2210	0.4543	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.2511	0.0795	0.0010	0.0088	0.0310	0.0715	0.2029	0.1230	0.0659	0.3415	0.0338	0.8443	
Pain Relievers	0.0253	0.0023	0.0001	0.0037	0.0102	0.0275	0.0321	0.0190	0.0307	0.0912	0.0267	0.6533	
OxyContin [®]	--	--	0.0094	0.0325	0.0083	0.0064	0.2293	0.0466	0.0035	0.0598	0.2795	0.0702	
Tranquilizers	0.3871	0.2143	0.4876	0.0879	0.7881	0.2155	0.8371	0.6691	0.3701	0.6681	0.0549	0.4474	
Stimulants ²	0.5116	0.3887	0.5153	0.5036	0.1299	0.9948	0.7030	0.7005	0.4881	0.4942	0.8255	0.5338	
Methamphetamine ²	0.3928	0.9308	0.5054	0.0963	0.7361	0.0479	0.0389	0.1225	0.0216	0.0517	0.0690	0.0291	
Sedatives	0.1456	0.2710	0.0739	0.0604	0.0344	0.2416	0.7633	0.9836	0.1786	0.7790	0.2950	0.5503	
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.7728	0.9477	0.2401	0.0778	0.7085	0.7176	0.5677	0.4506	0.8464	0.6976	0.4354	0.5034	
CIGARETTES	0.7346	0.4674	0.0430	0.9864	0.2172	0.0035	0.2607	0.7875	0.1024	0.0738	0.3115	0.4287	
Daily Cigarette Use ⁴	0.2035	0.2566	0.6464	0.4617	0.8829	0.5234	0.7544	0.0730	0.3626	0.5599	0.6116	0.7594	
SMOKELESS TOBACCO⁵	0.8449	0.6575	0.6921	0.5203	0.2248	0.4036	0.1117	0.3471	0.0940	0.1516	0.4340	0.9516	
CIGARS	0.0036	0.3482	0.0673	0.0031	0.2035	0.2725	0.5144	0.0892	0.1373	0.3677	0.6289	0.1123	
ALCOHOL	0.4297	0.5418	0.2983	0.6791	0.5170	0.5981	0.0893	0.4723	0.2285	0.7594	0.1209	0.8555	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.48C Past Year Initiation of Illicit Drugs and First Drug Initiated in the Past Year among Persons Aged 12 or Older Who Initiated Illicit Drug Use in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014

	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PAST YEAR INITIATION OF ILLICIT DRUGS^{1,2}	91	100	99	139	102	94	109	118	109	116	106	102	107
FIRST DRUG INITIATED³													
Marijuana and Hashish	58	60	67	116	67	64	75	69	80	88	83	77	82
Cocaine	10	8	31	7	11	6	9	4	2	3	3	2	10
Crack	*	*	*	1	*	*	*	*	*	*	*	*	*
Heroin	1	1	1	*	0	*	3	2	2	2	2	*	*
Hallucinogens	14	13	11	13	23	11	21	16	16	17	12	19	14
LSD	1	4	1	2	4	2	2	1	1	2	1	3	6
PCP	2	0	1	1	5	*	1	0	2	3	5	1	*
Ecstasy	11	10	8	6	21	6	7	10	11	14	6	9	10
Inhalants	19	30	23	28	22	24	23	25	23	21	20	19	24
Nonmedical Use of Psychotherapeutics ^{2,4}	60	68	67	73	67	64	68	90	70	64	57	60	60
Pain Relievers	50	60	55	54	55	44	60	50	43	53	45	42	39
OxyContin [®]	--	--	1	5	1	2	3	9	8	18	4	*	5
Tranquilizers	14	19	27	43	27	37	25	71	29	26	23	37	30
Stimulants ²	15	21	22	13	21	25	18	11	33	17	23	17	29
Methamphetamine ²	9	4	16	4	6	8	9	4	2	2	*	2	5
Sedatives	25	15	10	22	16	11	8	15	29	16	16	4	11
Illicit Drugs Other Than Marijuana ^{1,2}	67	79	76	80	74	70	76	95	75	70	61	66	68

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ First Drug Initiated is based on imputation-revised month and year and imputed day of first use for each drug. Respondents may be counted in more than one drug category if they reported initiating multiple drugs on the same day the first time they used drugs. Also, several drugs listed are subcategories of other drugs, and will have the same start date. Therefore, the sum of the estimates for the first drug or drugs initiated may not equal the total estimates for past year initiates.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.48D Past Year Initiation of Illicit Drugs and First Drug Initiated in the Past Year among Persons Aged 12 or Older Who Initiated Illicit Drug Use in the Past Year: Standard Errors of Percentages, 2002-2014

	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PAST YEAR INITIATION OF ILLICIT DRUGS^{1,2}	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
FIRST DRUG INITIATED³													
Marijuana and Hashish	1.67	1.93	1.87	2.39	1.77	1.82	1.80	1.98	1.87	1.73	1.65	1.79	1.75
Cocaine	0.38	0.29	1.10	0.26	0.40	0.21	0.32	0.13	0.08	0.09	0.09	0.07	0.34
Crack	*	*	*	0.04	*	*	*	*	*	*	*	*	*
Heroin	0.02	0.02	0.03	*	0.01	*	0.09	0.08	0.08	0.06	0.05	*	*
Hallucinogens	0.54	0.50	0.40	0.45	0.79	0.40	0.71	0.51	0.53	0.55	0.42	0.65	0.45
LSD	0.02	0.17	0.05	0.06	0.14	0.08	0.06	0.02	0.05	0.06	0.04	0.10	0.18
PCP	0.06	0.01	0.05	0.02	0.16	*	0.02	0.01	0.06	0.11	0.18	0.02	*
Ecstasy	0.42	0.38	0.30	0.21	0.73	0.21	0.25	0.31	0.36	0.46	0.22	0.30	0.33
Inhalants	0.69	1.09	0.83	0.94	0.78	0.87	0.81	0.79	0.77	0.70	0.69	0.67	0.77
Nonmedical Use of Psychotherapeutics ^{2,4}													
Pain Relievers	1.70	1.98	1.93	2.23	1.85	1.87	1.83	2.15	1.91	1.71	1.62	1.77	1.68
OxyContin [®]	--	--	0.05	0.18	0.03	0.08	0.09	0.28	0.27	0.59	0.14	*	0.18
Tranquilizers	0.54	0.72	0.95	1.42	0.95	1.31	0.85	2.09	0.94	0.82	0.78	1.26	0.94
Stimulants ²	0.54	0.77	0.77	0.46	0.74	0.92	0.62	0.34	1.08	0.53	0.76	0.60	0.92
Methamphetamine ²	0.33	0.16	0.57	0.13	0.20	0.30	0.29	0.14	0.06	0.06	*	0.08	0.16
Sedatives	0.91	0.58	0.35	0.74	0.57	0.41	0.27	0.48	0.94	0.51	0.54	0.13	0.35
Illicit Drugs Other Than Marijuana ^{1,2}	1.67	1.93	1.87	2.39	1.77	1.82	1.80	1.98	1.88	1.73	1.65	1.79	1.75

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ First Drug Initiated is based on imputation-revised month and year and imputed day of first use for each drug. Respondents may be counted in more than one drug category if they reported initiating multiple drugs on the same day the first time they used drugs. Also, several drugs listed are subcategories of other drugs, and will have the same start date. Therefore, the sum of the estimates for the first drug or drugs initiated may not equal the total estimates for past year initiates.⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.48P Past Year Initiation of Illicit Drugs and First Drug Initiated in the Past Year among Persons Aged 12 or Older Who Initiated Illicit Drug Use in the Past Year: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
PAST YEAR INITIATION OF ILLICIT DRUGS^{1,2}	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	
FIRST DRUG INITIATED³													
Marijuana and Hashish	0.0002	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0015	0.0294	0.9605	0.4030	0.2708	
Cocaine	0.2525	0.5678	0.3773	0.2694	0.1794	0.8527	0.5140	0.4743	0.2727	0.3495	0.2574	0.2300	
Crack	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	0.0871	0.6670	0.8500	0.7698	0.1945	0.9078	0.2151	0.9670	0.2034	0.3572	0.8185	0.5596	
LSD	0.1179	0.6324	0.1492	0.1772	0.6201	0.3962	0.2103	0.1099	0.2011	0.2279	0.1513	0.5466	
PCP	*	*	*	*	*	*	*	*	*	*	*	*	*
Ecstasy	0.0702	0.4103	0.9264	0.5993	0.3977	0.2860	0.9925	0.9358	0.4258	0.4890	0.4606	0.8617	
Inhalants	0.0011	0.0000	0.0000	0.0000	0.0003	0.0000	0.0006	0.0003	0.0045	0.1098	0.6777	0.6443	
Nonmedical Use of Psychotherapeutics ^{2,4}													
Pain Relievers	0.0003	0.0016	0.0000	0.0003	0.0000	0.0009	0.0000	0.0186	0.0102	0.4921	0.0205	0.9416	
OxyContin®	--	--	0.0558	0.5767	0.0513	0.1052	0.3326	0.2899	0.7431	0.3733	0.8135	*	
Tranquilizers	0.0051	0.0664	0.2365	0.6890	0.2693	0.5420	0.0717	0.1836	0.4913	0.3136	0.2567	0.8516	
Stimulants ²	0.3330	0.6156	0.4296	0.0564	0.7335	0.7261	0.1885	0.0101	0.1520	0.0737	0.4647	0.0986	
Methamphetamine ²	0.5853	0.8425	0.5240	0.9947	0.9500	0.3231	0.7435	0.9310	0.2177	0.3424	*	0.3669	
Sedatives	0.2712	0.9501	0.5829	0.4905	0.9916	0.7441	0.3469	0.6234	0.5139	0.9154	0.9101	0.0067	
Illicit Drugs Other Than Marijuana ^{1,2}	0.0002	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0015	0.0233	0.9547	0.3978	0.2733	

*Low precision; no estimate reported.

N/A: Not applicable.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.

³ First Drug Initiated is based on imputation-revised month and year and imputed day of first use for each drug. Respondents may be counted in more than one drug category if they reported initiating multiple drugs on the same day the first time they used drugs. Also, several drugs listed are subcategories of other drugs, and will have the same start date. Therefore, the sum of the estimates for the first drug or drugs initiated may not equal the total estimates for past year initiates.

⁴ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.49D Standard Errors of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49: 2002-2014

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	0.39	0.26	0.34	0.39	0.42	0.37	0.40	0.20	0.50	0.39	0.34	0.42	0.31
Marijuana and Hashish	0.19	0.14	0.21	0.22	0.21	0.34	0.35	0.13	0.48	0.18	0.32	0.26	0.28
Cocaine	0.33	0.36	0.45	0.38	0.51	0.52	0.44	0.42	0.96	0.47	0.32	0.36	0.48
Crack	1.35	1.01	1.35	1.28	1.47	2.54	1.68	0.66	2.45	1.05	1.03	2.70	2.78
Heroin	2.51	1.36	2.53	1.55	1.51	1.23	3.18	1.40	1.05	1.22	1.18	1.26	1.28
Hallucinogens	0.39	0.31	0.57	0.42	0.67	0.45	0.34	0.31	0.21	0.30	0.28	0.42	0.25
LSD	0.49	0.32	0.71	0.78	0.72	0.29	0.31	0.21	0.40	0.27	0.44	0.37	0.33
PCP	0.27	0.49	1.17	0.40	0.61	0.51	0.63	0.80	0.57	0.94	0.50	0.61	0.55
Ecstasy	0.63	0.50	0.61	0.73	0.76	0.57	0.61	0.39	0.44	0.44	0.42	0.58	0.47
Inhalants	0.17	0.26	0.29	0.34	0.24	0.54	0.25	0.67	0.35	0.33	0.32	0.95	0.80
Nonmedical Use of Psychotherapeutics ^{2,3}	0.50	0.48	0.40	0.52	0.56	0.52	0.51	0.42	0.56	0.60	0.54	0.64	0.40
Pain Relievers	0.50	0.45	0.42	0.47	0.57	0.45	0.48	0.45	0.45	0.72	0.58	0.62	0.48
OxyContin [®]	--	--	0.97	1.11	0.94	1.26	1.07	1.27	0.97	1.36	1.40	1.26	0.72
Tranquilizers	0.69	0.77	0.75	0.84	0.62	0.67	0.77	0.63	0.74	0.83	0.55	0.98	0.55
Stimulants ²	0.51	0.96	0.62	0.64	1.16	1.09	0.80	0.72	1.27	0.88	1.13	0.71	0.77
Methamphetamine ²	0.66	1.04	0.83	0.73	2.14	1.18	0.90	0.94	0.81	0.86	0.79	0.63	1.07
Sedatives	2.34	2.43	1.89	1.58	1.78	1.54	1.75	1.02	2.01	1.83	2.02	1.73	1.30
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.43	0.41	0.37	0.47	0.49	0.43	0.44	0.33	0.48	0.49	0.44	0.52	0.34
CIGARETTES	0.30	0.26	0.19	0.33	0.21	0.12	0.26	0.25	0.21	0.13	0.23	0.22	0.23
Daily Cigarette Use ⁴	0.72	0.62	0.40	0.71	0.53	0.45	0.78	0.69	0.34	0.44	0.51	0.65	0.49
SMOKELESS TOBACCO⁵	0.64	0.40	0.47	0.45	0.51	0.29	0.49	0.42	0.51	0.60	0.45	0.31	0.39
CIGARS	0.50	0.45	0.37	0.45	0.42	0.42	0.38	0.44	0.37	0.33	0.38	0.62	0.33
ALCOHOL	0.19	0.10	0.12	0.11	0.11	0.13	0.16	0.13	0.16	0.13	0.19	0.14	0.08

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.49P P Values from Tests of Differences of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 to 49: 2014 Versus 2002-2013

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS^{1,2}	0.0203	0.0001	0.0566	0.4617	0.9621	0.0278	0.6673	0.0001	0.9396	0.0793	0.4806	0.9654	
Marijuana and Hashish	0.0000	0.0000	0.0001	0.0015	0.0017	0.0305	0.1035	0.0000	0.8599	0.0023	0.1414	0.1891	
Cocaine	0.0006	0.0014	0.0095	0.0006	0.0389	0.0290	0.0035	0.0044	0.5843	0.0113	0.0026	0.0234	
Crack	0.6483	0.1244	0.1510	0.3267	0.2584	0.3989	0.8377	0.0494	0.6659	0.0657	0.0540	0.6075	
Heroin	0.0150	0.0003	0.2080	0.0054	0.0005	0.0010	0.1895	0.1493	0.0002	0.0013	0.0056	0.0511	
Hallucinogens	0.0975	0.0005	0.3235	0.2201	0.5932	0.7615	0.0899	0.2984	0.0022	0.1395	0.5102	0.2311	
LSD	0.0002	0.0000	0.1028	0.1137	0.7672	0.0014	0.0054	0.0009	0.2275	0.0158	0.2472	0.9288	
PCP	0.0104	0.6416	0.3627	0.0782	0.1002	0.0802	0.0393	0.3272	0.8644	0.9541	0.1275	0.4184	
Ecstasy	0.8079	0.0595	0.0536	0.7165	0.6514	0.3005	0.3491	0.2003	0.0155	0.0236	0.2470	0.4891	
Inhalants	0.0038	0.0080	0.0100	0.0136	0.0023	0.2470	0.0058	0.1885	0.0263	0.0320	0.1102	0.4603	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.4371	0.7087	0.8162	0.6001	0.1294	0.9856	0.7419	0.1485	0.4406	0.4373	0.1116	0.4268	
Pain Relievers	0.5537	0.7973	0.6149	0.9726	0.3386	0.9940	0.9022	0.4872	0.7144	0.5302	0.1690	0.5873	
OxyContin [®]	--	--	0.1532	0.1202	0.2237	0.0549	0.6592	0.4330	0.1768	0.2830	0.5718	0.0998	
Tranquilizers	0.6705	0.4835	0.2925	0.0253	0.4575	0.1913	0.3065	0.2479	0.1829	0.2285	0.7582	0.0804	
Stimulants ²	0.0127	0.9542	0.1208	0.1529	0.3045	0.7838	0.7968	0.9621	0.8248	0.5669	0.6729	0.9965	
Methamphetamine ²	0.0147	0.2683	0.2837	0.0101	0.9366	0.1035	0.0508	0.0608	0.0188	0.0032	0.0855	0.0124	
Sedatives	0.0568	0.9562	0.3556	0.4746	0.0223	0.1590	0.9313	0.2996	0.3906	0.8004	0.0479	0.1002	
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.1075	0.0344	0.1334	0.8330	0.8868	0.2819	0.6959	0.0057	0.4920	0.3981	0.2805	0.9957	
CIGARETTES	0.0000	0.0000	0.0000	0.0012	0.0000	0.0000	0.0004	0.0013	0.0000	0.0000	0.0197	0.0156	
Daily Cigarette Use ⁴	0.4453	0.3223	0.0060	0.2994	0.0175	0.0434	0.5911	0.8742	0.0149	0.0262	0.3499	0.3601	
SMOKELESS TOBACCO⁵	0.9065	0.1354	0.7486	0.2853	0.9515	0.0380	0.8525	0.8492	0.5959	0.2484	0.7495	0.2644	
CIGARS	0.0200	0.8441	0.6831	0.1369	0.3768	0.8087	0.4169	0.6012	0.8590	0.0908	0.8329	0.0916	
ALCOHOL	0.0005	0.0000	0.0000	0.0000	0.0000	0.0022	0.0667	0.0099	0.3825	0.1532	0.6983	0.9169	

*Low precision; no estimate reported.

-- Not available.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Estimates in these designated rows do not include data from new methamphetamine initiation items added in 2007 or new methamphetamine use items added in 2005 and 2006.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.⁵ Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.50C Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	224	206	210	203	221	231	210	224	245	204	258	239	220
Marijuana and Hashish	158	140	158	143	154	155	157	161	184	149	175	183	156
Cocaine	119	115	114	108	127	125	100	101	101	82	117	81	87
Heroin	45	37	49	36	66	43	46	64	55	63	70	62	63
Hallucinogens	38	34	44	36	41	42	38	41	44	41	46	40	34
Inhalants	31	27	28	38	27	22	28	20	28	19	61	26	19
Nonmedical Use of Psychotherapeutics ^{2,3}	139	116	115	108	115	129	112	131	152	123	158	144	122
Pain Relievers	126	102	86	97	104	116	100	115	139	115	144	133	113
Tranquilizers	76	56	73	51	53	65	50	62	85	52	79	56	52
Stimulants ²	68	41	48	44	46	63	43	46	51	45	77	74	51
Sedatives	57	37	27	20	43	36	27	34	39	26	36	27	35
ALCOHOL	415	387	417	400	411	409	398	419	412	377	419	415	362
BOTH ILLICIT DRUGS AND ALCOHOL¹	145	138	143	141	141	145	128	157	131	122	160	140	138
ILLCIT DRUGS OR ALCOHOL¹	439	410	435	426	455	456	426	457	467	407	470	460	389

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.50D Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.10	0.09	0.09	0.08	0.09	0.09	0.08	0.09	0.10	0.08	0.10	0.09	0.08
Marijuana and Hashish	0.07	0.06	0.07	0.06	0.06	0.06	0.06	0.06	0.07	0.06	0.07	0.07	0.06
Cocaine	0.05	0.05	0.05	0.04	0.05	0.05	0.04	0.04	0.04	0.03	0.05	0.03	0.03
Heroin	0.02	0.02	0.02	0.01	0.03	0.02	0.02	0.03	0.02	0.02	0.03	0.02	0.02
Hallucinogens	0.02	0.01	0.02	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.01
Inhalants	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01
Nonmedical Use of Psychotherapeutics ^{2,3}	0.06	0.05	0.05	0.04	0.05	0.05	0.04	0.05	0.06	0.05	0.06	0.05	0.05
Pain Relievers	0.05	0.04	0.04	0.04	0.04	0.05	0.04	0.05	0.05	0.04	0.06	0.05	0.04
Tranquilizers	0.03	0.02	0.03	0.02	0.02	0.03	0.02	0.02	0.03	0.02	0.03	0.02	0.02
Stimulants ²	0.03	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.03	0.03	0.02
Sedatives	0.02	0.02	0.01	0.01	0.02	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01
ALCOHOL	0.18	0.16	0.17	0.16	0.17	0.17	0.16	0.17	0.16	0.15	0.16	0.16	0.14
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.06	0.06	0.06	0.06	0.06	0.06	0.05	0.06	0.05	0.05	0.06	0.05	0.05
ILLCIT DRUGS OR ALCOHOL¹	0.19	0.17	0.18	0.18	0.18	0.18	0.17	0.18	0.18	0.16	0.18	0.18	0.15

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.50P Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0050	0.0865	0.0025	0.2355	0.1298	0.4203	0.2453	0.2021	0.2489	0.2429	0.2714	0.6367	
Marijuana and Hashish	0.0050	0.0220	0.0014	0.2023	0.1436	0.8612	0.1724	0.1054	0.0317	0.6127	0.3736	0.7610	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0659	0.2886	0.5723	0.1239	0.6800	
Heroin	0.0000	0.0000	0.0005	0.0000	0.0127	0.0000	0.0003	0.0323	0.0148	0.1039	0.2493	0.4789	
Hallucinogens	0.0000	0.0268	0.0000	0.0023	0.0035	0.0078	0.0089	0.0082	0.0023	0.0507	0.1155	0.5210	
Inhalants	0.0067	0.0094	0.0000	0.0016	0.0068	0.0093	0.0108	0.0072	0.0209	0.0686	0.2740	0.2391	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.4751	0.1266	0.3643	0.0984	0.2012	0.5894	0.5308	0.9955	0.7321	0.2207	0.2563	0.5527	
Pain Relievers	0.2360	0.0401	0.0089	0.1356	0.3313	0.6223	0.5307	0.7185	0.6172	0.5511	0.3356	0.9157	
Tranquilizers	0.3083	0.8764	0.0953	0.8410	0.6281	0.9837	0.9056	0.7238	0.4724	0.4193	0.0780	0.5586	
Stimulants ²	0.8650	0.4284	0.5728	0.6632	0.4183	0.6069	0.1311	0.2836	0.1691	0.0474	0.4589	0.9782	
Sedatives	0.6692	0.5294	0.9741	0.3694	0.8283	0.6755	0.8587	0.8097	0.6099	0.1554	0.9147	0.3350	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0015	0.7572	0.0575	0.3824	
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.0000	0.0001	0.0000	0.0000	0.0000	0.0001	0.0003	0.0001	0.0274	0.6584	0.1558	0.8997	
ILLCIT DRUGS OR ALCOHOL¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0002	0.0052	0.6319	0.0654	0.6153	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.51C Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	50	46	46	47	46	41	46	44	47	47	45	41	46
Marijuana and Hashish	44	38	40	41	40	36	39	40	40	41	41	38	40
Cocaine	16	12	11	14	14	13	13	9	8	9	9	6	9
Heroin	4	4	7	4	4	3	5	6	3	10	8	4	7
Hallucinogens	19	13	16	17	15	15	14	14	18	13	13	10	11
Inhalants	12	13	15	15	14	14	14	12	13	12	9	8	10
Nonmedical Use of Psychotherapeutics ^{2,3}	25	25	26	28	26	24	25	20	22	24	19	17	23
Pain Relievers	22	22	24	27	23	20	21	19	20	22	16	15	19
Tranquilizers	13	15	11	13	11	11	12	8	11	11	11	9	12
Stimulants ²	13	13	13	13	16	12	11	10	9	11	9	9	13
Sedatives	8	10	6	10	6	6	6	5	5	4	5	4	10
ALCOHOL	51	50	50	51	49	48	49	49	49	45	40	36	42
BOTH ILLICIT DRUGS AND ALCOHOL¹	34	30	33	34	33	29	29	29	31	30	25	21	29
ILLCIT DRUGS OR ALCOHOL¹	60	61	61	60	59	57	59	58	60	58	55	51	55

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.51D Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.20	0.18	0.18	0.19	0.18	0.16	0.18	0.18	0.19	0.19	0.18	0.17	0.18
Marijuana and Hashish	0.18	0.15	0.16	0.16	0.16	0.14	0.16	0.16	0.16	0.16	0.16	0.15	0.16
Cocaine	0.07	0.05	0.05	0.05	0.05	0.05	0.05	0.04	0.03	0.04	0.04	0.02	0.03
Heroin	0.02	0.01	0.03	0.01	0.01	0.01	0.02	0.02	0.01	0.04	0.03	0.02	0.03
Hallucinogens	0.08	0.05	0.06	0.07	0.06	0.06	0.06	0.06	0.07	0.05	0.05	0.04	0.04
Inhalants	0.05	0.05	0.06	0.06	0.06	0.05	0.06	0.05	0.05	0.05	0.04	0.03	0.04
Nonmedical Use of Psychotherapeutics ^{2,3}	0.10	0.10	0.10	0.11	0.10	0.09	0.10	0.08	0.09	0.10	0.08	0.07	0.09
Pain Relievers	0.09	0.09	0.10	0.11	0.09	0.08	0.08	0.08	0.08	0.09	0.07	0.06	0.08
Tranquilizers	0.05	0.06	0.04	0.05	0.04	0.04	0.05	0.03	0.05	0.04	0.04	0.04	0.05
Stimulants ²	0.05	0.05	0.05	0.05	0.06	0.05	0.05	0.04	0.04	0.05	0.03	0.03	0.05
Sedatives	0.03	0.04	0.02	0.04	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.01	0.04
ALCOHOL	0.20	0.20	0.20	0.20	0.19	0.19	0.20	0.20	0.20	0.18	0.16	0.14	0.17
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.14	0.12	0.13	0.13	0.13	0.12	0.12	0.12	0.13	0.12	0.10	0.09	0.12
ILLCIT DRUGS OR ALCOHOL¹	0.24	0.25	0.24	0.24	0.23	0.23	0.24	0.23	0.25	0.23	0.22	0.20	0.22

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.51P Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0000	0.0017	0.0000	0.0000	0.0494	0.9051	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0001	0.0010	0.0431	0.0013	0.0019	0.0001	0.0002	0.0160	0.3803	
Cocaine	0.0000	0.0001	0.0000	0.0001	0.0000	0.0000	0.0004	0.1428	0.5641	0.2128	0.3269	0.6439	
Heroin	0.5978	0.4938	0.8030	0.2696	0.4709	0.2314	0.8953	0.9366	0.1764	0.3098	0.8318	0.3710	
Hallucinogens	0.0001	0.0029	0.0006	0.0004	0.0006	0.0011	0.0178	0.0018	0.0001	0.0784	0.0564	0.5718	
Inhalants	0.0001	0.0001	0.0000	0.0001	0.0001	0.0005	0.0008	0.0006	0.0021	0.0101	0.1543	0.8930	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0071	0.0005	0.0001	0.0252	0.0037	0.0362	0.0055	0.4184	0.0706	0.0814	0.1951	0.0264	
Pain Relievers	0.0161	0.0001	0.0000	0.0019	0.0018	0.0163	0.0026	0.0496	0.0071	0.0058	0.4652	0.1711	
Tranquilizers	0.0423	0.0212	0.1679	0.3074	0.5658	0.5916	0.1923	0.9276	0.2816	0.5140	0.9041	0.4089	
Stimulants ²	0.0427	0.0443	0.1706	0.5359	0.0992	0.5065	0.6741	0.9067	0.6788	0.6849	0.5130	0.2344	
Sedatives	0.9311	0.4609	0.5505	0.7898	0.5716	0.6111	0.5765	0.3734	0.2619	0.1123	0.3327	0.0920	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0024	0.7517	
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0326	0.3272	0.4807	
ILLCIT DRUGS OR ALCOHOL¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0007	0.5021	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.52C Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	74	73	82	83	81	79	84	83	85	83	94	88	92
Marijuana and Hashish	63	64	73	70	70	69	71	74	76	73	80	76	78
Cocaine	29	28	33	34	31	33	32	28	26	24	27	28	24
Heroin	12	10	12	14	12	14	16	15	18	22	22	23	24
Hallucinogens	24	17	23	22	23	21	25	23	24	22	17	23	20
Inhalants	8	10	11	14	8	8	10	14	13	10	5	13	6
Nonmedical Use of Psychotherapeutics ^{2,3}	34	33	38	44	40	44	43	43	44	43	54	45	45
Pain Relievers	31	30	30	37	32	40	38	36	40	39	47	40	41
Tranquilizers	16	19	17	21	20	18	18	20	19	20	27	21	17
Stimulants ²	16	19	18	21	19	17	17	19	19	15	25	24	19
Sedatives	6	5	7	6	9	6	9	12	7	7	5	10	8
ALCOHOL	112	109	119	119	122	116	116	117	125	116	114	123	120
BOTH ILLICIT DRUGS AND ALCOHOL¹	53	55	60	60	56	59	62	60	58	56	58	58	58
ILLCIT DRUGS OR ALCOHOL¹	121	117	129	132	132	127	125	128	139	128	133	138	134

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.52D Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.24	0.23	0.26	0.26	0.25	0.24	0.25	0.25	0.25	0.24	0.27	0.25	0.26
Marijuana and Hashish	0.20	0.20	0.23	0.22	0.21	0.21	0.22	0.22	0.22	0.21	0.23	0.22	0.22
Cocaine	0.09	0.09	0.10	0.10	0.10	0.10	0.10	0.08	0.08	0.07	0.08	0.08	0.07
Heroin	0.04	0.03	0.04	0.04	0.04	0.04	0.05	0.05	0.05	0.06	0.06	0.07	0.07
Hallucinogens	0.08	0.05	0.07	0.07	0.07	0.06	0.08	0.07	0.07	0.06	0.05	0.07	0.06
Inhalants	0.02	0.03	0.03	0.04	0.02	0.03	0.03	0.04	0.04	0.03	0.01	0.04	0.02
Nonmedical Use of Psychotherapeutics ^{2,3}	0.11	0.11	0.12	0.13	0.12	0.13	0.13	0.13	0.13	0.13	0.16	0.13	0.13
Pain Relievers	0.10	0.09	0.09	0.11	0.10	0.12	0.12	0.11	0.12	0.11	0.14	0.11	0.12
Tranquilizers	0.05	0.06	0.05	0.07	0.06	0.06	0.05	0.06	0.06	0.06	0.08	0.06	0.05
Stimulants ²	0.05	0.06	0.06	0.06	0.06	0.05	0.05	0.06	0.06	0.04	0.07	0.07	0.05
Sedatives	0.02	0.02	0.02	0.02	0.03	0.02	0.03	0.04	0.02	0.02	0.02	0.03	0.02
ALCOHOL	0.36	0.34	0.37	0.37	0.37	0.35	0.35	0.35	0.37	0.34	0.33	0.35	0.34
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.17	0.17	0.19	0.19	0.17	0.18	0.19	0.18	0.17	0.16	0.17	0.17	0.16
ILLCIT DRUGS OR ALCOHOL¹	0.39	0.37	0.40	0.41	0.40	0.39	0.38	0.38	0.41	0.37	0.39	0.40	0.38

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.52P Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 18 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0006	0.0000	0.0000	0.0004	0.0003	0.0008	0.0025	0.0008	0.0160	0.0018	0.0515	
Marijuana and Hashish	0.0002	0.0004	0.0004	0.0014	0.0094	0.0168	0.0138	0.0331	0.0069	0.0102	0.0590	0.0989	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.1524	0.5112	0.3420	0.1936	
Heroin	0.0000	0.0000	0.0009	0.0122	0.0004	0.0027	0.0359	0.0412	0.1310	0.4346	0.8256	0.6425	
Hallucinogens	0.0000	0.0808	0.0004	0.0076	0.0073	0.0289	0.0065	0.0170	0.0136	0.1891	0.8760	0.3735	
Inhalants	0.0801	0.0175	0.0020	0.0156	0.1398	0.1454	0.2835	0.0081	0.0349	0.0476	0.9702	0.0297	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0818	0.8773	0.0141	0.0026	0.0197	0.0042	0.0006	0.0011	0.0006	0.0250	0.0000	0.0843	
Pain Relievers	0.4340	0.3891	0.2276	0.0078	0.0994	0.0047	0.0013	0.0046	0.0008	0.0048	0.0003	0.3197	
Tranquilizers	0.0175	0.0355	0.0118	0.0028	0.0259	0.1236	0.0865	0.0275	0.0940	0.2395	0.0122	0.1039	
Stimulants ²	0.4515	0.4247	0.0569	0.1000	0.5865	0.7082	0.6966	0.5019	0.7554	0.0766	0.2414	0.2259	
Sedatives	0.3663	0.8246	0.5740	0.8669	0.2849	0.8775	0.2425	0.1937	0.5351	0.9129	0.7054	0.4471	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.1249	
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0018	0.0054	0.0397	
ILLCIT DRUGS OR ALCOHOL¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0749	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.53C Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	202	180	189	175	197	205	176	202	214	177	234	219	187
Marijuana and Hashish	130	114	131	118	131	127	124	134	158	118	141	164	126
Cocaine	114	109	110	100	124	118	93	97	98	81	113	76	84
Heroin	43	34	47	33	65	41	42	62	52	58	66	56	56
Hallucinogens	20	26	34	21	30	32	24	32	32	35	40	30	23
Inhalants	28	17	21	33	21	15	22	9	21	11	61	22	16
Nonmedical Use of Psychotherapeutics ^{2,3}	131	107	105	94	105	119	98	121	142	110	147	134	112
Pain Relievers	118	94	76	84	96	107	88	108	129	103	134	126	105
Tranquilizers	74	52	70	42	47	61	46	58	81	48	73	50	49
Stimulants ²	65	33	43	36	38	59	39	39	42	42	74	69	48
Sedatives	56	34	25	16	41	35	25	32	38	25	36	25	32
ALCOHOL	391	356	376	355	382	386	368	396	388	354	386	392	324
BOTH ILLICIT DRUGS AND ALCOHOL¹	129	117	118	121	127	128	102	138	110	106	144	126	117
ILLCIT DRUGS OR ALCOHOL¹	406	374	395	376	415	423	387	431	431	380	427	429	348

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.53D Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLICIT DRUGS¹	0.11	0.10	0.10	0.09	0.11	0.11	0.09	0.10	0.11	0.09	0.12	0.11	0.09
Marijuana and Hashish	0.07	0.06	0.07	0.06	0.07	0.07	0.06	0.07	0.08	0.06	0.07	0.08	0.06
Cocaine	0.06	0.06	0.06	0.05	0.07	0.06	0.05	0.05	0.05	0.04	0.06	0.04	0.04
Heroin	0.02	0.02	0.03	0.02	0.03	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03
Hallucinogens	0.01	0.01	0.02	0.01	0.02	0.02	0.01	0.02	0.02	0.02	0.02	0.01	0.01
Inhalants	0.02	0.01	0.01	0.02	0.01	0.01	0.01	0.00	0.01	0.01	0.03	0.01	0.01
Nonmedical Use of Psychotherapeutics ^{2,3}	0.07	0.06	0.06	0.05	0.06	0.06	0.05	0.06	0.07	0.06	0.07	0.07	0.05
Pain Relievers	0.07	0.05	0.04	0.05	0.05	0.06	0.05	0.06	0.07	0.05	0.07	0.06	0.05
Tranquilizers	0.04	0.03	0.04	0.02	0.03	0.03	0.02	0.03	0.04	0.02	0.04	0.02	0.02
Stimulants ²	0.04	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.04	0.03	0.02
Sedatives	0.03	0.02	0.01	0.01	0.02	0.02	0.01	0.02	0.02	0.01	0.02	0.01	0.02
ALCOHOL	0.22	0.20	0.21	0.19	0.20	0.20	0.19	0.20	0.20	0.18	0.19	0.19	0.16
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.07	0.06	0.06	0.07	0.07	0.07	0.05	0.07	0.06	0.05	0.07	0.06	0.06
ILLICIT DRUGS OR ALCOHOL¹	0.23	0.21	0.22	0.20	0.22	0.22	0.20	0.22	0.22	0.19	0.21	0.21	0.17

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.53P Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 26 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLICIT DRUGS¹	0.4269	0.1327	0.4207	0.0147	0.2479	0.1155	0.1345	0.4357	0.1651	0.0002	0.5223	0.1406	
Marijuana and Hashish	0.2483	0.1439	0.7536	0.0295	0.2584	0.0414	0.3345	0.6619	0.8706	0.0173	0.3827	0.4144	
Cocaine	0.0039	0.0016	0.0032	0.0071	0.0005	0.0040	0.0217	0.4517	0.5149	0.3689	0.2058	0.4133	
Heroin	0.0027	0.0003	0.0097	0.0001	0.1443	0.0003	0.0022	0.1070	0.0512	0.1014	0.1733	0.3743	
Hallucinogens	0.5744	0.9858	0.1995	0.7550	0.9507	0.7335	0.9764	0.7881	0.9336	0.4376	0.2639	0.9287	
Inhalants	0.6767	0.5769	0.7057	0.4723	0.9660	0.8919	0.6355	0.1915	0.8954	0.3940	0.4194	0.9094	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0547	0.0095	0.0029	0.0001	0.0021	0.0308	0.0049	0.1571	0.2323	0.0080	0.8791	0.3609	
Pain Relievers	0.0457	0.0052	0.0000	0.0003	0.0175	0.0369	0.0075	0.3086	0.2707	0.0200	0.8534	0.7931	
Tranquilizers	0.9856	0.1997	0.4501	0.0668	0.1161	0.5535	0.4052	0.6857	0.9146	0.1367	0.3416	0.2717	
Stimulants ²	0.5713	0.0403	0.5249	0.1214	0.0728	0.5393	0.1020	0.1379	0.1051	0.1099	0.6482	0.7911	
Sedatives	0.7728	0.7172	0.9591	0.2174	0.7014	0.6086	0.6797	0.9627	0.5178	0.2840	0.8503	0.3958	
ALCOHOL	0.1731	0.5582	0.0928	0.1303	0.1477	0.1727	0.5548	0.0870	0.8764	0.0781	0.7613	0.7025	
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.6044	0.9150	0.4472	0.4784	0.6204	0.7779	0.6548	0.2291	0.4344	0.0821	0.8965	0.4687	
ILLICIT DRUGS OR ALCOHOL¹	0.4705	0.8103	0.3441	0.9815	0.5446	0.7115	0.9647	0.4215	0.7312	0.0028	0.9098	0.8487	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.54C Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	179	161	167	173	184	195	183	183	205	177	215	211	186
Marijuana and Hashish	109	104	115	119	115	127	122	117	154	119	136	164	124
Cocaine	97	89	101	101	112	106	86	81	83	70	99	73	66
Heroin	45	36	45	36	65	40	44	61	49	57	70	57	59
Hallucinogens	17	24	25	21	18	22	22	34	23	26	27	20	19
Inhalants	9	8	21	11	10	11	23	13	20	13	10	10	13
Nonmedical Use of Psychotherapeutics ^{2,3}	113	97	93	89	88	114	96	116	133	109	135	129	103
Pain Relievers	103	88	76	83	82	104	88	99	108	102	125	121	96
Tranquilizers	61	44	43	39	36	52	40	49	75	37	58	47	44
Stimulants ²	38	32	44	35	38	60	36	42	42	41	55	71	44
Sedatives	56	31	23	14	18	21	25	33	36	26	35	23	33
ALCOHOL	283	265	297	262	284	284	296	295	304	260	300	286	246
BOTH ILLICIT DRUGS AND ALCOHOL¹	105	75	93	92	94	98	92	102	89	98	121	87	93
ILLCIT DRUGS OR ALCOHOL¹	323	300	320	302	329	327	330	340	365	293	353	336	291

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.54D Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.08	0.07	0.07	0.07	0.07	0.08	0.07	0.07	0.08	0.07	0.08	0.08	0.07
Marijuana and Hashish	0.05	0.04	0.05	0.05	0.05	0.05	0.05	0.05	0.06	0.05	0.05	0.06	0.05
Cocaine	0.04	0.04	0.04	0.04	0.05	0.04	0.03	0.03	0.03	0.03	0.04	0.03	0.02
Heroin	0.02	0.01	0.02	0.01	0.03	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.02
Hallucinogens	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Inhalants	0.00	0.00	0.01	0.00	0.00	0.00	0.01	0.01	0.01	0.00	0.00	0.00	0.00
Nonmedical Use of Psychotherapeutics ^{2,3}	0.05	0.04	0.04	0.04	0.04	0.05	0.04	0.05	0.05	0.04	0.05	0.05	0.04
Pain Relievers	0.04	0.04	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.04	0.05	0.05	0.04
Tranquilizers	0.03	0.02	0.02	0.02	0.01	0.02	0.02	0.02	0.03	0.01	0.02	0.02	0.02
Stimulants ²	0.02	0.01	0.02	0.01	0.02	0.02	0.01	0.02	0.02	0.02	0.02	0.03	0.02
Sedatives	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
ALCOHOL	0.12	0.11	0.12	0.11	0.12	0.11	0.12	0.12	0.12	0.10	0.12	0.11	0.09
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.04	0.03	0.04	0.04	0.04	0.04	0.04	0.04	0.03	0.04	0.05	0.03	0.04
ILLCIT DRUGS OR ALCOHOL¹	0.14	0.13	0.13	0.12	0.13	0.13	0.13	0.14	0.14	0.11	0.14	0.13	0.11

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.54P Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.3894	0.6647	0.1711	0.5640	0.4078	0.9054	0.4870	0.5599	0.7851	0.2631	0.5681	0.9361	
Marijuana and Hashish	0.0955	0.3632	0.1058	0.2671	0.3227	0.8850	0.2414	0.4643	0.1627	0.7209	0.6425	0.3900	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0535	0.1720	0.7809	0.0311	0.5481	
Heroin	0.0001	0.0000	0.0002	0.0000	0.0181	0.0000	0.0004	0.0242	0.0096	0.0568	0.3164	0.2273	
Hallucinogens	0.0291	0.0838	0.0022	0.0367	0.2082	0.4999	0.1929	0.0126	0.0211	0.1035	0.3311	0.7463	
Inhalants	0.5732	0.9034	0.0360	0.6798	0.6030	0.7807	0.1087	0.2117	0.3084	0.8042	0.5870	0.8302	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0066	0.0048	0.0338	0.0122	0.0021	0.2546	0.1184	0.8817	0.9621	0.2547	0.4841	0.6752	
Pain Relievers	0.0029	0.0010	0.0006	0.0204	0.0071	0.3120	0.1537	0.8521	0.7657	0.4963	0.5617	0.9808	
Tranquilizers	0.9901	0.3233	0.9289	0.7576	0.1188	0.3301	0.5526	0.8953	0.6323	0.0844	0.6147	0.4936	
Stimulants ²	0.4764	0.4221	0.3609	0.5136	0.5216	0.8894	0.1349	0.7389	0.2376	0.2132	0.4737	0.6721	
Sedatives	0.6447	0.8336	0.6923	0.1280	0.1355	0.7142	0.6836	0.8935	0.6840	0.2845	0.9553	0.1876	
ALCOHOL	0.0028	0.3316	0.0052	0.0961	0.0100	0.0112	0.0011	0.0007	0.1034	0.8248	0.0624	0.9627	
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.0057	0.4073	0.0007	0.0033	0.0116	0.0067	0.0183	0.0103	0.2932	0.4373	0.1834	0.8575	
ILLCIT DRUGS OR ALCOHOL¹	0.0282	0.7225	0.0244	0.3900	0.0522	0.1562	0.0103	0.0135	0.2162	0.2540	0.1369	0.9503	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.55C Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	38	34	36	37	34	30	36	33	34	37	32	29	34
Marijuana and Hashish	33	29	29	32	27	25	29	29	29	32	29	27	26
Cocaine	8	10	10	10	9	10	10	7	6	7	5	4	5
Heroin	3	3	4	2	2	3	4	4	3	9	5	2	6
Hallucinogens	7	9	11	12	9	8	9	10	12	8	7	8	6
Inhalants	7	7	11	7	7	7	9	6	6	6	5	5	6
Nonmedical Use of Psychotherapeutics ^{2,3}	19	21	21	23	19	20	18	15	18	17	13	12	19
Pain Relievers	16	17	19	22	16	15	16	14	16	15	12	11	15
Tranquilizers	7	9	6	8	7	9	5	5	8	5	8	7	7
Stimulants ²	9	11	9	8	12	10	9	8	7	10	7	5	8
Sedatives	6	9	6	6	6	5	3	3	4	4	4	2	9
ALCOHOL	31	31	31	35	28	31	29	31	30	26	25	22	25
BOTH ILLICIT DRUGS AND ALCOHOL¹	21	17	19	23	17	16	16	15	16	14	12	13	15
ILLCIT DRUGS OR ALCOHOL¹	45	44	43	45	40	41	43	43	42	43	39	35	39

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.55D Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.16	0.14	0.14	0.14	0.13	0.12	0.14	0.13	0.14	0.15	0.13	0.12	0.14
Marijuana and Hashish	0.13	0.11	0.11	0.13	0.11	0.10	0.12	0.12	0.12	0.13	0.12	0.11	0.10
Cocaine	0.03	0.04	0.04	0.04	0.04	0.04	0.04	0.03	0.02	0.03	0.02	0.01	0.02
Heroin	0.01	0.01	0.02	0.01	0.01	0.01	0.02	0.02	0.01	0.04	0.02	0.01	0.02
Hallucinogens	0.03	0.04	0.04	0.05	0.03	0.03	0.04	0.04	0.05	0.03	0.03	0.03	0.02
Inhalants	0.03	0.03	0.04	0.03	0.03	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02
Nonmedical Use of Psychotherapeutics ^{2,3}	0.08	0.08	0.08	0.09	0.08	0.08	0.07	0.06	0.07	0.07	0.05	0.05	0.08
Pain Relievers	0.06	0.07	0.07	0.09	0.06	0.06	0.06	0.06	0.07	0.06	0.05	0.04	0.06
Tranquilizers	0.03	0.03	0.02	0.03	0.03	0.04	0.02	0.02	0.03	0.02	0.03	0.03	0.03
Stimulants ²	0.03	0.04	0.04	0.03	0.05	0.04	0.04	0.03	0.03	0.04	0.03	0.02	0.03
Sedatives	0.02	0.03	0.02	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.01	0.03
ALCOHOL	0.12	0.12	0.12	0.14	0.11	0.12	0.12	0.13	0.12	0.10	0.10	0.09	0.10
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.08	0.07	0.08	0.09	0.07	0.07	0.06	0.06	0.07	0.06	0.05	0.05	0.06
ILLCIT DRUGS OR ALCOHOL¹	0.18	0.17	0.17	0.18	0.16	0.16	0.17	0.17	0.17	0.17	0.15	0.14	0.16

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.55P Substance Dependence for Specific Substances in the Past Year among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0004	0.0128	0.0001	0.0091	0.0002	0.0005	0.3516	0.5862	
Marijuana and Hashish	0.0000	0.0000	0.0000	0.0000	0.0003	0.0427	0.0001	0.0003	0.0000	0.0001	0.0209	0.0374	
Cocaine	0.0001	0.0002	0.0000	0.0000	0.0004	0.0000	0.0075	0.1505	0.2162	0.1358	0.9785	0.9737	
Heroin	0.4503	0.5354	0.8172	0.1939	0.3034	0.4221	0.7654	0.8985	0.4279	0.4199	0.5563	0.1218	
Hallucinogens	0.0296	0.0124	0.0048	0.0046	0.0507	0.0755	0.1065	0.0101	0.0009	0.0942	0.3279	0.5738	
Inhalants	0.0224	0.1353	0.0152	0.0578	0.0546	0.1602	0.0882	0.0873	0.2450	0.4629	0.6653	0.9377	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.1117	0.0089	0.0261	0.2268	0.1724	0.2170	0.2560	0.8325	0.2256	0.5618	0.1453	0.0310	
Pain Relievers	0.1683	0.0091	0.0087	0.0348	0.1912	0.2543	0.1058	0.4655	0.0452	0.2160	0.4676	0.3493	
Tranquilizers	0.3365	0.1991	0.8527	0.4395	0.9952	0.3537	0.7724	0.8749	0.3512	0.6596	0.7270	0.7024	
Stimulants ²	0.0731	0.0625	0.2045	0.7833	0.0745	0.4709	0.6151	0.2890	0.9612	0.4463	0.9646	0.4292	
Sedatives	0.5252	0.4448	0.5420	0.4558	0.4530	0.4996	0.1324	0.1451	0.4130	0.2859	0.2125	0.0513	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0010	0.0452	0.5138	
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.0000	0.0001	0.0000	0.0000	0.0008	0.0001	0.0010	0.0016	0.0029	0.0385	0.4489	0.3866	
ILLCIT DRUGS OR ALCOHOL¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0707	0.5765	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.56C Substance Dependence for Specific Substances in the Past Year among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	58	59	68	70	68	70	70	68	76	72	78	73	80
Marijuana and Hashish	47	51	60	57	57	61	57	59	65	59	64	61	67
Cocaine	23	22	26	29	27	26	23	21	20	21	22	24	19
Heroin	12	9	10	14	11	13	15	14	17	22	21	20	23
Hallucinogens	14	12	15	16	13	9	13	16	16	15	10	12	12
Inhalants	3	4	6	5	7	3	6	8	6	2	3	7	3
Nonmedical Use of Psychotherapeutics ^{2,3}	29	28	31	33	31	35	36	37	41	39	46	40	39
Pain Relievers	24	23	25	28	26	33	33	33	38	37	41	35	36
Tranquilizers	10	10	13	16	12	10	11	12	14	12	21	17	13
Stimulants ²	13	16	16	17	17	13	16	16	17	13	18	21	17
Sedatives	5	4	6	3	7	5	7	11	5	7	5	9	8
ALCOHOL	71	70	73	79	81	79	79	78	82	74	77	78	78
BOTH ILLICIT DRUGS AND ALCOHOL¹	28	32	37	35	37	37	39	38	40	36	38	38	38
ILLCIT DRUGS OR ALCOHOL¹	87	82	88	97	100	99	95	94	104	93	104	102	104

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.56D Substance Dependence for Specific Substances in the Past Year among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.19	0.19	0.21	0.21	0.21	0.21	0.21	0.20	0.22	0.21	0.22	0.21	0.23
Marijuana and Hashish	0.15	0.16	0.19	0.17	0.17	0.19	0.17	0.18	0.19	0.17	0.19	0.18	0.19
Cocaine	0.08	0.07	0.08	0.09	0.08	0.08	0.07	0.06	0.06	0.06	0.07	0.07	0.06
Heroin	0.04	0.03	0.03	0.04	0.03	0.04	0.05	0.04	0.05	0.06	0.06	0.06	0.07
Hallucinogens	0.05	0.04	0.05	0.05	0.04	0.03	0.04	0.05	0.05	0.04	0.03	0.04	0.03
Inhalants	0.01	0.01	0.02	0.02	0.02	0.01	0.02	0.02	0.02	0.01	0.01	0.02	0.01
Nonmedical Use of Psychotherapeutics ^{2,3}	0.09	0.09	0.10	0.10	0.10	0.11	0.11	0.11	0.12	0.11	0.13	0.11	0.11
Pain Relievers	0.08	0.07	0.08	0.09	0.08	0.10	0.10	0.10	0.11	0.11	0.12	0.10	0.10
Tranquilizers	0.03	0.03	0.04	0.05	0.04	0.03	0.03	0.04	0.04	0.03	0.06	0.05	0.04
Stimulants ²	0.04	0.05	0.05	0.05	0.05	0.04	0.05	0.05	0.05	0.04	0.05	0.06	0.05
Sedatives	0.02	0.01	0.02	0.01	0.02	0.02	0.02	0.03	0.02	0.02	0.01	0.03	0.02
ALCOHOL	0.23	0.22	0.23	0.24	0.25	0.24	0.24	0.23	0.24	0.22	0.22	0.22	0.22
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.09	0.10	0.12	0.11	0.11	0.11	0.12	0.11	0.12	0.11	0.11	0.11	0.11
ILLCIT DRUGS OR ALCOHOL¹	0.28	0.26	0.27	0.30	0.31	0.30	0.29	0.28	0.30	0.27	0.30	0.29	0.30

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.56P Substance Dependence for Specific Substances in the Past Year among Persons Aged 18 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.0100	0.1186	0.0108	0.0005	0.0245	0.0248	0.0028	0.0251	0.0381	0.0782	0.0126	0.1528	
Marijuana and Hashish	0.0180	0.0060	0.0371	0.0227	0.0903	0.0890	0.0174	0.0753	0.0660	0.0905	0.1881	0.2337	
Cocaine	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0114	0.3273	0.3426	0.4154	0.3760	
Heroin	0.0001	0.0000	0.0002	0.0202	0.0004	0.0012	0.0332	0.0213	0.1767	0.5486	0.8640	0.9092	
Hallucinogens	0.0153	0.3269	0.0175	0.0835	0.2034	0.8085	0.3407	0.0818	0.1387	0.4009	0.8587	0.9935	
Inhalants	0.6876	0.4034	0.0568	0.6685	0.1952	0.6442	0.6282	0.0374	0.1101	0.6807	0.7547	0.1718	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.5013	0.0560	0.3710	0.1574	0.5807	0.2112	0.0301	0.0156	0.0099	0.0560	0.0008	0.1616	
Pain Relievers	0.2073	0.0099	0.8992	0.4730	0.8124	0.1436	0.0469	0.0113	0.0134	0.0095	0.0023	0.4159	
Tranquilizers	0.8552	0.5705	0.2867	0.0900	0.6484	0.2912	0.9123	0.7076	0.3660	0.8749	0.2008	0.3018	
Stimulants ²	0.8488	0.6230	0.2105	0.2461	0.6639	0.3475	0.8005	0.7313	0.8648	0.0878	0.7581	0.3363	
Sedatives	0.7748	0.7707	0.9003	0.4135	0.8696	0.8921	0.7513	0.3925	0.8777	0.8731	0.7907	0.6233	
ALCOHOL	0.0001	0.0009	0.0000	0.0000	0.0000	0.0000	0.0000	0.0035	0.0112	0.3311	0.3071	0.5922	
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.0051	0.0571	0.0001	0.0010	0.0001	0.0034	0.0001	0.0031	0.0563	0.1244	0.3000	0.1941	
ILLCIT DRUGS OR ALCOHOL¹	0.0000	0.0019	0.0001	0.0000	0.0000	0.0000	0.0000	0.0045	0.0053	0.1276	0.0227	0.8585	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.57C Substance Dependence for Specific Substances in the Past Year among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	162	143	150	150	167	171	154	166	184	152	197	197	157
Marijuana and Hashish	88	82	89	101	97	102	99	94	135	95	113	152	100
Cocaine	95	85	98	95	108	102	80	78	81	69	96	68	63
Heroin	43	33	43	33	64	38	41	60	46	52	66	52	53
Hallucinogens	8	18	18	*	8	18	16	28	13	20	23	14	13
Inhalants	*	*	17	*	*	8	21	9	18	11	*	*	11
Nonmedical Use of Psychotherapeutics ^{2,3}	105	89	82	79	79	105	87	109	124	98	127	121	94
Pain Relievers	98	82	66	75	75	98	80	94	99	93	117	115	89
Tranquilizers	60	42	41	35	34	50	39	47	72	35	54	42	42
Stimulants ²	35	25	42	30	31	57	31	37	36	37	53	68	41
Sedatives	56	28	21	13	15	20	24	31	35	24	34	21	31
ALCOHOL	278	250	284	241	266	270	280	285	289	247	288	278	228
BOTH ILLICIT DRUGS AND ALCOHOL¹	100	63	79	79	86	88	77	90	74	87	112	78	80
ILLCIT DRUGS OR ALCOHOL¹	308	281	302	273	301	307	303	324	340	271	330	319	264

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.57D Substance Dependence for Specific Substances in the Past Year among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS¹	0.09	0.08	0.08	0.08	0.09	0.09	0.08	0.09	0.09	0.08	0.10	0.10	0.08
Marijuana and Hashish	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.07	0.05	0.06	0.07	0.05
Cocaine	0.05	0.05	0.05	0.05	0.06	0.05	0.04	0.04	0.04	0.03	0.05	0.03	0.03
Heroin	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.03	0.02	0.03	0.03	0.03	0.03
Hallucinogens	0.00	0.01	0.01	*	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Inhalants	*	*	0.01	*	*	0.00	0.01	0.00	0.01	0.01	*	*	0.01
Nonmedical Use of Psychotherapeutics ^{2,3}	0.06	0.05	0.05	0.04	0.04	0.06	0.05	0.06	0.06	0.05	0.06	0.06	0.05
Pain Relievers	0.05	0.05	0.04	0.04	0.04	0.05	0.04	0.05	0.05	0.05	0.06	0.06	0.04
Tranquilizers	0.03	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.04	0.02	0.03	0.02	0.02
Stimulants ²	0.02	0.01	0.02	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.03	0.03	0.02
Sedatives	0.03	0.02	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.02	0.01	0.02
ALCOHOL	0.15	0.14	0.15	0.13	0.14	0.14	0.15	0.15	0.15	0.12	0.14	0.14	0.11
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.06	0.03	0.04	0.04	0.05	0.05	0.04	0.05	0.04	0.04	0.06	0.04	0.04
ILLCIT DRUGS OR ALCOHOL¹	0.17	0.16	0.16	0.15	0.16	0.16	0.16	0.17	0.17	0.14	0.16	0.16	0.13

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.57P Substance Dependence for Specific Substances in the Past Year among Persons Aged 26 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Past Year Dependence	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLICIT DRUGS¹	0.0838	0.0110	0.2536	0.0283	0.3334	0.1502	0.1253	0.3193	0.1403	0.0022	0.4896	0.4040	
Marijuana and Hashish	0.0475	0.0116	0.2368	0.0853	0.2793	0.1226	0.2294	0.1440	0.6165	0.0479	0.3497	0.9231	
Cocaine	0.0018	0.0049	0.0002	0.0013	0.0003	0.0009	0.0096	0.2346	0.2885	0.8935	0.0437	0.7307	
Heroin	0.0050	0.0003	0.0073	0.0001	0.1844	0.0002	0.0027	0.1047	0.0249	0.0478	0.2749	0.2055	
Hallucinogens	0.3424	0.8105	0.5603	*	0.3804	0.8778	0.9299	0.3373	0.6398	0.4440	0.3522	0.8717	
Inhalants	*	*	0.7514	*	*	0.3615	0.3778	0.4166	0.8661	0.9696	*	*	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0025	0.0027	0.0010	0.0002	0.0001	0.0509	0.0041	0.2933	0.2212	0.0301	0.7411	0.5108	
Pain Relievers	0.0025	0.0014	0.0000	0.0008	0.0010	0.0713	0.0075	0.1950	0.0956	0.0444	0.6714	0.8485	
Tranquilizers	0.9133	0.2517	0.6470	0.2490	0.0686	0.3567	0.4828	0.8239	0.8667	0.0869	0.9832	0.2955	
Stimulants ²	0.2511	0.0716	0.8810	0.1573	0.1266	1.0000	0.0921	0.4359	0.1588	0.3824	0.5348	0.8338	
Sedatives	0.6035	0.9835	0.7514	0.2159	0.1365	0.8502	0.8610	0.8602	0.5782	0.3982	0.8054	0.2636	
ALCOHOL	0.2126	0.4712	0.3834	0.5412	0.7407	0.6867	0.1571	0.0398	0.6561	0.3705	0.1407	0.8803	
BOTH ILLICIT DRUGS AND ALCOHOL¹	0.3575	0.3284	0.3609	0.4177	0.6813	0.3166	0.8343	0.2926	0.6877	0.8535	0.3420	0.5571	
ILLICIT DRUGS OR ALCOHOL¹	0.8894	0.0769	0.9421	0.0429	0.7030	0.4442	0.7028	0.3407	0.7299	0.0123	0.5772	0.8267	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.58C Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	136	119	134	137	157	151	124	157	140	134	141	157	141
Alcohol	146	144	174	171	178	167	183	179	175	174	174	174	158
Illicit Drugs or Alcohol ^{1,2}	178	169	198	191	214	193	211	213	196	194	198	211	186
Both Illicit Drugs and Alcohol ¹	111	100	111	120	128	125	96	128	109	107	108	123	113
Illicit Drugs but not Alcohol ¹	76	60	76	69	93	76	82	80	78	76	89	81	84
Alcohol but not Illicit Drugs ¹	98	100	127	120	111	112	152	124	138	140	134	128	104
SPECIALTY FACILITY													
Illicit Drugs ¹	121	91	114	108	125	114	94	124	116	109	117	123	108
Alcohol	124	102	135	139	125	131	138	127	126	124	127	125	121
Illicit Drugs or Alcohol ^{1,2}	151	118	165	157	162	166	159	168	155	150	155	160	148
Both Illicit Drugs and Alcohol ¹	88	71	73	89	82	76	63	87	79	77	83	75	73
Illicit Drugs but not Alcohol ¹	77	51	75	56	94	74	67	76	73	72	79	78	77
Alcohol but not Illicit Drugs ¹	87	69	102	100	88	102	116	89	94	97	97	101	82

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.58D Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	0.06	0.05	0.06	0.06	0.06	0.06	0.05	0.06	0.06	0.05	0.05	0.06	0.05
Alcohol	0.06	0.06	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.06
Illicit Drugs or Alcohol ^{1,2}	0.08	0.07	0.08	0.08	0.09	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.07
Both Illicit Drugs and Alcohol ¹	0.05	0.04	0.05	0.05	0.05	0.05	0.04	0.05	0.04	0.04	0.04	0.05	0.04
Illicit Drugs but not Alcohol ¹	0.03	0.03	0.03	0.03	0.04	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03
Alcohol but not Illicit Drugs ¹	0.04	0.04	0.05	0.05	0.05	0.05	0.06	0.05	0.05	0.05	0.05	0.05	0.04
SPECIALTY FACILITY													
Illicit Drugs ¹	0.05	0.04	0.05	0.04	0.05	0.05	0.04	0.05	0.05	0.04	0.04	0.05	0.04
Alcohol	0.05	0.04	0.06	0.06	0.05	0.05	0.06	0.05	0.05	0.05	0.05	0.05	0.05
Illicit Drugs or Alcohol ^{1,2}	0.06	0.05	0.07	0.06	0.07	0.07	0.06	0.07	0.06	0.06	0.06	0.06	0.06
Both Illicit Drugs and Alcohol ¹	0.04	0.03	0.03	0.04	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03
Illicit Drugs but not Alcohol ¹	0.03	0.02	0.03	0.02	0.04	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03
Alcohol but not Illicit Drugs ¹	0.04	0.03	0.04	0.04	0.04	0.04	0.05	0.04	0.04	0.04	0.04	0.04	0.03

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.58P Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	0.4146	0.0267	0.9048	0.7236	0.3394	0.6218	0.2568	0.9532	0.4933	0.0652	0.4229	0.3401	
Alcohol	0.9004	0.8177	0.3193	0.0886	0.2340	0.2918	0.1050	0.0227	0.1926	0.8732	0.9301	0.7453	
Illicit Drugs or Alcohol ^{1,2}	0.4184	0.0993	0.9177	0.6288	0.5108	0.8532	0.5798	0.2247	0.4900	0.3134	0.8065	0.8436	
Both Illicit Drugs and Alcohol ¹	0.7223	0.8688	0.2535	0.1804	0.1356	0.6147	0.8856	0.2812	0.7381	0.2543	0.2295	0.5827	
Illicit Drugs but not Alcohol ¹	0.0592	0.0002	0.0771	0.0076	0.6798	0.1063	0.1101	0.0922	0.4862	0.0967	0.8479	0.2235	
Alcohol but not Illicit Drugs ¹	0.8359	0.8664	0.7463	0.2787	0.8587	0.3054	0.0271	0.0266	0.0392	0.2200	0.3895	0.3246	
SPECIALTY FACILITY													
Illicit Drugs ¹	0.9486	0.0120	0.8540	0.1937	0.5750	0.3678	0.0310	0.8079	0.9186	0.1773	0.7971	0.5221	
Alcohol	0.3441	0.4575	0.5287	0.3013	0.5730	0.5425	0.6415	0.2062	0.6071	0.7452	0.7740	0.4371	
Illicit Drugs or Alcohol ^{1,2}	0.8653	0.0094	0.8586	0.6894	0.5909	0.9748	0.4424	0.4911	0.6311	0.3188	0.7755	0.6008	
Both Illicit Drugs and Alcohol ¹	0.3474	0.8558	0.3211	0.2754	0.3506	0.8826	0.4900	0.3660	0.8159	0.3850	0.7345	0.2112	
Illicit Drugs but not Alcohol ¹	0.2479	0.0001	0.0948	0.0006	0.9350	0.2166	0.0215	0.1169	0.7262	0.1894	0.8499	0.5454	
Alcohol but not Illicit Drugs ¹	0.3500	0.6997	0.7861	0.3463	0.8875	0.1481	0.2204	0.3210	0.1905	0.4935	0.6813	0.5157	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.59C Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	22	20	23	20	24	20	21	19	20	23	22	17	20
Alcohol	22	20	20	19	19	18	18	18	18	18	20	16	16
Illicit Drugs or Alcohol ^{1,2}	26	24	26	24	27	24	27	23	24	26	24	20	23
Both Illicit Drugs and Alcohol ¹	20	16	18	17	18	17	16	16	15	16	19	14	15
Illicit Drugs but not Alcohol ¹	11	11	12	12	16	12	12	11	13	16	13	9	13
Alcohol but not Illicit Drugs ¹	9	10	9	8	8	7	9	7	8	7	7	6	5
SPECIALTY FACILITY													
Illicit Drugs ¹	18	14	16	15	18	13	14	13	13	17	14	12	12
Alcohol	17	12	15	14	14	12	11	13	11	11	12	13	12
Illicit Drugs or Alcohol ^{1,2}	19	17	18	17	20	15	16	16	16	18	16	15	16
Both Illicit Drugs and Alcohol ¹	16	9	12	12	12	9	9	11	8	9	10	10	6
Illicit Drugs but not Alcohol ¹	8	9	8	9	14	9	9	8	9	13	11	7	11
Alcohol but not Illicit Drugs ¹	7	6	8	6	6	5	5	5	5	5	5	4	7

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.59D Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	0.09	0.08	0.09	0.08	0.09	0.08	0.08	0.08	0.08	0.09	0.09	0.07	0.08
Alcohol	0.09	0.08	0.08	0.07	0.07	0.07	0.07	0.07	0.07	0.07	0.08	0.06	0.06
Illicit Drugs or Alcohol ^{1,2}	0.11	0.10	0.10	0.09	0.11	0.10	0.11	0.09	0.10	0.11	0.10	0.08	0.09
Both Illicit Drugs and Alcohol ¹	0.08	0.06	0.07	0.07	0.07	0.07	0.07	0.07	0.06	0.07	0.07	0.06	0.06
Illicit Drugs but not Alcohol ¹	0.04	0.04	0.05	0.05	0.06	0.05	0.05	0.04	0.05	0.06	0.05	0.04	0.05
Alcohol but not Illicit Drugs ¹	0.04	0.04	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.02	0.02
SPECIALTY FACILITY													
Illicit Drugs ¹	0.07	0.05	0.06	0.06	0.07	0.05	0.06	0.05	0.06	0.07	0.06	0.05	0.05
Alcohol	0.07	0.05	0.06	0.05	0.06	0.05	0.04	0.05	0.05	0.04	0.05	0.05	0.05
Illicit Drugs or Alcohol ^{1,2}	0.08	0.07	0.07	0.07	0.08	0.06	0.06	0.06	0.07	0.07	0.07	0.06	0.06
Both Illicit Drugs and Alcohol ¹	0.06	0.04	0.05	0.05	0.05	0.04	0.04	0.04	0.03	0.04	0.04	0.04	0.03
Illicit Drugs but not Alcohol ¹	0.03	0.04	0.03	0.04	0.05	0.03	0.04	0.03	0.04	0.05	0.05	0.03	0.04
Alcohol but not Illicit Drugs ¹	0.03	0.03	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.59P Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	0.0030	0.0080	0.0004	0.0045	0.0050	0.0341	0.0607	0.1076	0.1375	0.0443	0.0577	0.7487	
Alcohol	0.0000	0.0000	0.0000	0.0000	0.0004	0.0007	0.0012	0.0020	0.0691	0.1279	0.0260	0.1741	
Illicit Drugs or Alcohol ^{1,2}	0.0000	0.0001	0.0000	0.0005	0.0007	0.0011	0.0031	0.0531	0.0376	0.0359	0.0846	0.8573	
Both Illicit Drugs and Alcohol ¹	0.0011	0.0017	0.0002	0.0013	0.0116	0.0078	0.0153	0.0200	0.2893	0.3336	0.0682	0.2953	
Illicit Drugs but not Alcohol ¹	0.6335	0.7554	0.3037	0.6875	0.1612	0.9688	0.9845	0.6984	0.3223	0.0724	0.4794	0.3731	
Alcohol but not Illicit Drugs ¹	0.0002	0.0008	0.0001	0.0012	0.0015	0.0108	0.0114	0.0131	0.0372	0.0787	0.1281	0.2553	
SPECIALTY FACILITY													
Illicit Drugs ¹	0.0017	0.0385	0.0034	0.0008	0.0060	0.0508	0.0435	0.0162	0.1511	0.0167	0.0142	0.3555	
Alcohol	0.0015	0.0170	0.0002	0.0007	0.0165	0.1018	0.1624	0.0162	0.3110	0.6302	0.2136	0.2913	
Illicit Drugs or Alcohol ^{1,2}	0.0021	0.0128	0.0019	0.0031	0.0068	0.0723	0.1090	0.0460	0.1492	0.1219	0.0376	0.5349	
Both Illicit Drugs and Alcohol ¹	0.0009	0.0074	0.0001	0.0000	0.0061	0.0155	0.0099	0.0007	0.1473	0.1162	0.0241	0.0433	
Illicit Drugs but not Alcohol ¹	0.4969	0.7540	0.9679	0.6705	0.1768	0.8782	0.7302	0.6951	0.6596	0.1003	0.1998	0.4350	
Alcohol but not Illicit Drugs ¹	0.0699	0.1912	0.0657	0.3547	0.3077	0.6818	0.8802	0.7895	0.8944	0.8184	0.9286	0.5471	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.60C Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	31	37	37	35	40	35	36	39	40	43	45	37	42
Alcohol	30	35	39	37	42	39	39	42	44	41	32	36	36
Illicit Drugs or Alcohol ^{1,2}	38	45	48	47	49	45	46	49	52	50	52	46	52
Both Illicit Drugs and Alcohol ¹	23	27	29	26	34	26	28	32	31	34	23	31	28
Illicit Drugs but not Alcohol ¹	19	25	23	25	19	23	23	24	26	25	35	21	30
Alcohol but not Illicit Drugs ¹	20	24	26	26	24	29	25	27	30	22	23	19	21
SPECIALTY FACILITY													
Illicit Drugs ¹	25	29	28	29	33	29	26	32	29	36	34	31	34
Alcohol	24	26	28	25	31	25	28	29	29	32	19	26	27
Illicit Drugs or Alcohol ^{1,2}	30	33	36	34	38	33	33	39	37	40	37	35	41
Both Illicit Drugs and Alcohol ¹	18	22	19	17	27	19	19	19	19	26	13	22	20
Illicit Drugs but not Alcohol ¹	16	20	20	22	17	20	18	22	22	24	31	19	26
Alcohol but not Illicit Drugs ¹	16	15	20	18	16	16	19	18	21	18	13	13	18

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.60D Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	0.10	0.12	0.12	0.11	0.12	0.11	0.11	0.12	0.12	0.12	0.13	0.11	0.12
Alcohol	0.10	0.11	0.12	0.11	0.13	0.12	0.12	0.13	0.13	0.12	0.09	0.10	0.10
Illicit Drugs or Alcohol ^{1,2}	0.12	0.14	0.15	0.15	0.15	0.14	0.14	0.15	0.15	0.15	0.15	0.13	0.15
Both Illicit Drugs and Alcohol ¹	0.08	0.09	0.09	0.08	0.10	0.08	0.09	0.10	0.09	0.10	0.07	0.09	0.08
Illicit Drugs but not Alcohol ¹	0.06	0.08	0.07	0.08	0.06	0.07	0.07	0.07	0.08	0.07	0.10	0.06	0.08
Alcohol but not Illicit Drugs ¹	0.06	0.08	0.08	0.08	0.07	0.09	0.07	0.08	0.09	0.06	0.07	0.05	0.06
SPECIALTY FACILITY													
Illicit Drugs ¹	0.08	0.09	0.09	0.09	0.10	0.09	0.08	0.09	0.08	0.10	0.10	0.09	0.10
Alcohol	0.08	0.08	0.09	0.08	0.10	0.08	0.08	0.09	0.09	0.09	0.05	0.07	0.08
Illicit Drugs or Alcohol ^{1,2}	0.10	0.10	0.11	0.10	0.12	0.10	0.10	0.12	0.11	0.12	0.11	0.10	0.12
Both Illicit Drugs and Alcohol ¹	0.06	0.07	0.06	0.05	0.08	0.06	0.06	0.06	0.05	0.08	0.04	0.06	0.06
Illicit Drugs but not Alcohol ¹	0.05	0.06	0.06	0.07	0.05	0.06	0.05	0.07	0.06	0.07	0.09	0.06	0.07
Alcohol but not Illicit Drugs ¹	0.05	0.05	0.06	0.06	0.05	0.05	0.06	0.05	0.06	0.05	0.04	0.04	0.05

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.60P Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 18 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	0.6439	0.0849	0.1126	0.0532	0.1174	0.8051	0.3962	0.0316	0.1335	0.1178	0.3276	0.7867	
Alcohol	0.0862	0.0000	0.0004	0.0001	0.0000	0.0000	0.0001	0.0000	0.0004	0.0196	0.8383	0.4007	
Illicit Drugs or Alcohol ^{1,2}	0.9005	0.0026	0.0119	0.0002	0.0208	0.0247	0.0340	0.0004	0.0020	0.0889	0.2507	0.9703	
Both Illicit Drugs and Alcohol ¹	0.2499	0.0006	0.0175	0.0155	0.0008	0.0509	0.0153	0.0004	0.0587	0.0519	0.5780	0.1450	
Illicit Drugs but not Alcohol ¹	0.0495	0.2873	0.8516	0.7481	0.0820	0.0950	0.1644	0.4621	0.8543	0.9693	0.0792	0.1972	
Alcohol but not Illicit Drugs ¹	0.1872	0.0081	0.0059	0.0017	0.0103	0.0000	0.0004	0.0007	0.0009	0.1803	0.7617	0.5200	
SPECIALTY FACILITY													
Illicit Drugs ¹	0.9095	0.3197	0.2428	0.2936	0.4568	0.9881	0.9247	0.1792	0.4215	0.1065	0.3985	0.7569	
Alcohol	0.1180	0.0208	0.0098	0.0616	0.0297	0.0782	0.0026	0.0026	0.0621	0.0732	0.0616	0.7881	
Illicit Drugs or Alcohol ^{1,2}	0.7031	0.2349	0.0287	0.0972	0.2307	0.3419	0.2428	0.0155	0.1512	0.0658	0.9829	0.8445	
Both Illicit Drugs and Alcohol ¹	0.1143	0.0181	0.1161	0.3277	0.0319	0.3600	0.1078	0.0557	0.3222	0.1301	0.2086	0.5916	
Illicit Drugs but not Alcohol ¹	0.0818	0.2014	0.8150	0.9256	0.1363	0.2568	0.0663	0.6169	0.9399	0.5724	0.1611	0.4399	
Alcohol but not Illicit Drugs ¹	0.3974	0.5776	0.0462	0.0559	0.5615	0.0844	0.0118	0.0658	0.1629	0.2348	0.1628	0.3132	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.61C Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	130	108	124	129	152	145	115	152	131	122	132	149	131
Alcohol	139	136	168	162	172	161	178	172	169	169	170	168	151
Illicit Drugs or Alcohol ^{1,2}	169	160	189	180	208	186	201	205	187	184	191	204	172
Both Illicit Drugs and Alcohol ¹	106	93	104	113	123	122	89	124	102	100	104	118	108
Illicit Drugs but not Alcohol ¹	76	52	73	63	92	71	77	76	72	69	81	75	74
Alcohol but not Illicit Drugs ¹	96	96	124	116	108	107	151	121	135	137	131	126	101
SPECIALTY FACILITY													
Illicit Drugs ¹	116	84	107	101	120	107	88	120	111	100	113	115	100
Alcohol	121	96	131	135	121	127	132	124	121	120	127	121	115
Illicit Drugs or Alcohol ^{1,2}	145	110	157	150	158	160	151	163	148	141	153	153	139
Both Illicit Drugs and Alcohol ¹	84	67	69	86	78	73	59	85	77	71	83	70	69
Illicit Drugs but not Alcohol ¹	77	45	72	48	92	68	63	72	69	66	72	74	68
Alcohol but not Illicit Drugs ¹	86	67	99	98	86	99	114	88	91	95	97	100	79

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.61D Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	0.07	0.06	0.07	0.07	0.08	0.08	0.06	0.08	0.07	0.06	0.07	0.07	0.06
Alcohol	0.08	0.08	0.09	0.09	0.09	0.08	0.09	0.09	0.09	0.09	0.08	0.08	0.07
Illicit Drugs or Alcohol ^{1,2}	0.09	0.09	0.10	0.10	0.11	0.10	0.10	0.11	0.10	0.09	0.10	0.10	0.08
Both Illicit Drugs and Alcohol ¹	0.06	0.05	0.06	0.06	0.07	0.06	0.05	0.06	0.05	0.05	0.05	0.06	0.05
Illicit Drugs but not Alcohol ¹	0.04	0.03	0.04	0.03	0.05	0.04	0.04	0.04	0.04	0.03	0.04	0.04	0.04
Alcohol but not Illicit Drugs ¹	0.05	0.05	0.07	0.06	0.06	0.06	0.08	0.06	0.07	0.07	0.07	0.06	0.05
SPECIALTY FACILITY													
Illicit Drugs ¹	0.06	0.05	0.06	0.05	0.06	0.06	0.05	0.06	0.06	0.05	0.06	0.06	0.05
Alcohol	0.07	0.05	0.07	0.07	0.06	0.07	0.07	0.06	0.06	0.06	0.06	0.06	0.06
Illicit Drugs or Alcohol ^{1,2}	0.08	0.06	0.09	0.08	0.08	0.08	0.08	0.08	0.08	0.07	0.08	0.08	0.07
Both Illicit Drugs and Alcohol ¹	0.05	0.04	0.04	0.05	0.04	0.04	0.03	0.04	0.04	0.04	0.04	0.03	0.03
Illicit Drugs but not Alcohol ¹	0.04	0.02	0.04	0.03	0.05	0.04	0.03	0.04	0.04	0.03	0.04	0.04	0.03
Alcohol but not Illicit Drugs ¹	0.05	0.04	0.05	0.05	0.05	0.05	0.06	0.05	0.05	0.05	0.05	0.05	0.04

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.61P Received Substance Use Treatment at Any Treatment Location or at a Specialty Facility in the Past Year among Persons Aged 26 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Location/Substance for Which Treatment Was Received in Past Year	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ANY TREATMENT LOCATION													
Illicit Drugs ¹	0.2191	0.0005	0.2130	0.1554	0.9079	0.3684	0.0708	0.3612	0.1441	0.0038	0.1335	0.2541	
Alcohol	0.4216	0.0648	0.7433	0.6570	0.8989	0.7642	0.6464	0.3652	0.7868	0.5853	0.7745	0.9853	
Illicit Drugs or Alcohol ^{1,2}	0.1446	0.0014	0.2100	0.3325	0.7401	0.4169	0.7389	0.9147	0.6605	0.0647	0.4182	0.8124	
Both Illicit Drugs and Alcohol ¹	0.6582	0.0978	0.9792	0.7272	0.7748	0.7609	0.2140	0.9087	0.2900	0.0508	0.1575	0.2736	
Illicit Drugs but not Alcohol ¹	0.1444	0.0001	0.0379	0.0013	0.8509	0.2346	0.2035	0.1264	0.2962	0.0228	0.5073	0.4416	
Alcohol but not Illicit Drugs ¹	0.4839	0.3157	0.6594	0.7757	0.6032	0.9103	0.1341	0.1529	0.2023	0.3422	0.4485	0.2839	
SPECIALTY FACILITY													
Illicit Drugs ¹	0.6471	0.0008	0.3497	0.0296	1.0000	0.2393	0.0107	0.3479	0.5942	0.0172	0.3930	0.3745	
Alcohol	0.7692	0.1094	0.7694	0.7051	0.8573	0.8911	0.7797	0.7159	0.9974	0.4074	0.9686	0.3941	
Illicit Drugs or Alcohol ^{1,2}	0.8042	0.0006	0.2847	0.2494	0.9648	0.6801	0.2099	0.9099	0.9645	0.0762	0.5983	0.5729	
Both Illicit Drugs and Alcohol ¹	0.9155	0.2270	0.9581	0.6903	0.9634	0.5111	0.1265	0.9512	0.5194	0.1081	0.7480	0.0862	
Illicit Drugs but not Alcohol ¹	0.4339	0.0001	0.0845	0.0001	0.7742	0.3481	0.0524	0.1356	0.6262	0.0487	0.3222	0.7519	
Alcohol but not Illicit Drugs ¹	0.4924	0.5543	0.7998	0.5741	0.9940	0.2405	0.4103	0.5332	0.2997	0.6236	0.5027	0.3787	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s); therefore, the substance-specific estimates may not sum to this combined row.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.62C Substances for Which Last or Current Treatment Was Received among Persons Aged 12 or Older Who Received Substance Use Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014

Substance for Which Last or Current Treatment Was Received¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	179	169	202	192	216	192	210	211	199	196	200	210	188
Marijuana	91	71	78	80	105	91	74	102	89	76	84	72	87
Cocaine	95	72	89	89	112	103	67	91	84	70	81	77	90
Heroin	55	50	57	63	78	73	55	78	73	72	63	69	75
Hallucinogens	42	38	48	57	73	46	42	65	57	50	59	48	53
Inhalants	29	26	39	33	54	39	34	51	53	39	34	39	39
Pain Relievers	59	51	46	55	64	70	61	84	83	78	99	85	72
Tranquilizers	36	43	45	43	74	43	50	66	61	47	68	57	65
Stimulants ²	39	51	56	55	85	43	51	78	59	52	56	73	60
Sedatives	60	27	28	24	58	17	32	50	51	32	37	41	38
Alcohol	143	137	172	157	176	156	179	176	173	166	171	168	150

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.62D Substances for Which Last or Current Treatment Was Received among Persons Aged 12 or Older Who Received Substance Use Treatment in the Past Year: Standard Errors of Percentages, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Marijuana	2.20	1.85	2.03	1.88	2.10	1.96	1.75	2.01	1.89	1.89	1.94	1.68	1.79
Cocaine	2.30	1.95	2.03	2.01	2.40	2.21	1.59	1.89	1.78	1.73	1.82	1.78	1.89
Heroin	1.51	1.40	1.44	1.53	1.77	1.74	1.30	1.68	1.65	1.77	1.47	1.57	1.64
Hallucinogens	1.17	1.08	1.23	1.38	1.70	1.12	1.00	1.40	1.30	1.24	1.40	1.15	1.19
Inhalants	0.84	0.77	1.01	0.84	1.28	0.95	0.82	1.13	1.24	1.00	0.83	0.94	0.91
Pain Relievers	1.60	1.47	1.24	1.35	1.51	1.66	1.43	1.72	1.87	1.93	2.18	1.90	1.58
Tranquilizers	1.02	1.26	1.18	1.07	1.66	1.06	1.18	1.46	1.41	1.20	1.58	1.35	1.48
Stimulants ²	1.12	1.46	1.42	1.38	1.86	1.05	1.23	1.66	1.36	1.32	1.32	1.66	1.37
Sedatives	1.65	0.79	0.70	0.61	1.34	0.44	0.78	1.13	1.21	0.84	0.91	1.00	0.89
Alcohol	2.35	2.28	2.41	2.19	2.46	2.30	2.25	2.16	2.21	2.32	2.49	2.29	2.19

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.62P Substances for Which Last or Current Treatment Was Received among Persons Aged 12 or Older Who Received Substance Use Treatment in the Past Year: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	
Marijuana	0.2848	0.0895	0.4803	0.4198	0.0404	0.7265	0.5709	0.1443	0.9513	0.4995	0.6951	0.0974	
Cocaine	0.1768	0.4455	0.1040	0.5956	0.1635	0.4896	0.3165	0.8113	0.4860	0.0407	0.3631	0.0896	
Heroin	0.0020	0.0028	0.0007	0.0034	0.1721	0.0104	0.0022	0.1337	0.0430	0.1505	0.0985	0.4008	
Hallucinogens	0.8646	0.9012	0.5528	0.6043	0.1720	0.8192	0.5165	0.2247	0.9568	0.8214	0.5923	0.6813	
Inhalants	0.6409	0.6718	0.4368	0.7983	0.2645	0.8763	0.9815	0.3913	0.6064	0.7069	0.5458	0.9624	
Pain Relievers	0.0003	0.0046	0.0002	0.0012	0.0216	0.0667	0.0801	0.5532	0.9071	0.7823	0.0344	0.9362	
Tranquilizers	0.0011	0.0415	0.0244	0.0144	0.3783	0.0100	0.0649	0.3620	0.1363	0.1045	0.9532	0.2551	
Stimulants ²	0.1533	0.9670	0.9510	0.5022	0.1848	0.1951	0.3002	0.3901	0.3182	0.2843	0.4852	0.5968	
Sedatives	0.5891	0.7035	0.6068	0.2794	0.5195	0.0347	0.6964	0.6505	0.5630	0.6990	0.8869	0.9218	
Alcohol	0.0723	0.0265	0.0833	0.1305	0.1198	0.1199	0.0117	0.0024	0.1228	0.1313	0.5804	0.2035	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.63C Substances for Which Last or Current Treatment Was Received among Persons Aged 12 to 17 Who Received Substance Use Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	26	24	27	24	27	25	27	23	24	27	24	20	23
Marijuana	20	17	21	17	20	18	17	17	17	19	21	14	16
Cocaine	11	7	9	7	11	7	6	7	8	5	8	7	3
Heroin	3	3	4	4	2	3	4	3	2	6	5	*	5
Hallucinogens	12	8	7	8	10	7	8	8	7	9	9	7	4
Inhalants	9	7	6	6	6	4	5	7	9	4	6	4	4
Pain Relievers	10	9	11	9	12	8	8	9	10	10	9	7	6
Tranquilizers	10	4	6	6	10	7	5	6	7	8	12	5	7
Stimulants ²	10	6	8	6	10	6	5	5	7	7	5	6	7
Sedatives	4	5	4	3	4	3	1	2	*	4	4	*	6
Alcohol	19	17	19	17	17	17	16	16	16	13	17	14	13

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.63D Substances for Which Last or Current Treatment Was Received among Persons Aged 12 to 17 Who Received Substance Use Treatment in the Past Year: Standard Errors of Percentages, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Marijuana	3.22	3.55	3.46	3.34	3.70	3.66	3.93	3.78	4.02	4.02	3.86	4.25	5.06
Cocaine	2.64	1.74	2.22	1.89	2.77	2.03	1.79	2.39	2.52	1.71	2.69	2.76	1.47
Heroin	0.83	0.84	1.08	1.17	0.51	0.99	1.25	1.00	0.65	1.91	1.84	*	2.31
Hallucinogens	2.97	2.13	1.70	2.16	2.71	1.97	2.27	2.72	2.39	2.87	2.87	2.82	1.72
Inhalants	2.18	1.84	1.46	1.69	1.76	1.04	1.56	2.23	2.85	1.33	2.10	1.79	1.73
Pain Relievers	2.53	2.21	2.46	2.37	2.99	2.30	2.36	2.94	3.05	3.19	3.08	2.77	2.59
Tranquilizers	2.51	1.12	1.48	1.69	2.58	1.97	1.44	1.98	2.14	2.60	3.70	2.11	3.03
Stimulants ²	2.50	1.63	1.83	1.61	2.67	1.81	1.37	1.79	2.22	2.34	1.81	2.34	2.82
Sedatives	1.05	1.22	1.04	0.88	1.01	0.93	0.45	0.82	*	1.16	1.37	*	2.47
Alcohol	3.44	3.48	3.35	3.57	3.61	3.62	3.71	3.90	4.09	3.87	4.36	4.34	4.58

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.63P Substances for Which Last or Current Treatment Was Received among Persons Aged 12 to 17 Who Received Substance Use Treatment in the Past Year: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	
Marijuana	0.5446	0.9186	0.6757	0.7950	0.7234	0.8649	0.6815	0.5241	0.7783	0.6040	0.0278	0.6385	
Cocaine	0.0046	0.0159	0.0148	0.0170	0.0014	0.0092	0.1449	0.0245	0.0443	0.3151	0.1027	0.1110	
Heroin	0.1390	0.1998	0.2832	0.3789	0.0929	0.2661	0.3524	0.2425	0.0814	0.8505	0.4476	*	
Hallucinogens	0.0617	0.0773	0.2600	0.0565	0.0216	0.2511	0.1280	0.0459	0.1640	0.0386	0.1140	0.1521	
Inhalants	0.3861	0.2581	0.2951	0.2986	0.6827	0.5891	0.7973	0.1229	0.0241	0.7456	0.4596	0.8268	
Pain Relievers	0.9592	0.7315	0.1802	0.2877	0.1099	0.5235	0.3233	0.1503	0.0717	0.0819	0.1581	0.4958	
Tranquilizers	0.4360	0.0649	0.3620	0.2818	0.9043	0.3863	0.2208	0.4999	0.6005	0.9141	0.2865	0.5890	
Stimulants ²	0.8431	0.5131	0.9196	0.5535	0.3017	0.5688	0.1631	0.5787	0.8124	0.9513	0.4321	0.8995	
Sedatives	0.4623	0.4475	0.6048	0.2835	0.4935	0.2452	0.1192	0.2453	*	0.4725	0.5483	*	
Alcohol	0.0058	0.0323	0.0174	0.0147	0.2162	0.0307	0.1325	0.0425	0.2416	0.9149	0.2066	0.1118	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.64C Substances for Which Last or Current Treatment Was Received among Persons Aged 18 to 25 Who Received Substance Use Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	38	46	49	48	50	46	47	50	52	51	54	47	52
Marijuana	24	28	29	27	36	28	29	31	29	29	31	26	29
Cocaine	15	17	19	17	21	18	19	18	18	19	20	21	19
Heroin	8	11	10	15	13	14	13	15	13	18	17	19	22
Hallucinogens	13	15	19	14	23	18	15	20	13	16	16	18	17
Inhalants	7	9	12	11	15	14	14	16	8	11	10	14	13
Pain Relievers	11	17	19	17	21	18	21	25	21	25	31	22	26
Tranquilizers	11	13	14	13	23	15	14	16	13	19	16	17	21
Stimulants ²	11	20	19	15	25	17	15	18	13	17	25	17	20
Sedatives	6	6	6	7	8	7	8	11	6	4	4	10	9
Alcohol	28	34	35	35	40	39	37	40	39	38	29	33	34

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.64D Substances for Which Last or Current Treatment Was Received among Persons Aged 18 to 25 Who Received Substance Use Treatment in the Past Year: Standard Errors of Percentages, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Marijuana	2.72	2.48	2.54	2.36	2.94	2.55	2.62	2.45	2.47	2.74	2.87	2.73	3.07
Cocaine	1.99	1.76	1.96	1.69	2.08	1.90	2.00	1.72	1.68	2.05	2.14	2.46	2.35
Heroin	1.16	1.17	1.09	1.49	1.42	1.48	1.40	1.48	1.35	1.89	1.85	2.17	2.49
Hallucinogens	1.79	1.60	1.96	1.40	2.35	1.83	1.65	1.91	1.32	1.76	1.65	2.16	2.06
Inhalants	1.04	0.99	1.30	1.07	1.56	1.51	1.51	1.54	0.83	1.21	1.11	1.79	1.60
Pain Relievers	1.59	1.80	1.96	1.68	2.09	1.84	2.18	2.25	1.99	2.51	2.93	2.41	2.81
Tranquilizers	1.48	1.43	1.50	1.33	2.32	1.60	1.50	1.61	1.28	2.06	1.78	2.00	2.52
Stimulants ²	1.59	2.10	1.97	1.52	2.51	1.77	1.61	1.74	1.30	1.78	2.57	1.99	2.47
Sedatives	0.87	0.71	0.69	0.77	0.85	0.75	0.96	1.13	0.64	0.49	0.51	1.21	1.14
Alcohol	2.71	2.70	2.61	2.47	2.54	2.58	2.55	2.49	2.77	2.76	2.95	2.87	3.15

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.64P Substances for Which Last or Current Treatment Was Received among Persons Aged 18 to 25 Who Received Substance Use Treatment in the Past Year: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	
Marijuana	0.0736	0.0269	0.0340	0.3919	0.0044	0.7464	0.2442	0.1573	0.3849	0.4743	0.6401	0.3426	
Cocaine	0.0926	0.1675	0.0233	0.2799	0.0153	0.1883	0.0926	0.3142	0.2481	0.4404	0.7307	0.2719	
Heroin	0.0000	0.0000	0.0001	0.0041	0.0006	0.0022	0.0007	0.0071	0.0025	0.1468	0.1032	0.8388	
Hallucinogens	0.4929	0.3334	0.1723	0.9913	0.2355	0.4367	0.4694	0.2390	0.7420	0.7310	0.8443	0.4216	
Inhalants	0.8994	0.8954	0.2126	0.8270	0.2910	0.2908	0.2608	0.3649	0.4231	0.8968	0.9919	0.5423	
Pain Relievers	0.0001	0.0096	0.0225	0.0033	0.1040	0.0094	0.2565	0.4260	0.1433	0.5437	0.1814	0.7925	
Tranquilizers	0.1510	0.1719	0.1843	0.1081	0.8698	0.1926	0.2141	0.3632	0.0957	0.9576	0.9515	0.7960	
Stimulants ²	0.1731	0.5657	0.8743	0.3837	0.7336	0.7901	0.2087	0.4735	0.0824	0.3189	0.8366	0.8880	
Sedatives	0.9128	0.8530	0.6632	0.7374	0.7446	0.8511	0.6761	0.8855	0.4927	0.3658	0.3430	0.7198	
Alcohol	0.0860	0.0227	0.0841	0.1377	0.0005	0.0000	0.0003	0.0017	0.1430	0.2455	0.1199	0.5146	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.65C Substances for Which Last or Current Treatment Was Received among Persons Aged 26 or Older Who Received Substance Use Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014

Substance for Which Last or Current Treatment Was Received¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	170	159	191	181	210	184	199	203	190	186	192	203	173
Marijuana	84	62	68	72	98	84	65	97	83	66	76	63	79
Cocaine	92	70	87	87	110	101	63	89	81	66	78	73	87
Heroin	54	49	54	61	77	71	53	76	72	69	60	66	70
Hallucinogens	37	33	43	54	68	42	37	61	56	45	55	42	51
Inhalants	26	24	38	31	51	37	32	48	52	36	32	35	38
Pain Relievers	57	47	40	51	60	68	57	80	80	69	93	82	64
Tranquilizers	32	41	40	40	70	40	48	64	59	41	65	53	61
Stimulants ²	35	46	52	53	81	40	49	76	58	47	48	70	56
Sedatives	60	26	27	23	57	16	31	49	51	32	37	39	36
Alcohol	138	131	166	150	170	149	174	170	167	162	167	164	144

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.65D Substances for Which Last or Current Treatment Was Received among Persons Aged 26 or Older Who Received Substance Use Treatment in the Past Year: Standard Errors of Percentages, 2002-2014

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Marijuana	3.04	2.61	2.58	2.50	2.83	2.66	2.16	2.74	2.56	2.38	2.43	1.96	2.17
Cocaine	3.20	2.99	3.05	2.92	3.37	3.05	2.15	2.63	2.48	2.36	2.42	2.24	2.40
Heroin	2.12	2.18	2.08	2.24	2.49	2.42	1.80	2.30	2.32	2.46	1.96	2.01	2.02
Hallucinogens	1.50	1.51	1.70	1.98	2.29	1.48	1.27	1.90	1.83	1.64	1.85	1.36	1.50
Inhalants	1.07	1.11	1.48	1.16	1.76	1.29	1.08	1.52	1.73	1.34	1.09	1.11	1.16
Pain Relievers	2.22	2.17	1.64	1.90	2.03	2.33	1.90	2.34	2.60	2.49	2.84	2.43	1.90
Tranquilizers	1.28	1.88	1.60	1.50	2.25	1.43	1.60	2.00	1.97	1.51	2.12	1.68	1.84
Stimulants ²	1.42	2.08	2.01	1.98	2.52	1.40	1.65	2.28	1.91	1.75	1.60	2.12	1.69
Sedatives	2.34	1.22	1.02	0.87	1.90	0.57	1.06	1.56	1.71	1.19	1.25	1.27	1.12
Alcohol	3.33	3.34	3.44	3.06	3.35	3.13	3.06	2.95	2.89	3.09	3.16	2.85	2.71

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.65P Substances for Which Last or Current Treatment Was Received among Persons Aged 26 or Older Who Received Substance Use Treatment in the Past Year: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance for Which Last or Current Treatment Was Received ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	
Marijuana	0.8623	0.8277	0.2960	0.8369	0.5345	0.3073	0.1299	0.5241	0.3899	0.0770	0.1740	0.0208	
Cocaine	0.2587	0.4211	0.1569	0.5716	0.3546	0.5736	0.1758	0.6764	0.3555	0.0296	0.3249	0.0321	
Heroin	0.0649	0.1587	0.0253	0.0385	0.7935	0.0881	0.0367	0.4790	0.2626	0.3050	0.2504	0.4505	
Hallucinogens	0.4032	0.4544	0.9368	0.7798	0.4596	0.4592	0.2090	0.5541	0.8914	0.4469	0.7705	0.3583	
Inhalants	0.4733	0.4502	0.8721	0.5549	0.4348	0.8568	0.6229	0.7188	0.6802	0.6712	0.4386	0.8392	
Pain Relievers	0.0080	0.0259	0.0003	0.0075	0.0266	0.2434	0.1006	0.5461	0.9218	0.8200	0.0924	0.8187	
Tranquilizers	0.0026	0.1565	0.0597	0.0518	0.3707	0.0264	0.1628	0.5445	0.3196	0.0567	0.7858	0.2161	
Stimulants ²	0.2423	0.8272	0.9532	0.7064	0.2561	0.1972	0.6256	0.2381	0.5791	0.3936	0.4193	0.5613	
Sedatives	0.5001	0.9124	0.7698	0.4161	0.4712	0.0441	0.8072	0.5332	0.3711	0.9838	0.8658	0.9120	
Alcohol	0.1575	0.0357	0.1173	0.1827	0.3666	0.6055	0.0745	0.0203	0.1592	0.0770	0.2512	0.3031	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If respondents reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

² Estimates for stimulants do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.66C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	179	169	202	192	216	192	210	211	199	196	200	210	188
Hospital - Inpatient	103	66	88	80	97	100	81	88	85	90	101	93	85
Rehabilitation Facility - Inpatient	112	80	88	111	95	109	75	115	96	102	101	107	85
Rehabilitation Facility - Outpatient	124	94	137	132	124	139	115	149	127	125	116	134	118
Mental Health Center - Outpatient	108	68	111	116	104	93	108	109	93	94	95	109	94
Emergency Room	68	35	63	55	56	79	55	62	72	76	88	74	60
Private Doctor's Office	81	45	61	58	74	68	81	82	76	83	93	86	78
Self-Help Group	137	131	157	144	167	148	161	164	166	154	148	172	136
Prison/Jail	39	37	50	54	67	53	56	60	46	60	65	42	51

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.66D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Percentages, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Hospital - Inpatient	2.47	1.79	2.04	1.73	2.12	2.22	1.85	1.89	1.90	2.13	2.18	2.04	1.80
Rehabilitation Facility - Inpatient	2.54	2.08	1.94	2.33	2.13	2.27	1.78	2.23	2.07	2.32	2.17	2.29	1.87
Rehabilitation Facility - Outpatient	2.59	2.34	2.41	2.45	2.49	2.44	2.33	2.48	2.50	2.48	2.33	2.47	2.01
Mental Health Center - Outpatient	2.47	1.80	2.34	2.44	2.09	2.05	2.25	2.16	1.97	2.07	2.09	2.29	1.88
Emergency Room	1.81	1.05	1.48	1.34	1.36	1.79	1.28	1.39	1.61	1.80	2.02	1.69	1.37
Private Doctor's Office	2.06	1.30	1.53	1.43	1.73	1.58	1.79	1.74	1.70	1.93	2.08	1.91	1.65
Self-Help Group	2.52	2.36	2.57	2.43	2.76	2.50	2.54	2.45	2.43	2.30	2.41	2.44	2.16
Prison/Jail	1.11	1.07	1.27	1.34	1.59	1.30	1.32	1.33	1.05	1.50	1.54	1.03	1.17

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.66P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Hospital - Inpatient	0.4231	0.0731	0.5158	0.3073	0.4766	0.4346	0.0352	0.3951	0.0890	0.7340	0.8071	0.8500	
Rehabilitation Facility - Inpatient	0.0815	0.2373	0.7297	0.4431	0.3310	0.7514	0.0033	0.4950	0.4652	0.8205	0.8091	0.9439	
Rehabilitation Facility - Outpatient	0.4714	0.1557	0.4814	0.4512	0.6091	0.4988	0.0602	0.1276	0.8221	0.7389	0.1782	0.6308	
Mental Health Center - Outpatient	0.6797	0.0218	0.5065	0.6755	0.8796	0.0609	0.5161	0.4125	0.1695	0.8604	0.3023	0.7013	
Emergency Room	0.6882	0.0038	0.7620	0.2104	0.1532	0.7182	0.0758	0.5895	0.5760	0.2363	0.3349	0.2882	
Private Doctor's Office	0.1512	0.0004	0.0092	0.0012	0.1350	0.1209	0.4443	0.4179	0.2042	0.9388	0.8678	0.9365	
Self-Help Group	0.1510	0.1932	0.5358	0.9464	0.7988	0.4607	0.7921	0.2629	0.3290	0.2593	0.9137	0.3092	
Prison/Jail	0.3929	0.0992	0.7151	0.9730	0.4070	0.5157	0.8410	0.9618	0.7352	0.1494	0.6503	0.1357	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.67C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 to 17 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	26	24	27	24	27	25	27	23	24	27	24	20	23
Hospital - Inpatient	12	9	10	11	13	10	9	9	9	10	7	9	12
Rehabilitation Facility - Inpatient	15	11	12	10	11	11	9	10	10	11	10	9	10
Rehabilitation Facility - Outpatient	16	13	13	14	15	13	12	12	9	11	16	13	11
Mental Health Center - Outpatient	13	11	13	11	16	9	11	9	9	13	9	11	11
Emergency Room	6	10	8	9	12	9	8	8	8	9	6	9	12
Private Doctor's Office	9	9	11	8	11	11	8	7	9	8	7	7	11
Self-Help Group	20	14	15	14	17	17	12	12	13	16	17	12	13
Prison/Jail	8	7	7	5	10	5	6	7	9	6	8	6	8

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.67D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 to 17 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Percentages, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Hospital - Inpatient	2.76	2.48	2.17	2.86	3.18	2.75	2.52	2.93	2.89	2.87	2.30	3.54	4.39
Rehabilitation Facility - Inpatient	3.48	2.76	2.51	2.68	2.83	2.94	2.64	3.11	3.05	3.41	3.12	3.56	4.09
Rehabilitation Facility - Outpatient	3.21	3.03	2.76	3.26	3.58	3.36	3.17	3.35	2.98	3.24	4.27	4.29	4.35
Mental Health Center - Outpatient	3.01	2.58	2.71	2.78	3.73	2.52	2.91	2.85	2.92	3.89	2.90	4.04	4.34
Emergency Room	1.64	2.51	1.91	2.33	3.06	2.44	2.16	2.68	2.59	2.66	2.13	3.42	4.41
Private Doctor's Office	2.22	2.32	2.61	2.15	2.85	2.95	2.26	2.37	2.77	2.54	2.41	2.86	4.14
Self-Help Group	3.60	3.32	3.17	3.34	3.66	3.69	3.26	3.46	3.69	3.99	4.54	4.31	4.63
Prison/Jail	2.12	1.95	1.77	1.41	2.70	1.56	1.76	2.36	2.76	2.04	2.61	2.52	3.22

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.67P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 12 to 17 Who Received Illicit Drug or Alcohol Treatment in the Past Year: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Hospital - Inpatient	0.5732	0.1981	0.1566	0.7940	0.4870	0.3653	0.2378	0.4297	0.2080	0.1950	0.0631	0.4716	
Rehabilitation Facility - Inpatient	0.7238	0.6144	0.9521	0.9608	0.3053	0.9824	0.5839	0.4486	0.3703	0.7783	0.9546	0.9176	
Rehabilitation Facility - Outpatient	0.1178	0.3804	0.4394	0.1889	0.4653	0.2253	0.8916	0.2483	0.6451	0.6677	0.0109	0.0433	
Mental Health Center - Outpatient	0.8529	0.6602	0.7644	0.7873	0.3623	0.2469	0.3853	0.4483	0.5090	0.8855	0.4717	0.4745	
Emergency Room	0.0094	0.2347	0.0553	0.1713	0.3464	0.1772	0.0468	0.0841	0.0878	0.1617	0.0356	0.6352	
Private Doctor's Office	0.1625	0.3754	0.5002	0.1850	0.3581	0.3483	0.0218	0.0301	0.1599	0.0935	0.1019	0.1782	
Self-Help Group	0.0068	0.3576	0.6225	0.5219	0.5195	0.0249	0.3991	0.9642	0.7678	0.5531	0.4820	0.5815	
Prison/Jail	0.7510	0.8142	0.8313	0.3476	0.8439	0.4354	0.5512	0.8915	0.9118	0.5603	0.9789	0.9936	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.68C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 18 to 25 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	38	46	49	48	50	46	47	50	52	51	54	47	52
Hospital - Inpatient	16	17	16	19	26	18	18	22	16	24	26	22	22
Rehabilitation Facility - Inpatient	19	22	19	20	28	24	17	24	24	29	28	24	29
Rehabilitation Facility - Outpatient	23	27	30	29	33	30	29	34	30	34	31	31	34
Mental Health Center - Outpatient	20	18	20	22	28	19	20	26	21	25	21	22	24
Emergency Room	11	14	12	13	16	15	14	18	14	18	19	16	16
Private Doctor's Office	13	12	13	16	20	19	20	20	15	21	23	19	18
Self-Help Group	27	36	34	32	41	31	35	37	39	37	42	31	38
Prison/Jail	10	11	11	16	17	16	11	16	15	20	9	11	13

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.68D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 18 to 25 Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Percentages, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Hospital - Inpatient	2.14	1.76	1.71	1.83	2.52	1.89	1.98	2.01	1.59	2.36	2.62	2.42	2.52
Rehabilitation Facility - Inpatient	2.39	2.20	1.96	1.91	2.66	2.32	1.90	2.18	2.22	2.65	2.74	2.60	3.11
Rehabilitation Facility - Outpatient	2.55	2.36	2.57	2.49	2.81	2.59	2.66	2.64	2.57	2.80	2.77	2.91	3.20
Mental Health Center - Outpatient	2.46	1.81	2.13	2.00	2.60	1.96	2.12	2.27	2.04	2.44	2.21	2.41	2.68
Emergency Room	1.49	1.44	1.25	1.31	1.74	1.59	1.56	1.74	1.35	1.90	2.09	1.94	2.00
Private Doctor's Office	1.75	1.31	1.37	1.50	1.99	1.98	2.12	1.95	1.53	2.14	2.36	2.13	2.10
Self-Help Group	2.77	2.63	2.62	2.36	2.63	2.67	2.72	2.51	2.55	2.63	3.18	2.79	3.21
Prison/Jail	1.41	1.25	1.22	1.54	1.78	1.76	1.23	1.56	1.46	2.04	1.10	1.37	1.66

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.68P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 18 to 25 Who Received Illicit Drug or Alcohol Treatment in the Past Year: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	
Hospital - Inpatient	0.3331	0.0385	0.0844	0.1364	0.5221	0.1823	0.2469	0.4357	0.0117	0.5907	0.4316	0.8192	
Rehabilitation Facility - Inpatient	0.2559	0.0669	0.1249	0.0081	0.1595	0.1391	0.0189	0.1543	0.1020	0.8290	0.3440	0.7319	
Rehabilitation Facility - Outpatient	0.3930	0.0714	0.8293	0.1884	0.6347	0.6548	0.8519	0.5085	0.3789	0.5412	0.1757	0.4843	
Mental Health Center - Outpatient	0.5295	0.0049	0.1835	0.2746	0.5450	0.0137	0.1982	0.4567	0.0459	0.9708	0.0557	0.3579	
Emergency Room	0.3706	0.1405	0.3689	0.1769	0.9543	0.9239	0.7610	0.8717	0.1632	0.3110	0.7260	0.7949	
Private Doctor's Office	0.3464	0.0369	0.1131	0.3809	0.3620	0.6523	0.2662	0.8245	0.4706	0.2008	0.2100	0.4546	
Self-Help Group	0.7400	0.9401	1.0000	0.1474	0.1214	0.1423	0.8941	0.5928	0.5363	0.7177	0.5922	0.3213	
Prison/Jail	0.3717	0.7751	0.4240	0.0352	0.0294	0.0134	0.7152	0.0658	0.0764	0.0017	0.9514	0.5959	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.69C Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 26 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Numbers in Thousands, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	170	159	191	181	210	184	199	203	190	186	192	203	173
Hospital - Inpatient	100	63	85	76	96	97	77	86	84	87	98	91	79
Rehabilitation Facility - Inpatient	107	76	84	108	91	105	70	112	92	97	97	101	76
Rehabilitation Facility - Outpatient	119	88	130	127	118	134	107	144	123	117	114	128	110
Mental Health Center - Outpatient	104	66	107	113	99	91	104	105	91	90	93	105	89
Emergency Room	67	31	60	54	53	76	52	58	70	74	86	71	57
Private Doctor's Office	79	43	59	56	69	65	78	80	74	79	90	83	75
Self-Help Group	131	126	151	140	162	145	156	156	162	151	141	169	130
Prison/Jail	37	34	48	52	64	49	54	58	42	56	64	40	49

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.69D Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 26 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: Standard Errors of Percentages, 2002-2014

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Hospital - Inpatient	3.37	2.79	2.96	2.48	2.99	3.06	2.51	2.61	2.66	2.98	2.91	2.65	2.20
Rehabilitation Facility - Inpatient	3.43	3.14	2.82	3.35	2.94	3.11	2.39	3.09	2.84	3.20	2.90	2.89	2.26
Rehabilitation Facility - Outpatient	3.54	3.55	3.42	3.48	3.50	3.32	3.12	3.37	3.51	3.39	3.14	3.17	2.52
Mental Health Center - Outpatient	3.43	2.82	3.36	3.53	2.89	2.88	3.09	2.98	2.73	2.89	2.82	2.95	2.35
Emergency Room	2.53	1.49	2.15	1.94	1.83	2.47	1.75	1.85	2.24	2.53	2.71	2.19	1.69
Private Doctor's Office	2.91	2.01	2.25	2.05	2.34	2.15	2.46	2.38	2.36	2.66	2.79	2.48	2.08
Self-Help Group	3.48	3.62	3.69	3.53	3.86	3.50	3.43	3.28	3.37	3.24	3.19	3.07	2.71
Prison/Jail	1.49	1.57	1.85	1.92	2.18	1.74	1.82	1.82	1.39	2.02	2.09	1.30	1.47

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.69P Locations Received Illicit Drug or Alcohol Treatment in the Past Year among Persons Aged 26 or Older Who Received Illicit Drug or Alcohol Treatment in the Past Year: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Location of Treatment ¹	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL POPULATION	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Hospital - Inpatient	0.2208	0.3895	0.8938	0.6113	0.6820	0.7602	0.0836	0.6044	0.3829	0.6414	0.8075	0.9635	
Rehabilitation Facility - Inpatient	0.0253	0.6671	0.8047	0.0782	0.7134	0.3900	0.0185	0.2729	0.9148	0.7650	0.9547	0.9919	
Rehabilitation Facility - Outpatient	0.3210	0.4461	0.3833	0.6867	0.7352	0.3877	0.0494	0.1750	0.8571	0.7154	0.1921	0.9221	
Mental Health Center - Outpatient	0.4550	0.2131	0.8847	0.9933	0.9557	0.2999	0.8848	0.6160	0.4825	0.8868	0.6757	0.5880	
Emergency Room	0.2838	0.0087	0.7727	0.4880	0.1399	0.5441	0.1381	0.7377	0.9359	0.2374	0.2264	0.2680	
Private Doctor's Office	0.3120	0.0044	0.0327	0.0038	0.1026	0.1210	0.5011	0.6074	0.4084	0.9423	0.8185	0.9374	
Self-Help Group	0.1339	0.0758	0.2986	0.4438	0.9930	0.2553	0.4965	0.2362	0.3283	0.2094	0.9226	0.1966	
Prison/Jail	0.2063	0.0723	0.6179	0.6426	0.7796	0.1833	0.8810	0.5959	0.3475	0.5031	0.6096	0.0878	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Received Illicit Drug or Alcohol Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. It includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

¹ Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.70C Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	231	215	228	222	240	249	219	244	264	223	264	261	237
Received Treatment at a Specialty Facility	121	91	114	108	125	114	94	124	116	109	117	123	108
Did Not Receive Treatment at a Specialty Facility	205	192	203	198	203	217	196	213	231	191	243	226	214
NEEDED TREATMENT FOR ALCOHOL	411	389	431	413	419	421	407	427	416	390	424	429	375
Received Treatment at a Specialty Facility	124	102	135	139	125	131	138	127	126	124	127	125	121
Did Not Receive Treatment at a Specialty Facility	403	370	409	386	407	398	387	410	400	372	406	406	351
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	436	415	461	443	463	470	439	474	476	423	477	477	404
Received Treatment at a Specialty Facility ²	151	118	165	157	162	166	159	168	155	150	155	160	148
Did Not Receive Treatment at a Specialty Facility	429	390	426	412	441	434	411	441	453	397	453	444	373

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.70D Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	0.10	0.09	0.09	0.09	0.10	0.10	0.09	0.10	0.10	0.09	0.10	0.10	0.09
Received Treatment at a Specialty Facility	0.05	0.04	0.05	0.04	0.05	0.05	0.04	0.05	0.05	0.04	0.04	0.05	0.04
Did Not Receive Treatment at a Specialty Facility	0.09	0.08	0.08	0.08	0.08	0.09	0.08	0.08	0.09	0.07	0.09	0.09	0.08
NEEDED TREATMENT FOR ALCOHOL	0.17	0.16	0.18	0.17	0.17	0.17	0.16	0.17	0.16	0.15	0.16	0.16	0.14
Received Treatment at a Specialty Facility	0.05	0.04	0.06	0.06	0.05	0.05	0.06	0.05	0.05	0.05	0.05	0.05	0.05
Did Not Receive Treatment at a Specialty Facility	0.17	0.16	0.17	0.16	0.17	0.16	0.16	0.16	0.16	0.14	0.16	0.15	0.13
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	0.19	0.17	0.19	0.18	0.19	0.19	0.18	0.19	0.19	0.16	0.18	0.18	0.15
Received Treatment at a Specialty Facility ²	0.06	0.05	0.07	0.06	0.07	0.07	0.06	0.07	0.06	0.06	0.06	0.06	0.06
Did Not Receive Treatment at a Specialty Facility	0.18	0.16	0.18	0.17	0.18	0.18	0.16	0.18	0.18	0.15	0.17	0.17	0.14

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.70P Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	0.0157	0.3768	0.0041	0.3041	0.1688	0.5693	0.6232	0.2570	0.2547	0.1420	0.3799	0.5846	
Received Treatment at a Specialty Facility	0.9486	0.0120	0.8540	0.1937	0.5750	0.3678	0.0310	0.8079	0.9186	0.1773	0.7971	0.5221	
Did Not Receive Treatment at a Specialty Facility	0.0062	0.0269	0.0010	0.0674	0.2069	0.2680	0.1074	0.1574	0.1815	0.3434	0.2759	0.7775	
NEEDED TREATMENT FOR ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0015	0.6466	0.0801	0.3021	
Received Treatment at a Specialty Facility	0.3441	0.4575	0.5287	0.3013	0.5730	0.5425	0.6415	0.2062	0.6071	0.7452	0.7740	0.4371	
Did Not Receive Treatment at a Specialty Facility	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0015	0.5520	0.0528	0.1783	
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	0.0000	0.0003	0.0000	0.0000	0.0000	0.0002	0.0005	0.0002	0.0056	0.6508	0.1029	0.4797	
Received Treatment at a Specialty Facility ²	0.8653	0.0094	0.8586	0.6894	0.5909	0.9748	0.4424	0.4911	0.6311	0.3188	0.7755	0.6008	
Did Not Receive Treatment at a Specialty Facility	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0002	0.0055	0.9200	0.0658	0.3377	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.71C Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	51	46	48	48	48	42	47	45	47	48	45	41	46
Received Treatment at a Specialty Facility	18	14	16	15	18	13	14	13	13	17	14	12	12
Did Not Receive Treatment at a Specialty Facility	46	45	45	45	45	40	45	43	45	44	43	39	45
NEEDED TREATMENT FOR ALCOHOL	51	50	52	52	49	48	49	50	50	46	40	37	42
Received Treatment at a Specialty Facility	17	12	15	14	14	12	11	13	11	11	12	13	12
Did Not Receive Treatment at a Specialty Facility	49	49	49	50	48	47	48	47	48	44	39	35	40
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	61	61	63	61	60	57	59	59	61	58	55	51	55
Received Treatment at a Specialty Facility ²	19	17	18	17	20	15	16	16	16	18	16	15	16
Did Not Receive Treatment at a Specialty Facility	57	61	60	58	58	56	58	56	58	55	53	49	54

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.71D Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	0.20	0.18	0.19	0.19	0.19	0.17	0.19	0.18	0.19	0.19	0.18	0.17	0.19
Received Treatment at a Specialty Facility	0.07	0.05	0.06	0.06	0.07	0.05	0.06	0.05	0.06	0.07	0.06	0.05	0.05
Did Not Receive Treatment at a Specialty Facility	0.19	0.18	0.18	0.18	0.18	0.16	0.18	0.17	0.18	0.18	0.17	0.16	0.18
NEEDED TREATMENT FOR ALCOHOL	0.21	0.20	0.20	0.21	0.19	0.19	0.20	0.20	0.20	0.18	0.16	0.15	0.17
Received Treatment at a Specialty Facility	0.07	0.05	0.06	0.05	0.06	0.05	0.04	0.05	0.05	0.04	0.05	0.05	0.05
Did Not Receive Treatment at a Specialty Facility	0.20	0.20	0.20	0.20	0.19	0.19	0.19	0.19	0.20	0.18	0.15	0.14	0.16
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	0.25	0.24	0.25	0.24	0.23	0.23	0.24	0.24	0.25	0.23	0.22	0.21	0.22
Received Treatment at a Specialty Facility ²	0.08	0.07	0.07	0.07	0.08	0.06	0.06	0.06	0.07	0.07	0.07	0.06	0.06
Did Not Receive Treatment at a Specialty Facility	0.23	0.24	0.24	0.23	0.23	0.22	0.23	0.23	0.24	0.22	0.21	0.19	0.22

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.71P Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 12 to 17: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0006	0.0000	0.0009	0.0000	0.0000	0.0260	0.8400	
Received Treatment at a Specialty Facility	0.0017	0.0385	0.0034	0.0008	0.0060	0.0508	0.0435	0.0162	0.1511	0.0167	0.0142	0.3555	
Did Not Receive Treatment at a Specialty Facility	0.0000	0.0000	0.0000	0.0000	0.0001	0.0030	0.0000	0.0057	0.0000	0.0002	0.1176	0.9524	
NEEDED TREATMENT FOR ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0012	0.5313	
Received Treatment at a Specialty Facility	0.0015	0.0170	0.0002	0.0007	0.0165	0.1018	0.1624	0.0162	0.3110	0.6302	0.2136	0.2913	
Did Not Receive Treatment at a Specialty Facility	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0026	0.7468	
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0004	0.4602	
Received Treatment at a Specialty Facility ²	0.0021	0.0128	0.0019	0.0031	0.0068	0.0723	0.1090	0.0460	0.1492	0.1219	0.0376	0.5349	
Did Not Receive Treatment at a Specialty Facility	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0025	0.5612	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.72C Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	78	75	84	84	84	82	84	85	87	86	95	90	93
Received Treatment at a Specialty Facility	25	29	28	29	33	29	26	32	29	36	34	31	34
Did Not Receive Treatment at a Specialty Facility	72	70	82	81	77	77	81	80	82	79	90	83	88
NEEDED TREATMENT FOR ALCOHOL	113	110	121	121	122	117	116	117	127	118	115	123	120
Received Treatment at a Specialty Facility	24	26	28	25	31	25	28	29	29	32	19	26	27
Did Not Receive Treatment at a Specialty Facility	111	107	118	118	122	114	113	114	123	115	114	120	117
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	122	117	130	133	133	128	125	129	140	130	134	138	135
Received Treatment at a Specialty Facility ²	30	33	36	34	38	33	33	39	37	40	37	35	41
Did Not Receive Treatment at a Specialty Facility	120	115	127	130	131	124	121	125	137	127	131	133	130

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.72D Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 18 to 25: Standard Errors of Percentages, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	0.25	0.24	0.26	0.26	0.26	0.25	0.26	0.25	0.26	0.25	0.28	0.26	0.27
Received Treatment at a Specialty Facility	0.08	0.09	0.09	0.09	0.10	0.09	0.08	0.09	0.08	0.10	0.10	0.09	0.10
Did Not Receive Treatment at a Specialty Facility	0.23	0.22	0.25	0.25	0.23	0.23	0.25	0.24	0.24	0.23	0.26	0.24	0.25
NEEDED TREATMENT FOR ALCOHOL	0.36	0.35	0.38	0.37	0.37	0.36	0.35	0.35	0.37	0.34	0.33	0.35	0.34
Received Treatment at a Specialty Facility	0.08	0.08	0.09	0.08	0.10	0.08	0.08	0.09	0.09	0.09	0.05	0.07	0.08
Did Not Receive Treatment at a Specialty Facility	0.36	0.34	0.37	0.36	0.37	0.35	0.34	0.34	0.36	0.34	0.33	0.35	0.33
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	0.39	0.37	0.40	0.41	0.41	0.39	0.38	0.39	0.41	0.38	0.39	0.40	0.39
Received Treatment at a Specialty Facility ²	0.10	0.10	0.11	0.10	0.12	0.10	0.10	0.12	0.11	0.12	0.11	0.10	0.12
Did Not Receive Treatment at a Specialty Facility	0.39	0.36	0.39	0.40	0.40	0.38	0.37	0.37	0.40	0.37	0.38	0.38	0.37

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.72P Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 18 to 25: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	0.0000	0.0006	0.0000	0.0000	0.0004	0.0002	0.0019	0.0020	0.0010	0.0060	0.0019	0.0861	
Received Treatment at a Specialty Facility	0.9095	0.3197	0.2428	0.2936	0.4568	0.9881	0.9247	0.1792	0.4215	0.1065	0.3985	0.7569	
Did Not Receive Treatment at a Specialty Facility	0.0000	0.0011	0.0000	0.0000	0.0005	0.0001	0.0010	0.0058	0.0013	0.0227	0.0030	0.0857	
NEEDED TREATMENT FOR ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0919	
Received Treatment at a Specialty Facility	0.1180	0.0208	0.0098	0.0616	0.0297	0.0782	0.0026	0.0026	0.0621	0.0732	0.0616	0.7881	
Did Not Receive Treatment at a Specialty Facility	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0732	
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0913	
Received Treatment at a Specialty Facility ²	0.7031	0.2349	0.0287	0.0972	0.2307	0.3419	0.2428	0.0155	0.1512	0.0658	0.9829	0.8445	
Did Not Receive Treatment at a Specialty Facility	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0701	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.73C Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	213	187	209	196	217	224	186	220	232	197	242	238	207
Received Treatment at a Specialty Facility	116	84	107	101	120	107	88	120	111	100	113	115	100
Did Not Receive Treatment at a Specialty Facility	178	168	181	170	182	193	164	190	201	163	221	209	180
NEEDED TREATMENT FOR ALCOHOL	389	358	389	369	390	395	377	405	393	369	392	405	337
Received Treatment at a Specialty Facility	121	96	131	135	121	127	132	124	121	120	127	121	115
Did Not Receive Treatment at a Specialty Facility	380	341	367	345	376	374	355	385	375	349	372	385	318
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	406	378	424	395	423	437	399	448	439	397	434	443	363
Received Treatment at a Specialty Facility ²	145	110	157	150	158	160	151	163	148	141	153	153	139
Did Not Receive Treatment at a Specialty Facility	397	357	385	366	400	402	370	414	416	370	409	418	335

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.73D Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 26 or Older: Standard Errors of Percentages, 2002-2014

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	0.12	0.10	0.11	0.11	0.12	0.12	0.10	0.11	0.12	0.10	0.12	0.12	0.10
Received Treatment at a Specialty Facility	0.06	0.05	0.06	0.05	0.06	0.06	0.05	0.06	0.06	0.05	0.06	0.06	0.05
Did Not Receive Treatment at a Specialty Facility	0.10	0.09	0.10	0.09	0.10	0.10	0.09	0.10	0.10	0.08	0.11	0.10	0.09
NEEDED TREATMENT FOR ALCOHOL	0.22	0.20	0.21	0.20	0.21	0.21	0.20	0.21	0.20	0.19	0.20	0.20	0.16
Received Treatment at a Specialty Facility	0.07	0.05	0.07	0.07	0.06	0.07	0.07	0.06	0.06	0.06	0.06	0.06	0.06
Did Not Receive Treatment at a Specialty Facility	0.21	0.19	0.20	0.19	0.20	0.20	0.18	0.20	0.19	0.18	0.19	0.19	0.15
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	0.23	0.21	0.23	0.21	0.23	0.23	0.21	0.23	0.22	0.20	0.22	0.22	0.18
Received Treatment at a Specialty Facility ²	0.08	0.06	0.09	0.08	0.08	0.08	0.08	0.08	0.08	0.07	0.08	0.08	0.07
Did Not Receive Treatment at a Specialty Facility	0.22	0.20	0.21	0.20	0.21	0.21	0.19	0.21	0.21	0.19	0.20	0.21	0.16

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.73P Need for and Receipt of Treatment at a Specialty Facility for a Substance Use Problem in the Past Year among Persons Aged 26 or Older: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance/Substance Treatment Status	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
NEEDED TREATMENT FOR ILLICIT DRUGS¹	0.2802	0.0200	0.3957	0.0172	0.2476	0.0851	0.0420	0.3880	0.2346	0.0001	0.3879	0.1608	
Received Treatment at a Specialty Facility	0.6471	0.0008	0.3497	0.0296	1.0000	0.2393	0.0107	0.3479	0.5942	0.0172	0.3930	0.3745	
Did Not Receive Treatment at a Specialty Facility	0.3230	0.3897	0.6598	0.1378	0.1753	0.1818	0.3531	0.6636	0.2826	0.0008	0.6091	0.2663	
NEEDED TREATMENT FOR ALCOHOL	0.2476	0.9418	0.0947	0.1229	0.1399	0.1830	0.6366	0.1063	0.8909	0.1013	0.8869	0.6248	
Received Treatment at a Specialty Facility	0.7692	0.1094	0.7694	0.7051	0.8573	0.8911	0.7797	0.7159	0.9974	0.4074	0.9686	0.3941	
Did Not Receive Treatment at a Specialty Facility	0.2703	0.5597	0.0604	0.1327	0.1088	0.1735	0.5451	0.1134	0.8841	0.1488	0.8696	0.4236	
NEEDED TREATMENT FOR ILLICIT DRUGS OR ALCOHOL¹	0.5993	0.3748	0.3966	0.9960	0.5240	0.7894	0.7776	0.4329	0.7576	0.0035	0.7547	0.9362	
Received Treatment at a Specialty Facility ²	0.8042	0.0006	0.2847	0.2494	0.9648	0.6801	0.2099	0.9099	0.9645	0.0762	0.5983	0.5729	
Did Not Receive Treatment at a Specialty Facility	0.5194	0.7857	0.1719	0.6389	0.4849	0.6475	0.8341	0.3707	0.7547	0.0143	0.8977	0.7619	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents were classified as needing treatment for a substance use problem if they met at least one of three criteria during the past year: (1) dependent on the substance; (2) abuse of the substance; or (3) received treatment for substance use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.74C Type of Substance Treatment Received at Any Location in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014

Substance for Which Treatment Was Received/Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS BUT NOT ALCOHOL													
12-17	11	11	12	12	16	12	12	11	13	16	13	9	13
18-25	18	25	23	25	19	23	23	24	26	26	36	21	30
26 or Older	76	52	73	63	92	71	77	76	72	69	81	75	74
ALCOHOL BUT NOT ILLICIT DRUGS													
12-17	9	10	9	8	8	7	9	7	8	7	7	6	5
18-25	20	25	27	27	25	29	25	27	31	22	24	19	22
26 or Older	96	96	124	117	108	107	151	122	136	138	132	126	101
ILLCIT DRUGS AND ALCOHOL													
12-17	20	16	19	17	18	17	16	16	15	16	19	14	15
18-25	23	28	29	26	34	26	29	32	31	34	23	31	28
26 or Older	106	92	104	112	123	122	88	123	103	100	104	118	108

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.74D Type of Substance Treatment Received at Any Location in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use in the Past Year, by Age Group: Standard Errors of Percentages, 2002-2014

Substance for Which Treatment Was Received/Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS BUT NOT ALCOHOL													
12-17	1.99	1.69	1.92	1.68	2.06	1.71	1.85	1.67	1.76	1.91	2.00	1.78	1.81
18-25	2.63	2.84	2.77	2.93	3.63	3.16	3.21	3.28	3.69	4.21	3.93	3.56	4.78
26 or Older	2.34	2.45	2.24	2.22	2.01	2.35	2.29	2.18	2.39	2.60	2.97	2.36	3.00
	2.83	2.35	2.76	2.28	2.90	2.31	2.48	2.25	2.33	2.54	2.54	2.20	2.13
ALCOHOL BUT NOT ILLICIT DRUGS													
12-17	2.39	2.41	2.59	2.36	2.22	2.39	2.56	2.34	2.50	2.74	2.56	2.54	2.05
18-25	2.38	2.61	2.09	2.20	2.17	1.96	2.45	2.42	2.73	2.24	2.36	2.46	2.07
26 or Older	2.50	2.45	2.50	2.26	2.41	2.63	2.35	2.30	2.56	2.23	2.58	2.21	2.53
	3.39	3.61	3.67	3.36	3.08	3.30	3.47	3.28	3.36	3.71	3.35	3.27	2.63
ILLCIT DRUGS AND ALCOHOL													
12-17	2.47	2.32	2.30	2.37	2.39	2.39	2.15	2.32	2.19	2.39	2.26	2.44	2.09
18-25	3.53	3.45	3.15	3.63	3.69	3.62	3.68	3.84	3.96	4.31	4.47	4.17	4.96
26 or Older	2.60	2.51	2.63	2.39	2.74	2.44	2.53	2.49	2.59	2.94	2.37	2.97	2.98
	3.47	3.50	3.33	3.37	3.32	3.36	2.85	3.23	2.97	3.24	3.05	3.15	2.60

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.74P Type of Substance Treatment Received at Any Location in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance for Which Treatment Was Received/Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS BUT NOT ALCOHOL	0.1021	0.0015	0.0494	0.0017	0.3949	0.0552	0.0379	0.0101	0.2452	0.2248	0.7169	0.2173	
12-17	0.0524	0.0541	0.0815	0.1081	0.6163	0.0670	0.0754	0.1035	0.8004	0.4798	0.7426	0.2400	
18-25	0.0119	0.0026	0.0557	0.0485	0.0005	0.0010	0.0030	0.0028	0.0740	0.2651	0.1362	0.1053	
26 or Older	0.3899	0.0088	0.1155	0.0027	0.9783	0.3866	0.2261	0.0782	0.3705	0.1394	0.7745	0.4743	
ALCOHOL BUT NOT ILLICIT DRUGS	0.7728	0.3639	0.7322	0.2961	0.8144	0.2662	0.0081	0.0603	0.0275	0.0234	0.2103	0.1784	
12-17	0.0109	0.0141	0.0217	0.0294	0.0453	0.1481	0.0951	0.0548	0.1238	0.2863	0.3100	0.2600	
18-25	0.1597	0.2568	0.1030	0.1685	0.1554	0.0001	0.0070	0.1015	0.0461	0.6825	0.7328	0.4621	
26 or Older	0.8506	0.2534	0.7574	0.2857	0.6928	0.6592	0.0182	0.0951	0.0446	0.0078	0.1143	0.1326	
ILLCIT DRUGS AND ALCOHOL	0.2837	0.2842	0.1648	0.1702	0.1489	0.5966	0.5492	0.6269	0.3624	0.4866	0.1848	0.5764	
12-17	0.3053	0.4797	0.4772	0.2127	0.6473	0.4448	0.5663	0.1504	0.7908	0.6793	0.2885	0.1968	
18-25	0.1717	0.0521	0.2719	0.7725	0.0069	0.4406	0.1435	0.0703	0.8383	0.1927	0.1418	0.0616	
26 or Older	0.6237	0.7789	0.3367	0.2374	0.5256	0.8567	0.2148	0.8219	0.3123	0.2520	0.2327	0.2216	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at Any Treatment Location includes treatment received at any location, such as a hospital (inpatient), rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.75C Type of Substance Treatment Received at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014

Substance for Which Treatment Was Received/Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS BUT NOT ALCOHOL													
12-17	77	51	76	56	94	73	67	75	73	72	80	77	77
18-25	8	9	8	9	14	9	9	8	9	13	11	7	*
26 or Older	16	20	20	23	17	20	18	22	22	24	31	19	26
	77	45	72	48	92	68	63	71	69	66	73	73	68
ALCOHOL BUT NOT ILLICIT DRUGS													
12-17	87	69	102	100	88	101	116	90	94	97	97	101	82
18-25	7	6	8	6	6	5	5	5	5	5	5	4	*
26 or Older	16	15	21	18	16	16	19	18	21	18	13	13	18
	86	67	99	98	86	99	114	88	91	95	97	101	79
ILLCIT DRUGS AND ALCOHOL													
12-17	88	71	73	88	82	76	63	86	79	77	84	75	73
18-25	16	9	13	12	12	9	9	11	8	9	10	10	*
26 or Older	18	22	19	17	27	19	18	19	18	26	13	22	20
	84	67	69	86	78	72	58	84	77	71	83	70	69

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.75D Type of Substance Treatment Received at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use at a Specialty Facility in the Past Year, by Age Group: Standard Errors of Percentages, 2002-2014

Substance for Which Treatment Was Received/Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS BUT NOT ALCOHOL													
12-17	2.78	2.45	2.83	2.27	3.04	2.66	2.68	2.45	2.44	2.74	2.80	2.75	2.56
18-25	4.22	4.55	3.94	4.35	5.63	4.87	4.96	4.56	5.44	5.60	5.60	5.16	*
26 or Older	3.07	3.24	3.03	3.33	2.99	3.42	3.11	3.15	3.23	3.56	3.77	3.41	3.86
	3.81	3.31	3.98	2.80	4.08	3.36	3.53	3.24	3.15	3.66	3.41	3.42	2.95
ALCOHOL BUT NOT ILLICIT DRUGS													
12-17	3.12	2.94	3.31	3.19	2.91	3.19	3.48	2.92	2.88	3.22	3.11	3.30	2.58
18-25	3.63	3.55	3.84	3.14	3.29	3.30	3.45	3.19	3.60	3.03	2.95	3.31	*
26 or Older	3.08	2.81	3.11	3.02	2.93	3.13	3.22	2.73	3.19	2.90	2.69	2.62	3.28
	4.18	4.33	4.58	4.40	3.86	4.20	4.61	4.01	3.74	4.42	4.02	4.18	3.18
ILLCIT DRUGS AND ALCOHOL													
12-17	3.05	2.93	2.70	3.03	2.79	2.67	2.48	2.69	2.56	2.85	2.90	2.67	2.28
18-25	5.46	4.33	4.61	4.68	5.25	4.86	5.07	4.97	4.71	4.91	5.42	5.80	*
26 or Older	3.35	3.59	3.05	2.79	3.91	3.10	2.97	2.88	3.01	3.65	2.57	3.71	3.44
	4.04	4.33	3.75	4.25	3.63	3.47	3.27	3.67	3.39	3.84	3.82	3.37	2.79

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.75P Type of Substance Treatment Received at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Received Treatment for Substance Use at a Specialty Facility in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance for Which Treatment Was Received/Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
ILLCIT DRUGS BUT NOT ALCOHOL	0.1426	0.0078	0.0734	0.0004	0.7793	0.1639	0.0419	0.0211	0.4489	0.4380	0.9945	0.7589	
12-17	*	*	*	*	*	*	*	*	*	*	*	*	
18-25	0.0080	0.0082	0.0335	0.2050	0.0034	0.0239	0.0011	0.0102	0.2432	0.3855	0.0269	0.3895	
26 or Older	0.4581	0.0228	0.2142	0.0002	0.7110	0.4413	0.1907	0.1003	0.6001	0.2952	0.4423	0.9689	
ALCOHOL BUT NOT ILLICIT DRUGS	0.3135	0.1883	0.6431	0.1342	0.8624	0.0629	0.0284	0.4743	0.1933	0.1271	0.4910	0.2558	
12-17	*	*	*	*	*	*	*	*	*	*	*	*	
18-25	0.4590	0.9322	0.3150	0.2462	0.9263	0.1932	0.0324	0.5779	0.4328	0.8464	0.1191	0.3048	
26 or Older	0.3194	0.0678	0.6847	0.1122	0.9831	0.0700	0.0399	0.4134	0.1837	0.0564	0.2248	0.1349	
ILLCIT DRUGS AND ALCOHOL	0.2935	0.1368	0.1915	0.1027	0.4464	0.8764	0.7771	0.5135	0.5670	0.6692	0.8167	0.2484	
12-17	*	*	*	*	*	*	*	*	*	*	*	*	
18-25	0.0951	0.0342	0.6495	0.9160	0.0408	0.6062	0.2390	0.5195	0.8006	0.4340	0.1386	0.4237	
26 or Older	0.7822	0.5267	0.4094	0.2111	0.9790	0.6053	0.3395	0.8830	0.4548	0.3910	0.9458	0.0941	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Substance Use Treatment refers to treatment received in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use. Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.76C Receipt of Illicit Drug Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug Treatment, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	121	90	114	108	126	113	94	123	116	109	118	122	109
12-17	18	14	16	15	18	13	14	14	13	17	15	12	12
18-25	25	29	28	29	33	29	26	31	28	36	35	31	35
26 or Older	116	83	107	101	121	107	88	119	111	100	114	114	100
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	210	198	211	201	204	216	199	217	229	193	250	231	221
12-17	48	47	47	47	46	42	46	45	47	47	45	40	46
18-25	80	76	92	87	83	84	88	86	89	86	103	87	91
26 or Older	180	172	186	172	183	192	165	194	200	164	224	213	184

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.76D Receipt of Illicit Drug Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug Treatment, by Age Group: Standard Errors of Percentages, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	1.39	1.12	1.26	1.28	1.38	1.32	1.12	1.39	1.28	1.31	1.34	1.41	1.23
12-17	1.17	1.02	1.06	1.12	1.40	1.15	1.11	1.17	1.09	1.32	1.34	1.27	1.34
18-25	0.86	1.03	0.96	0.97	1.13	0.98	0.95	1.09	0.96	1.19	1.14	1.08	1.31
26 or Older	2.66	2.24	2.44	2.57	2.64	2.51	2.10	2.57	2.45	2.57	2.51	2.52	1.93
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	1.39	1.12	1.26	1.28	1.38	1.32	1.12	1.39	1.28	1.31	1.34	1.41	1.23
12-17	1.17	1.02	1.06	1.12	1.40	1.15	1.11	1.17	1.09	1.32	1.34	1.27	1.34
18-25	0.86	1.03	0.96	0.97	1.13	0.98	0.95	1.09	0.96	1.19	1.14	1.08	1.31
26 or Older	2.66	2.24	2.44	2.57	2.64	2.51	2.10	2.57	2.45	2.57	2.51	2.52	1.93

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.76P Receipt of Illicit Drug Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug Treatment, by Age Group: P
Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	0.2558	0.0015	0.1373	0.0576	0.9921	0.1985	0.0089	0.4281	0.4932	0.4037	0.4796	0.6462	
12-17	0.3180	0.8978	0.4361	0.0836	0.1296	0.3688	0.5490	0.1823	0.9658	0.2360	0.0771	0.3671	
18-25	0.0930	0.8249	0.5269	0.5121	0.6294	0.1790	0.2125	0.8250	0.6609	0.4962	0.7710	0.7361	
26 or Older	0.9097	0.0132	0.5681	0.2590	0.4793	0.7820	0.0806	0.5577	0.8840	0.7583	0.6721	0.8878	
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	0.2558	0.0015	0.1373	0.0576	0.9921	0.1985	0.0089	0.4281	0.4932	0.4037	0.4796	0.6462	
12-17	0.3180	0.8978	0.4361	0.0836	0.1296	0.3688	0.5490	0.1823	0.9658	0.2360	0.0771	0.3671	
18-25	0.0930	0.8249	0.5269	0.5121	0.6294	0.1790	0.2125	0.8250	0.6609	0.4962	0.7710	0.7361	
26 or Older	0.9097	0.0132	0.5681	0.2590	0.4793	0.7820	0.0806	0.5577	0.8840	0.7583	0.6721	0.8878	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug use, or for medical problems associated with illicit drug use.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs; (2) abuse of illicit drugs; or (3) received treatment for illicit drug use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.77C Receipt of Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Alcohol Treatment, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	123	102	136	140	124	130	138	127	126	125	127	125	121
12-17	17	12	15	14	14	12	11	13	11	11	12	13	12
18-25	24	26	29	25	31	26	28	29	29	32	19	26	28
26 or Older	120	96	132	135	121	126	132	123	121	120	127	121	115
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	430	387	423	408	425	427	408	429	435	397	433	436	364
12-17	51	52	52	53	49	49	50	48	51	46	40	35	40
18-25	143	129	141	148	149	145	143	142	150	148	148	142	133
26 or Older	401	354	372	362	388	394	368	395	402	364	390	408	328

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.77D Receipt of Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Alcohol Treatment, by Age Group: Standard Errors of Percentages, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	0.65	0.54	0.66	0.68	0.62	0.64	0.69	0.63	0.65	0.68	0.67	0.65	0.64
12-17	1.11	0.78	0.91	0.91	0.99	0.80	0.88	1.06	0.97	1.06	1.26	1.61	1.57
18-25	0.42	0.45	0.48	0.43	0.52	0.44	0.46	0.51	0.52	0.62	0.37	0.55	0.61
26 or Older	1.02	0.82	1.03	1.05	0.96	0.98	1.04	0.93	0.96	1.00	0.97	0.91	0.86
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	0.65	0.54	0.66	0.68	0.62	0.64	0.69	0.63	0.65	0.68	0.67	0.65	0.64
12-17	1.11	0.78	0.91	0.91	0.99	0.80	0.88	1.06	0.97	1.06	1.26	1.61	1.57
18-25	0.42	0.45	0.48	0.43	0.52	0.44	0.46	0.51	0.52	0.62	0.37	0.55	0.61
26 or Older	1.02	0.82	1.03	1.05	0.96	0.98	1.04	0.93	0.96	1.00	0.97	0.91	0.86

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.77P Receipt of Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Alcohol Treatment, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	0.5131	0.0336	0.2904	0.5801	0.2775	0.3952	0.4473	0.9063	0.6876	0.6352	0.4169	0.2603	
12-17	0.9090	0.3758	0.9389	0.9018	0.6907	0.2618	0.3431	0.8805	0.3419	0.4443	0.7605	0.3562	
18-25	0.3001	0.8544	0.9733	0.4679	0.7668	0.5777	0.6997	0.3315	0.8824	0.4749	0.0046	0.4870	
26 or Older	0.9426	0.0867	0.4138	0.9499	0.5127	0.7778	0.6604	0.8798	0.9611	0.7194	0.9304	0.2918	
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	0.5131	0.0336	0.2904	0.5801	0.2775	0.3952	0.4473	0.9063	0.6876	0.6352	0.4169	0.2603	
12-17	0.9090	0.3758	0.9389	0.9018	0.6907	0.2618	0.3431	0.8805	0.3419	0.4443	0.7605	0.3562	
18-25	0.3001	0.8544	0.9733	0.4679	0.7668	0.5777	0.6997	0.3315	0.8824	0.4749	0.0046	0.4870	
26 or Older	0.9426	0.0867	0.4138	0.9499	0.5127	0.7778	0.6604	0.8798	0.9611	0.7194	0.9304	0.2918	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.78C Receipt of Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment, by Age Group: Standard Errors of Numbers in Thousands, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	150	117	166	157	163	165	158	167	154	151	157	158	149
12-17	20	17	18	17	20	16	16	16	16	18	17	15	16
18-25	30	34	37	34	38	34	33	39	37	40	37	36	41
26 or Older	145	109	158	150	158	159	150	161	147	142	154	151	139
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	466	416	450	437	458	463	437	467	488	426	486	481	393
12-17	61	65	65	63	61	59	61	58	63	58	56	50	55
18-25	161	145	157	166	167	163	160	159	168	170	177	162	150
26 or Older	425	377	399	384	413	420	387	430	443	386	430	448	351

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.78D Receipt of Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment, by Age Group: Standard Errors of Percentages, 2002-2014

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	0.65	0.50	0.65	0.63	0.65	0.66	0.64	0.66	0.63	0.65	0.64	0.66	0.61
12-17	0.81	0.74	0.74	0.77	0.93	0.77	0.79	0.85	0.83	0.94	1.01	1.06	1.21
18-25	0.43	0.47	0.50	0.46	0.52	0.47	0.45	0.54	0.52	0.60	0.54	0.55	0.67
26 or Older	1.03	0.79	1.01	1.00	1.04	1.03	0.99	1.01	0.97	0.99	0.97	0.95	0.84
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	0.65	0.50	0.65	0.63	0.65	0.66	0.64	0.66	0.63	0.65	0.64	0.66	0.61
12-17	0.81	0.74	0.74	0.77	0.93	0.77	0.79	0.85	0.83	0.94	1.01	1.06	1.21
18-25	0.43	0.47	0.50	0.46	0.52	0.47	0.45	0.54	0.52	0.60	0.54	0.55	0.67
26 or Older	1.03	0.79	1.01	1.00	1.04	1.03	0.99	1.01	0.97	0.99	0.97	0.95	0.84

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.78P Receipt of Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment, by Age Group: P Values from Tests of Differences of Percentages, 2014 Versus 2002-2013

Substance Treatment Status/ Age Group	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
RECEIVED TREATMENT AT A SPECIALTY FACILITY	0.1399	0.0001	0.0595	0.0635	0.3262	0.2004	0.0535	0.5945	0.6347	0.3588	0.3828	0.4181	
12-17	0.8690	0.4664	0.7773	0.9082	0.8921	0.5392	0.4738	0.9658	0.5350	0.9474	0.3227	0.6796	
18-25	0.0324	0.2631	0.7484	0.3144	0.2485	0.2220	0.2561	0.6307	0.7001	0.5107	0.2204	0.4921	
26 or Older	0.6496	0.0008	0.1411	0.2143	0.7771	0.5835	0.2106	0.6741	0.9400	0.4134	0.6560	0.5261	
DID NOT RECEIVE TREATMENT AT A SPECIALTY FACILITY	0.1399	0.0001	0.0595	0.0635	0.3262	0.2004	0.0535	0.5945	0.6347	0.3588	0.3828	0.4181	
12-17	0.8690	0.4664	0.7773	0.9082	0.8921	0.5392	0.4738	0.9658	0.5350	0.9474	0.3227	0.6796	
18-25	0.0324	0.2631	0.7484	0.3144	0.2485	0.2220	0.2561	0.6307	0.7001	0.5107	0.2204	0.4921	
26 or Older	0.6496	0.0008	0.1411	0.2143	0.7771	0.5835	0.2106	0.6741	0.9400	0.4134	0.6560	0.5261	

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop illicit drug or alcohol use, or for medical problems associated with illicit drug or alcohol use.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 7.79D Marijuana Use in the Past Month, by Age Group: Standard Errors of Percentages, 1971-2014

Survey Year	Aged 12-17	Aged 18-25	Aged 26-34
1971	0.98	1.74	1.13
1972	1.00	2.02	1.57
1974	1.23	1.86	1.21
1976	1.22	1.82	1.41
1977	1.22	1.44	1.66
1979	2.51	5.03	3.34
1982	1.82	3.95	3.06
1985	1.82	3.15	2.96
1988	0.99	2.38	2.22
1990	0.93	2.01	1.74
1991	0.60	1.82	1.16
1992	0.60	1.62	1.45
1993	0.67	1.70	1.22
1994	0.50	0.81	0.47
1995	0.55	0.71	0.47
1996	0.54	0.80	0.42
1997	0.56	0.88	0.54
1998	0.57	0.68	0.46
1999	0.20	0.38	0.32
2000	0.21	0.34	0.33
2001	0.24	0.39	0.45
2002	0.24	0.36	0.45
2003	0.24	0.37	0.44
2004	0.23	0.37	0.47
2005	0.22	0.37	0.51
2006	0.21	0.35	0.47
2007	0.22	0.37	0.44
2008	0.22	0.37	0.48
2009	0.24	0.38	0.47
2010	0.25	0.38	0.54
2011	0.24	0.39	0.54
2012	0.22	0.39	0.52
2013	0.23	0.39	0.62
2014	0.27	0.45	0.45

*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Standard errors for 1971 through 1977 were calculated by multiplying the standard error of the percentage (assuming simple random sampling) by the square root of the estimated "design effect" (i.e., 1.25). See Appendix A of the *National Household Survey on Drug Abuse: Main Findings 1979* for further information on these procedures.

NOTE: Estimates for 1971 through 1977 for youths aged 12 to 17 and for adults aged 26 to 34 may differ from estimates for these age groups for these survey years that were published in other NSDUH reports. The estimates shown here for these survey years have been adjusted to improve their comparability with estimates based on later versions of the NSDUH instrument. Estimates were adjusted by applying the ratio of the adjusted and unadjusted rate from 1979 for each age group to the original estimate.

NOTE: Estimates for 1979 through 1993 have been adjusted for comparability with estimates for 1994 through 1998. Estimates were adjusted by formally modeling the effect of the new methodology, relative to the old methodology, using 1994 data. See Appendix E of the *National Household Survey on Drug Abuse: Main Findings 1998* (<http://oas.samhsa.gov/nsduh/reports.htm>) for information on the adjustment procedures.

NOTE: Estimates for 1999 through 2014 have not been adjusted for comparability. See Appendix C of the *Results from the 2004 National Survey on Drug Use and Health: National Findings* for information on the comparability of the 1999 and later estimates with pre-1999 estimates.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 1971-2014.

Table 7.80D Cocaine Use in the Past Year, by Age Group: Standard Errors of Percentages, 1971-2014

Survey Year	Aged 12-17	Aged 18-25	Aged 26-34
1971	--	--	--
1972	0.48	--	--
1974	0.61	1.09	--
1976	0.56	1.01	--
1977	0.51	0.91	--
1979	0.60	2.11	1.23
1982	0.80	2.10	1.41
1985	0.59	1.72	1.48
1988	0.43	1.54	1.02
1990	0.56	0.98	0.83
1991	0.22	0.82	0.55
1992	0.19	0.75	0.55
1993	0.15	0.64	0.52
1994	0.20	0.37	0.32
1995	0.26	0.44	0.27
1996	0.19	0.43	0.34
1997	0.29	0.42	0.38
1998	0.24	0.41	0.35
1999	0.10	0.21	0.22
2000	0.12	0.18	0.18
2001	0.10	0.23	0.28
2002	0.13	0.24	0.31
2003	0.11	0.23	0.36
2004	0.11	0.25	0.30
2005	0.11	0.23	0.27
2006	0.11	0.24	0.35
2007	0.11	0.24	0.32
2008	0.10	0.21	0.33
2009	0.09	0.20	0.30
2010	0.09	0.20	0.33
2011	0.08	0.21	0.25
2012	0.08	0.21	0.31
2013	0.06	0.21	0.32
2014	0.09	0.23	0.23

*Low precision; no estimate reported.

-- Not available.

NOTE: Cocaine use includes the use of crack.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Standard errors for 1971 through 1977 were calculated by multiplying the standard error of the percentage (assuming simple random sampling) by the square root of the estimated "design effect" (i.e., 1.25). See Appendix A of the *National Household Survey on Drug Abuse: Main Findings 1979* for further information on these procedures.

NOTE: Estimates for 1971 through 1977 for youths aged 12 to 17 may differ from estimates for these age groups for these survey years that were published in other NSDUH reports. The estimates shown here for these survey years have been adjusted to improve their comparability with estimates based on later versions of the NSDUH instrument. Estimates have been adjusted by applying the ratio of the adjusted and unadjusted rate from 1979 for each age group to the original estimate.

NOTE: Estimates for 1979 through 1993 have been adjusted for comparability with estimates for 1994 through 1998. Estimates were adjusted using a simple ratio correction factor that measured the effect of the new methodology, relative to the old methodology, estimated at the total population level using pooled 1993 and 1994 data. See Appendix E of the *National Household Survey on Drug Abuse: Main Findings 1998* (<http://oas.samhsa.gov/nsduh/reports.htm>) for further information on the adjustment procedures.

NOTE: Estimates for 1999 through 2014 have not been adjusted for comparability. See Appendix C of the *Results from the 2004 National Survey on Drug Use and Health: National Findings* for information on the comparability of the 1999 and later estimates with pre-1999 estimates.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 1971-2014.

Table 8.1A Numbers (in Thousands) of Persons Aged 12 or Older, by Gender and Detailed Age Category: 2013 and 2014

Age Category	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL	262,391	265,123	127,120	128,423	135,272	136,700
12	3,750	3,823	1,929	2,001	1,821	1,822
13	4,221	4,080	2,191	2,113	2,030	1,968
14	4,386	4,171	2,284	2,114	2,103	2,056
15	4,133	4,289	2,058	2,152	2,075	2,137
16	4,275	4,339	2,138	2,162	2,136	2,177
17	4,127	4,173	2,106	2,140	2,020	2,033
18	4,733	4,605	2,465	2,407	2,268	2,198
19	4,166	4,140	2,197	2,126	1,969	2,014
20	4,295	4,361	2,157	2,231	2,138	2,130
21	4,237	4,347	2,051	2,204	2,185	2,142
22	4,475	4,512	2,242	2,353	2,233	2,159
23	4,377	4,394	2,215	2,151	2,162	2,244
24	4,392	4,325	2,180	2,050	2,212	2,275
25	4,110	4,250	1,954	2,007	2,157	2,243
26-29	17,210	17,026	8,649	8,481	8,561	8,545
30-34	20,136	20,854	9,780	10,211	10,356	10,642
35-39	19,046	19,559	8,974	9,691	10,072	9,867
40-44	21,472	20,176	10,508	9,779	10,965	10,397
45-49	19,992	20,535	10,154	10,041	9,838	10,494
50-54	23,077	23,757	11,221	11,273	11,856	12,484
55-59	20,549	20,868	9,375	10,381	11,174	10,487
60-64	17,778	17,546	9,065	8,375	8,713	9,171
65 or Older	43,452	44,992	19,225	19,978	24,227	25,014

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.1N Survey Sample Size for Respondents Aged 12 or Older, by Gender and Detailed Age Category: 2013 and 2014

Age Category	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL	67,838	67,901	32,843	32,402	34,995	35,499
12	3,537	2,640	1,824	1,347	1,713	1,293
13	3,812	2,821	1,963	1,433	1,849	1,388
14	3,891	2,934	2,026	1,491	1,865	1,443
15	3,750	2,934	1,882	1,483	1,868	1,451
16	3,830	2,936	1,940	1,467	1,890	1,469
17	3,674	2,742	1,914	1,392	1,760	1,350
18	2,933	2,051	1,479	1,048	1,454	1,003
19	2,617	1,941	1,294	949	1,323	992
20	2,746	2,035	1,348	1,017	1,398	1,018
21	2,758	2,057	1,323	1,015	1,435	1,042
22	2,839	2,087	1,376	1,005	1,463	1,082
23	2,806	2,109	1,315	1,016	1,491	1,093
24	2,823	2,120	1,302	993	1,521	1,127
25	2,692	2,049	1,234	936	1,458	1,113
26-29	2,979	4,630	1,376	2,122	1,603	2,508
30-34	3,331	5,622	1,529	2,579	1,802	3,043
35-39	2,879	4,561	1,317	2,107	1,562	2,454
40-44	3,108	4,535	1,437	2,075	1,671	2,460
45-49	3,071	4,494	1,440	2,078	1,631	2,416
50-54	1,788	2,370	837	1,079	951	1,291
55-59	1,620	2,124	711	987	909	1,137
60-64	1,393	1,824	674	837	719	987
65 or Older	2,961	4,285	1,302	1,946	1,659	2,339

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.2A Numbers (in Thousands) of Persons Aged 12 or Older, by Age Group and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	262,391	265,123	24,893	24,875	34,786	34,935	202,713	205,313
GENDER								
Male	127,120	128,423	12,708	12,682	17,461	17,530	96,951	98,211
Female	135,272	136,700	12,185	12,193	17,325	17,405	105,762	107,102
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	221,130	222,651	19,329	19,228	27,535	27,570	174,267	175,853
White	169,985	170,234	13,632	13,482	19,496	19,363	136,857	137,390
Black or African American	31,233	31,680	3,442	3,485	5,006	5,033	22,786	23,162
American Indian or Alaska Native	1,401	1,442	159	166	230	203	1,012	1,073
Native Hawaiian or Other Pacific Islander	871	1,021	98	103	201	177	573	741
Asian	13,351	13,856	1,245	1,227	1,870	2,010	10,235	10,619
Two or More Races	4,290	4,417	754	766	733	784	2,804	2,867
Hispanic or Latino	41,261	42,472	5,564	5,646	7,251	7,365	28,446	29,461
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	82,764	82,942	6,956	6,887	9,767	9,763	66,042	66,291
Female, White, Not Hispanic	87,220	87,292	6,676	6,594	9,729	9,599	70,816	71,099
Male, Black, Not Hispanic	14,289	14,487	1,754	1,767	2,423	2,417	10,113	10,303
Female, Black, Not Hispanic	16,944	17,193	1,688	1,718	2,583	2,616	12,673	12,859
Male, Hispanic	20,703	21,214	2,842	2,873	3,744	3,770	14,118	14,571
Female, Hispanic	20,558	21,258	2,722	2,774	3,507	3,595	14,329	14,889
EDUCATION¹								
< High School	32,168	32,225	N/A	N/A	5,589	4,946	26,579	27,280
High School Graduate	69,976	68,918	N/A	N/A	11,971	11,974	58,005	56,944
Some College	63,397	65,581	N/A	N/A	12,180	12,443	51,218	53,139
College Graduate	71,958	73,523	N/A	N/A	5,046	5,573	66,912	67,951
CURRENT EMPLOYMENT¹								
Full-Time	118,952	122,934	N/A	N/A	13,513	14,255	105,439	108,679
Part-Time	33,749	33,015	N/A	N/A	9,667	9,477	24,082	23,538
Unemployed	11,805	10,793	N/A	N/A	4,157	3,888	7,648	6,905
Other ²	72,993	73,506	N/A	N/A	7,449	7,314	65,544	66,192

N/A: Not applicable.

¹ Estimates for education and current employment are shown only for persons aged 18 or older.² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.2N Survey Sample Size for Respondents Aged 12 or Older, by Age Group and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	67,838	67,901	22,494	17,007	22,214	16,449	23,130	34,445
GENDER								
Male	32,843	32,402	11,549	8,613	10,671	7,979	10,623	15,810
Female	34,995	35,499	10,945	8,394	11,543	8,470	12,507	18,635
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	56,238	56,158	18,042	13,415	18,184	13,249	20,012	29,494
White	41,495	41,525	12,740	9,259	13,023	9,390	15,732	22,876
Black or African American	8,607	8,106	3,066	2,255	2,996	2,130	2,545	3,721
American Indian or Alaska Native	865	1,040	298	272	300	281	267	487
Native Hawaiian or Other Pacific Islander	328	334	101	84	123	90	104	160
Asian	2,622	2,853	773	651	968	762	881	1,440
Two or More Races	2,321	2,300	1,064	894	774	596	483	810
Hispanic or Latino	11,600	11,743	4,452	3,592	4,030	3,200	3,118	4,951
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	20,184	19,929	6,505	4,732	6,298	4,556	7,381	10,641
Female, White, Not Hispanic	21,311	21,596	6,235	4,527	6,725	4,834	8,351	12,235
Male, Black, Not Hispanic	3,964	3,739	1,531	1,169	1,383	989	1,050	1,581
Female, Black, Not Hispanic	4,643	4,367	1,535	1,086	1,613	1,141	1,495	2,140
Male, Hispanic	5,699	5,542	2,366	1,771	1,918	1,566	1,415	2,205
Female, Hispanic	5,901	6,201	2,086	1,821	2,112	1,634	1,703	2,746
EDUCATION¹								
< High School	6,544	6,755	N/A	N/A	3,573	2,365	2,971	4,390
High School Graduate	14,414	15,160	N/A	N/A	7,768	5,752	6,646	9,408
Some College	13,812	14,940	N/A	N/A	7,753	5,792	6,059	9,148
College Graduate	10,574	14,039	N/A	N/A	3,120	2,540	7,454	11,499
CURRENT EMPLOYMENT¹								
Full-Time	22,133	27,383	N/A	N/A	8,858	6,925	13,275	20,458
Part-Time	8,949	8,477	N/A	N/A	6,095	4,395	2,854	4,082
Unemployed	3,545	3,058	N/A	N/A	2,555	1,742	990	1,316
Other ²	10,717	11,976	N/A	N/A	4,706	3,387	6,011	8,589

N/A: Not applicable.

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

¹ Estimates for education and current employment are shown only for persons aged 18 or older.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.3A Numbers (in Thousands) of Persons Aged 18 or Older, by Age Group and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26-49 (2013)	Aged 26-49 (2014)	Aged 50+ (2013)	Aged 50+ (2014)
TOTAL	237,499	240,248	34,786	34,935	97,857	98,150	104,856	107,163
GENDER								
Male	114,412	115,741	17,461	17,530	48,065	48,204	48,886	50,007
Female	123,087	124,507	17,325	17,405	49,792	49,946	55,970	57,156
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	201,802	203,422	27,535	27,570	79,264	79,141	95,003	96,711
White	156,353	156,753	19,496	19,363	58,813	58,161	78,044	79,229
Black or African American	27,791	28,195	5,006	5,033	11,919	12,108	10,867	11,055
American Indian or Alaska Native	1,242	1,276	230	203	609	560	403	514
Native Hawaiian or Other Pacific Islander	773	918	201	177	353	447	219	294
Asian	12,106	12,629	1,870	2,010	6,165	6,442	4,070	4,177
Two or More Races	3,537	3,651	733	784	1,405	1,424	1,399	1,443
Hispanic or Latino	35,697	36,826	7,251	7,365	18,593	19,009	9,854	10,452
EDUCATION								
< High School	32,168	32,225	5,589	4,946	12,025	11,911	14,554	15,369
High School Graduate	69,976	68,918	11,971	11,974	25,364	24,278	32,641	32,666
Some College	63,397	65,581	12,180	12,443	26,233	26,720	24,985	26,419
College Graduate	71,958	73,523	5,046	5,573	34,235	35,242	32,676	32,709
CURRENT EMPLOYMENT								
Full-Time	118,952	122,934	13,513	14,255	65,539	68,194	39,900	40,485
Part-Time	33,749	33,015	9,667	9,477	12,147	11,211	11,935	12,327
Unemployed	11,805	10,793	4,157	3,888	5,222	4,522	2,426	2,383
Other ¹	72,993	73,506	7,449	7,314	14,949	14,223	50,595	51,969

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.3N Survey Sample Size for Respondents Aged 18 or Older, by Age Group and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26-49 (2013)	Aged 26-49 (2014)	Aged 50+ (2013)	Aged 50+ (2014)
TOTAL	45,344	50,894	22,214	16,449	15,368	23,842	7,762	10,603
GENDER								
Male	21,294	23,789	10,671	7,979	7,099	10,961	3,524	4,849
Female	24,050	27,105	11,543	8,470	8,269	12,881	4,238	5,754
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	38,196	42,743	18,184	13,249	12,876	19,835	7,136	9,659
White	28,755	32,266	13,023	9,390	9,765	14,924	5,967	7,952
Black or African American	5,541	5,851	2,996	2,130	1,801	2,680	744	1,041
American Indian or Alaska Native	567	768	300	281	198	356	69	131
Native Hawaiian or Other Pacific Islander	227	250	123	90	84	128	20	32
Asian	1,849	2,202	968	762	688	1,142	193	298
Two or More Races	1,257	1,406	774	596	340	605	143	205
Hispanic or Latino	7,148	8,151	4,030	3,200	2,492	4,007	626	944
EDUCATION								
< High School	6,544	6,755	3,573	2,365	1,907	2,872	1,064	1,518
High School Graduate	14,414	15,160	7,768	5,752	4,087	6,049	2,559	3,359
Some College	13,812	14,940	7,753	5,792	4,191	6,524	1,868	2,624
College Graduate	10,574	14,039	3,120	2,540	5,183	8,397	2,271	3,102
CURRENT EMPLOYMENT								
Full-Time	22,133	27,383	8,858	6,925	10,281	16,399	2,994	4,059
Part-Time	8,949	8,477	6,095	4,395	1,914	2,832	940	1,250
Unemployed	3,545	3,058	2,555	1,742	805	1,081	185	235
Other ¹	10,717	11,976	4,706	3,387	2,368	3,530	3,643	5,059

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.4A Numbers (in Thousands) of Persons Aged 12 or Older, by Youth and Adult Age Groups and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18+ (2013)	Aged 18+ (2014)
TOTAL	262,391	265,123	24,893	24,875	237,499	240,248
GENDER						
Male	127,120	128,423	12,708	12,682	114,412	115,741
Female	135,272	136,700	12,185	12,193	123,087	124,507
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	221,130	222,651	19,329	19,228	201,802	203,422
White	169,985	170,234	13,632	13,482	156,353	156,753
Black or African American	31,233	31,680	3,442	3,485	27,791	28,195
American Indian or Alaska Native	1,401	1,442	159	166	1,242	1,276
Native Hawaiian or Other Pacific Islander	871	1,021	98	103	773	918
Asian	13,351	13,856	1,245	1,227	12,106	12,629
Two or More Races	4,290	4,417	754	766	3,537	3,651
Hispanic or Latino	41,261	42,472	5,564	5,646	35,697	36,826
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	82,764	82,942	6,956	6,887	75,809	76,055
Female, White, Not Hispanic	87,220	87,292	6,676	6,594	80,544	80,698
Male, Black, Not Hispanic	14,289	14,487	1,754	1,767	12,535	12,720
Female, Black, Not Hispanic	16,944	17,193	1,688	1,718	15,256	15,475
Male, Hispanic	20,703	21,214	2,842	2,873	17,861	18,341
Female, Hispanic	20,558	21,258	2,722	2,774	17,836	18,485
EDUCATION¹						
< High School	32,168	32,225	N/A	N/A	32,168	32,225
High School Graduate	69,976	68,918	N/A	N/A	69,976	68,918
Some College	63,397	65,581	N/A	N/A	63,397	65,581
College Graduate	71,958	73,523	N/A	N/A	71,958	73,523
CURRENT EMPLOYMENT¹						
Full-Time	118,952	122,934	N/A	N/A	118,952	122,934
Part-Time	33,749	33,015	N/A	N/A	33,749	33,015
Unemployed	11,805	10,793	N/A	N/A	11,805	10,793
Other ²	72,993	73,506	N/A	N/A	72,993	73,506

N/A: Not applicable.

¹ Estimates for education and current employment are shown only for persons aged 18 or older.² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.4N Survey Sample Size for Respondents Aged 12 or Older, by Youth and Adult Age Groups and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18+ (2013)	Aged 18+ (2014)
TOTAL	67,838	67,901	22,494	17,007	45,344	50,894
GENDER						
Male	32,843	32,402	11,549	8,613	21,294	23,789
Female	34,995	35,499	10,945	8,394	24,050	27,105
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	56,238	56,158	18,042	13,415	38,196	42,743
White	41,495	41,525	12,740	9,259	28,755	32,266
Black or African American	8,607	8,106	3,066	2,255	5,541	5,851
American Indian or Alaska Native	865	1,040	298	272	567	768
Native Hawaiian or Other Pacific Islander	328	334	101	84	227	250
Asian	2,622	2,853	773	651	1,849	2,202
Two or More Races	2,321	2,300	1,064	894	1,257	1,406
Hispanic or Latino	11,600	11,743	4,452	3,592	7,148	8,151
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	20,184	19,929	6,505	4,732	13,679	15,197
Female, White, Not Hispanic	21,311	21,596	6,235	4,527	15,076	17,069
Male, Black, Not Hispanic	3,964	3,739	1,531	1,169	2,433	2,570
Female, Black, Not Hispanic	4,643	4,367	1,535	1,086	3,108	3,281
Male, Hispanic	5,699	5,542	2,366	1,771	3,333	3,771
Female, Hispanic	5,901	6,201	2,086	1,821	3,815	4,380
EDUCATION¹						
< High School	6,544	6,755	N/A	N/A	6,544	6,755
High School Graduate	14,414	15,160	N/A	N/A	14,414	15,160
Some College	13,812	14,940	N/A	N/A	13,812	14,940
College Graduate	10,574	14,039	N/A	N/A	10,574	14,039
CURRENT EMPLOYMENT¹						
Full-Time	22,133	27,383	N/A	N/A	22,133	27,383
Part-Time	8,949	8,477	N/A	N/A	8,949	8,477
Unemployed	3,545	3,058	N/A	N/A	3,545	3,058
Other ²	10,717	11,976	N/A	N/A	10,717	11,976

N/A: Not applicable.

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

¹ Estimates for education and current employment are shown only for persons aged 18 or older.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.5A Numbers (in Thousands) of Persons Aged 12 or Older, by Underage and Legal Drinking Age Groups and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
TOTAL	262,391	265,123	38,087	37,981	224,305	227,142
GENDER						
Male	127,120	128,423	19,527	19,446	107,593	108,977
Female	135,272	136,700	18,560	18,535	116,712	118,165
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	221,130	222,651	29,654	29,277	191,476	193,373
White	169,985	170,234	20,856	20,482	149,128	149,752
Black or African American	31,233	31,680	5,337	5,323	25,896	26,357
American Indian or Alaska Native	1,401	1,442	246	247	1,156	1,195
Native Hawaiian or Other Pacific Islander	871	1,021	161	160	710	861
Asian	13,351	13,856	1,996	1,966	11,355	11,890
Two or More Races	4,290	4,417	1,059	1,100	3,231	3,318
Hispanic or Latino	41,261	42,472	8,432	8,704	32,829	33,768
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	82,764	82,942	10,701	10,474	72,063	72,468
Female, White, Not Hispanic	87,220	87,292	10,155	10,008	77,065	77,284
Male, Black, Not Hispanic	14,289	14,487	2,731	2,699	11,558	11,788
Female, Black, Not Hispanic	16,944	17,193	2,606	2,624	14,338	14,569
Male, Hispanic	20,703	21,214	4,305	4,495	16,398	16,719
Female, Hispanic	20,558	21,258	4,127	4,209	16,430	17,049

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.5N Survey Sample Size for Respondents Aged 12 or Older, by Underage and Legal Drinking Age Groups and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
TOTAL	67,838	67,901	30,790	23,034	37,048	44,867
GENDER						
Male	32,843	32,402	15,670	11,627	17,173	20,775
Female	34,995	35,499	15,120	11,407	19,875	24,092
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	56,238	56,158	24,748	18,122	31,490	38,036
White	41,495	41,525	17,461	12,528	24,034	28,997
Black or African American	8,607	8,106	4,199	3,026	4,408	5,080
American Indian or Alaska Native	865	1,040	407	383	458	657
Native Hawaiian or Other Pacific Islander	328	334	139	115	189	219
Asian	2,622	2,853	1,142	923	1,480	1,930
Two or More Races	2,321	2,300	1,400	1,147	921	1,153
Hispanic or Latino	11,600	11,743	6,042	4,912	5,558	6,831
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	20,184	19,929	8,871	6,358	11,313	13,571
Female, White, Not Hispanic	21,311	21,596	8,590	6,170	12,721	15,426
Male, Black, Not Hispanic	3,964	3,739	2,092	1,552	1,872	2,187
Female, Black, Not Hispanic	4,643	4,367	2,107	1,474	2,536	2,893
Male, Hispanic	5,699	5,542	3,127	2,429	2,572	3,113
Female, Hispanic	5,901	6,201	2,915	2,483	2,986	3,718

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.6A Numbers (in Thousands) of Persons Aged 12 or Older, by Age Group and Geographic Characteristics: 2013 and 2014

Geographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	262,391	265,123	24,893	24,875	34,786	34,935	202,713	205,313
GEOGRAPHIC DIVISION								
Northeast	47,388	47,632	4,187	4,156	6,149	6,150	37,052	37,325
New England	12,483	12,565	1,092	1,086	1,630	1,644	9,761	9,836
Middle Atlantic	34,905	35,067	3,096	3,071	4,519	4,506	27,291	27,489
Midwest	56,215	56,462	5,398	5,372	7,407	7,428	43,410	43,663
East North Central	38,980	39,111	3,757	3,729	5,104	5,109	30,120	30,273
West North Central	17,234	17,351	1,641	1,643	2,303	2,318	13,290	13,390
South	97,513	98,844	9,356	9,411	12,858	12,943	75,299	76,490
South Atlantic	51,455	52,161	4,657	4,680	6,563	6,591	40,235	40,890
East South Central	15,495	15,594	1,475	1,474	2,040	2,050	11,980	12,070
West South Central	30,563	31,089	3,225	3,257	4,254	4,301	23,084	23,531
West	61,276	62,185	5,951	5,936	8,372	8,414	46,952	47,835
Mountain	18,697	19,008	1,875	1,892	2,532	2,559	14,290	14,557
Pacific	42,579	43,176	4,076	4,043	5,840	5,856	32,662	33,277
COUNTY TYPE								
Large Metro	141,041	146,556	13,559	14,025	18,746	19,175	108,736	113,356
Small Metro	78,898	77,412	7,448	7,072	10,871	10,636	60,579	59,704
250K – 1 Mil. Pop.	53,389	52,372	5,122	4,867	6,990	6,876	41,277	40,629
< 250K Pop.	25,509	25,041	2,326	2,206	3,881	3,759	19,302	19,076
Nonmetro	42,452	41,154	3,886	3,777	5,168	5,124	33,398	32,253
Urbanized	19,000	17,244	1,660	1,538	2,473	2,394	14,868	13,312
Less Urbanized	18,986	19,700	1,783	1,872	2,222	2,342	14,981	15,486
Completely Rural	4,466	4,211	443	367	473	389	3,549	3,455

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.6N Survey Sample Size for Respondents Aged 12 or Older, by Age Group and Geographic Characteristics: 2013 and 2014

Geographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	67,838	67,901	22,494	17,007	22,214	16,449	23,130	34,445
GEOGRAPHIC DIVISION								
Northeast	13,661	12,999	4,556	3,267	4,432	3,098	4,673	6,634
New England	5,448	5,791	1,818	1,458	1,782	1,386	1,848	2,947
Middle Atlantic	8,213	7,208	2,738	1,809	2,650	1,712	2,825	3,687
Midwest	18,822	15,825	6,207	3,908	6,265	3,794	6,350	8,123
East North Central	12,468	9,142	4,116	2,289	4,163	2,081	4,189	4,772
West North Central	6,354	6,683	2,091	1,619	2,102	1,713	2,161	3,351
South	20,782	22,781	6,885	5,814	6,692	5,580	7,205	11,387
South Atlantic	10,801	12,740	3,583	3,248	3,386	3,193	3,832	6,299
East South Central	3,616	3,765	1,271	940	1,164	892	1,181	1,933
West South Central	6,365	6,276	2,031	1,626	2,142	1,495	2,192	3,155
West	14,573	16,296	4,846	4,018	4,825	3,977	4,902	8,301
Mountain	7,296	7,790	2,439	1,905	2,443	1,956	2,414	3,929
Pacific	7,277	8,506	2,407	2,113	2,382	2,021	2,488	4,372
COUNTY TYPE								
Large Metro	30,126	30,393	10,070	7,833	9,721	7,181	10,335	15,379
Small Metro	23,290	23,361	7,653	5,627	7,957	5,963	7,680	11,771
250K – 1 Mil. Pop.	15,031	14,907	4,981	3,666	5,012	3,648	5,038	7,593
< 250K Pop.	8,259	8,454	2,672	1,961	2,945	2,315	2,642	4,178
Nonmetro	14,422	14,147	4,771	3,547	4,536	3,305	5,115	7,295
Urbanized	6,667	5,813	2,116	1,387	2,212	1,473	2,339	2,953
Less Urbanized	6,140	6,542	2,087	1,650	1,906	1,496	2,147	3,396
Completely Rural	1,615	1,792	568	510	418	336	629	946

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.7A Numbers (in Thousands) of Persons Aged 18 or Older, by Age Group and Geographic Characteristics: 2013 and 2014

Geographic Characteristic	Total (2013)	Total (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26-49 (2013)	Aged 26-49 (2014)	Aged 50+ (2013)	Aged 50+ (2014)
TOTAL	237,499	240,248	34,786	34,935	97,857	98,150	104,856	107,163
GEOGRAPHIC DIVISION								
Northeast	43,201	43,476	6,149	6,150	17,427	17,377	19,625	19,948
New England	11,391	11,480	1,630	1,644	4,496	4,474	5,265	5,362
Middle Atlantic	31,810	31,996	4,519	4,506	12,931	12,904	14,360	14,586
Midwest	50,817	51,091	7,407	7,428	20,360	20,253	23,050	23,410
East North Central	35,223	35,382	5,104	5,109	14,134	14,041	15,985	16,232
West North Central	15,593	15,709	2,303	2,318	6,226	6,212	7,064	7,178
South	88,157	89,433	12,858	12,943	36,480	36,684	38,819	39,806
South Atlantic	46,798	47,481	6,563	6,591	18,959	19,036	21,277	21,853
East South Central	14,020	14,120	2,040	2,050	5,640	5,615	6,340	6,455
West South Central	27,338	27,832	4,254	4,301	11,882	12,032	11,202	11,498
West	55,325	56,249	8,372	8,414	23,589	23,835	23,363	24,000
Mountain	16,822	17,116	2,532	2,559	7,022	7,090	7,268	7,467
Pacific	38,503	39,133	5,840	5,856	16,568	16,745	16,095	16,532
COUNTY TYPE								
Large Metro	127,482	132,531	18,746	19,175	56,240	57,144	52,496	56,212
Small Metro	71,451	70,340	10,871	10,636	27,825	27,617	32,754	32,087
250K – 1 Mil. Pop.	48,267	47,505	6,990	6,876	19,201	18,578	22,076	22,050
< 250K Pop.	23,183	22,835	3,881	3,759	8,624	9,039	10,678	10,037
Nonmetro	38,566	37,378	5,168	5,124	13,791	13,390	19,606	18,864
Urbanized	17,340	15,706	2,473	2,394	6,300	6,013	8,568	7,300
Less Urbanized	17,204	17,827	2,222	2,342	6,180	6,220	8,801	9,266
Completely Rural	4,022	3,844	473	389	1,312	1,157	2,237	2,298

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.7N Survey Sample Size for Respondents Aged 18 or Older, by Age Group and Geographic Characteristics: 2013 and 2014

Geographic Characteristic	Total (2013)	Total (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26-49 (2013)	Aged 26-49 (2014)	Aged 50+ (2013)	Aged 50+ (2014)
TOTAL	45,344	50,894	22,214	16,449	15,368	23,842	7,762	10,603
GEOGRAPHIC DIVISION								
Northeast	9,105	9,732	4,432	3,098	3,027	4,609	1,646	2,025
New England	3,630	4,333	1,782	1,386	1,158	2,033	690	914
Middle Atlantic	5,475	5,399	2,650	1,712	1,869	2,576	956	1,111
Midwest	12,615	11,917	6,265	3,794	4,175	5,591	2,175	2,532
East North Central	8,352	6,853	4,163	2,081	2,795	3,318	1,394	1,454
West North Central	4,263	5,064	2,102	1,713	1,380	2,273	781	1,078
South	13,897	16,967	6,692	5,580	4,789	7,870	2,416	3,517
South Atlantic	7,218	9,492	3,386	3,193	2,546	4,372	1,286	1,927
East South Central	2,345	2,825	1,164	892	790	1,318	391	615
West South Central	4,334	4,650	2,142	1,495	1,453	2,180	739	975
West	9,727	12,278	4,825	3,977	3,377	5,772	1,525	2,529
Mountain	4,857	5,885	2,443	1,956	1,668	2,758	746	1,171
Pacific	4,870	6,393	2,382	2,021	1,709	3,014	779	1,358
COUNTY TYPE								
Large Metro	20,056	22,560	9,721	7,181	7,179	10,971	3,156	4,408
Small Metro	15,637	17,734	7,957	5,963	5,058	8,194	2,622	3,577
250K – 1 Mil. Pop.	10,050	11,241	5,012	3,648	3,322	5,230	1,716	2,363
< 250K Pop.	5,587	6,493	2,945	2,315	1,736	2,964	906	1,214
Nonmetro	9,651	10,600	4,536	3,305	3,131	4,677	1,984	2,618
Urbanized	4,551	4,426	2,212	1,473	1,478	2,009	861	944
Less Urbanized	4,053	4,892	1,906	1,496	1,315	2,113	832	1,283
Completely Rural	1,047	1,282	418	336	338	555	291	391

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.8A Numbers (in Thousands) of Persons Aged 12 or Older, by Underage and Legal Drinking Age Groups and Geographic Characteristics: 2013 and 2014

Geographic Characteristic	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
TOTAL	262,391	265,123	38,087	37,981	224,305	227,142
GEOGRAPHIC DIVISION						
Northeast	47,388	47,632	6,380	6,503	41,009	41,129
New England	12,483	12,565	1,675	1,783	10,808	10,782
Middle Atlantic	34,905	35,067	4,704	4,719	30,201	30,347
Midwest	56,215	56,462	8,218	8,115	47,997	48,348
East North Central	38,980	39,111	5,724	5,609	33,257	33,502
West North Central	17,234	17,351	2,494	2,505	14,740	14,846
South	97,513	98,844	14,071	14,076	83,442	84,768
South Atlantic	51,455	52,161	6,994	6,957	44,461	45,204
East South Central	15,495	15,594	2,215	2,248	13,280	13,345
West South Central	30,563	31,089	4,862	4,871	25,701	26,218
West	61,276	62,185	9,418	9,287	51,857	52,897
Mountain	18,697	19,008	2,834	2,828	15,863	16,180
Pacific	42,579	43,176	6,584	6,459	35,995	36,718
COUNTY TYPE						
Large Metro	141,041	146,556	20,631	21,100	120,410	125,456
Small Metro	78,898	77,412	11,549	11,129	67,350	66,283
250K – 1 Mil. Pop.	53,389	52,372	7,673	7,498	45,716	44,874
< 250K Pop.	25,509	25,041	3,876	3,631	21,634	21,409
Nonmetro	42,452	41,154	5,906	5,752	36,545	35,403
Urbanized	19,000	17,244	2,585	2,445	16,415	14,799
Less Urbanized	18,986	19,700	2,668	2,808	16,319	16,892
Completely Rural	4,466	4,211	653	499	3,812	3,712

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.8N Survey Sample Size for Respondents Aged 12 or Older, by Underage and Legal Drinking Age Groups and Geographic Characteristics: 2013 and 2014

Geographic Characteristic	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
TOTAL	67,838	67,901	30,790	23,034	37,048	44,867
GEOGRAPHIC DIVISION						
Northeast	13,661	12,999	6,225	4,456	7,436	8,543
New England	5,448	5,791	2,511	2,016	2,937	3,775
Middle Atlantic	8,213	7,208	3,714	2,440	4,499	4,768
Midwest	18,822	15,825	8,588	5,272	10,234	10,553
East North Central	12,468	9,142	5,705	3,021	6,763	6,121
West North Central	6,354	6,683	2,883	2,251	3,471	4,432
South	20,782	22,781	9,269	7,803	11,513	14,978
South Atlantic	10,801	12,740	4,765	4,339	6,036	8,401
East South Central	3,616	3,765	1,680	1,260	1,936	2,505
West South Central	6,365	6,276	2,824	2,204	3,541	4,072
West	14,573	16,296	6,708	5,503	7,865	10,793
Mountain	7,296	7,790	3,339	2,605	3,957	5,185
Pacific	7,277	8,506	3,369	2,898	3,908	5,608
COUNTY TYPE						
Large Metro	30,126	30,393	13,698	10,436	16,428	19,957
Small Metro	23,290	23,361	10,586	7,842	12,704	15,519
250K – 1 Mil. Pop.	15,031	14,907	6,818	5,022	8,213	9,885
< 250K Pop.	8,259	8,454	3,768	2,820	4,491	5,634
Nonmetro	14,422	14,147	6,506	4,756	7,916	9,391
Urbanized	6,667	5,813	2,946	1,905	3,721	3,908
Less Urbanized	6,140	6,542	2,820	2,221	3,320	4,321
Completely Rural	1,615	1,792	740	630	875	1,162

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.9A Numbers (in Thousands) of Persons Aged 12 or Older, by Youth and Adult Age Groups and Gender: 2002-2014

Demographic Characteristic	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL	235,143	237,682	240,515	243,220	246,022	247,845	249,815	251,816	253,619	257,599	260,057	262,391	265,123
Male	113,602	114,985	116,483	117,923	119,362	120,265	121,261	122,291	123,430	124,624	125,893	127,120	128,423
Female	121,541	122,697	124,032	125,297	126,659	127,581	128,554	129,524	130,189	132,975	134,164	135,272	136,700
12-17	24,754	24,995	25,214	25,355	25,392	25,241	24,892	24,609	24,347	24,974	24,933	24,893	24,875
Male	12,647	12,770	12,877	12,950	12,969	12,891	12,708	12,568	12,461	12,765	12,738	12,708	12,682
Female	12,107	12,225	12,337	12,405	12,423	12,351	12,185	12,041	11,885	12,208	12,195	12,185	12,193
12-20	37,201	37,517	37,737	38,323	38,180	38,455	38,081	38,241	37,978	38,498	38,206	38,087	37,981
18 or Older	210,390	212,687	215,300	217,865	220,630	222,604	224,923	227,207	229,273	232,625	235,124	237,499	240,248
18-25	31,024	31,728	32,194	32,486	32,740	32,731	32,938	33,580	34,072	34,302	34,590	34,786	34,935
26 or Older	179,365	180,958	183,106	185,379	187,890	189,873	191,985	193,627	195,200	198,324	200,534	202,713	205,313
26-49	100,287	99,992	100,102	100,112	100,422	100,144	99,833	99,381	98,566	98,033	97,947	97,857	98,150
26-34	35,163	34,961	34,975	34,858	35,110	35,300	35,634	36,215	36,524	36,411	36,912	37,346	37,880
35-49	65,124	65,031	65,128	65,255	65,312	64,844	64,199	63,166	62,043	61,622	61,035	60,511	60,270
50 or Older	79,079	80,966	83,004	85,267	87,468	89,729	92,152	94,246	96,634	100,290	102,588	104,856	107,163

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 8.9N Survey Sample Size for Respondents Aged 12 or Older, by Youth and Adult Age Groups and Gender: 2002-2014

Demographic Characteristic	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL	68,126	67,784	67,760	68,308	67,491	67,377	67,928	68,007	67,804	70,109	68,309	67,838	67,901
Male	32,767	32,611	32,696	32,786	32,568	32,570	32,714	32,922	32,837	33,778	32,861	32,843	32,402
Female	35,359	35,173	35,064	35,522	34,923	34,807	35,214	35,085	34,967	36,331	35,448	34,995	35,499
12-17	23,645	22,665	22,301	22,534	22,766	22,286	22,250	22,398	21,960	23,510	22,473	22,494	17,007
Male	12,053	11,637	11,363	11,378	11,675	11,452	11,373	11,398	11,140	12,028	11,376	11,549	8,613
Female	11,592	11,028	10,938	11,156	11,091	10,834	10,877	11,000	10,820	11,482	11,097	10,945	8,394
12-20	32,787	31,475	31,235	31,282	31,178	30,926	31,228	31,547	30,917	32,366	30,910	30,790	23,034
18 or Older	44,481	45,119	45,459	45,774	44,725	45,091	45,678	45,609	45,844	46,599	45,836	45,344	50,894
18-25	23,066	22,738	22,829	22,511	21,848	22,006	22,875	22,686	22,793	22,876	22,529	22,214	16,449
26 or Older	21,415	22,381	22,630	23,263	22,877	23,085	22,803	22,923	23,051	23,723	23,307	23,130	34,445
26-49	15,994	16,401	16,525	16,955	15,964	16,767	16,234	16,207	16,448	15,692	15,560	15,368	23,842
26-34	6,374	6,570	6,574	6,856	6,609	6,861	6,560	6,591	6,780	6,543	6,484	6,310	10,252
35-49	9,620	9,831	9,951	10,099	9,355	9,906	9,674	9,616	9,668	9,149	9,076	9,058	13,590
50 or Older	5,421	5,980	6,105	6,308	6,913	6,318	6,569	6,716	6,603	8,031	7,747	7,762	10,603

NOTE: Some 2006 to 2010 sample sizes may differ from previously published sample sizes due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 8.10A Numbers (in Thousands) of Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total¹ (2011-2012)	Total¹ (2013-2014)	Pregnant (2011-2012)	Pregnant (2013-2014)	Not Pregnant (2011-2012)	Not Pregnant (2013-2014)
TOTAL	62,478	63,357	2,206	2,305	60,032	60,758
AGE						
15-17	6,257	6,289	60	41	6,161	6,231
18-25	17,184	17,365	821	807	16,280	16,464
26-44	39,038	39,703	1,325	1,457	37,591	38,064
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	50,345	50,718	1,753	1,882	48,396	48,622
White	36,123	35,887	1,239	1,266	34,739	34,487
Black or African American	8,581	8,847	340	387	8,203	8,406
American Indian or Alaska Native	405	362	15	13	390	343
Native Hawaiian or Other Pacific Islander	261	266	13	14	248	252
Asian	3,834	4,132	109	155	3,720	3,960
Two or More Races	1,141	1,224	36	48	1,097	1,174
Hispanic or Latino	12,133	12,639	453	423	11,635	12,137
TRIMESTER²						
First	N/A	N/A	707	703	N/A	N/A
Second	N/A	N/A	795	842	N/A	N/A
Third	N/A	N/A	697	730	N/A	N/A

N/A: Not applicable.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 8.10N Survey Sample Size for Female Respondents Aged 15 to 44, by Pregnancy and Demographic Characteristics: 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total¹ (2011-2012)	Total¹ (2013-2014)	Pregnant (2011-2012)	Pregnant (2013-2014)	Not Pregnant (2011-2012)	Not Pregnant (2013-2014)
TOTAL	48,608	46,904	1,846	1,759	46,517	44,929
AGE						
15-17	11,514	9,788	125	79	11,315	9,671
18-25	23,704	20,013	1,234	1,019	22,353	18,885
26-44	13,390	17,103	487	661	12,849	16,373
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	40,136	38,230	1,503	1,421	38,429	36,647
White	29,215	27,396	1,026	964	28,048	26,331
Black or African American	6,608	6,257	333	275	6,233	5,942
American Indian or Alaska Native	698	642	27	39	666	598
Native Hawaiian or Other Pacific Islander	198	245	12	19	186	225
Asian	1,807	2,094	44	67	1,757	2,021
Two or More Races	1,610	1,596	61	57	1,539	1,530
Hispanic or Latino	8,472	8,674	343	338	8,088	8,282
TRIMESTER²						
First	N/A	N/A	566	519	N/A	N/A
Second	N/A	N/A	653	659	N/A	N/A
Third	N/A	N/A	613	558	N/A	N/A

N/A: Not applicable.

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 8.11A Numbers (in Thousands) of Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	Full-Time College Students (2013)	Full-Time College Students (2014)	Other Persons Aged 18-22² (2013)	Other Persons Aged 18-22² (2014)
TOTAL	21,906	21,965	8,659	8,757	13,244	13,203
AGE						
18	4,733	4,605	1,586	1,666	3,147	2,934
19	4,166	4,140	2,065	1,948	2,101	2,192
20	4,295	4,361	1,981	1,957	2,314	2,404
21	4,237	4,347	1,730	1,849	2,506	2,498
22	4,475	4,512	1,298	1,337	3,175	3,175
GENDER						
Male	11,113	11,322	4,016	4,130	7,097	7,186
Female	10,793	10,643	4,643	4,627	6,147	6,016
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	17,172	17,177	7,221	7,189	9,947	9,988
White	12,149	12,009	5,201	5,017	6,945	6,992
Black or African American	3,112	3,192	1,022	1,064	2,090	2,128
American Indian or Alaska Native	130	138	25	45	104	93
Native Hawaiian or Other Pacific Islander	132	106	53	34	78	72
Asian	1,143	1,211	711	809	432	402
Two or More Races	506	522	208	220	298	301
Hispanic or Latino	4,734	4,788	1,437	1,568	3,297	3,215
CURRENT EMPLOYMENT						
Full-Time	6,301	6,928	1,378	1,499	4,923	5,429
Part-Time	7,195	6,996	3,745	3,861	3,446	3,135
Unemployed	3,001	2,734	734	681	2,267	2,053
Other ³	5,409	5,307	2,802	2,716	2,607	2,586

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.11N Survey Sample Size for Respondents Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	Full-Time College Students (2013)	Full-Time College Students (2014)	Other Persons Aged 18-22² (2013)	Other Persons Aged 18-22² (2014)
TOTAL	13,893	10,171	5,447	4,001	8,444	6,168
AGE						
18	2,933	2,051	1,015	724	1,918	1,326
19	2,617	1,941	1,286	918	1,331	1,023
20	2,746	2,035	1,198	900	1,548	1,135
21	2,758	2,057	1,117	865	1,640	1,192
22	2,839	2,087	831	594	2,007	1,492
GENDER						
Male	6,820	5,034	2,439	1,827	4,381	3,205
Female	7,073	5,137	3,008	2,174	4,063	2,963
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	11,272	8,077	4,666	3,340	6,604	4,737
White	8,017	5,657	3,458	2,388	4,557	3,269
Black or African American	1,875	1,328	606	454	1,269	874
American Indian or Alaska Native	175	176	26	40	149	136
Native Hawaiian or Other Pacific Islander	74	55	21	17	53	38
Asian	594	464	358	299	236	165
Two or More Races	537	397	197	142	340	255
Hispanic or Latino	2,621	2,094	781	661	1,840	1,431
CURRENT EMPLOYMENT						
Full-Time	4,171	3,311	898	701	3,273	2,609
Part-Time	4,542	3,252	2,403	1,789	2,137	1,463
Unemployed	1,831	1,211	417	281	1,414	930
Other ³	3,349	2,397	1,729	1,230	1,620	1,166

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.12A Numbers (in Thousands) of Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	237,499	240,248	4,501	4,406	232,446	235,284
AGE						
18-49	132,643	133,085	3,780	3,697	128,533	128,975
18-25	34,786	34,935	1,497	1,309	33,186	33,508
26-34	37,346	37,880	1,089	1,263	36,209	36,474
35-49	60,511	60,270	1,195	1,125	59,138	58,992
GENDER/AGE						
Male 18 or Older	114,412	115,741	3,231	3,004	110,871	112,407
Male 18-49	65,526	65,734	2,688	2,466	62,669	63,012
Female 18 or Older	123,087	124,507	1,270	1,403	121,575	122,877
Female 18-49	67,117	67,351	1,092	1,231	65,864	65,963

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.12N Survey Sample Size for Respondents Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	45,344	50,894	1,442	1,340	43,789	49,425
AGE						
18-49	37,582	40,291	1,390	1,257	36,093	38,923
18-25	22,214	16,449	987	655	21,163	15,745
26-34	6,310	10,252	210	342	6,092	9,880
35-49	9,058	13,590	193	260	8,838	13,298
GENDER/AGE						
Male 18 or Older	21,294	23,789	998	881	20,240	22,834
Male 18-49	17,770	18,940	959	823	16,762	18,052
Female 18 or Older	24,050	27,105	444	459	23,549	26,591
Female 18-49	19,812	21,351	431	434	19,331	20,871

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.13A Numbers (in Thousands) of Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	237,499	240,248	1,679	1,452	235,319	238,303
AGE						
18-49	132,643	133,085	1,340	1,167	131,011	131,558
18-25	34,786	34,935	400	310	34,298	34,530
26-34	37,346	37,880	428	416	36,865	37,348
35-49	60,511	60,270	513	441	59,848	59,680
GENDER/AGE						
Male 18 or Older	114,412	115,741	1,349	1,098	112,801	114,366
Male 18-49	65,526	65,734	1,053	872	64,335	64,651
Female 18 or Older	123,087	124,507	330	354	122,518	123,937
Female 18-49	67,117	67,351	288	295	66,676	66,907

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.13N Survey Sample Size for Respondents Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	45,344	50,894	462	419	44,792	50,363
AGE						
18-49	37,582	40,291	432	384	37,075	39,811
18-25	22,214	16,449	274	159	21,895	16,247
26-34	6,310	10,252	77	124	6,225	10,105
35-49	9,058	13,590	81	101	8,955	13,459
GENDER/AGE						
Male 18 or Older	21,294	23,789	355	306	20,891	23,420
Male 18-49	17,770	18,940	330	279	17,398	18,606
Female 18 or Older	24,050	27,105	107	113	23,901	26,943
Female 18-49	19,812	21,351	102	105	19,677	21,205

NOTE: Because of a change in the sample allocation by age group, the sample size distribution for 2014 is different than the distribution for prior years. In 2013, for example, the sample was allocated equally among those aged 12 to 17, 18 to 25, and 26 or older. In 2014, the sample was allocated to these three age groups in proportions of 25, 25, and 50 percent, respectively. See Section A.1 in the 2014 NSDUH summary of methods for national estimates and supporting information (CBHSQ, 2015).

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.1C Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Gender and Detailed Age Category: 2013 and 2014

Age Category	Total (2013)	Total (2014)	Male (2013)	Male (2014)	Female (2013)	Female (2014)
TOTAL	0	0	0	0	0	0
12	95	97	66	69	63	68
13	106	99	73	72	68	67
14	106	96	76	71	68	69
15	94	105	65	73	65	72
16	102	109	71	73	67	72
17	97	104	66	74	67	70
18	157	166	94	97	103	119
19	129	128	86	92	82	85
20	129	138	89	94	83	96
21	123	140	81	93	87	92
22	130	142	88	102	84	86
23	122	122	85	86	83	80
24	120	125	87	87	81	88
25	121	131	78	88	83	92
26-29	485	349	332	250	311	211
30-34	557	413	352	271	369	266
35-39	488	394	326	267	336	265
40-44	554	388	366	275	368	256
45-49	474	395	369	279	297	268
50-54	771	632	547	427	517	415
55-59	733	561	496	383	495	381
60-64	667	507	477	370	432	338
65 or Older	0	0	0	0	0	0

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.2C Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Age Group and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	0	0	0	0	0	0	0	0
GENDER								
Male	0	0	0	0	0	0	0	0
Female	0	0	0	0	0	0	0	0
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0	0	0	0	0	0	0	0
White	0	0	247	221	343	337	1,901	1,594
Black or African American	0	0	118	124	169	180	813	641
American Indian or Alaska Native	120	130	19	22	26	22	108	115
Native Hawaiian or Other Pacific Islander	102	110	16	17	29	24	91	100
Asian	703	613	85	83	133	130	635	552
Two or More Races	239	174	41	43	46	46	230	159
Hispanic or Latino	0	0	0	0	0	0	0	0
GENDER/RACE/HISPANIC ORIGIN								
Male, White, Not Hispanic	1,209	1,026	153	142	201	213	1,126	939
Female, White, Not Hispanic	1,315	1,111	142	140	207	215	1,236	1,034
Male, Black, Not Hispanic	530	440	72	78	93	116	491	382
Female, Black, Not Hispanic	602	507	72	78	112	122	537	439
Male, Hispanic	0	0	108	99	141	142	580	462
Female, Hispanic	0	0	98	93	120	127	530	425
EDUCATION¹								
< High School	780	657	N/A	N/A	154	139	746	629
High School Graduate	1,091	897	N/A	N/A	242	252	1,048	836
Some College	998	830	N/A	N/A	262	265	947	788
College Graduate	1,443	1,125	N/A	N/A	184	170	1,410	1,094
CURRENT EMPLOYMENT¹								
Full-Time	1,629	1,269	N/A	N/A	245	257	1,576	1,226
Part-Time	692	543	N/A	N/A	221	214	648	491
Unemployed	360	284	N/A	N/A	119	117	335	254
Other ²	1,216	1,119	N/A	N/A	220	205	1,201	1,111

N/A: Not applicable.

¹ Estimates for education and current employment are shown only for persons aged 18 or older.² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.3C Standard Errors of Numbers (in Thousands) of Persons Aged 18 or Older, by Age Group and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26-49 (2013)	Aged 26-49 (2014)	Aged 50+ (2013)	Aged 50+ (2014)
TOTAL	0	0	0	0	0	0	0	0
GENDER								
Male	0	0	0	0	0	0	0	0
Female	0	0	0	0	0	0	0	0
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0	0	0	0	0	0	0	0
White	1,977	1,695	343	337	986	726	1,479	1,312
Black or African American	882	720	169	180	438	350	607	474
American Indian or Alaska Native	115	122	26	22	78	67	75	75
Native Hawaiian or Other Pacific Islander	96	105	29	24	58	64	70	74
Asian	673	577	133	130	371	363	490	353
Two or More Races	234	166	46	46	122	79	198	139
Hispanic or Latino	0	0	0	0	0	0	0	0
EDUCATION								
< High School	780	657	154	139	419	317	628	514
High School Graduate	1,091	897	242	252	577	415	850	707
Some College	998	830	262	265	571	451	735	641
College Graduate	1,443	1,125	184	170	822	661	1,071	834
CURRENT EMPLOYMENT								
Full-Time	1,629	1,269	245	257	1,041	827	1,085	837
Part-Time	692	543	221	214	388	276	521	426
Unemployed	360	284	119	117	245	175	227	184
Other ¹	1,216	1,119	220	205	418	306	1,133	1,045

¹ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.4C Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Youth and Adult Age Groups and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18+ (2013)	Aged 18+ (2014)
TOTAL	0	0	0	0	0	0
GENDER						
Male	0	0	0	0	0	0
Female	0	0	0	0	0	0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0	0	0	0	0	0
White	0	0	247	221	1,977	1,695
Black or African American	0	0	118	124	882	720
American Indian or Alaska Native	120	130	19	22	115	122
Native Hawaiian or Other Pacific Islander	102	110	16	17	96	105
Asian	703	613	85	83	673	577
Two or More Races	239	174	41	43	234	166
Hispanic or Latino	0	0	0	0	0	0
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	1,209	1,026	153	142	1,162	992
Female, White, Not Hispanic	1,315	1,111	142	140	1,272	1,081
Male, Black, Not Hispanic	530	440	72	78	511	415
Female, Black, Not Hispanic	602	507	72	78	575	481
Male, Hispanic	0	0	108	99	603	498
Female, Hispanic	0	0	98	93	559	467
EDUCATION¹						
< High School	780	657	N/A	N/A	780	657
High School Graduate	1,091	897	N/A	N/A	1,091	897
Some College	998	830	N/A	N/A	998	830
College Graduate	1,443	1,125	N/A	N/A	1,443	1,125
CURRENT EMPLOYMENT¹						
Full-Time	1,629	1,269	N/A	N/A	1,629	1,269
Part-Time	692	543	N/A	N/A	692	543
Unemployed	360	284	N/A	N/A	360	284
Other ²	1,216	1,119	N/A	N/A	1,216	1,119

N/A: Not applicable.

¹ Estimates for education and current employment are shown only for persons aged 18 or older.

² The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.5C Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Underage and Legal Drinking Age Groups and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
TOTAL	0	0	470	429	2,312	1,908
GENDER						
Male	0	0	284	267	1,444	1,183
Female	0	0	274	281	1,483	1,211
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0	0	405	368	2,181	1,838
White	0	0	335	306	1,950	1,651
Black or African American	0	0	161	181	853	687
American Indian or Alaska Native	120	130	28	27	112	120
Native Hawaiian or Other Pacific Islander	102	110	22	24	94	103
Asian	703	613	134	121	657	570
Two or More Races	239	174	52	56	232	163
Hispanic or Latino	0	0	223	209	879	745
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	1,209	1,026	204	190	1,149	970
Female, White, Not Hispanic	1,315	1,111	192	197	1,258	1,059
Male, Black, Not Hispanic	530	440	98	111	502	403
Female, Black, Not Hispanic	602	507	100	108	559	467
Male, Hispanic	0	0	141	143	589	482
Female, Hispanic	0	0	129	125	547	453

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.6C Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Age Group and Geographic Characteristics: 2013 and 2014

Geographic Characteristic	Total (2013)	Total (2014)	Aged 12-17 (2013)	Aged 12-17 (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26+ (2013)	Aged 26+ (2014)
TOTAL	0	0	0	0	0	0	0	0
GEOGRAPHIC DIVISION								
Northeast	0	0	0	0	0	0	0	0
New England	0	0	0	0	0	0	0	0
Middle Atlantic	0	0	0	0	0	0	0	0
Midwest	0	0	0	0	0	0	0	0
East North Central	0	0	0	0	0	0	0	0
West North Central	0	0	0	0	0	0	0	0
South	0	0	0	0	0	0	0	0
South Atlantic	0	0	0	0	0	0	0	0
East South Central	0	0	0	0	0	0	0	0
West South Central	0	0	0	0	0	0	0	0
West	0	0	0	0	0	0	0	0
Mountain	0	0	0	0	0	0	0	0
Pacific	0	0	0	0	0	0	0	0
COUNTY TYPE								
Large Metro	2,051	1,737	270	252	397	360	1,817	1,564
Small Metro	1,706	1,360	201	164	296	276	1,472	1,191
250K – 1 Mil. Pop.	1,437	1,098	177	134	227	215	1,215	974
< 250K Pop.	1,093	878	109	99	212	172	952	741
Nonmetro	1,182	1,117	127	126	194	191	1,035	938
Urbanized	794	796	87	83	156	142	686	665
Less Urbanized	947	862	93	104	122	132	812	705
Completely Rural	449	397	60	37	62	45	370	336

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.7C Standard Errors of Numbers (in Thousands) of Persons Aged 18 or Older, by Age Group and Geographic Characteristics: 2013 and 2014

Geographic Characteristic	Total (2013)	Total (2014)	Aged 18-25 (2013)	Aged 18-25 (2014)	Aged 26-49 (2013)	Aged 26-49 (2014)	Aged 50+ (2013)	Aged 50+ (2014)
TOTAL	0	0	0	0	0	0	0	0
GEOGRAPHIC DIVISION								
Northeast	0	0	0	0	0	0	0	0
New England	0	0	0	0	0	0	0	0
Middle Atlantic	0	0	0	0	0	0	0	0
Midwest	0	0	0	0	0	0	0	0
East North Central	0	0	0	0	0	0	0	0
West North Central	0	0	0	0	0	0	0	0
South	0	0	0	0	0	0	0	0
South Atlantic	0	0	0	0	0	0	0	0
East South Central	0	0	0	0	0	0	0	0
West South Central	0	0	0	0	0	0	0	0
West	0	0	0	0	0	0	0	0
Mountain	0	0	0	0	0	0	0	0
Pacific	0	0	0	0	0	0	0	0
COUNTY TYPE								
Large Metro	1,938	1,640	397	360	1,082	893	1,361	1,204
Small Metro	1,581	1,283	296	276	770	553	1,089	895
250K – 1 Mil. Pop.	1,321	1,037	227	215	664	447	865	761
< 250K Pop.	1,028	823	212	172	416	362	743	510
Nonmetro	1,114	1,042	194	191	445	414	811	654
Urbanized	748	745	156	142	300	295	516	438
Less Urbanized	883	787	122	132	331	305	616	495
Completely Rural	407	369	62	45	162	119	264	251

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.8C Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Underage and Legal Drinking Age Groups and Geographic Characteristics: 2013 and 2014

Geographic Characteristic	Total (2013)	Total (2014)	Aged 12-20 (2013)	Aged 12-20 (2014)	Aged 21+ (2013)	Aged 21+ (2014)
TOTAL	0	0	470	429	2,312	1,908
GEOGRAPHIC DIVISION						
Northeast	0	0	163	187	940	706
New England	0	0	84	106	467	304
Middle Atlantic	0	0	139	154	817	641
Midwest	0	0	183	157	880	741
East North Central	0	0	156	137	708	645
West North Central	0	0	96	85	504	338
South	0	0	302	248	1,647	1,176
South Atlantic	0	0	222	172	1,385	849
East South Central	0	0	99	91	555	391
West South Central	0	0	170	158	779	690
West	0	0	273	235	1,185	983
Mountain	0	0	118	99	543	414
Pacific	0	0	246	215	1,058	910
COUNTY TYPE						
Large Metro	2,051	1,737	378	358	1,897	1,619
Small Metro	1,706	1,360	281	253	1,537	1,244
250K – 1 Mil. Pop.	1,437	1,098	237	203	1,285	1,009
< 250K Pop.	1,093	878	179	156	989	789
Nonmetro	1,182	1,117	197	179	1,078	1,008
Urbanized	794	796	146	126	720	721
Less Urbanized	947	862	136	144	851	753
Completely Rural	449	397	86	48	390	360

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.9C Standard Errors of Numbers (in Thousands) of Persons Aged 12 or Older, by Youth and Adult Age Groups and Gender: 2002-2014

Demographic Characteristic	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
TOTAL	0	0	0	0	0	0	0	0	0	0	0	0	0
Male	0	0	0	0	0	0	0	0	0	0	0	0	0
Female	0	0	0	0	0	0	0	0	0	0	0	0	0
12-17	0	0	0	0	0	0	0	0	0	0	0	0	0
Male	0	0	0	0	0	0	0	0	0	0	0	0	0
Female	0	0	0	0	0	0	0	0	0	0	0	0	0
12-20	437	428	393	420	403	458	449	465	467	508	501	470	429
18 or Older	0	0	0	0	0	0	0	0	0	0	0	0	0
18-25	0	0	0	0	0	0	0	0	0	0	0	0	0
26 or Older	0	0	0	0	0	0	0	0	0	0	0	0	0
26-49	0	0	0	0	0	0	0	0	0	0	0	0	0
26-34	0	0	0	0	0	0	0	0	0	0	0	0	0
35-49	0	0	0	0	0	0	0	0	0	0	0	0	0
50 or Older	0	0	0	0	0	0	0	0	0	0	0	0	0

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: National Findings*).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2014.

Table 8.10C Standard Errors of Numbers (in Thousands) of Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Annual Averages Based on 2011-2012 and 2013-2014

Demographic/Pregnancy Characteristic	Total¹ (2011-2012)	Total¹ (2013-2014)	Pregnant (2011-2012)	Pregnant (2013-2014)	Not Pregnant (2011-2012)	Not Pregnant (2013-2014)
TOTAL	667	553	90	82	650	541
AGE						
15-17	92	92	7	6	91	91
18-25	0	0	33	32	245	205
26-44	538	467	82	75	529	458
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	568	487	78	76	555	474
White	470	392	64	59	457	384
Black or African American	256	218	31	37	248	208
American Indian or Alaska Native	29	31	7	3	28	30
Native Hawaiian or Other Pacific Islander	39	27	6	5	39	27
Asian	178	170	25	25	177	163
Two or More Races	59	58	8	12	58	57
Hispanic or Latino	290	262	42	34	285	258
TRIMESTER²						
First	N/A	N/A	47	47	N/A	N/A
Second	N/A	N/A	52	50	N/A	N/A
Third	N/A	N/A	49	46	N/A	N/A

N/A: Not applicable.

¹ Estimates in the Total column represent all females in the specified subgroup, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011-2014.

Table 8.11C Standard Errors of Numbers (in Thousands) of Persons Aged 18 to 22, by College Enrollment Status and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	Full-Time College Students (2013)	Full-Time College Students (2014)	Other Persons Aged 18-22² (2013)	Other Persons Aged 18-22² (2014)
TOTAL	372	359	280	281	234	229
AGE						
18	157	166	124	128	95	108
19	129	128	97	95	82	87
20	129	138	94	96	85	96
21	123	140	88	106	85	91
22	130	142	70	80	103	115
GENDER						
Male	225	232	155	164	160	170
Female	229	232	172	177	141	148
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	329	318	253	250	188	194
White	256	261	199	190	151	164
Black or African American	115	137	68	84	87	102
American Indian or Alaska Native	17	19	7	12	15	14
Native Hawaiian or Other Pacific Islander	24	18	14	10	19	16
Asian	103	101	91	89	40	44
Two or More Races	37	37	25	26	28	26
Hispanic or Latino	157	160	77	91	131	124
CURRENT EMPLOYMENT						
Full-Time	146	171	63	71	135	152
Part-Time	179	181	137	146	103	106
Unemployed	100	98	52	52	84	85
Other ³	192	184	166	158	91	97

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown enrollment status.

² Other Persons include respondents aged 18 to 22 not enrolled in school, enrolled in college part time, enrolled in other grades either full or part time, or enrolled with no other information available.

³ The Other Employment category includes students, persons keeping house or caring for children full time, retired or disabled persons, or other persons not in the labor force.

Table 8.12C Standard Errors of Numbers (in Thousands) of Persons Aged 18 or Older, by Past Year Probation Status and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Probation (2013)	On Probation (2014)	Not on Probation (2013)	Not on Probation (2014)
TOTAL 18 OR OLDER	0	0	220	175	2,340	1,907
AGE						
18-49	0	0	173	139	1,501	1,115
18-25	0	0	70	64	490	445
26-34	0	0	97	83	788	569
35-49	0	0	112	94	944	729
GENDER/AGE						
Male 18 or Older	0	0	184	149	1,460	1,190
Male 18-49	0	0	146	118	914	689
Female 18 or Older	0	0	110	90	1,498	1,219
Female 18-49	0	0	87	74	919	689

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown probation status in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.

Table 8.13C Standard Errors of Numbers (in Thousands) of Persons Aged 18 or Older, by Past Year Parole/Supervised Release Status and Demographic Characteristics: 2013 and 2014

Demographic Characteristic	Total¹ (2013)	Total¹ (2014)	On Parole/Supervised Release² (2013)	On Parole/Supervised Release² (2014)	Not on Parole/Supervised Release² (2013)	Not on Parole/Supervised Release² (2014)
TOTAL 18 OR OLDER	0	0	136	105	2,344	1,922
AGE						
18-49	0	0	103	83	1,502	1,122
18-25	0	0	33	31	500	450
26-34	0	0	63	47	795	573
35-49	0	0	73	56	942	734
GENDER/AGE						
Male 18 or Older	0	0	128	95	1,462	1,192
Male 18-49	0	0	94	71	918	691
Female 18 or Older	0	0	47	49	1,498	1,225
Female 18-49	0	0	42	42	919	696

¹ Estimates in the Total column represent all persons in the specified subgroup, including those with unknown parole/supervised release status in the past year.

² Respondents were asked if they were on parole, supervised release, or other conditional release from prison during the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014.